Multiscale modeling of presynaptic dynamics from molecular to mesoscale

Authors

Jonathan W. Garcia[1,2,#*], Thomas M. Bartol[2], Terrence J. Sejnowski[1,2]

Affiliations

[1] Division of Biological Sciences, University of California San Diego, La Jolla, California, United States of America

[2] Computational Neurobiology Laboratory, Salk Institute for Biological Studies, La Jolla, California, United States of America

[#] Current Affiliation: Independent Researcher

* Corresponding author

E-mail: soiens24@gmail.com (JWG)

Abstract

Chemical synapses exhibit a diverse array of internal mechanisms that affect the dynamics of transmission efficacy. Many of these processes, such as release of neurotransmitter and vesicle recycling, depend strongly on activity-dependent influx and accumulation of Ca^{2+}. To model how each of these processes may affect the processing of information in neural circuits, and how their dysfunction may lead to disease states, requires a computationally efficient modelling framework, capable of generating accurate phenomenology without incurring a heavy computational cost per synapse. Constructing a phenomenologically realistic model requires the precise characterization of the timing and probability of neurotransmitter release. Difficulties arise in that functional forms of instantaneous release rate can be difficult to extract from noisy data without running many thousands of trials, and in biophysical synapses, facilitation of per-vesicle release probability is confounded by depletion. To overcome this, we obtained traces of free Ca^{2+} concentration in response to various action potential stimulus trains from a molecular MCell model of a hippocampal mossy fiber axon. Ca^{2+} sensors were placed at varying distance from a voltage-dependent calcium channel (VDCC) cluster, and Ca^{2+} was buffered by calbindin. Then, using the calcium traces to drive deterministic state vector models of synaptotagmin 1 and 7 (Syt-1/7), which respectively mediate synchronous and asynchronous release in excitatory hippocampal synapses, we obtained high-resolution profiles of instantaneous release rate, to which we applied functional fits. Synchronous vesicle release occurred predominantly within half a micron of the source of spike-evoked Ca^{2+} influx, while asynchronous release occurred more consistently at all distances. Both fast and slow mechanisms exhibited multi-exponential release rate curves, whose magnitudes decayed exponentially with distance from the Ca^{2+} source. Profile parameters facilitate on different time scales according to a single, general facilitation function. These functional descriptions lay the groundwork for efficient mesoscale modelling of vesicular release dynamics.

Author Summary

Most information transmission between neurons in the brain occurs via release of neurotransmitter from synaptic vesicles. In response to a presynaptic spike, calcium influx at the active zone of a synapse can trigger the release of neurotransmitter with a certain probability. These stochastic release events may occur immediately after a spike or with some delay. As calcium accumulates from one spike to the next, the probability of release may increase (facilitate) for subsequent spikes. This process, known as short-term plasticity, transforms the spiking code to a release code, underlying much of the brain's information processing. In this paper, we use an accurate, detailed model of presynaptic molecular physiology to characterize these processes at high precision in response to various spike trains. We then apply model reduction to the results to obtain a phenomenological model of release timing, probability, and facilitation, which can perform as accurately as the molecular model but with far less computational cost. This mesoscale model of spike-evoked release and facilitation helps to bridge the gap between microscale molecular dynamics and macroscale information processing in neural circuits. It can thus benefit large scale modelling of neural circuits, biologically inspired machine learning models, and the design of neuromorphic chips.

Introduction

Chemical synapses constitute the primary means of direct communication between neurons throughout the nervous system [1-3]. Neurotransmitters are stored in synaptic vesicles, which are docked to the plasma membrane of the axon terminal by soluble N-ethylmaieimide-sensitive factor attachment protein receptor (SNARE) complexes. Vesicle-membrane-bound synaptobrevin (v-SNARE) and the target-membrane-bound syntaxin and SNAP-25 (t-SNAREs) form energetic SNAREpin complexes where the α-helices of the v-SNAREs entwine with those of the t-SNAREs [4-6]. Synaptotagmin (Syt) proteins

embedded in both membranes associate with the SNARE complex and act as Ca^{2+}-sensitive triggers for

vesicle fusion. When the action potential of the presynaptic neuron reaches the axon terminal, it

triggers a sudden influx of Ca^{2+} through voltage-dependent Ca^{2+} channels (VDCCs), and when a sufficient

number Ca^{2+} ions binds to the C_2 domains of synaptotagmin, it undergoes a conformational change that

triggers the associated SNAREpin to zipper completely, causing the vesicle to fuse with the membrane

and to release its neurotransmitter through the newly opened fusion pore [7-11] (see Fig 1).

Fig 1. SNARE Complex Structure and Dynamics.
73   A: SNAREpins prior to vesicle fusion. B: Binding of Ca^{2+} to synaptotagmin (Syt-1 here, Syt-7 attaches to
74   target membrane [12, 13]) triggers full zippering of SNARE complex and, in turn, vesicle fusion [14, 15].
75

Very often, discussion of the activity in a network tends to focus on the action potentials (spikes) and

subthreshold fluctuations in membrane potential [16-18]. The utility of these measurements, however,

depends on the relevance of the spike code to neural information processing. How neurons integrate

their inputs and generate signals in the context of larger neural circuits largely determines the sorts of

computations that the network can perform [19, 20]. Biological neural networks need to represent

information in a way that confers behavioral utility, but since so much of the information in the

environment is irrelevant to survival, synapses may not be optimized to transmit all information

faithfully, but rather selectively.

Significantly, neurons do not directly see the spiking activity of their neighbors at chemical synapses, but

only detect presynaptic activation upon the release of neurotransmitter, which is a stochastic process

[21]. Synapses form the basis for learning and information processing, and short-term plasticity (STP)

defines a transformation from a spiking code to a neurotransmitter release code. All spiking activity is

filtered through the dynamics of probabilistic synaptic release before the rest of the network can see it.

This implies that one must first have an accurate model of release dynamics in order to understand the

true nature of information processing of brain circuits. Such a model could, for instance, provide a

crucial preprocessing step of motor cortex for training BCI-based prosthetics [22-24], or it could enable

more accurate computation of the information capacity of sensory cortex by studying the "language"

that neurons actually receive rather than simply the output that they generate [19, 25-28].

Synaptic dysfunction has been implicated in numerous psychological disorders, including schizophrenia

[29, 30], bipolar disorder [29], ASD [31], and fragile X syndrome [32]. To ascertain exactly what role

synapses play and what specific mechanisms might be causing or exacerbating these diseases,

controlled experiments would need to be performed on the brain circuits of interest, testing which

changes to synaptic function might push the network into a pathological state. Doing this in humans

would pose significant problems, both technical and ethical. However, with a computational model that

exhibits sufficient realism and scalability, such experiments become possible in large simulated

networks, which could provide important insight into what sorts of targeted therapies to explore for

treating these diseases.

The molecular simulator MCell can track the kinetics and interactions of thousands of molecules and

ions in a three-dimensional model of the synapse, achieving a high degree of realism and elucidating

how complex bimolecular systems may function in the absence of experimental interventions [33-

37].When properly constrained by experimental data, MCell will not only automatically reproduce

observed features such as asynchronous vesicle release, facilitation, and depression in the probability of

release [35], but it can also make surprising predictions that are later confirmed through experiment

[38, 39]. However, it quickly becomes too computationally expensive to scale up to the many synapses

that exist even in relatively simple neural circuits.

The ultimate goal, therefore, is to develop a presynaptic model that captures realistic phenomenology

while maintaining computational scalability. To that end, in this paper we develop a mathematical

model that describes the phenomenology of presynaptic dynamics, using the MCell model of [35] as a

reasonable approximation to ground truth. Our simplified model takes an arbitrary spike train and uses

it to approximate the single-vesicle release rate histograms that would emerge from running an infinite

number of MCell simulations, taking into account the facilitation in release rate of both synchronous and

asynchronous release dynamics. The results are easily amenable to event-driven models, producing the

generating distributions for temporally asynchronous vesicle release times with respect to arbitrary

sequences of action potentials. Thus, it provides an avenue for large-scale simulations of spiking neural

networks with many realistically performing synapses without incurring high computational costs,

enabling investigations into how various presynaptic mechanisms can affect computation at the circuit

level.

124 Results

125 **Ca^{2+}-Evoked Dynamics of Vesicle Release**

Diffusion plays a key role in presynaptic processes. Simplified models of Ca^{2+}-dependent presynaptic

dynamics may assume that the axon terminal is locally well mixed, equivalent to saying that diffusion

happens infinitely fast, at least relative to the spatial and temporal scales being studied. However, MCell

allows one to add a spatial component to molecular simulations, which can account for certain

phenomena that well mixed molecular kinetics models cannot capture [33, 34, 36]. To characterize the

process of Ca^{2+}-dependent neurotransmitter release, we based two models on the presynaptic model of

[35]: a spatially explicit model implemented in MCell, and an equivalent "well-mixed" model. Both

models contained voltage-dependent Ca^{2+} channels (VDCCs) that let Ca^{2+} ions into the presynaptic

volume in response to an action potential stimulus, a calbindin (CB) buffer that moderated diffusion of

the ions via its binding kinetics, and plasma membrane Ca^{2+}-ATPase (PMCA) pumps that helped

intracellular Ca^{2+} concentration return to equilibrium over time (see S1 Fig for state transition diagrams).

However, the two models differed in that the MCell model relied heavily on diffusion of molecular

species through space and discrete state transition events occurring over time, while the well-mixed

model treated all molecular interactions as occurring within the same point in space and tracked

continuous state probabilities over time (see Methods).

Comparing these models, we found that the diffusion of Ca^{2+} and calbindin through the axonal volume

affects both the timing and the probability of spike-evoked vesicle release, depending on the distance

from the Ca^{2+} source. Fig 2 compares the well-mixed simulation without diffusion to the equivalent

MCell simulations performed at multiple distances from the VDCC Ca^{2+} source. The shape of the Ca^{2+}

transient measured in MCell displays marked qualitative differences from that obtained without

diffusion: Ca^{2+} sensors near the VDCC source see a much higher peak concentration with an extra

component of decay immediately following the peak; those farther away progressively lose the fast

peak until nothing is left but an extremely small distance-independent component. The extra

component of the proximal Ca^{2+} curve, which does not appear in the well-mixed model, likely arises

from local saturation in the nanodomains near the VDCC cluster, where the very high free $[Ca^{2+}]_i$

temporarily saturates both the calbindin buffer and the PMCA pumps [40]. Farther out, the MCell model

qualitatively matches the well-mixed model more closely, until at very large distances, the fast

components almost completely disappear. The distance-independent components represent a sustained

global elevation in $[Ca^{2+}]_i$ that persists due to the excess Ca^{2+} that has yet to unbind from the calbindin

buffer. The slowest component has a magnitude comparable to resting $[Ca^{2+}]_i$ and a time constant of

around 1 second.

Fig 2. Spatial Modeling of Spike-Evoked Ca^{2+} Transients.

A: Ca^{2+} sensors (dark yellow through dark blue filled circles) at vesicle cluster centers, displaced linearly

from cluster of Ca^{2+} channels (blue half-disk on the left); distance in µm, $d_n=0.160+0.105n$ for

$n\in\{0,...,16\}$. B: [Ca^{2+}]$_i$ measured over time in MCell (dark yellow through dark blue) and in the

deterministic well-mixed model (maroon). MCell traces averaged from 2000 trials of MCell simulations

with Δt=0.1 ms. Color transitions from yellow for vesicles proximal to the VDCC Ca^{2+} source to blue for

vesicles far away, as in A. Proximally (distally) measured [Ca^{2+}]$_i$ displays more (fewer) components of

decay than are evident in the deterministic model. C: Logarithmic plots of peak [Ca^{2+}]$_i$ (blue) and peak

time (red) as a function of distance from Ca^{2+} source; peak [Ca^{2+}]$_i$ drops off exponentially with distance

from VDCC cluster; amplitude of latent Ca^{2+} dominates over the initial action-potential-evoked influx

after 1.4 µm.

One would expect the strength of spike-evoked neurotransmitter release to diminish with increasing

distance from the Ca^{2+} source, where Ca^{2+} has more time to diffuse and bind to buffer molecules before

reaching the sensor. In fact, numerous studies have found that vesicles of the readily releasable pool

(RRP) fall into one of two subpopulations, depending on their physical location of vesicles within the

synapse: vesicles located very near Ca^{2+} channels release quickly in response to spikes, while those

farther away are more reluctant [41-44]. To explore how the release rate profiles vary with distance, we

established a linear array of Ca^{2+} sensors along the length of the model axon, with a cluster of 50 VDCCs

arranged in a half-disk at one end (Fig 2 A). Reflective boundaries on the ends of a 2-μm tube effectively

simulated the effects of having one cluster of 100 VDCCs every 4 μm, consistent with previous models of

the Schaffer collateral axon [35, 45]. Running the model again for 2000 trials, with a single action

potential stimulus applied at the beginning, we obtained Ca^{2+} traces measured at each point along the

axon. For the first 1.4 μm, free Ca^{2+} from the initial influx dominated, and the peak concentration

declined exponentially with distance (length constant 0.204 μm; Fig 2 C). Farther out, global

accumulation and depletion of Ca^{2+} dominates, which, although spike-evoked, does not vary in

magnitude with distance and acts over a much longer time scale and at a much lower level than most of

the spike-triggered Ca^{2+}.

Running these simulations in MCell, rather than as a much simpler well-mixed model, was essential for

capturing both distance-dependent effects and temporal features of the Ca^{2+} waveform. The well-mixed

assumption, which ignores diffusion and treated all chemical processes as occurring at the same point in

space, does not hold at the spatial and temporal scales of interest in the synapse [46, 47]. As seen in Fig

2 C, peak Ca^{2+} dropped precipitously even over fractions of a micron away from the VDCC cluster, and

the shape of the response changed dramatically over this same scale, transitioning from a

predominantly synchronous to a predominantly asynchronous profile, even before the Ca^{2+} sensors

started responding. These trends, elucidated by the spatial MCell model, were completely absent in the

space-less well-mixed simulation (Fig 2 B, maroon line), even when all other aspects of the model

remained the same, such as the number of VDCCs, calbindin buffer molecules, and PMCA pumps and

the set of all state transitions for each molecular species (see Methods for details). Note also from Fig 2

B that the transition in time from the fast synchronous component to the extended asynchronous

component was much sharper in the case without space. The extra Ca^{2+} decay component arose from

local saturation effects. After the initial rapid influx, the calbindin buffer immediately around the VDCC

cluster became saturated, causing the high free Ca^{2+} that remains to overwhelm the PMCA pumps'

ability to evacuate it from the area. The pumps removed it at a constant maximum rate, leading to a

short linear decay only evident very near the VDCCs (yellow traces, S4 Fig A) or when all calbindin is

removed from the simulation (S5 Fig B). Such effects did not appear in the well-mixed case because all

buffer molecules and pumps were simultaneously available to all the free Ca^{2+}, preventing any local

saturation from occurring. Thus, in light of all these effects, the spatial MCell model was crucial for the

task of properly characterizing the Ca^{2+} transient in the synapse.

Although many more proteins are involved in coordinating release kinetics at active zones [14, 15, 44,

48-52], for validation purposes we restrict the scope of this paper to the function of synaptotagmins.

The model of synaptotagmin-mediated release used in our simulations followed the dual Ca^{2+}-sensor

model of Sun et al. [53], which includes mechanisms for both fast/synchronous and slow/asynchronous

release. In excitatory hippocampal synapses, these synchronous and asynchronous modes of release

may correspond to the roles of synaptotagmin-1 (Syt-1) and synaptotagmin-7 (Syt-7), respectively [54-

59]. The model incorporates cooperative binding of Ca^{2+} to multiple sites on the sensor, requiring five

Ca^{2+} ions before triggering synchronous release and two Ca^{2+} ions for asynchronous release (see Fig 3).

Because both binding and unbinding rates for the synchronous mechanism are substantially higher than

those for the asynchronous mechanism, Syt-1 produces rapid release over a very narrow window

relative to spike arrival time, while Syt-7 produces slow release over a much more extended window.

Table 1 contains the values used in this model for Ca^{2+}-binding and unbinding rates with each release

mechanism, along with the rates of vesicle fusion from the fully bound states (γ_S and γ_A) and the time

constant for the post-release refractory period (ε) [60, 61], which features in the Nadkarni et al. [35]

MCell model.

Fig 3. Model of Ca^{2+}-Evoked, Synaptotagmin-Mediated Neurotransmitter Release.

A, B: Model adapted from Sun et al. [53]. γ_S and γ_A represent rates of vesicle fusion from the releasable states of the synchronous and asynchronous mechanisms, respectively. A: Ca^{2+}-bound states for Syt-1 (synchronous release); S_n indicates n Ca^{2+} ions bound to the synchronous release mechanism. B: Ca^{2+}-bound states for Syt-7 (asynchronous release); A_n indicates n Ca^{2+} ions bound to the asynchronous release mechanism. C, D: Action-potential-like stimulus delivered to model axon starting at 0 ms. Diffusion is assumed to be instantaneous, and molecular state probabilities are tracked deterministically over time. C: Free [Ca^{2+}]$_i$ in response to single action potential. D: Instantaneous vesicle release rate in response to buffered Ca^{2+} from both synaptotagmin-mediated release mechanisms.

Table 1. SNARE Release State Transition Parameters.

Values taken from Nadkarni et al. [35], adapted from Sun et al. [53].

synchronous		asynchronous		other parameters	
k_{S+}	6.12×10^7 M^{-1}s^{-1}	k_{A+}	3.82×10^6 M^{-1}s^{-1}	b	0.25
k_{S-}	2.32×10^3 s^{-1}	k_{A-}	13 s^{-1}	ε	6.34 ms
γ_S	6.0×10^3 s^{-1}	γ_A	50 s^{-1}		

The MCell model, because it tracks thousands of individual particles through Markov chain Monte Carlo

simulations [33, 34, 36], can both capture very realistic synaptic dynamics and uncover their underlying

molecular causes, which would be difficult to obtain through other methods. Unfortunately, this realism

can also obscure the patterns necessary for building simplified models. First, many processes, such as

asynchronous or "mini" release events [62-64], occur slowly enough that many thousands or millions of

simulated trials would be required to uncover precise functional descriptions, which could become

computationally prohibitive. For instance, the histograms of synchronous release obtained from 2000

trials of MCell in Fig 4 offer little information on spontaneous release from the Syt-1 mechanism

between action potentials, and synchronous release far from the VDCC cluster (blue) hardly occurs at all.

Second, the fact that vesicles deplete upon release hides how the instantaneous single-vesicle release

rate actually changes with time. The tails of the release distributions fall off too quickly as vesicles are

removed from the simulation over time, and any paired-pulse facilitation (PPF) in single-vesicle release

probability is countered by the release-dependent depletion in the model (Fig 4 D).

Fig 4. Synchronous and Asynchronous Release in MCell.

Color indicates distance from VDCC source, with yellow representing a nearby Ca^{2+} sensor and dark blue a distant one (as in Fig 2 A-B). Action-potential-like stimulus delivered at 0 ms (left), followed by another at 20 ms (center) and 100 ms (right). A: Spike-evoked Ca^{2+} traces that drive release. B: Synchronous release raster. C: Asynchronous release raster. D: Synchronous (tall, thin bars) and asynchronous (short, wide bars) release stacked histogram. Most synchronous releases happen close to the Ca^{2+} source; asynchronous releases distributed across all distances.

The only way to avoid these depletion effects in MCell would be to run many millions of trials with a

single vesicle to track how the vesicle's alacrity for release fluctuates with the Ca^{2+} history detected at its

position. For these reasons, we decided not to depend on the release histograms generated by many

trials of MCell for building a phenomenological model. Instead, we used the Ca^{2+} traces generated by

MCell, which do not suffer from the aforementioned problems, to drive deterministic simulations of the

SNARE state probability dynamics (as described in Methods), effectively producing what an infinite

number of trials would produce in MCell with the same Ca^{2+} data. Thus, using the deterministic release

rates driven by the stochastic MCell Ca^{2+} data balances the necessary realism of MCell with the

smoothness and insights required for designing a versatile phenomenological model.

Reducing from Molecular Simulations to Phenomenological Model

269   By driving a deterministic simulation of SNARE dynamics with the $[Ca^{2+}]_i$ waveform obtained from MCell,

270   one can see that each release mechanism induces vesicle fusion with a histogram that essentially

271   follows a multi-exponential form (Fig 5). The release rate profiles ($r(t)$, where $r \in \{S, A\}$ may refer to

272   synchronous or asynchronous release rate) rise quickly from baseline after the spike and decay with

273   several exponential components, approximated as

$$r(t) = r_0 + \sum_{c=1}^{N} \frac{P_c}{\tau_c} e^{-t/\tau_c} u(t), \tag{1}$$

where r_0 is the spontaneous release rate (related to "mini"-EPSCs [63-65]; $S_0 = 5.70 \times 10^{-9} ms^{-1}$;

$A_0 = 1.84 \times 10^{-5} ms^{-1}$), t is the time since the last spike, $u(t)$ is the Heaviside step function (so that

release occurs only for $t \geq 0$), N is the number of exponential decay components, τ_c are the time

constants of exponential decay, and P_c are the expected number of releases from each component for a

single vesicle. Note that because the release rate profile is not a probability distribution, but rather it

represents the instantaneous rate of release conditioned on having not released yet (see Methods), its

integral P_c can potentially exceed one (just as the integral over the spontaneous component r_0 for

$t = 0 \dots \infty$ is infinite). The probability that the exponential component causes release at any point in

time, ignoring the other components, is $p_{rc} = 1 - \exp(-P_c)$ (and the integrated release probability for

the spontaneous component is $p_{r0} = 1 - \exp(-\infty) = 1$). The existence of multiple exponential

components is apparent from the linear segments visible in log-linear space for synchronous and

asynchronous release rates in Fig 5 B. To calculate the time constants of exponential decay (Fig 5 C), we

used the slope of the logarithm of the release rate curve according to

$$\tau(t) = -\left(\frac{d}{dt}\left[\ln\left(r^*(t) - r^*(0)\right)\right]\right)^{-1}, \tag{2}$$

287 where $\tau(t)$ is the instantaneous time constant and $r^*(t)$ is the observed instantaneous release rate. We

288 used the well-mixed model for the derivative calculation because it had no noise in the release rate

289 profiles.

Fig 5. Multi-Exponential Shape of Ca^{2+}-Driven Vesicle Release Rate.
A, B: Plots given as semi-log to highlight exponential decay components (straight line segments of
profiles). A: A single, spike-evoked [Ca^{2+}]$_i$ transient, which drives B: the synchronous and asynchronous
release rates. C: Instantaneous time constants for Ca^{2+}, synchronous, and asynchronous curves,
calculated from the well-mixed model (see Eq (2)). Long release rate time constants (around 80 ms and
1000 ms; dashed lines) follow Ca^{2+} curve due to slow un-buffering of latent Ca^{2+}. Asynchronous starts
high because fast and slow components have comparable magnitude and become conflated; it goes up
to infinity where additive effects cause the curve to flatten.

Most of the Ca^{2+} that enters the axon following an action potential quickly binds with the calbindin

buffer before diffusing to the SNARE complex, causing a narrow spike in the free [Ca^{2+}]$_i$ available to the

release mechanism. Therefore, most of the spike-evoked release occurs in response to this narrow

window of influx. To test how each release mechanism responds to transient Ca^{2+} spikes, we supplied an

instantaneous burst of Ca^{2+} to a single time step of the deterministic model, allowing us to measure the

impulse-response function. These simulations were repeated for various resting Ca^{2+} levels ([Ca^{2+}]$_{i0}$),

ranging from 0 to 10 μM, to see how the presence of Ca^{2+} at equilibrium affects the response to spike-

evoked transients. As Fig 6 shows, when there is no resting $[Ca^{2+}]_i$, the rate of release for both

synchronous and asynchronous mechanisms rises quickly in response to a sudden influx before dropping

exponentially with a single exponential component (black). However, when $[Ca^{2+}]_{i0}$ settles at some level

greater than zero, an extra exponential component emerges for both mechanisms (blue and red lines).

The exponential decay time constants seem to be mostly independent of resting $[Ca^{2+}]_{i0}$ at low levels,

but they drop off more quickly as spontaneous release rates begin to overtake the spike-evoked rates at

high concentrations. The extra component emerges as a result of the back-and-forth Ca^{2+}-binding and

unbinding processes, where finite baseline $[Ca^{2+}]_{i0}$ likely provides a "floor" to "bounce off of" in terms of

the number of Ca^{2+} ions bound to the release mechanism. Note, however, that even though it depends

on equilibrium $[Ca^{2+}]_{i0}$, this secondary release component is still purely spike-evoked and arises due to

the nonlinearity of the system. The baseline rate $r^*(0)$ was subtracted off to ensure that the function

approached zero prior to taking the logarithm (dotted lines). Time constants in Fig 6 C were calculated

using Eq (2).

Fig 6. Synchronous and Asynchronous Release Rates in Response to Ca^{2+} Impulse at

Different Resting Concentrations.

Instantaneous impulse of Ca^{2+} delivered at 10 ms. Solid lines represent true release rate; dotted lines
have spontaneous rates subtracted off to show secondary exponential components. Black lines show
release rate decaying with a single exponential component with no baseline [Ca^{2+}]$_i$. For other curves,
[Ca^{2+}]$_{i0}$ ranges from 0.001 μM to 10 μM. A: Synchronous release rate over time: S(t). B: Asynchronous
release rate over time: A(t). C: Instantaneous release rate decay time constants for synchronous and
asynchronous mechanisms. Fast components (lower blue and red lines) determined from profiles with
[Ca^{2+}]$_{i0}$=0 (black lines in A and B). Slower components (upper blue and red curves) determined from
cases with small [Ca^{2+}]$_{i0}$.

From the above, it would seem that each mechanism should have three components to its release

histogram: a constant spontaneous rate that increases with [Ca^{2+}]$_{i0}$, a fast exponential component that

acts in response to an impulse of spike-evoked Ca^{2+}, and a slower spike-evoked component that results

from a "rebound" interaction with the Ca^{2+} floor. However, the profiles of the release rate histograms

display more complexity than this, which will be discussed in more detail below. Significantly, $[Ca^{2+}]_i$

does not drop instantly to baseline after the initial influx, but some leftover Ca^{2+} continues to have a

small effect over a long time window as it slowly unbinds from the calbindin buffer (see S5 Fig). This

allows a small but noticeably enhanced rate of release efficacy to continue out to hundreds or

thousands of milliseconds before returning fully to baseline (within noise). Fig 5 C shows the effect that

this latent Ca^{2+} has on producing longer time constants in the decay of the release rate profiles, using

the smooth curves obtained from the well-mixed model.

Of course, neurotransmitter release cannot begin at exactly the moment of the spike, both because the

action potential itself is not an instantaneous process and because it takes finite time for Ca^{2+} to diffuse

from the VDCC source, through the buffer, to the Ca^{2+} sensor in the SNARE complex. MCell represents

this complex process with a Markov chain Monte Carlo method (MCMC). Because of this, the release

process cannot begin until the spike-evoked Ca^{2+} arrives, which time may vary randomly relative to the

timing of the spike. Thus, the process of buffered diffusion acts as a temporal filter on the release

dynamics, transforming the equation of release to

$$r(t) = r_0 + \sum_{c=1}^{N} \frac{P_c}{\tau_c}\left(e^{-t/\tau_c}u(t)\right) * a(t; k_c, \mu_c, \sigma_c), \tag{3}$$

where $a(\cdot)$ is the temporal filter and k_c, μ_c, and σ_c are parameters to be discussed below. The

convolution operation effectively smears the start time of the average release profile in a way that

accounts for random temporal jitter across trials.

Importantly, the release-start-time filter $a(\cdot)$ must integrate to one over all real numbers. That way, it

does not affect the probability of release, only its timing. The temporal filter chosen is an ex-Gaussian

distribution, resulting from the convolution of an exponential distribution of rate k with a normal

distribution of mean μ and standard deviation σ:

$$a(t; k, \mu, \sigma) = \left(k\mathrm{e}^{-kt}\mathrm{u}(t)\right) * \left(\frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(t-\mu)^2}{2\sigma^2}}\right)$$

$$= \int_{-\infty}^{t} k\mathrm{e}^{-k(t-t')} \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(t'-\mu)^2}{2\sigma^2}} \mathrm{d}t' \tag{4}$$

$$= k\mathrm{e}^{-k\left(t-\left(\mu+\frac{\sigma^2}{2}k\right)\right)} \Phi\left(\frac{t-(\mu+\sigma^2 k)}{\sigma}\right),$$

where $\Phi(\cdot)$ represents the CDF of the zero-mean, unit-variance normal distribution. In the limit where

$\sigma \to 0$, this CDF simply becomes the shifted Heaviside step function $\mathrm{u}(t-\mu)$, and $a(t; k, \mu, \sigma) \to$

$k\mathrm{e}^{-k(t-\mu)}\mathrm{u}(t-\mu)$, which is just a rightward shift in time of the exponential distribution by μ. The values

of μ and σ result from the sum of the delays caused by numerous random processes, including the

timing of Ca^{2+} entry relative to the spike, the accumulation of collision events during Brownian motion,

and the binding/unbinding events with the calbindin buffer and SNARE complex. Assuming that the

individual events of the buffered diffusion process are numerous and similar enough for a given spike,

the central limit theorem states that the sum of their delays should approximate a normal distribution

[66]. The value of k represents the rate of some limiting step in the process of buffered diffusion, and it

slows with increasing distance between the VDCC source and the Ca^{2+} sensor in the SNARE complex.

Keep in mind that these parameters constitute only a phenomenological approximation to the exact

filter, but they work well enough for the purposes of this paper.

For an event-driven model, which this paper is working towards, the convolutional operation can be

implemented by sampling a normally distributed random number (with mean and standard deviation μ_c

and σ_c) and an exponentially distributed random number (with rate k_c) and adding them to the spike

time to determine the start time for the release response. In other words, combine the spike time with

an ex-Gaussian random delay to determine when the release component begins to respond to the spike,

following Eq (1). In aggregate, across many trials with the same spike time, the release histogram will

approach Eq (3).

With the mathematical description of the release rate profiles in mind, we ran a fitting algorithm (see

Methods) to determine the values of the parameters for each profile. Initially, we used release profiles

driven by Ca^{2+} measured at 400 nm from the cluster of 100 VDCCs, which provides a physiologically

realistic probability of release for a single vesicle (around 0.04) [67]. The synchronous release

mechanism exhibits more exponential decay components in its release rate histogram than does the

asynchronous mechanism (4 versus 3), likely because it has more Ca^{2+} binding sites (5 versus 2) and

because it operates on a faster time scale. Fig 7 shows how the fitted parametric release profiles match

the simulated release profiles to within noise across multiple orders of magnitude; the noise in the

simulated profiles was due to fluctuations in $[Ca^{2+}]$ at the Ca^{2+} sensor. Table 2 lists the values of the best-

fit profile parameters.

Fig 7. Fitted Release Rate Histogram Profiles for a Single Spike.

Parameter values given in Table 2. A: Synchronous release rate: true histogram (blue) with estimated histogram (black). B: Asynchronous release rate: true histogram (red) with estimated histogram (black).

Table 2. Spike-Evoked Release Rate Parameters.

Parameter values calculated for a single spike following a period of low activity. Valid for Ca^{2+}-sensitive synchronous and asynchronous release mechanisms located 400 nm from a cluster of 100 VDCCs.

component	P	τ	k	μ	σ
S_1	0.0175	0.163 ms	1.79 ms^{-1}	3.41 ms	0.168 ms
S_2	0.0220	6.50 ms	18.0 ms^{-1}	3.56 ms	0.0977 ms
S_3	1.70×10^{-5}	80.0 ms	0.526 ms^{-1}	10.0 ms	4.44 ms
S_4	1.10×10^{-5}	1000 ms	0.142 ms^{-1}	50.0 ms	11.5 ms
A_1	3.72×10^{-3}	17.7 ms	1.60 ms^{-1}	3.05 ms	0.243 ms
A_2	0.0111	76.9 ms	0.0759 ms^{-1}	4.00 ms	1.14 ms
A_3	0.0136	1000 ms	0.0337 ms^{-1}	76.5 ms	21.9 ms

Keep in mind that the μ values are somewhat arbitrary in that they depend on exactly when during the

action potential that the spike time is taken to occur. Action potential waveforms last a couple of

400   milliseconds (see S2 Fig) [68]; the values for μ above used a point on the action potential waveform

401   immediately prior to the rising phase as the spike time. Using the peak of the action potential would

402   take away about 2 ms from all values of μ. Again, the time point along the action potential where the

403   spike is counted is arbitrary, but it must be consistent across all components.

Combining Release Profiles for Multiple Spikes

405   Having established the shape of the release profile for a single spike, we considered how the release

406   profiles of multiple spikes in a train would combine. Consider first the case of two spike times, t_{s1} and t_{s2}

407   (where $t_{s2} > t_{s1}$). After the first spike, the release mechanism will respond to a Ca^{2+} influx after some

408   delay with the release profile of Eq (1) shifted in time by t_{Ca1}, where the delay between the spike time

409   and the arrival of the Ca^{2+} influx is distributed according to the ex-Gaussian temporal delay filter:

$$t_{Ca1} - t_{s1} \sim a(t - t_{s1}; k, \mu, \sigma). \qquad (5)$$

Fig 8 A-C shows visually how this temporal delay filter affects a given release profile component.

Fig 8. Convolutional Filter Applied to a Component of a Release Rate Function.

Toy model with P=5, τ=10 ms, k=0.5 ms^{-1}, μ=5 ms, and σ=1 ms. A: Unfiltered release rate component. B:

MCMC ex-Gaussian filter shape. C: Filtered release profile produced by convolving the release rate

profile with the temporal delay filter. D-F: Release rates in response to spike trains without applying

delay filter. G-I: Release rates in response to spike trains with delay filter applied. D,G: Response to one

spike. E,H: Response to two spikes. F,I: Response to multiple spikes. Dotted lines show how the

histogram of response to one spike falls off with interference from the response to the following spike.

Spike times at 0, 15, 20, 30, and 50 ms.

When the second spike arrives at the release site, the VDCCs produce another influx of Ca^{2+} that can

again propagate to the SNARE complex, building on the Ca^{2+} from the first spike. The buffered diffusion

again involves an ex-Gaussian-distributed delay, after which the release mechanism starts responding to

the second spike at time t_{Ca2}. With a future event-driven simulator in mind, we treated the arrival of Ca^{2+}

from the second spike as a transition point between release-time generating functions. That is, the

synapse stops generating release times in response to the first spike (whose release profile was shifted

by t_{Ca1}) and starts generating release times in response to the second spike (with a release profile shifted

by t_{Ca2}), according to

$$r(t; \{t_{Ca1}, t_{Ca2}\}) = r_0 + \sum_{c=1}^{N} \frac{P_c}{\tau_c} e^{-(t-t_{Ca1})/\tau_c} u(t - t_{Ca1})\left(1 - u(t - t_{Ca2})\right)$$

$$+ \sum_{c=1}^{N} \frac{P_c}{\tau_c} e^{-(t-t_{Ca2})/\tau_c} u(t - t_{Ca2}) \qquad (6)$$

$$= r(t - t_{Ca1})\left(1 - u(t - t_{Ca2})\right) + r(t - t_{Ca2}),$$

429   where r(t) is the unfiltered release profile from Eq (1). Notice that Eq (6) does not account for facilitation

430   yet. In this section, we focus on the interaction of individual spikes' release profiles, reserving the

431   discussion of facilitation for the next section. For a spike train with an arbitrary number of spikes, this

432   becomes

$$r(t; \mathcal{T}_{Ca}) = \sum_{t_{Cai} \in \mathcal{T}_{Ca}} \left(r(t - t_{Cai}) \prod_{\substack{t_{Caj} \in \mathcal{T}_{Ca} \\ t_{Caj} > t_{Cai}}} \left(1 - u\left(t - t_{Caj}\right)\right) \right), \qquad (7)$$

where $\mathcal{T}_{Ca} = \{t_{Ca1}, t_{Ca2}, \dots\}$ is the set of all Ca^{2+} arrival times, each resulting from the combination of a

spike time with an ex-Gaussian-distributed delay. Fig 8 D-F shows what these profiles should look like for

a certain set of parameters in response to various spike trains.

While the above formulation relies on fixed delay times, an event-driven simulator will need to sample

new delay times on every trial for a given spike train, as well as different delay times for each release

component. Therefore, we calculated the probability $D_c(t)$ that the Ca^{2+} sensor has begun responding

to the Ca^{2+} from the latest spike for release component c, allowing for a gradual switch from one spike-

evoked release profile to the next, taking into account the variation of delay among all possible trials.

For two spike times, $t_{s1} < t_{s2}$, the net release profile for each component of Eq (6) becomes

$$r_c(t; \{t_{s1}, t_{s2}\}) = r_c(t - t_{s1})\big(1 - D_c(t - t_{s2})\big) + r_c(t - t_{s2}). \tag{8}$$

442   In other words, the response to the first spike is cut short after the second spike by $D_c(\cdot)$ to give way to

443   the new release response. And every time another spike arrives, it decreases the probability of release

444   relative to the first spike multiplicatively, such that each component of Eq (7) becomes

$$r_c(t; \mathcal{T}_S) = \sum_{t_{si} \in \mathcal{T}_S} \left(r_c(t - t_{si}) \prod_{\substack{t_{sj} \in \mathcal{T}_S \\ t_{sj} > t_{si}}} \big(1 - D_c(t - t_{sj})\big) \right), \tag{9}$$

445   where $\mathcal{T}_S = \{t_{s1}, t_{s2}, \dots\}$ is now the set of all spike times.

446   Because this formulation now uses spike times rather than delayed Ca^{2+} arrival times, the step functions

447   u(t–t_{Caj}) of Eq (6) and (7) have been smeared out in time by the temporal delay filter of Eq (4) to become

448    $D_c(t–t_{sj})$. Assuming the latest spike arrives at t=0, $D_c(t)$ is simply the cumulative distribution of the

449   temporal delay filter:

$$D_c(t) = \int_{-\infty}^{t} a(t; k_c, \mu_c, \sigma_c)\mathrm{d}t$$

$$= \Phi\left(\frac{t - \mu_c}{\sigma_c}\right) - \mathrm{e}^{-k_c\left(t - \left(\mu_c + \frac{\sigma_c^2}{2}k_c\right)\right)}\Phi\left(\frac{t - (\mu_c + \sigma_c^2 k_c)}{\sigma_c}\right). \tag{10}$$

450   More intuitively, by letting $\sigma_c \to 0$, the Gaussian component becomes a delta function, and the first-

451   release distribution function above becomes much more simply

$$D_c(t) = \big(1 - \mathrm{e}^{-k_c(t - \mu_c)}\big)\mathrm{u}(t - \mu_c). \tag{11}$$

452   Thus, after the second spike, the histogram of releases from the first spike drops off exponentially, while

453   those due to the second spike rise and fall as for the first spike. When σ_c is small relative to the median

454   interspike interval, third spikes have an almost imperceptible effect at cutting the first profile short

455   relative to the second spike's effect. Fig 8 G-I shows how this transition works in response to the same

456   spike trains as in Fig 8 D-F.

Characterizing Facilitation in Vesicle Release Rates

The discussion above has focused on the release response of a single vesicle with a constant probability

of release across spikes. However, many synapses display a facilitation in release probability from one

spike to another [20, 60, 69-73]. This results both from an accumulation of Ca^{2+} in the presynaptic space

[69] and from a stochastic accumulation of Ca^{2+} on the sensor (Syt) of the SNARE complex. Simulations

with the MCell model demonstrate how nonlinear binding cooperativity in the Ca^{2+} sensors induces

facilitation in excess of what would be expected from cytoplasmic Ca^{2+} buildup alone (S9 Fig). This

happens because on some trials, Ca^{2+} accumulates on the sensor, not enough to trigger vesicle fusion on

the first spike, but enough to increase the probability of reaching the releasable state after subsequent

spikes. As can be seen in S8 Fig, Ca^{2+} entry from one AP can predispose the distribution of bound states

of the sensor to trigger release with greater alacrity on subsequent spikes.

Furthermore, the level of facilitation depends to some extent on the full history of spiking activity in the

synapse. In the simplest case, the probability of release on one spike should depend only on the

probability for the previous spike and on the time since the previous spike. However, the level of

facilitation is not a simple function of the most recent activity but depends on the rate of stimulation

prior to the last spike. To explore the space of facilitation dynamics more fully, we applied spike trains

with spike ramps of different rates and durations, to see how quickly facilitation builds up, followed by

single probe spikes at increasing interspike intervals (ISI), to see how quickly it decays back to baseline

(see Methods). Fig 9 shows examples of how these different spike trains affect the rates of synchronous

and asynchronous release.

477
Fig 9. Empirical Facilitation in Synchronous and Asynchronous Release Rates.
Synchronous release rate shown in blue; asynchronous shown in red. Dark colors represent initial spike
ramp (common to all traces on a plot); light colors represent single probe spikes from different
simulations. Associated Ca^{2+} traces omitted for clarity. A: PPF decays with increasing ISI (probe ISIs of 2,
5, 10, 20, 50, 100, 200 ms). B: 5-spike ramp with a 5-ms ISI shows strong facilitation. C: 5-spike ramp
with a 20-ms ISI shows weaker facilitation. Note the change in scale from A to B and C.
484

485   Facilitation does not affect all components of release equally. Therefore, we derived a general

486   facilitation function $F_c(\cdot)$ that affects each release component c independently. The area under the

487   curve of each component of the release rate profile (see Eq (1) and (3)) depends on the facilitation

488   factor according to

$$P_c(n) = P_{c0} \cdot F_c(n),\tag{12}$$

489   where P_{c0} is the baseline value and n is the index of the current spike. To ensure that the function works

490   for arbitrary spike trains, the factor $F_c(n)$ needs both to grow somehow from spike to spike and to

491   decay back toward one for large ISIs. This growth can happen in a highly nonlinear fashion, so to account

492   for this, we take $F_c(n)$ to be a nonlinear combination of linear facilitation factors $f_{ci}(n)$ such that

$$F_c(n) = \prod_{i=1}^{M_c} f_{ci}(n)^{\xi_{ci}},\tag{13}$$

where M_c represents the number of facilitation components (either one or two for all functions

explored below), and ξ_{ci} represents the nonlinearity applied to facilitation component i of release

component c. Each $f_{ci}(n)$ accounts for some aspect of the internal state of the SNARE complex, in terms

of how the expected number of Ca^{2+} ions bound changes with time, that helps determine the probability

of release on subsequent spikes.

In the simplest case, each $f_{ci}(n)$ would decay exponentially from its previous value $f_{ci}(n-1)$ before

incrementing by one:

$$f_{ci}(n) = f_{ci}(n-1)e^{-\Delta t/\tau_{ci}} + 1,\tag{14}$$

where Δt is the delay from the previous spike to the current one. The increment of one is meant to

account for the influx of about the same amount of Ca^{2+} during each action potential. This formulation

ensures that even after infinitely long intervals, the facilitation factor will equal a value no less than

$F_c(0) = 1$, allowing the release components to return to their baseline values of $P_c = P_{c0}$ during long

periods of inactivity, as expected.

However, this formula implies that for an infinitely fast rate of stimulation, $f_{ci}(n)$ could grow toward

infinity, producing an infinitely fast rate of release, all of which are impossible. More realistically, there

should exist some finite saturation level, L_{ci}, such that the facilitation function could never theoretically

exceed

$$F_c(\infty) = \prod_{i=1}^{M_c} L_{ci}. \tag{15}$$

The value of this upper limit is constrained by the rates of vesicle fusion from the fully bound states of

the SNARE complex (γ_S and γ_A in Table 1) and by the maximum level of Ca^{2+} buildup in the presynaptic

space. When facilitation is still well below this level, it should continue to increment by approximately

one on every spike, but this increment should fall to zero quickly enough that facilitation never exceeds

saturation. Setting a maximum number of equal-sized steps to saturation for each component,

$N_{ci} = L_{ci}^{1/\xi_{ci}}$, the value of $f_{ci}(n)$ becomes

$$f_{ci}(n) = f_{ci}(n-1)e^{-\Delta t/\tau_{ci}} + 1 - \left(\frac{f_{ci}(n-1)e^{-\Delta t/\tau_{ci}}}{N_{ci}}\right)^{N_{ci}}. \tag{16}$$

The new term subtracted off at the end ensures that $f_{ci}(n)$ never exceeds N_{ci}, just as Ca^{2+} cannot

accumulate to infinite concentrations but is limited by the electrochemical gradient across the cell

membrane [74]. An alternative would be simply to set $f_{ci}(n) = N_{ci}$ whenever a step size of one would

cause it to exceed this limit, but the formula in Eq (16) allows for a smoother approach.

With the model for facilitation established, we sought to fit it to the empirical changes observed in

release rate for complex spike trains. For simplicity, we took facilitation to apply only to the P_c

parameters, which control for the magnitude of each release component, although in principle the

parameters of the temporal filter (k_c, μ_c, and σ_c) might also increase ($\xi_{ci} > 0$) or decrease ($\xi_{ci} < 0$)

with spike history. As discussed in Methods, we explored 136 unique spike trains for how both spike rate

(along the spike ramp) and ISI (of the probe spike) affect the release rate in response to the last spike.

For the fitting algorithm, we used a simplex method for gradient descent, since the derivatives of the

error function are difficult to compute (see Methods). The values of the P_c parameters were allowed to

vary within bounds, while the profile time constants and the temporal filter parameters were held

constant. The best-fit set of values for P_c were found for the spike-response profile at the end of each

spike train, after which the meta-parameters of the facilitation functions could be fitted to the patterns

in P_c. Fig 10 A shows some examples of fitted release rate functions fit to baseline (blue) and facilitated

(yellow) profiles. Fig 10 B depicts how release magnitudes varied for the release magnitude components

(first (P_{S1}) and third (P_{S3}) components of synchronous release and first component (P_{A1}) of

asynchronous release) across all 136 facilitation spike trains (see S10 Fig for a depiction of how stimulus

cases are ordered).

Fig 10. Release Rate Parameters and Facilitation Metaparameters Fitted to Empirical Histogram Profiles.

A: Synchronous and asynchronous profiles fitted for baseline (un-facilitated) case, and for highly facilitated case (probe spike 5 ms after 5-spike ramp of 5-ms ISIs). B, C: Release fidelity values fitted case-by-case (dark colors) overlaid with predictions from best-fit facilitation functions (light colors) for synchronous (B) and asynchronous (C) components.

Through trial-and-error, we found that the two components of synchronous release with the fastest

time constants, along with the fastest component of asynchronous release, each required two

facilitation components to explain their patterns of change from case to case. The synchronous and

asynchronous release components with "medium" time constants (each close to 80 ms) could each be

fitted with a single facilitation component. The slowest release components, with time constants of

1000 ms due to latent $[Ca^{2+}]_i$ released from the buffer (see Fig 5 C), were constrained not to facilitate,

since changes in these components seemed to have a negligible effect on fitting error. Table 3 records

the facilitation meta-parameters obtained from the fits, along with the baseline values for release

fidelity for each component (P_0). During each step of the fitting algorithm, these metaparameters were

used to generate predictions for the state of facilitation across all 136 spike-train cases, and error was

calculated as the fraction of the variance of the "true" release fidelity values unexplained by the

predicted pattern (S12 Fig; see Methods).

Table 3. Metaparameters for Facilitation of Release Fidelity.

First column shows baseline magnitudes of integrated release rate, duplicated from Table 2. First and second components of synchronous release and first component of asynchronous release facilitate with two time constants each. Smallest component of both release mechanisms does not facilitate. P_0 is integrated release rate for the un-facilitated case (baseline), τ is the time constant of decay for each facilitation component, N is the number of linear facilitation steps to saturation, ξ is the nonlinearity parameter, and L is the maximum facilitation factor contributed by each component. Valid for Ca^{2+}-sensitive synchronous and asynchronous release mechanisms located 400 nm from a cluster of 100 VDCCs.

component	P_0	τ	N	ξ	$L = N^\xi$
S_{11}	0.0175	95.9 ms	7.00	1.27	11.8
S_{12}		7.66 ms	2.32	2.93	11.8
S_{21}	0.0220	13.1 ms	10.0	1.23	17.0
S_{22}		114 ms	17.6	1.68	125
S_3	1.70×10^{-5}	199 ms	12.5	2.67	846
S_4	1.10×10^{-5}	–	1	0	1
A_{11}	3.72×10^{-3}	141 ms	12.2	1.48	40.0
A_{12}		17.2 ms	12.5	0.996	12.4
A_2	0.0111	126 ms	12.1	1.67	64.4
A_3	0.0136	–	1	0	1

Discussion

Advantages and Limiting Assumptions of Treating MCell as Ground Truth

Basing the new model on MCell has distinct advantages over biological experiments in terms of both flexibility and precision when it comes to model validation. As an example, estimates of $[Ca^{2+}]_i$ in neurons obtained from fluorescent reporters in physiological experiments may provide accurate

estimates of slow (tens of milliseconds) Ca^{2+} transients [75, 76], but the buffering kinetics of the calcium

reporters can act as a low-pass filter, obscuring the faster (0-5 milliseconds) components of Ca^{2+}

dynamics [37]. Molecular simulations like MCell, on the other hand, can capture these fast transients,

since they track every particle, which may be crucial for correctly modeling fast, Ca^{2+}-dependent

synaptic processes like synchronous vesicular release of neurotransmitter [65] (see Figure 6 from [37]).

Furthermore, neurotransmitter release may occur asynchronously with respect to the arrival time of

action potentials, following some time-dependent distribution [65]. Experimental methods for

determining release rate would offer far less control of presynaptic conditions over the number of trials

that would be required to tease out the same resolution of detail as is possible with controlled

simulations. Therefore, we chose to constrain ourselves to validating the model developed in this paper

against an MCell model, which has itself been validated already against hippocampal Schaffer collateral

synapses [35].

However, this approach is limited on several levels. First, it assumes that MCell can correctly replicate

the dynamics of diffusion and molecular interactions of biophysical systems through its Markov chain

Monte Carlo framework [33, 34, 36]. Second, it assumes that the molecular kinetics of the included

species match their true kinetics. Finally, it assumes that the molecular species and biological systems

modelled are the only ones present in the presynaptic compartment, or at least that any other systems

would produce only negligible changes to the phenomenology of the synapse. MCell has been

sufficiently well validated to satisfy the first assumption [36], at least for the temporal and spatial scales

of interest here (hundreds of microseconds and several microns).

The second assumption is valid insofar as the molecular models used by MCell correctly represent

reality, both in terms of the molecular state diagrams and in terms of the binding and interaction

kinetics reported by other groups: Sun et al. [53] for the descriptions of Ca^{2+}-driven SNARE kinetics for

vesicular release; Bischofberger et al. [68] for the VDCC dynamics for spike-evoked Ca^{2+} influx; Nägerl et

al. [77] for the high- and medium-affinity sites of the calbindin Ca^{2+} buffer; and Sneyd et al. [78] for the

kinetics of the PMCA pumps. The state diagrams and kinetic parameters for these species are

summarized in S1 Fig and S1 Table. Simplifying assumptions inevitably go into models such as these,

which limit the accuracy of any model based on them. However, for the purposes of this paper, we

assume that these models reproduce experimental results sufficiently well to use them.

The greatest limitations to model accuracy come from the third assumption in that the true variety of

systems and molecular species in biological synapses far exceeds what MCell represents [79]. For

instance, the Nadkarni et al. [35] model did not include any endoplasmic reticulum (ER), which stores

intracellular Ca^{2+} and has a significant effect on neuronal signaling [80]. Ryanodine receptors (RyR) [81-

83] and inositol 1,4,5-trisphosphate receptors (IP3R) [84, 85] can unleash these Ca^{2+} stores. Inclusion of

such an ER would likely alter the shape of the Ca^{2+} transient and increase the probability of

neurotransmitter release, possibly over longer time scales, as in synaptic augmentation [67, 86].

Another system that would significantly affect Ca^{2+} dynamics over multiple spikes is the inactivation and

facilitation of Ca^{2+} channels mediated by Ca^{2+}-calmodulin (CaM) and Ca^{2+} binding proteins (CaBP1) [62,

87-90]. These interactions might help control release-independent depression (RID) and the frequency-

dependent recovery (FDR) from depression [91] by restricting Ca^{2+} influx over extended spike trains.

Furthermore, the presynaptic Ca^{2+} buffer includes more than just calbindin [92, 93] and diffusion is

limited by the plethora of intracellular microstructures [94, 95]. Because location and movement

through space plays a crucial role in phenomenology (see Fig 2, S4 Fig, S6 Fig, and S7 Fig), investigating

how these features affect release dynamics would require rerunning the MCell model with them

included.

The advantages of using MCell as ground truth, we believe, outweigh the limitations enumerated above.

A simulated synaptic model allows for much finer experimental control and consistency from trial to

trial, while yielding far more precise results than physiological experiments. Measuring release rate,

probability, facilitation, and depression at biological synapses is difficult and requires a number of

problematic assumptions [96]. Using MCell allows for precise measurements of unmodified Ca^{2+} traces

[37] and of single-vesicle release rates, even controlling for such confounding processes as the post-

release refractory period [35]. Any new features, including arbitrary numbers of vesicle pools and

synaptic processes, can easily be included in future work.

Comparison to Other Models of Short-Term Synaptic Dynamics

In response to a given spike train, this model produces a release rate profile that represents the rate of a

time-varying Poisson process, the average release activity of an infinite number of synapses (or trials)

responding to the same spike train. The multi-exponential form of the release rate profile and the

facilitation factors that modify it evolve deterministically for a given sequence of spike times.

Incorporating this as a deterministic synaptic model into a spiking neural network simulation would

allow for comparison with other deterministic models of short-term plasticity, such as that introduced

by Tsodyks and Markram [97]. Their model uses a relatively simple representation of the utilization of

synaptic resources, tracking the fraction of resources in recovered, active, and inactivated states, to

flexibly model both short-term depression and facilitation [20], and has proven useful in simulated

neural network contexts for producing complex behavior [98]. On the other hand, the predictive power

of the Tsodyks-Markram model is limited by how it abstracts away all the internal processes of the

synapse, conflating presynaptic (vesicle availability, release probability) with postsynaptic

(neurotransmitter receptor saturation and desensitization) resources. In contrast, our model more

closely tracks the vesicle release phenomenology that results directly from Ca^{2+}-evoked molecular

dynamics. This allows us to explore the contribution of more fine-grained synaptic features to whole-

network computations.

Another limitation of the Tsodyks-Markram model is its deterministic formulation, representing average

or aggregate synaptic behavior rather than single-trial or single-synapse behavior. However, the

presence of trial-to-trial stochasticity in synaptic transmission may have important implications for the

learning and information processing performed in neural circuits [27, 99-103]. Our model can capture

such trial-to-trial and synapse-to-synapse variability if we sample release events from the time-varying

Poisson process defined by the release rate profile.

An example of a presynaptic model that captures probabilistic release is that by Maass and Zador [104].

The Maass-Zador model tracks short-term facilitation in response to spike history and short-term

depression in response to release history, calculating the probability of release at each spike time and

generating Boolean release events according to this probability. However, these release events only

occur exactly at spike times, in contrast to the spontaneous and asynchronous release that occurs in real

synapses [105, 106]. Furthermore, the phenomenology of the Maass-Zador model, while elegant, arises

from mathematical abstractions rather than from physiologically grounded mechanisms, making it

susceptible to producing unrealistic behavior and limiting its utility as a testable model.

Kandaswamy et al. [67] present another presynaptic model that aims for more physiologically grounded

realism, employing multiple mechanisms of vesicle recycling, facilitation, augmentation, and release-

dependent depression, each with its own set of parameters. Some parameters were constrained to

values derived from earlier studies, while other parameters were fit to experimental measurements of

changes in synaptic strength in response to various constant-frequency spike train stimuli. Although the

resulting model does qualitatively well at matching the experimental data, it lacks generalizability. In

particular, the free parameters of their model, which they adjusted to fit the experimental data, depend

empirically on the frequency of stimulation but without any discernible pattern that would provide

insight into their physiologic origin (see Table 2 in [67]). Furthermore, the model for facilitation treats

the first spike as a special case relative to all subsequent spikes, excluding the first spike from facilitation

to fit the data. A realistic short-term facilitation model should scale the probability of release on the first

spike as though it occurred in the middle of a train after an infinite ISI, since the synapse should return

to its baseline state after a sufficiently long interval of no spiking activity, which our model

accomplishes.

Future work with our model will involve implementing it in an event-driven framework, where release

events are sampled from the time-varying Poisson process defined by the release rate profile. This

approach will provide a powerful way to include a highly generalizable facilitation function and sub-

millisecond vesicular release phenomenology, grounded in molecular kinetics, into a highly scalable and

computationally efficient presynaptic model. In contrast to the models reviewed above, this model can

achieve both the stochasticity and the asynchronicity that are characteristic of real synapses, while

maintaining a clear mapping between its parameters and the underlying molecular mechanisms. This

approach will be crucial for exploring the impact of different presynaptic mechanisms on the

computational performance of large neural circuits.

The main advantage of our model is its balance of computational efficiency with physiologically

grounded realism. It accounts for both asynchronous and spontaneous release events, which the Maass-

Zador model lacks. For neural network simulations where the presence of stochasticity in synaptic

transmission is more important than reproducing true dynamics, the Maass-Zador model may suffice.

However, for investigations into how presynaptic mechanisms of vesicle release affect information

transmission and network behavior, our model provides and indispensable layer of flexibility.

Furthermore, our model provides a highly flexible and explanatory framework. Each of the parameters

for describing the profile of the release histogram (P, τ, k, μ, σ; see Eq (1), (3), (4)) has an almost direct

link to the underlying physiology, whether to the Ca^{2+}-binding and vesicle fusion kinetics of the SNARE

complex (P and τ) or to the stochastic delay in response to the spike caused by buffered diffusion of Ca^{2+}

(k, μ, and σ). Furthermore, the facilitation function has sufficient complexity to account for the changes

seen in neurotransmitter release fidelity of a wide variety of spike train patterns. Importantly, all spikes

are treated equally: Eq (13) and (16) apply as consistently to the first spike as to the n-th. While it falls

short in terms of computational efficiency relative to the Kandaswamy et al. model, it makes up for it in

terms of biophysical plausibility and its utility for testing hypotheses regarding synaptic function.

Importance of Facilitation Function Parameterization

Different synapse types in different regions of the brain employ different short-term plasticity functions,

including various forms of facilitation and depression, transforming spike timing codes into

neurotransmitter release timing codes [97]. We do not know what the precise computational roles of

facilitation and depression play in neural circuits, but it seems likely to involve more than just high-pass

or low-pass filtering of spike trains [19, 26, 27, 107-109]. The model of facilitation presented in this

paper enables the exploration of this question both because of its grounding in explainable molecular

physiology via MCell and because of its amenability to efficient simulation of complex neural circuits.

One possible shortcoming of the facilitation function presented in this paper is that we did not explore

the steady-state behavior in our simulations. Although the steady-state facilitation factors can be

calculated from the fitted parameters, we will need to run MCell simulations with longer spike ramps to

confirm their values for long spike trains of different frequencies. Since the release rates explored in this

paper deal with single-vesicle release probabilities, release-dependent depression (i.e. depletion-

dependent depression) should occur naturally when applying the model to simulations of synapses with

finite readily releasable vesicle pools. However, other forms of release-independent depression may be

more relevant to natural spike frequencies [91, 110] and could be incorporated into an extra facilitation

factor within the current model by using a negative ξ as the exponent in Eq (13). Thus, the facilitation

function can be naturally extended to include release-independent depression mechanisms, although

further research is necessary to confirm the kinetics of the other active zone molecules of non-

hippocampal synapses that underlie such mechanisms.

Future Refinements

Because so much about release probability and timing depends on the precise magnitude and time

course of the Ca^{2+} transient, accurate modelling of Ca^{2+} dynamics using MCell is essential to future

refinements of our model. For instance, intracellular calcium stores, including ER and its associated

receptors and channels [80-86, 111] and mitochondria and its associated calcium uniporter [112], can

impact the Ca^{2+} signaling over the long term. The diffusion of Ca^{2+} through the presynaptic space can be

affected in ways not captured by our current model by buffering molecules other than calbindin and

signaling molecules such as CaM and CaBP1 [62, 87-90, 92, 93] and by the geometry of cytoskeletal

microstructures [94, 95]. Additional mechanisms that can influence the Ca^{2+} transient include the shape

of the presynaptic action potential, which can vary in cases such as Fragile X syndrome [113], spatially

localized Ca^{2+} spikes mediated by presynaptic NMDA receptors [114-116], and retrograde signaling via

endocannabinoid receptors [117-119].

Other considerations of synaptic physiology, those which do not affect the shape of the spike-evoked

Ca^{2+} transient, can be characterized without running full MCell simulations. These include the structure

and dynamics of the SNARE complex. In our model, we assumed that each vesicle employs two Ca^{2+}

sensors for triggering vesicle fusion, Syt-1/2 for synchronous release and Syt-7 for asynchronous release,

as characterized by the Sun et al. [53] model. Each mechanism acted independently, and together they

served as the sole mechanism of release. However, many more molecules comprise the SNARE complex,

each affecting release fidelity in complex ways [14, 15, 44, 48-52]. In fact, the inclusion of extra

molecules in the SNARE assembly may play a crucial "superpriming" step in enhancing the release

alacrity of already-primed vesicles [44, 69]. Furthermore, although a single SNARE complex is sufficient

to induce spike-evoked release [120], each vesicle may have multiple SNARE complexes associated with

it, which is necessary for fast vesicle fusion [121]. We predict that multiplying SNARE complexes would

simply multiply release rate in proportion. It is less clear what effects that other molecules, such as

complexins and Muncs, would have on release kinetics. A quantitative understanding of their molecular

interaction kinetics is required before they can be applied to this model. Once obtained, however, we

can apply deterministic simulations of state probabilities (see Methods) for this more complex SNARE

model, similar to what we did in this paper. If these simulations are driven by the improved Ca^{2+}

transients obtained from the more mechanistically exhaustive MCell simulations described above, we

can derive much more biologically accurate phenomenology.

Although our work focused on characterizing hippocampal Schaffer collateral synapses, our approach

can apply just as well to other synapse types. Additionally, other internal synaptic processes such as

vesicle recycling may be combined with our model within an event-driven framework. In this way, future

refinements of our model will have the ability to capture physiologically realistic phenomenology, trial-

to-trial variability, asynchronicity, and internal synaptic dynamics for a wide variety of synapses at very

low computational cost. It can help to uncover the contributions of single synapses or synaptic features

to network computations, establishing a connection from molecular kinetics through synaptic

phenomenology up to whole-network dynamics. This can be applied to create predictive models of

biological networks from known physiological parameters or to design neuromorphic chips or spiking

neural networks with dynamical properties relevant to future developments in brain-computer

interfaces and artificial intelligence.

754   Methods

755 Ca^{2+}-Evoked Vesicle Release Model

The detailed model of molecular reaction-diffusion dynamics was developed in the modeling

environment known as MCell [33, 34, 36]. The MCell model used as a basis for the design and validation

of the presynaptic model presented in this paper comes from Nadkarni et al. [35]. It includes

mechanisms for voltage-sensitive Ca^{2+} influx and for Ca^{2+} buffering in the presynaptic space, with pumps

and channels in the membranes to maintain a steady-state average free Ca^{2+} concentration of 100 nM

[74].

When an action potential arrives at the presynaptic membrane, voltage-dependent Ca^{2+} channels

(VDCCs) open stochastically, traversing through four unopened states via voltage-dependent state

transition rates [68] (see S1 Fig A), producing a Ca^{2+} influx due to the steep electrochemical gradient

[74]. The VDCCs very quickly shut off after the membrane potential returns to baseline (S2 Fig), and the

newly introduced Ca^{2+} ions diffuse randomly in the presynaptic space. Vesicle fusion occurs when a

sufficient number of Ca^{2+} ions have diffused over and bound to the release machinery associated with

the SNARE complex of a docked vesicle [15, 35, 53]. Cytoplasmic calbindin (CB) with a concentration of

45 μM acts as a buffer that modulates the magnitude and duration of the free Ca^{2+} (S1 Fig B) [77], and

plasma membrane Ca^{2+}-ATPase (PMCA) pumps (S1 Fig C) actively remove Ca^{2+} ions over a time course of

seconds [78] to the baseline $[Ca^{2+}]_i$ of 100 nM [35, 74]. Parameter values for these molecular

mechanisms are given in S1 Table.

Calcium concentration as a function of time is measured at various spatial locations within the

presynaptic space (see Fig 2 for synaptic structure and "Estimating $[Ca^{2+}]_i$ from Collision Events" in the

Supplemental Information for a description of how local $[Ca^{2+}]$ is measured). For each spike train, local

$[Ca^{2+}]_i$ in the vicinity of the SNARE complex was averaged over 2000 trials at a resolution of 0.1 msec.

These calcium transient profiles were then used to drive deterministic simulations of SNARE state

probabilities, as described below.

Deterministic Simulations of State Probabilities

With the Ca^{2+} profiles obtained for each spike-evoked simulation in MCell and with the state transitions

for the release mechanisms defined in Fig 3 A-B and Table 1, the corresponding vesicle release-rate

profiles become computable. While the MCell simulations we ran do generate their own sets of release

times, it would be impossible to fit precise phenomenological functions to the release histograms

without running an infeasibly large number of trials. Furthermore, vesicle depletion following release

events confounds the representation of release rate, its functional form, and its facilitation dynamics

(see Fig 4). Therefore, instead of running millions of trials of MCell (or more) to produce temporally

precise single-vesicle release rate histograms, we used the averaged calcium profiles from 2000 trials (at

a resolution of 0.1 msec) to drive a deterministic simulation of the SNARE complex, tracking the

probabilities of being in each state as functions of time. This approach, in effect, produced the average

release histograms equivalent to an infinite number of trials acting on the averaged calcium traces.

This method tracked the probabilities of a particular release mechanism being in each possible state at

every time step. That is, each state represents the number of Ca^{2+} ions bound to the release molecule (0

through 5 for Syt-1 (synchronous; S) and 0 through 2 for Syt-7 (asynchronous; A)). Each mechanism

$X \in \{S, A\}$ has a state probability vector $\mathbf{s}^X(t)$ associated with it that tracks the probability of being in

each possible state, that is, the probability of having n Ca^{2+} ions currently bound for $n \in \{0 \dots N_X\}$,

where $N_S = 5$ and $N_A = 2$. State probabilities add to unity, and they update on each time step

according to a [Ca^{2+}]-dependent state transition rate matrix $\mathbf{T}^X$, whose superdiagonal terms are the

unbinding rates, moving from a higher to a lower-bound state, and whose subdiagonal terms are the

binding rates, moving from a lower to a higher-bound state. Specifically, for mechanism $X \in \{S, A\}$ with

$N_X \in \{5,2\}$ calcium ions needed for release to occur, the binding rate is

$$T^X_{n+1,n} = (N_X - n) \cdot k_{X+} \cdot [\text{Ca}^{2+}], \tag{17}$$

801   and the unbinding rate is

$$T^X_{n-1,n} = n \cdot b^{n-1} \cdot k_{X-} \tag{18}$$

for $n \in \{0 \dots N_X\}$ ions currently bound, where $b = 0.25$ acts as a binding cooperativity factor (see Fig 3

and S8 Fig for state transition diagrams and Table 1 for parameter values). Notice that we index the

rows and columns of $\mathbf{T}^X$ (as well as the dimensions of $\mathbf{s}^X$) starting from 0 rather than 1 for convenience

in representing the number of calcium ions bound in the current state (column) and in the next state

(row). To enforce conservation of mass, the diagonal terms must equal the combined rate of leaving the

current state through both binding and unbinding:

$$
\begin{aligned}
T^X_{n,n} &= -\left(T^X_{n+1,n} + T^X_{n-1,n}\right) \\
&= -\left((N_X - n) \cdot k_{X+} \cdot [\text{Ca}^{2+}] + n \cdot b^{n-1} \cdot k_{X-}\right).
\end{aligned}
\tag{19}
$$

809   The above formulation does not yet take into account vesicle fusion. Recall that each mechanism X

810   induces vesicle release at a certain rate γ_X from its releasable state (all Ca^{2+} ions bound). Therefore, in

811   addition to the unbinding rate, the diagonal term for the fully bound state of each mechanism also

812   includes the rate of transition to the release state:

$$T^X_{N_X,N_X} = -(\gamma_X + N_X \cdot b^{N_X-1} \cdot k_{X-}), \tag{20}$$

where γ_X is the mechanism-specific release rate defined in Table 1. By including the release rate, the

sum of state probabilities would slowly decay towards zero as probability mass leaks into the

(untracked) release state, since when release occurs, the vesicle can no longer participate in further

activity. This depletion of vesicle probability would obscure the true shape of the single-vesicle release

rate. To account for this, the state vector is renormalized at each time step by the probability of no

release event having yet occurred, such that the occupancies of each state again add to one:

$$\mathbf{s}^X(t) \leftarrow \frac{\mathbf{s}^X(t)}{\sum_{n=0}^{N_X} s^X_n(t)}. \tag{21}$$

If one considers the deterministic simulation to represent a state histogram averaged over an infinite

number of trials, this normalization step effectively "zooms in" on the fraction of trials at each time step

for which no release occurred. Thus, the model tracks the instantaneous rate of release, given that no

release has yet occurred since the start of the simulation. This permits the calculation, for example, of

the equilibrium state probability distribution (see S8 Fig, left pie charts), driven by the steady-state Ca^{2+}

concentration (100 nM in MCell: [35]). These equilibrium state vectors are essential for initializing all

other simulations. From these, it is possible to determine the steady-state spontaneous release rates for

each mechanism that result ($S_0 = 5.70 \times 10^{-9} ms^{-1}$; $A_0 = 1.84 \times 10^{-5} ms^{-1}$; see S3 Fig).

Because the matrix $\mathbf{T}^X$ represents transition rates rather than transition probabilities, it acts as the

infinitesimal generator for a continuous-time finite state Markov process [122] rather than as a discrete

transition matrix. Converting this exactly to a discrete transition probability matrix $\mathbf{P}^X$ using a time step

of Δt requires an infinite sum of matrix products, according to the Taylor series

$$\mathbf{P}^X([Ca^{2+}];\Delta t) = \exp(\Delta t \cdot \boldsymbol{T}^X([Ca^{2+}])) = \sum_{m=0}^{\infty} \frac{(\Delta t \cdot \mathbf{T}^X([Ca^{2+}]))^m}{m!}. \tag{22}$$

This is akin to the probability of a Poisson process with transition rate λ remaining in the same state for

a time duration Δt, which follows the exponential form $p(t > \Delta t) = \exp(-\Delta t \cdot \lambda)$, acting just like the

diagonal terms of the transition rate matrix and sharing the same Taylor series expansion. The time step

Δt needs to be small enough such that the level of [Ca^{2+}] can be regarded as constant (on the order of

0.1 msec for the calcium transients investigated for this paper). However, in order to avoid too many

matrix multiplications and sums, we chose a time step of 0.005 msec, which is small enough to use the

linear approximation to Eq (22) while still maintaining numerical stability, even at very high levels of

[Ca^{2+}]:

$$\mathbf{P}^X([Ca^{2+}];\Delta t) \approx \mathbf{I} + \Delta t \cdot \boldsymbol{T}^X([Ca^{2+}]). \tag{23}$$

839   Thus, the state vector at time $t + \Delta t$ is the product of the state transition probability matrix with the

840   state vector at time t, according to

$$\begin{aligned} \mathbf{s}^X(t + \Delta t) &= \mathbf{P}^X([Ca^{2+}](t);\Delta t)\mathbf{s}^X(t) \\ &= \left(\mathbf{I} + \Delta t \cdot \boldsymbol{T}^X([Ca^{2+}](t))\right)\mathbf{s}^X(t), \end{aligned} \tag{24}$$

841   followed by the renormalization described above in Eq (21).

842   Similarly, the transitions in state probabilities for VDCCs, calbindin, and PMCA pumps in the well-mixed

843   model were calculated according to

$$\mathbf{s}^M(t + \Delta t) = (\mathbf{I} + \Delta t \cdot \mathbf{T}^M)\mathbf{s}^M(t), \tag{25}$$

where $M \in \{VDCC, Cb, PMCA\}$ refers to the molecular species. Furthermore, the transition rate matrix

$\mathbf{T}^{VDCC}$ is a function of membrane potential, while the matrices $\mathbf{T}^{Cb}$ and $\mathbf{T}^{PMCA}$ are functions of [Ca^{2+}]$_i$

(see S1 Fig and S1 Table for state transition diagrams and transition rate parameters). Within the well-

mixed model simulations, diffusion occurs instantaneously, effectively eliminating space from

consideration and allowing $[Ca^{2+}]_i$ to be the same for all molecular mechanisms within the synapse.

Stimulus Protocols for Exploring Facilitation

Whereas simulations with single action potentials can elucidate the functional form of synchronous and

asynchronous release, stimulus trains of multiple spikes can reveal the dynamics of facilitation in release

probability, which is well documented experimentally [20, 60, 69-73]. Short-term facilitation in release

probability is more pronounced for spikes closer together in time than for those separated by long

intervals. To investigate how delay affects probability of release, we studied paired-pulse facilitation

(PPF) for interspike intervals (ISIs) of exponentially increasing delay. Specifically, we stimulated the

MCell model with paired pulses of action-potential-like waveforms separated by 2, 5, 10, 20, 50, 100,

and 200 ms and measured the local $[Ca^{2+}]_i$ at a point within the axon (see Fig 2 for synaptic structure and

"Estimating $[Ca^{2+}]_i$ from Collision Events" in the Supplemental Information for a description of how local

$[Ca^{2+}]$ is measured). These Ca^{2+} traces then drove deterministic simulations of synchronous and

asynchronous release rate, as described in the previous section. This permitted us to determine a

functional form to describe PPF (see Results).

Realistic spike trains, however, involve patterns much more complex than paired pulses, and the recent

history of presynaptic activity can have a strong effect on future changes in release probability. To see

how facilitation evolves in more complex trains of action potentials, we designed a protocol to explore a

larger space of possible facilitated states, assuming that the level of facilitation experienced on one

spike depends exclusively on the delay since the previous spike (the interspike interval, or ISI) and on the

state of some internal facilitation parameter from the previous spike. The spike trains generally consist

of two phases: a spiking ramp and a probe spike. The ramp phase explores how facilitation develops

with multiple spikes at fixed ISIs and having anywhere from one to five spikes with an ISI of 2, 5, 10, or

20 ms (time prevented the exploration of ramps with more spikes). The probe phase explores how

facilitation wears off with increasing delay between spikes and consists of a single spike at 2, 5, 10, 20,

50, 100, or 200 ms after the end of the ramp, as in the PPF protocol above. All these combinations of

ramps and probes add up to $5 \times 4 \times (7 + 1) = 160$ cases (including those cases without a probe spike)

or 136 unique spike trains (discounting the repeats with one spike in the ramp at different ISIs). Future

work can explore the steady-state behavior of the facilitation factors by testing longer spike ramps of

each ISI.

Algorithms for Fitting Parameters and Metaparameters

Fitting parameter values to the shapes of the release-rate histograms involved two steps: first, obtaining

an initial guess, and second, optimizing the parameter values to a best-fit set. For the first step, the time

constants for rate decay (see Eq (1), (3), and Table 2) were found from the slopes of the logarithms of

the profiles (see Eq (2), Fig 5, and Fig 6) in response to both Ca^{2+} impulse and the Ca^{2+} traces derived

from MCell. Other parameters were initialized through trial and error. For the second step, we applied

the Nelder-Mead simplex method of function optimization using the fminsearch() function in MATLAB

2016 [123-125] to minimize the cost function over the parameters. This method does not require a

measure of the gradient of the cost function, which was not computable analytically. The cost function

uses the fraction of the variance unexplained (FVU) by the model, as in

$$\text{FVU}\big(y(t), f(t)\big) = \frac{\sum_{n=1}^{N_t}\big(y(t_n) - f(t_n)\big)^2}{\sum_{n=1}^{N_t}(y(t_n) - \bar{y})^2}, \tag{26}$$

887   where $\bar{y}$ is the mean release rate, $y(t)$ is the true shape of the release rate profile, and $f(t)$ is the

888   model profile at the same N_t time points. More precisely, the cost function ϵ is a linear combination of

889   the FVU for the function and for the logarithm of the function:

$$\epsilon\big(y(t), f(t)\big) = \alpha \cdot \mathrm{FVU}\big(y(t), f(t)\big) + \beta \cdot \mathrm{FVU}\big(\log\big(y(t)\big), \log\big(f(t)\big)\big), \qquad (27)$$

where α and β are constants ($\alpha = \beta = 1$ in the fits run for this paper). The FVU for the functions in

linear space is more sensitive to the high-amplitude peaks that occur for the fast release components,

while the FVU for the functions in logarithmic space is more sensitive to the slopes (and therefore the

time constants) of the exponential components.

The metaparameters of the facilitation functions (see Eq (13), (16), and Table 3) were fitted after the

parameters were fitted to the release profiles in response to each spike of the trains described in the

previous section of Methods. As described in Results, we took facilitation to apply only to the P_c

parameters, allowing them to vary within bounds for each spike in the train, while the profile time

constants and the temporal filter parameters were held constant. The best-fit set of values for P_c were

found for the final spike-response profile of each spike train. The fitted parameters were taken as true,

and the space of the logarithms of the metaparameters τ, N, and $L = N^\xi$ was explored, using the same

error function and optimization as above.

Acknowledgements

903   We would like to thank Donald Spencer and Margot Wagner for all the perspectives, insights, and

904   suggestions they offered as we prepared this manuscript for publication. MCell development is

905   supported by the NIGMS-funded (P41GM103712) National Center for Multiscale Modeling of Biological

906   Systems (MMBioS). This work was supported by Howard Hughes Medical Institute, by The Swartz

907   Foundation, and by the National Science Foundation (NSF DBI-1707356 NeuroNex).

References

1. Abbott, L.F. and W.G. Regehr, *Synaptic computation.* Nature, 2004. **431**(7010): p. 796-803.
2. Alabi, A.A. and R.W. Tsien, *Synaptic vesicle pools and dynamics.* Cold Spring Harb Perspect Biol, 2012. **4**(8): p. a013680.
3. Gross, O.P. and H. von Gersdorff, *Recycling at synapses.* eLife, 2016. **5**: p. e17692.
4. Li F, P.F., Perez E, Eng WS, Melia TJ, Rothman JE, Tareste D, *Energetics and dynamics of SNAREpin folding across lipid bilayers.* Nat Struct Mol Biol, 2007. **14**(10): p. 890-896.
5. Sutton, R.B., et al., *Crystal structure of a SNARE complex involved in synaptic exocytosis at 2.4 A resolution.* Nature, 1998. **395**(6700): p. 347-53.
6. Pobbati, A.V., A. Stein, and D. Fasshauer, *N- to C-terminal SNARE complex assembly promotes rapid membrane fusion.* Science, 2006. **313**(5787): p. 673-6.
7. Li, F., et al., *A half-zippered SNARE complex represents a functional intermediate in membrane fusion.* J Am Chem Soc, 2014. **136**(9): p. 3456-64.
8. Ubach, J., et al., *The C2B domain of synaptotagmin I is a Ca2+-binding module.* Biochemistry, 2001. **40**(20): p. 5854-60.
9. Ubach, J., et al., *Ca2+ binding to synaptotagmin: how many Ca2+ ions bind to the tip of a C2-domain?* EMBO J, 1998. **17**(14): p. 3921-30.
10. Fernandez, I., et al., *Three-dimensional structure of the synaptotagmin 1 C2B-domain: synaptotagmin 1 as a phospholipid binding machine.* Neuron, 2001. **32**(6): p. 1057-69.
11. Zhu, Y. and C.F. Stevens, *Probing synaptic vesicle fusion by altering mechanical properties of the neuronal surface membrane.* Proc Natl Acad Sci U S A, 2008. **105**(46): p. 18018-22.
12. Sugita, S., et al., *Synaptotagmin VII as a plasma membrane Ca(2+) sensor in exocytosis.* Neuron, 2001. **30**(2): p. 459-73.
13. Schiavo, G., et al., *Binding of the synaptic vesicle v-SNARE, synaptotagmin, to the plasma membrane t-SNARE, SNAP-25, can explain docked vesicles at neurotoxin-treated synapses.* Proc Natl Acad Sci U S A, 1997. **94**(3): p. 997-1001.
14. Südhof, T.C., *Neurotransmitter release: the last millisecond in the life of a synaptic vesicle.* Neuron, 2013. **80**(3): p. 675-90.
15. Südhof, T.C., *A molecular machine for neurotransmitter release: synaptotagmin and beyond.* Nat Med, 2013. **19**(10): p. 1227-31.
16. Buzsáki, G., C.A. Anastassiou, and C. Koch, *The origin of extracellular fields and currents--EEG, ECoG, LFP and spikes.* Nat Rev Neurosci, 2012. **13**(6): p. 407-20.
17. Hahnloser, R.H., A.A. Kozhevnikov, and M.S. Fee, *An ultra-sparse code underlies the generation of neural sequences in a songbird.* Nature, 2002. **419**(6902): p. 65-70.
18. Morrison, A., et al., *Exact subthreshold integration with continuous spike times in discrete-time neural network simulations.* Neural Comput, 2007. **19**(1): p. 47-79.
19. Scott, P., A.I. Cowan, and C. Stricker, *Quantifying impacts of short-term plasticity on neuronal information transfer.* Phys Rev E Stat Nonlin Soft Matter Phys, 2012. **85**(4 Pt 1): p. 041921.
20. Tsodyks, M., K. Pawelzik, and H. Markram, *Neural networks with dynamic synapses.* Neural Comput, 1998. **10**(4): p. 821-35.
21. Branco, T. and K. Staras, *The probability of neurotransmitter release: variability and feedback control at single synapses.* Nat Rev Neurosci, 2009. **10**(5): p. 373-83.
22. Klaes, C., et al., *A cognitive neuroprosthetic that uses cortical stimulation for somatosensory feedback.* J Neural Eng, 2014. **11**(5): p. 056024.
23. Collinger, J.L., et al., *High-performance neuroprosthetic control by an individual with tetraplegia.* Lancet, 2013. **381**(9866): p. 557-64.
24. Hochberg, L.R., et al., *Neuronal ensemble control of prosthetic devices by a human with tetraplegia.* Nature, 2006. **442**(7099): p. 164-71.
25. Salmasi, M., et al., *Information Rate Analysis of a Synaptic Release Site Using a Two-State Model of Short-Term Depression.* Neural Comput, 2017. **29**(6): p. 1528-1560.

26. Rotman, Z., P.Y. Deng, and V.A. Klyachko, *Short-term plasticity optimizes synaptic information*
transmission. J Neurosci, 2011. **31**(41): p. 14800-9.
27. Rosenbaum, R., J. Rubin, and B. Doiron, *Short term synaptic depression imposes a frequency dependent*
filter on synaptic information transfer. PLoS Comput Biol, 2012. **8**(6): p. e1002557.
28. Veletić, M., et al., *On the Upper Bound of the Information Capacity in Neuronal Synapses.* IEEE
Transactions on Communications, 2016. **64**(12): p. 5205-5036.
29. Vawter, M.P., et al., *Reduction of synapsin in the hippocampus of patients with bipolar disorder and*
schizophrenia. Mol Psychiatry, 2002. **7**(6): p. 571-8.
30. Crabtree, G.W. and J.A. Gogos, *Synaptic plasticity, neural circuits, and*
the emerging role of altered short-term information processing in schizophrenia. Front Synaptic
Neurosci, 2014. **6**: p. 28.
31. Giovedi, S., et al., *Involvement of synaptic genes in the pathogenesis of autism spectrum disorders: the*
case of synapsins. Front Pediatr, 2014. **2**: p. 94.
32. Deng, P.Y., D. Soika, and V.A. Klyachko, *Abnormal presynaptic short-term plasticity and information*
processing in a mouse model of fragile X syndrome. J Neurosci, 2011. **31**(30): p. 10971-82.
33. Stiles, J.R. and T.M. Bartol, *Monte Carlo methods for simulating realistic synaptic microphysiology using*
MCell, in *Computational Neuroscience: Realistic Modeling for Experimentalists*, E. De Schutter, Editor.
2001, CRC Press: Boca Raton. p. 87-127.
34. Stiles, J.R., et al., *Miniature endplate current rise times less than 100 microseconds from improved dual*
recordings can be modeled with passive acetylcholine diffusion from a synaptic vesicle. Proc Natl Acad
Sci U S A, 1996. **93**(12): p. 5747-52.
35. Nadkarni, S., et al., *Modelling vesicular release at hippocampal synapses.* PLoS Comput Biol, 2010.
6(11): p. e1000983.
36. Kerr, R.A., et al., *FAST MONTE CARLO SIMULATION METHODS FOR BIOLOGICAL REACTION-DIFFUSION*
SYSTEMS IN SOLUTION AND ON SURFACES. SIAM J Sci Comput, 2008. **30**(6): p. 3126.
37. Bartol, T.M., et al., *Computational reconstitution of spine calcium transients from individual proteins.*
Front Synaptic Neurosci, 2015. **7**: p. 17.
38. Coggan, J.S., et al., *Evidence for ectopic neurotransmission at a neuronal synapse.* Science, 2005.
309(5733): p. 446-51.
39. Sargent, P.B., *Nicotinic receptors concentrated in the subsynaptic membrane do not contribute*
significantly to synaptic currents at an embryonic synapse in the chicken ciliary ganglion. J Neurosci,
2009. **29**(12): p. 3749-59.
40. Fioravante, D. and W.G. Regehr, *Short-term forms of presynaptic plasticity.* Curr Opin Neurobiol, 2011.
21(2): p. 269-74.
41. Moulder, K.L. and S. Mennerick, *Reluctant vesicles contribute to the total readily releasable pool in*
glutamatergic hippocampal neurons. J Neurosci, 2005. **25**(15): p. 3842-50.
42. Sakaba, T. and E. Neher, *Calmodulin mediates rapid recruitment of fast-releasing synaptic vesicles at a*
calyx-type synapse. Neuron, 2001. **32**(6): p. 1119-31.
43. Lee, J.S., W.K. Ho, and S.H. Lee, *Actin-dependent rapid recruitment of reluctant synaptic vesicles into a*
fast-releasing vesicle pool. Proc Natl Acad Sci U S A, 2012. **109**(13): p. E765-74.
44. Lee, J.S., et al., *Superpriming of synaptic vesicles after their recruitment to the readily releasable pool.*
Proc Natl Acad Sci U S A, 2013. **110**(37): p. 15079-84.
45. Nadkarni, S., et al., *Short-term plasticity constrains spatial organization of a hippocampal presynaptic*
terminal. Proc Natl Acad Sci U S A, 2012. **109**(36): p. 14657-62.
46. Warren, C.S., et al., *A cautionary note on implications of the well-mixed compartment assumption as*
applied to mass balance models of chemical fate in flowing systems. Environ Toxicol Chem, 2009.
28(9): p. 1858-65.
47. Franks, K.M. and T.J. Sejnowski, *Complexity of calcium signaling in synaptic spines.* Bioessays, 2002.
24(12): p. 1130-44.
48. Varoqueaux, F., et al., *Total arrest of spontaneous and evoked synaptic transmission but normal*
synaptogenesis in the absence of Munc13-mediated vesicle priming. Proc Natl Acad Sci U S A, 2002.
99(13): p. 9037-42.

49.   Imig, C., et al., *The Morphological and Molecular Nature of Synaptic Vesicle Priming at Presynaptic*
Active Zones. Neuron, 2014. **84**: p. 416-431.
50.   Sudhof, T.C., *The synaptic vesicle cycle.* Annu Rev Neurosci, 2004. **27**: p. 509-47.
51.   Südhof, T.C. and J.E. Rothman, *Membrane fusion: grappling with SNARE and SM proteins.* Science,
2009. **323**(5913): p. 474-7.
52.   Tang, J., et al., *A complexin/synaptotagmin 1 switch controls fast synaptic vesicle exocytosis.* Cell, 2006.
126(6): p. 1175-87.
53.   Sun, J., et al., *A dual-Ca2+-sensor model for neurotransmitter release in a central synapse.* Nature, 2007.
450(7170): p. 676-82.
54.   Luo, F., T. Bacaj, and T.C. Südhof, *Synaptotagmin-7 Is Essential for Ca2+-Triggered Delayed*
Asynchronous Release But Not for Ca2+-Dependent Vesicle Priming in Retinal Ribbon Synapses. J
Neurosci, 2015. **35**(31): p. 11024-33.
55.   Bacaj, T., et al., *Synaptotagmin-1 and -7 Are Redundantly Essential for Maintaining the Capacity of the*
Readily-Releasable Pool of Synaptic Vesicles. PLoS Biol, 2015. **13**(10): p. e1002267.
56.   Bacaj, T., et al., *Synaptotagmin-1 and synaptotagmin-7 trigger synchronous and asynchronous phases of*
neurotransmitter release. Neuron, 2013. **80**(4): p. 947-59.
57.   Kochubey, O., X. Lou, and R. Schneggenburger, *Regulation of transmitter release by Ca(2+) and*
synaptotagmin: insights from a large CNS synapse. Trends Neurosci, 2011. **34**(5): p. 237-46.
58.   Schonn, J.S., et al., *Synaptotagmin-1 and -7 are functionally overlapping Ca2+ sensors for exocytosis in*
adrenal chromaffin cells. Proc Natl Acad Sci U S A, 2008. **105**(10): p. 3998-4003.
59.   Maximov, A., et al., *Genetic analysis of synaptotagmin-7 function in synaptic vesicle exocytosis.* Proc Natl
Acad Sci U S A, 2008. **105**(10): p. 3986-91.
60.   Stevens, C.F. and Y. Wang, *Facilitation and depression at single central synapses.* Neuron, 1995. **14**(4):
p. 795-802.
61.   Dobrunz, L.E., E.P. Huang, and C.F. Stevens, *Very short-term plasticity in hippocampal synapses.* Proc
Natl Acad Sci U S A, 1997. **94**(26): p. 14843-7.
62.   Nanou, E., et al., *Calcium sensor regulation of the CaV2.1 Ca2+ channel contributes to short-term*
synaptic plasticity in hippocampal neurons. Proc Natl Acad Sci U S A, 2016. **113**(4): p. 1062-7.
63.   Malgaroli, A. and R.W. Tsien, *Glutamate-induced long-term potentiation of the frequency of miniature*
synaptic currents in cultured hippocampal neurons. Nature, 1992. **357**(6374): p. 134-9.
64.   Kato, K., et al., *Increase in AMPA receptor-mediated miniature EPSC amplitude after chronic NMDA*
receptor blockade in cultured hippocampal neurons. Neurosci Lett, 2007. **418**(1): p. 4-8.
65.   Kaeser, P.S. and W.G. Regehr, *Molecular mechanisms for synchronous, asynchronous, and spontaneous*
neurotransmitter release. Annu Rev Physiol, 2014. **76**: p. 333-63.
66.   DasGupta, A., *Normal Approximations and the Central Limit Theorem*, in *Fundamentals of Probability:*
A First Course. 2010, Springer. p. 213-242.
67.   Kandaswamy, U., et al., *The role of presynaptic dynamics in processing of natural spike trains in*
hippocampal synapses. J Neurosci, 2010. **30**(47): p. 15904-14.
68.   Bischofberger, J., J.R. Geiger, and P. Jonas, *Timing and efficacy of Ca2+ channel activation in*
hippocampal mossy fiber boutons. J Neurosci, 2002. **22**(24): p. 10593-602.
69.   Neher, E. and T. Sakaba, *Multiple roles of calcium ions in the regulation of neurotransmitter release.*
Neuron, 2008. **59**(6): p. 861-72.
70.   Rosahl, T.W., et al., *Short-term synaptic plasticity is altered in mice lacking synapsin I.* Cell, 1993.
75(4): p. 661-70.
71.   Thomson, A.M. and A.P. Bannister, *Release-independent depression at pyramidal inputs onto specific*
cell targets: dual recordings in slices of rat cortex. J Physiol, 1999. **519 Pt 1**: p. 57-70.
72.   Varela, J.A., et al., *A quantitative description of short-term plasticity at excitatory synapses in layer 2/3*
of rat primary visual cortex. J Neurosci, 1997. **17**(20): p. 7926-40.
73.   Zucker, R.S. and W.G. Regehr, *Short-term synaptic plasticity.* Annu Rev Physiol, 2002. **64**: p. 355-405.
74.   Simons, T.J., *Calcium and neuronal function.* Neurosurg Rev, 1988. **11**(2): p. 119-29.
75.   Grewe, B.F. and F. Helmchen, *High-speed two-photon calcium imaging of neuronal population activity*
using acousto-optic deflectors. Cold Spring Harb Protoc, 2014. **2014**(6): p. 618-29.
76.   Grienberger, C. and A. Konnerth, *Imaging calcium in neurons.* Neuron, 2012. **73**(5): p. 862-85.

77.   Nägerl, U.V., et al., *Binding kinetics of calbindin-D(28k) determined by flash photolysis of caged Ca(2+).*
1065   Biophys J, 2000. **79**(6): p. 3009-18.
1066   78.   Sneyd, J., et al., *A model of calcium waves in pancreatic and parotid acinar cells.* Biophys J, 2003. **85**(3):
1067   p. 1392-405.
1068   79.   Rizzoli, S.O. and L. Tabares, *Editorial: Molecular Nanomachines of the Presynaptic Terminal.* Front
1069   Synaptic Neurosci, 2016. **8**: p. 27.
1070   80.   Verkhratsky, A., *The endoplasmic reticulum and neuronal calcium signalling.* Cell Calcium, 2002. **32**(5-
1071   6): p. 393-404.
1072   81.   Bouchard, R., R. Pattarini, and J.D. Geiger, *Presence and functional significance of presynaptic
1073   ryanodine receptors.* Prog Neurobiol, 2003. **69**(6): p. 391-418.
1074   82.   Lanner, J.T., et al., *Ryanodine receptors: structure, expression, molecular details, and function in calcium
1075   release.* Cold Spring Harb Perspect Biol, 2010. **2**(11): p. a003996.
1076   83.   Otsu, K., et al., *Molecular cloning of cDNA encoding the Ca2+ release channel (ryanodine receptor) of
1077   rabbit cardiac muscle sarcoplasmic reticulum.* J Biol Chem, 1990. **265**(23): p. 13472-83.
1078   84.   Nixon, G.F., G.A. Mignery, and A.V. Somlyo, *Immunogold localization of inositol 1,4,5-trisphosphate
1079   receptors and characterization of ultrastructural features of the sarcoplasmic reticulum in phasic and
1080   tonic smooth muscle.* J Muscle Res Cell Motil, 1994. **15**(6): p. 682-700.
1081   85.   Mikoshiba, K., *The IP3 receptor/Ca2+ channel and its cellular function.* Biochem Soc Symp, 2007(74):
1082   p. 9-22.
1083   86.   de Juan-Sanz, J., et al., *Axonal Endoplasmic Reticulum Ca(2+) Content Controls Release Probability in
1084   CNS Nerve Terminals.* Neuron, 2017. **93**(4): p. 867-881.e6.
1085   87.   Hardie, J. and A. Lee, *Decalmodulation of Cav1 channels by CaBPs.* Channels (Austin), 2016. **10**(1): p.
1086   33-7.
1087   88.   Lee, A., et al., *Differential modulation of Ca(v)2.1 channels by calmodulin and Ca2+-binding protein 1.*
1088   Nat Neurosci, 2002. **5**(3): p. 210-7.
1089   89.   Catterall, W.A., K. Leal, and E. Nanou, *Calcium channels and short-term plasticity.* J Biol Chem, 2013.
1090   **288**(15): p. 10742-9.
1091   90.   Ben-Johny, M. and D.T. Yue, *Calmodulin regulation (calmodulation) of voltage-gated calcium channels.*
1092   J Gen Physiol, 2014. **143**(6): p. 679-92.
1093   91.   Fuhrmann, G., et al., *Multiple mechanisms govern the dynamics of depression at neocortical synapses of
1094   young rats.* J Physiol, 2004. **557**(Pt 2): p. 415-38.
1095   92.   Schwaller, B., *Cytosolic Ca2+ buffers.* Cold Spring Harb Perspect Biol, 2010. **2**(11): p. a004051.
1096   93.   Timofeeva, Y. and K.E. Volynski, *Calmodulin as a major calcium buffer shaping vesicular release and
1097   short-term synaptic plasticity: facilitation through buffer dislocation.* Front Cell Neurosci, 2015. **9**: p.
1098   239.
1099   94.   Blum, J.J., et al., *Effect of cytoskeletal geometry on intracellular diffusion.* Biophys J, 1989. **56**(5): p.
1100   995-1005.
1101   95.   Novak, I.L., P. Kraikivski, and B.M. Slepchenko, *Diffusion in cytoplasm: effects of excluded volume due to
1102   internal membranes and cytoskeletal structures.* Biophys J, 2009. **97**(3): p. 758-67.
1103   96.   Neher, E., *Merits and Limitations of Vesicle Pool Models in View of Heterogeneous Populations of
1104   Synaptic Vesicles.* Neuron, 2015. **87**(6): p. 1131-42.
1105   97.   Tsodyks, M.V. and H. Markram, *The neural code between neocortical pyramidal neurons depends on
1106   neurotransmitter release probability.* Proc Natl Acad Sci U S A, 1997. **94**(2): p. 719-23.
1107   98.   Cortes, J.M., et al., *Short-term synaptic plasticity in the deterministic Tsodyks-
1108   Markram model leads to unpredictable network dynamics.* Proc Natl Acad Sci U S A, 2013. **110**(41): p.
1109   16610-5.
1110   99.   Faisal, A.A., L.P. Selen, and D.M. Wolpert, *Noise in the nervous system.* Nat Rev Neurosci, 2008. **9**(4): p.
1111   292-303.
1112   100.   Faber, D.S., et al., *Intrinsic quantal variability due to stochastic properties of receptor-transmitter
1113   interactions.* Science, 1992. **258**(5087): p. 1494-8.
1114   101.   Otmakhov, N., A.M. Shirke, and R. Malinow, *Measuring the impact of probabilistic transmission on
1115   neuronal output.* Neuron, 1993. **10**(6): p. 1101-11.

1116   102.   Seung, H.S., *Learning in spiking neural networks by reinforcement of stochastic synaptic transmission.*
1117   Neuron, 2003. **40**(6): p. 1063-73.
1118   103.   Buesing, L., et al., *Neural dynamics as sampling: a model for stochastic computation in recurrent*
1119   *networks of spiking neurons.* PLoS Comput Biol, 2011. **7**(11): p. e1002211.
1120   104.   Maass, W. and A.M. Zador, *Dynamic stochastic synapses as computational units.* Neural Comput, 1999.
1121   **11**(4): p. 903-17.
1122   105.   Goda, Y. and C.F. Stevens, *Two components of transmitter release at a central synapse.* Proc Natl Acad
1123   Sci U S A, 1994. **91**(26): p. 12942-6.
1124   106.   Smith, S.M., et al., *Calcium regulation of spontaneous and asynchronous neurotransmitter release.* Cell
1125   Calcium, 2012. **52**(3-4): p. 226-33.
1126   107.   Fortune, E.S. and G.J. Rose, *Short-term synaptic plasticity as a temporal filter.* Trends Neurosci, 2001.
1127   **24**(7): p. 381-5.
1128   108.   Izhikevich, E.M., et al., *Bursts as a unit of neural information: selective communication via resonance.*
1129   Trends Neurosci, 2003. **26**(3): p. 161-7.
1130   109.   Lee, C.C., et al., *A kinetic model unifying presynaptic short-term facilitation and depression.* J Comput
1131   Neurosci, 2009. **26**(3): p. 459-73.
1132   110.   Neher, E., *What is Rate-Limiting during Sustained Synaptic Activity: Vesicle Supply or the Availability of*
1133   *Release Sites.* Front Synaptic Neurosci, 2010. **2**: p. 144.
1134   111.   Singh, N., et al., *Presynaptic Endoplasmic Reticulum Contributes Crucially to Short-term Plasticity in*
1135   *Small Hippocampal Synapses.* bioRxiv, 2018: p. 431866.
1136   112.   Pan, X., et al., *The physiological role of mitochondrial calcium revealed by mice lacking the*
1137   *mitochondrial calcium uniporter.* Nat Cell Biol, 2013. **15**(12): p. 1464-72.
1138   113.   Contractor, A., V.A. Klyachko, and C. Portera-Cailliau, *Altered Neuronal and Circuit Excitability in*
1139   *Fragile X Syndrome.* Neuron, 2015. **87**(4): p. 699-715.
1140   114.   Corlew, R., et al., *Presynaptic NMDA receptors: newly appreciated roles in cortical synaptic function and*
1141   *plasticity.* Neuroscientist, 2008. **14**(6): p. 609-25.
1142   115.   Rodriguez-Moreno, A. and O. Paulsen, *Spike timing-dependent long-term depression requires*
1143   *presynaptic NMDA receptors.* Nat Neurosci, 2008. **11**(7): p. 744-5.
1144   116.   Duguid, I.C. and T.G. Smart, *Retrograde activation of presynaptic NMDA receptors enhances GABA*
1145   *release at cerebellar interneuron-Purkinje cell synapses.* Nat Neurosci, 2004. **7**(5): p. 525-33.
1146   117.   Ohno-Shosaku, T., et al., *Endocannabinoid signalling triggered by NMDA receptor-mediated calcium*
1147   *entry into rat hippocampal neurons.* J Physiol, 2007. **584**(Pt 2): p. 407-18.
1148   118.   Navarrete, M., A. Diez, and A. Araque, *Astrocytes in endocannabinoid signalling.* Philos Trans R Soc
1149   Lond B Biol Sci, 2014. **369**(1654): p. 20130599.
1150   119.   Piomelli, D., *The molecular logic of endocannabinoid signalling.* Nat Rev Neurosci, 2003. **4**(11): p. 873-
1151   84.
1152   120.   van den Bogaart, G., et al., *One SNARE complex is sufficient for membrane fusion.* Nat Struct Mol Biol,
1153   2010. **17**(3): p. 358-64.
1154   121.   Mohrmann, R., et al., *Fast vesicle fusion in living cells requires at least three SNARE complexes.* Science,
1155   2010. **330**(6003): p. 502-5.
1156   122.   Albert, A., *Estimating the Infinitesimal Generator of a Continuous Time, Finite State Markov Process.*
1157   The Annals of Mathematical Statistics, 1962. **33**(2): p. 727-753.
1158   123.   Nelder, J.A. and R. Mead, *A Simplex Method for Function Minimization* Comput J, 1965. **7**(4): p. 308-
1159   313.
1160   124.   McKinnon, K.I.M., *Convergence of the Nelder--Mead Simplex Method to a Nonstationary Point.* SIAM J.
1161   Optim., 1998. **9**(1): p. 148-158.
1162   125.   Lagarias, J., et al., *Convergence Properties of the Nelder--Mead Simplex Method in Low Dimensions.*
1163   SIAM Journal on Optimization, 1998. **9**(1): p. 112-147.
1164   126.   Körber, C. and T. Kuner, *Molecular Machines Regulating the Release Probability of Synaptic Vesicles at*
1165   *the Active Zone.* Front Synaptic Neurosci, 2016. **8**: p. 5.
1166   127.   Sakaba, T., *Roles of the fast-releasing and the slowly releasing vesicles in synaptic transmission at the*
1167   *calyx of Held.* J Neurosci, 2006. **26**(22): p. 5863-71.

1168   128.   Schneggenburger, R. and E. Neher, *Intracellular calcium dependence of transmitter release rates at a*
1169   *fast central synapse.* Nature, 2000. **406**(6798): p. 889-93.
1170   129.   Wölfel, M., X. Lou, and R. Schneggenburger, *A mechanism intrinsic to the vesicle fusion machinery*
1171   *determines fast and slow transmitter release at a large CNS synapse.* J Neurosci, 2007. **27**(12): p. 3198-
1172   210.
1173   130.   Xu, J., et al., *Synaptotagmin-1 functions as a Ca2+ sensor for spontaneous release.* Nat Neurosci, 2009.
1174   **12**(6): p. 759-66.
1175   131.   Delvendahl, I., et al., *Reduced endogenous Ca2+ buffering speeds active zone Ca2+ signaling.* Proc Natl
1176   Acad Sci U S A, 2015. **112**(23): p. E3075-84.

1177