Adipose gene expression profiles reveal novel insights into the

adaptation of northern Eurasian semi-domestic reindeer

(*Rangifer tarandus*)

Short title: Reindeer adipose transcriptome

Melak Weldenegodguad[1, 2], Kisun Pokharel[1¶], Laura Niiranen[3¶], Päivi Soppela[4,]

Innokentyi Ammosov[5], Mervi Honkatukia[6], Heli Lindeberg[1], Jaana Peippo[1, 6], Tiina

Reilas[1], Nuccio Mazzullo[4], Kari A. Mäkelä[3], Tommi Nyman[7], Arja Tervahauta[2],

Karl-Heinz Herzig[8, 9, 10], Florian Stammler[4], Juha Kantanen[1*]

[1] Natural Resources Institute Finland (Luke), Jokioinen, Finland

[2] Department of Environmental and Biological Sciences, University of Eastern

Finland, Kuopio, Finland

[3] Research Unit of Biomedicine, Faculty of Medicine, University of Oulu, Oulu,

Finland

[4] Arctic Centre, University of Lapland, Rovaniemi, Finland

[5] Board of Agricultural Office of Eveno-Bytantaj Region, Batagay-Alyta, The Sakha

Republic (Yakutia), Russia

[6] NordGen—Nordic Genetic Resource Center, Ås, Norway

[7] Department of Ecosystems in the Barents Region, Norwegian Institute of

Bioeconomy Research, Svanvik, Norway

[8] Research Unit of Biomedicine, Medical Research Center, Faculty of Medicine,

University of Oulu

[9] Oulu University Hospital, Oulu, Finland

[10]Pediatric Institute, Poznan University of Medical Sciences, Poznan, Poland

* Corresponding author

Email: juha.kantanen@luke.fi (JK)

¶These authors contributed equally to this work.

Abstract

Reindeer (*Rangifer tarandus*) are semi-domesticated animals adapted to the

challenging arctic conditions of northern Eurasia. Adipose tissues play a crucial role

in animals living in northern environments by altering gene expression in their

tissues to regulate energy homeostasis and thermogenic activity. Here, we

performed transcriptome profiling by RNA sequencing of adipose tissues from three

different anatomical depots: metacarpal (bone marrow), perirenal, and prescapular

fat in Finnish and Even reindeer (in Sakha) during two seasonal time points (spring

and winter). On average 36.5 million pair-ended clean reads were obtained for each

sample, and a total of 16,362 genes were expressed in our data. Gene expression

profiles in metacarpal tissue were distinct and clustered separately from perirenal

and prescapular adipose tissues. Notably, metacarpal adipose tissue appeared to

have a significant role in the regulation of the energy metabolism of reindeer in

spring when their nutritional condition is poor after winter. During spring, when the

animals are in less optimal condition, genes associated with the immune system

(e.g., *CCL2, CCL11, CXCL14, IGSF3, IGHM, IGLC7, IGKC, JCHAIN,* and *IGSF10)*

were upregulated in the perirenal and prescapular adipose tissue, while genes

involved in energy metabolism (e.g., *ACOT2, APOA1, ANGPTL1, ANGPTL8,*

ELOVL7, *MSMO1*, *PFKFB1*, and *ST3GAL6*) were upregulated in metacarpal

tissue. Even reindeer harboured relatively fewer significantly differentially

expressed genes than Finnish reindeer, irrespective of the season, possibly owing

to climatic and management differences. Moreover, blood and tissue parameters

reflecting general physiological and metabolic status showed less seasonal

variation in Even reindeer than in Finnish reindeer. This study identified adipose

candidate genes potentially involved in immune response, fat deposition, energy

metabolism, development, cell growth, and organogenesis. Taken together, this

study provides new information on the mechanisms by which reindeer adapt to less

optimal arctic conditions.

Keywords: adaptation; immune process; *PRDM9*; lipid metabolism; metabolites;

metacarpal adipose tissue; mRNA; perirenal adipose tissue; prescapular adipose

tissue; *UCP1*

Introduction

Native to northern and subarctic regions of Eurasia, reindeer (*Rangifer tarandus*)

have societal, cultural, and ecological values for the livelihoods of arctic indigenous

people and pastoralists, and have multiple socio-economic roles, such as providing

meat, hides, milk, and serving as a means of transportation [1–3]. Reindeer survive

in challenging northern and extreme arctic environments characterized by low

temperatures, prolonged daylight during summers, and darkness and limited

availability of grazing resources during long winters [3,4]. Adipose tissues are vital

for reindeer to adapt to such extreme conditions [5,6]. Adipose tissues are important

organs for several functions in energy metabolism that are crucial for survival and

successful reproduction. White adipose tissue (WAT) stores energy in the form of

lipids and serves as a long-term energy reserve, whereas brown adipose tissue

(BAT) contributes to both thermohomeostasis and energy balance by producing heat

via the function of uncoupling protein 1 (*UCP1*) [7]. WAT is also an important

endocrine organ that secretes several hormones, including adipomyokines and

cytokines, which contribute to energy metabolism and immunity and act as signals

to the central nervous system [8,9]. The reindeer is a lean animal, but it also relies

on WAT as a source of energy and hormonal signals in changing environmental

conditions [5]. In newborn reindeer, BAT plays a crucial role in regulating non-

shivering thermogenesis, but its effect diminishes over time [10,11]. However,

browning of WAT has been observed in various species during later stages of life

after chronic cold exposure [12] and due to pharmacological and nutritional agents

[13]. Bone marrow has a specific type of adipose tissue (BMAT) that acts as an

energy reservoir, contributes to local and systemic metabolic processes [14,15], and

undergoes dynamic changes [16]. BMAT decreases as a result of starvation in freely

grazing animals [17–19].

The majority of adipose transcriptome studies using high-throughput RNA

sequencing (RNA-Seq) have been limited to mice [20,21], sheep [22–24], pigs [25–

27], and humans [28–30]. Adipose transcriptomes are affected by the type of

adipose, as well as by the sex and age of the animal [31]. Changes in gene

expression within adipose tissues in response to temperature fluctuations have been

studied mostly in mice so far [20,21,32]. However, gene expression profiles in

reindeer adipose tissues have thus far not been investigated. Adipose tissues from

various parts of the body have their own unique functions. In the present study, we

investigated gene expression profiles of three adipose tissue depots: metacarpal

(M), perirenal (P), and prescapular (S) tissues. These adipose depots were selected

because they represent visceral (P), peripheral (S), and bone marrow (M) fat. These

anatomical depots are also expected to reflect different metabolic functions. For

example, the prescapular area is a major BAT depot in newborn reindeer [11] and

thus an interesting target for regulating the expression of *UCP1*, as well as other

markers for cold adaptation. The samples were collected from the Finnish and Even

(Sakha Republic, Yakutia, Russia) reindeer breeds (or populations), which belong to

two different phylogenetic clusters of Eurasian reindeer (Pokharel K. et al.,

unpublished) and differ in present-day management and feeding practices. In

Yakutia, reindeer feed on natural pastures with expressed seasonal variation and

high migratory behaviour (high lichen content in winter, fresh leaves in spring, grass,

and mountain herbs in summer), whereas in Finland, extra fodder (concentrates) is

provided during peak winter (February and March). Reindeer herders observe that

the lichen-rich diet in winter helps reindeer keep their weight and survive the cold,

while the grassy diet in summer accounts for the principal annual gain in weight. In

Yakutia, reindeer have to survive in extreme climatic conditions where the annual

temperature fluctuates from -60℃ up to more than +30℃.

Here, we aim to obtain insights into the seasonal fluctuations in the transcriptome

profiles of three adipose tissues (metacarpal, perirenal, and prescapular) in two

reindeer populations (Even and Finnish). The tissue sampling was conducted during

(early) winter (November–December) and (early) spring (April), when the animals

were in the best and worst nutritional conditions, respectively. For the Even reindeer,

we also compared the transcriptome profiles of these adipose tissues between male

and female individuals. Moreover, to assess the health and physiological condition

of the reindeer in different seasons and geographical locations, and to complement

genetic analyses with phenotypic data, we analysed several blood metabolites along

with hormones regulating energy metabolism, such as blood insulin, leptin, and

hormone sensitive lipase (HSL), and proteins such as *UCP1* and *COX4* from adipose

tissue. We expect that assessing changes in gene expression in reindeer adipose

tissue due to seasonal variation may reveal adaptation mechanisms and help to

understand the evolution of this adaptive response in reindeer and other mammalian

species sharing northern Eurasian habitat conditions.

Results

RNA sequencing and mapping

A total of 220.5 gigabases (Gb) of RNA-seq data were generated from 56

adipose tissue samples collected from 19 Finnish and Even reindeer. As the

adapters were trimmed automatically, and the Phred quality scores of the reads from

all samples were greater than 30, we did not perform further trimming and quality

filtering. The number of reads per sample ranged from 26.6 million (M) (4.0 Gb) to

217 M (32.6 Gb), with a mean of high-quality 36.5 M, 2 × 75 bp pair-ended reads per

sample (S1 Table). As shown in S1 Table, the two samples (FR12_SCAP and

YR1_SCAP), revealing the highest numbers of reads (217 M and 165 M,

respectively), will be used to detect lncRNAs in a future study. The proportion of

reads mapped to the reindeer reference genome ranged from 81% to 92%, with, on

average, >90% of the reads from each sample uniquely mapped to the reindeer draft

genome assembly (S2 Table). Raw sequence reads in compressed fastq format

(fastq.gz) analyzed in this study have been deposited to the European Nucleotide

Archive (ENA) and are publicly available under accession XXXXXX

144   Gene expression overview

A total of 16,362 genes were expressed (cpm > 0.5 in at least two samples) in

all 56 samples (S3 Table), representing approximately 60% of the 27,332 reindeer

genes reported in the draft reindeer genome assembly annotation file [3]. The

highest number of genes were expressed in metacarpal adipose tissue (n = 15,761),

followed by prescapular (n = 15,087) and perirenal (n = 14,920) adipose tissues.

Moreover, we examined the expressed genes across these three tissues to search

for shared and uniquely expressed genes in the respective region and season (Fig

1). In all cohorts (spring and winter Finnish and Even samples), the highest number

of expressed genes was found in the metacarpal adipose tissue. This tissue also

displayed a higher number of tissue-specific expressed genes than did the perirenal

and prescapular adipose tissues (Fig 1). In various region-season comparisons

between the adipose tissues, >13,000 genes were commonly expressed. To assess

expression similarity among the samples, we performed Principal Component

Analysis PCA based on the top 500 most variable genes (Fig 2, S1 Fig). The PCA

plot (Fig 2) shows that the metacarpal tissue samples were clustered separately from

the other two tissues, and additional grouping of the samples along axis 2 based on

two reindeer breeds. Similarly, a hierarchal clustering based on the top 25 genes

with the highest variance across all samples also showed a similar pattern that

clearly separated metacarpal tissue and the other tissues (S2 Fig).

While more than 13,000 genes were commonly expressed in all tissues, several

tissue-specific genes were identified in this study. In line with the distinct cluster

observed in the PCA plot, more than 730 genes were uniquely expressed in the

metacarpal adipose tissue (Fig 1). We further explored the genes specific to this

tissue by removing the lowly expressed genes (TPM < 1). Out of 875 genes, 61

were commonly expressed in metacarpal adipose tissue in the different

experimental groups. The metacarpal adipose tissue of Finnish reindeer

collected in the winter had the largest number of uniquely expressed genes (n =

181), whereas Finnish metacarpal tissue collected in the spring possessed the

lowest (n = 71) number of uniquely expressed genes. By contrast, Even reindeer

samples collected in either season had roughly similar numbers of uniquely

expressed genes (138 in spring and 136 in winter samples) (Fig 3).

Among the 61 genes that were unique to metacarpal adipose tissue yet shared

by Finnish and Even reindeer, annotations were not available for 10 genes. The

metacarpal-specific genes included several homeobox proteins (eg. *HOXD13*,

HOXA11, *HOXD11*, *HOXA13*, *DLXC*), bone sialoprotein 2 (*IBSP*), osteomodulin

(*OMD*), carbonic anhydrase 3 (*CA3*), C-X-C motif chemokine 10 (*CXCL10*),

nuclear receptor-interacting protein 3 (*NRIP3*), and R-spondin-2 (*RSPO2)* (Table

S4).

Results from differential gene expression analyses

Seasonal differences in gene expression. For an insight into the effects of

seasonal conditions, we compared gene expression profiles of adipose tissues from

spring versus winter samples separately in the Finnish and Even reindeer. Finnish

reindeer showed a higher number of significant differentially expressed genes

(DEGs) than the Even reindeer (Table 2).

Table 2. The number of significantly identified differentially expressed genes (DEGs)

in Finnish (F) and Even (E) reindeer due to seasonal changes (S, spring, W, winter)

in each tissue.

Comparison	Total DEGs	Upregulated	Downregulated
FM-S vs. FM-W	346	195	151
FP-S vs. FP-W	583	273	310
FS-S vs. FS-W	611	325	286
EM-S vs. EM-W	156	57	99
EP-S vs. EP-W	103	45	58
ES-S vs. ES-W	176	126	50

Altogether 346 genes were differentially expressed between seasons in

metacarpal tissues of Finnish reindeer; of these, 195 were upregulated in spring

samples and the rest were upregulated in the winter samples (Table 2, S5 Table and

S4 Fig). Genes involved in metabolism (*ANGPTL8*, *BDH1*, *ASIP*, *QPRT*) and stress

response (*BOLA3*, *GPR158*) were strongly upregulated in spring (LFC > 4) and

those particularly associated with immune functions (*CCL11*, *CXCL8*, *CD33*, *IL1B*,

IRF4) were strongly downregulated (LFC < -3.8) in spring metacarpal tissue (S5

Table and S4 Fig).

Among 583 DEGs in the perirenal tissue, genes such as *CCL11, TRH, MT1A,*

RPL38, AREG, NPW, SRSF3, RPS29, RPL36A and *SLC11A1*, were highly

upregulated (LFC > 4) in the spring samples and *MP68, RPL39, SLC39A12, SCN3A,*

RGS9, HSPA6, TNFRSF10B, EDIL3, RTP1 and *ABCC4* were highly downregulated

(LFC < -3.7) (Table 2, S6 Table and S5 Fig). The DEGs upregulated in the spring

samples include genes participating in the immune system (*SLC11A1*, *COMMD6*,

BATF, *CCL19*), ribosomal and transcription processes (*RPL38*, *RPL36A*, *FKBP11*).

On the other hand, downregulated genes were associated with signalling or signal

transduction (*RGS9, RAPGEF5, COL4A5, PTGER2, MAML3, PDE5A*), cell

differentiation or organogenesis (*GJA5, THSD7A, DACH1, MMRN2, NHSL2,*

ZFHX3), and interaction of glucose and fatty acid metabolism (*PDK4*, SIK1).

The highest number (n = 611) of DEGs was found in prescapular tissue, 325 of

which were upregulated in the spring samples (Table 2, S7 Table and S6 Fig). In

prescapular adipose tissue from the spring, genes such as *SERPINA3-7, ESD,*

APOH, SERPINI2, SERPINC1, TCEB2, ITIH1, WNT9B, C8B, TRH, ASCL2, GSTA2

and *VPREB1* were among the top upregulated genes (LFC > 4.5) and *LHFP,*

TNFRSF10B, CRISP3, NEGR1, KCNA2, PTGER2, HSPA6, GTSF1 and *RPL30*

were among the top downregulated genes (LFC < -3.5) (S7 Table and S6 Fig). A

total of 30 genes, including *APOLD1, CYP26B1, IGFBP5, MICA, MICB, MT-ATP8,*

PDK4 and SCP2, were commonly differentially expressed in three tissues (S3 Fig).

Many of the genes in the prescapular tissue that were upregulated during spring

were associated with inflammatory or immunological responses (*FXYD5, CCL19,*

IFIT3, *CARD9*, *GMFG*, *SLC11A1*), feeding behaviour (*NPW*), adipogenesis (*DLK1,*

CITED4), cell growth or differentiation (*MGP*, *ECSCR*, *TIMP1*, *MT1A*), and

spermatogenesis (*SCHBP1L*, *MORN2*). Similarly, downregulated DEGs were

associated with signalling (*RGS5*, *DCBLD2*, *RASGEF1B*, *RAPGEF5*),

organogenesis (*PTGER2*, *KLF7*, *PHACTR2*, *KMT2A*, *DACH1*, *PDLIM5*), and lipid

metabolism (*ARFGEF3*, *PDK4*, *FOXO1*, *PITPNC1*).

Similar pairwise comparison in Even reindeer revealed 156, 103, and 176 DEGs

in metacarpal, perirenal, and prescapular tissues, respectively (Table 2, S8-S10

Tables, S8-S10 Figs). Prescapular adipose tissue had the highest number of unique

significant DEGs (n = 141), followed by metacarpal (n = 135), and perirenal (n = 68)

(S7 Fig). In both breeds, prescapular tissue harboured the highest (611 in Finnish

and 176 in Even reindeer) number of DEGs (S7 and S10 Tables). Interestingly, while

the number of upregulated genes was higher in the metacarpal tissue of Finnish

samples from the winter collection, the same was not observed in Even reindeer. In

the Even reindeer we, found seven common DEGs (*ORC5*, *ADI1*, *RPS15A*, *DAGLA*,

JCHAIN, ENSP00000353290, and *FEZ1*) among all tissues (S7 Fig). DEGs that

were upregulated in the spring samples of Even metacarpal adipose tissue appeared

to have roles in lipid/fatty acid metabolism (*ELOVL7*, *APOA1*, *ACOT2*, *ANGPTL1*),

cell structural functions (*CAPN6*, *MYL9*, *CLDN5*), and oxygen metabolism (*FMO1*,

FMO2, *AOC1*, *STEAP1*). Downregulated DEGs in Even metacarpal adipose tissue

were associated with development and organogenesis (*DMAP1*, *COL27A1*, *CA2*,

NRG3, *JAK3*, *EPB41L3*, *ESM1*) and immune system (*C4A*, *IL17RB*, *HLA-DOA*,

SIGLEC1, *CPA3*). In Even perirenal adipose tissue, several immunoglobulin related

genes were upregulated during spring, suggesting activation of the immune system

by pathogens. However, downregulated genes were mainly associated with

lipid/energy metabolism (*ACACB*, *PRLR*, *APOL6*) and functions related to growth

and development (*MEGF8*, *DAGLA*, *IGSF10*, *GRIP2*, *NSMF*). Among the three

adipose tissues in Even reindeer, immune related DEGs (e.g., *JCHAIN*, *IGHM*,

IGKC, *IGHV3-6*, *IGLC7*, *MUCM*) were predominantly upregulated in the prescapular

tissue, whereas downregulated genes in the prescapular tissue were associated with

lipid/energy metabolism (*TP53INP2*, *PPARGC1B*, *ACACB*) and growth and

development (*DAGLA*, *IGS10*, *BMP5*, *FGFR2*, *NRG3*, *CYP26B1*).

Gene expression differences between the Finnish and Even reindeer. We made

six pair-wise comparisons (three each for spring and winter samples) to identify

DEGs between Finnish and Even reindeer (Table 3, S11-S16 Tables) of which the

highest number (n = 504) of DEGs was present in metacarpal tissues collected

during winter, whereas the same tissue revealed the lowest number (n = 126) of

DEGs during spring (Table 2, S11 and S14 Tables). In comparisons involving spring

samples, the highest numbers of DEGs were found in prescapular adipose tissue.

Four genes, *FIBP*, *CREB3L3*, *CLDN4,* and *ALKBH3,* were exclusively

upregulated in the Finnish reindeer irrespective of the seasons. *FIBP* (*FGF1*

Intracellular Binding Protein) is known to promote mitogenic action to induce

morphogenesis and differentiation. *CREB3L3* has been linked to triglyceride

metabolism and growth suppression. *CLDN4* (Claudin 4) is a member of the claudin

gene family. Being integral membrane proteins, claudins, in general, have a vital role

in regulating the transport of solutes and ions through calcium-independent cell-

adhesion activity.

Similarly, *TMEM182*, *AACS*, *FAM159B,* and *C19ORF80* were always

upregulated in all five comparisons except between EM-S vs FM-S. Among the four

genes, we did not find any relevant information about the function of *FAM159B*.

There is relatively little information available on *TMEM182*, but its upregulation may

be associated with adipose growth and remodelling [33,34]. In the present context,

a greater abundance of *TMEM182* in Even reindeer males might be linked to

castration, as castrated males are known to accumulate more adipose tissues.

AACS (*Acetoacetyl-CoA Synthetase*) appears to be involved in ketone body

metabolism during adipose tissue development. *C19ORF80* (alternatively

ANGPTL8, *Angiopoietin-like 8*) is known to mediate the transition between fasting

and re-feeding and has an important role in the storage of fatty acids in adipose

tissue during the refeeding state [35]. Interestingly, *C19ORF80* was downregulated

in Even reindeer compared to Finnish reindeer among spring samples. We speculate

that the additional winter feeding for the Finnish reindeer might have a role in the

differential expression of *C19ORF80*.

Table 3. The number of significantly identified DEGs in the metacarpal (M),

perirenal (P), and prescapular (S) adipose tissues of Finnish (F) and Even (E)

reindeer that were compared separately for spring (-S) and winter (-W) samples.

Comparison	Total DEGs	Upregulated	Downregulated
EM-S vs FM-S	126	78	48
EP-S vs FP-S	301	189	112
ES-S vs FS-S	385	229	156
EM-W vs FM-W	504	336	168
EP-W vs FP-W	401	182	219
ES-W vs FS-W	365	154	221

Gender difference in gene expression. Our comparison of gene expression

between female and male Even reindeer samples collected during winter revealed a

total of 327 significant DEGs in the three adipose tissues (S17-S19 Tables).

Table 4. Summary of DEGS from male (-M) and female (-F) comparisons in

metacarpal (M), perirenal (P), and prescapular (S) adipose tissues of Even (E)

reindeer.

Comparison	Total DEGs	Upregulated	Downregulated

EM-F vs EM-M	225	78	147
EP-F vs EP-M	104	54	50
ES-F vs ES-M	49	27	22

298  We identified a total of 225 significant DEGs between female and male reindeer

299 in metacarpal tissue (EM-F vs. EM-M) (Table 4 and S17 Table). Of 225 significant

300 DEGs, 78 genes were upregulated, and 147 genes were downregulated in EM-F

301 (Table 4 and S17 Table). The upregulated genes in female samples included

302 *ANGPTL1, CX3CR1, CYP2B11, CYP2F3, CYP4B1, FABP6, ELOVL7, IGHV3-6,*

303 *SLC19A3, SLC35F1* and *SLC6A17*, whereas genes such as *BMP1, BMP, IBSP,*

304 *EFS, CCL19, IGDCC4, IL17RB, NAV3, NCAM1, NRL, OLFML2B, OLFML3,*

305 *PRDM9, SCUBE1* and *FLT4* were downregulated.

306  In perirenal tissue, 104 significantly DEGs were detected between female and

307 male reindeer, of which 54 and 50 genes were upregulated and downregulated in

308 EP-F, respectively (Table 4 and S18 Table). The upregulated genes in female

309 samples included *ABCG1, ACSL6, SLC14A1, SLC16A2, SLC4A10, SLC9A2,*

310 *ZNF219, PLCD3, PTER, PLA2G5, ETNPPL, PLCD3, S1PR3,* and *S1PR3*, while

311 *ATP5H, CCL3, COX7A1, CXCL9, NMB, NRL, PRDM9, SLC19A1, SLC22A5, ZFX,*

312 and *ZRSR2* were downregulated.

313  Furthermore, 49 significantly DEGs were detected between female and male

314 reindeer in prescapular tissue (ES-F vs. ES-M), of which 27 and 22 genes were

315 upregulated and downregulated in ES-F, respectively (Table 4 and S19 Table).

316 *GYS2, ARHGEF5, ABCC4, SUMO1, RPL39, SLC4A10,* and *ACSL6* were examples

317 of upregulated genes in female samples, and genes such as *TXLNG, DDX3Y,*

318 *USP9X, EIF2S3X, PRDM9, KDM6A, ZRSR2,* and *UCP1* were downregulated.

A total of 10 genes (*PRDM9, DDX3Y, ZRSR2, EIF2S3X, KDM6A, ZFX, UBA1,*

USP9X, TXLNG, and *NRL*) were always upregulated in males irrespective of the

adipose tissue type. These genes have diverse functions and may be linked to

speciation (*PRDM9*), male infertility or spermatogenic failure (*DDX3Y*), mRNA

splicing (*ZRSR2*), lipid metabolism (*EIF2S3X*), circadian rhythm (*USP9X*), cell cycle

regulation (*TXLNG*), and photoreceptor development/function (*NRL*).

We did not find any genes commonly upregulated in all tissues of female

samples; however, two upregulated DEGs (*CPE*, *RPL34*) were shared by

metacarpal and perirenal adipose tissues, while 12 (*ARHGEF5*, *SLC4A10*, *ABCC4*,

SGCG, *HAND1*, *CPXM2*, *GYS2*, *ACSL6*, *ANK1*, *NPR3*, *TBX20*, and *PGR*) were

shared among prescapular and perirenal adipose tissues.

330   Results of GO enrichment analyses

Enriched GO terms associated with DEGs resulting from seasonal

comparisons: Finnish reindeer. Enrichment analyses based on significantly

differentially expressed genes revealed several GO terms and KEGG pathways, thus

highlighting the important biological and physiological activities in each tissue. From

the 346 significant DEGs between FM-S and FM-W (Table 2 and S5 Table), 112

genes lacked GO annotations. Separate GO enrichment analyses were performed

for the downregulated (n = 117 with GO annotation) and upregulated (n = 117 with

GO annotation) genes. GO analysis indicated that 16 and 36 GO terms were

significantly associated with downregulated and upregulated DEGs, respectively

(S20 Table). Downregulated DEGs were represented in GO terms mainly associated

with immune processes (e.g., "immune system process", "immune response") and

stimulus (e.g., "response to stimulus", "response to chemical", "chemotaxis"),

whereas upregulated DEGs were represented in terms associated with metabolic

processes ("ATP metabolic process", "nucleoside metabolic process", "single-

organism metabolic process", "carbohydrate derivative metabolic process"), and

transport (e.g., "proton transport", "hydrogen transport", "cation transmembrane

transport") in the metacarpal adipose tissue of Finnish reindeer.

In contrast to metacarpal adipose tissue, both the perirenal and prescapular

adipose tissues had downregulated genes represented in a relatively higher number

of GO terms (n = 29, n = 55, respectively; S21 and S22 Tables) than upregulated

genes (n = 16, n = 7, respectively; S21 and S22 Tables). The DEGs upregulated

during spring in Finnish reindeer perirenal adipose tissue were associated with

metabolic ("cellular amide metabolic process", "organonitrogen compound metabolic

process") and biosynthetic ("amide biosynthetic process", "organonitrogen

compound biosynthetic process") processes, and those upregulated during winter

were associated with signalling ("signalling", "G-protein coupled receptor signalling

process", cell communication", "single organism signalling", "signal transduction"),

and regulation ("regulation of biological process", "regulation of cellular process",

"biological regulation") related biological processes. Altogether, 35 of 55 GO terms

associated with DEGs upregulated in the prescapular adipose tissue of Finnish

reindeer from winter sampling were categorised under "biological process", whereas

none of the seven GO terms enriched in upregulated DEGs of spring samples were

categorized under "biological process". Moreover, GO terms associated with the

regulation of several processes (e.g., "regulation of metabolic process," "regulation

of RNA biosynthetic process," "regulation of biological process," "regulation of gene

expression"), response (e.g., "cellular response to organic cyclic compound",

"response to steroid hormone", "response to lipid", "response to hormone", "cellular

response to stimulus) and transcription activities (e.g., "transcription coactivator

activity" and "transcription cofactor activity") categorised upregulated DEGs of

Finnish prescapular adipose tissue collected during winter.

Enriched GO terms associated with DEGs resulting from seasonal

comparisons: Even reindeer. The differential expression analysis between EM-S

and EM-W revealed 156 significantly DEGs (Table 2 and S8 Table), of which 55

lacked GO annotations. GO enrichment analysis of the downregulated genes (n =

77 with GO annotations) yielded seven significantly represented GO terms (S23

Table), including "metalloendopeptidase activity" and "metallopeptidase activity,"

whereas no significantly enriched GO terms were found in the upregulated genes.

In Even reindeer perirenal tissue, of the 103 significantly DEGs in EP-S vs EP-W

(Table 2 and S9 Table), 32 genes did not have GO annotations. Gene enrichment

analysis of the upregulated and downregulated DEGs revealed no statistically

significantly represented GO terms. From the list of 176 significantly DEGs between

ES-S and ES-W (Table 2 and S10 Table), 59 genes lacked GO annotation. Gene

enrichment analysis of the upregulated and downregulated DEGs revealed no

statistically significantly represented GO terms.

Enriched GO terms associated with DEGs resulting from location comparison.

DEGs resulting from the comparison of the metacarpal adipose tissue between Even and Finnish reindeer did not reveal any GO terms. Upregulated genes in the perirenal adipose tissue of Even reindeer from spring sampling were associated only with the GO term "cofactor binding," whereas downregulated genes were not enriched in any GO terms. Similar comparison of perirenal adipose tissue from winter sampling revealed eight and 46 GO terms associated with upregulated and downregulated genes, respectively, in Even reindeer (S24 Table).

In spring prescapular adipose tissues of Even reindeer, a total of 19 GO terms including signalling (e.g., "single organism signalling," "signal transduction," "G-protein coupled receptor signalling pathway"), and response to stimulus ("response to external stimulus," "cellular response to stimulus") were associated with upregulated genes (S25 Table), while none of the downregulated genes were enriched in any GO terms. Similar comparison of winter samples in Even reindeer revealed three overrepresented GO terms ("anion binding", "cofactor binding" and "pyridoxal phosphate binding") and 20 underrepresented GO terms (including six signalling terms, "G-protein coupled receptor activity", "receptor activity" and "cell communication") (S26 Table).

Enriched GO terms associated with DEGs resulting from gender comparison.

From the list of 225 significantly DEGs identified in EM-Fversus EM-M comparison (S17 Table), 59 genes did not have GO annotations. The GO analysis results of the upregulated (n = 58 with GO annotations) and downregulated (n = 108 with GO

annotation) DEGs in female metacarpal tissue revealed one and 31 significantly

represented GO terms, respectively (S27 Table). The upregulated DEGs in female

metacarpal tissue were significantly enriched in only one GO term: "oxidoreductase

activity, acting on paired donors, with incorporation or reduction of molecular

oxygen". The downregulated DEGs were significantly enriched in 18 biological

processes, 10 molecular functions, and three cellular component categories (S27

Table). Out of the 18 biological processes, nine GO terms were associated with

developmental   processes,   such   as   "circulatory   system   development,"

"cardiovascular system development," "blood vessel development," "vasculature

development," and "anatomical structure development" (S27 Table) Moreover,

downregulated DEGs in female metacarpal tissue were also significantly enriched in

the biological process represented by the GO terms "cell adhesion," "angiogenesis,"

"blood vessel morphogenesis," and "anatomical structure formation involved in

morphogenesis" (S27 Table).

From the list of 104 significantly DEGs identified in EP-F versus EP-M

comparison (S18 Table), 15 lacked GO annotations. The GO analysis results of the

upregulated (n = 48 with GO annotations) DEGs in female perirenal tissue revealed

six significantly represented cellular component categories, including "plasma

membrane", "cell periphery", "plasma membrane part," and "integral component of

membrane"; no significantly represented GO terms were associated with

downregulated DEGs (n = 41 with GO annotations) (S28Table).

From the list of 49 significantly DEGs identified in ES-F vs. ES-M comparison

(S19 Table), seven genes lacked GO annotation. The GO analysis result of the

upregulated (n = 23 with GO annotation) and downregulated (n = 19 with GO

annotation) DEGs in female prescapular tissue revealed three significant

represented molecular function categories and no significantly enriched GO terms,

respectively. The significantly represented GO terms associated with downregulated

DEGs in female reindeer prescapular tissue included "cation binding," "ion binding,"

and "metal ion binding," indicating their role in the acquisition of mineral nutrients:

iron, zinc, and calcium.

439   Results of KEGG pathway analyses

KEGG pathways associated with seasonal comparisons. Finnish reindeer. We

performed KEGG pathway analysis using GAGE to identify pathways that were

differentially regulated by season (FM-S vs FM-W). Pathway enrichment analysis

revealed 18 and 5 significantly downregulated and upregulated KEGG pathways,

respectively, in spring metacarpal adipose tissue (S29 Table). Most of the

downregulated pathways were associated with the immune system, environmental

information processing, and signal transduction (S29 Table). Pathways associated

with immune system included "cytokine-cytokine receptor interaction," "chemokine

signalling pathway," "Fc gamma R-mediated phagocytosis," "IL-17 signalling

pathway," and "T cell receptor signalling pathway" (S29 Table). In addition, the

significantly downregulated pathways in spring metacarpal tissue associated with

environmental information processing and signal transduction included: "TNF

signalling pathway," "MAPK signalling pathway," "NF-kappa B signalling pathway,"

"Jak-STAT signalling pathway," and "ErbB signalling pathway" (S29 Table).

Similarly, the significantly upregulated pathways in spring metacarpal tissue included

"ribosome," "oxidative phosphorylation," "biosynthesis of secondary metabolites,"

"microbial metabolism in diverse environments," and "biosynthesis of antibiotics"

(S29 Table).

KEGG pathway analysis using DEGs of FP-S vs FP-W comparison indicated

that 12 and three pathways were significantly downregulated and upregulated in

spring perirenal tissue, respectively (S30 Table). The downregulated pathways were

mainly associated with environmental information processing and signal

transduction, such as "cAMP signalling pathway," "cGMP-PKG signalling pathway,"

"Rap1 signalling pathway," "Hippo signalling pathway," and "MAPK signalling

pathway" (S30 Table). However, pathways such as "ribosome," "oxidative

phosphorylation," and "ribosome biogenesis in eukaryotes" (S30 Table) were

significantly upregulated in spring perirenal tissue.

KEGG pathway analysis for FS-S versus FS-W DEGs revealed two and nine

significantly upregulated and downregulated KEGG pathways in spring prescapular

tissue, respectively (S31 Table). Pathway analysis showed a significant upregulation

of two pathways, "ribosome" and "complement and coagulation cascades". The

downregulated pathways were mainly associated with environmental information

processing and signal transduction: "Hippo signalling pathway–fly", "cAMP signalling

pathway", "ErbB signalling pathway", "MAPK signalling pathway", "Hippo signalling

pathway" and "FoxO signalling pathway".

KEGG pathways associated with seasonal comparisons. Pathway analysis in

Even reindeer revealed three KEGG pathways significantly upregulated in the

metacarpal adipose tissues collected during spring ("ribosome", "spliceosome," and

"ribosome biogenesis in eukaryotes"), whereas no significantly downregulated

pathways were found. Analysis of DEGs from perirenal adipose tissue did not reveal

any KEGG pathways. Pathways such as "complement and coagulation cascades"

and "cytokine-cytokine receptor interaction" were significantly upregulated during

spring in prescapular adipose tissue, while no KEGG pathways were associated with

downregulated genes.

KEGG pathways associated with location comparisons. DEGs from the

comparison of the metacarpal adipose tissue between Even and Finnish reindeer

did not reveal any KEGG pathways associated with upregulated genes, whereas two

KEGG pathways, "oxidative phosphorylation" and "ribosome," were associated with

genes downregulated in Even reindeer in spring (S32 Table) and a similar

comparison from winter sampling revealed seven KEGG pathways, such as

"ribosome biogenesis in eukaryotes", "TNF signalling pathway", "IL-17 signalling

pathway," "NF-kappa B signalling pathway," and "cytokine-cytokine receptor

interaction" associated with downregulated genes (S33 Table).

Upregulated genes in the perirenal adipose tissue of Even reindeer from spring

sampling were associated with one KEGG pathway "ribosome," while downregulated

DEGs were not associated with any of the pathways. Similar comparison from winter

sampling revealed six and 13 KEGG pathways associated with upregulated and

downregulated genes, respectively, in Even reindeer (S34 Table).

While no KEGG pathways were associated with upregulated genes in

prescapular adipose tissues from spring samples, two ("ribosome" and "complement

and coagulation cascades") were linked to the downregulated genes. Similar

comparison in winter samples revealed 14 and one KEGG pathways associated with

upregulated and downregulated genes, respectively, in Even reindeer (S35 Table).

KEGG pathways associated withgender comparisons. KEGG pathway analysis

revealed no statistically significant enriched KEGG pathways between EM-F and

EM-M. "Ribosome" was the only KEGG pathway associated with DEGs from the

perirenal and prescapular adipose tissues. Interestingly, in both tissues, the

"ribosome" pathway was downregulated in female reindeer.

Physiological analyses

Immunoblotting of UCP1 and COX4. Immunoblotting on UCP1, a protein specific

to thermogenic adipocytes, was conducted on a cross-section of different adipose

tissues. Immunoreactivity at 32 kDa molecular weight characteristic of UCP1 was

evident, although in small amounts, in the total proteins of prescapular and perirenal

adipose tissues of both Even and Finnish reindeer (Fig 4). In general, prescapular

adipose tissues appeared to have slightly higher UCP1 expression than perirenal

adipose tissues. In Finnish reindeer, the relative expression of UCP1 was

significantly higher in winter compared with spring in both prescapular (p = 0.0016*)

and perirenal (p = 0.032*) adipose tissues (Fig 4c, f). The Even reindeer exhibited a

similar albeit not statistically significant trend of higher UCP1 expression in winter

compared with spring (Fig 4c, f).

The expression of COX4, an enzyme central to oxidative phosphorylation, was

undetectable from the adipose tissue total proteins. However, we analyzed its

expression in other metabolically active tissues—the liver and muscle (see Materials

and methods). The relative expression level of COX4 in the liver samples was

significantly higher in Finnish reindeer in winter (p = 0.008**) in comparison to spring

(Fig5). There were no statistically significant differences in muscle COX4 levels

between seasons.

Blood metabolites. Plasma leptin levels were at a similar range in Even and Finnish

reindeer in spring, but under the detection limit (1.56 ng/ml) in both male and female

Even reindeer in winter (Fig 6a). Plasma insulin levels of reindeer were at similar

level in both seasons and regions (Fig 6b). Plasma HSL was significantly higher in

winter in Finnish reindeer compared with spring (p = 0.016*), but there were no

significant differences in the Even reindeer between seasons (Fig 6c).

Serum glucose, creatinine and triglyceride concentrations were at a similar

range in both Even and Finnish reindeer in both seasons (Figures 6d, e, g). There

was a trend of higher serum urea (Fig 6f) in winter samples compared to spring

samples within the deer of both regions, albeit not statistically significant.

Discussion

Adipose tissues are vital for animals living in cold environments, promoting

adaptation by temperature regulation, energy homeostasis, regulation of fat

deposition, and metabolism [5,6,10,11,36–40]. Here, we have investigated gene

expression profiles of adipose tissues from three different anatomical depots in

Finnish and Even reindeer during two seasonal time points. To the best of our

knowledge, this is the first transcriptome study of adipose tissues in reindeer. Our

results indicated a clear difference in gene expression profiles in metacarpal adipose

tissue compared to perirenal and prescapular adipose tissues. We found that during

the less optimal circumstances in early spring, mainly characterized by

undernutrition, genes associated with the immune system were upregulated in

perirenal and prescapular adipose tissues, while genes involved in energy

metabolism were upregulated in metacarpal tissue. Interestingly, developmental,

growth, and adipogenesis processes were downregulated in all three tissues during

spring.

RNA-Seq is an efficient method to screen new genes, transcripts, gene

expression, and differentially expressed genes in various organisms, tissues, and

cells [41–43]. In this study, we identified a total of 16,362 expressed genes in focal

adipose tissues, which appeared to cover approximately 60% of the list of genes

available for the reindeer reference genome [3]. The present results revealed an

adequate number of expressed genes in reindeer adipose tissues, which were then

utilized in subsequent gene expression analysis to investigate genes associated with

seasonal (early spring *vs.* early winter), location (Even reindeer *vs.* Finnish reindeer),

and gender (male *vs.* female) differences.

In this study, highly abundant genes were associated with fat and lipid

metabolism, thermogenesis, and energy homeostasis that are critical for the survival

of reindeer during seasonal fluctuations. Several of the abundant genes involved in

fat and lipid metabolism, such as *FABP4*, *FABP5*, *MT-CYB* and *ADIPOQ* (Table

S36), were highly expressed in all adipose tissues in both Finnish and Even reindeer.

Fatty acid binding protein 4 (*FABP4*) was the most highly expressed gene in all

adipose tissues in both Finnish and Even reindeer. Three of the top expressed

genes, including *FABP4*, *FABP5,* and *ADIPOQ,* were also found to be highly

expressed in a previous adipose transcriptome profiling study in two fat-tailed sheep

breeds [22]. Moreover, *FABP4* and *FABP5* were also highly expressed in the

perirenal tissue of sheep [44]. Fatty acid binding proteins (FABPs) 4 and 5, encode

the fatty acid binding protein found in adipocytes and epidermal cells, respectively.

A previous study showed that *FABP4* and *FABP5* play an important role in

thermogenesis during cold exposure and starvation [45]. Moreover, previous studies

in cattle have suggested that *FABP4* plays a crucial role in fat deposition, fatty acid

transport, catabolism, and metabolism [22,46,47]. Adiponectin (*ADIPOQ*), secreted

by adipocytes and exclusively expressed in adipose tissues, plays an important role

in modulating the regulation of fatty acid oxidation, glucose levels, and insulin

sensitivity [22,48,49]. The top expressed genes associated with fat metabolism

observed in the present study may play a vital role in energy homeostasis,

thermoregulation, and promoting adaptation of reindeer to the challenging

environment. The blood lipid, glucose, and insulin levels of reindeer support the view

of homeostasis despite challenging conditions.

Tissue-wise, the gene expression profiles of metacarpal adipose tissue were

remarkably different from those of prescapular and perirenal adipose tissues. The

highest number of tissue-specific genes were found in metacarpal adipose tissue

(Fig 1). Similarly, the PCA plot based on all expressed genes revealed a cluster for

metacarpal adipose tissue, which was distinct from the samples representing other

tissues (Fig 2). We found that several genes from the homeobox (HOX) family of

proteins were uniquely expressed in metacarpal adipose tissue. These genes are

known to play important roles in the differentiation of adipocytes [50]. Furthermore,

several genes associated with cytokines and immune response appeared to be

downregulated in the metacarpal tissue during spring (Table S5 and S8). By

contrast, DEGs associated with cytokines and immune response were upregulated

in the perirenal and prescapular tissues during spring (Tables S6, S7, S9, and S10).

Hence, these results indicate the unique biological functions of the metacarpal tissue

compared to the other two tissues.

The distinctiveness of the metacarpal adipose tissue may be due to its local

niche functions in bone marrow, although it also contributes to systemic metabolism

[14,16]. Bone marrow adipose tissue has many properties in common with white

adipose tissue, but it is an adipose type of its own that is currently being actively

studied [14,16]. In addition to its role as a local energy reservoir and its contribution

to haematopoiesis and osteogenesis [15], metacarpal adipose tissue also secretes

a variety of hormones and proteins, such as adiponectin and leptin, which have an

important function in the regulation of energy metabolism [16]. The metabolic profile

of metacarpal adipose tissue is composed of both white and brown fat, indicating its

plasticity in performing different functions [16].

We observed more seasonal differences in the number of DEGs in the adipose

tissues of Finnish reindeer (n = 1229) compared to Even reindeer (n = 386, also see

Table 2). The Finnish reindeer specifically exhibited several significant DEGs

associated with ATPase and ATP synthase, whereas no ATP-related genes were

detected in Even reindeer. The significant DEGs associated with ATPase and ATP

synthase identified in Finnish reindeer include *ATP12A, ATP5L, ATP5O, ATP6AP1L*

and *ATP8* in metacarpal, *ATP1A2, ATP1B2, ATP6V0C, ATP7A* and *ATP8* in

perirenal, and *ATP1A2, ATP1B2, ATP7A, ATP8B1, ATP8B3* and *ATP8* in

prescapular adipose tissue. A previous study on zebrafish (*Danio rerio*) reported that

feeding altered the expression of ATP-related genes [51]. The differences between

Even and Finnish reindeer could be due to the influence of management (extra

feeding in Finland *versus* no additional feeding in Yakutia), vegetation (relatively

sparser in Yakutia), and temperature (relatively warmer in Finland). A further

possible explanation for the high expression of ATP genes in Finnish reindeer is that

they were fed concentrates with a high protein content (~10% protein content)

compared to their natural winter food, lichens (2–3% protein content).

In terms of seasonal comparison, we observed that the gene expression profiles

of metacarpal tissue were mainly enriched for energy metabolism instead of their

typical role in immune systems. Adipose tissue has been previously reported to play

a role in immune and inflammatory systems [52,53]. Earlier studies reported that

adipocytes of both peripheral and bone marrow fat secrete a variety of hormones

and proteins, such as pro-inflammatory and anti-inflammatory cytokines [54–56].

However, in the present Even reindeer metacarpal tissue, the downregulated genes

in spring included genes associated with cytokines, such as *CCL19*, *IGDCC4*,

IL17RB, *JCHAIN*, *LECT1* and *IGLV1-51*. Moreover, several of the downregulated

DEGs in spring in Finnish and Even reindeer metacarpal adipose tissues revealed

genes associated with immune system. On the other hand, the upregulated DEGs

in spring in Finnish and Even reindeer metacarpal adipose tissue revealed genes

associated with lipid and energy metabolism, supporting the view that bone marrow

fat acts as a source of energy when reindeer are in poor condition [18,19]. In

reindeer, the proportions of unsaturated fatty acids, oleic, and linoleic acid are

significantly decreased in metatarsal bone marrow fat in poor conditions in spring

[19]. This may be related to their use for oxidation or other, synthetic processes. For

instance, angiopoietin-like protein 8 (*ANGPTL8*) and angiopoietin-like protein 1

(*ANGPTL1*) were among the upregulated genes detected in the metacarpal tissue

of Finnish and Even reindeer, respectively, in spring. *ANGPTL8* is a member of the

angiopoietin-like protein (*ANGPTL*) family involved in the metabolic transition from

fasting to re-feeding and plays a key role in lipid metabolism [57–59]. Depending on

the location in the body, adipose tissues differ in terms of cellular composition,

quantity, and proportion of adipocytes, and their capacity to produce adipocytokines

[9]. During starvation, adipose tissue often limits the cytokine levels to reduce the

consumption of resource/energy by the immune systems to actively decrease energy

usage; subsequently, the activity of immune cells is limited [60–62]. Reindeer in both

locations, and particularly Even reindeer, are in their poorest nutritional condition

during spring. In this study, all cytokine genes identified in the metacarpal tissue

were downregulated in spring. This might be due to the metacarpal adipose tissue

being exclusively involved in thermogenesis and energy metabolism by suppressing

the function of the immune system. Hence, when these reindeer experience the

worst conditions during spring, the stored fat depots from metacarpal adipose tissues

may be exclusively reserved for energy usage.

As revealed by the top downregulated genes in the perirenal and prescapular

adipose tissues of Finnish reindeer and in all three adipose tissues of Even reindeer,

biological processes associated with development, cell growth, and organogenesis

were repressed during spring. The downregulation of genes involved in

organogenesis, cell growth, and development in spring may indicate that the animals

give less priority for growth- and development-related processes during extreme

conditions; thus, the animals spend no extra energy for growth related processes.

As mentioned above, during extreme conditions, the animals reduce several

metabolic activities to increase the efficiency of energy usage.

Furthermore, gene expression analysis of gender differences in even reindeer

adipose tissues indicated that genes associated with fatty acid metabolism and male

sterility were upregulated in female and male reindeer, respectively. The upregulated

genes associated with fatty acid metabolism in females include *ELOVL7* and *FABP6*

in metacarpal, *ABCG1*, *ACSL6*, *PLCD3,* and *PLA2G5* in perirenal tissue, and *ACSL6*

and *PLA2R1* in prescapular tissue. This result is consistent with previous studies in

humans and mice [63,64], which suggested that females show higher levels of fat

deposition than males. By contrast, the upregulated genes in male reindeer revealed

10 shared genes among the three tissues, such as *UBA1*, *TXLNG*, *PRDM9*,

EIF2S3X, *NRL*, *DDX3Y*, *KDM6A*, *ZRSR2*, *USP9X* and *ZFX*. Six of these genes,

PRDM9 [65,66], *UBA1* [67], *EIF2S3X* [68,69]*, DDX3Y* [70–72], *ZRSR2* [57], and

USP9X [74], have been shown to be associated with male sterility. Moreover, it

should be noted that the reindeer reference genome lacked gene annotations for the

Y chromosome, and that many of the Y chromosome-specific RNA-Seq reads may

have aligned to the paralog genes of the X chromosome.

We also analyzed adipose *UCP1* and *COX4* protein levels as indicators of the

potential thermogenesis and metabolic state of the reindeer. We anticipated that

adipose tissues of the adult reindeer are white adipose tissue but may potentially

have some brown adipose tissue characteristics, considering the extreme long-term

exposure of the reindeer to cold, especially in Siberia. There is no previous evidence

of BAT or 'browning' of adipose tissues in adult reindeer, but it is well established

that new-born reindeer have active BAT at birth and during their first month of life

[10,11].

Our results show the presence of *UCP1* protein in two different adipose tissues of

adult reindeer, prescapular, and perirenal depots (Fig 4). Relative *UCP1* expression

was significantly higher in winter compared with spring in Finnish reindeer in both

prescapular and perirenal adipose tissues, likely reflecting colder weather conditions

in winter. A similar trend was observed in the Even reindeer. Due to the low amount

of protein, it is likely that *UCP1* does not have major thermogenic relevance.

However, the findings are still interesting, as they show that *UCP1* was present in

the white adipose tissues of adult reindeer. They also refer to plasticity of adipose

tissue, that is, the potential to adjust its functions according to prevailing conditions

[75]. In general, reindeer can cope with very low ambient temperatures in winter (-

30°C) without increasing their heat production [76], and thus non-shivering

thermogenesis is usually not necessary. This is mainly due to the good insulation

capacity of the winter coat of reindeer, which effectively prevents heat loss. Liver

COX4 expression was significantly higher in Finnish reindeer in winter as compared

to spring, and a similar trend was apparent in Even reindeer (Fig 5). This indicates

an increase of oxidative phosphorylation and ATP synthesis in the liver to match the

increased energy demands caused by the colder season.

Plasma leptin and insulin levels were low (Fig 6) and agree with earlier findings

in reindeer [5]. Leptin is a hormone secreted by adipose tissues and plays a role in

the regulation of body weight [77]. The low leptin levels suggest that the adipose

tissues of reindeer were small and that the animals were striving to preserve their

adipose tissues. Low leptin also agrees with adipose transcriptomics results referring

to fat mobilization. Plasma hormone-sensitive lipase (HSL) was significantly higher

in Finnish reindeer in the early winter than in the spring group, suggesting the

mobilization of storage lipids to accommodate increased energy expenditure of male

reindeer related to the breeding season.

Serum glucose, triglyceride, creatinine, and urea concentrations were similar

between seasons (Fig 6), indicating that the harsh winter season is relatively well

tolerated by the reindeer without severe muscle catabolism, which would be

indicated by changes in the blood parameters.

Conclusions

Collectively, our mRNA-Seq data uncovered variations in the transcriptome

profiles of three adipose tissues in relation to seasonality, location, and gender

differences. In general, our study showed that highly expressed genes in adipose

tissues were associated with fat and lipid metabolism and thermal and energy

homeostasis, promoting the adaptation of reindeer to challenging Northern Eurasian

environments. Further, our results indicated a distinct gene expression profile in

metacarpal adipose tissue compared to perirenal and prescapular adipose tissues.

Metacarpal Adipose tissue appeared to have a greater role in metabolic activities

compared to the other two tissues especially during spring when the animals are

experiencing the worst nutritional conditions. Moreover, reindeer from Finland and

Yakutia displayed different gene expression profiles, in part owing to climatic and

management differences. Thermogenic UCP1 protein was present in adipose

tissues of both Even and Finnish reindeer, although in low amounts, showing that

the reindeer have an option for extra heat production and thermal and energy

homeostasis if needed. Taken together, the results and resources from this study

will be useful for elucidating the genetics and physiology of adipose tissue for

adaption to northern Eurasian conditions.

Materials and Methods

Sample collection for transcriptome analysis

This study includes RNA-Seq of 56 tissue samples from 19 reindeer individuals

(three adult females and 16 adult males) that were randomly collected at slaughter

from two different geographical regions (Inari, northern Finland and Eveno-Bytantay,

Sakha, Yakutia, the Russian Federation) at two seasonal time points: winter

(November–December) and spring (April) (Table 1 and S1 Table). Perirenal samples

were taken from the adipose tissue around the kidneys, prescapular samples from

the adipose tissue located beneath the cervical muscles in front of the scapula, and

metacarpal samples from the bone marrow in the diaphysis of the metacarpal bone

(left front leg). For convenience, throughout the text, the sample groups are

abbreviated using reindeer location (F, E), tissue type (P, S, M), and seasonal time

points (S, W). For example, FM-S represents the metacarpal tissue of Finnish

reindeer collected during spring. The samples were stored in RNAlater® Solution

(Ambion/QIAGEN, Valencia, CA, USA). It should be noted that three male reindeer

(spring) from Yakutia were castrated, whereas those from Finland (n = 10) and

autumn Yakutian males (n = 3) were uncastrated. The animals grazed on natural

pastures throughout the year before the sampling. However, the Finnish reindeer

were fed concentrates (Poroherkku, Raisio, Finland) for 2–8 weeks in February and

March 2016 and kept in feeding pens prior to slaughter. The animals were exposed

to seasonal ambient temperatures and photoperiod. The mean daily atemperature

in Inari, Finland varied between -16.1℃ and 5.2℃ before the sampling in winter (14

hours light, 10 hours dark) and between -13.2℃ and 4℃ before the sampling in

spring (16 hours light, 8 hours dark). In northern Sakha, the daily temperature varied

between -13℃ and -24℃ during the winter sampling (6.5 hours daylight, 17.5 hours

dark) and between -9℃ and - 0℃ during the spring sampling (14 hours of daylight,

10 hours dark). Serum and plasma samples were also collected from Sodankylä,

Finland in the spring (15 hours light, 9 hours dark), where the mean daily

temperatures varied between -11.4℃ and 3.7℃. All protocols and sample collections

were performed in accordance with the legislations approved by the Russian

authorization board (FS/UVN 03/163733/07.04.2016) and the Animal Experiment

Board in Finland (ESAVI/7034/04.10.07.2015).

Table 1. Summary of adipose tissue samples used for RNA-seq and physiological

studies (the latter in brackets).Blood samples were collected from all the animals and

two additional male Finnish reindeer in spring.

	Finnish reindeer		Even reindeer		
	Spring male	Winter male	Spring male*	Winter	
				male	female
Metacarpal	5	5	3	3	3
Perirenal	4 (4-5)	5 (5)	3 (3)	3 (3)	3
Prescapular	5 (5)	5 (5)	3 (3)	3 (3)	3

782 *Castrated males (Spring samples)

Sample collection for physiological analysis

Blood samples were taken before slaughter by a jugular venipuncture into

vacuum serum and EDTA K3 tubes. The blood samples were centrifuged, and the

separated serum and plasma were stored at -80°C until analysis. A total of 18 males,

Finnish (n = 12) and Even (n = 6) reindeer, were examined, of which 16 were

included in the RNA-seq analysis (see S1 Table). The aforementioned adipose

tissues as well as additional liver and muscle (*M. gluteobiceps femoris*) samples

were stored in RNAlater® solution and then used for immunoblotting analysis. We

used RNAlater for preservation instead of liquid nitrogen due to the long storage of

the samples in field conditions. The use of RNAlater was validated by testing both

RNAlater and liquid nitrogen-preserved samples from western blotting (data not

shown here).

RNA extraction, library preparation, and sequencing

RNA extraction, library preparation, and sequencing were performed at The

Finnish Functional Genomic Center (FFGC), Turku, Finland. Total RNA was

extracted from adipose tissues (ca <30mg/sample) using the Qiagen AllPrep

DNA/RNA/miRNA kit according to the manufacturer's protocol. The quality of the

obtained RNA was ensured with an Agilent Bioanalyzer 2100 (Agilent Technologies,

Waldbronn, Germany), and the concentration of each sample was measured with a

Nanodrop ND-2000 (Thermo Scientific; Wilmington, USA) and a Qbit(R)

Fluorometric Quantification, Life Technologies. All samples revealed an RNA

integrity number (RIN) above 7.5.

Library preparation was done according to Illumina TruSeq® Stranded mRNA

Sample Preparation Guide (part #15031047). Unique Illumina TruSeq indexing

adapters were ligated to each sample to pool several samples later in one flow cell

lane. Library quality was inferred with an Advanced Analytical Fragment Analyzer

and concentration with a Qubit fluorometer, and only good-quality libraries were

sequenced.

The samples were normalized and pooled for automated cluster preparation,

which was carried out with Illumina cBot station. Libraries prepared for sample

YR1_SCAP_322D and FR12_SCAP_402D (S1 Table) were pooled together and run

in one lane to generate a deep sequence to detect long non-coding RNAs (lncRNAS)

for a future study. The remaining 61 libraries were combined in one pool and run on

seven lanes of an Illumina HiSeq 3000 platform. Paired-end sequencing with 2 × 75

bp read length was used with a 8 + 8 bp dual index run. Two samples

(FR9_SCAP_369D and FR13_PREN_425C) (S1 Table) suffering fed pooling error

and low amounts of reads and were therefore resequenced in an extra lane. Base

calling and adapter trimming were performed using Illumina's standard bcl2fastq2

software.

Bioinformatics analyses

The overall quality of the raw RNA-seq reads in fastq and aligned reads in BAM

format were assessed using FastQC software v0.11.7 [78]. FastQC reports were

summarized using MultiQC v1.7 [79].High quality RNA-seq reads for each sample

were mapped against the reindeer draft assembly [3] using Spliced Transcripts

Alignment to a Reference (STAR) (version 2.6.0a) [80] with default parameters. We

next generated read counts from the aligned files using the featureCounts software

(version 1.6.1) from the Subread package [81] to assign reads to genes. The GTF-

format annotation file associated with the reindeer draft assembly was used for gene

coordinate information.

To examine the shared and uniquely expressed genes across the three adipose

tissues, we used the cpm function from the edgeR library [82] to generate count-per-

million (CPM) values; lowly expressed transcripts with a CPM < 0.5 were discarded.

Adipose transcriptomes are affected by the type of adipose as well as by the sex

of the animal [31]. We also hypothesized that there could be differences in seasonal

gene expression profiles due to changes in ambient temperature and other climatic

factors and subsequent changes in body condition. Hence, we conducted differential

gene expression analysis between the spring and winter sampling for each tissue

and region using only male reindeer samples. In addition, we compared gene

expression in samples collected from female and male Even reindeer in winter.

Furthermore, to explore regional (and population) differences in gene expression,

we compared expression in each tissue between E and F male reindeer. In our study,

the analyzed group of animals for each tissue included at least three animals from

each geographical region and season (Table 1 and S1 Table)

Raw read counts were processed using the R Bioconductor package DESeq2

[83] to perform differential gene expression and related quality control analysis. Prior

to running DESeq2, lowly expressed (rowSums < 1) genes were discarded. Raw

gene expression counts were normalized for differences in library size and

sequencing depth using DESeq2, to enable gene expression comparisons across

samples. We performed principal component analysis (PCA) to assess sample

similarity using the variance stabilizing transformation (VST) method. In this study,

we used fold-change and false discovery rate (FDR) filtering criteria to identify

significantly differentially expressed genes (DEGs). We set absolute value of log2-

fold change (LFC) to be greater than or equal to 1.5 ($|log2FoldChange| > 1.5$) and

an adjusted p-value of 0.05 (*padj < 0.05*) to screen for significant DEGs. The

Benjamini-Hochberg FDR method was used to calculate adjusted p-values.

To gain insight into the biological functions and relevance of the identified DEGs,

a functional enrichment analysis was conducted using AgriGO v2.0 [84]. In the

AgriGO analysis toolkit, to detect the significantly enriched GO terms, default

parameters were used in the "Advanced options": Fisher as the statistical test

method, Yekutieli for multiple test correction at a significance level threshold 0.05

(FDR < 0.05), and minimum number of mapping entries 5. In this analysis, the GO

annotation file from the *de novo* assembled reindeer genome [3] was used as a

background reference. Furthermore, to explore the biological pathways associated

with the DEGs, we performed Kyoto Encyclopaedia of Genes and Genomes (KEGG)

pathway analysis using the GAGE [85] Bioconductor package. The significantly

enriched pathways were identified based on the q-values obtained from a Fisher's

exact test (*q-value < 0.1*).

Physiological analyses

Immunoblotting analysis of proteins. Adipose tissue samples were homogenized and dissolved in lysis buffer (25 mM Tris [pH 7.4], 0.1 mM EDTA, 1 mM DTT, 15µl/ml protease inhibitor cocktail (Sigma, St Luis, MO, USA) to extract total protein content. Insoluble material was removed from the extracts by centrifugation (13,000 g, 10 min, +4˚C). Mitochondrial proteins were extracted from reindeer muscle, liver, reindeer calf prescapular brown adipose tissue (BAT), and mouse BAT samples as described previously [86]. Both total and mitochondrial protein concentrations were determined using the Bradford method (Bio-Rad protein assay, Bio-Rad Laboratories GmbH, München, Germany). Protein extract volumes equivalent to 50–75 µg of total protein were concentrated into smaller volumes by lyophilizing the samples with a Savant Speed Vac Plus SC210A centrifugal evaporator (Thermo Fisher Scientific, Rockford, USA) in cooled conditions for 1 hour. Due to the low number of mitochondria in adipose tissue, total protein extractions were used for the immunoblotting of UCP1.

The proteins were separated electrophoretically using a 4–12% gradient gel and transferred to a nitrocellulose membrane (Bio-Rad, Trans-Blot® Transfer Medium, Pure Nitrocellulose Membrane [0.2 µm], Bio-Rad Laboratories, Hercules, CA, USA). The membranes were incubated overnight with UCP1 antibody and loading control protein alpha tubulin antibody for the total protein adipose tissue samples (1:1.000 UCP1 Polyclonal Antibody, cat no. PA1-24894, 1:500 alpha Tubulin Polyclonal Antibody, cat no. PA5-16891, Invitrogen, Thermo Fisher Scientific, Rockford, USA). Membranes with liver and muscle mitochondrial protein samples were incubated

overnight with COX4 antibody (1:1.000 COX4 Polyclonal Antibody, cat no. PA5-

17511, Thermo Fisher Scientific, Rockford, USA) and loading control protein ATP5A

antibody (1:1.000 anti-ATP5A antibody, cat no. ab151229, Abcam, Cambridge, UK).

After the primary antibody treatments, the membranes were incubated with a

secondary antibody (1:25.000, Goat anti-Rabbit IgG (H+L) horseradish peroxidase

conjugate, cat no. 31460, Invitrogen, Thermo Fisher Scientific, Rockford, USA) for 1

hour. Chemiluminescence for UCP1 and COX4 was detected with SuperSignal West

Femto Maximum Sensitivity Substrate (cat no. 34095, Thermo Fisher Scientific,

Rockford, USA) according to the manufacturer's instructions. Blots were visualized

with Odyssey® Fc imaging system (LI-COR Biosciences, Ltd, Cambridge, UK).

Positive immunoreactivity for UCP1 with mouse BAT mitochondria and with

prescapular BAT mitochondria from newborn reindeer were used as reference

samples. Results were normalized with the loading control optical density for alpha

tubulin and ATP5A for UCP1 and COX4, respectively.

Blood metabolites. Plasma leptin concentration was assayed using a multispecies

leptin RIA kit (Cat#XL-85K, Millipore, Billerica, Massachusetts, USA). The validity

test for reindeer showed a linear correlation between the label and sample

concentration. A sensitive bovine, ovine, rat, and mouse insulin RIA kit (Cat#SRI-

13K, Millipore, Billerica, Massachusetts, USA) was used to measure plasma insulin.

Plasma hormone-sensitive lipase levels (HSL) were estimated by bovine hormone

sensitive ELISA Kit (Cat#MBS033124, MyBioSource, San Diego, CA, USA). Serum

glucose, triglyceride, creatinine, and urea concentrations were determined with

enzymatic colorimetric analyses in NordLab, Oulu, Finland.

Statistical analyses for immunoblotting and blood metabolite analyses.

Statistical analysis for the multiple comparisons was performed with an independent-

samples Kruskal-Wallis test followed by an independent samples Mann-Whitney U-

test. Significance values were adjusted with the Bonferroni correction for multiple

tests. Statistical analyses were performed using the IBM SPSS Statistics 21 Data

Editor software (IBM, Armonk, NY, USA). P-values below 0.05 were considered

statistically significant. The results of the relative peptide expressions are presented

as the mean ± SD.

Figure captions

Fig 1. Venn diagram showing overlap of expressed genes (CPM ≥ 0.5 for at

least two samples) among tissues in each region and season. Shared and

uniquely expressed genes across tissues (M, P, and S) in (A) Finnish reindeer in the

summer, (B) Finnish reindeer in the winter, (C) Even reindeer in the summer, and

(D) Even reindeer in the winter.

Fig 2. PCA plots of the analyzed samples based on expression profiles,

with dot colours indicating tissue and region (see legend).

Fig 3. **Distribution of uniquely expressed genes in the metacarpal adipose**

tissue of Finnish (FM-W, FM-S) and Even (EM-W, EM-S) reindeer.

Fig 4. The western blots and relative expression of UCP1 from (a–c) reindeer

prescapular and (d–f) perirenal adipose tissue total proteins. The upper blots

on the left side show UCP1 content in prescapular adipose tissue (a) in spring in

Even reindeer (n = 3) and Finnish reindeer (n = 5), (b) in winter in Even reindeer (n

= 3) and Finnish reindeer (n = 5), and the lower graph shows (c) their relative

expressions. The upper blots on the right side show UCP1 content in perirenal

adipose tissue (d) in spring in Even reindeer (n = 3) and Finnish reindeer (n = 4), (e)

in spring in Even reindeer (n = 3) and Finnish reindeer (n = 4), and the lower graph

shows (f) their relative expressions. Samples contained 50 µg of total protein per

lane, except the reindeer BAT (brown adipose tissue; BAT and BATr) samples with

5 µg and mouse BAT (BATm) with 1 µg of protein. Alpha tubulin was used as a

loading control. The relative expression of UCP1 was normalized to alpha tubulin

and presented as mean arbitrary units (a.u.) ± SD c,f). Significant differences

between seasons are indicated with a bar and an asterisk (*P < 0.05).

Fig 5. The western blots and relative expression of *COX4* from (a–c) reindeer

muscle (*M. gluteobiceps femoris*) and (d–f) liver mitochondrial proteins. The

upper blots on the left side show COX4 content in muscle (a) in spring in Even

reindeer (n = 3) and Finnish reindeer (n = 5), (b) in winter in Even reindeer (n = 3)

and Finnish reindeer (n = 5), and the lower graph shows (c) their relative

expressions. The upper blots on the right side show *COX4* in liver (d) in spring in

Even reindeer (n = 3) and Finnish reindeer (n = 5), e) in winter in Even reindeer (n =

3) and Finnish reindeer (n = 5), and the lower graph shows (f) their relative

expressions. Samples contained 10 µg of mitochondrial protein per lane, except the

reindeer BAT (brown adipose tissue) sample with 1 µg of mitochondrial protein.

Mitochondrial ATP synthase (ATP5A) was used as a loading control. The relative

expression of *COX4* was normalized to *ATP5A* and presented as mean arbitrary

units (a.u.) ± SD c,f). Significant difference between seasons is indicated with a bar

and an asterisk (**P ≤ 0.01).

Fig 6. Plasma hormone and serum metabolite levels in Finnish male reindeer

in winter and spring (n = 7), in male Even reindeer in winter and spring (n=3),

and in Even female (F) reindeer in winter (n = 3). Plasma hormones (a) leptin, (b)

insulin, (c) hormone sensitive lipase (HSL) and serum metabolites (d) glucose, (e)

creatinine, (f) urea, and (g) triglyceride levels are presented as mean ± SD.

Significant difference between seasons is indicated with a bar and an asterisk (**P

≤ 0.01).

FIG 1

Fig 2

Fig 3.

Supporting information

S1 Fig. PCA plot based on the region for each tissue. (A) metacarpal, (B) perirenal,

and (C) prescapular.

S2 Fig. Heatmap plot of the top 25 genes with the highest genetic variance across

all samples.

S3 Fig. The number of shared and uniquely significant DEGs between three adipose

tissues in Finnish reindeer due to seasonal differences. Significant DEGs detected

in Finnish reindeer for three adipose tissues due to seasonal change: FM-S *vs.* FM-

W, FP-S *vs.* FP-W, and FS-S *vs.* FS-W.

S4 Fig. Volcano plot of differentially expressed genes between spring and winter for

metacarpal adipose tissue in Finnish reindeer (FM-S *vs.* FM-W).

S5 Fig. Volcano plot of differentially expressed genes between spring and winter for

perirenal adipose tissue in Finnish reindeer (FP-S *vs.* FP-W).

S6 Fig. Volcano plot of differentially expressed genes between spring and winter for

prescapular adipose tissue in Finnish reindeer (FS-S *vs.* FS-W).

S7 Fig. The number of shared and unique significant DEGs between three adipose

tissues in Even reindeer due to seasonal differences. Significant DEGs detected in

Even reindeer for three adipose tissues due to seasonal change (EM-S *vs.* EM-W,

EP-S *vs.* EP-W and ES-S *vs.* ES-W).

S8 Fig. Volcano plot of differentially expressed genes between early spring and early

winter for metacarpal adipose tissue in Even reindeer (EM-S *vs.* EM-W).

S9 Fig. Volcano plot of differentially expressed genes between early spring and early

winter for metacarpal adipose tissue in Even reindeer (EM-S *vs.* EM-W).

S10 Fig. Volcano plot of differentially expressed genes between early spring and

early winter for metacarpal adipose tissue in Even reindeer (EM-S *vs.* EM-W).

S11 Fig. The number of shared and unique significant DEGs in the three adipose

tissues detected between Even reindeer and Finnish reindeer due to regional

differences in early spring. Significant DEGs detected in three adipose tissues due

to regional differences in early spring: EM-S *vs.* EM-S, EP-S *vs.* EP-S and ES-S *vs.*

ES-S.

S12 Fig. The number of shared and unique significant DEGs in the three adipose

tissues detected between Even reindeer and Finnish reindeer due to regional

differences in early winter. Significant DEGs detected in three adipose tissues due

to regional differences in early winter: EM-W *vs.* EM-W, EP-W *vs.* EP-W and ES-W

vs. ES-W.

1011

S1 Table. Statistics of clean data.

S2 Table. STAR mapping statistics.

S3 Table. Summary of expressed genes in each tissue (sheet). List of expressed

genes in Finnish reindeer metacarpal adipose tissue in spring (FM-S) (sheet 2). List

of expressed genes in Finnish reindeer perirenal adipose tissue in spring (FP-S)

(sheet 3). List of expressed genes in Finnish reindeer prescapular adipose tissue in

spring (FS-S) (sheet 4). List of expressed genes in Finnish reindeer metacarpal

adipose tissue in winter (FM-W) (sheet 5). List of expressed genes in Finnish

reindeer perirenal adipose tissue in winter (FP-W) (sheet 6). List of expressed genes

in Finnish reindeer prescapular adipose tissue in winter (FS-W) (sheet 7). List of

expressed genes in Even reindeer metacarpal adipose tissue in spring (EM-S)

(sheet 8). List of expressed genes in Even reindeer perirenal adipose tissue in spring

(EP-S) (sheet 9). List of expressed genes in Even reindeer prescapular adipose

tissue in spring (ES-S) (sheet 10). List of expressed genes in Even reindeer

metacarpal adipose tissue in winter (EM-W) (sheet 11). List of expressed genes in

Even reindeer perirenal adipose tissue in winter (EP-W) (sheet 12). List of expressed

genes in Even reindeer prescapular adipose tissue in winter (ES-W) (sheet 13).

S4 Table. Uniquely expressed genes in metacarpal adipose tissue shared by Finnish

and Even reindeer in both seasons.

S5 Table. Significantly differentially expressed genes between spring and winter in

Finnish reindeer metacarpal tissue (FM-S *vs.* FM-W).

S6 Table. Significantly differentially expressed genes between spring and winter in

Finnish reindeer perirenal tissue (FP-S *vs.* FP-W).

S7 Table. Significantly differentially expressed genes between spring and winter in

Finnish reindeer prescapular tissue (FS-S vs. FS-W).

S8 Table. Significantly differentially expressed genes between spring and winter in

Even reindeer metacarpal tissue (EM-S *vs.* EM-W).

S9 Table. Significantly differentially expressed genes between spring and winter in

Even reindeer perirenal tissue (EP-S *vs.* EP-W).

S10 Table. Significantly differentially expressed genes between spring and winter in

Even reindeer prescapular tissue (ES-S *vs.* ES-W).

S11 Table. Significantly differentially expressed genes between Even and Finnish

reindeer in metacarpal adipose tissue in spring (EM-S *vs.* FM-S).

S12 Table. Significantly differentially expressed genes between Even and Finnish

reindeer in perirenal adipose tissue in spring (EP-S *vs.* FM-S).

S13 Table. Significantly differentially expressed genes between Even and Finnish

reindeer in prescapular adipose tissue in spring (ES-S *vs.* FS-S).

S14 Table. Significantly differentially expressed genes between Even and Finnish

reindeer in metacarpal adipose tissue in winter (EM-W *vs.* FM-W).

S15 Table. Significantly differentially expressed genes between Even and Finnish

reindeer in perirenal adipose tissue in winter (EP-W *vs.* FPW).

S16 Table. Significantly differentially expressed genes between Even and Finnish

reindeer in prescapular adipose tissue in winter (ES-W *vs.* FS-W).

S17 Table. Significantly differentially expressed genes between female and male

Even reindeer metacarpal adipose tissue (EM-F *vs.* EM-M).

S18 Table. Significantly differentially expressed genes between female and male

Even reindeer perirenal adipose tissue (EP-F *vs.* EP-M).

S19 Table. Significantly differentially expressed genes between female and male

Even reindeer prescapular adipose tissue (ES-F *vs.* ES-M).

S20 Table. List of significantly enriched GO terms associated with significantly

downregulated DEGs (sheet 1) and upregulated DEGs (sheet 2) in Finnish reindeer

metacarpal adipose tissue in spring compared to winter.

S21 Table. List of significantly enriched GO terms associated with significantly

downregulated DEGs (sheet 1) and upregulated DEGs (sheet 2) in Finnish reindeer

perirenal adipose tissue in spring compared to winter.

S22 Table. List of significantly enriched GO terms associated with significantly

downregulated DEGs (sheet 1) and upregulated DEGs (sheet 2) in Finnish reindeer

prescapular adipose tissue in spring compared to winter.

S23 Table. List of significantly enriched GO terms associated with significantly

downregulated DEGs in Even reindeer metacarpal adipose tissue in spring

compared to winter.

S24 Table. List of significantly enriched GO terms associated with significantly

downregulated DEGs (sheet 1) and upregulated DEGs (sheet 2) in Even reindeer

perirenal adipose tissue compared to Finnish reindeer perirenal adipose tissue in

winter.

S25 Table. List of significantly enriched GO terms associated with significantly

upregulated DEGs in Even reindeer prescapular adipose tissue compared to Finnish

reindeer prescapular adipose tissue in spring.

S26 Table. List of significantly enriched GO terms associated with significantly

downregulated DEGs (sheet 1) and upregulated DEGs (sheet 2) in Even reindeer

prescapular adipose tissue compared to Finnish reindeer prescapular adipose tissue

in winter.

S27 Table. List of significantly enriched GO terms associated with significantly

downregulated DEGs in female Even reindeer metacarpal adipose tissue compared

to male Even reindeer metacarpal adipose tissue.

S28 Table. List of significantly enriched GO terms associated with significantly

upregulated DEGs in female Even reindeer perirenal adipose tissue compared to

male Even reindeer perirenal adipose tissue.

S29 Table. List of significantly enriched KEGG pathways associated with

significantly downregulated DEGs (sheet 1) and upregulated DEGs (sheet 2) in

Finnish reindeer metacarpal adipose tissue in spring compared to winter.

S30 Table. List of significantly enriched KEGG pathways associated with

significantly downregulated DEGs (sheet 1) and upregulated DEGs (sheet 2) in

Finnish reindeer perirenal adipose tissue in spring compared to winter.

S31 Table. List of significantly enriched KEGG pathways associated with

significantly downregulated DEGs (sheet 1) and upregulated DEGs (sheet 2) in

Finnish reindeer prescapular adipose tissue in spring compared to winter.

S32 Table. List of significantly enriched KEGG pathways associated with

significantly downregulated DEGs in Even reindeer metacarpal adipose tissue

compared to Finnish reindeer metacarpal adipose tissue in spring.

S33 Table. List of significantly enriched KEGG pathway associated with significantly downregulated DEGs in Even reindeer metacarpal adipose tissue compared to Finnish reindeer metacarpal adipose tissue in winter.

S34 Table. List of significantly enriched KEGG pathways associated with significantly downregulated DEGs (sheet 1) and upregulated DEGs (sheet 2) in Even reindeer perirenal adipose tissue compared to Finnish reindeer perirenal adipose tissue in winter.

S35 Table. List of significantly enriched KEGG pathways associated with significantly downregulated DEGs (sheet 1) and upregulated DEGs (sheet 2) in Even reindeer prescapular adipose tissue compared to Finnish reindeer prescapular adipose tissue in winter.

S36 Table. List of the top 25 most abundant genes expressed in the three adipose tissues based on mean TPM in Finnish reindeer (sheet1) and Even reindeer (sheet2).

Author contributions

Conceptualization: Juha Kantanen

Data curation: Melak Weldenegodguad, Kisun Pokharel, Laura Niranen

Formal analysis: Melak Weldenegodguad, Laura Niranen

Funding acquisition: Juha Kantanen

Investigation: Melak Weldenegodguad, Kisun Pokharel, Laura Niiranen, Päivi Soppela

Methodology: Melak Weldenegodguad, Kisun Pokharel, Laura Niiranen, Päivi Soppela

Project Administration: Juha Kantanen

Resources: Päivi Soppela, Innokentyi Ammosov, Mervi Honkatukia, Heli Lindeberg, Jaana Peippo, Tiina Reilas, Nuccio Mazzullo, Florian Stammler, Juha Kantanen

Supervision: Juha Kantanen, Tommi Nyman, Arja Tervahauta, Päivi Soppela;

Visualization: Melak Weldenegodguad, Kisun Pokharel, Laura Niiranen, Tommi Nyman

Writing – original draft preparation: Melak Weldenegodguad, Kisun Pokharel, Laura Niiranen, Juha Kantanen

Writing – review and editing: Melak Weldenegodguad, Kisun Pokharel, Laura Niiranen, Päivi Soppela, Innokentyi Ammosov, Mervi Honkatukia, Heli Lindeberg, Jaana Peippo, Tiina Reilas, Nuccio Mazzullo, Kari A. Mäkelä, Tommi Nyman, Arja Tervahauta, Karl-Heinz Herzig, Florian Stammler, Juha Kantanen

Funding disclosure

This study was funded by the Academy of Finland in the Arctic Research Programme ARKTIKO (decision number 286040).

Acknowledgements

We thank reindeer herders from the Sallivaara and Oraniemi reindeer herding districts in Finland and from the Eveno-Bytantay District in Sakha, Russia for providing reindeer for our research and for helping with the field work. We thank Tuula-Marjatta Hamama for RNA extractions. We thank the CSC-IT Center for Science, Finland, for computational resources, and the Finnish Functional Genomics Centre supported by the University of Turku, Åbo Akademi University, and Biocenter Finland for RNA sequencing.

Competing interest

The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

1. Forbes BC, Bölter M, Müller-Wille L, Hukkinen J, Müller F, Gunslay N, et al. Reindeer management in northernmost Europe: linking practical and scientific knowledge in social-ecological systems. Springer Science & Business Media; 2006.

2. Røed KH, Flagstad Ø, Nieminen M, Holand Ø, Dwyer MJ, Røv N, et al. Genetic analyses reveal independent domestication origins of Eurasian reindeer. Proceedings of the Royal Society B: Biological Sciences. 2008;275: 1849–1855. doi:10.1098/rspb.2008.0332

3. Weldenegodguad M, Pokharel K, Ming Y, Honkatukia M, Peippo J, Reilas T, et al. Genome sequence and comparative analysis of reindeer (Rangifer tarandus) in northern Eurasia. Scientific Reports. 2020;10: 8980. doi:10.1038/s41598-020-65487-y

4. Lin Z, Chen L, Chen X, Zhong Y, Yang Y, Xia W, et al. Biological adaptations in the Arctic cervid, the reindeer (Rangifer tarandus). Science. 2019;364. doi:10.1126/science.aav6312

5. Soppela P, Saarela S, Heiskari U, Nieminen M. The effects of wintertime undernutrition on plasma leptin and insulin levels in an arctic ruminant, the reindeer. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology. 2008;149: 613–621. doi:10.1016/j.cbpb.2007.12.008

6. Chouchani ET, Kajimura S. Metabolic adaptation and maladaptation in adipose tissue. Nature Metabolism. 2019;1: 189–200. doi:10.1038/s42255-018-0021-8

7. Cannon B, Nedergaard J. Brown Adipose Tissue: Function and Physiological Significance. Physiological Reviews. 2004;84: 277–359. doi:10.1152/physrev.00015.2003

8. Kershaw EE, Flier JS. Adipose Tissue as an Endocrine Organ. J Clin Endocrinol Metab. 2004;89: 2548–2556. doi:10.1210/jc.2004-0395

9. Trzeciak-Ryczek A, Tokarz-Deptuła B, Niedźwiedzka-Rystwej P, Deptuła W. Adipose tissue – component of the immune system. 2011 [cited 15 Jun 2020]. Available: /paper/Adipose-tissue-%E2%80%93-component-of-the-immune-system-Trzeciak-Ryczek-Tokarz-Deptu%C5%82a/4ffd92a4b2cdc1f0c80dc139f8e146fd06ef4c7d

10. Soppela P, Nieminen M, Saarela S, Keith JS, Morrison JN, Macfarlane F, et al. Brown fat-specific mitochondrial uncoupling protein in adipose tissues of newborn reindeer. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology. 1991;260: R1229–R1234. doi:10.1152/ajpregu.1991.260.6.R1229

11. Soppela P, Sormunen R, Saarela S, Huttunen P, Nieminen M. Localization, cellular morphology and respiratory capacity of "brown" adipose tissue in newborn reindeer. Comparative Biochemistry and Physiology Part A: Physiology. 1992;101: 281–293. doi:10.1016/0300-9629(92)90534-W

12. Giordano A, Smorlesi A, Frontini A, Barbatelli G, Cinti S. MECHANISMS IN ENDOCRINOLOGY: White, brown and pink adipocytes: the extraordinary plasticity of the adipose organ. European Journal of Endocrinology. 2014;170: R159–R171. doi:10.1530/EJE-13-0945

13. Bonet ML, Oliver P, Palou A. Pharmacological and nutritional agents promoting browning of white adipose tissue. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids. 2013;1831: 969–985. doi:10.1016/j.bbalip.2012.12.002

14. Li Z, Hardij J, Bagchi DP, Scheller EL, MacDougald OA. Development, regulation, metabolism and function of bone marrow adipose tissues. Bone. 2018;110: 134–140. doi:10.1016/j.bone.2018.01.008

15. Hawkes CP, Mostoufi-Moab S. Fat-bone interaction within the bone marrow milieu: Impact on hematopoiesis and systemic energy metabolism. Bone. 2019;119: 57–64. doi:10.1016/j.bone.2018.03.012

16. Lecka-Czernik B. Marrow fat metabolism is linked to the systemic energy metabolism. Bone. 2012;50: 534–539. doi:10.1016/j.bone.2011.06.032

17. Thouzeau C, Massemin S, Handrich Y. Bone marrow fat mobilization in relation to lipid and protein catabolism during prolonged fasting in barn owls. J Comp Physiol B. 1997;167: 17–24. doi:10.1007/s003600050043

18. Davis JL, Valkenburg P, Reed DJ. Correlations and Depletion Patterns of Marrow Fat in Caribou Bones. The Journal of Wildlife Management. 1987;51: 365–371. doi:10.2307/3801018

19. Soppela P, Nieminen M. The effect of wintertime undernutrition on the fatty acid composition of leg bone marrow fats in reindeer (Rangifer tarandus tarandus L.). Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology. 2001;128: 63–72. doi:10.1016/S1096-4959(00)00297-9

20. Shore AM, Karamitri A, Kemp P, Speakman JR, Graham NS, Lomax MA. Cold-Induced Changes in Gene Expression in Brown Adipose Tissue, White Adipose Tissue and Liver. Lenburg M, editor. PLoS ONE. 2013;8: e68933–e68933. doi:10.1371/journal.pone.0068933

21. Marcher A-B, Loft A, Nielsen R, Vihervaara T, Madsen JGS, Sysi-Aho M, et al. RNA-Seq and Mass-Spectrometry-Based Lipidomics Reveal Extensive Changes of Glycerolipid Pathways in Brown Adipose Tissue in Response to Cold. Cell reports. 2015;13: 2000–13. doi:10.1016/j.celrep.2015.10.069

22. Li B, Qiao L, An L, Wang W, Liu J, Ren Y, et al. Transcriptome analysis of adipose tissues from two fat-tailed sheep breeds reveals key genes involved in fat deposition. BMC Genomics. 2018;19: 1–13. doi:10.1186/s12864-018-4747-1

23. Ma L, Zhang M, Jin Y, Erdenee S, Hu L, Chen H, et al. Comparative Transcriptome Profiling of mRNA and lncRNA Related to Tail Adipose Tissues of Sheep. Front Genet. 2018;9. doi:10.3389/fgene.2018.00365

24. Su A, Fuente LFD, Mateo J, Guti B. Early adipose deposits in sheep : comparative analysis of the perirenal fat transcriptome of Assaf and Churra suckling lambs. 2018; 605–617. doi:10.1111/age.12725

25. MIAO Z, WEI P, KHAN MA, ZHANG J, GUO L, LIU D, et al. Transcriptome analysis reveals differential gene expression in intramuscular adipose tissues of Jinhua and Landrace pigs. J Vet Med Sci. 2018;80: 953–959. doi:10.1292/jvms.18-0074

26. Xing K, Wang K, Ao H, Chen S, Tan Z, Wang Y, et al. Comparative adipose transcriptome analysis digs out genes related to fat deposition in two pig breeds. Scientific Reports. 2019;9: 12925. doi:10.1038/s41598-019-49548-5

27. Xu Y, Qi X, Hu M, Lin R, Hou Y, Wang Z, et al. Transcriptome Analysis of Adipose Tissue Indicates That the cAMP Signaling Pathway Affects the Feed Efficiency of Pigs. Genes. 2018;9: 336. doi:10.3390/genes9070336

28. Ferguson JF, Xue C, Hu Y, Li M, Reilly MP. Adipose tissue RNASeq reveals novel gene-nutrient interactions following n-3 PUFA supplementation and

evoked inflammation in humans. J Nutr Biochem. 2016;30: 126–132. doi:10.1016/j.jnutbio.2015.12.010

29. Haffa M, Holowatyj AN, Kratz M, Toth R, Benner A, Gigic B, et al. Transcriptome Profiling of Adipose Tissue Reveals Depot-Specific Metabolic Alterations Among Patients with Colorectal Cancer. The Journal of Clinical Endocrinology & Metabolism. 2019;104: 5225–5237. doi:10.1210/jc.2019-00461

30. Schleinitz D, Krause K, Wohland T, Gebhardt C, Linder N, Stumvoll M, et al. Identification of distinct transcriptome signatures of human adipose tissue from fifteen depots. European Journal of Human Genetics. 2020;28: 1714–1725. doi:10.1038/s41431-020-0681-1

31. Zhou Y, Sun J, Li C, Wang Y, Li L, Cai H, et al. Characterization of Transcriptional Complexity during Adipose Tissue Development in Bovines of Different Ages and Sexes. te Pas MFW, editor. PLoS ONE. 2014;9: e101261. doi:10.1371/journal.pone.0101261

32. Keipert S, Kutschke M, Ost M, Schwarzmayr T, van Schothorst EM, Lamp D, et al. Long-Term Cold Adaptation Does Not Require FGF21 or UCP1. Cell Metabolism. 2017;26: 437-446.e5. doi:10.1016/J.CMET.2017.07.016

33. Wu Y, Smas CM. Expression and regulation of transcript for the novel transmembrane protein Tmem182 in the adipocyte and muscle lineage. BMC Res Notes. 2008;1: 85. doi:10.1186/1756-0500-1-85

34. Kumar H, Srikanth K, Park W, Lee S-H, Choi B-H, Kim H, et al. Transcriptome analysis to identify long non coding RNA (lncRNA) and characterize their functional role in back fat tissue of pig. Gene. 2019;703: 71–82. doi:10.1016/j.gene.2019.04.014

35. Dang F, Wu R, Wang P, Wu Y, Azam MS, Xu Q, et al. Fasting and Feeding Signals Control the Oscillatory Expression of Angptl8 to Modulate Lipid Metabolism. Scientific Reports. 2016;6: 36926. doi:10.1038/srep36926

36. Lu X, Solmonson A, Lodi A, Nowinski SM, Sentandreu E, Riley CL, et al. The early metabolomic response of adipose tissue during acute cold exposure in mice. Scientific Reports. 2017;7: 3455. doi:10.1038/s41598-017-03108-x

37. Hiroshima Y, Yamamoto T, Watanabe M, Baba Y, Shinohara Y. Effects of cold exposure on metabolites in brown adipose tissue of rats. Molecular Genetics and Metabolism Reports. 2018;15: 36–42. doi:10.1016/j.ymgmr.2018.01.005

38. Lynes MD, Shamsi F, Sustarsic EG, Leiria LO, Wang C-H, Su S-C, et al. Cold-Activated Lipid Dynamics in Adipose Tissue Highlights a Role for Cardiolipin in Thermogenic Metabolism. Cell reports. 2018;24: 781–790. doi:10.1016/j.celrep.2018.06.073

39. Pond CM. An evolutionary and functional view of mammalian adipose tissue. Proceedings of the Nutrition Society. 1992;51: 367–377. doi:10.1079/PNS19920050

40. Pond CM. The Evolution of Mammalian Adipose Tissues. In: Symonds ME, editor. Adipose Tissue Biology. Cham: Springer International Publishing; 2017. pp. 1–59. doi:10.1007/978-3-319-52031-5_1

41. Marioni JC, Mason CE, Mane SM, Stephens M, Gilad Y. RNA-seq: an assessment of technical reproducibility and comparison with gene expression arrays. Genome research. 2008;18: 1509–17. doi:10.1101/gr.079558.108

42. Wang Z, Gerstein M, Snyder M. RNA-Seq: a revolutionary tool for transcriptomics. Nat Rev Genet. 2009;10: 57–63. doi:10.1038/nrg2484

43. Song C, Huang Y, Yang Z, Ma Y, Chaogetu B, Zhuoma Z, et al. RNA-Seq Analysis Identifies Differentially Expressed Genes in Subcutaneous Adipose Tissue in Qaidaford Cattle, Cattle-Yak, and Angus Cattle. Animals. 2019;9: 1077. doi:10.3390/ani9121077

44. Basse AL, Dixen K, Yadav R, Tygesen MP, Qvortrup K, Kristiansen K, et al. Global gene expression profiling of brown to white adipose tissue transformation in sheep reveals novel transcriptional components linked to adipose remodeling. BMC Genomics. 2015;16: 215. doi:10.1186/s12864-015-1405-8

45. Syamsunarno MRAA, Iso T, Yamaguchi A, Hanaoka H, Putri M, Obokata M, et al. Fatty Acid Binding Protein 4 and 5 Play a Crucial Role in Thermogenesis under the Conditions of Fasting and Cold Stress. PLOS ONE. 2014;9: e90825–e90825.

46. Avilés C, Polvillo O, Peña F, Juárez M, Martínez AL, Molina A. Associations between DGAT1, FABP4, LEP, RORC, and SCD1 gene polymorphisms and fat deposition in Spanish commercial beef. J Anim Sci. 2013;91: 4571–4577. doi:10.2527/jas.2013-6402

47. Wei S, Zan LS, Wang HB, Cheng G, Du M, Jiang Z, et al. Adenovirus-mediated interference of FABP4 regulates mRNA expression of ADIPOQ, LEP and LEPR in bovine adipocytes. Genet Mol Res. 2013;12: 494–505. doi:10.4238/2013.January.4.21

48. Kern PA, Gregorio GBD, Lu T, Rassouli N, Ranganathan G. Adiponectin Expression From Human Adipose Tissue: Relation to Obesity, Insulin Resistance, and Tumor Necrosis Factor-α Expression. Diabetes. 2003;52: 1779–1785. doi:10.2337/diabetes.52.7.1779

49. Dall'Olio S, Davoli R, Buttazzoni L, Zambonelli P, Russo V. Study of porcine adiponectin (ADIPOQ) gene and association of a missense mutation with EBVs for production and carcass traits in Italian Duroc heavy pigs. Livestock Science. 2009;125: 101–104. doi:10.1016/j.livsci.2009.03.003

50. Kumar V, Sekar M, Sarkar P, Acharya KK, Thirumurugan K. Dynamics of HOX gene expression and regulation in adipocyte development. Gene. 2021;768: 145308. doi:10.1016/j.gene.2020.145308

51. Masuda Y, Oku H, Okumura T, Nomura K, Kurokawa T. Feeding restriction alters expression of some ATP related genes more sensitively than the RNA/DNA ratio in zebrafish, Danio rerio. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology. 2009;152: 287–291. doi:10.1016/j.cbpb.2008.12.008

52. Mohamed-Ali V, Pinkney JH, Coppack SW. Adipose tissue as an endocrine and paracrine organ. International Journal of Obesity. 1998;22: 1145–1158. doi:10.1038/sj.ijo.0800770

53. Pénicaud L, Cousin B, Laharrague P, Leloup C, Lorsignol A, Casteilla L. Adipose Tissues as Part of the Immune System: Role of Leptin and Cytokines. In: Kordon C, Robinson I, Hanoune J, Dantzer R, Christen Y, editors. Brain Somatic Cross-Talk and the Central Control of Metabolism. Berlin, Heidelberg: Springer; 2003. pp. 81–89. doi:10.1007/978-3-642-18999-9_6

54. Schäffler A, Schölmerich J. Innate immunity and adipose tissue biology. Trends in Immunology. 2010;31: 228–235. doi:10.1016/j.it.2010.03.001

55. Lin Y-W, Wei L-N. Innate immunity orchestrates adipose tissue homeostasis. Hormone Molecular Biology and Clinical Investigation. 2017;31. doi:10.1515/hmbci-2017-0013

56. Tseng Y-H, Ke P-Y, Liao C-J, Wu S-M, Chi H-C, Tsai C-Y, et al. Chromosome 19 open reading frame 80 is upregulated by thyroid hormone and modulates autophagy and lipid metabolism. Autophagy. 2014;10: 20–31. doi:10.4161/auto.26126

57. Luo M, Peng D. ANGPTL8: An Important Regulator in Metabolic Disorders. Front Endocrinol. 2018;9. doi:10.3389/fendo.2018.00169

58. Siddiqa A, Cirillo E, Tareen SHK, Ali A, Kutmon M, Eijssen LMT, et al. Biological Pathways Leading From ANGPTL8 to Diabetes Mellitus–A Co-expression Network Based Analysis. Front Physiol. 2018;9. doi:10.3389/fphys.2018.01841

59. Wing EJ, Magee DM, Barczynski LK. Acute starvation in mice reduces the number of T cells and suppresses the development of T-cell-mediated immunity. Immunology. 1988;63: 677–682.

60. Leibel RL. Molecular physiology of weight regulation in mice and humans. International Journal of Obesity. 2008;32: S98–S108. doi:10.1038/ijo.2008.245

61. Wensveen FM, Valentić S, Šestan M, Turk Wensveen T, Polić B. Interactions between adipose tissue and the immune system in health and malnutrition. Seminars in Immunology. 2015;27: 322–333. doi:10.1016/j.smim.2015.10.006

62. Lee M-J, Fried and SK. Sex-dependent Depot Differences in Adipose Tissue Development and Function; Role of Sex Steroids. Journal of Obesity & Metabolic Syndrome. 2017;26: 172–180. doi:10.7570/jomes.2017.26.3.172

63. Valencak TG, Osterrieder A, Schulz TJ. Sex matters: The effects of biological sex on adipose tissue biology and energy metabolism. Redox Biology. 2017;12: 806–813. doi:10.1016/j.redox.2017.04.012

64. Mukaj A, Piálek J, Fotopulosova V, Morgan AP, Odenthal-Hesse L, Parvanov ED, et al. Prdm9 Intersubspecific Interactions in Hybrid Male Sterility of House Mouse. Mol Biol Evol. [cited 4 Nov 2020]. doi:10.1093/molbev/msaa167

65. Davies B, Hatton E, Altemose N, Hussin JG, Pratto F, Zhang G, et al. Re-engineering the zinc fingers of PRDM9 reverses hybrid sterility in mice. Nature. 2016;530: 171–176. doi:10.1038/nature16931

66. Kulkarni M, Smith HE. E1 Ubiquitin-Activating Enzyme UBA-1 Plays Multiple Roles throughout C. elegans Development. PLOS Genetics. 2008;4: e1000131. doi:10.1371/journal.pgen.1000131

67. Yamauchi Y, Riel JM, Stoytcheva Z, Ward MA. Two Y Genes Can Replace the Entire Y Chromosome for Assisted Reproduction in the Mouse. Science. 2014;343: 69–72. doi:10.1126/science.1242544

68. Ruthig VA, Nielsen T, Riel JM, Yamauchi Y, Ortega EA, Salvador Q, et al. Testicular abnormalities in mice with Y chromosome deficiencies. Biol Reprod. 2017;96: 694–706. doi:10.1095/biolreprod.116.144006

69. Lardone MC, Parodi DA, Valdevenito R, Ebensperger M, Piottante A, Madariaga M, et al. Quantification of DDX3Y, RBMY1, DAZ and TSPY mRNAs in testes of patients with severe impairment of spermatogenesis. Mol Hum Reprod. 2007;13: 705–712. doi:10.1093/molehr/gam057

70. Ramathal C, Angulo B, Sukhwani M, Cui J, Durruthy-Durruthy J, Fang F, et al. DDX3Y gene rescue of a Y chromosome AZFa deletion restores germ cell formation and transcriptional programs. Scientific Reports. 2015;5: 15041. doi:10.1038/srep15041

71. Matsumura T, Endo T, Isotani A, Ogawa M, Ikawa M. An azoospermic factor gene, Ddx3y and its paralog, Ddx3x are dispensable in germ cells for male fertility. J Reprod Dev. 2019;65: 121–128. doi:10.1262/jrd.2018-145

72. Horiuchi K, Perez-Cerezales S, Papasaikas P, Ramos-Ibeas P, López-Cardona AP, Laguna-Barraza R, et al. Impaired Spermatogenesis, Muscle, and Erythrocyte Function in U12 Intron Splicing-Defective Zrsr1 Mutant Mice. Cell Reports. 2018;23: 143–155. doi:10.1016/j.celrep.2018.03.028

73. Kishi K, Uchida A, Takase HM, Suzuki H, Kurohmaru M, Tsunekawa N, et al. Spermatogonial deubiquitinase USP9X is essential for proper spermatogenesis in mice. Reproduction. 2017;154: 135–143. doi:10.1530/REP-17-0184

74. Nilssen KJ, Sundsfjord JA, Blix AS. Regulation of metabolic rate in Svalbard and Norwegian reindeer. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology. 1984;247: R837–R841. doi:10.1152/ajpregu.1984.247.5.R837

75. Fantuzzi G. Adipose tissue, adipokines, and inflammation. Journal of Allergy and Clinical Immunology. 2005;115: 911–919. doi:10.1016/j.jaci.2005.02.023

76. Andrews S. A quality control analysis tool for high throughput sequencing data: s-andrews/FastQC. 2019. Available: https://github.com/s-andrews/FastQC

77. Ewels P, Magnusson M, Käller M, Lundin S. MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics. 2016;32: 3047–3048. doi:10.1093/bioinformatics/btw354

78. Dobin A, Davis CA, Zaleski C, Schlesinger F, Drenkow J, Chaisson M, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2012;29: 15–21. doi:10.1093/bioinformatics/bts635

79. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. 2014;30: 923–930. doi:10.1093/bioinformatics/btt656

80. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 2010;26: 139–140. doi:10.1093/bioinformatics/btp616

81. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology. 2014;15: 550. doi:10.1186/s13059-014-0550-8

82. Tian T, Liu Y, Yan H, You Q, Yi X, Du Z, et al. agriGO v2.0: a GO analysis toolkit for the agricultural community, 2017 update. Nucleic Acids Research. 2017;45: W122–W129. doi:10.1093/nar/gkx382

83. Luo W, Friedman MS, Shedden K, Hankenson KD, Woolf PJ. GAGE: generally applicable gene set enrichment for pathway analysis. BMC Bioinformatics. 2009;10: 161–161. doi:10.1186/1471-2105-10-161

84. Cadenas S, Echtay KS, Harper JA, Jekabsons MB, Buckingham JA, Grau E, et al. The Basal Proton Conductance of Skeletal Muscle Mitochondria from Transgenic Mice Overexpressing or Lacking Uncoupling Protein-3. Journal of Biological Chemistry. 2002;277: 2773–2778. doi:10.1074/jbc.M109736200