The critical role of natural history museums in advancing eDNA for biodiversity studies: a

case study with Amazonian fishes

C. David de Santana[1*], Lynne R. Parenti[1], Casey Dillman[2], Jonathan Coddington[3], D. A. Bastos[4],

Carole Baldwin[1], Jansen Zuanon[5], Gislene Torrente-Vilara[6], Raphaël Covain[7], Naércio A.

Menezes[8], Aléssio Datovo[8], T. Sado[9], M. Miya[9]

1 Division of Fishes, Department of Vertebrate Zoology, MRC 159, National Museum of

Natural History, PO Box 37012, Smithsonian Institution, Washington, DC 20013-7012, USA

2 Cornell University Museum of Vertebrates, Department of Ecology and Evolutionary Biology,

Cornell University, Ithaca, NY, 14850, USA

3 Global Genome Initiative, National Museum of Natural History, PO Box 37012, Smithsonian

Institution, Washington, DC 20013-7012, USA

4 Programa de Pós☐Graduação em Ciências Biológicas (BADPI), Instituto Nacional de

Pesquisas da Amazônia, Manaus, Brazil

5 Coordenac☐ão de Biodiversidade, Instituto Nacional de Pesquisas da Amazo☐nia, Manaus,

Amazonas, Brazil

6 Instituto do Mar, Universidade Federal de São Paulo, Campus Baixada Santista, Santos, São

Paulo, Brazil

7 Muséum d'histoire naturelle, Département d'herpétologie et d'ichtyologie, route de Malagnou

1, case postale 6434, CH-1211, Genève 6, Switzerland

8 Museu de Zoologia da Universidade de São Paulo (MZUSP), Av. Nazaré, 481, 04263-000, São

Paulo, SP, Brazil

9 Natural History Museum and Institute, Chuo-ku, Chiba 260-8682, Japan

*Corresponding author

Keywords: Metabarcoding; Archives; Vouchers; Mitochondrial DNA; Reference libraries;

Scientific collections; Outreach.

Short title: Megadiversity of Amazonian Fishes from a Drop of Water

ABSTRACT. Ichthyological surveys have traditionally been conducted using whole specimen

capture-based sampling with varied, but conventional fishing gear. Recently, environmental

DNA (eDNA) metabarcoding has emerged as a complementary, and possible alternative,

approach to whole specimen approaches. In the tropics, where much of the diversity remains

undescribed, vast reaches remain unexplored, and anthropogenic activities are constant threats;

there have been few eDNA attempts for ichthyological inventories. We tested the discriminatory

power of eDNA using the 12S rRNA MiFish primers with existing public reference libraries and

compared this with capture-based methods in two distinct ecosystems in the megadiverse

Amazon basin. eDNA provided an accurate snapshot of the fishes at higher taxonomic levels and

corroborated the effectiveness of eDNA to detect specialized fish assemblages. Some flaws in

fish metabarcoding studies are routine issues addressed in natural history museums. Thus, by

expanding their archives to include eDNA and adopting a series of initiatives linking collection-

based research, training and outreach, natural history museums can enable the effective use of

eDNA to survey Earth's hotspots of biodiversity before taxa go extinct. Our project surveying

poorly explored rivers and using DNA vouchered archives to build metabarcoding libraries for

Neotropical fishes can serve as a model of this protocol.

INTRODUCTION. Historical ichthyological surveys in freshwater ecosystems globally, were

conducted with whole specimen capture-based sampling using conventional fishing methods

such as gill nets, cast nets, hook and line, dipnets, seines, and rotenone – the last a chemical

ichthyocide. Although the use of ichthyocides is considered advantageous in the tropics where

unknown quantities of diversity remain to be described (e.g., Lundberg *et al*. 2000), collecting

fishes with rotenone has been banned as a sampling method in many regions due to its

extraordinary power to kill fishes and associated fauna (e.g., Relyea, 2005). Capture-based

methods other than rotenone are less powerful, especially for collecting small cryptobenthic

species, and all capture-based methods may result in low capture rates in hard-to-sample

environments, such as rapids, waterfalls, and deep-water reaches. Yet, by undersampling we

increase the likelihood of overlooking, and in the sense of the biodiversity crisis globally,

missing heretofore unaccounted-for species diversity.

Recently, DNA barcodes from environmental samples (eDNA), a non-invasive and

quickly developing methodology that captures genetic material of multiple organisms, has

emerged as a complementary, and a possible alternative approach, to repeated whole specimen

capture methods. Universal eDNA metabarcoding primers based on short variable DNA regions

(typically ribosomal RNA – 12S rRNA, e.g., Miya *et al.*, 2015) were developed to detect

multiple species of fishes through next generation sequencing (NGS) of free DNA molecules that

exist in nature, e.g., lost scales, excrement, and mucosal secretions in the water (Clare *et al*.

2019; Tsuji *et al*., 2020; Miya *et al*., 2020).

The Amazon rainforest and basin maintain the most diverse riverine ichthyofauna on

Earth, with more than 2,700 species classified in 18 orders and 60 families (Dagosta & de Pinna,

2019; Jézéquel *et al*., 2020). Such numbers are underestimates as many undescribed taxa await

discovery and formal description (Reis *et al*., 2003). The evolution of this fauna, one-fifth of the

world's freshwater fishes (Tedesco *et al*., 2017), dates to at least the upper Cretaceous and lower

Cenozoic, ~ 120-150 million years before present (mybp; Brito *et al*., 2007). The Characiphysae

—catfishes (Siluriformes), piranhas and allies (Characiformes) and electric fishes

(Gymnotiformes)— represent more than 75% of the fishes in Amazonian aquatic ecosystems

(Dagosta & de Pinna, 2019). That overwhelming fish diversity is also represented by, among

other taxa, cichlids (Cichliformes), killifishes (Cyprinodontiformes), river stingrays

(Myliobatiformes), pufferfishes (Tetraodontiformes), and silver croakers (Perciformes) (Lowe-

McConnell, 1987; Bloom & Lovejoy, 2017; Dagosta & de Pinna, 2019). Over time these fishes

have diversified under a wildly varied set of environmental conditions to inhabit myriad aquatic

systems (Carvalho *et al.*, 2007). In contrast to the proposed ancient age of those lineages, most

species-level diversification is hypothesized to have occurred relatively recently, less than 10

mybp (e.g., Bloom & Lovejoy, 2017; de Santana *et al.*, 2019).

83   Accurate and thorough sampling is the critical first step towards a more complete

knowledge of biodiversity, a path that also requires the proper identification of collected

samples. Specimens of Amazonian fishes have been identified almost exclusively based on

morphology, but given that molecular evolutionary rates can far outpace divergence in

phenotypes, recent studies that integrate molecular and morphological data have greatly

improved our understanding of species diversity, including that of fishes (Cardoso *et al.*, 2018;

de Santana *et al.*, 2019). DNA barcoding – which typically uses the mitochondrial COI

(Cytochrome Oxidase subunit I) gene to identify candidate species (Hebert *et al.* 2003) – is now

a common molecular method used in taxonomic studies of fishes and has been valuable in

revealing cryptic species diversity and in helping to resolve complex taxonomic issues (Baldwin

et al., 2011; Robertson *et al.*, 2017; de Santana *et al.*, 2019).

94   Accordingly, the demand for samples appropriate for DNA barcoding, i.e., properly

preserved and vouchered in ichthyological collections by the scientific community, has increased

significantly. This demand is correlated directly with efforts to collect DNA-worthy samples

during biodiversity surveys along with museum vouchers and has become a common practice

among scientists worldwide (Weigt *et al.*, 2012; Seberg *et al.*, 2016; Parenti *et al.*, 2020).

Concomitant with sampling and curating efforts, new public platforms have been created to help

close gaps in shared sample information (e.g., Global Genome Biodiversity Network, Droege *et

al.*, 2016) and facilitate access to the sequences (e.g., BOLD). In contrast to this trend,

historically few efforts to collect a substantial number of tissue samples during ichthyological

surveys – possibly because of the lack of infrastructure to maintain such a collection – results in

a lack of robust reference libraries for Amazonian fishes (e.g., Marques *et al.*, 2020). In addition,

despite recognizing that Genbank is a reliable resource (Leray *et al*., 2019), several samples of

Amazonian fishes are poorly identified in GenBank and/or lack of properly preserved voucher

specimens – a problem that extends to other fishes (e.g., Dillman *et al*., 2014; Sidharthan *et al*.,

2021).

Most metabarcoding inventories of freshwater fishes have been conducted in temperate

habitats with well-characterized species diversity (Jerde *et al*., 2019). Yet there have been only a

few attempts to use eDNA metabarcoding in ichthyological surveys in the Neotropical region

(Nobile *et al*., 2019; Cilleros *et al*., 2019; Sales *et al*., 2019; Jackman et al., 2021), an area where

understanding species-level diversity is more complex. For example, Cilleros *et al*. (2019) built

12S eDNA metabarcoding primers based on a reference library for over 130 species known to

occur in the rivers and streams of the French Guiana and the eDNA results were compared with

capture-based sampling methodologies. They recovered a similar number of species, with a

partial match to species identification, using both capture-based and eDNA approaches.

Conversely, Jackman *et al*. (2021) used Miya *et al*. (2015) MiFish primers in three localities in

the central Amazon and suggested that a new approach would be necessary to evaluate the

Neotropical fish fauna using eDNA metabarcoding.

Despite the problems inherent in the development of new methodologies, eDNA

technology and bioinformatics is evolving at accelerated rates and will soon play a central role in

the inventory of fish diversity (Valentini *et al.,* 2016; Jerde *et al*., 2019; McElroy *et al*., 2020;

Miya *et al*., 2020). Freshwater aquatic ecosystems, many of which are poorly explored, are under

severe and fast-pacing threat due to anthropogenic activities (Dudgeon, 2020). Thus, the next

decade or so will be pivotal to survey these habitats to secure vouchers, DNA, and eDNA

samples to build reference libraries and archive the samples as well as to engage society in

protection and preservation as these environments reach their tipping point. Natural history

museums are the sound common ground where key flaws and gaps in those two inversely

proportional trends can be addressed and filled. Here, we tested the discriminatory power of the

MiFish primers using the existing public reference libraries by surveying two distinct

ecosystems, river and stream, during a scientific expedition to the heretofore largely unexplored

Javari River basin in Brazil-Peru-Colombia border. The results of eDNA were compared with the

capture-based methodology and discussed in the context of the critical role of natural history

museums in the development of eDNA metabarcoding as a tool for biodiversity studies.

Results

1. Ichthyological survey – Capture-based sampling (CBS)

In total, 443 species classified in 236 genera, 49 families, and 15 orders were collected

using traditional methods from 46 stations in multiple environments during the Javari River

expedition (Table S1). Among these collections are over 60 species that are new to science.

More specifically, in the three stations sampled using traditional and eDNA

methodologies (Figure 1), we collected: 145 species, 101 genera, 32 families and nine orders in

the main Javari River (station 1); 56 species, 38 genera, 21 families, and six orders in a stream

(station 2); and 67 species, 58 genera, 27 families, and seven orders in the Quixito River (station

3; Table S1).

2. eDNA data analyses and assessment of taxonomic resolution of public reference

database

A total of 1,903,160 reads were assigned to the 11 libraries (station 1 = 5 libraries; station 2 =

5 libraries; station 3 = 1 library), and the number of raw reads for each library ranged from

135,818 to 213,952 with an average of 173,015 reads (Table S2). The final reference database

with 1,671,871 fish reads (99.7% of the denoised reads) yielded 222 MOTUs assigned to 104

genera, 41 families, and 9 orders of fishes (Figure 2 and Tables S3, S4).

Matching sequences identity of >98.5% for 58 species (26%) of 222 species detected by

eDNA were found in the reference library database (Table 1). It represented 36.3% of

Siluriformes; 27.1% of Characiformes; 11.7% of Cichliformes; and 10.8% of Gymnotiformes in

the most species-rich orders detected by eDNA. From these, six species (10.3%) were identified

as "sp." in reference libraries. Only 17species (7.6%) were also identified in the CBS (Table 1).

3. Species composition among eDNA samples: distinguishing between river versus

stream-dwelling communities

Six eDNA samples were collected in the river (stations 1 and 3) and five in the stream

(station 2) and a clear split is seen between these two fish communities (Figure 3). The number

of species detected per sample ranged from 33 to 87 (for details see Supp information) with an

abrupt differentiation between species composition in the stream (samples 1-5) and river samples

(samples 6-11), as detected by the Pearson correlation coefficients (Figure 3A). That is, stream

and river-dwelling communities are clearly distinct on the species composition Habitat axis.

Pearson coefficients are varying from 0.5 to 1.0 in stream versus 0.0 to -0.5 in river. Thus,

species composition is more similar within each community, except for a clear distinction

between the river assemblages at Javari (samples 6-10) and Quixito Rivers (sample 11).

To assess whether the difference in species composition between stream and river

communities observed in the Pearson correlation coefficients were significant, we calculated

Jaccard's dissimilarities indices through a NMDS analysis. The original position of the 222

detected species in river, stream, and in both habitats were represented in a three-dimensional

NMDS space (Figure 3B). The Stress = 0.0524 of the NMDS plot indicated that its first three

axes provided an appropriate three-dimensional representation of the habitats according to their

species composition (Clarke, 1993), and NMDS significantly distinguished between the river and

stream communities (ANOSIM R = 0.4327; $p < 0.0001$; Figure 3B).

Based on the species frequency detected per order we determined the composition of the

stream and river habitats (Figures 3B and C). Of note is the difference in the species composition

between the five samples from Javari River (Samples 1 to 5) and the single sample (Sample 6)

collected in the Quixito River (Figures 3A and D). The interrelationships between habitat and

species diversity and composition per order are represented in the chord diagram in Figure 3E.

4. Comparing Capture-based sampling (CBS) and Molecular-based sampling (MBS) –

eDNA metabarcoding species richness

4.1. Javari River (station 1). CBS captured a total of 145 species, 101 genera, 32 families and

nine orders in the main Javari River. Conversely, MBS found 107 species, 28 genera, 20

families, and seven orders (Figure 4A, B; Tables S5, S6). With 13 species detected by both CBS

and MBS (Table 1). The rarefaction sampling curve illustrating the accumulation of unique

species with the number of individuals collected by CBS does not reach an asymptote (Figure

4C), indicating that several species remain to be detected. This is also corroborated by the Chao

II species richness bias-corrected estimator for MBS, which predicated 216 species (95%

confidence interval: 163–318).

4.2. Stream (station 2). CBS caught 56 species, 38 genera, 21 families, and 6 orders. In contrast,

in the stream, MBS detected 126 species, 22 genera, 17 families and 4 orders (Figure 4D, E;

Tables S7, S8). Six species were detected by the two methodologies (Table 1). The rarefaction

curve for CBS extrapolates to slightly over 60 species the diversity in the stream (Figure 4F).

Conversely, MBS Chao II bias- corrected estimator calculated 145 species in the stream (95%

confidence interval: 134–172).

Discussion

1. Can eDNA provide an accurate snapshot of the Amazonian megadiverse freshwater

ichthyofauna considering current public reference libraries for 12s rRNA?

The Javari River basin contains a considerable fraction of Amazonian fish diversity, ca. 15% of

species, 37% of genera, 60% of families, and 83%. It is, therefore, an excellent testing ground for

eDNA metabarcoding effectiveness for the Amazonian fish fauna. Based on the current public

reference libraries, i.e., Genbank and MiFish DB, MBS provided an accurate snapshot of the

Amazonian megadiverse freshwater ichthyofauna at the Javari River basin when we consider

higher taxonomic levels, i.e., order.

The detection of 222 species in 11 samples from three stations confirms that eDNA is

highly sensitive. However, the low number (28%) of matching sequences with identity of

>98.5% in the public reference libraries suggests severe gaps in the library for Amazonian fishes.

It corroborates a recent global gap analysis of reference databases (Marques *et al*., 2020: figure

1), which revealed that 13% of the over 33,000 known teleostean fish species are sequenced for

12S, representing 38% of genera, 80% of families and 98.5% of orders. For freshwater fishes,

among all continents, South America and Africa had by far the lowest coverage. Not

surprisingly, we found the lowest eDNA identification match at the species level.

Conversely, studies that built reference libraries for highly diverse fish communities
considerably improved the match ratio to species identification between capture-based and
eDNA approaches. Cilleros *et al*. (2019), for example, identified 65% of 203 species of Guianese
fishes. Likewise, Milan *et al*. (2020) detected and correctly assigned all 67 species with 12S
previously designed primers and reference library in the São Francisco River, Brazil. In contrast,
Jackman *et al.* (2021) assigned only 4 of 84 MOTUs to species, demonstrating problems on the
taxonomic resolution in the target gene and general threshold used for species assignment.
The DNA barcoding and eDNA metabarcoding both rely on short, variable, standardized
DNA regions, which can be amplified by PCR, sequenced, and analyzed to identify taxa. The
eDNA approach for vertebrates does not efficiently employ the COI gene because interspecific
genetic variation prevents the use of universal primers (Deagle *et al.,* 2014) and can result in
non-specific amplifications (Collins *et al.*, 2019; but see, Antich *et al.*, 2021). Instead, rRNA
genes used in DNA metabarcoding, such as 12S rRNA (e.g., Miya *et al*., 2015), have the
acceptable resolution at the species level and an elevated copy number per cell due to the number
of mitochondria per cell. Similarly, rRNA genes are preferable over single-copy nuclear DNA
which is less likely to be detected in the environment. Yet, due to the low substitution rate of
rRNA genes it will compromise the identification of fast evolved and complex fish assemblages
such as that in the Neotropical region. Thus, it is likely that in the near future, DNA barcode and
eDNA metabarcode will converge to use large portions of the mitochondrial genome. Therefore,
regardless of the fragment or the threshold used to delimit species (e.g., Jackman *et al.* 2021), it
is essential that studies involving eDNA fish diversity move towards building robust reference
libraries for mitochondrial DNA based on vouchered specimens.
242
2. eDNA species detection across heterogeneous aquatic environments
Amazonian aquatic environments are characterized by specialized fish communities
segregated across a variety of habitats, such as streams, rivers, and their microhabitats (Carvalho
et al., 2007; Vieira *et al*., 2018). In streams, diverse microhabitats are home to leaf-dwelling,
sand-dwelling, and pool-dwelling fish communities (Zuanon *et al*., 2006; Sazima *et al*., 2006;
Espírito-Santo & Zuanon, 2017). Similarly, rivers have specialized fish groups living in high-

energy or deep water (>5 meters) environments. It is critical that fishes inhabiting all aquatic

environments are sampled in biodiversity inventories. Perhaps unsurprisingly, it is incredibly

difficult to sample and therefore assess some microhabitats by CBS. For example, some species

are buried deep in the roots of plants in the riparian zone (e.g., de Pinna *et al.,* 2018), leaf litter,

or in the sand of streams that are extremely difficult to collect with traditional sampling gear.

These life history strategies naturally underestimate the number of fish species living in these

areas due to microhabitat partitioning and undescribed diversity. We corroborate the potential

effectiveness of MBS to detect specialized fish assemblages across heterogeneous aquatic

environments. More intensive sampling efforts might be required to detect low-occurrence taxa

as well as to appropriately sample microhabitats, e.g., filtering a higher amount of water or

collecting water from temporary pools and the river bottom.

2.1. River. In the Javari River (station 1), seven orders were detected by CBS and MBS

(Characiformes, Cichliformes, Clupeiformes, Gymnotiformes, Perciformes, Pleuronectiformes,

Siluriformes). In addition, CBS detected Beloniformes and Myliobatiformes; and MBS found

Osteoglossiformes. The absence of Beloniformes and Myliobatiformes in the MBS could be due

to the poor reference library for comparisons. In contrast, Osteoglossiformes (*Arapaima gigas*) is

well known to occur in the region and specimens were found in the local market. Thus, the

absence of *Arapaima* in the CBS was circumstantial.

The difference in species composition between the two methodologies was detected,

perhaps because of sampling bias in MBS. Water samples for MBS were only collected at the

river surface, detecting mostly free DNA of fish assemblages occurring at midwater and near the

surface, where species-diverse Characiformes are the dominant assemblage. Despite that, as

aforementioned, MBS was also able to take a snapshot of the benthic fish fauna by detecting

many catfish species typically restricted to river channels (e.g., López-Rojas *et al.*,1984; Marrero

& Taphorn, 1991). For example, MBS detected river-dwelling fishes living near the surface, as

well as some deep-water (> 5m depth) inhabitants e.g., *Brachyplatystoma* spp. – goliath

catfishes; Pleuronectiformes – flatfishes; and a large number of unidentified species of electric

fishes (sequences identities within the range of 80–98.5%) belonging to the families

Apteronotidae (10 species) and Sternopygidae (15 species) – common, but often underestimated

components of rivers (e.g., Cox-Fernandes *et al*., 2004; Peixoto *et al*., 2015; de Santana & Vari,

2010).

In addition, the sole sample collected in the mouth of the Quixito River (station 3) was

substantially different from the five samples collected in the Javari River reflecting the different

milieu where the samples were collected. The Javari samples were dominated by Characiformes

whereas Gymnotiformes dominated in the Quixito River sample. In the Javari River, samples

were collected in fast-flowing water along the edge between a shallow peat bog and the main

channel. The Quixito River sample was collected at the mouth of the river, characterized by

small slow-flowing channel.

2.2. Stream. Typically, Characiformes, Siluriformes, Gymnotiformes, Cichliformes,

Cyprinodontiformes, Beloniformes, and Synbranchiformes are the dominant orders in

Amazonian streams (e.g., Castro, 1999). At station 2, both approaches detected species

belonging to Characiformes, Siluriformes, Gymnotiformes, and Cichliformes. In addition, CBS

found Beloniformes and Cyprinodontiformes whereas MBS detected Synbranchiformes for a

total of seven orders. The absence of Beloniformes in the MBS may be due to the poor reference

library for comparisons, and the absence of Synbranchiformes in the CBS here could be due to

the difficulty in collecting cryptobiotic species. We were able to detect at a fine-scale specialized

species assemblage restricted to microhabitats. For example, we captured members of the leaf-

dwelling (e.g., *Apistogramma* spp. – dwarf cichlids) and sand-dwelling (e.g.,

Gymnorhamphichthys spp. – sand knifefishes) fish communities. It remains to be determined

whether eDNA failed to detect fishes that are residents in the temporary pools (e.g., killifishes –

Rivulidae) because of the limitation of its radius of action, or due to the poor reference library for

Neotropical fishes.

Species diversity in Amazonian Terra firme streams ranges from ca. 30 to 170 species

(Mojica *et al*., 2009) with Characiformes and Siluriformes being the most species-rich orders

(e.g., Oliveira *et al*., 2009). Quantification of fish richness in these streams depends upon the

sampling methodology employed and its substrate composition (for reviews see Caramaschi *et*

al., 1999; Anjos & Zuanon, 2007; Mojica *et al*., 2014). For example, in litter banks-rich streams,

Gymnotiformes species diversity can surpass Siluriformes (e.g., Barros *et al*., 2011). In station 2,

according to CBS, Characiformes and Siluriformes were the dominant orders. In contrast,

Characiformes followed by Gymnotiformes were the more species-diverse groups. The

extremely high number of species detected by MBS in the sampled stream, more than twice that

of CBS, primarily in the two dominant orders, Characiformes and Gymnotiformes, is likely

related to four different issues. First, MBS was collected near the confluence between the river

and stream, which may have resulted in occasional, wandering river fishes. Second, the CBS was

conducted with a standardized sampling effort in a restrict (50-m) stretch of the stream (e.g.,

Anjos & Zuanon, 2007), not including its headwaters and areas near its mouth. Third,

Characiformes hidden species diversity. This is corroborated by the historical difficulty in

identification of small tetra species, wherein one named species may represent several

undescribed species, such as in *Astyanax* (e.g., Escobar-Camacho *et al*., 2015; Ramirez *et al*.,

2017). Fourth, diversity is also underestimated for the Gymnotiformes (e.g., Crampton *et al*.,

2016), for which difficulties in capturing species with cryptobiotic habits possibly play a critical

role in the underestimation of their diversity by CBS methods (sub-estimative may reach three

times the local species richness and up to 10 times the specimens abundance; JZ, unpublished

data).

3.   The role of natural history museums in the advance of eDNA studies

The biodiversity crisis is one of the grand challenges of the 21st century (Abel, 2002; Dudgeon,

2010) with the next two decades critical for the conservation of freshwater environments.

Freshwater ecosystems worldwide hold ca. 30% of vertebrate diversity, including ca. 50% of all

fish species diversity, and are one of the most vulnerable environments on Earth (Jenkins,

2003; Bunn *et al*., 2010; Dudgeon, 2010; Albert *et al*., 2020; Dudgeon, 2020).

330   Combining specimens, DNA sampling and taxonomic identification is required to obtain

a comprehensive assessment of biodiversity. Yet, DNA samples correspond to less than 10% of

the specimens deposited in most fish repositories. Since most fish specimens deposited in

museums and other repositories were collected before using PCR, a vast majority were fixed in

formalin, a standard method of fixation for over a century. Despite the advances in the

techniques of DNA extraction from formalin-fixed materials, the success of these techniques is

still limited especially for specimens stored for long periods in unbuffered solutions (Gilbert *et*

al., 2007; Campos & Gilbert, 2012; Hykin *et al*., 2015). Thus, well-identified vouchered DNA

tissue samples are critical for the identification of unknown DNA in environmental samples.

These DNA tissues may be stored as dried, frozen, or alcohol-fixed samples or as cryopreserved

living samples that have broad potential applications (e.g., Hagedorn, *et al*., 2018). However,

scientific collections in regions holding most of the fish diversity, such as the Neotropics, often

lack the ideal infrastructure to hold long-term genetic resources (e.g. ultrafreezers, liquid

nitrogen storage, cryo-facilities). Nevertheless, GGBN has targeted and sometimes funded

Neotropical institutions to build biorepository capacity and to make their collections globally

discoverable.

These limitations are particularly worrisome given the stark reality of anthropogenic

destruction, climate change and the great extent of predicted unknown diversity that remains to

be described in the Amazon rainforest (Albert & Reis, 2011; Sabaj Perez, 2015). These factors

make this area and Earth's other hotspots of biodiversity priority targets for complete species

inventories in the next decade before suffering irreversible damage (e.g., Amigo, 2020). Another

advantage of eDNA is the long-term biodiversity monitoring in preserved areas/conservation

units (e.g., Murienne *et al*., 2019). The use of eDNA is a highly valuable and cost-effective way

to monitor biodiversity, especially in areas with low anthropogenic threats (McDevitt *et al.,*

2019). This would allow a better prioritization of scarce resources for research and/or

conservation actions.

In the face of these challenges, natural history museums should play a primary role in the

development of eDNA as a tool of biodiversity inventories as well as to track changes in

biodiversity hotspots by:

(1) prioritizing expeditions to jointly secure DNA samples, vouchers, and eDNA in Earth's

hotspots of biodiversity; (2) adapting their biorepositories to archive eDNA samples, which as a

consequence, would provide samples not only for analysis with current but heretofore unseen

technologies; (3) creating reference libraries for the mitochondrial genome; (4) backing up DNA

samples with species-level accuracy on the identification of vouchered specimens; (5) expanding

and improving their tissue biobanks. It is crucial that these modifications for eDNA storage also

occur in museums in Neotropical and Afrotropical countries which host most of freshwater fish

diversity yet lack the resources to build and maintain these tissue collections in perpetuity

(Fernandes *et al*., 2017; Alves *et al*., 2018; Overbeck *et al*., 2018). These efforts would maximize

the information extracted from eDNA metabarcoding and DNA samples, facilitate the design of

sets of universal primers for broader biodiversity inventories, monitor hotspots of biodiversity,

and support taxon-specific surveys; (6) improving public platforms to close gaps in sampling

information and making possible access to DNA sequences; (7) training students and researchers

to use CBS, MBS, morphology and molecular-based taxonomy to survey and identify

biodiversity. By combining eDNA with tissues associated with museum curated-voucher

specimens we can continue to fill gaps currently missing in our knowledge of biodiversity.

Anytime we end with "sp.", species undetermined, in a diagnosis via a new technology the need

for highly trained taxonomic specialists is highlighted. Finally, (8) using eDNA research as a

gateway to inspire and engage the society in natural history and the race against time to survey

and protect Earth's hotspots of biodiversity through education and citizen science programs.

Considering the simplicity of implementing MBS in certain aquatic environments, such as rivers

(see Methods), scientific communities at natural history museums can launch regional/ global

outreach and human resource training initiatives involving citizen scientists, K-12 students, and

professional scientists. Likewise, it would create niches for large-scale natural history museums

to work with regional-scale scientific institutions worldwide, such as in the training of human

resources (e.g., technicians to curate genetic resources) and promoting horizontal transfer of

technology in South America and Africa (e.g., eDNA methodology). In sum, activities involving

eDNA have the potential to fulfill the priorities of natural history museums in the 21st century:

research, collections, training, and outreach.

One successful initiative is the DNA barcoding and metabarcoding libraries for

Amazonian fishes supported by Smithsonian's Global Genome Initiative (GGI), DNA Barcode

Alliance, and São Paulo Research Foundation (FAPESP). The current project is the first of many

scientific expeditions planned over the next three years to survey fishes in poorly explored areas

of the Amazon basin supported by these three initiatives. DNA and eDNA samples and vouchers

are being used to develop a robust, well-documented, mitochondrial DNA reference database.

This eDNA database is validated by morphological (phenotypic) vouchers. Additional eDNA
samples have been collected and deposited in the Smithsonian Institution's National Museum of
Natural History Biorepository. We aim to make available an online platform of all orders and
families, most of the genera, and a significant number of species of Amazonian fishes. Likewise,
GGI is also supporting an initiative for African freshwater fishes. These actions together with the
ongoing development of eDNA technology and bioinformatics will enable the use of eDNA
metabarcoding in fish inventories and the more effective monitoring of hotspots of biodiversity
worldwide.
402

Acknowledgements
The expedition to the Javari River was funded by São Paulo State Research Foundation grant to
GTV (FAPESP #2016/07910-0) and Global Genome Initiative grant to CDS, CBD, and LRP
(GGI-Peer-2017-149). A Global Genome Initiative grant to CDS to survey African rivers and
build DNA barcoding and metabarcoding archives for freshwater fishes (GGI-Peer-2020-258).
JZ received a productivity grant from Brazil's CNPq (#313183/2014-7). AD, CDS, and NAM are
funded by FAPESP (#2016/19075-9). MM and TS were supported by Environment Research and
Technology Development Fund (4-1602) of the Ministry of the Environment, Japan.

DATA AVAILABILITY
Data will be made public on the DRYAD repository upon acceptance.
414

METHODS
Study area. The Javari River encompass an area of 109.202 km^2 with a 1180 km of a main white
water river channel (*sensu* Sioli, 1967; i.e., pH-neutral low-transparency, alluvial sediment-laden
tributary of the Amazon River forming the border between Brazil, Peru and Colombia for ca. 800
km. The first formal records for the Javari river basin were obtained during the Thayer
Expedition to Brazil, in 1865. Most of region remained largely unexplored until our survey
conducted along the Javari River basin during the low water season in July-August of 2017.
422

Specimens sampling and identification. All samples were collected under Jansen Zuanon

permanent permit (SISBIO # 10199-3). Capture-based specimens were sampled at 46 localities

along the Javari River basin (Figure 1) during the low water season in July-August, 2017, using

gill nets, cast nets, hand nets, and trawl nets in rivers, rapids, beaches, streams, and lakes (Table

S1). All fish specimens collected were identified to species level and deposited at the Instituto

Nacional de Pesquisas da Amazônia (INPA) under the numbers INPA-ICT 055148 to INPA-ICT

057159, in Brazil.

Water sampling sites and on-site filtration. Eleven water samples were collected from water

surface at three stations to represent the Javari fish fauna: Station 1, Figure 1; JAV2017081606

(5 samples) – Javari River, below Limoeiro (-4.176, -70.779); Station 2, Figure 1;

JAV2017082108 (5 samples) – "Terra firme" clearwater stream (locally named as "igarapés"),

i.e., acid, highly-transparent and shallow (depth <2 meters), Palmari community (-4.293, -

70.291); and Station 3, QUI 2017082906 (1 Sample) – Quixito River (-4.428, -70.260). We used

low-tech bucket-sampling to collect freshwater using a 10L polypropylene bucket fastened to a 5

m rope (nylon rope, 6 mm in diameter) to collect 5L of water. Before the water sampling, we

wore disposable gloves on both hands and assembled two sets of on-site filtration kits consisting

of a Sterivex filter cartridge (pore size 0.45 µm; Merck Millipore, MA, USA) and a 50 mL

disposable syringe. Then we thoroughly decontaminated the bucket with a foam-style 10%

bleach solution and brought the equipment to the sampling point. We fastened one end of the 5 m

rope to the bucket and collected surface freshwater by tossing and retrieving it. We repeated

collection of fresh water three times to minimize sampling biases at each station.

We performed on-site filtration using two pairs of the filtration kits described above

(filter cartridge + syringe) to obtain duplicate samples. With each collection of fresh water, we

removed the filter cartridge from the syringe, drew approximately 50 ml freshwater into the

syringe by pulling the plunger, reattached the filter cartridge to the syringe, and pushed the

plunger to filter the water. We repeated this step twice in each toss of the bucket sampling so that

the final filtration volume reached 100 ml $\times$ 2 with three tosses of the bucket. When the filter

was clogged before reaching 100-ml filtration, we recorded the total volume of water filtered

(70–100 mL from three stations).

After on-site filtration, we sealed an outlet port of the filter cartridge with Parafilm (Bemis

NA, Wisconsin, USA), added 2 ml of RNAlater (Thermo Fisher Scientific, DE, USA) into the

cartridge from an inlet port of the cartridge using a disposable capillary pipette (Kinglate, USA)

to prevent eDNA degradation, and then sealed the inlet port either with Parafilm or a cap for

preservation. Filtered cartridges filled with RNAlater were kept in –20˚C freezers until shipment

to MM's lab at Natural History Museum and Institute, Chiba, Japan. Samples shipped under

export for biological material permit at room temperature using an overseas courier service.

DNA extraction. All DNA experiments were conducted in MM's lab. We sterilized the

workspace and all equipment before DNA extraction. We used filtered pipette tips and conducted

all eDNA-extractions and manipulations in a dedicated room that is physically separated from

pre- and post-PCR rooms to safeguard against cross-contamination from PCR products.

We extracted eDNA from the filter cartridges using a DNeasy Blood & Tissue kit

(Qiagen, Hilden, Germany) following the methods developed and visualized by Miya *et al.*

(2016) with slight modifications.

We connected an inlet port of each filter cartridge with a 2.0-ml collection tube and

tightly sealed the connection between the cartridge and collection tube with Parafilm. We

inserted the combined unit into a 15-ml conical tube and centrifuged the capped conical tube at

6,000xg for 1 min to remove freshwater and RNAlater. After centrifugation we discarded the

collection tube and used an aspirator (QIAvac 24 Plus, Qiagen, Hilden, Germany) to completely

remove liquid remaining in the cartridge.

We subjected the filter cartridge to lysis using proteinase K. Before the lysis, we mixed

PBS (220 μl), proteinase K (20 μl) and buffer AL (200 μl), and gently pipetted the mixed

solution into the cartridge from an inlet port of the filter cartridge. We again sealed the inlet port

and then placed the cartridge in a 56˚C preheated incubator for 20 min while stirring the

cartridge using a rotator (Mini Rotator ACR-100, AS ONE, Tokyo, Japan) with a rate of 10 rpm.

After the incubation, we removed the film from the inlet port and connected the port with a 2-ml

tube (DNA LowBind tube, SARSTEDT, Tokyo, Japan) for DNA collection. We placed the

combined unit in a 50-ml conical tube and centrifuged the capped tube at 6,000xg for 1 min to

collect the DNA extract.

We purified the collected DNA extract (*ca*. 900 µl) using the DNeasy Blood and Tissue

kit following the manufacture's protocol with a final elution volume of 200 µl. We completed

DNA extraction in one round and used one more premix for the extraction blank (EB) to monitor

contamination. All DNA extracts were frozen at −20°C until paired-end library preparation.

Paired-end library preparation and sequencing. We sterilized the workspace and equipment

in the pre-PCR area before library preparation. We used filtered pipette tips and performed pre-

and post-PCR manipulations in two different, dedicated rooms to safeguard against cross

contamination.

We employed a two-step PCR for paired-end library preparation on the MiSeq platform

(Illumina, CA, USA) and generally followed the methods developed by Miya *et al*. (2015). For

the first-round PCR (1st PCR), we used a mixture of the following four primers: MiFish-U-

forward (5´–ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT NNN NNN GTC GGT

AAA ACT CGT GCC AGC–3´), MiFish-U-reverse (5´–GTG ACT GGA GTT CAG ACG TGT

GCT CTT CCG ATC TNN NNN NCA TAG TGG GGT ATC TAA TCC CAG TTT G–3´),

MiFish-E-forward-v2 (5´–ACA CTC TTT CCC TAC ACG ACG CTC TTC CGA TCT NNN

NNN RGT TGG TAA ATC TCG TGC CAG C–3´) and MiFish-E-reverse-v2 (5´–GTG ACT

GGA GTT CAG ACG TGT GCT CTT CCG ATC TNN NNN NGC ATA GTG GGG TAT CTA

ATC CTA GTT TG–3´). These primer pairs amplify a hypervariable region of the mitochondrial

12S rRNA gene (*ca*. 172 bp; hereafter called "MiFish sequence") and append primer-binding

sites (5´ ends of the sequences before six Ns) for sequencing at both ends of the amplicon. We

used the six random bases (Ns) in the middle of those primer to enhance cluster separation on the

flow cells during initial base call calibrations on the MiSeq platform.

We carried out the 1st PCR with 35 cycles in a 12-µl reaction volume containing 6.0-µl

2 × KAPA HiFi HotStart ReadyMix (KAPA Biosystems, MA, USA), 2.8 µl of a mixture of the

four MiFish primers in an equal volume (U/E forward and reverse primers; 5 µM), 1.2-µl sterile

distilled H_2O and 2.0-µl eDNA template (a mixture of the duplicated eDNA extracts in an equal

volume). To minimize PCR dropouts during the 1st PCR, we performed 8 replications for the

same eDNA template using a strip of 8 tubes (0.2 µl). The thermal cycle profile after an initial 3

min denaturation at 95˚C was as follows: denaturation at 98˚C for 20 sec, annealing at 65˚C for

15 sec and extension at 72˚C for 15 sec with the final extension at the same temperature for 5

min. We also made a 1st PCR blank (1B) during this process in addition to EB. Note that we did

not perform 8 replications and used a single tube for each of the two blanks (EB, 1B) to

minimize cost of the experiments.

After completion of the 1st PCR, we pooled an equal volume of the PCR products from

the 8 replications in a single 1.5-ml tube and purified the pooled products using a GeneRead Size

Selection kit (Qiagen, Hilden, Germany) following the manufacturer's protocol for the

GeneRead DNA Library Prep I Kit. This protocol repeats the column purification twice to

completely remove adapter dimers and monomers. Subsequently we quantified the purified

target products (ca. 172 bp) using TapeStation 2200 (Agilent Technologies, Tokyo, Japan),

diluted it to 0.1 ng/µl using Milli Q water and used the diluted products as templates for the

second-round PCR (2nd PCR). For the two blanks (EB, 1B), we purified the 1st PCR products in

the same manner, but did not quantify the purified PCR products, diluted them with an average

dilution ratio for the positive samples, and used the diluted products as templates for the 2nd

PCR.

For the 2nd PCR, we used the following two primers to append dual-index sequences (8

nucleotides indicated by Xs) and flowcell-binding sites for the MiSeq platform (5´ ends of the

sequences before eight Xs): 2nd-PCR-forward (5´–AAT GAT ACG GCG ACC ACC GAG ATC

TAC ACX XXX XXX XAC ACT CTT TCC CTA CAC GAC GCT CTT CCG ATC T–3´); and

2nd-PCR-reverse (5´–CAA GCA GAA GAC GGC ATA CGA GAT XXX XXX XXG TGA

CTG GAG TTC AGA CGT GTG CTC TTC CGA TCT–3´).

We carried out the 2nd PCR with 10 cycles of a 15-µl reaction volume containing 7.5-µl

$2 \times$ KAPA HiFi HotStart ReadyMix, 0.9-µl each primer (5 µM), 3.9-µl sterile distilled H_2O and

1.9-µl template (0.1 ng/µl with the exceptions of the three blanks). The thermal cycle profile

after an initial 3 min denaturation at 95˚C was as follows: denaturation at 98˚C for 20 sec,

annealing and extension combined at 72°C (shuttle PCR) for 15 sec with the final extension at

the same temperature for 5 min. We also made a 2nd PCR blank (2B) during this process in

addition to EB and 1B.

To monitor for contamination during the DNA extraction, 1st and 2nd PCRs of the 11

samples, we made a total of 3 blanks (EB, 1B, 2B) and subjected them to the above library

preparation procedure.

We pooled each individual library in an equal volume into a 1.5-ml tube. Then we

electrophoresed the pooled dual-indexed libraries using a 2% E-Gel Size Select agarose gel

(Invitrogen, CA, USA) and excised the target amplicons (*ca*. 370 bp) by retrieving them from the

recovery wells using a micropipette. The concentration of the size-selected libraries was

measured using a Qubit dsDNA HS assay kit and a Qubit fluorometer (Life Technologies, CA,

USA), diluted them at 12.0 pM with HT1 buffer (Illumina, CA, USA) and sequenced on the

MiSeq platform using a MiSeq v2 Reagent Kit for 2×150 bp PE (Illumina, CA, USA) following

the manufacturer's protocol. We subjected the pooled dual-indexed libraries a MiSeq run with a

PhiX Control library (v3) spike-in (expected at 5%).

Data preprocessing and taxonomic assignment. We performed data preprocessing and

analysis of MiSeq raw reads using USEARCH v10.0.240 (Edgar, 2010) according to the

following steps: 1) Forward (R1) and reverse (R2) reads were merged by aligning the two reads

using the *fastq_mergepairs* command. During this process, low-quality tail reads with a cut-off

threshold set at a quality (Phred) score of 2, too short reads (<100 bp) after tail trimming and

those paired reads with too many differences (>5 positions) in the aligned region (*ca*. 65 bp)

were discarded; 2) primer sequences were removed from those merged reads using the

fastx_truncate command; 3) those reads without the primer sequences underwent quality filtering

using the *fastq_filter* command to remove low quality reads with an expected error rate of >1%

and too short reads of <120 bp; 4) the preprocessed reads were dereplicated using the

fastx_uniques command and all singletons, doubletons, and tripletons were removed from the

subsequent analysis following the recommendation by the author of the program (Edgar, 2010);

5) the dereplicated reads were denoised using the *unoise3* command to generate amplicon

sequence variants (ASVs) that remove all putatively chimeric and erroneous sequences (Callahan

et al., 2017); 6) finally ASVs were subjected to taxonomic assignments to species names

(metabarcoding operational taxonomic units; MOTUs) using the *usearch_global* command with

a sequence identity of >98.5% with the reference sequences and a query coverage of ≥90% (two

nucleotide differences allowed).

Those ASVs with the sequence identities of 80–98.5% were tentatively assigned "U98.5"

labels before the corresponding species name with the highest identities (*e.g.*,

U98.5_*Synbranchus marmoratus*) and they were subjected to clustering at the level of 0.985

using *cluster smallmem* command. An incomplete reference database necessitates this clustering

step that enables detection of multiple MOTUs under an identical species name. We annotated

such multiple MOTUs with "gotu1, 2, 3…" and tabulated all the outputs (MOTUs plus

U98.5_MOTUs) with read abundances. We excluded those ASVs with sequence identities of

<80% (saved as "no_hit") from the above taxonomic assignments and downstream analyses,

because all of them were found to be non-fish organisms. For a reference database, we used

MiFish DB ver. 36 for taxa assignment, which contained 7,973 species distributed across 464

families and 2,675 genera. In addition, we downloaded all the fish whole mitochondrial genome

and 12S rRNA gene sequences from Genbank as of 15 December 2020.

We refined the above automatic taxonomic assignments with reference to a family-level

phylogeny based on MiFish sequences from both MOTUs and the reference database. For each

family, we assembled representative sequences (most abundant reads) from MOTUs (including

U98.5) and added all reference sequences from that family and an outgroup (a sequence from a

closely-related family) in FASTA format. We subjected the FASTA file to multiple alignment

using MAFFT (Katoh & Standley, 2013) with a default set of parameters. We constructed a

neighbor-joining (NJ) tree with the aligned sequences in MEGA7 (Kumar *et al.*, 2016) using

pairwise deletion of gaps and the Kimura two-parameter distances (Kimura 1980) with the

among-site rate variations modeled with gamma distributions (shape parameter = 1). We

assessed statistical support for internal branches of the NJ tree using the bootstrap resampling

technique (100 resamplings). In addition, aligned sequences were submitted to Bayesian

Inference (BI) analyses run for 10 million generations sampling every 1000 generations to

determine posterior probability for each MOTU and reference sequences. Models were obtained

on JModeltest2 (Darriba *et al*., 2012). BI analyses were run in the Mr. Bayes v3.2.7. (Ronquist *et*

al., 2012). Some of the BI analyses were conducted on the CIPRES science gateway v3.3.

(Miller *et al*., 2015). Trees were analyzed and rendered in iTOL v5.7 (Ciccarelli *et al*., 2006).

The MiSeq paired-end sequencing (2×150 bp) of the 11 libraries, together with an

additional 88 libraries (total = 99), yielded a total of 5,274,381 reads, with an average of 96.5%

base calls, with Phred quality scores of $\geq$30.0 (Q30; error rate = 0.1% or base call accuracy =

99.9%). This run was highly successful considering the manufacture's guidelines (Illumina

Publication no. 770-2011-001 as of 27 May 2014) are >80% bases $\geq$Q30 at 2×150 bp.

Of the 5,274,381 reads, a total of 1,903,160 reads were assigned to the 11 libraries, and

the number of raw reads for each library ranged from 135,818 to 213,952 with an average of

173,015 reads (Table S8). After merging the two overlapping paired-end FASTq files (1,826,828

reads [96.0%]), the primer-trimmed sequences were subjected to quality filtering to remove low-

quality reads (1,802,098 reads [94.7%]). The remaining reads were dereplicated for subsequent

analysis, and single- to tripletons were removed from the unique sequences as recommended by

the author of the program (Edgar, 2010). Then, reads were denoised to remove putatively

erroneous and chimeric sequences, and the remaining 1,677,402 reads (88.1% of the raw reads)

were subjected to taxon assignments. Of these, 1,671,871 reads (99.7% of the denoised reads)

were putatively considered as sequences for fishes, and BLAST searches indicated that non-fish

sequences (5,531 reads [0.3%]) mostly consisted of mammals (i.e., cows, pigs, and humans) and

a few unknown sequences. The three negative controls (i.e., EB, 1B, and 2B) were subjected to

the same analysis pipeline and yielded only 103 denoised reads in total (only 0.006% of the total

raw reads), which were not taken into consideration in the subsequent analyses as their

subtraction from the corresponding species did not affect the presence/absence data matrix of

sequences assignable to fishes. Contamination from non-Amazonian fishes at Miya's lab was

detected and removed (Table S9).

Statistical analyses

All statistical analyses were conducted in R v.4.0.2. (R Core Team, 2020).

Community structure - Molecular-based sampling (MBS)

Evaluation of species richness for eDNA included all 11 samples from the river and stream

localities. Specifically for river: five samples from station 1 (JAV2017081606) and one sample

from station 3 (QUI 2017082906); stream: five samples from station 2 (JAV2017082108).

Species richness between CBS and MBS was performed by comparing fish assemblages

captured and detected in stations 1 and 2 only.

Species abundance per order was evaluated by heatmaps produced in *ggplot2* (Wickham,

2016). Composition per Similarity among all 11 samples, three stations, versus stream and river

assemblages were calculated using the Pearson correlation coefficient. Then, we calculated

Jaccard's dissimilarities, and the coefficient values were ordinated using non-metric

multidimensional scaling (NMDS) to visualize how replicated eDNA data discriminate sites and

habitat (streams vs. rivers) patterns and to determine the sampling effort needed to identify

community changes among sites in the VEGAN package version 2.4–4 (Oksanen *et al*., 2013). A

3D graph was produced in CAR (Fox and Weisberg, 2019) and GLR version 0.103.5 (Adler *et*

al. 2013) packages. Differences in species compositions between sites and habitat types were

statistically tested by permutational analysis of similarities (ANOSIM). It allowed for test of the

statistical significance of similarity between groups comparing to the within groups similarity

using the rank of similarity values (Clarke, 1993). A chord diagram showing the inter-

relationship between species composition and habitat (river versus stream) was produced using

the Circlize package (Gu, 2014). Fish silhouettes were produced in Fishsualize v. 0.2.1

(Schiettekatte *et al*., 2019) with the addition of a species of Gymnotiformes.

Species richness

Water samples station 1 and station 2: The number of detected taxa between CBS and MBS were

represented by Venn diagrams. Graphs were plotted using *ggplot2* and expected species richness

and confidence intervals for species accumulation curves were built using the function

specaccum in VEGAN package v2.5.4 (Oksanen *et al*., 2013) were calculated for stations 1 and

2 (Chao, 1989). The dissimilarity species composition among samples in stations 1 and 2 were

assessed by calculating pairwise Jaccard's distances with the function *vegdist*. Bias-corrected

estimators Chao II (Chao *et al*., 2005) was applied to calculate species richness detected by

MBS, as suggested by Olds *et al*. (2016). It was calculated in SpadeR package in R (Chao *et al*.,

2016).

Molecular-based sampling species accumulation curves were calculated using iNEXT

package in R (Hsieh *et al*., 2020) for Hill number with order $q = 0$ (species richness) with 1000

bootstraps.

References

Abel R. 2002. Conservation biology for the biodiversity crisis: a freshwater follow-up.

Conservation Biology, 5: 1435–1437.

Adler D., O. Nenadic, & W. Zucchini. 2013. rgl: 3D visualization device system (OpenGL). *R*

package version 0.93.945, URL http://CRAN.R-project.org/package=rgl.

Albert J. & R. E. Reis. 2011. *Historical biogeography of Neotropical freshwater fishes*.

University of California Press, Berkeley and Los Angeles, California.

Albert J. S., *et al.* 2020. Scientists' warning to humanity on the freshwater biodiversity crisis.

Ambio, 50: 85–94.

Alves R. J. V., M. Weksler, J. A. Oliveira, P. A. Buckup, H. R. Santana, A. L. Peracchi & A. L.

Albernaz. 2018. Brazilian legislation on genetic heritage harms Biodiversity Convention

goals and threatens basic biology research and education. *Anais da Academia Brasileira de*

Ciências, 90: 1279–1284.

Amigo I. 2020. When will the Amazon hit a tipping point? *Nature,* 578: 505–507.

Anjos M. B. & J. Zuanon. 2007. Sampling effort and fish species richness in small Terra firme

forest streams of central Amazonia, Brazil. *Neotropical Ichthyology*, 5: 45–52.

Antich A., *et al.* 2021. To denoise or to cluster, that is not the question: optimizing pipelines for

COI metabarcoding and metaphylogeography. *BMC Bioinformatics* 22**:** 177.

Baldwin, C. C., C. I. Castillo, L. A. Weigt, & B. C. Victor. 2011. Seven new species within

western Atlantic *Starksia atlantica*, *S. lepicoelia*, and *S. sluiteri* (Teleostei, Labrisomidae),

with comments on congruence of DNA barcodes and species. *ZooKeys*, 79: 21—27.

Barros D. F., J. Zuanon, F. P. Mendonça, H. M. V. Espirito-Santo, A. V. Galuch, & A. L. M

Albernaz. 2011. The fish fauna of streams in the Madeira-Purus interfluvial region,

Brazilian Amazon. *Check List*, 7: 768–773.

Bloom D. D. & N. R. Lovejoy. 2017. On the origins of marine derived fishes in South

America. *Journal of Biogeography.* 44: 1927–1938.

Brito P. M., F. J. Meunier, & M. E. C. Leal. 2007. Origine

et diversification de líchthyofaune Neotropical: une revue. Cybium, 31: 139-153.

Bunn S. E., C. A. Sullivan, C. R. Liermann, P. M. Davies, C. J. Vörösmarty, P. B. McIntire, M.

O. Gessner, D. Dudgeon, A. Prusevich, P. Green & S. Glidden. 2010. Global threats to

human water security and river biodiversity. *Nature*, 467: 555–561.

Callahan B. J., P. J. McMurdie & S. P. Holmes. 2017. Exact sequence variants should replace

operational taxonomic units in marker-gene data analysis. *The ISME Journal,* 11: 2639–

2643.

Campos P. F., & T. M. Gilbert. 2012. DNA extraction from formalin-fixed material. In *Ancient*

DNA (pp. 81–85). Humana Press.

Caramaschi E., P. R. Mazzoni, C. R. S. F. Bizerril, & P. R. Peres-Neto. 1999. Ecologia de Peixes

de Riachos: Estado Atual e Perspectivas. Oecologia Brasiliensis, v. VI, Rio de Janeiro.

Cardoso Y. P., J. J. Rosso, E. Mabragaña, M. Gonzalez-Castro, M. Delpiani, & E. Avigliano.

2018. A continental-wide molecular approach unraveling mtDNA diversity and geographic

distribution of the Neotropical genus *Hoplias*. *Plos One*, 13: e0202024.

Carvalho, L. N., J. Zuanon, & I. Sazima. 2007. Natural history of Amazon fishes. *In*: K. Del-

Claro (Ed.), Tropical Biology and Natural Resources Theme, *In*: K. Del- Claro & R. J.

Marquis (Session Eds. the Natural History Session), Encyclopedia of Life Support Systems

(EOLSS). Eolss Publishers, Oxford.

Castro R. M. C. 1999. Evolução da ictiofauna de riachos sul-americanos: padrões gerais e

possíveis processos causais. In: Caramaschi EP, Mazzoni R, Peres-Neto PR. (Eds) *Ecologia*

de peixes de riachos. Série Oecologia Brasiliensis volume VI, PPGE-UFRJ, Rio de Janeiro,

139–155.

Chao A. 1987. Estimating population size for capture-recapture data with unequal catchability.
Biometrics, 43: 783–91.

Chao A. 1989. Estimating Population Size for Sparse Data in Capture-Recapture Experiments.
Biometrics, 45: 427.

Chao A., R. L. Chazdon, R. K. Colwell, & T.-J. Shen. 2005. A new statistical approach for
assessing compositional similarity based on incidence and abundance data. *Ecology*
Letters, 8: 148–15.

Chao A., K. H. Ma, T. C. Hsieh, T. C. & C. H. Chiu. 2016. SpadeR (Species-richness Prediction
And Diversity Estimation in R): an R package in CRAN. Program and User's Guide also
published at http://chao.stat.nthu.edu.tw/wordpress/software_download/.

Ciccarelli F. D., T. Goerks, C. von Mering, C. J. Creevery, B. Snel, & P. Bork. 2006. Toward
Automatic Reconstruction of a Highly Resolved Tree of Life. *Science,* 311: 1283–1287.

Cilleros K., A. Valentini, A. Cerdan, T. dejean, A. Iribar, P. Taberlet, R. Vigouroux, & R.
Brosse. 2019. Unlocking biodiversity and conservation studies in high diversity
environments using environmental DNA (eDNA): a text with Guianese freshwater fishes.
Molecular Ecology Resources, 19: 27–46.

Clare A. I. M., M. Knapp, N. J. Gemmell, J. Gert-Jan, M. Bunce, M. D. Lamare, H. R. Taylor, R.
Helen. 2019. Beyond Biodiversity: Can Environmental DNA (eDNA) Cut It as a Population
Genetics Tool? *Genes,* 10: 192.

Clarke K. R. 1993. Non-parametric multivariate analyses of changes in community structure.
Austral Ecology, 18: 117–143.

Collins R. A., J. Bakker, O. S. Wangensteen, A. Z. Soto, L. Corrigan, D. W. Sims, M. J. Genner,
& S. Mariani. 2019. Non□specific amplification compromises environmental DNA
metabarcoding with COI. *Methods in Ecology and Evolution*, 10: 1985–2001.

Cox-Fernandes C., J. Podos, & J. G. Lundberg. 2004. Amazonian Ecology: tributaries enhance
the diversity of electric fishes. *Science*, 305: 1960–1962.

Crampton W. G. R., C. D. de Santana, J. C. Waddell, & N. R. Lovejoy. 2016. The Neotropical
electric fish genus *Brachyhypopomus* (Ostariophysi: Gymnotiformes: Hypopomidae):
taxonomy and biology, with descriptions of 15 new species. *Neotropical Ichthyology*, 14:
639–790.

Dagosta F. C. P. & M. C. C. de Pinna. 2019. The fishes of the Amazon: distribution and
biogeographical patterns, with a comprehensive list of species. Bulletin of the American
Museum of Natural History, 431.

Darriba D., G. L. Taboada, R. Doallo, & D. Posada. 2012. jModelTest 2: more models, new
heuristics and parallel computing. *Nature Methods,* 9: 772.

Deagle B. E., S. N. Jarman, E. Coissac, F. Pompanon, & P. Taberlet. 2014. DNA metabarcoding
and the cytochrome c oxidase subunit I marker: Not a perfect match. *Biology Letters*, 10:
20140562.

Dillman C. B., P. Zhuang, T. Zhang, L.-Z. Zhang, N. Mugue, E. J. Hilton. 2014. Forensic
investigations into a GenBank anomaly: endangered taxa and the importance of voucher
specimens in molecular studies. *Journal of Applied Ichthyology*, 30: 1300–1309.

Droege G. *et al.* 2016. The Global Genome Biodiversity Network (GGBN) Data Standard
specification. *Database*: 10.1093/database/baw125.

Dudgeon D. 2010. Prospects for sustaining freshwater biodiversity in the 21st century: linking
ecosystem structure and function. *Current Opinion in Environmental Sustainability*, 5:
422–430.

Dubgeon D. 2020. Freshwater Biodiversity: Status, Threats and Conservation. Cambridge
University Press.

Edgar R. C. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*,
26: 2460–2461.

Escobar-Camacho D., R. Barriga, & S. R. Ron. 2015. Discovering Hidden Diversity of
Characins (Teleostei: Characiformes) in Ecuador's Yasuní National Park. *Plos One*, 10:
e0135569.

Espírito-Santo H. M. V., & J. Zuanon. 2017. Temporary pools provide stability to fish
assemblages in Amazon headwater streams. *Ecology of Freshwater Fish*, 26: 475–483.

Fernandes G. W., M. M. Vale, G. E. Overbeck, M. M. Bustamante, C. E. Grelle, H. G. Bergallo

& J. Araújo. 2017. Dismantling Brazil's science threatens global biodiversity heritage.

Perspectives in Ecology and Conservation, 15: 239–243.

Fox J. & S. Weisberg. 2019. *An R Companion to Applied Regression*, Third edition. Sage,

Thousand Oaks CA.

Gilbert M. T. P., T. Haselkorn, M. Bunce, J. J. Sanchez, S. B. Lucas, L. D. Jewell, & M.

Worobey. 2007. The isolation of nucleic acids from fixed, paraffin-embedded tissues–

which methods are useful when? *Plos One*, 2: e537.

Gu Z. 2014. Circlize implements and enhances circular visualization in R. *Bioinformatics*, 30:

2811–2812.

Hagedorn M. M., J. P. Daly, V. L. Carter, K. S. Cole, Z. Jaafar, C. V. Lager & L. R. Parenti.

2018. Cryopreservation of fish spermatogonial cells: the future of natural history

collections. *Scientific Reports,* 8: 6149.

Hebert P. D. N., A. Cywinska, S. L. Ball, & J. R. de Waard. 2003. Biological identifications

through DNA barcodes. *Proceedings of the Royal Society London. Series B: Biological*

Sciences, 270: 313–321.

Hsieh T. C., K. H. Ma, A. Chao. 2020. *iNEXT: Interpolation and Extrapolation for Species*

Diversity. R package version 2.0.20

Hykin S. M., K. Bi, & J. A. McGuire. 2015. Fixing formalin: a method to recover genomic-scale

DNA sequence data from formalin-fixed museum specimens using high-throughput

sequencing. *Plos One*, 10: e0141579.

Jackman J. M. C. *et al.* 2021. eDNA in a bottleneck: obstacles to fish metabarcoding studies in

megadiverse freshwater systems. Environmental DNA: 10.1002/edna3.191

Jenkins M. 2003. Prospects for biodiversity. *Science*, 302: 1175–1177.

Jerde C. L., E. A. Wilson, & T. L. Dressler. 2019. Measuring global fish species richness with

eDNA metabarcoding. *Molecular Ecology Resources*, 19: 19–22.

Jézéquel C., P. A. Tedesco, & R. Bigorne. 2020. A database of freshwater fish species of the

Amazon Basin. *Scientific Data,* 7: 96.

Katoh K. & D. M. Standley. 2013. MAFFT multiple sequence alignment software version 7:

794   Improvements in performance and usability. *Molecular Biology and Evolution,* 30: 772–

795   780.

796   Kimura M. 1980. A simple method for estimating evolutionary rates of base substitutions

797   through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, 16:

798   111–120.

799   Kumar S., G. Stecher, & K. Tamura. 2016. MEGA7: Molecular evolutionary genetics analysis

800   version 7.0 for bigger datasets. *Molecular Biology and Evolution,* 33: 1870–1874.

801   Leray, M., N. Knowlton, Shien-Lei, H., B. N. Nguyen, & R. J. Machida. 2019. GenBank is a

802   reliable resource for 21[st] biodiversity research. *Proceedings National Academy of Sciences,*

803   116: 22651–22656*.*

804   López-Rojas H., J. G. Lundberg, & E. Marsh. 1984. Design and operation of a small trawling

805   apparatus for use with dugout canoes. *North American Journal of Fisheries Management,*

806   4: 331–334.

807   Lowe-McConnell R. H. 1987. Ecological studies in tropical fish communities. Cambridge

808   Tropical Biology Series. Cambridge University Press, Cambridge.

809   Lundberg J. G., M. Kottelat, G. R. Smith, M. L. J. Stiassny, & A. C. Gill. 2000. So Many Fishes, So

810   Little Time: An Overview of Recent Ichthyological Discovery in Continental Waters. *Annals of*

811   *the Missouri Botanical Garden,* 87: 26–62.

812   Marques, V., P. É. Guérin, M. Rocle, A. Valentini, S. Manel, D. Mouillot, & T. Dejean,

813   2020. Blind assessment of vertebrate taxonomic diversity across spatial scales by clustering

814   environmental DNA metabarcoding sequences. *Ecography*, 43: 1779–1790.

815   Marrero C. & D. C. Taphorn. 1991. Notas sobre la historia natural y la distribution de los peces

816   Gymnotiformes in la cuenca del Rio Apure y otros rios de la Orinoquia. *Biollania,* 8: 123–142.

817   McDevitt A. D., *et al*. 2019. Environmental DNA metabarcoding as an effective and rapid tool

818   for fish monitoring in canals. *Journal of Fish Biology*, 95: 679–682.

819   McElroy M. E., *et al.* 2020. Calibrating environmental DNA metabarcoding to conventional

820   surveys for measuring fish species richness. *Frontiers in Ecology and Evolution*, 8: 276.

Milan D. T., I. S. Mendes & D. C. Carvalho. 2020. New 12S metabarcoding primers for
enhanced Neotropical freshwater fish biodiversity assessment. *Scientific Reports,* 10:
17966.
Miller M. A., *et al.* 2015. A RESTful API for Access to Phylogenetic Tools via the CIPRES
Science Gateway. *Evolutionary Bioinformatics,* 11: 43–48.
Miya M., T. Minamoto, H. Yamanaka, S. Oka, K. Sato, S. Yamamoto, T. Sado, & H. Doi. 2016.
Use of a filter cartridge for filtration of water samples and extraction of environmental
DNA. *Journal of Visualized Experiments*, 117: 54741.
Miya M., Y. Sato, T. Sado, J. Y. Poulsen, K. Sato, T. Minamoto, S. Yamamoto, H. Yamanaka,
H. Araki, M. Kondoh, & W. Ywasaki. 2015. MiFish, a set of universal PCR primers for
metabarcoding environmental DNA from fishes: detection of more than 230 subtropical
marine species. *Royal Society Open Science,* 2: 150088.
Miya M., R. O. Gotoh, & T. Sado. 2020. MiFish metabarcoding: a high-throughput approach for
simultaneous detection of multiple fish species from environmental DNA and other
samples. *Fisheries Science,* 86: 939–970.
Mojica J. I., Castellanos C. & J. Lobón -Cerviá. 2009. High temporal species turnover enhances
the complexity of fish assemblages in Amazonian Terra firme streams. *Ecology of
Freshwater Fish*, 18: 518–526.
Mojica J. I., J. Lobón-Cerviá, & C. Castellanos. 2014. Quantifying fish species richness and
abundance in Amazonian streams: assessment of a multiple gear method suitable for Terra
firme stream fish assemblages. *Fisheries Management and Ecology*, 21: 220–233.
Murienne J., I. Cantera, A. Cerdan, J.B. Decotte, T. Dejean, R. Vigouroux, & S. Brosse. 2019.
Aquatic DNA for monitoring French Guiana biodiversity. *Biodiversity Data Journal*, 7:
e37518.
Nobile A. B., D. Freitas-Souza, F.J. Ruiz-Ruano, M. L. M. O. Nobile, G. O. Costa, F. P. de
Lima, J. P. M. Camacho, F. Foresti, & C. Oliveira. 2019. DNA metabarcoding of
Neotropical ichthyoplankton: enabling high accuracy with lower cost. *Metabarcoding and
Metagenomics*, 3: e.35060.

Oksanen J., R. Kindt, & B. O'Hara. 2013. Package VEGAN. *Community Ecology Package*,
Version 2.

Olds B. P., C. L. Jerde, M. A. Renshaw, L. Yiuan, N. T. Evans, C. R. Turner, K. Deiner, A. R.
Mahon, M. A. Brueseke, P. D. Shirey, M. E. Pfrende, D. M. Lodge, & G. A. Lambert.
2016. Estimating species richness using environmental DNA. *Ecology and Evolution,* 6:
4214–4226**.**

de Oliveira R. R., M. M. Rocha, M. B. Anjos, J. Zuanon, & L. H. Rapp Py-Daniel. 2009. Fish
fauna of small streams of the Catua-Ipixuna Extractive Reserve, State of Amazonas,
Brazil. *Check List,* 5: 154–172.

Overbeck G. E., H. G. Bergallo, C. E. Grelle, A. Akama, F. Bravo, G. R. Colli & G. W.
Fernandes. 2018. Global biodiversity threatened by science budget cuts in Brazil.
BioScience, 68: 11–12.

Parenti L. R., D. Pitassy, Z. Jaafar, K. Vinnikov, N. E. Redmond & K. S. Cole. 2020. Fishes
collected during the 2017 MarineGEO assessment of Kāneʻohe Bay, Oʻahu, Hawaiʻi.
Journal of the Marine Biological Association of the United Kingdom, 100: 607–637.

Peixoto L. A. W., G. M. Dutra, & W. B. Wosiack. 2015. The electric. Glassknife fishes of the
Eigenmannia trilineata group (Gymnotiformes: Sternopygidae): monophyly and description
of seven new species. *Zoological Journal of the Linnean Society*, 175: 384–414.

de Pinna, M. C. C., J. Zuanon, L. R. Rapp-Py-Daniel, & P. Petry. 2018. A new family of
neotropical freshwater fishes from deep fossorial Amazonian habitat, with a reappraisal of
morphological characiform phylogeny (Teleostei: Ostariophysi). *Zoological Journal of the*
Linnean Society, 182: 76–106.

Ramirez J. L, J. L. Birindelli, D. C. Carvalho, P. R. A. M. Affonso, P. C. Venere, H. Ortega, M.
Carrillo-Avila, J.A. Rodríguez-Pulido, & P. M. Galetti Jr. 2017. Revealing Hidden Diversity
of the Underestimated Neotropical Ichthyofauna: DNA Barcoding in the Recently Described
Genus *Megaleporinus* (Characiformes: Anostomidae). *Frontiers in Genetics,* 8: 149.

Reis R. E., S. O. Kullander, & C. J. Ferraris. 2003. Check list of the freshwater fishes of South
and Central America. Edipucrs.

Relyea R. A. 2005. The impact of insecticides and herbicides on the biodiversity and

productivity of aquatic communities. *Ecological Applications*, 15: 618–627.

R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.R project.org/

Robertson D. R., A. Angulo, C. C. Baldwin, C. Pitassy, E. Diane, A. W. Driskell, L. A. Weigt,

A. Lee & I. J. F. Navarro. 2017. Deep-water bony fishes collected by the B/O Miguel Oliver

on the shelf edge of Pacific Central America: an annotated, illustrated and DNA-barcoded

checklist. *Zootaxa,* 4348: 1–125.

Ronquist F., M. Teslenko, P. van der Mark, D.L. Ayres, A. Darling, S. Höhna, B. Larget, L. Liu,

M.A.Suchard & J. P. Huelsenbeck. 2012. MrBayes 3.2: efficient Bayesian phylogenetic

inference and model choice across a large model space. *Systematics Biology,* 61: 539–542.

Sabaj Pérez M. H. 2015. Where the Xingu bends and will soon break. *American Scientist*, 103:

395–403.

Sales N. G., O. S. Wangensteen, D. C. Carvalho, & S. Mariani. 2019. Influence of preservation

methods, sample medium and sampling time on eDNA recovery in a neotropical river.

Environmental DNA, 1: 119–130.

de Santana C. D. & R. P. Vari. 2010. Electric fishes of the genus *Sternarchorhynchus* (Teleostei,

Ostariophysi, Gymnotiformes); phylogenetic and revisionary studies. *Zoological Journal of*

the Linnean Society, 159: 223–371.

de Santana C. D., *et al.* 2019. Unexpected species diversity in electric eels with a description of

the strongest living bioelectricity generator. *Nature Communications*, 10: 4000.

Sazima I., Carvalho L. N., Mendonça F. P., & Zuanon J. 2006. Fallen leaves on the water-bed:

diurnal camouflage of three night-active fish species in an Amazonian streamlet.

Neotropical Ichthyology, 4: 119–122.

Schiettekatte N. M. D., S. J. Brandl, & J. M. Casey. 2019. Fishualize: color palettes based on

fish species. CRAN version 0.2.0.

Seberg O., G. Droege, K. Barker, J. A. Coddington, V. Funk, M. Gostel., G. Petersen, & P. P.

Smith. 2016. Global Genome Biodiversity Network: saving a blueprint of the Tree of Life -

a botanical perspective. *Annals of Botany,* 118: 393–399.

Sidhartan A., R. Raghavan, V. K. Anoop, A. Keskar, & N. Dahanukar. 2021. Phylogenetic

position and relationships of mountain loaches (Teleostei: Balitoridae) of the Western

Ghats as revealed by CO1 sequences. *Zootaxa,* 4926: 79–92.

Tedesco P., *et al.* 2017. A global database on freshwater fish species occurrence in drainage

basins. *Scientific Data* 4: 170141.

Tsuji S., N. Shibata, H. Sawada, & M. Ushio. 2020. Quantitative evaluation of intraspecific

genetic diversity in a natural fish population using environmental DNA. *Molecular Ecology*

Resources, 20: 1323–1332.

Valentini A., *et al.* 2016. Next-generation monitoring of aquatic biodiversity using

environmental DNA metabarcoding. *Molecular Ecology*, 25: 929–942.

Vieira T. B., C. S. Pavanelli, L. Casatti, W.S. Smith, E. Benedito, & R. Mazzoni. 2018. A

multiple hypothesis approach to explain species richness patterns in neotropical stream-

dweller fish communities. *Plos One*, 13: e0204114.

Weigt L. A., C. C. Baldwin, A. Driskell, D.G. Smith, A. Ormos, & E. A. Reyier. 2012. Using

DNA Barcoding to Assess Caribbean Reef Fish Biodiversity: Expanding Taxonomic and

Geographic Coverage. *Plos One*, 7: e41059.

Wickham H. 2016. Ggplot2: Elegant graphics for data analysis. *Houston, TX: Springer.*

Zuanon J., F. A. Bockmann, & I. Sazima. 2006. A remarkable sand-dwelling fish assemblage

from central Amazonia, with comments on the evolution of psammophily in South

American freshwater fishes. *Neotropical Ichthyology*, 4: 107–118.

Table

Table 1. List of 58 species identity of >98.5% detected in the public reference libraries for 12S rRNA in the Javari River basin. S = Station. S1 = Javari River; S2 = Terra firme stream; S3 = Quixito River. Vouchers numbers for species detected with capture-based sampling are provided.

Figures

Figure 1. Map of Javari River basin showing 46 sampling stations (white and red dots). Red dots represent stations in two distinct ecosystems (River: stations 1 – Javari, 3 – Quixito; Stream: station 2 – Terra firme stream) sampled by capture and molecular based methodologies. In the map one dot can represent more than one station. Illustration of 12 orders detected by capture-based sampling (CBS) and molecular based sampling (MBS) in the three stations: (A) Clupeiformes – *Anchoviella jamesi*, INPA-ICT 055391, stations 1&3(CBS); (B) Perciformes – *Plagioscion squamosissimus,* INPA-ICT 055328, stations 1(CBS, MBS), 3(CBS); (C) Cichliformes – *Crenicichla reticulata*, INPA-ICT 055413, station 1(CBS); (D) Osteoglossiformes – *Osteoglossum bicirrhosum*, INPA-ICT 056354; (E) Siluriformes – *Brachyplatystoma vaillantii*, INPA-ICT 056703, station 1(MBS); (F) Characiformes – *Chalceus erythrurus*, INPA-ICT 055360, stations 1(CBS, MBS), 2(MBS); (G) Gymnotiformes – *Eigenmannia limbata*, INPA-ICT 055420, stations 1 & 3(CBS, MBS), 2(MBS); (H) Pleuronectiformes – *Apionichthys nattereri*, INPA-ICT 055487, stations 1(CBS, MBS), 3(MBS); (I) Cyprinodontiformes – *Laimosemion* sp., INPA-ICT 056039, station 2(CBS); (J) Synbranchiformes – *Synbranchus* sp., INPA-ICT 055815, station 2(MBS); (K) Beloniformes – *Potamorrhaphis guianensis*, INPA-ICT 055254, station 2(CBS); (L) Myliobatiformes – newborn specimens of *Potmotrygon orbigny*.

Figure 2. Illustrative cladogram with reads and frequencies for each of 222 metabarcoding operational taxonomic units (MOTUs) and reference sequences included in nine orders and 41 families detected by 11 eDNA samples in the Javari River basin. Color highlighting MOTUs names corresponds to each of the nine orders. In the left side, species richness, key color, and

general bauplan silhouettes for each order. At the center, spherical view of species diversity

detected by eDNA.

Figure 3. Ichthyofauna segregation into river and stream at Javari Basin as detected by 11

samples of eDNA in the three stations: stream (station 2): samples 1 to 6; Javari river (station 1):

Samples 6 to 10; and Quixito River (station 3): sample 11. (A) Heatmap based on the Pearson

correlation coefficients between species composition and habitat. Note the difference in the

species composition along the river, i.e., Javari versus Quixito rivers (B) Non-metric

multidimensional scaling (NMDS) based on Jaccard's dissimilarities coefficients discriminating

habitat (streams vs. rivers). The Stress of the NMDS plot 0.052 indicates that its first three axes

provided an appropriate three dimensional representation of the habitats according to their

species composition. Each dot represents a species and the relative distance between two points

represents the dissimilarity. The ANOSIM $p < 0.0001$ suggests that NMDS significantly

distinguished between the river and stream communities; (C) Heatmap for species abundance for

each of the five orders detected in the stream; (D) Heatmap showing species abundance for each

of the eight orders detected in the Javari river and Quixito River. Note the alteration in species

abundance between samples 5 to 10 (Javari) and 11 (Quixito). In samples 5 to 10 Characiformes

and Siluriformes are more abundant. Conversely, Gymnotiformes and Siluriformes are more

species rich in the sample 6. (E) Chord diagram showing the directional relationship between

habitat and species richness distributed into the nine detected orders.

Figure 4. Comparison between Capture-Based Sampling (CBS) in blue and Molecular-Based

Sampling (MBS) in red in two sampled localities. Javari River: (A) Histogram comparing the

number of species per order detected by CBS and MBS; (B) Venn diagram of the number of

orders detected by CBS and MBS; (C) Rarefaction species accumulation curve for CBS with

95% confidence interval and extrapolation for twice the number of individuals sampled. Terra

firme stream: (D) Histogram comparing the number of species per order detected by CBS and

MBS; (E) Venn diagram of the number of orders detected by CBS and MBS; (F) Rarefaction

species accumulation curve for CBS with 95% confidence interval and extrapolation for twice

the number of individuals sampled.

Station 1 - Javari River

Station 2 - Terra firme stream

Station 3 - Quixito River

A

B

C

D

E

F

G

H

I

J

K

L

Station 1

Station 2

Station 3

Javari River

Quixito River

Itaquai River

COLOMBIA

PERU

BRAZIL

Leticia

Tabatinga

Tabatinga Airport

Alfredo Vásquez Cobo Airport

Benjamin Constant

-4°3'0"

-4°30'0"

-70°48'0"

-70°24'0"

-70°0'0"

0   10   20 km

MOTUS richness in 9 orders

Reads 1,671,871
Frequencies 1 to 11
MOTUs & Orders 222   9
Families 41

Characiformes 45.9%
Siluriformes 24.7%
Gymnotiformes 16.6%
Cichliformes 8.1%
Clupeiformes 1.8%
Perciformes 0.9%
Synbrachiformes 0.9%
Pleuronectiformes 0.45%
Osteoglossiformes 0.45%

Order	Family	Species	Read abundance	Frequency	CBS	Station
Osteoglossiformes	Arapaimidae	*Arapaima gigas*	7,050	4		
Clupeiformes	Engraulidae	*Anchoviella* sp. 1	18,117	4	INPA-ICT 055766	S1
Characiformes	Crenuchidae	*Characidium* sp. 1	69	2		
Characiformes	Erythrinidae	*Hoplerythrinus unitaeniatus*	1,332	5		
Characiformes	Erythrinidae	*Hoplias malabaricus*	324	2	INPA-ICT 055204	S2
Characiformes	Cynodontidae	*Cynodon meionactis*	1,769	1		
Characiformes	Cynodontidae	*Hydrolycus scomberoides*	9,118	4		
Characiformes	Serrasalmidae	*Serrasalmus eigenmanni*	1,222	2		
Characiformes	Hemiodontidae	*Anodus* sp.	4,483	3		
Characiformes	Anostomidae	*Abramites hypselonotus*	1,304	1		
Characiformes	Anostomidae	*Leporinus apollo*	846	3		
Characiformes	Anostomidae	*Leporinus fasciatus*	31	1		
Characiformes	Anostomidae	*Scheizodon knerii*	46,775	6		
Characiformes	Chilodontidae	*Caenotropus labyrinthicus*	14	1	INPA-ICT 055404	S2
Characiformes	Curimatidae	*Potamorhina* sp.	18,470	3		
Characiformes	Prochilodontidae	*Prochilodus harttii*	90,591	6		
Characiformes	Prochilodontidae	*Semaprochilodus* sp. 1	15,460	5		
Characiformes	Acestrorhynchidae	*Acestrorhynchus falcatus*	11,546	5	INPA-ICT 055991	S2
Characiformes	Chalceidae	*Chalceus erythrurus*	8,741	5	INPA-ICT 055360	S1, S2
Characiformes	Chalceidae	*Chalceus macrolepidotus*	4,158	2		
Characiformes	Characidae	*Charax pauciradiatus*	10,186	5		
Characiformes	Characidae	*Charax* sp. 1	32	1		
Characiformes	Characidae	*Ctenobrycon hauxwellianus*	491	3		
Characiformes	Characidae	*Jupiaba ocellata*	1,150	4		
Characiformes	Characidae	*Moenkhausia lepidura*	782	3		
Characiformes	Serrasalmidae	*Myloplus rubripinnis*	232	4		
Characiformes	Serrasalmidae	*Pygocentrus nattereri*	19,766	9	INPA-ICT 055811	S1, S2
Characiformes	Iguanodectidae	*Bryconops affinis*	9,344	5		

Order	Family	Species			Voucher	
Characiformes	Triportheidae	*Triportheus* sp. 1	232	3		
Siluriformes	Loricariidae	*Dekeyseria amazonica*	86	1	INPA-ICT 055419	S1
Siluriformes	Loricariidae	*Hypostomus plecostomus*	113	1		
Siluriformes	Loricariidae	*Lasiancistrus saetiger*	1,837	1		
Siluriformes	Doradidae	*Hemidoras morrisi*	14	1	INPA-ICT 055422	S1
Siluriformes	Doradidae	*Liosomadoras morrowi*	25	1		
Siluriformes	Doradidae	*Megalodoras uranoscopus*	9	1	INPA-ICT 055491	S1
Siluriformes	Doradidae	*Platydoras costatus*	40	2		
Siluriformes	Doradidae	*Pterodoras granulosus*	94	1	INPA-ICT 055463	S1
Siluriformes	Doradidae	*Trachydoras nattereri*	34	1	INPA-ICT 055660	S3
Siluriformes	Pimelodidae	*Hypophthalmus edentatus*	472	2		
Siluriformes	Heptapteridae	*Pimelodella cristata*	1,359	7		
Siluriformes	Pimelodidae	*Pseudoplatystoma eticulatum*	2,064	6		
Siluriformes	Pimelodidae	*Pinirampus pinirampu*	4	1	INPA-ICT 055316	S1
Siluriformes	Pimelodidae	*Pseudoplatystoma tigrinum*	183	2	INPA-ICT 055379	S1
Siluriformes	Pimelodidae	*Hemisorubim platyrhynchos*	110	1		
Siluriformes	Pimelodidae	*Brachyplatystoma vaillantii*	25	1		
Siluriformes	Pimelodidae	*Sorubimichthys planiceps*	431	1		
Siluriformes	Pimelodidae	*Sorubim elongatus*	282	1		
Siluriformes	Pimelodidae	*Phractocephalus hemioliopterus*	316	4		
Siluriformes	Pimelodidae	*Zungaro jahu*	31	1		
Gymnotiformes	Gymnotidae	*Electrophorus varii*	1,234	5		
Gymnotiformes	Gymnotidae	*Gymnotus carapo*	121	1		
Gymnotiformes	Sternopygidae	*Eigenmannia limbata*	10,443	7	INPA-ICT 055420	S1, S2, S3
Gymnotiformes	Apteronotidae	*Apteronotus albifrons*	567	5		
Characiformes	Ctenoluciidae	*Boulengerella maculata*	126	1	INPA-ICT 055263	S1
Cichliformes	Cichlidae	*Cichla ocellaris*	52	1		
Cichliformes	Cichlidae	*Hypselecara temporalis*	38	1		

| Pleuronectiformes | Achiridae | *Hypoclinemus mentalis* | 50,324 | 1 | INPA-ICT 055269 | S1 |
| Perciformes | Sciaenidae | *Plagioscion squamosissimus* | 1,220 | 2 | INPA-ICT 055328 | S1 |