An Immune Cell Atlas Reveals Dynamic COVID-19 Specific Neutrophil Programming Amenable to Dexamethasone Therapy

Authors: Sarthak Sinha[1*], Nicole L. Rosin[1*+], Rohit Arora[1], Elodie Labit[1], Arzina Jaffer[1], Leslie Cao[1], Raquel Farias[3,4], Angela P. Nguyen[3,4], Luiz G. N. de Almeida[2,5], Antoine Dufour[2,5], Amy Bromley[6], Braedon McDonald[3,4], Mark Gillrie[3,7,8], Marvin J. Fritzler[3,8], Bryan Yipp[3,4 +], Jeff Biernaskie[1,9,10 +]

Affiliations:

[1] Department of Comparative Biology and Experimental Medicine, Faculty of Veterinary Medicine, University of Calgary;

[2] Department of Physiology & Pharmacology;

[3] Calvin, Phoebe and Joan Snyder Institute for Chronic Diseases, Cumming School of Medicine, University of Calgary;

[4] Department of Critical Care Medicine, Cumming School of Medicine, University of Calgary;

[5] McCaig Institute for Bone and Joint Health, University of Calgary;

[6] Department of Pathology and Laboratory Medicine, University of Calgary;

[7] Department of Microbiology, Immunology and Infectious Diseases, University of Calgary;

[8] Department of Medicine, Cumming School of Medicine, University of Calgary, Calgary, AB, Canada;

[9] Hotchkiss Brain Institute, University of Calgary;

[10] Alberta Children's Hospital Research Institute, University of Calgary

[*] Equal contribution

[+] Corresponding authors

Summary

SARS-CoV-2 is a novel coronavirus that causes acute respiratory distress syndrome (ARDS), death and long-term sequelae. Innate immune cells are critical for host defense but are also the primary drivers of ARDS. The relationships between innate cellular responses in ARDS resulting from COVID-19 compared to other causes of ARDS, such as bacterial sepsis is unclear. Moreover, the beneficial effects of dexamethasone therapy during severe COVID-19 remain speculative, but understanding the mechanistic effects could improve evidence-based therapeutic interventions. To interrogate these relationships, we developed an scRNA-Seq and plasma proteomics atlas (biernaskielab.ca/COVID_neutrophil). We discovered that compared to bacterial ARDS, COVID-19 was associated with distinct neutrophil polarization characterized by either interferon (IFN) or prostaglandin (PG) active states. Neutrophils from bacterial ARDS had higher expression of antibacterial molecules such as PLAC8 and CD83. Dexamethasone therapy in COVID patients rapidly altered the IFN^{active} state, downregulated interferon responsive genes, and activated IL1R2^{+ve} neutrophils. Dexamethasone also induced the emergence of immature neutrophils expressing immunosuppressive molecules ARG1 and ANXA1, which were not present in healthy controls. Moreover, dexamethasone remodeled global cellular interactions by changing neutrophils from information receivers into information providers. Importantly, male patients had higher proportions of IFN^{active} neutrophils, a greater degree of steroid-induced immature neutrophil expansion, and increased mortality benefit compared to females in the dexamethasone era. Indeed, the highest proportion of IFN^{active} neutrophils was associated with mortality. These results define neutrophil states unique to COVID-19 when contextualized to other life-threatening infections, thereby enhancing the relevance of our findings at the bedside. Furthermore, the molecular benefits of dexamethasone therapy are also defined, and the identified pathways and plasma proteins can now be targeted to develop improved therapeutics.

COVID-19 ARDS host responses contextualized to bacterial ARDS.

A broad array of infections including SARS-CoV-2 and bacterial sepsis can induce acute respiratory distress syndrome (ARDS), respiratory failure and death[1-3]. Neutrophils are thought to be key drivers of both COVID-19 and bacterial ARDS[4-6], yet it is unclear if this is related to intrinsic and/or irreversible cellular responses. While recent studies have leveraged single-cell transcriptomics to dissect peripheral[7-9] and bronchoalveolar fluid [10-12]immune landscapes driving COVID-19 pathogenesis, the protocols used can inadvertently exclude the majority of polymorphonuclear granulocytes, including neutrophils, as they are highly sensitive cells with low RNA (and high RNase) content. Here, we employ whole-blood-preserving protocols that capture all major immune cell types from critically ill patients admitted to intensive care units (ICUs) (Extended Fig 1). All samples taken from COVID-19 patients were assessed by bacterial culture and tested negative. All COVID-19 patients tested positive by PCR for SARS-CoV-2, and we previously confirmed an absence of viral mRNA in any circulating immune cells in a subset of patients[13]. However, a plasma proteomic screen for SARS-CoV-2 specific viral proteins in all samples revealed detection of one or more viral proteins in COVID-19 patient serum (Extended Fig 2a). Furthermore, we compared patient samples from COVID-19 ARDS to bacterial sepsis with ARDS (herein referred to as bacterial ARDS) (Extended Fig 2b), as there were unusually low admissions to ICU with viral pneumonias/ARDS during the period studied, likely due to COVID-19 public health measures.

Patient cohorts had comparable ages, sex, days on life support and time in hospital, but COVID-19 patients had broader racial diversity (Extended Fig 2c,d, Extended Data Table 1). Bacterial ARDS induced significant neutrophilia, and relative thrombocytopenia compared to the near normal circulating neutrophil numbers in COVID-19, while both had similar degrees of lymphopenia (Extended Fig 2e). Both cohorts had comparable PaO2 / FiO2 (P/F) ratios, which is an indicator of the severity of ARDS[14], but bacterial ARDS patients had significantly more kidney injury demonstrated by higher levels of serum creatinine (Extended Fig 2f). We further compared families of soluble inflammatory markers (Extended Fig 2g) used to distinguish prototypical states, including those identified during cytokine storm (Extended

Fig 2h) and cytokine release syndrome (Extended Fig 2i)[15], which demonstrated similar soluble cytokine

and chemokine responses between the infections. Therefore, in the context of life-threatening bacterial

ARDS, COVID-19 ARDS patients had normal neutrophil counts, comparable IL-6 levels, and less organ

failure as indicated by serum creatinine levels, all of which have been previously proposed as markers of

COVID disease severity[16,17]. This prompted us to further investigate immune states and composition in

response to COVID-19 compared to bacterial ARDS.

The online companion atlas (biernaskielab.ca/COVID_neutrophil) contains accessible scRNAseq data

performed on freshly obtained whole blood at timepoint 1 (t1, <72h after ICU admission) and at timepoint

2 (t2, 7 days after t1) (Fig 1a). Cellular identity was mapped to 30 immune cell types/states using a

UMAP projection from 21 patients and 86,935 cells (Fig 1b, Extended Figure 3a). Global magnitude of

gene expression was directly compared between COVID-19 and bacterial ARDS patients (Extended Data

Table 3), which revealed a more globally altered distribution of differential expression at t1 than at t2.

Altered regulation of genes was most pronounced in neutrophils at t1, with lower neutrophil gene

expression in COVID-19 compared to bacterial ARDS (Fig 1c; Extended Fig 3b-c). At t2, the global

alterations in gene expression when comparing COVID-19 to bacterial ARDS were most pronounced in

plasmablasts (Fig 1d; Extended Fig 3d-e). We further compared and quantified the proportions of known

peripheral blood cellular constituents, which highlighted significant differences in CD4 T cells, CD8 T

cells and NK cells (Extended Fig 3f). These data highlight that significant global differences in immune

cell gene expression exist between COVID-19 ARDS and bacterial ARDS.

COVID-19 drives specific neutrophil maturation states.

Neutrophils are a primary participant in the development of ARDS[18]; yet despite similar severity of

ARDS between our bacterial and COVID-19 cohorts, the numbers of circulating neutrophils from clinical

cell counts were significantly different (Extended Fig 2d). We hypothesized that neutrophil qualitative

states may be important determinants of disease. Neutrophils were subjected to velocity analysis[19,20] to

reconstruct maturation dynamics. Louvain clusters (Fig 1e), clinical cohort, individual patient, and

velocity length were overlayed on velocity vector fields (Extended Fig 4a-d). The proportions of distinct

neutrophil states were compared at t1 and this revealed a divergent expansion of IFN^{active} neutrophils

(clusters 2, 4 and 5) marked by IFITM1 expression in COVID-19, which became similar to bacterial

ARDS at t7 (Fig 1f-h). Expression of IFITM1 in neutrophils from COVID-19 patients at t1 was

confirmed by immunofluorescent staining for IFITM1, colocalized with S100A8/9 and typical neutrophil

nuclear morphology. Relative to healthy donors, the IFN^{active} population in both COVID-19 and bacterial

ARDS patients were elevated (Extended Fig 4h-k), suggesting that infections dramatically alter neutrophil

dynamics and that comparing COVID-19 neutrophils to healthy neutrophils may only reveal broad

features separating pathogen-challenged versus non-challenged (homeostatic) neutrophils. Hence, to map

pathogen-activated neutrophils dynamics with high resolution, subsequent analyses employ principal

components with top loading genes that distinguish different pathogen-activated states arising during

COVID-19 and bacterial sepsis for downstream dimensionality reduction.

Figure 1 – COVID-19 alters neutrophil maturation. a. Schematic summarizing patients with COVID-
19 and bacterial sepsis profiled at t1 and t2. **b.** UMAP projection of 86,935 whole blood cells from 21

patient samples, coloured by Azimuth reference-mapped immune cell states. **c-d.** Kernel density estimates
depicting magnitude of molecular response elicited by immune cell subsets during COVID-19 compared
to Bacterial ARDS at t1 (c) and t2 (d) calculated by summing DEG fold changes for each cell state shown
in Panel a. **e.** UMAP plotting RNA velocity analysis of 29,653 subclustered neutrophils undergoing state
transitions, coloured by cluster ID. **f.** Stacked bar plot depicting cluster composition of clinical cohorts
examined. **g.** UMAP coloured by neutrophil clusters and overlaid with summary path curves based on
vector fields and neutrophil state compositions in Panel d and e, respectively to determine neutrophil
states. **h.** Immunocytochemistry for S100A8/A9 (red) and IFITM1 (green) expression on leukocyte-rich
preparation from COVID-19 donor at tl. **i-k.** Transcriptional kinetics driving expansion of IFN^{active} (i),
Bacterial ARDS-enriched (j), and PG^{active} (k) neutrophils. Latent time distribution of trajectory-associated
louvain clusters (left), phase portraits with equilibrium slopes of spliced–unspliced ratios (center), and
RNA velocity and gene expression (right) of selected genes driving divergent maturation trajectories.
Phase portraits are coloured by clinical cohort.

Classically, peripheral neutrophils are considered terminally differentiated and non-dividing, however the

increase in velocity length suggested the ability to alter phenotypic states once in circulation along

specific paths or 'lineages'. COVID-19 neutrophils followed unique maturation paths compared to

bacterial ARDS, culminating in three distinct terminal states: Interferon active (IFN^{active}), prostaglandin

active (PG^{active}) or bacterial ARDS enriched (Fig 1e-g; Extended Fig 4e). Interestingly, the apex of this

trajectory was marked by high velocity lengths, characteristic of cells undergoing differentiation

(Extended Fig 4c, d). COVID-19 neutrophils preferentially transitioned from the apex of the trajectory,

which was an immature state (TOP2A expressing; Extended Fig 4e) to an IFN responsive state

characterized by IFITM1, IFITM2 and IFI6 expression (Cluster 1 to 4 and 5; Fig 1i; Online Atlas). This is

clearly illustrated in Extended Video 1. This immature state was not present in healthy controls, though it

is present in both comparator groups, suggesting these states are liberated into circulation upon pathogen

exposure (Extended Fig 4h-k). The lineage relationship was less clear for COVID-19 enriched PG^{active}

clusters defined by prostaglandin responsive genes (clusters 2, 6 and 8), with notable increases in

PTGER4 and PTGS2 (or COX2), a proposed therapeutic target in COVID-19[21] (Fig 1k; Extended Fig 4f,

g, Online Atlas). The dominant conventional bacterial ARDS state was characterized by antibacterial

proteins CD83[22], CD177, and PLAC8[23] (cluster 3 to 0; Fig 1j; Online Atlas). Taken together, this data

demonstrated that peripheral neutrophils have dynamic programming abilities which result in COVID-19

specific neutrophil polarization defined by the emergence of IFN^{active} and PG^{active} neutrophil states.

Unique transcriptional regulatory pathways drive neutrophil maturation in COVID-19.

Rapid and robust IFN responses protect against COVID-19 severe disease, while delayed responses could exacerbate systemic and pulmonary inflammation[24,25]. Moreover, neutrophil IFN responses are not traditionally considered during infections and neutrophils are generally considered to be homogenous, with a uniform proinflammatory capacity. Global neutrophil expression aligned with neutrophil state specific markers, such as interferon response genes (IFITM1, RSAD2, IFI6, and ISG10), being more highly expressed in COVID-19 neutrophils (Fig 2a; Extended Fig 4f). The inverse was the case for anti-bacterial proteins like PLAC8 (Fig 2a; Online Atlas). However, the discovery of differential neutrophil states prompted further exploration of the factors driving neutrophil state polarization. Gene regulatory network reconstruction using SCENIC analysis[26] revealed differentially activated transcription factors STAT1, IRF2 and PRDM1 in COVID-19 (Fig 2b), while bacterial ARDS neutrophils had increased prototypical granulocyte transcription factors such as CEBPA, CEBPB, STAT5B and less defined factors such as NFE2 (Fig 2b, Online Atlas). PRDM1 activation was most pronounced in the IFN^{active} neutrophil population and was likely responsible for driving expression of interferon response elements (IFIT1, ISG15, IFI6) and antiviral signaling, such as RSAD2 and STAT1 (Fig 2c; Online Atlas). A hallmark of PG^{active} neutrophil polarization was the activation of an E2F4 pathway (Fig 2d), while neutrophil programming during bacterial ARDS included STAT5B (Fig 2e). To summarize, in response to COVID-19, neutrophils were polarized by unique transcriptional regulation towards one of two main populations, either an IFN^{active} population or a PG^{active} population (Fig 2f).

Figure 2 – Distinct regulatory programs drive divergent neutrophil maturation. a. Consensus neutrophil DEGs upregulated (positive FC) or suppressed (negative FC) during COVID-19 in at least 3 of 8 patients at t1 relative to Bacterial ARDS. **b.** Differentially activated consensus transcription factors

(TFs) in neutrophils from patients with COVID-19 relative to bacterial ARDS at t1. Stacked bars depict
logFC contributions of each COVID-19 patient. **c-e.** Gene-regulatory networks preferentially driving
IFN^{active} (PRDM1, c), PG^{active} (E2F4, d), and bacterial ARDS-enriched (STAT5B, e) neutrophil states.
Scale bars depict kernel density estimates approximating magnitude of TF activation inferred by
SCENIC-calculated AUCell scores. **f.** Schematic summarizing neutrophil fates favoured during COVID-
19 versus bacterial ARDS.

Dexamethasone alters immune cell dynamics and plasma proteomic milieu.

Conventional therapeutics have limited efficacy for COVID-19, and while dexamethasone offers a

moderate benefit, the RECOVERY trial reported the benefit was greatest in the most severely affected

patients[27]. However, the mechanisms underlying this benefit are unclear and not universal, so opportunity

exists to optimize or better target this therapy. In our cohort, median time between dexamethasone

administration to t1 blood draw (within 72 hours of ICU admission) was 31 hours (Fig 3a, Extended

Figure 5a, Extended Table 1). Global differences in transcription were apparent at t1 with clear

upregulation of genes in neutrophils and some T cell subsets in COVID-19 patients treated with

dexamethasone versus those that were not treated (Fig 3b-d, Extended Figure 5b, Extended Data Table 4).

Dexamethasone globally downregulated genes at t1, including in naïve B cells, plasmablasts and some T

cells (Extended Figure 5b-d). At t2 gene upregulation occurred in adaptive immune cells, including naïve

and effector CD8 T cells, with limited alterations in the innate myeloid cell lineages including

neutrophils. However, neutrophils demonstrated clear down regulation of genes at t2, as did CD4 naïve

and central memory T cells (Extended Figure 5e, f). Proportionally, at t1, dexamethasone administration

was associated with an increase in cytotoxic CD4 T cells, naïve B cells, plasmablasts, and decreased

proliferating NK cells, and CD4 effector memory cells (Extended Fig 5g). By t2, dexamethasone was

associated with suppressed neutrophil proportions in circulation compared to untreated controls (13% vs

41%, Extended Fig 5g). Plasma proteomics from the same cohort revealed that dexamethasone

suppressed 10 host proteins (S100A8, S100A9, SERPINA1, SERPINA3, ORM1, LBP, VWF, PIGR,

AZGP1, CRP) that others have previously identified as biomarkers distinguishing severe COVID-19

cases from mild to moderate counterparts (full host proteome quarriable via Online Atlas; Extended Table

2)[28-31]. Suppression of calprotectin (S100A8/S100A9) and neutrophil serine proteases (SERPINA1 and

SERPINA3), paired with depletion of neutrophil proportions, implicates the modulation of neutrophil-

related inflammatory processes as a method of action for dexamethasone treatment.

Figure 3 – Dexamethasone suppresses IFN programs and depletes IFN[active] neutrophils in COVID-19. a. Schematic summarizing COVID-19 patients treated with or without dexamethasone profiled at t1 and t2. **b.** UMAP projection of 80,994 whole blood cells from 21 patient samples, coloured by Azimuth reference-mapped immune cell states. **c-d.** Kernel density estimates depicting magnitude of molecular response elicited by immune cell subsets following Dexamethasone treatment t1 (c) and t2 (d) calculated by summing DEG fold changes for each cell state shown in Panel A. **e.** Neutrophil states overlaid on a UMAP of 23,193 subclustered neutrophils from Dexamethasone- and non-Dexamethasone-treated COVID-19 patients, colored by cluster ID. **f.** Magnitude of molecular response elicited by each neutrophil state post-Dexamethasone treatment calculated by summing DEG fold changes for each cell state shown in Panel d. **g.** RNA velocity vector length (indicating rate of differentiation/state transition) in Dexamethasone- and non-Dexamethasone-treated neutrophils at t1 and t2. **h.** Consensus neutrophil DEGs upregulated (positive FC) or suppressed (negative FC) post-Dexamethasone in at least 3 of 6 COVID-19 patients at t1 relative to non-Dexamethasone COVID-19 controls. Stacked bars depict logFC contribution of each Dexamethasone-treated patient. **i-j.** Differential splicing kinetics drives activation of IL1R2 (i) and suppression of IFITM1 expression (j) post-Dexamethasone treatment. Phase portraits show equilibrium slopes of spliced–unspliced mRNA ratios. Green denotes most upregulated and red denotes most down regulated differentially expressed genes with COVID-19 (f).

Dexamethasone therapy restrains neutrophil IFN programs

Due to the early and sustained effects of dexamethasone on gene expression in neutrophils, the effects of

dexamethasone therapy on neutrophil functional states were evaluated. Neutrophil reclustering again

identified immature neutrophils at the apex of the maturation trajectory, accelerating and exhibiting

maximal divergence prior to PG[active] and IFN[active] state commitments (Fig. 3 d, Extended Fig 6a-e).

Interestingly, we also identified IL7R[+ve] neutrophils (comprising roughly 8% of total neutrophils) whose

trajectories remained completely separate (Fig. 3 d, Extended Fig 6g, j) suggesting an entirely distinct

neutrophil state. Initially, dexamethasone was associated with increased global transcription in PG[active]

neutrophils, while ongoing therapy resulted in the emergence of a PG[active] neutrophils concomitant with

high IL1R2 expression (IL1R2[+ve]) (Fig. 3 e). Conversely, dexamethasone had a pronounced attenuation of

global transcription of IFN[active] neutrophils at t1 and t2 (Fig 3 e, f). Remarkably, dexamethasone

administration at t1 halted dynamic state changes in IFN[active] and IL7R[+ve] neutrophils, followed by

preferential depletion of IFN[active] subsets (Fig 3 g). Indeed, dexamethasone was associated with a

reduction in IFN[active] neutrophils to a proportion more similar to that detected in healthy controls (9%

post-Dex at t2 versus 10% in healthy controls) (Fig. 4a, Extended Fig 4h-k). Although collection of

airway samples (i.e. bronchoalveolar lavage fluid; BALF) was not feasible at our institution, we leveraged

two recent BALF scRNA-Seq datasets[11,32] to assess whether IFN^{active} neutrophils dominate the

bronchoalveolar landscape during severe COVID-19. Projection of CSF3R$^+$S100A8$^+$S100A9$^+$ BALF

neutrophils onto our reference revealed: a. 1.5 FC expansion of IFN^{active} neutrophils in severe COVID-19

relative to moderate disease (77% vs 52%, Extended Fig 7a-b), b. preferential activation of IFN-

stimulated genes such as IFITM1, IFITM2, IFI6, IRF7, and ISG20 in severe COVID-19 neutrophils

(Extended Fig 7c), and c. 4.7 FC greater IFN^{active} neutrophils in COVID-19 relative to bacterial

pneumonia patients (14% vs 3%, Extended Fig 7d-f). Albeit anecdotal, in our whole blood cohort, the

IFN^{active} neutrophil state was dominant in patient S7 [32], an 80-year-old male with remarkably high viral

titers who succumbed to COVID-19 complications within 3-4 days of sampling (Extended Fig 7f).

Consensus DEG analysis highlighted that upregulation of IL1R2, a decoy receptor that sequesters IL-1,

and downregulation of IFITM1 were the most prominent discriminating features of treatment with

steroids (Fig. 3h). Additionally, dexamethasone attenuated neutrophil expression of IFN pathways more

broadly, including the reduction of IFITM1-3, IFIT1, ISG15 and RSAD2 (Fig 3h). Examination of

unspliced pre-mRNA to mature spliced mRNA ratios supported the notion that induction of

immunoregulatory systems (i.e., IL-1R2; Fig 3 i) and suppression of IFN (i.e., IFITM1; Fig 3 j) programs

were driven by differential splicing kinetics.

Dexamethasone therapy intensifies neutrophil immunosuppressive function

Corticosteroid therapy shifted neutrophil state compositions. While IFN^{active} neutrophils were significantly

depleted by seven days of therapy, there was >2-fold expansion in immature neutrophils relative to non-

treated COVID-19 controls (Fig 4a; Extended Fig 6 h, i), which were absent in the healthy controls.

Albeit anecdotal, the dominance of IFN^{active} neutrophils at t1 in the patient who succumbed to COVID-19

in the non-dexamethasone cohort further supports depletion of IFN^{active} neutrophils as a mechanism by

which dexamethasone is protective (Extended Fig 8 g-j). Assessment of gene regulatory networks

demonstrated that IRF7 and MEF2A exhibited opposing activation patterns, with IRF7 being the most

suppressed and MEF2A the most enhanced transcription factors identified with dexamethasone, which

correlates with the emergence of PG^{active} and IL1R2^{+ve} states and attenuation of the IFN^{active} neutrophil

states (Fig 4b, Extended Fig 6k-m). To assess the generalizability of the dexamethasone regulated DEGs

identified in our cohort, we asked whether they accurately predicted mortality due to COVID-19 in a

larger validation cohort. By leveraging a whole blood bulk RNA-Seq dataset from 103 COVID-19

patients[33][34], we scored each sample by the aggregated expression of dexamethasone suppressed DEGs at

t1 and t2 (Extended Data Table 3). Interestingly, suppressed DEGs at t2 (but not t1) proved to be a far

superior predictor of 28-day mortality (AUC: 0.78, CI: 0.67 -0.89) compared to clinical severity scales

such as sequential organ failure assessment (SOFA) (AUC: 0.67, CI: 0.51-0.82) across all classification

thresholds (Fig 4c).

Unexpectedly, steroid administration was associated with an increase in circulating immature neutrophils,

which highly expressed TOP2A, and activated ATF4 and JDP2, transcription factors seen in

undifferentiated cells or those undergoing nuclear reprogramming (Extended Fig 6h). Interestingly, these

immature neutrophils expressed high levels of ARG1, ANXA1 (Fig 4d), and CD24 (both mRNA and

protein; Extended Fig 6 i), also suggesting an immunomodulatory role[35,36,37-39] that was expanded with

dexamethasone treatment. Both ARG1 and ANXA1 express glucocorticoid response elements, supporting

direct regulation by dexamethasone treatment[40,41].

To further understand the role of neutrophils during COVID-19 and the effects of dexamethasone, we

investigated cellular connectomes. Cellular interactions between many cell types (including highly

interactive neutrophils) were noted (Extended Fig 8a), and dexamethasone altered the globally predicted

interactions by suppressing intercellular signalling, in both number and strength of interactions (Extended

Fig 8b, c). Dexamethasone enhanced (Fig 4e) and suppressed (Extended Fig 8d) a number of unique

neutrophil-driven signalling networks. Of note, annexin family signalling, which was enhanced in the

immature neutrophils and represent powerful immunomodulators, were augmented between neutrophils

and the other circulating immune cells when patients received dexamethasone (Fig 4e). Of note is the

direction of annexin family signaling, which switched from incoming toward neutrophils without

dexamethasone treatment to being almost entirely outgoing from neutrophils toward B intermediate and

memory cells and MAIT cells following dexamethasone (Fig 4f, g, Extended Fig 8e, f). Therefore,

dexamethasone directly altered neutrophil functional states, by promoting expansion of an

ARG1+/ANXA1+ immature state with immunosuppressive features and altered the global

communication structure such that neutrophils became active instructors of some peripheral immune

cells.

Neutrophil response to dexamethasone is sexually dimorphic

Given the apparent clinical benefit of dexamethasone is more evident in males[27], and since males are

predisposed to more severe COVID-19 presentations and outcomes[42] , we surmised that dexamethasone

incites sexually dimorphic immunosuppressive effects. Our retrospective province-wide audit comparing

72 pre-dexamethasone (51 M, 21 F) versus 1,581 post-dexamethasone (1013 M, 568 F) treated ICU-

admitted patients confirmed a preferential mortality benefit in male COVID-19 patients (Extended Fig 9a,

b). While dexamethasone modulated 525 neutrophil DEGs across both sexes, while 892 were uniquely

modulated in either males or females (Extended Data Table 5). Amongst the jointly modulated DEGs, a

subset (24 of 525) exhibited statistically significant dimorphism in either magnitude or direction of

regulation (Extended Fig 9c, d). Interestingly, while neutrophils were depleted in both sexes post-

dexamethasone, this was particularly pronounced in males (1.9 FC higher in males at t1 and 3.4 FC

higher in males at t2, Extended Fig 9e). Of the two salient neutrophil state alterations, an immature

(ARG1^{+ve} immunosuppressive) state was preferentially expanded with dexamethasone in males (Extended

Fig. 9e), whereas ISGs were preferentially suppressed (Extended Fig. 9f) and IFN^{active} states were

depleted in females (Extended Fig. 9g-h) at both t1 and t2 (Fig 4h, i). Sexually dimorphic effects of

dexamethasone on neutrophil maturation kinetics may in part explain these state alterations. Dynamo-

reconstructed vector dynamics revealed that dexamethasone slowed IFN^{active} transitions (Extended Fig. 9i)

whilst accelerating immature (ARG1^{+ve} immunosuppressive) neutrophil differentiation in females

(Extended Fig. 9j) ultimately leading to a diminished immature neutrophil progenitor pool.

Figure 4 – Dexamethasone expands immunosuppressive neutrophils and their interactions in COVID-19. a. Neutrophil states mapped onto Louvain-clustered UMAP, with comparison of neutrophil composition between dexamethasone- and non-dexamethasone-treated samples at t1 and t2. **b.** Consensus

TFs activated or suppressed post-dexamethasone in at least 3 of 6 patients at t1 and predicted activity of
MEF2A and IRF7, two of the most differentially regulated TFs post-dexamethasone. **c.** Receiver
operating characteristic (ROC) curves assessing the discriminatory capacity of dexamethasone suppressed
DEGs at t1, t2, and sequential organ failure assessment (SOFA) scores for predicting 28-day mortality in
a validation cohort of 103 bulk whole blood RNA-Seq samples where 17 cases were fatal. **d.** Immature
and IL1R2[+ve] neutrophil subsets express high levels of immunosuppressive neutrophil marker ARG1 and
ANXA1. **e.** Neutrophil-driven signaling pathways induced post-dexamethasone, identified using CellChat
(MHC-I signalling filtered out). **f, g**. Topology of annexin signalling without (e) and with dexamethasone
(f) treatment (edges filtered to those where neutrophils function as senders or recipients of annexin
signals). **h.** Neutrophil state composition separated by sex and dexamethasone status at t1 and t2. **i.**
Schematic summarizing the effects of dexamethasone on neutrophil fates and function in COVID-19
following dexamethasone treatment.

Conclusions

Surviving SARS-CoV-2 infection depends on striking a temporal balance between inciting viral clearance

immune programs during the early stage and subsequently restraining those same programs at later stages

to limit immunity-induced tissue damage. IFN signaling stands at the nexus between antiviral immunity

and over active effector immune programs that inadvertently compromise tissue function and threaten

survival[43]. Our work uncovered downstream IFN signalling as a signature of a stable neutrophil state that

is selectively expanded during late stage COVID-19 infection from a common pool of immature

progenitors. Given that inborn errors [25] and suppressed *early stage* [6] IFN signalling predicts COVID-19

severity, increased IFN[active] neutrophils in females correlated with decreased mortality[44], and early

initiation of IFN therapy has been suggested to mitigate disease severity [45,46], one may posit that IFN

activity in neutrophils represents a concerted host antiviral program.

Interestingly, immunosuppression with dexamethasone, a corticosteroid known to improve mortality in

hospitalized COVID-19 patients[27], was associated with suppressed COVID19-specific IFN regulatory

networks and depleted COVID19-enriched IFN[active] neutrophils in favour of expanding immature (ARG1[+]

immunosuppressive) neutrophils. These altered neutrophil states shared striking resemblances to bacterial

ARDS, suggesting installation of generalized microbicidal programs ameliorate the overzealous

neutrophil responses during COVID-19 (and perhaps during other viral infections). While neutrophil ISG

activation may promote anti-viral immunity during early stages of SARS-CoV-2 infection, sustained IFN

activation during *late stages* (e.g., critically ill patients requiring intensive care) could drive

immunopathology of COVID-19. Indeed, positive correlation between neutrophil Type 1 IFN programs

and COVID-19 severity[7,47] paired with our observation that IFN^{active} neutrophils dominate the

bronchoalveolar microenvironment during severe COVID-19 [11]directly support this view.

Immunotherapies that support the innate antiviral immune response by decoupling IFN-exaggerated

neutrophil response whilst reinforcing acquisition of suppressor states may limit the pathogenic potential

of neutrophils and provide tremendous clinical benefit for treating severe COVID-19.

There are three major limitations of our study. First, <u>non-random group allocation (since the timing of the</u>

<u>RECOVERY trial made dexamethasone standard of care overnight) and small sample size may</u>

<u>inadvertently introduce selection bias and limit generalizability of dexamethasone findings</u>. Second,

comparisons were against bacterial ARDS, and not related respiratory viral infections (i.e., H1N1

influenza) since public health measures eradicated such cases; this precludes assessment of whether the

dynamics defined are specific to SARS-CoV-2. Finally, a subset of patients sampled at t1 were

discharged from ICU prior to t2 collection (non-random or non-ignorable missing data), precluding

unbiased estimation of temporal changes between timepoints.

Methods

Patient enrolment. All patients were enrolled following admission to any of the four adult intensive care

units at South Health Campus, Rockyview General Hospital, Foothills Medical Center or Peter Lougheed

Center in Calgary, Alberta, Canada (Extended Fig 1). Patient admission to the ICU was determined by the

attending ICU physician based on the need for life sustaining interventions, monitoring and life-support.

The research teams did not participate in clinical decisions. Study inclusion required a minimal age of 18,

the ability to provide consent, or for most participants, the ability of a surrogate decision maker to provide

regained capacity consent. All participants required an arterial catheter for blood draws, but the insertion

of this catheter was at the discretion of the attending medical team. Participants required a positive

clinical RNA COVID-19 test prior to enrolment, and evidence of bilateral lung infiltrates and hypoxemia

consistent with ARDS. At the time of sample collections, all COVID-19⁺ enrolled individuals were

culture negative for concurrent bacterial infections in the blood, urine, and sputum. The bacterial ARDS

cohort required a negative COVID-19 test and a definitive microbiological diagnosis of bacterial

pneumonia with chest imaging consistent with a diagnosis of ARDS. Patients were excluded from our

study if they: 1. were on immunosuppressive therapies, 2. had established autoimmune disease, or 3. had

active malignancy. Since tocilizumab or other immunomodulatory agents were not approved for use in

patients with severe COVID-19 in Alberta over the timespan of this study, none of them received these

medications. While bacterial sepsis patients received appropriate antibiotic treatments, none were

prescribed immunosuppressive or steroid therapy. All bacterial sepsis patients had lung infections caused

by gram-positive cocci (4 *Staphylococcus aureus* and 2 *Streptococcus pneumoniae*). Participants were

required to have a definitive diagnosis and appropriate consent and samples collected within 72hrs of

admission to the ICU in order to be included. Timepoint 1 (T1) refers to the first blood draw, while T2

was a repeat blood draw taken 7 days after T1, if the participant remained in the ICU, and had an arterial

catheter. For each participant, whole blood was collected via the arterial catheter and immediately

processed for analysis. Healthy blood donors were recruited by university-wide advertisement and

required that participants were: 1. not on immunomodulatory medications, 2. were asymptomatic for

SARS-CoV-2, 3. did not receive vaccination against SARS-CoV-2, and 4. did not have underlying

immune disorders.

Epidemiological analysis. We used the Alberta provincial eCRITICAL oracle-based analytics database

(Tracer) to query and extract Alberta COVID-19 ICU cases and volumes for this study[48]. Aggregate data

from sixteen individual adult ICUs was obtained over the study periods. The administration of

dexamethasone was not possible to capture at an aggregate level; therefore, we queried the database for

patients admitted to ICU prior to dexamethasone becoming standard of care in our Province (pre-

dexamethasone era; January 2020 till May 31ˢᵗ, 2020) versus dexamethasone as standard of care for

severe COVID-19 (June 1ˢᵗ, 2020, till May 31ˢᵗ, 2021). Tocilizumab was approved for use in Alberta

March 11 2021, and a small supply (150 doses) was obtained for severe COVID-19 patients after this

date.

Human Study Ethics. All work with humans was approved by the Conjoint Health Research Ethics

Board (CHREB) at the University of Calgary (Ethics ID: REB20-0481) and is consistent with the

Declaration of Helsinki.

Serum cytokine assessment. Cytokines, chemokines and soluble cytokine receptors were quantitated on

multiplex arrays that included a 65 MIlliPLEX cytokine/chemokine (6Ckine, BCA-1, CTACK, EGF,

ENA-78, Eotaxin, Eotaxin-2, Eotaxin-3, FGF-2, Flt-3L, Fractalkine, G-CSF, GM-CSF, GRO, I-309,

IFNα2, IFNγ, IL-1α, IL-1β, IL-1ra, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12 (p40), IL-

12 (p70), IL-13, IL-15, IL-16, IL-17A, IL-18, IL-20, IL-21, IL-23, IL-28a, IL-33, IP-10, LIF, MCP-1,

MCP-2, MCP-3, MCP-4, MDC, MIP-1α, MIP-1β, MIP-1d, PDGF-AA, PDGF-AB/BB, RANTES, SDF-1

a+b, sCD40L, SCF,TARC, TGFa, TNFa, TNFb, TPO, TRAIL, TSLP, VEGF) and a 14 MilliPLEX

soluble cytokine (sCD30, sEGFR, sgp130, sIL-1RI, sIL-1RII, sIL-2Ra, sIL-4R, sIL-6R, sRAGE, sTNF

RI, sTNF RII, sVEGF R1, sVEGF R2, sVEGF R3) arrays (Millipore Sigma, Oakville, ON, Canada) on a

Luminex Model 200 Luminometer (Luminex Corporation, Austin, TX). EDTA-plasma samples were

collected from each patient by venipuncture following a standard operating protocol (SOP) and stored at -

80C until tested. Each run included a full range of calibrators. The Mann-Whitney U test was used to

compare groups and p-values were adjusted for multiple comparisons using Holm-Šídák stepdown

method with alpha set to 0.05.

Shotgun proteomics using Liquid Chromatography and Mass Spectrometry (LC-MS/MS)

The serum of COVID-19 patients (COVID-19 = 9, dexamethasone-treated = 4) and bacterial ARDS

controls (N = 6) were collected and subjected to quantitative proteomics. The total protein concentrations

were determined by Pierce™ BCA Protein Assay Kit (23225, ThermoFisher). A trichloroacetic acid

(TCA)/acetone protocol was used to pellet 100μg of proteins per sample. Samples were subjected to a

quantitative proteomics workflow as per supplier (Thermo Fisher) recommendations. Samples were

reduced in 200mM tris(2-carboxyethyl)phosphine (TCEP), for 1h at 55°C, reduced cysteines were

alkylated by incubation with iodoacetamide solution (50mM) for 20min at room temperature. Samples

were precipitated by acetone/methanol, and 600μL ice-cold acetone was added followed by incubation at

-20°C overnight. A protein pellet was obtained by centrifugation ($8,000g$, 10min, 4°C) followed by

acetone drying (2min). Precipitated pellet was resuspended in100 μL of 50mM triethylammonium

bicarbonate (TEAB) buffer followed by tryptase digestion (5μg trypsin per 100μg of protein) overnight at

37°C. TMT-6plex™ Isobaric Labeling Reagents (90061, Thermo Fisher) were resuspended in anhydrous

acetonitrile and added to each sample (41μL TMT-6plex™ per 100μL sample) and incubated at room

temperature for 1h. The TMT labeling reaction was quenched by 2.5% hydroxylamine for 15min at room

temperature. TMT labeled samples were combined and acidified in 100% trifluoroacetic acid to pH < 3.0

and subjected to C18 chromatography (Sep-Pak) according to manufacturer recommendations. Samples

were stored at -80°C before lyophilization, followed by resuspension in 1% formic acid before liquid

chromatography and tandem mass spectrometry analysis.

Tryptic peptides were analyzed on an Orbitrap Fusion Lumos Tribrid mass spectrometer (Thermo

Scientific) operated with Xcalibur (version 4.0.21.10) and coupled to a Thermo Scientific Easy-nLC

(nanoflow liquid chromatography) 1200 System. Tryptic peptides (2μg) were loaded onto a C18 trap

(75μm x 2cm; Acclaim PepMap 100, P/N 164946; ThermoFisher) at a flow rate of 2μL/min of solvent A

(0.1% formic acid in LC-MS grade H2O). Peptides were eluted using a 120min gradient from 5 to 40%

(5% to 28% in 105min followed by an increase to 40% B in 15min) of solvent B (0.1% formic acid in

80% LC-MS grade acetonitrile) at a flow rate of 0.3μL/min and separated on a C18 analytical column

(75μm x 50cm; PepMap RSLC C18; P/N ES803A; ThermoScientific). Peptides were then electrosprayed

using 2.1kV voltage into the ion transfer tube (300°C) of the Orbitrap Lumos operating in positive mode.

For LC-MS/MS measurements with the FAIMS Pro (Thermo Fisher Scientific), multiple compensation

voltages (CV) were applied, -40V, -60V, and -80V with a cycle time of 1 second. FAIMS was used to

generate technical replicates from plex 1 to 6. The Orbitrap first performed a full MS scan at a resolution

of 120,000 FWHM to detect the precursor ion having a m/z between 375 and 1,575 and a +2 to +4 charge.

The Orbitrap AGC (Auto Gain Control) and the maximum injection time were set at 4 x 10^5 and 50ms,

respectively. The Orbitrap was operated using the top speed mode with a 3 second cycle time for

precursor selection. The most intense precursor ions presenting a peptidic isotopic profile and having an

intensity threshold of at least 2 x 10^4 were isolated using the quadrupole (Isolation window (m/z) of 0.7)

and fragmented using HCD (38% collision energy) in the ion routing multipole. The fragment ions (MS2)

were analyzed in the Orbitrap at a resolution of 15,000. The AGC and the maximum injection time were

set at 1 x 10^5 and 105ms, respectively. The first mass for the MS2 was set at 100 to acquire the TMT

reporter ions. Dynamic exclusion was enabled for 45 seconds to avoid of the acquisition of same

precursor ion having a similar m/z (plus or minus 10ppm).

Proteomic data and bioinformatics analysis

Spectral data acquired from the mass spectrometer were matched to peptide sequences using MaxQuant

software (v.1.6.14)[49]. Due to a lack of direct compatibility with Maxquant, spectra generated using the

FAIMS pro was first converted to MzXML using the FAIMS MzXML Generator from the Coon's lab

(https://github.com/coongroup/FAIMS-MzXML-Generator). Next, peptide sequences from the human

proteome and Sars-CoV-2 proteins were obtained from the UniProt database (May 2021) and matched

using the Andromeda[50] algorithm at a peptide-spectrum match false discovery rate (FDR) of 0.05. Search

parameters included a mass tolerance of 20 p.p.m. for the parent ion, 0.5 Da for the fragment ion,

carbamidomethylation of cysteine residues (+57.021464 Da), variable N-terminal modification by

acetylation (+42.010565 Da), and variable methionine oxidation (+15.994915 Da). Relative quantification

was set as TMT 6-plex labels 126 to 131. The cleavage site specificity was set to Trypsin/P, with up to

two missed cleavages allowed. Next, the evidence.txt and proteinGroups.txt were loaded into the R

software (v4.0.2) for statistical analysis. The normalization and identification of differentially expressed

proteins was performed using the MSstatsTMT package[51]. Multiple comparisons were corrected using the

Benjamini-Hochberg approach.

Leukocyte and lymphocyte isolation. For lymphocyte isolation, whole blood (2mL) was collected in

5mL polystyrene round-bottom heparinized vacutubes. To isolate lymphocytes by immunomagnetic

negative selection, 100μL of Isolation Cocktail and 100μL of Rapid Spheres (EasySep[TM] Direct Human

Total Lymphocytes Isolation Kit: 19655, StemCell Technologies) were added to 2 mL of whole blood.

After mixing and 5min incubation at RT, the sample volumes were topped up to 2.5mL with 0.04%

bovine serum albumin (BSA) in PBS. The diluted sample was incubated in the magnet without lid for

5min, at RT and negatively selected lymphocytes were decanted into a new 5 mL polystyrene tube.

Except the addition of Isolation Cocktail, all steps were repeated once. The final lymphocyte cell

suspension was transferred to a 15 mL polypropylene tube and a volume of 5mL 0.04% BSA in PBS was

added to the sample. Lymphocytes were precipitated by centrifugation for 5 min at 2000rpm, supernatant

was discarded, and cells were resuspended in 5 mL of 0.04% BSA in PBS. This last step was repeated

once, and cells were then resuspended in 100 μL of PBS+0.04% BSA. Cell density was quantified with a

hemacytometer, cell viability was assessed with Trypan Blue staining (T8154; Sigma Aldrich), and 7500

live lymphocytes were transferred to a sterile 1.5 mL microcentrifuge tube.

For leukocyte isolation, 1 mL of whole blood from heparin containing vacutubes was transferred to 5 mL

polystyrene round-bottom tubes and 12μL of 0.5M EDTA was added. 2% FBS in PBS (1mL) and 50μL

of EasySep RBC Depletion spheres (EasySep[TM] RBC Depletion Reagent: 18170, Stem Cell

Technologies) were added to immunomagnetically deplete red blood cells. After 5 min of magnet

incubation at RT, cell suspension containing leukocytes was decanted into a new 5mL polystyrene tube.

To ensure complete removal of red blood cells, RBC depletion was repeated, and cell suspension

containing leukocytes was decanted into a new 15mL polypropylene tube. Leukocytes were precipitated

by centrifugation at 2000rpm for 5 min at 20°C and resuspended in 5mL of 0.04% BSA in PBS. This last

step was repeated once, and leukocytes were resuspended in 2 mL of 0.04% BSA in PBS. Cell viability

and cell density were assessed, and 7500 live leukocytes were transferred to the microcentrifuge tube

containing the lymphocyte cell suspension. The volume of the cell suspension containing 7500

lymphocytes and 7500 leukocytes in a total of 50 μL of 0.04% BSA in PBS.

Immunocytochemistry and immunohistochemistry

Isolated leukocyte and lymphocyte samples were fixed in 4% paraformaldahyde in PBS (0.2mM and

pH7.4), and spun in a cytocentrifuge (8min at 300g) onto coated slides. Pathological lung sections (FFPE

fixed and sectioned at 5um) were deparaffinized in Slide Brite (Fisher Scientific NC968653) and

rehydrated. Slides were permeabilized and blocked with 10% normal donkey serum in PBS (with 0.5%

triton X-100), primary antibodies (S100A8/9 Abcam ab22506; IFITM1 Abcam ab233545) were

incubated at 4°C overnight, followed by incubation with donkey anti-rabbit-Alexa488 (Invitrogen

A32790) or anti-mouse-Alexa555 (Invitrogen A31570) for 1h at room temperature (RT). Cytospun slides

were sequentially stained with CD24 (Abcam ab202073) on the same slides for 1h at RT, followed by

donkey anti-rabbit-Alexa647 (Invitrogen A31573). Imaging was done using a VS-120 slide scanner

(Olympus) and high resolution image imaging was done using an SP8 spectral confocal microscope

(Leica). Image processing was completed in Fiji [52].

Single-cell RNA-Seq library construction, alignment, and quality control. A total of 15,000 single

cells (containing an equal proportion of leukocytes and lymphocytes) were loaded for partitioning using

10X Genomics NextGEM Gel Bead emulsions (Version 3.1). All samples were processed as per

manufacturer's protocol (with both PCR amplification steps run 12X). Quality control of resulting

libararies and quantification was performed using TapeStation D1000 ScreenTape assay (Agilent).

Sequencing was performed using Illumina NovaSeq S2 and SP 100 cycle dual lane flow cells over

multiple rounds to ensure each sample received approximately 32,000 reads per cell. Sequencing reads

were aligned using CellRanger 3.1.0 pipeline[53] to the standard pre-built GRCh38 reference genome.

Samples that passed alignment QC were aggregated into single datasets using CellRanger aggr with

between-sample normalization to ensure each sample received an equal number of mapped reads per cell.

Aggregated non-dexamethasone-treated COVID-19 (n = 12) and bacterial ARDS (n = 9) samples

recovered 1,872,659 cells that were sequenced to 38,410 post-normalization reads per cell. Likewise,

aggregated COVID-19 samples with (n = 9) or without (n = 12) dexamethasone recovered 1,748,551

single cells sequenced to 51,415 post-normalization reads per cell. Aggregated healthy samples recovered

19,816 cells, including 1,912 post-QC neutrophils (n = 5).

Single-cell RNA-Seq computational analyses and workflows. Filtered feature-barcode HDF5 matrices

from aggregated datasets were imported into the R package Seurat v.3.9 for normalization, scaling,

integration, multi-modal reference mapping, louvain clustering, dimensionality reduction, differential

expression analysis, and visualization [54]. Briefly, cells with abnormal transcriptional complexity (fewer

than 500 UMIs, greater than 25,000 UMIs, or greater than 25% of mitochondrial reads) were considered

artifacts and were removed from subsequent analysis. Since granulocytes have relatively low RNA

content (due to high levels of RNases), QC thresholds were informed by [8] as they recently defined several

rodent and human neutrophil subsets from scRNA-Seq samples. Cell identity was classified by mapping

single cell profiles to the recently published PBMC single-cell joint RNA/CITE-Seq multi-omic reference

[55].

Annotation of neutrophil states. Since no published reference automates granulocyte annotations,

neutrophil clusters were manually annotated by querying known markers (i.e. CSF3R, S100A8, S100A9,

MMP8, MMP9, ELANE, MPO)[56] and were corroborated using the R package SingleR[57]. Neutrophil

states were defined by grouping unsupervised (louvain at default resolution) subclusters based on two

overlapping criteria: scVelo-inferred neutrophil maturity, and 2. by corroborating gene expression and

SCENIC-inferred GRN signatures with previous human and rodent neutrophil scRNA-Seq studies.

Immature neutrophils were defined as $CD24^+ARG1^+ELANE^+MPO^+ATF4^{GRN\text{-}active}JDP2^{GRN\text{-}active}$

neutrophils [7,8,47,58] that were reproducibly assigned as 'root cells' in scVelo-based latent time pseudo-

ordering. IFN^{active} neutrophils were defined by preferential mRNA splicing (positive velocity) and

expression of IFN-stimulated genes such as IFITM1/2, IFIT1/2/3, ISG15/20, and IFI6/27/44/44L [6,44,59].

PG^{active} neutrophils were distinguished by preferential splicing of PTGS2/COX2 (as well as expression for

prostaglandin transport LST1) [44] and included a subset that expressed high levels of IL1β decoy receptor

IL1R2 [33]. Lastly, IL7R$^+$ neutrophils (a small but distinct subset that maybe of thymic origin [60] expressed

high levels of ribosomal subunit genes (e.g. RPL5/7A/8/13/18/19/23/24/27/P0) that are highly

reminiscent of 'ribosomal^{hi}-specific cluster 7' identified previously [47]..

Statistical approach for comparing cell proportions. To test whether cell composition was changed

due to infection type (COVID-19 versus Bacterial ARDS) or treatment group (dexamethasone versus

non-dexamethasone), a generalized linear mixed-effects model was employed where infection type and

treatment group were considered fixed and individual patients were considered random effect. Fitting was

done with Laplace approximation using the 'glmer' function in the 'lme4' R package [61] and p-values were

calculated using the R package 'car'. Boxplots comparing cell type composition were generated using the

ggplot2 package. Since a subset of patients sampled at t1 were discharged from ICU prior to t2 collection

(non-random or non-ignorable missing data), we limit statistical comparisons to between group

comparisons within one time point (e.g., COVID-19 72h vs Bacterial ARDS 72hr, dexamethasone-treated

72h vs non-dexamethasone-treated 72h) and do not estimate temporal differences across t1 and t2.

Inferring cell communication networks. Differential cell-cell interaction networks were reconstructed

using the Connectome R toolkit v0.2.2[62] and CellChat v1.0.0 [63]. Briefly, *DifferentialConnectome* queried

Seurat R objects housing datasets integrated by infection type and dexamethasone status to define nodes

and edges for downstream network analysis. Total number of interactions and interaction strengths were

calculated using CellChat's *compareInteractions* function. Differential edge list was passed through

CircosDiff (a wrapper around the R package 'circlize') and CellChat's *netVisual_chord_gene* to filter

receptor-ligand edges and generate Circos plots.

Consensus DEGs and perturbation scores. Differentially expressed genes (DEGs) were those with an

average log fold change (FC) greater than 0.25 (p-adjusted < 0.05) as determined by Seurat's Wilcoxon

rank-sum test. Consensus stacked bars showing cumulative log fold changes (colored by individual

sample contributions) were generated using *constructConsensus* function [7] for genes exhibiting

reproducible changes across patients (>3 for 72-hour comparisons, > 2 for 7-day comparisons). Gene Set

Enrichment analyses of consensus DEGs were performed using gProfiler's g:GOSt (p-value cutoff

<0.05). A cell state-specific 'perturbation score' was calculated to reflect the magnitude of response

elicited by factoring in number and cumulative FC of consensus DEGs. Perturbation scores were

visualized using Nebulosa-generated density plots [64].

Constructing cellular trajectories using RNA velocity. Analysis of neutrophil trajectories was

performed by realigning CellRanger count-generated BAMs with RNA velocity command-line tool [20]

using the *run10x* command and human (GRCh38) annotations. The output loom files containing spliced

and unspliced counts were combined to compare neutrophils in COVID-19 with Bacterial ARDS controls

and dexamethasone-treated with non-treated COVID-19 patients. For both analyses, combined looms

were imported into Seurat v.3.9 using the *ReadVelocity* function in SeuratWrappers v.0.2.0, normalized

using *SCTransform* v.0.3.2 [65], reduced and projected onto a UMAP, and exported as a .h5 file using the

SaveH5Seurat function. Counts stored in H5 files were imported, filtered, and normalized as

recommended in the scVelo v.0.2.1 workflow [19]. RNA velocities were estimated using stochastic and

dynamical models. Since both models yielded comparable results, stochastic model was used as default

for all subsequent analyses. Calculations stored in AnnData's metadata were exported as CSVs and kernel

density lines depicting Velocity-inferred latent time distribution were plotted with ggplot2 v.3.1.1.

Gene Regulatory Network reconstruction. Single-cell regulatory network inference and clustering

(SCENIC)[26] was employed to infer regulatory interactions between transcription factors (TFs) and their

targetome by calculating and pruning co-expression modules. Briefly, neutrophils were subsetted from

scVelo-realigned Seurat object and processed using default and recommended parameters specified in

SCENIC's vignette (https://github.com/aertslab/SCENIC) using the hg19 RcisTarget reference. Regulon

activity scores (in '3.4_regulonAUC.Rds', an output of the SCENIC workflow) were added to scVelo

object (using *CreateAssayObject* function) to jointly project trajectory and TF activity onto the same

UMAP embeddings. Consensus stacked bars showing cumulative logFC of AUCell scores for each TF

(colored by individual sample contributions) were generated by modifying the *constructConsensus*

function[7] for SCENIC assay. Targetome of TFs predicted as drivers of neutrophil states (stored in

'2.6_regulons_asGeneSet.Rds') was profiled using g:Profiler's functional enrichment analysis and genes

intersecting with the Interferon pathway were plotted using iRegulon (Cytoscape plugin)[66].

Comparing scRNA-Seq findings with published datasets. To test whether dexamethasone-suppressed

neutrophil genes at t1 and t2 (Extended Data Table 4) predicted COVID-19 mortality, we repurposed

methods described in [33] and employed whole blood bulk RNA-Seq datasets generated by [34] as a validation

cohort of 103 samples (where 17 were fatal). Briefly, each of the 103 samples were scored by the

aggregated expression of dexamethasone-suppressed neutrophil consensus genes at t1 and t2 using

Seurat's AddModuleScore(). Dexamethasone-suppressed module scores were used as the predictor

variable and 28-day mortality was used as the response variable to construct an ROC curve using pROC's

roc() function. To infer bronchoalveolar neutrophil composition in severe and moderate COVID-19 [11]and

across bacterial pneumonia and COVID-19 [32], neutrophils ($CSF3R^+$, $S100A8^+$, $S100A9^+$) captured in

BALF scRNA-Seq datasets were projected onto our peripheral blood reference using mutual nearest

neighbor anchoring (FindTransferAnchors) and identity transferring (TransferData and AddMetaData)

strategy implemented in Seurat v4 [54].

COVID Neutrophil Atlas. To enable intuitive exploration of single-cell datasets, a web portal

(http://biernaskielab.ca/covid_neutrophil or http://biernaskielab.com/covid_neutrophil) was built using

RShiny v1.1.0, shinyLP v.1.1.2, and shinythemes v.1.1.2 packages.

Data availability. Single cell RNA-Seq datasets are available at NCBI GEO (which automatically makes

SRA deposit) at the following accession: GSE157789. Single-cell datasets can be further explored on our

companion portal at http://biernaskielab.ca/COVID_neutrophil or

http://biernaskielab.com/COVID_neutrophil. Velocyto-generated LOOM files and processed R objects

are available for reanalysis from: http://doi.org/10.6084/m9.figshare.14330795. Whole blood bulk RNA-

Seq datasets employed as an independent validation cohort were downloaded from GSE157103. BALF

scRNA-Seq datasets from severe and moderate COVID-19 were downloaded from GSE145926.

Processed BALF scRNA-Seq objects from patients with bacterial pneumonia and COVID-19 (archived at

GSE167118) were downloaded from authors' archive: https://figshare.com/articles/dataset/_/13608734.

Mass spectrometry datasets will be available via PRIDE Archive (http://www.ebi.ac.uk/pride/archive), it

has been submitted (submission #: 1-20210702-114055) and is pending accessioning.

Proteomics data will be available at PRIDE (https://www.ebi.ac.uk/pride/), it has been submitted

(submission #: 1-20210702-114055) and is pending accessioning.

Code availability. All analyses were performed using publicly available software as described in the

methods section. Raw scripts are available upon request.

Supplementary Information is available for this paper.

Acknowledgements: This work was funded by a FastGrant from the Thistledown Foundation (JB and

BY) and Calgary Firefighters Burn Treatment Society (JB). S Sinha received CIHR Vanier, Alberta

Innovates, and Killam doctoral scholarships. E.L received an Alberta Children's Hospital Research

Institute postdoctoral fellowship. B.G.Y is a tier II Canada Research Chair in Pulmonary Immunology,

Inflammation and Host Defence. We acknowledge the assistance of the nurse practitioners, Charissa

Elton-Lacasse, Kirsten Deemer and Robert Ralph as well as the healthcare teams from the Calgary Adult

ICU's at South Health Campus, Rockyview General Hospital, Foothills Medical Center and Peter

Lougheed Center. We thank Dr. Kirsten Fiest and the ICU study coordinators Cassidy Codan, Zdenka

Slavikova and Olesya Dmitrieva. We acknowledge Dan Jones, Cathy Curr and the eCritical team (Alberta

Health Services in Alberta, Canada) for their help in data acquisition and extraction via eCritical

databases. Mortality predictions using dexamethasone-suppressed gene signatures were completed by

repurposing computational workflows kindly shared by Aaron Wilk and Dr. Catherine Blish (Stanford

University).

Author contributions: SS performed scRNAseq analyses, figure preparation, and co-wrote the paper.

NLR contributed to experimental design, performed scRNAseq experiments, figure preparation and co-

wrote the paper. AJ, RA, and LC performed bioinformatics and created the online atlas. EL, RF and APN

contributed to sample preparation and scRNAseq processing. MG and BM contributed to patient consent

and sample collection. LGA and AD conducted proteomics and related analyses. AB provided clinical

biospecimens. MJF provided serum cytokine assays. JB and BY conceived of all experiments,

experimental design, wrote the paper and supervised all experiments.

The authors have no competing interests.

References:

1   Matthay, M. A. *et al.* Acute respiratory distress syndrome. *Nat Rev Dis Primers* **5**, 18,
doi:10.1038/s41572-019-0069-0 (2019).
2   Wang, D. *et al.* Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel
Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA* **323**, 1061-1069,
doi:10.1001/jama.2020.1585 (2020).
3   Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med*
382, 727-733, doi:10.1056/NEJMoa2001017 (2020).
4   Middleton, E. A. *et al.* Neutrophil extracellular traps contribute to immunothrombosis in COVID-
19 acute respiratory distress syndrome. *Blood* **136**, 1169-1179, doi:10.1182/blood.2020007008
(2020).
5   Veras, F. P. *et al.* SARS-CoV-2-triggered neutrophil extracellular traps mediate COVID-19
pathology. *J Exp Med* **217**, doi:10.1084/jem.20201129 (2020).
6   Combes, A. J. *et al.* Global absence and targeting of protective immune states in severe COVID-
19. *Nature*, doi:10.1038/s41586-021-03234-7 (2021).
7   Wilk, A. J. *et al.* A single-cell atlas of the peripheral immune response in patients with severe
COVID-19. *Nat Med* **26**, 1070-1076, doi:10.1038/s41591-020-0944-y (2020).
8   Xie, X. *et al.* Single-cell transcriptome profiling reveals neutrophil heterogeneity in homeostasis
and infection. *Nature Immunology* **21**, 1119-1133, doi:10.1038/s41590-020-0736-z (2020).
9   Lee, J. S. *et al.* Immunophenotyping of COVID-19 and influenza highlights the role of type I
interferons in development of severe COVID-19. *Science Immunology* **5**, eabd1554,
doi:10.1126/sciimmunol.abd1554 (2020).
10   Xu, G. *et al.* The differential immune responses to COVID-19 in peripheral and lung revealed by
single-cell RNA sequencing. *Cell Discovery* **6**, 73, doi:10.1038/s41421-020-00225-2 (2020).
11   Liao, M. *et al.* Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19.
Nature Medicine **26**, 842-844, doi:10.1038/s41591-020-0901-9 (2020).
12   Wauters, E. *et al.* Discriminating mild from critical COVID-19 by innate and adaptive immune
single-cell profiling of bronchoalveolar lavages. *Cell Research* **31**, 272-290, doi:10.1038/s41422-
020-00455-9 (2021).
13   Rosin, N. L. *et al.* SARS-CoV-2 infection of circulating immune cells is not responsible for virus
dissemination in severe COVID-19 patients. *bioRxiv*, 2021.2001.2019.427282,
doi:10.1101/2021.01.19.427282 (2021).
14   Ranieri, V. M. *et al.* Acute respiratory distress syndrome: the Berlin Definition. *JAMA* **307**, 2526-
2533, doi:10.1001/jama.2012.5669 (2012).
15   Mehta, P. *et al.* COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet*
395, 1033-1034, doi:10.1016/s0140-6736(20)30628-0 (2020).
16   Feng, Z. *et al.* Early prediction of disease progression in COVID-19 pneumonia patients with
chest CT and clinical characteristics. *Nat Commun* **11**, 4968, doi:10.1038/s41467-020-18786-x
(2020).
17   Wu, C. *et al.* Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in
Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med*,
doi:10.1001/jamainternmed.2020.0994 (2020).
18   Barnes, B. J. *et al.* Targeting potential drivers of COVID-19: Neutrophil extracellular traps. *J Exp*
Med **217**, doi:10.1084/jem.20200652 (2020).
19   Bergen, V., Lange, M., Peidli, S., Wolf, F. A. & Theis, F. J. Generalizing RNA velocity to transient
cell states through dynamical modeling. *Nature Biotechnology* **38**, 1408-1414,
doi:10.1038/s41587-020-0591-3 (2020).

20   La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498, doi:10.1038/s41586-018-
0414-6 (2018).

21   Chen, J. S. *et al.* Non-steroidal anti-inflammatory drugs dampen the cytokine and antibody
response to SARS-CoV-2 infection. *Journal of Virology*, JVI.00014-00021, doi:10.1128/JVI.00014-
21 (2021).

22   Takashima, A. & Yao, Y. Neutrophil plasticity: acquisition of phenotype and functionality of
antigen-presenting cell. *J Leukoc Biol*, doi:10.1189/jlb.1MR1014-502R (2015).

23   Ledford, J. G., Kovarova, M. & Koller, B. H. Impaired Host Defense in Mice Lacking ONZIN. *The*
Journal of Immunology **178**, 5132, doi:10.4049/jimmunol.178.8.5132 (2007).

24   Zhang, Q. *et al.* in *Med (N Y)* Vol. 1   14-20 (© 2020 Elsevier Inc., 2020).

25   Zhang, Q. *et al.* Inborn errors of type I IFN immunity in patients with life-threatening COVID-19.
Science **370**, doi:10.1126/science.abd4570 (2020).

26   Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat Methods* **14**,
1083-1086, doi:10.1038/nmeth.4463 (2017).

27   Horby, P. *et al.* Dexamethasone in Hospitalized Patients with Covid-19 - Preliminary Report. *N*
Engl J Med, doi:10.1056/NEJMoa2021436 (2020).

28   Mahler, M., Meroni, P. L., Infantino, M., Buhler, K. A. & Fritzler, M. J. Circulating Calprotectin as a
Biomarker of COVID-19 Severity. *Expert Rev Clin Immunol* **17**, 431-443,
doi:10.1080/1744666X.2021.1905526 (2021).

29   Demichev, V. *et al.* A time-resolved proteomic and prognostic map of COVID-19. *Cell Syst*,
doi:10.1016/j.cels.2021.05.005 (2021).

30   Park, J. *et al.* In-depth blood proteome profiling analysis revealed distinct functional
characteristics of plasma proteins between severe and non-severe COVID-19 patients. *Sci Rep*
10, 22418, doi:10.1038/s41598-020-80120-8 (2020).

31   Shu, T. *et al.* Plasma Proteomics Identify Biomarkers and Pathogenesis of COVID-19. *Immunity*
53, 1108-1122 e1105, doi:10.1016/j.immuni.2020.10.008 (2020).

32   Zhao, Y. *et al.* Clonal expansion and activation of tissue-resident memory-like T$_H$17
cells expressing GM-CSF in the lungs of patients with severe COVID-19. *Science Immunology* **6**,
eabf6692, doi:10.1126/sciimmunol.abf6692 (2021).

33   Wilk, A. J. *et al.* Multi-omic profiling reveals widespread dysregulation of innate immunity and
hematopoiesis in COVID-19. *Journal of Experimental Medicine* **218**, doi:10.1084/jem.20210582
(2021).

34   Overmyer, K. A. *et al.* Large-Scale Multi-omic Analysis of COVID-19 Severity. *Cell Systems* **12**, 23-
40.e27, doi:https://doi.org/10.1016/j.cels.2020.10.003 (2021).

35   Vago, J. P. *et al.* Annexin A1 modulates natural and glucocorticoid-induced resolution of
inflammation by enhancing neutrophil apoptosis. *Journal of Leukocyte Biology* **92**, 249-258,
doi:https://doi.org/10.1189/jlb.0112008 (2012).

36   Oliveira, L. G. *et al.* Annexin A1 Is Involved in the Resolution of Inflammatory Responses during
Leishmania braziliensis Infection. *The Journal of Immunology* **198**, 3227,
doi:10.4049/jimmunol.1602028 (2017).

37   Uhel, F. *et al.* Early Expansion of Circulating Granulocytic Myeloid-derived Suppressor Cells
Predicts Development of Nosocomial Infections in Patients with Sepsis. *Am J Respir Crit Care*
Med **196**, 315-327, doi:10.1164/rccm.201606-1143OC (2017).

38   Arlauckas, S. P. *et al.* Arg1 expression defines immunosuppressive subsets of tumor-associated
macrophages. *Theranostics* **8**, 5842-5854, doi:10.7150/thno.26888 (2018).

39   Derakhshani, A. *et al.* Arginase 1 (Arg1) as an Up-Regulated Gene in COVID-19 Patients: A
Promising Marker in COVID-19 Immunopathy. *J Clin Med* **10**, doi:10.3390/jcm10051051 (2021).

40   Kelly-Scumpia, K. M. *et al.* ER Stress Regulates Immunosuppressive Function of Myeloid Derived
Suppressor Cells in Leprosy that Can Be Overcome in the Presence of IFN-γ. *iScience* **23**, 101050,
doi:https://doi.org/10.1016/j.isci.2020.101050 (2020).
41   Okun, J. G. *et al.* Molecular regulation of urea cycle function by the liver glucocorticoid receptor.
Mol Metab **4**, 732-740, doi:10.1016/j.molmet.2015.07.006 (2015).
42   Peckham, H. *et al.* Male sex identified by global COVID-19 meta-analysis as a risk factor for death
and ITU admission. *Nature Communications* **11**, 6317, doi:10.1038/s41467-020-19741-6 (2020).
43   Park, A. & Iwasaki, A. Type I and Type III Interferons - Induction, Signaling, Evasion, and
Application to Combat COVID-19. *Cell host & microbe* **27**, 870-878,
doi:10.1016/j.chom.2020.05.008 (2020).
44   Gupta, S. *et al.* Sex differences in neutrophil biology modulate response to type I interferons and
immunometabolism. *Proceedings of the National Academy of Sciences* **117**, 16481,
doi:10.1073/pnas.2003603117 (2020).
45   Monk, P. D. *et al.* Safety and efficacy of inhaled nebulised interferon beta-1a (SNG001) for
treatment of SARS-CoV-2 infection: a randomised, double-blind, placebo-controlled, phase 2
trial. *The Lancet Respiratory Medicine* **9**, 196-206, doi:10.1016/S2213-2600(20)30511-7 (2021).
46   Wang, N. *et al.* Retrospective Multicenter Cohort Study Shows Early Interferon Therapy Is
Associated with Favorable Clinical Responses in COVID-19 Patients. *Cell Host & Microbe* **28**, 455-
464.e452, doi:https://doi.org/10.1016/j.chom.2020.07.005 (2020).
47   Schulte-Schrepping, J. *et al.* Severe COVID-19 Is Marked by a Dysregulated Myeloid Cell
Compartment. *Cell* **182**, 1419-1440.e1423, doi:10.1016/j.cell.2020.08.001 (2020).
48   Brundin-Mather, R. *et al.* Secondary EMR data for quality improvement and research: A
comparison of manual and electronic data collection from an integrated critical care electronic
medical record system. *J Crit Care* **47**, 295-301, doi:10.1016/j.jcrc.2018.07.021 (2018).
49   Cox, J. & Mann, M. MaxQuant enables high peptide identification rates, individualized p.p.b.-
range mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* **26**, 1367-
1372, doi:10.1038/nbt.1511 (2008).
50   Cox, J. *et al.* Andromeda: a peptide search engine integrated into the MaxQuant environment. *J
Proteome Res* **10**, 1794-1805, doi:10.1021/pr101065j (2011).
51   Huang, T. *et al.* MSstatsTMT: Statistical Detection of Differentially Abundant Proteins in
Experiments with Isobaric Labeling and Multiple Mixtures. *Mol Cell Proteomics* **19**, 1706-1723,
doi:10.1074/mcp.RA120.002105 (2020).
52   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**,
676-682, doi:10.1038/nmeth.2019 (2012).
53   Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nature
Communications* **8**, 14049, doi:10.1038/ncomms14049 (2017).
54   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e1821,
doi:https://doi.org/10.1016/j.cell.2019.05.031 (2019).
55   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *bioRxiv*, 2020.2010.2012.335331,
doi:10.1101/2020.10.12.335331 (2020).
56   Zilionis, R. *et al.* Single-Cell Transcriptomics of Human and Mouse Lung Cancers Reveals
Conserved Myeloid Populations across Individuals and Species. *Immunity* **50**, 1317-1334.e1310,
doi:https://doi.org/10.1016/j.immuni.2019.03.009 (2019).
57   Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a transitional
profibrotic macrophage. *Nature Immunology* **20**, 163-172, doi:10.1038/s41590-018-0276-y
(2019).
58   Reusch, N. *et al.* Neutrophils in COVID-19. *Frontiers in immunology* **12**, 652470-652470,
doi:10.3389/fimmu.2021.652470 (2021).

59   Shaath, H., Vishnubalaji, R., Elkord, E. & Alajez, N. M. Single-Cell Transcriptome Analysis
Highlights a Role for Neutrophils and Inflammatory Macrophages in the Pathogenesis of Severe
COVID-19. *Cells* **9**, doi:10.3390/cells9112374 (2020).
60   Schlenner, S. M. *et al.* Fate Mapping Reveals Separate Origins of T Cells and Myeloid Lineages in
the Thymus. *Immunity* **32**, 426-436, doi:https://doi.org/10.1016/j.immuni.2010.03.005 (2010).
61   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4.
Journal of Statistical Software; Vol 1, Issue 1 (2015), doi:10.18637/jss.v067.i01 (2015).
62   Raredon, M. S. B. *et al.* Connectome: computation and visualization of cell-cell
signaling topologies in single-cell systems data. *bioRxiv*, 2021.2001.2021.427529,
doi:10.1101/2021.01.21.427529 (2021).
63   Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nature
Communications* **12**, 1088, doi:10.1038/s41467-021-21246-9 (2021).
64   Alquicira-Hernandez, J. & Powell, J. E. Nebulosa recovers single-cell gene expression signals by
kernel density estimation. *Bioinformatics*, doi:10.1093/bioinformatics/btab003 (2021).
65   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq data
using regularized negative binomial regression. *Genome Biology* **20**, 296, doi:10.1186/s13059-
019-1874-1 (2019).
66   Janky, R. s. *et al.* iRegulon: From a Gene List to a Gene Regulatory Network Using Large Motif
and Track Collections. *PLOS Computational Biology* **10**, e1003731,
doi:10.1371/journal.pcbi.1003731 (2014).