Genetic lineage of the Amami islanders inferred from classical genetic markers

Yuri Nishikawa and Takafumi Ishida

Department of Biological Sciences, Graduate School of Science, The University of

Tokyo, Tokyo, Japan

Correspondence: Yuri Nishikawa, Department of Biological Sciences, Graduate

School of Science, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo

113-0033, Japan.

E-mail address: yuri.nishikawa@gmail.com

Abstract

The genetic structure of the people of mainland Japan and Okinawa has

been gradually unveiled in recent years. However, previous anthropological

studies dealing with people in the Amami islands, located between mainland

Japan and Okinawa, were less informative because of the lack of genetic data. In

this study, we collected DNAs from 104 subjects in two of the Amami islands,

Amami-Oshima island and Kikai island. We analyzed the D-loop region of mtDNA,

four Y-STRs, and four autosomal nonsynonymous SNPs to clarify the Amami

islanders' genetic structure compared with peoples in Okinawa, mainland Japan,

and other regions of East Asia. We found that the Amami islanders showed a

genetically intermediate position between mainland Japan and Okinawa in

mtDNA and Y-STR. However, the frequencies of several autosomal SNPs in the

Amami islanders indicated a significant difference from mainland Japanese,

which may be because of the gene flow from Okinawa but not natural selection.

Moreover, extremely high or low frequencies of several alleles implied a founder

effect in Kikai islanders. Note that there is room for the interpretation of the

results because of the small sample size and number of alleles in the present study.

Geographically broad and detailed samplings and genome-wide analyses are

awaited.

Keywords: Japanese, Amami islanders, Classical genetic markers

1. Introduction

The genetic structure of the people of Japan has been gradually unveiled

in recent years. The dual structure model (Hanihara, 1991) is a well-known

hypothesis that predicts that the modern Japanese population was formed by the

admixture of two populations, Jomon and Yayoi; Jomon peoples dwelled in the

Japanese Archipelago since about 18,000 years ago, and Yayoi peoples immigrated

from the Asian continent 2,000–3,000 years ago (Habu, 2004; Nakagome et al.,

2015). This model is also supported largely by genomic/genetic studies (Jinam et

al., 2012; Nakagome et al., 2015). According to this model, there are two

populations assumed to be strongly contributed from Jomon peoples. One of these

populations is the Ainus, indigenous in Hokkaido, the most northern part of

Japan. The other population comprises people living in the Ryukyu Archipelago,

southern end of Japan (Fig. 1).

This model has been supported by previous genetic studies among the

Ryukyuans, Ainus, mainland Japanese, and other East Asian populations.

Genetic similarity between the Ryukyuans and Ainus has been observed

(Hammer et al., 2006; Koganebuchi et al. 2012), whereas the Ryukyuans are

believed to be differentiated from mainland Japanese by the analyses of the Y

chromosome (Hammer & Horai, 1995; Hammer et al., 2006), mtDNA (Horai et al.,

1996; Tanaka et al., 2004), autosomal markers (Omoto & Saitou, 1997), and

genome-wide SNP data (Yamaguchi-Kabata et al., 2008). Additionally, modern

Ryukyuans are believed to have no direct genetic relations with the Taiwan

aborigines (Matsukusa et al., 2010) or human skeletons in the late Pleistocene

(Sato et al., 2014; Mizuno et al., 2021) that are older than Jomon peoples found in

several sites in the Ryukyu Archipelago (Kobayashi et al., 1974; Nakagawa et al.,

2010). Furthermore, rich genetic variations among the residents on the islands of
the Ryukyu Archipelago have been observed (Sato et al., 2014; Matsunami et al.,
2021).

However, in those studies mentioned above, much attention was not paid
to the Amami islands located in the northern Ryukyu Archipelago (Fig. 1). There
have been historical differences between the Amami islands and the rest of the
Ryukyu Archipelago (Okinawa prefecture). Some researchers claim that the
Amami islands played important roles in the trade among East Asia, including
mainland Japan and China, around the 7th–11th century, which distinguished
the Amami islands from Okinawa (Takanashi, 2011; Yoshinari, 2011). In Kikai
island, one of the Amami islands, there is the Gusuku Site group which is
supposed to be the local agency of the government of mainland Japan in the
9th–13th century (Yoshinari, 2011; Takamiya et al., 2019), and this corresponds to
the hypothesis that there was population migration into the Ryukyu Archipelago
from the north via the Amami islands around the 11th–12th century (Takamiya,
2013). In the 15th century, the Amami islands were conquered by the Ryukyu
Kingdom, which was an independent country centered on main-island Okinawa
(Yoshinari & Fuku, 2007), but since 1609, the Amami islands were directly ruled
by the Satsuma-Han, one of the feudal domains in mainland Japan; this has
brought about administrative and cultural differences between the Amami islands
and Okinawa (Tsuha, 2012).

Despite these non-negligible historical and cultural features, few
anthropological studies, including genetic studies about the Amami islanders,
have been conducted. Dodo et al. (1998) analyzed 22 cranial nonmetric traits of
early modern Amami and Okinawa islanders and found that they shared
morphological similarities. Nishiyama et al. (2012) indicated a low but significant

level of genetic difference between the Amami islanders and mainland Japanese

using SNP data, but their research was limited to only two islands, Tokuno and

Okinoerabu islands (Fig. 1), in the southern part of the Amami islands, and they

concluded that studies that cover further north areas should be indispensable.

We then newly focused on two of the Amami islands, Amami-Oshima and

Kikai islands, in this research. Amami-Oshima island (Fig. 1) is the largest island

and has the largest population in the Amami islands (Kagoshima Prefecture,

2018a). Kikai island (Fig. 1) is a relatively small island with a population of about

7,000 (Kagoshima Prefecture, 2018b), located on the east of Amami-Oshima

island, and assumed to have played a very important role in the relationship

between the Ryukyu Archipelago and mainland Japan around the 11th–12th

century (Takanashi, 2009; Yoshinari, 2011; Takamiya, 2013; Takamiya et al.,

2019). On the basis of the previous studies and historical backgrounds described

above, we hypothesized that people in these two islands might genetically be

located in the intermediate position between Okinawa and mainland Japan. This

study aims to clarify the genetic structure of the Amami islanders compared with

peoples in Okinawa, mainland Japan, and other regions in East Asia.

2. Materials and methods

2.1. Subjects

A total of 104 Amami islanders (Fig. 1), 78 in Amami-Oshima island (49 males and 29 females) and 26 in Kikai island (15 males and 11 females), who were not genetically related, were the subjects of this study. This research was approved by the Research Ethics Committee of the University of Tokyo, and all subjects provided informed consent. We collected nail or oral mucosa samples from the subjects. Their origins were confirmed by an interview that at least one of the grandparents of each subject was born in the Amami islands.

2.2. DNA extraction

We extracted DNA from nail samples using the ISOHAIR (Nippon Gene) and oral mucosa samples using the High Pure PCR Template Preparation Kit (Roche).

2.3. Genotyping

We amplified the D-loop region (16024–16569 and 1–41 of the rCRS; Anderson et al., 1981; Andrews et al., 1999) of mtDNA by nested PCR to avoid nuclear insert contaminations. The first PCR for 9 kb of mtDNA including the D-loop region was conducted (primers: GENBANK_MT_26F & GENBANK_MT_5R) with an initial denaturation at 94°C for 2 min, followed by 25 cycles of denaturation at 98°C for 10 s, annealing at 72°C for 30 s, extension at 68°C for 10 min. The second PCR for the D-loop region was conducted (primers: L15996F & H408R) with an initial denaturation at 94°C for 2 min, followed by 25 cycles of denaturation at 98°C for 10 s, annealing at 56°C for 30 s, extension at

68°C for 1 min. After confirmation by 2% agarose gel electrophoresis, PCR

products were purified using the PCR Clean-Up Mini Kit (FAVORGEN) and

sequenced by outsourcing those to Fasmac (Kanagawa, Japan) or Eurofins

Genomics (Tokyo, Japan). The primers used for PCR amplification and sequencing

are listed in Supp. Table 1.

Four Y-STRs (DYS393, DYS19, DYS391, and DYS438) of which genetic

diversities differed between Honshu (Fig. 1) and Okinawa (Uchihi et al., 2003)

were selected. PCRs for these loci were conducted with an initial denaturation at

95°C for 10 min, followed by 45 cycles of denaturation at 95°C for 15 s, annealing

at 53°C–59°C for 30 s, extension at 72°C for 1 min, followed by a final extension of

5 min at 72°C. After confirmation by 2% agarose gel electrophoresis, PCR

products were purified using the PCR Clean-Up Mini Kit (FAVORGEN) and

sequenced by outsourcing those to Fasmac (Kanagawa, Japan) or Eurofins

Genomics (Tokyo, Japan). The primers and annealing temperatures for each PCR

amplification are listed in Supp. Table 2.

We genotyped four nonsynonymous autosomal SNPs (rs3827760,

rs17822931, rs2070235, and rs14103), of which genotype frequencies differed

between the Hondo cluster, which includes most of the individuals in mainland

Japan and the Ryukyu cluster, which includes most of the individuals in Okinawa

(Yamaguchi-Kabata et al., 2008) by PCR-RFLP. PCRs were conducted with an

initial denaturation at 95°C for 10 min, followed by 40 cycles of denaturation at

95°C for 15 s, annealing at 54°C–65°C for 30 s, extension at 72°C for 1 min,

followed by a final extension of 5 min at 72°C. After confirmation by 2% agarose

gel electrophoresis, PCR products were purified using the PCR Clean-Up Mini Kit

(FAVORGEN) and digested with an appropriate restriction enzyme at 37°C for 1 h.

Thereafter, we electrophoresed these digested products on 4% agarose gel for

genotype determination. The primers, annealing temperatures, and restriction

enzymes for each locus are listed in Supp. Table 3.

2.4. Data analyses

The mtDNA sequences were aligned using MEGA X (Kumar et al., 2018)

together with sequences of other Asian populations downloaded from the NCBI

(National Center for Biotechnology Information) database. Net genetic distances

(d_A) between populations were calculated from these sequences under the

Tamura-Nei model (Tamura & Nei, 1993) with gamma distribution. Phylogenetic

networks between populations based on d_A distances were constructed in the

Neighbor-Net method (Bryant & Moulton, 2004) by using SplitsTree4 (Huson &

Bryant, 2006). Pairwise F_{ST} between regions and nucleotide diversities within

each population were calculated by using Arlequin ver 3.5.2.2 (Excoffier & Lischer,

2010), and Tajima's D (Tajima, 1989) within each population were calculated

using DnaSP 5.10 (Librado & Rozas, 2009). The probabilities of Tajima's D were

computed through 5,000 coalescence simulations.

We aligned the Y-STR sequences using MEGA X and counted the number

of repeats. Pairwise R_{ST} between regions was calculated by using Arlequin ver

3.5.2.2 as well as gene diversities.

Fisher's exact test was used to compare the allele frequencies of the SNPs

between populations.

3. Results

3.1. mtDNA

D-loop sequences (587 bp) were determined for a total of 102 subjects

whose maternal grandmothers had been born in the Amami islands. Fig. 2 and

Supp. Fig. 1 show the phylogenetic networks for 78 Amami-Oshima islanders, 24

Kikai islanders, and other Asian populations. Amami-Oshima and Kikai islands

were close to each other in both networks, and the Amami islanders did not form a

clear cluster with the residents of Honshu, Kyushu, the Okinawa islands, or any

other populations. Table 1 shows the nucleotide diversities and Tajima's D in each

population. Nucleotide diversities in the Amami islands were almost the same as

those in other regions in Japan. Tajima's D values were significantly negative in

the Amami islands (−1.613, P = 0.024) and Amami-Oshima island (−1.539, P =

0.038). Tajima's D value in Kikai island was negative but not significant (−1.494,

P = 0.050). Table 2 shows the pairwise F_{ST} values between regions. F_{ST} value

between Amami-Oshima and Kikai islands was not significant. F_{ST} values

between Amami-Oshima island and Honshu, between Amami-Oshima island and

Okinawa, between Amami-Oshima and Ishigaki islands, and between Kikai and

Ishigaki islands were significant. Additionally, F_{ST} values between Korea and four

regions in Japan (Amami-Oshima island, Kikai island, Kyushu, and Honshu)

were not significant.

3.2. Y-STR

Y-STR sequences were determined for 58 male subjects whose paternal

grandfathers had been born in the Amami islands. Twenty-five different

haplotypes were observed in the Amami islands (Supp. Table 4). Three of them

were observed in Kikai island but not in Amami-Oshima island. The paternal

grandfather of one subject who had one of the three haplotypes (Ht11) was born in

Amami-Oshima island. This subject was counted as Amami-Oshima samples in

the Y-STR analyses. Comparing with other regions in Japan (Hashiyada et al.,

2008), six haplotypes observed in Amami-Oshima island were not in Okinawa,

and one haplotype was found only in Amami-Oshima island and Okinawa but not

in any other regions. The allele frequency distributions of four loci were similar

among regions (Supp. Table 5). Gene diversities of the four loci are shown in Supp.

Table 6. In Amami-Oshima island, gene diversities of any loci did not significantly

differ with Honshu or Okinawa. In Kikai island, gene diversities of the loci except

DYS438 did not differ with other regions, and that of DYS438 was significantly

higher than any other regions. Table 3 shows the pairwise R_{ST} values between

regions. R_{ST} value between Amami-Oshima and Kikai islands was significant. R_{ST}

values between Amami-Oshima island and Okinawa and between Kikai island

and Okinawa were not significant.

3.3. Autosomal SNPs

Genotypes of 91 subjects whose all four grandparents were born in the

Amami islands were determined, and allele frequencies were calculated.

The frequency of the T allele in rs3827760 (*EDAR*) in Amami-Oshima

island (0.385) was significantly higher than in the Hondo cluster (0.222) ($P <$

0.001) and not significantly different from the Ryukyu cluster (0.398) ($P = 0.426$).

The frequency of the T allele in Kikai island (0.647) was significantly higher than

in Amami-Oshima island, the Ryukyu cluster, and the Hondo cluster ($P = 0.005$, P

= 0.004, and $P < 0.001$, respectively) (Table 4).

The frequency of the G allele in rs17822931 (*ABCC11*) in Amami-Oshima

island (0.311) was significantly higher than in the Hondo cluster (0.121) ($P <$

0.001) but not significantly different from that in the Ryukyu cluster (0.258) ($P =$

0.123). The frequency of the G allele in Kikai island (0.118) was significantly

lower than in the Ryukyu cluster ($P = 0.044$) and Amami-Oshima island ($P =$

0.015) but not significantly different from that in the Hondo cluster ($P = 0.612$)

(Table 4).

The frequency of the G allele in rs2070235 (*MYBL2*) in Amami-Oshima

island (0.324) was significantly higher than in the Hondo cluster (0.185) ($P <$

0.001) but not significantly different from that in the Ryukyu cluster (0.291) ($P =$

0.228). The frequency of the G allele in Kikai island was not significantly different

from those in Amami-Oshima island, the Ryukyu cluster, and the Hondo cluster

($P = 0.324$, $P = 0.456$, and $P = 0.167$, respectively) (Table 4).

The frequency of the A allele in rs14103 (*C1orf212*) in Amami-Oshima

island (0.791) was significantly lower than in the Hondo cluster (0.855) ($P = 0.022$)

and not significantly different from that in the Ryukyu cluster (0.753) ($P = 0.187$),

whereas that in Kikai island was not different from those in Amami-Oshima

island, the Ryukyu cluster, and the Hondo cluster ($P = 0.287$, $P = 0.126$, and $P =$

0.562, respectively) (Table 4).

4. Discussion

Amami islanders showed a trend of Tajima's D values of mtDNA to be

negative. This trend is commonly observed in the populations of East Asia,

including main-island Okinawa, Miyako island, Ishigaki island, and Honshu

(Oota et al., 2002; Matsukusa et al., 2010). Purifying selection or recent

population expansion would generally be a reason for the negative Tajima's D

values (Tajima, 1989), but the former is unsuitable in this case because D-loop is a

non-coding region. Therefore, this result suggests the past population expansion

and/or gene flow from the surrounding regions, *i.e.,* the Amani islanders were

genetically not isolated for long.

Pairwise F_{ST} values and phylogenetic networks represent close genetic

relations in the female lineages between Amami-Oshima and Kikai islands. The

female lineages in Amami-Oshima island differed from Honshu, main-island

Okinawa, and Ishigaki island, and those in Kikai island differed from Ishigaki

island. In the phylogenetic network, the female lineages in Amami-Oshima and

Kikai islands were apart from those in main-island Okinawa and Ishigaki island.

Conversely, the female lineages in Amami-Oshima and Kikai islands did not differ

from those in Miyako island, and those three were located close to each other in

the phylogenetic network; however, no historical records that link geographically

separated people in the Amami islands and Miyako island have been recognized.

Furthermore, female lineages of Korea were not different from those of

Amami-Oshima island, Kikai island, Kyushu, and Honshu, whereas female

lineages of Ainu, Taiwan Han, and Taiwan aborigines were different from any

other populations; this indirectly suggests that female lineages in the Amami

islanders have been genetically affected by mainland Japanese strongly compared

with the genetic effect of mainland Japanese on female lineages of Okinawa.

Gene diversities and R_{ST} values of the four Y-STRs indicated that the male

lineages in Amami-Oshima island were located between those in Honshu and

Okinawa, but those in Kikai island showed some genetic deviation from these

three populations, including Amami-Oshima islanders; this is assumed to be due

to the founder effect that refers to the phenomenon that gene frequencies of a

small population migrated to a new location sometimes differ from those of the

original population because of chance bias (Mayr, 1954). In Kikai islanders, the

small population size (Kagoshima Prefecture, 2018b) and the degree of contacts

with mainland Japanese different from Amami-Oshima islanders (Takanashi,

2009; Yoshinari, 2011; Takamiya, 2013; Takamiya et al., 2019) may have caused

that.

Comparing the maternal (mtDNA) and paternal (Y-STR) genes (Table 5),

we found that the mean of pairwise R_{ST} values of Y-STR had a trend that the

larger the value is, the smaller population size is (0.034 in entire Japan, 0.055 in

the Ryukyu Archipelago and Kyushu, 0.073 in the Ryukyu Archipelago, and 0.113

in the Amami islands). Conversely, the mean F_{ST} values of mtDNA did not change

so much (0.020 in entire Japan, 0.021 in the Ryukyu Archipelago and Kyushu,

0.019 in the Ryukyu Archipelago, and − 0.007 in the Amami islands). These values

suggest that males are genetically more divergent than females because of

patrilocality; in fact, female migration was suggested frequently in the Ryukyu

Archipelago (Matsukusa et al., 2010), and we confirmed this in the Amami

islands.

Four autosomal SNPs whose allele frequencies differed between Okinawa

and the rest of Japan (Yamaguchi-Kabata et al., 2008) were selected to identify

the Amami islanders' genealogic position and search for the traits of selective

pressure if it ever exerted in these populations. The allele frequencies of the four

SNPs in Amami-Oshima islanders were similar to the Ryukyu cluster and

significantly different from the Hondo cluster, which may be because of gene flow

from Okinawa. Conversely, in Kikai islanders, the T allele frequency of rs3827760

(*EDAR*) was significantly higher than those in the other three populations, and

the G allele frequency of rs17822931 (*ABCC11*) was significantly lower than those

in the Ryukyu cluster and Amami-Oshima islanders but not lower than that in

the Hondo cluster. These can be explained by the founder effect, as observed in

Y-STR: DYS438. The presence of selective pressure is hard to recognize in these

loci for the following reasons.

The C allele of rs3827760 on the EDAR gene is associated with thicker

hair (Fujimoto et al., 2008a, b), straighter hair (Tan et al., 2013), multiple dental

traits (Kimura et al., 2009; Park et al., 2012; Kamberov et al., 2013), and high

density of sweat gland (Kamberov et al., 2013). The frequencies of the C allele are

very high in East Asians and Native Americans, whereas that allele is almost

absent in Europeans and Africans, and this implies strong positive selection in

Asia (Sabeti et al., 2007; Bryk et al., 2008; Xue et al., 2009; Grossman et al., 2010).

Adaptation to high humidity is assumed to be selective pressure (Kamberov et al.,

2013; Tan et al., 2013), but frequencies of the C allele in the Ryukyu cluster,

Amami-Oshima island, and Kikai island were significantly lower than that in the

Hondo cluster, despite hot and humid climate in the Ryukyu Archipelago (Table 4).

This suggests that selective pressure on the C allele has not been so strong in the

Ryukyu Archipelago, and low frequencies of the C allele in the Amami islands

may be due to gene flow from Okinawa.

Secondly, G/G and G/A genotypes of rs17822931 located on the ABCC11

gene are associated with wet earwax (Yoshiura et al., 2006) and axillary

osmidrosis (Nakano et al., 2009), which are assumed to be due to their enhanced

function of the apocrine gland. The frequencies of the A allele are very high in

East Asia. They decline when the latitude is getting lower in Asian, African, and

European populations, and this implies strong positive selection and can be

explained by adaptation to cold climate, such as less sweating (Yoshiura et al.,

2006; Ohashi et al., 2011). Therefore, the frequencies of the A allele are

significantly lower in the Ryukyu cluster and Amami-Oshima island than in the

Hondo cluster (Table 4) because selective pressure should not have existed under

hotter circumstances in the Ryukyu Archipelago.

In conclusion, Amami-Oshima and Kikai island populations showed a

genetically intermediate position between mainland Japan and Okinawa in

mtDNA and Y-STR. However, the frequencies of several autosomal SNPs in the

Amami islanders indicated a significant difference from mainland Japanese. The

EDAR and *ABCC11* of mainland Japanese are assumed to have been under

natural selection; however, those of the Amami islanders may have been affected

by the gene flow from Okinawa but not natural selection. Furthermore, the

extremely high or low frequencies of several alleles implied a founder effect in

Kikai island. It should be noted that we could not say with strong evidence

whether mtDNA and Y-STRs of the Amami islanders were close to Okinawa or

mainland Japan and whether autosomal SNPs of the Amami islanders have been

affected by gene flow from Okinawa because of the small sample size (especially in

Kikai island) and the number of alleles in the present study. Geographically broad

and detailed samplings and genome-wide analyses are awaited.

Declarations of interest: none

Acknowledgments

We would like to thank all anonymous subjects in Naze, Amami-shi,

Kagoshima, in Kasari-cho, Amami-shi, Kagoshima, in Yamato-son, Oshima-gun,

Kagoshima, and in Kikai-cho, Oshima-gun, Kagoshima for participating in this

study. We are grateful to Dr. M. Suzuki (Ministry of the Environment, Japan),

Prof. H. Takamiya (Kagoshima University), Ms. R. Koike, and anonymous

cooperators for their help in sampling. We thank Enago (www.enago.jp) for the

English language review.

References

Anderson, S., Bankier, A.T., Barrell, B.G., de Bruijn, M.H.L., Coulson, A.R.,

Drouin, J., Eperon, I.C., Nierlich, D.P., Roe, B.A., Sanger, F., Schreier, P.H.,

Smith, A.J.H., Staden, R., Young, I.G., 1981. Sequence and organization of the

human mitochondrial genome. Nature 290, 457–465.

https://doi.org/10.1038/290457a0

Andrews, R.M., Kubacka, I., Chinnery, P.F., Lightowlers, R.N., Turnbull, D.M.,

Howell, N., 1999. Reanalysis and revision of the Cambridge reference

sequence for human mitochondrial DNA. Nat. Genet. 23, 147–147.

https://doi.org/10.1038/13779

Ayub, Q., 2000. Identification and characterisation of novel human

Y-chromosomal microsatellites from sequence database information. Nucleic

Acids Res. 28, 8e – 8. https://doi.org/10.1093/nar/28.2.e8

Bryant, D., Moulton, V., 2004. Neighbor-Net: An Agglomerative Method for the

Construction of Phylogenetic Networks. Mol. Biol. Evol. 21, 255–265.

https://doi.org/10.1093/molbev/msh018

Bryk, J., Hardouin, E., Pugach, I., Hughes, D., Strotmann, R., Stoneking, M.,

Myles, S., 2008. Positive Selection in East Asians for an EDAR Allele that

Enhances NF-κB Activation. PLoS One 3, e2209.

https://doi.org/10.1371/journal.pone.0002209

Butler, J.M., Schoske, R., Vallone, P.M., Kline, M.C., Redd, A.J., Hammer, M.F.,

2002. A novel multiplex for simultaneous amplification of 20 Y chromosome

STR markers. Forensic Sci. Int. 129, 10–24.

https://doi.org/10.1016/S0379-0738(02)00195-0

de Knijff, P., Kayser, M., Caglià, A., Corach, D., Fretwell, N., Gehrig, C., Graziosi,

G., Heidorn, F., Herrmann, S., Herzog, B., Hidding, M., Honda, K., Jobling,

M., Krawczak, M., Leim, K., Meuser, S., Meyer, E., Oesterreich, W., Pandya,

A., Parson, W., Penacino, G., Perez-Lezaun, A., Piccinini, A., Prinz, M.,

Schmitt, C., Schneider, P.M., Szibor, R., Teifel-Greding, J., Weichhold, G.,

Roewer, L., 1997. Chromosome Y microsatellites: population genetic and

evolutionary aspects. Int. J. Legal Med. 110, 134–140.

https://doi.org/10.1007/s004140050052

Dodo, Y., Doi, N., Kondo, O., 1998. Ainu and Ryukyuan cranial nonmetric

variation: Evidence which disputes the Ainu-Ryukyu common origin theory.

Anthropol. Sci. 106, 99–120. https://doi.org/10.1537/ase.106.99

Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: a new series of

programs to perform population genetics analyses under Linux and Windows.

Mol. Ecol. Resour. 10, 564–567.

https://doi.org/10.1111/j.1755-0998.2010.02847.x

Fujimoto, A., Kimura, R., Ohashi, J., Omi, K., Yuliwulandari, R., Batubara, L.,

Mustofa, M.S., Samakkarn, U., Settheetham-Ishida, W., Ishida, T., Morishita,

Y., Furusawa, T., Nakazawa, M., Ohtsuka, R., Tokunaga, K., 2008a. A scan for

genetic determinants of human hair morphology: EDAR is associated with

Asian hair thickness. Hum. Mol. Genet. 17, 835–843.

https://doi.org/10.1093/hmg/ddm355

Fujimoto, A., Ohashi, J., Nishida, N., Miyagawa, T., Morishita, Y., Tsunoda, T.,

Kimura, R., Tokunaga, K., 2008b. A replication study confirmed the EDAR

gene to be a major contributor to population differentiation regarding head

hair thickness in Asia. Hum. Genet. 124, 179–185.

https://doi.org/10.1007/s00439-008-0537-1

Grossman, S.R., Shylakhter, I., Karlsson, E.K., Byrne, E.H., Morales, S., Frieden,

G., Hostetter, E., Angelino, E., Garber, M., Zuk, O., Lander, E.S., Schaffner,

414  S.F., Sabeti, P.C., 2010. A Composite of Multiple Signals Distinguishes Causal

415  Variants in Regions of Positive Selection. Science (80-.). 327, 883 LP – 886.

416  https://doi.org/10.1126/science.1183863

417 Habu, J., 2004. Ancient Jomon of Japan. Cambridge University Press, Cambridge.

418 Hammer, M.F., Horai, S., 1995. Y chromosomal DNA variation and the peopling of

419  Japan. Am. J. Hum. Genet. 56, 951–962.

420 Hammer, M.F., Karafet, T.M., Park, H., Omoto, K., Harihara, S., Stoneking, M.,

421  Horai, S., 2006. Dual origins of the Japanese: Common ground for

422  hunter-gatherer and farmer Y chromosomes. J. Hum. Genet. 51, 47–58.

423  https://doi.org/10.1007/s10038-005-0322-0

424 Hanihara, K., 1991. Dual Structure Model for the Population History of the

425  Japanese. Japan Rev. 2, 1–33.

426 Hashiyada, M., Umetsu, K., Yuasa, I., Tamura, A., Matsusue, A., Suzuki, K.,

427  Kashimura, S., Funayama, M., 2008. Population genetics of 17

428  Y-chromosomal STR loci in Japanese. Forensic Sci. Int. Genet. 2, 69–70.

429  https://doi.org/10.1016/j.fsigen.2008.01.001

430 Hayashida, M., Ota, T., Iwao-Koizumi, K., Murata, S., 2010. Simple SNP

431  genotyping of hair thickness-determining gene EDAR and human

432  earwax-type gene ABCC11 for education of genetic tests. Bull. Mukogawa

433  Women's Univ. Nat. Sci. 58, 43–47. https://doi.org/10.14993/00000508

434 Horai, S., Murayama, K., Hayasaka, K., Matsubayashi, S., Hattori, Y., Fucharoen,

435  G., Harihara, S., Park, K.S., Omoto, K., Pan, I.H., 1996. mtDNA

436  polymorphism in East Asian populations, with special reference to the

437  peopling of Japan. Am. J. Hum. Genet. 59, 579–590.

438 Huson, D.H., Bryant, D., 2006. Application of Phylogenetic Networks in

439  Evolutionary Studies. Mol. Biol. Evol. 23, 254–267.

https://doi.org/10.1093/molbev/msj030

Jinam, T., Nishida, N., Hirai, M., Kawamura, S., Oota, H., Umetsu, K., Kimura,

R., Ohashi, J., Tajima, A., Yamamoto, T., Tanabe, H., Mano, S., Suto, Y.,

Kaname, T., Naritomi, K., Yanagi, K., Niikawa, N., Omoto, K., Tokunaga, K.,

Saitou, N., 2012. The history of human populations in the Japanese

Archipelago inferred from genome-wide SNP data with a special reference to

the Ainu and the Ryukyuan populations. J. Hum. Genet. 57, 787–795.

https://doi.org/10.1038/jhg.2012.114

Kagoshima Prefecture. (2018a) Amami-Oshima Kakeroma shotō no gaiyō.

http://www.pref.kagoshima.jp/ac07/pr/shima/gaiyo/amami/amami_top.html

(accessed 19 April 2021). (in Japanese)

Kagoshima Prefecture. (2018b) Kikaijima no gaiyō.

http://www.pref.kagoshima.jp/ac07/pr/shima/gaiyo/kikai/kikai_top.html

(accessed 19 April 2021). (in Japanese)

Kamberov, Y.G., Wang, S., Tan, J., Gerbault, P., Wark, A., Tan, L., Yang, Y., Li, S.,

Tang, K., Chen, H., Powell, A., Itan, Y., Fuller, D., Lohmueller, J., Mao, J.,

Schachar, A., Paymer, M., Hostetter, E., Byrne, E., Burnett, M., McMahon,

A.P., Thomas, M.G., Lieberman, D.E., Jin, L., Tabin, C.J., Morgan, B.A.,

Sabeti, P.C., 2013. Modeling Recent Human Evolution in Mice by Expression

of a Selected EDAR Variant. Cell 152, 691–702.

https://doi.org/10.1016/j.cell.2013.01.016

Kayser, M., Caglià, A., Corach, D., Fretwell, N., Gehrig, C., Graziosi, G., Heidorn,

F., Herrmann, S., Herzog, B., Hidding, M., Honda, K., Jobling, M., Krawczak,

M., Leim, K., Meuser, S., Meyer, E., Oesterreich, W., Pandya, A., Parson, W.,

Penacino, G., Perez-Lezaun, A., Piccinini, A., Prinz, M., Schmitt, C.,

Schneider, P.M., Szibor, R., Teifel-Greding, J., Weichhold, G., De Knijff, P.,

Roewer, L., 1997. Evaluation of Y-chromosomal STRs: A multicenter study.

Int. J. Legal Med. 110, 125–133. https://doi.org/10.1007/s004140050051

Kimura, R., Yamaguchi, T., Takeda, M., Kondo, O., Toma, T., Haneji, K., Hanihara,

T., Matsukusa, H., Kawamura, S., Maki, K., Osawa, M., Ishida, H., Oota, H.,

2009. A Common Variation in EDAR Is a Genetic Determinant of

Shovel-Shaped Incisors. Am. J. Hum. Genet. 85, 528–535.

https://doi.org/10.1016/j.ajhg.2009.09.006

Kobayashi, H., Hirose, T., Sugino, M., Watanabe, N., 1974. University of Tokyo

Radiocarbon Measurements V. Radiocarbon 16, 381–387.

https://doi.org/10.1017/S0033822200059671

Koganebuchi, K., Katsumura, T., Nakagome, S., Ishida, H., Kawamura, S., Oota,

H., Ohashi, J., Yamamoto, T., Nishida, N., Tajima, A., Tanabe, H., Umetsu, K.,

Saitou, N., Tokunaga, K., Suto, Y., Hirai, M., Kimura, R., Mano, S.,

Kawamura, S., Oota, H., Omoto, K., 2012. Autosomal and Y-chromosomal

STR markers reveal a close relationship between Hokkaido Ainu and Ryukyu

islanders. Anthropol. Sci. 120, 199–208. https://doi.org/10.1537/ase.120322

Kumar, S., Stecher, G., Li, M., Knyaz, C., Tamura, K., 2018. MEGA X: Molecular

evolutionary genetics analysis across computing platforms. Mol. Biol. Evol. 35,

1547–1549. https://doi.org/10.1093/molbev/msy096

Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of

DNA polymorphism data. Bioinformatics 25, 1451–1452.

https://doi.org/10.1093/bioinformatics/btp187

Matsukusa, H., Oota, H., Haneji, K., Toma, T., Kawamura, S., Ishida, H., 2010. A

genetic analysis of the sakishima islanders reveals no relationship with

taiwan aborigines but shared ancestry with ainu and main-island Japanese.

Am. J. Phys. Anthropol. 142, 211–223. https://doi.org/10.1002/ajpa.21212

Matsunami, M., Koganebuchi, K., Imamura, M., Ishida, H., Kimura, R., Maeda, S.,

2021. Fine-scale genetic structure and demographic history in the Miyako

Islands of the Ryukyu Archipelago. Mol. Biol. Evol.

https://doi.org/10.1093/molbev/msab005

Mayr, E., 1954. Change of environment and evolution, in: Huxley, J., Hardy, A.C.,

Ford, E.B. (Eds.), Evolution as a Process. Allen and Unwin, London, pp.

157-180.

Mizuno, F., Gojobori, J., Kumagai, M., Baba, H., Taniguchi, Y., 2021. Population

dynamics in the Japanese Archipelago since the Pleistocene revealed by the

complete mitochondrial genome sequences. Sci. Rep. 1–11.

https://doi.org/10.1038/s41598-021-91357-2

Nakagawa, R., Doi, N., Nishioka, Y., Nunami, S., Yamauchi, H., Fujita, M.,

Yamazaki, S., Yamamoto, M., Katagiri, C., Mukai, H., Matsuzaki, H.,

Gakuhari, T., Takigami, M., Yoneda, M., 2010. Pleistocene human remains

from Shiraho-Saonetabaru Cave on Ishigaki Island, Okinawa, Japan, and

their radiocarbon dating. Anthropol. Sci. 118, 173–183.

https://doi.org/10.1537/ase.091214

Nakagome, S., Sato, T., Ishida, H., Hanihara, T., Yamaguchi, T., Kimura, R., Mano,

S., Oota, H., 2015. Model-Based Verification of Hypotheses on the Origin of

Modern Japanese Revisited by Bayesian Inference Based on Genome-Wide

SNP Data. Mol. Biol. Evol. 32, 1533–1543.

https://doi.org/10.1093/molbev/msv045

Nakano, M., Miwa, N., Hirano, A., Yoshiura, K.I., Niikawa, N., 2009. A strong

association of axillary osmidrosis with the wet earwax type determined by

genotyping of the ABCC11 gene. BMC Genet. 10, 1–5.

https://doi.org/10.1186/1471-2156-10-42

Nishiyama, T., Kishino, H., Suzuki, S., Ando, R., Niimura, H., Uemura, H., Horita,

M., Ohnaka, K., Kuriyama, N., Mikami, H., Takashima, N., Mastuo, K.,

Guang, Y., Wakai, K., Hamajima, N., Tanaka, H., 2012. Detailed Analysis of

Japanese Population Substructure with a Focus on the Southwest Islands of

Japan. PLoS One 7, e35000. https://doi.org/10.1371/journal.pone.0035000

Ohashi, J., Naka, I., Tsuchiya, N., 2011. The impact of natural selection on an

ABCC11 SNP determining earwax type. Mol. Biol. Evol. 28, 849–857.

https://doi.org/10.1093/molbev/msq264

Omoto, K., Saitou, N., 1997. Genetic origins of the Japanese: A partial support for

the dual structure hypothesis. Am. J. Phys. Anthropol. 102, 437–446.

https://doi.org/10.1002/(SICI)1096-8644(199704)102:4<437::AID-AJPA1>3.0.

CO;2-P

Oota, H., Kitano, T., Jin, F., Yuasa, I., Wang, L., Ueda, S., Saitou, N., Stoneking,

M., 2002. Extreme mtDNA homogeneity in continental Asian populations. Am.

J. Phys. Anthropol. 118, 146–153. https://doi.org/10.1002/ajpa.10056

Park, J.-H., Yamaguchi, T., Watanabe, C., Kawaguchi, A., Haneji, K., Takeda, M.,

Kim, Y.-I., Tomoyasu, Y., Watanabe, M., Oota, H., Hanihara, T., Ishida, H.,

Maki, K., Park, S.-B., Kimura, R., 2012. Effects of an Asian-specific

nonsynonymous EDAR variant on multiple dental traits. J. Hum. Genet. 57,

508–514. https://doi.org/10.1038/jhg.2012.60

Roewer, L., Epplen, J.T., 1992. Rapid and sensitive typing of forensic stains by

PCR amplification of polymorphic simple repeat sequences in case work.

Forensic Sci. Int. 53, 163–171.

https://doi.org/https://doi.org/10.1016/0379-0738(92)90193-Z

Sabeti, P.C., Varilly, P., Fry, B., Lohmueller, J., Hostetter, E., Cotsapas, C., Xie, X.,

Byrne, E.H., McCarroll, S.A., Gaudet, R., Schaffner, S.F., Lander, E.S., 2007.

Genome-wide detection and characterization of positive selection in human

populations. Nature 449, 913–918. https://doi.org/10.1038/nature06250

Sato, T., Nakagome, S., Watanabe, C., Yamaguchi, K., Kawaguchi, A.,

Koganebuchi, K., Haneji, K., Yamaguchi, T., Hanihara, T., Yamamoto, K.,

Ishida, H., Mano, S., Kimura, R., Oota, H., 2014. Genome-wide SNP analysis

reveals population structure and demographic history of the ryukyu islanders

in the southern part of the japanese archipelago. Mol. Biol. Evol. 31,

2929–2940. https://doi.org/10.1093/molbev/msu230

Stanford Genome Technology Center, 2002. List of primers used for resequencing

the mitochondrial genome. http://insertion.stanford.edu/primers.html

(accessed 19 April 2021).

Tajima, A., Sun, C.-S., Pan, I.-H., Ishida, T., Saitou, N., Horai, S., 2003.

Mitochondrial DNA polymorphisms in nine aboriginal groups of Taiwan:

implications for the population history of aboriginal Taiwanese. Hum. Genet.

113, 24–33. https://doi.org/10.1007/s00439-003-0945-1

Tajima, F., 1989. Statistical method for testing the neutral mutation hypothesis

by DNA polymorphism. Genetics 123, 585–595.

Takamiya, H., 2013. Amami Okinawa shoto eno hito no idō, in: Intoh, H. (Eds.),

Jinrui no idōshi. Rinsen Shoten, Kyoto, pp. 182–197. (in Japanese)

Takamiya, H., Katagiri, C., Yamasaki, S., Fujita, M., 2019. Human Colonization of

the Central Ryukyus (Amami and Okinawa Archipelagos), Japan. J. Isl.

Coast. Archaeol. 14, 375–393. https://doi.org/10.1080/15564894.2018.1501443

Takanashi, O., 2009. Doki dōtai kara kangaeru "Nihon bunka no nanzen", in:

Takanashi, O., Abe, M., Nakamoto, K., Yoshinari, N, Okinawa bunka wa doko

kara kitaka: Gusuku jidai to iu kakki. Shinwasha, Tokyo, pp. 47–132. (in

Japanese)

Takanashi, O., 2011. Nampō sekai eno hirogari: Ritsuryō kokka to Ryukyu-ko, in:

Arano, Y., Ishii, M., Murai, S. (Eds.), Ritsuryō kokka to Higashi Ajia.

Yoshikawakobunkan, Tokyo, pp. 319–333. (in Japanese)

Tamura, K., Nei, M., 1993. Estimation of the number of nucleotide substitutions

in the control region of mitochondrial DNA in humans and chimpanzees. Mol.

Biol. Evol. 10, 512–526.

https://doi.org/10.1093/oxfordjournals.molbev.a040023

Tan, J., Yang, Y., Tang, K., Sabeti, P.C., Jin, L., Wang, S., 2013. The adaptive

variant EDARV370A is associated with straight hair in East Asians. Hum.

Genet. 132, 1187–1191. https://doi.org/10.1007/s00439-013-1324-1

Tanaka, M., Cabrera, V.M., González, A.M., Larruga, J.M., Takeyasu, T., Fuku, N.,

Guo, L.J., Hirose, R., Fujita, Y., Kurata, M., Shinoda, K.I., Umetsu, K.,

Yamada, Y., Oshida, Y., Sato, Y., Hattori, N., Mizuno, Y., Arai, Y., Hirose, N.,

Ohta, S., Ogawa, O., Tanaka, Y., Kawamori, R., Shamoto-Nagai, M.,

Maruyama, W., Shimokata, H., Suzuki, R., Shimodaira, H., 2004.

Mitochondrial genome variation in Eastern Asia and the peopling of Japan.

Genome Res. 14, 1832–1850. https://doi.org/10.1101/gr.2286304

Tsuha, T., 2012. Okinawa gawa kara mita Amami no bunka henyō. Daiichi Shobo,

Tokyo. (in Japanese)

Uchihi, R., Yamamoto, T., Usuda, K., Yoshimoto, T., Tanaka, M., Tokunaga, S.,

Kurihara, R., Tokunaga, K., Katsumata, Y., 2003. Haplotype analysis with 14

Y-STR loci using 2 multiplex amplification and typing systems in 2 regional

populations in Japan. Int. J. Legal Med. 117, 34–38.

https://doi.org/10.1007/s00414-002-0319-6

Vigilant, L., Pennington, R., Harpending, H., Kocher, T.D., Wilson, A.C., 1989.

Mitochondrial DNA sequences in single hairs from a southern African

population. Proc. Natl. Acad. Sci. U. S. A. 86, 9350–9354.

https://doi.org/10.1073/pnas.86.23.9350

Xue, Y., Zhang, X., Huang, N., Daly, A., Gillson, C.J., MacArthur, D.G.,

Yngvadottir, B., Nica, A.C., Woodwark, C., Chen, Y., Conrad, D.F., Ayub, Q.,

Mehdi, S.Q., Li, P., Tyler-Smith, C., 2009. Population Differentiation as an

Indicator of Recent Positive Selection in Humans: An Empirical Evaluation.

Genetics 183, 1065 LP – 1077. https://doi.org/10.1534/genetics.109.107722

Yamaguchi-Kabata, Y., Nakazono, K., Takahashi, A., Saito, S., Hosono, N., Kubo,

M., Nakamura, Y., Kamatani, N., 2008. Japanese Population Structure,

Based on SNP Genotypes from 7003 Individuals Compared to Other Ethnic

Groups: Effects on Population-Based Association Studies. Am. J. Hum. Genet.

83, 445–456. https://doi.org/10.1016/j.ajhg.2008.08.019

Yamaguchi-Kabata, Y., Tsunoda, T., Kumasaka, N., Takahashi, A., Hosono, N.,

Kubo, M., Nakamura, Y., Kamatani, N., 2012. Genetic differences in the two

main groups of the Japanese population based on autosomal SNPs and

haplotypes. J. Hum. Genet. 57, 326–334. https://doi.org/10.1038/jhg.2012.26

Yoshinari, N., 2011. Ryukyu no seiritsu: Ijū to kōeki no rekishi. Nanpou Shinsha,

Kagoshima. (in Japanese)

Yoshinari, N., Fuku, H., 2007. Ryukyu ōkoku tanjō: Amami shotō shi kara.

Shinwasha, Tokyo. (in Japanese)

Yoshiura, K., Kinoshita, A., Ishida, T., Ninokata, A., Ishikawa, T., Kaname, T.,

Bannai, M., Tokunaga, K., Sonoda, S., Komaki, R., Ihara, M., Saenko, V.A.,

Alipov, G.K., Sekine, I., Komatsu, K., Takahashi, H., Nakashima, M.,

Sosonkina, N., Mapendano, C.K., Ghadami, M., Nomura, M., Liang, D.-S.,

Miwa, N., Kim, D.-K., Garidkhuu, A., Natsume, N., Ohta, T., Tomita, H.,

Kaneko, A., Kikuchi, M., Russomando, G., Hirayama, K., Ishibashi, M.,

Takahashi, A., Saitou, N., Murray, J.C., Saito, S., Nakamura, Y., Niikawa, N.,

2006. A SNP in the ABCC11 gene is the determinant of human earwax type.

Nat. Genet. 38, 324–330. https://doi.org/10.1038/ng1733

Table 1. Nucleotide diversities and Tajima's D values of mtDNA

	Size	Nucleotide diversity	Tajima's D	P value	Ref.
Amami islands	102	0.019 ± 0.010	-1.613	0.024	Present study
Amami-Oshima island	78	0.019 ± 0.010	-1.516	0.042	Present study
Kikai island	24	0.020 ± 0.011	-1.494	0.050	Present study
Main-island Okinawa	95	0.019 ± 0.010	-1.972	0.004	Matsukusa et al., 2010
Miyako island	66	0.015 ± 0.008	-1.916	0.008	Matsukusa et al., 2010
Ishigaki island	63	0.016 ± 0.008	-1.582	0.039	Matsukusa et al., 2010
Honshu	89	0.015 ± 0.008	-2.050	0.002	Oota et al., 2002

For the analyses of mtDNA, one subject from Kikai island whose maternal grandmother was born in Amami-Oshima island was excluded from Kikai island samples and included in Amami-Oshima island samples.

Table 2. Pairwise F_{ST} values of mtDNA

	Amami-Oshima island	Kikai island	Main-island Okinawa	Miyako island	Ishigaki island	Kyushu	Honshu	Ainu	Korea	Taiwan Han	Taiwan aborigines
Amami-Oshima island (n=78)	0										
Kikai island (n=24)	-0.007	0									
Main-island Okinawa (n=95)	**0.026**	0.003	0								
Miyako island (n=66)	0.004	-0.001	**0.034**	0							
Ishigaki island (n=63)	**0.044**	**0.028**	**0.013**	**0.043**	0						
Kyushu (n=104)	-0.001	-0.004	**0.036**	**0.013**	**0.055**	0					
Honshu (n=89)	**0.009**	0.010	**0.043**	**0.012**	**0.045**	**0.008**	0				
Ainu (n=51)	**0.043**	**0.027**	**0.043**	**0.062**	**0.068**	**0.042**	**0.074**	0			
Korea (n=64)	0.004	-0.002	**0.036**	**0.014**	**0.048**	-0.001	0.004	**0.040**	0		
Taiwan Han (n=52)	**0.033**	**0.037**	**0.081**	**0.053**	**0.102**	**0.030**	**0.033**	**0.078**	**0.039**	0	
Taiwan aborigines (n=180)	**0.074**	**0.057**	**0.097**	**0.103**	**0.120**	**0.060**	**0.092**	**0.080**	**0.069**	**0.084**	0

Data of Miyako, Ishigaki and Main-island Okinawa were taken from Matsukusa et al. (2010). Data of Honshu were taken from Oot a et al. (2002) 2002). Data of Ainu, Korea and Taiwan Han were taken from Horai et al. (1996). Data of Taiwan aborigines were taken from Tajima et al. (2003). Significant differences are in bold (p < 0.05).

Table 3. Pairwise R_{ST} values of Y-STR

| | Amami-Oshima island | Kikai island | Okinawa | Kyushu | | Honshu | | |
				Fukuoka	Osaka	Tottori	Yamagata	Miyagi
Amami-Oshima island (n=48)	0							
Kikai island (n=10)	**0.113**	0						
Okinawa (n=83)	0.027	0.079	0					
Kyushu Fukuoka (n=212)	-0.002	**0.102**	0.010	0				
Honshu Osaka (n=131)	-0.010	**0.140**	**0.031**	0.001	0			
Tottori (n=243)	-0.008	**0.117**	**0.017**	-0.003	-0.003	0		
Yamagata (n=201)	-0.011	**0.161**	**0.047**	**0.010**	-0.004	0.002	0	
Miyagi (n=296)	-0.004	**0.120**	**0.031**	0.001	-0.004	0.000	0.002	0

Data of the regions except Amami-Oshima island and Kikai island were taken from Hashiyada et al. (2008). Significant differences are in bold (p < 0.05).

Table 4. Allele frequencies of four autosamal SNPs

SNP	Gene	Allele frequencies								P value					
		Amami-Oshima island		Kikai island		Ryukyu		Hondo		Between Amami-Oshima and Kikai	Between Amami-Oshima and Ryukyu	Between Amami-Oshima and Hondo	Between Kikai and Ryukyu	Between Kikai and Hondo	Between Ryukyu and Hondo
		Allele 1	Allele 2	Allele 1	Allele 2	Allele 1	Allele 2	Allele 1	Allele 2						
rs3827760	*EDAR*	0.385	0.615	0.647	0.353	0.398	0.602	0.222	0.778	0.005	0.426	< 0.001	0.004	< 0.001	< 0.001
rs17822931	*ABCC11*	0.311	0.689	0.118	0.882	0.258	0.742	0.121	0.879	0.015	0.123	< 0.001	0.044	0.612	< 0.001
rs2070235	*MYBL2*	0.324	0.676	0.265	0.735	0.291	0.709	0.185	0.815	0.324	0.228	< 0.001	0.456	0.167	< 0.001
rs14103	*C1orf212*	0.791	0.209	0.853	0.147	0.753	0.247	0.855	0.145	0.287	0.187	0.022	0.126	0.562	< 0.001

Data of Ryukyu and Hondo clusters were taken from Yamaguchi-Kabata et al. (2008) for rs3827760 and rs17822931, and Yamaguchi-Kabata et al. (2012) for rs2070235 and rs14103.

Table 5. Mean of pairwise F_{ST} values of mtDNA and pairwise R_{ST} values of Y-STR

	mtDNA	Y-STR
	F_{ST}	R_{ST}
Amami islands	-0.007	0.113
Ryukyu Archipelago	0.019	0.073
Ryukyu and Kyushu	0.021	0.055
Japan	0.020	0.034

Data used for calculation were taken from Matsukusa et al. (2010) and Hashiyada et al. (2008) except those of the Amami islands.

Fig. 1. Map of East Asia. Amami-Oshima island and Kikai island are placed on the northern edge of the Ryukyu Archipelago. Main-island Okinawa, Miyako island and Ishigaki island are included in Okinawa. Hokkaido, Honshu and Kyushu are included in mainland Japan (Hondo).

Fig. 2. Phylogenetic network between populations based on d_A distances calculated from 337 bp of D-loop region. Data of Miyako, Ishigaki and Main-island Okinawa were taken from Matsukusa et al. (2010). Data of Honshu were taken from Oota et al. (2002). Data of Ainu, Korea and Taiwan Han were taken from Horai et al. (1996). Data of Taiwan aborigines were taken from Tajima et al. (2003).