Genome organisation and evolution of a eukaryotic nicotinate co-inducible pathway

Eszter Bokor[1+], Michel Flipphi[2#], Sándor Kocsubé[1], Judit Ámon[1], Csaba Vágvölgyi[1], Claudio Scazzocchio[3]* and Zsuzsanna Hamari[1]*

[1]University of Szeged Faculty of Science and Informatics, Department of Microbiology, Szeged, Hungary

[2] Institute de Génétique et Microbiologie, Université Paris-Sud, Orsay, France

[3]Department of Microbiology, Imperial College, London, United Kingdom and Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198, Gif-sur-Yvette, France.

#Current Address: Department of Biochemical Engineering, Faculty of Science and Technology, University of Debrecen, Debrecen, Hungary

+ These authors contributed equally

* Corresponding authors

hamari@bio.u-szeged.hu,

c.scazzocchio@imperial.ac.uk

Abstract

In *Aspergillus nidulans* a regulon including 11 *hxn* genes (*hxnS, T, R, P, Y, Z, X, W, V, M* and *N*) is inducible by a nicotinate metabolic derivative, repressible by ammonium and under stringent control of the nitrogen-state sensitive GATA factor AreA and the specific transcription factor HxnR. This is the first report in a eukaryote of the genomic organisation of a possibly complete pathway of nicotinate utilisation. In *A. nidulans* the regulon is organised in three distinct clusters, this organisation is variable in the *Ascomycota*. In some *Pezizomycotina* species all the 11 genes map in a single cluster, in other in two clusters. This variable organisation sheds light on cluster evolution. Instances of gene duplication, followed by, or simultaneous with, integration in the cluster; partial or total cluster loss; horizontal gene transfer of several genes, including an example of whole cluster re-acquisition in *Aspergillus* of section *Flavi* were detected, together with the incorporation in some clusters of genes not found in the *A. nidulans* co-regulated regulon, which underlie both the plasticity and the reticulate character of metabolic cluster evolution. This study provides a comprehensive phylogeny of six members of the cluster across representatives of all *Ascomycota* classes.

Introduction

Nicotinic acid (niacin, vitamin B3), a precursor of NAD and NADP, can be utilised by some bacteria as sole nitrogen and carbon source. The common first step in all investigated prokaryotes is the hydroxylation of nicotinic acid (NA) to 6-hydroxynicotinic acid (6-NA). The further fate of 6-NA is variable; in *Pseudomonas sp.* [1, 2] it is converted to 2,5-dihydroxypyridine (2,5-DP), in *Bacillus* sp. to 2,6-dihydroxynicotinic acid (2,6-NA) [3] and anaerobically to 1,4,5,6-tetrahydro-6-

oxonicotinic acid in *Eubacterium barkeri* (formerly *Clostridium barkeri*) [4]. The

detailed and variable further bacterial metabolic steps, whether aerobic or anaerobic

have been reviewed in [5].

The ascomycete fungus *Aspergillus nidulans* can utilise NA as sole nitrogen source. In

common with bacteria, a molybdenum cofactor (MOCO)-containing flavoprotein

catalyses the conversion of NA to 6-NA (Purine hydroxylase II, previously called

xanthine dehydrogenase II, HxnS [6-9]. The *hxnS* gene is a paralogue of *hxA,* encoding

a canonical xanthine dehydrogenase (HxA, Purine hydroxylase I, [10, 11]) which is co-

regulated with most other genes of the purine utilisation pathway ([12, 13] and

references therein). The substrate specificities of HxA and HxnS have been studied in

detail ([11] and references therein). In *A. nidulans* a NA-inducible co-regulated gene

cluster is extant (*hxn*1/VI cluster, for cluster I in chromosome VI) comprising six genes,

namely, *hxnS, hxnR* (encoding the pathway-specific transcription factor), *hxnP* and

hxnZ (encoding transporters of the Major Facilitator Superfamily, which could play a

role in the uptake of NA and/or NA-derivatives), a putative flavin oxidoreductase

(*hxnT)* and a α-ketoglutarate-dependent dioxygenase (*hxnY*) both which may be

involved in the further metabolism of 6-NA [11]. In the 1970s, NA non-utilizer mutants

were isolated and genetically characterised [6]. These map in *hxnS* and *hxnR*, but also in

a second gene cluster in chromosome VI (see below).

The *hxn1/VI* genes are specifically induced by a metabolite of NA catabolism but also

by nitrogen starvation ([11] and RNASeq data [14] available at FungiDB). Expression

of the *hxn* genes requires both the pathway-specific Zn-finger factor HxnR and the

wide-domain GATA transcription factor AreA [11]. The latter mediates de-repression

of a wide range of genes in the absence of preferred nitrogen sources (such as

ammonium, L-glutamate and L-glutamine) [15-17]. The *hxnR* gene is defined by loss of

function mutations which are non-inducible for the six genes of the cluster (including

hxnR itself) and by constitutive mutations where transcription of all *hxn1/VI* genes

occurs in the absence of inducer compounds [11]. The physiological involvement of the

hxn1/VI cluster in nicotinate metabolism is further shown by the phenotype of null

mutations in the *hxnR* gene, which result in inability to utilise nicotinate, and two of its

downstream metabolic derivatives as nitrogen sources [11].

Hereby we complete the description of the genomic organisation of the nicotinate-

inducible *hxn* genes by the identification of five additional, HxnR-dependent genes in *A.*

nidulans, and we describe variations in the genomic organisation of the eleven *hxn*

genes throughout the *Ascomycota* phylum.

The evolution of gene clustering in primary metabolism has been a subject of

discussion. Specifically, we do not know which are the factors, which lead to clustering

of previously unclustered genes, those involved in clustering maintenance and those

eventually leading to de-clustering [18]. Rokas and co-workers have proposed that

clustering confers a specific advantage when, in a given metabolic pathway one or more

intermediates are toxic as single gene loss, leading to accumulation of a toxic

metabolism will be minimised [19, 20]. Toxic intermediates, such as 2,5-DP have been

identified in the nicotinate degradation pathway of number of bacteria. Our own work to

be published elsewhere (Bokor E., Amon J., Flipphi M., Vagvolgyi C., Scazzocchio C.

and Hamari Z.) indicates that these intermediates also occur in *A. nidulans*.

Investigating the diverse organisation and evolution of the nicotinate regulon may

contribute to this debate.

Results and Discussion

Three HxnR dependent, co-inducible gene clusters are extant in *A. nidulans*.

In order to search for additional genes involved in nicotinate metabolism we investigated the cluster structure in available ascomycete genomes (see below for a thorough description). Strikingly, in *Cyphellophora europaea* (*Pezizomycotina*, *Eurotiomycetes*, *Chaetothyriales*), five additional genes (to be called *hxnV, hxnW, hxnX, hxnM* and *hxnN* see below) are positioned between *hxnP* and *hxnR* orthologues, forming a single, 11-gene cluster that includes all orthologues of the *A. nidulans hxnZ, hxnY, hxnP, hxnR, hxnT* and *hxnS* genes [11] (Fig 1, *A. nidulans* cluster 1/VI, table 1). In *A. terreus* (and several other Aspergilli, see below) *hxnV, hxnW* and *hxnX* are directly adjacent to *hxnS* (Fig 1). In *A. nidulans* a cluster including *hxnX, hxnW* and *hxnV* (cluster 2/VI) is ~40 kb distant from *hxnZ* (deduced from the re-assembled genomic sequences [21]) while *hxnM* and *hxnN* are adjacent to each other in chromosome I (cluster 3/I). While this article was being written, Martins et al. [22] suggested the clustered organisation we described for *A. terreus* and *A. nidulans* and drew comparisons with a number of other species. However, these authors did not investigate the co-regulation by nicotinate or its metabolites of the putative new *hxn* genes.

 Fig 1. Expanded clusters in *Eurotiomycetes* uncover new *hxn* genes.

Comparison of the organisation of known [11] and putative novel *hxn* genes in three

species: *A. nidulans*, *A. terreus* and *Cyphellophora europaea.* Each orthologous gene is

symbolised by a thick arrow of a different colour, which also indicates relative

orientation. Colour-coded double headed arrows connect the 5 new putative *C.*

europaea hxn genes to orthologues in the *A. nidulans* genome. Dashed lines connect

similarly arranged cluster segments in the three species. For *A. nidulans,* a double

vertical line indicates separation of clusters in different chromosomes (Super-scaffold

BN001306 for Chromosome VI, BN001301 for Chromosome I). For *A. terreus*, a single

vertical line separates two distinct contigs (Contig AAJN01000215 for the 9-gene

cluster, AAJN01000156 for the 2-gene cluster). In *C. europaea*, the 11-gene cluster is

contained in contig AOBU01000059.

Table 1. Results of i*n silico* domain analysis of modelled Hxn enzymes

gene name / annotation no. / protein length	cDNA accession number (NCBI)	name of identified domains[*] (identification code / AA interval / e-value)	proposed enzyme class
HxnY (AN11188) (349 AAs)	MT707473 this work	- PcbC (COG3491, 1-320 AAs, 3.85e-97) / DIOX_N (PF14226, 7-131 AAs, 9.5e-30) / 2OG-FeII_Oxy (PF03171, 179-282 AAs, 5.8e-22)	α-ketoglutarate dependent dioxygenase
HxnT (AN9177) (388 AAs)	MT707472 this work	- OYE-like FMN (cd02933, 9-368 AAs, 0e+00); - FadH (COG1902, 6-387 AAs, 1.12e-118)	Old Yellow Enzyme
HxnX	MN718567	-UbiH (COG0654, 17-414 AAs, 5.82e-44)	FAD dependent

(AN9161) (461 AAs)	this work	-FAD_binding_3 (PF01494, 16-235 AAs, 1.0e-09);	oxidoreductase
HxnW (AN11172) (254 AAs)	MN718568 this work	- adh_short_C2 (PF13561, 13-251 AAs, 1.2e-57);	Enoyl-(acyl carrier protein) reductase-like
HxnV (AN11187) (620 AAs)	MN718569 this work**	- PRK08294 (PRK08294, 7-620 AAs, 2.68e-93) - FAD_binding_3 (PF01494, 23-380 AAs, 3.6e-76) / UbiH (COG0654, 24-373 AAs, 9.70e-43); - PHOX_C (cd02979, 435-616 AAs, 7.26e-18) / Phe_hydrox_dim (PF07976, 404-574 AAs, 2.8e-26)	Phenol 2-monooxygenase-like enzyme
HxnM (AN6518) (307 AAs)	MN718566 this work	- CE4_HpPgdA_like (cd10938, 8-287 AAs, 1.+0e-133) - CDA1 (COG0726, 43-145 AAs; 4.03e-21)	C-N bond cleaving hydrolase-like
HxnN (AN10833) (543 AAs)	MN718565 this work	- Amidase (PF01425, 78-531 AAs, 8.5e-108)	Amidase
HxnP (AN11189) (491 AAs)	KX585439 this work**	- MFS1 (PF07690.13, 49-417 AAs, 3.2e-37)	Transporter
HxnZ (AN11196) (533 AAs)	MT707474 this work**	- MFS1 (PF07690.13, 89-513 AAs, 4.0e-24)	Transporter
HxnR (AN11197)	MT707475 this work	- two C2H2 zinc finger domains (PF00096, 8-32 AAs and 41-63 AAs, 0.029 and 0.7)	Transcription factor

		- Fungal transcription specific domain (PF04082, 394-668 AAs, 2.0e-36) [11]	[11]
HxnS (AN9178) (1396 AAs)	KX585438 Amon et al. 2017	- Fer2 (PF14111, 14-82 AAs, 1.5e-06) - Fer2_2 (PF01799, 92-174 AAs, 3.8e-25) - FAD_binding_5 (PF00941, 286-471 AAs, 8.9e-43) - CO_deh_flav_C (PF03450, 480-586 AAs, 1.5e-30) - Ald_Xan_dh_C2 (PF02738, 755-1296 AAs, 6.2e-203) ([11] and refs there in)	Xanthine dehydrogenase-type nicotinate dehydrogenase ([11] and refs there in)

* Description of the abbreviated names of protein domains: **PcbC**: Isopenicillin N
synthase and related dioxygenases; **DIOX_N**: non-haem dioxygenase in morphine
synthesis N-terminal; **2OG-FeII_Oxy**: 2OG-Fe(II) oxygenase superfamily; **OYE-like**
FMN: Old yellow enzyme (OYE)-like FMN binding domain; **FadH**: 2,4-dienoyl-CoA
reductase or related NADH-dependent reductase; **UbiH**: 2-polyprenyl-6-methoxyphenol
hydroxylase and related FAD-dependent oxidoreductases; **FAD_binding-3**: FAD
binding domain; **PRK08163**: salicylate hydroxylase; **adh_short_C2**: Enoyl-(Acyl
carrier protein) reductase; **PRK08294**: phenol 2-monooxygenase; **PHOX_C**: FAD-
dependent Phenol hydoxylase (PHOX) family, C-terminal TRX-fold domain;
Phe_hydrox_dim: Phenol hydroxylase, C-terminal dimerisation domain;
CE4_HpPgdA_like: Catalytic domain of *Helicobacter pylori* peptidoglycan
deacetylase (HpPgdA) (proposed as cyclic imidase) and similar proteins; **CDA1**:
deacetylase, PgdA/CDA1 family; **MFS1:** Major Facilitator Superfamily; **Fer2** and
Fer2_2: [2Fe-2S] binding domain; **FAD_binding_5**: FAD-binding domain;
CO_deh_flav_C: CO dehydrogenase flavoprotein C-terminal domain;
Ald_Xan_dh_C2: Molybdopterin-binding domain of aldehyde dehydrogenase.
** cDNA analysis revealed that automatic annotation was erroneous and the
experimental determination of cDNA resulted in a corrected gene model
147

The genomic organisation in Chromosome VI confirms data obtained with a mutagenic

screen, which yielded besides mutations in *hxnS* and *hxnR* [11] additional mutants

unable to grow on either NA or 6-NA as sole nitrogen sources. A number of tightly

linked mutations, of which only two (*hxn6* and *hxn7*) are still available, mapped in

chromosome VI at about ≈10 cM from mutations in the *hxnS* and *hxnR* genes, which is

coherent with the genomic organisation described above (Kelly and Scazzocchio,

personal communication).

We isolated from a genomic DNA library [23] a plasmid able to complement *hxn6* for

growth on 6-NA as sole nitrogen source. The 8256 bp insert comprises *hxnV, hxnW,*

hxnX and partial flanking sequences of the AN9159 and AN9162 loci. The *hxn6*

mutation is a G1171A transition within the *hxnV* ORF (see below for correction of the

hxnV gene model in S1 Fig) resulting in W296STOP (amber). Southern blots showed

hxn7 to be a chromosomal aberration (possibly an insertion) interrupting the *hxnV* open

reading frame (S2 Fig). In *A. nidulans*, the *hxnX* gene (cluster 2/VI) is at 40,748 bps

from *hxnZ* (based on re-assembly data [21], while *hxnN* and *hxnM* are adjacent to each

other and transcribed from the same strand in Chromosome I (Cluster 3/I) (Fig 1). We

obtained cDNAs of the genes in the three clusters, and confirmed that, as gathered by

manual inspection and comparative genomics, the database gene models (proposed by

automated annotation) for *hxnP, hxnZ* and *hxnV* are erroneous (S1, S3 and S4 Figs for

the correct gene models, Table 1 for accession numbers). HxnX, HxnW, HxnV are

oxidoreductases, while HxnM and HxnN are hydrolases. A summary of the predicted

activities of all the encoded Hxn proteins are shown in table 1. All the genes in cluster

2/VI and 3/I show an HxnR-dependent induction by 6-NA. In an $hxnR^c7$ strain, the

genes in clusters 2/VI and 3/I show variable levels of constitutive expression (Fig 2A),

as shown before for cluster 1/VI [11]. The limits of the newly detected clusters are

demarcated by the completely different pattern of expression of the flanking genes (loci

AN9159 and AN9162 for cluster 2/VI, and loci AN6517 and AN10825 for cluster 3/I;

Fig 2B).

Fig 2. HxnR dependent co-induction by 6-NA and ammonium repression of genes

in clusters 2/VI and 3/I.

All genes in clusters 2/VI and 3/I (Panel A) and the cognate cluster-flanking genes

(Panel B) were tested together with *hxnS* (in cluster VI/1), which was included as a

positive control of expression. The relative mRNA levels were measured by RT-qPCR

and data were processed according to the relative standard curve method [24] with the

γ-actin transcript (*actA*/AN6542) as reference. Mycelia were grown on 10 mM

acetamide as sole N-source for 8 h at 37 °C. They were either kept on the same medium

for a further 2 h (non-induced, NI) or induced with 1 mM 6-NA (as the sodium salt, I)

or induced as above together with 5 mM of L-(+)di-ammonium-tartrate (induced-

repressed, IR), also for 2 h. Strains used were *hxnR*[+] (FGSC A26), *hxnR*Δ (HZS.136)

and *hxnR*[c]7 (FGSC A872) (S1 Table). Standard deviations of three independent

experiments are shown. Primers are listed in S2 Table.

As previously shown for the genes in cluster 1/VI, these five newly identified *hxn* genes

are strongly ammonium repressible (Fig 2A) and with one exception (see below),

strictly dependent on the AreA GATA factor, mediating nitrogen metabolite de-

repression (Fig 3).

Fig 3. The GATA factor AreA is essential for expression of all *hxn* genes with the exception of *hxnN*.

Relative mRNA levels in *areA*[+] (FGSC A26), a supposedly derepressed *areA* mutant (*xprD1*, HZS.216) and an *areA* null mutant (*areA600*, CS3095) strains were determined (S1 Table). Non-induced conditions (NI): Strains were grown on MM media with 5 mM L-(+)di-ammonium-tartrate as sole N-source for 8 h, then the mycelia were transferred to MM with 10 mM acetamide for further 2 h. Induced conditions (I): as above but transferred to 10 mM nicotinic acid as sole N-source. Induced repressed conditions (IR): transferred to 10 mM nicotinic acid and 5 mM L-(+)di-ammonium tartrate. N-starvation conditions (St): transferred to nitrogen-source-free medium. RT-qPCR data were processed according to the standard curve method [24] with the γ-actin transcript (*actA/*AN6542) as reference. Standard deviations based on three biological replicates are shown. Primers are listed in S2 Table.

The *xprD1* is usually considered to be the most extreme de-repressed allele of the *areA* regulatory gene [25], however, it did not behave as a de-repressed allele for the expression of any *hxn* gene but rather as a partial loss of function allele for *hxnS* and *hxnP* expression [11] while being variable in its effects on the genes in clusters 2/VI and 3/I (Fig 3). A similar behaviour was reported for *ureA* (a urea transporter gene) expression [26], which strongly suggests that the phenotypes resulting from this specific mutation are promoter-dependent. The amidase-encoding *hxnN* gene shows a paradoxical pattern of expression. While it is clearly subject to repression by ammonium, it is drastically over-expressed in *areA600* background under neutral (non-induced, non-repressed conditions, see legend to Fig 3), as well as under induced and nitrogen starvation conditions (Fig 3). As *areA600* is a null mutation due to a chain

termination mutation upstream of the DNA binding domain [27], we must conclude that

AreA does not act as a transcriptional activator but as a repressor for *hxnN*. The

apparently paradoxical susceptibility of *hxnN* to ammonium repression is most probably

due to its complete dependence on HxnR, whose expression is drastically repressed by

ammonium ([11] and Figs 2 and 3). We searched the genes in the three clusters for the

consensus AreA 5'HGATAR DNA binding sites [28] (Fig 4). The *hxnV* gene upstream

sequence does not feature canonical AreA sites; nevertheless its expression is

repressible by ammonium, most likely due to indirect repression via repression of *hxnR*

transcription. The *hxnR* upstream region shows both canonical AreA sites and one

putative HxnR binding site (see below). This is consistent with this gene being

inducible, self-regulated and subject to nitrogen metabolite repression ([11] and Fig 2).

The negative effect of AreA on *hxnN* expression may be due to the presence of a

canonical GATA binding motif (5'AGATAA on the non-coding strand at position -14

to -19), interfering with the start or progress of transcription.

Fig 4. AreA and putative HxnR binding sites are extant in the 11 genes of the *hxn* regulon.

(A) Sequence logo of the DNA binding motif of the HxnR transcription factor generated by the "DNA binding site predictor for Cys2His2 Zinc Finger Proteins" application (http://zf.princeton.edu/) [29]. (B) Distribution of 5'HGATAR AreA binding sites (orange boxes) [28] and putative canonical 5'GHGGGG HxnR binding sites (dark green lozenges) in *hxn* gene promoters and also in the promoter of the *hxB* gene. The latter encodes a trans-sulphurylase necessary for the activity of the MOCO cofactor in enzymes of the xanthine oxidoreductase group (including HxnS and HxA). UaY binding sites on the *hxB* promoter are marked by blue coloured ovals [30]. Sequences conforming to the consensus 5'GHGGGG sequence are present in all HxnR-regulated genes, except *hxnN*. Nevertheless, Fig 2 shows clearly that *hxnN* is under the control of HxnR. Thus, the physiological binding sites may have a more relaxed consensus sequence. We propose 5'GNGGDG motif as a non-canonical consensus binding site

that can be found in *hxnN* as well as in other *hxn* promoters. Light green lozenges

indicate the location of the more relaxed consensus 5'GNGGDG motif. Note that the

hxnT/*hxnS*, *hxnP*/*hxnY* and *hxnX*/*hxnW* gene couples share bidirectional promoters.

The binding sites of HxnR have not been experimentally determined, however, they

could be predicted with reasonable probability [29]. Besides the consensus 5'HGATAR

AreA binding sites, Fig 4 shows also the distribution of the putative canonical and non-

canonical HxnR binding sites (5'GHGGGG and 5'GNGGDG, respectively) in all 11

hxn genes as well as in the *hxB* gene (AN1637*),* encoding a MOCO sulphurylase ([31]

for review) necessary for the enzymatic activity of both HxA and HxnS [30]. Two

putative canonical HxnR binding sites are extant in the *hxB* promoter (Fig 4). This gene

is under the independent and additive control of UaY (the transcription factor regulating

the purine utilisation pathway) and HxnR [30].

Chromosome rearrangements lead to separation of clusters 1/VI and 2/VI in *A.*

***nidulans* and other Aspergilli.**

The organisation described above for *A. terreus* (section *Terrei*) is most probably

ancestral to the Aspergilli, as is it seen in species belonging to diverging sections of this

genus (Fig 5 and S5 Fig) namely in *A. carbonarius* (section *Nigri*) and in *A. unguis,* an

early diverging species of section *Nidulantes* (Fig 5 and S5 Fig).

Fig 5. Genomic arrangement of the *hxn* gene clusters.

(A) *Aspergillaceae* (B) Other *Eurotiomycetes* compared with *Hymenoscyphus repandus* (*Leotiomycetes*, *Helotiales*). Orthologues found in different species are indicated by arrows of the same colour as in Fig 1. A single vertical line symbolises physical separation of genes on different contigs.

This organisation closely resembles the one seen in species of the basal section *Aspergillus* (*A. ruber*, *A. glaucus*, *A. chevalieri* and *A. cristatus*), where, however, the *hxnT* gene is absent. Within section *Nidulantes*, a first chromosomal inversion resulted in the separation of the clusters 1/VI and 2/VI, inverting the position and orientation of *hxnX, hxnW,* and *hxnV* (in cluster 2/VI) in relation to *hxnS* (in cluster 1/VI). This results in *A. mulundensis* in a gap of 41,992 bp between the two clusters (shown in Fig 5, section *Nidulantes*, series *Multicolores* [32]). A second inversion within this taxon led

to the configuration seen in *A. nidulans* and its closest sequenced relative *A.*

spinulosporus (section *Nidulantes*, series *Nidulantes*) [32] leaving respectively gaps of

40,876 bp and 49,972 bp between *hxnZ* and *hxnX*. *A. sydowii* and *A. versicolor* (section

Nidulantes, series *Versicolores*) [32] (S5A Fig), also show separation of clusters 1/VI

and 2/VI, however, the relative gene orientation and phylogenetic position of the latter

two species strongly suggest that this organisation arose from events independent to

those described above for *A. nidulans* and *A. spinulosporus*. Two distinct independent

inversions, like the one described above for *A. mulundensis* must have occurred within

section *Nigri,* leading to the organisation seen in *A. aculeatus* and the *A. niger* clade

(Fig 5 and S5A Fig); in *A. niger* and allied species, *hxnS* and *hxnX* are abutting

neighbours; in *A. aculeatus* (also in section *Nigri*), where *hxnT* is absent there is a ~32

kb gap between these genes.

In two species (*A. steynii* and *A. westerdijkiae*) of two closely related series (ser.

Steyniorum and ser. *Circumdati*, respectively)*,* clusters 1 and 2 are separated without

any relative change of gene orientation (S5A Fig). This could be formally described as

an insertion, however partial DNA identity and gene synteny in the inter-cluster

sequence rather suggest two successive inversions. In *A. wentii* (section *Cremei*) a

rearrangement associated with the loss of *hxnS* separates from the original cluster, a

sub-cluster including *hxnZ*, *hxnY*, and a pseudogenised *hxnP*; while *hxnV*, *hxnW* and

hxnX are still included in the main cluster together with the neighbouring *hxnT* and

hxnR.

312   **In the *Pezizomycotina*, with the exception of the Aspergilli, the *hxnN* and *hxnM***

313   **genes are included in the *hxn* cluster.**

314   The enzymes encoded in clusters 1/VI and 2/VI are all oxidoreductase enzymes,

315   however to release ammonium from NA-derived metabolites, hydrolytic enzymes are

316   necessary [2]. Within the putative *hxn* clusters of many *Pezizomycotina* species, two

317   genes encoding respectively a putative cyclic-imide hydrolase (*hxnM* EC 3.5.2.16,

318   >60% identity with AAY98498, the cyclic imide hydrolase from *Pseudomonas putida*,

319   [33]) and a putative amidase (*hxnN* EC 3.5.1.4) are extant. The cognate genes of *A.*

320   *nidulans* have been described above. In *C. europaea, hxnN* and *hxnM* lie in between

321   *hxnX* and *hxnV*, and are separated by *hxnW* constituting two neighbouring, divergently

322   transcribed gene couples, *hxnV-hxnN* and *hxnW-hxnM* within the cluster (Fig 1). It

323   should be stressed that these two divergently transcribed couples are conserved across

324   different classes of the *Pezizomycotina* (S5B Fig), however, not in the genus

325   *Aspergillus*, where with the exception of section *Flavi*, *hxnN* and *hxnM* are separated

326   from the main cluster.

327   In *Monascus ruber*, (*Eurotiomycetes*, *Eurotiales*, *Aspergillaceae* - same family as

328   *Aspergillus*) where *hxnS* is not included in a 10-strong gene cluster (see below), the two

329   divergently transcribed couples are conserved. In *Talaromyces stipitatus* and *T.*

330   *islandicus* (*Eurotiales*, *Trichocomaceae*), where *hxnS* is altogether missing) *hxnM* and

331   *hxnN* are at one terminus of the 10-strong gene cluster. Fig 5 and S5B Fig show a

332   variety of cluster organisations in species of the *Pezizomycotina* and *Saccharomycotina*

333   subphyla, with the *hxnN* and *hxnM* genes showing different patterns of integration

334   within the *hxn* cluster, with however a remarkable conservation of the *<hxnN-hxnV>*

335   and *<hxnM-hxnW>* divergently transcribed couples in classes of *Pezizomycotina*.

In the genome of *C. europaea*, besides the divergently transcribed couples mentioned above, two other couples are extant, <*hxnS-hxnT*> and <*hxnP-hxnY*>. These couples are mostly conserved in the *Pezizomycotina*, irrespective of whether all eleven genes are included in a single cluster. Noticeably, in *A. nidulans*, cluster 1/VI comprises <*hxnS-hxnT*> and <*hxnP-hxnY*>. In *Hymenoscyphus repandus* (*Leotiomycetes*, *Helotiales*), similarly to *C. europaea*, all 11 genes are included in a single mega-cluster, albeit in a different arrangement; nevertheless, two divergent couples are conserved (<*hxnS-hxnT*> and <*hxnM-hxnW*>). A similar conservation of divergently transcribed genes is seen in other gene clusters, such as the DAL cluster of the *Saccharomycetales*, where the <*DAL4-DAL1*> pair is conserved between *S. cerevisiae* and *Naumovia castellii* in spite of two inversions affecting the budding yeast DAL cluster in chromosome IX [34], and in the biotin biosynthesis cluster of the *Pezizomycotina* (<*bioF-bioDA*> [35]. The persistence of these divergently transcribed couples could be due to the fact that they share a bi-directional promoter, as established for *GAL10* and *GAL1* in *S. cerevisiae* ([36, 37] and refs therein) and for *niiA - niaD* in *A. nidulans* [38, 39].

Evolution of the *hxn* gene cluster(s) in the *Ascomycetes*.

Previous work has shown that HxnS is restricted to the *Pezizomycotina* [11]. Thus, it is unlikely that other fungi could hydroxylate NA and thus utilise it as a nitrogen source. However, it is possible that an *hxnS* gene was incorporated into a pre-existent metabolic pathway, whether catabolic or detoxifying, whether or not organised as a cluster. We thus investigated the presence of putative *hxn* clustered genes throughout the fungal kingdom. No putative *hxn* clusters are present in any early divergent fungal lineages in the *Basidiomycota* or in the *Taphrynomycotina,* except that *hxnT*, *hxnN* and *hxnM*

unlinked orthologues are present in the early diverging *Taphrinomycotina*, *Saitoella*

complicata (see S5 Fig).

Clusters comprising *hxn* genes are present in several scattered species of

Saccharomycotina (S5B Fig), however not in the *Saccharomycetaceae* and

Debaryomycetaceae families. All species of *Lipomyces,* an early divergent genus of the

Saccahromycotina, include *hxnN* and *hxnM* genes (see above). The genomes of fourteen

scattered species of *Saccharomycotina* (S5B Fig) comprise clusters with the *hxn* gene

complement, always including the transcription factor *hxnR* and never including *hxnS,*

hxnZ, and *hxnN*, even if the latter gene could be found unlinked to the cluster in an early

divergent species (*Trigonopsis variabilis*). A phylogeny of *hxnR* is shown in

supplementary S6 Fig and is consistent with a monophyletic origin of this gene in the

Saccharomycotina and *Pezizomycotina*. It seems most unlikely that the clusters of the

Saccharomycotina have a single origin. The *Lipomyces hxnM-hxnN* divergent gene pair

is found only in this genus from where all other *hxn* genes are absent. Among other

families, the occurrence of clusters with variable organisations does not follow any

obvious evolutionary pattern. In the fourteen species of *Saccharomycotina* where we

found an *hxn* cluster, the *hxnT*, *hxnR* and *hxnV* genes, are monophyletic (S5-S8 Figs).

Notwithstanding the above, the phylogeny of *hxnM* suggests several different origins of

clustered *hxnM*s within the *Saccharomycetales* from an un-clustered paralogue, possibly

acquired by HGT (see below, and S9 Fig). One clustering event occurred in the

Phaffomycetaceae, possibly two in the *Pichiaceae*, while only one species of the CUG-

Ala clade, *Pachysolen tannophilus* [40] includes an *hxn* cluster, with an *hxnM* gene.

Among the *Pichiaceae*, in the genus *Ogatea*, the monophyletic origin of clustered and

un-clustered *hxnM* genes is supported by their intron exon organisation (S9 Fig).

Several instances of gene loss, gene duplication and cluster reorganisation have

occurred in the *Pezizomycotina*. In some *Aspergillus* species, *hxnT* (encoding an FMN

dependent oxidoreductase) is missing from the cluster (S5A Fig) and indeed from the

genome. In many taxa of *Sordariomycetes* duplication of *hxnV* and subsequent loss of

the *hxn* cluster genes can be observed, leaving just the *hxnV* copy and *hxnM*.

It is striking that in the *Aspergillus* section *Flavi,* in *Talaromyces* species and in most

species of *Penicillium* the *hxnS* gene is absent and the organisation of the whole cluster

is completely identical in some species of *Talaromyces*, in most of Penicillia and in

Aspergillus section *Flavi* (S5 Fig). This coincidence indicates possible HGTs between

these taxons (see below, HGT between *Talaromyces* and *Aspergillus* section *Flavi*). As

the transcription factor-encoding gene *hxnR* is conserved, the implication is that these

organisms should be able to utilise 6-NA but not NA.

Insertion of additional genes within the *hxn* clusters

We define as "additional genes" those that appear sporadically within the *hxn* clusters of

some taxa. The insertion of a gene encoding a nitro reductase (*nfsA*) originally

horizontally transmitted from a cyanobacterium has been discussed previously [11]; the

insertion occurred after the divergence of *A. carbonarius* from other members of section

Nigri [41] (Fig 5 and S5A Fig).

In the *hxn* cluster of *Aspergillu*s section *Flavi*, and in a number of *Penicillium* and

Talaromyces species (S5, S10 and S11 Figs), a gene of unknown function, to be called

pfdB, for putative **p**eroxisomal **F**MN-dependent **d**ehydrogenase (see below) lies

between *hxnZ* and *hxnM*. This is a paralogue of *pfdA*, a gene universally present in the

Pezizomycotina, which is never included in an *hxn* cluster. The encoded proteins

include PF01070.18 (FMN-dependent dehydrogenase) and PF00173.28 (Cytochrome

b5-like binding domain) domains and have a canonical PST1 (peroxisomal entry signal

[42]. The phylogeny of PfdA and PfdB clearly suppports a scenario of gene duplication

of *pfdA* in the ancestor of Penicillia with simultaneous of subsequent cluster integration

(mean similarity between *A* and *B* paralogues 65% compared with 88% of *A*

orthologues among themselves) (S11 Fig). *pfdA* has a second, distinct paralogue, *pfdC*,

too, which however lost the PST1 signal in some cases and are only present in section

Flavi, and in a number of *Talaromyces* and *Penicilium* species and in a few species of

other clades (S10 and S11 Figs). The occurrence of PfdC in taxons is consistent with the

duplication of the PfdA ancestor in an early diverging species followed by several

episodes of loss completely unrelated to the evolution of the *hxn* cluster.

In *P. paxilli*, *P. citrinum* and *P. steckii,* a gene encoding a protein of 467-469 residues,

comprising a PF00781.24, diacylglycerol kinase catalytic domain, (orthologues

annotated as sphingoid long chain kinases) lies between the *hxnZ* and *hxnM* genes. This

gene is duplication of a gene present elsewhere in these organisms and omnipresent in

the *Eurotiomycetes*. In *Talaromyces stipitatus* a *pfdB* pseudogene is extant between

hxnZ and *hxnM*, and additionally, an intron-less gene encoding 751 residue-

multidomain protein, comprising an N terminal PF0104820.11 (phosphorylase

superfamily N-terminal, most similar to nucleoside phophorylases) domain and a C-

terminal PF05960.11 (bacterial protein of unknown function) domain is located between

hxnN and *hxnM*, the nearest homologues of the inserted gene being present and

unlinked to any *hxn* gene in *T. verruculosus.*

In *Kregervanrija fluxuum* (*Saccahromycotina*, *Pichiaceae*) a putative amidase gene is

inserted in the cluster between *hxnM* and *hxnT* (S5B Fig). The encoded protein has only

35% identity with *hxnN* of *A. nidulans*, compared with the 51% identity shown by the

genuine HxnN proteins of *Lipomyces starkei*, *Trigonopsis variabilis* and *Saitoella*

complicata. Its nearest homologue is a putative amidase from *Ogatea parapolymorpha*

(56% identity). It is tempting to speculate that this amidase has been recruited to the

cluster to carry out a similar catalytical function to that afforded by HxnN.

HGT events involving *hxn* genes

In most Penicillia and *Talaromyces* species the *hxnS* gene is absent. The un-clustered

hxnS genes of *T. islandicus*, *T. piceae*, and *T. wortmanii* have as sister clade the *hxnS* of

Monascus species, consistent with standard phylogeny. The phylogeny [11] (S5 and

S12 Figs) together with *hxnS* sequence identity strongly suggests episodes of *hxnS* de-

clustering for these three species. A different situation occurs in *P. citrinum*, *P. paxilli*

and *P. steckii*. These sister species (Section *Citrina*, [43]) have reacquired an un-

clustered *hxnS* gene by HGT from either a *Fusarium* or a *Colletotrichum* species (both

are *Sordariomycetes*, S12 Fig) [11].

In all investigated dikarya, HxnM paralogues, presumably non-related to NA

metabolism are extant. Based on comprehensive phylogeny of *hxnM* and its paralogues

(S9 Fig) subjected to reconciliation with the species tree (using GeneRax), we

confirmed HGTs amongst *Ascomycota* taxons and HGT from *Ascomycota* to

Basidiomycota. *Dothideomycetes* acquired clustered *hxnM* from *Symbiotaphrina*

(*Xylonomycetes*). Additionally, within the large clade of unclustered *hxnM* genes of the

Pezizomycotina a group containing *P. brasillianum* was found as the donor of *hxnM* to

Aspergillus section *Usti*. The common ancestor of the *Panellus stipticus* and *Mycena*

galopus (*Basidiomycota*) acquired *hxnM* from the common ancestor of the Fusaria

(*Ascomycota*). Since these two *Basidiomycetes* have only a single, Fusaria-derived

hxnM gene, the *Basidiomycota hxnM* must necessarily be lost from these species.

S9 Fig is consistent with a vertical inheritance of *hxnM* homologues in the dikarya,

excluding a recent HGT from bacteria. The phylogeny of HxnM is compatible with an

originally un-clustered *hxnM* homologue being duplicated, one copy being recruited in

an *hxn* cluster. Within the *Pezizomycotina*, there are two clades including un-clustered

hxnM homologues, one (shown by light green highlighting in S9 Fig) is basal to all the

hxnM clustered genes in both *Pezizomycotina* and *Saccharomycotina*. In the early

diverging *Lipomyces* (*Lipomycetaceae*, *Saccharomycotina*) genus, *hxnM* and *hxnN* are

clustered and divergently transcribed, and no other putative *hxn* genes are extant.

While the clustered *hxnM* genes appear monophyletic, originating from the same clade

of un-clustered genes, clustering in the *Pezizomycotina* occurred independently from

that within the *Saccharomycotina*, followed by several independent instances of

separation of an *hxnN-hxnM* minicluster (such as detailed above for the *Aspergilli*) and

presence of an *hxnM* un-clustered homologue, as it occurred in the *Leotiomycetes*.

The clade comprising the HxnM homologues of the *Saccharomycotina* seems

monophyletic. However, it does not occur as expected as a sister clade of all the

homologues of the *Pezizomycotina*, but within the different *Pezizomycotina* clades. The

low aLRT value at the relevant node, however, does not support *Saccharomycotina*

acquiring an *hxnM* gene by HGT from *Pezizomycotina*.

Reconciliation of the phylogeny of the nicotinate catabolism non-related PfdB found in

Eurotiomycetes with the species tree (by using GeneRax) confirmed that the *pfdB* of

Talaromyces which was acquired by HGT from an ancestral species of Penicillia was

further transferred from a *Talaromyces* by HGT to an ancestor of *Aspergillus* section

Flavi (Fig 6). Since all Penicillia and *Aspergillus* section *Flavi* share an identical cluster

organisation with some species of *Talaromyces*, the HGT events probably involved two

episodes of HGT of the whole *hxn* cluster. This outlines a scenario by which, after the

appearance of *pfdB* by a single gene duplication of *pfdA* in the ancestral species of

Penicillia, *pfdB* subsequently integrated into the cluster in this genus. An HGT of the

whole cluster to an early diverging species of *Talaromyces* would have occurred

followed by a further HGT from *Talaromyces* to the ancestor of *Aspergillus* section

Flavi. This scenario implies that the putative acceptor ancestor *Aspergillus* must have

lost previously the cluster present in other Aspergilli. This is strikingly confirmed by

genomes of early diverging species of section *Flavi* (*A. leporis*, *A. aliaceus*, *A.*

albertensis and *A. bertholletius*), which show both instances of *hxn* gene loss and

presence of *hxn* pseudogenes (S5A Fig). The most extreme case being that of *A.*

coremiiformis, where no *hxn* genes are present. In *A. bertholletius* a cluster of 7 *hxn*

pseudogenes is extant, where the only intact gene is *hxnT* (S5A Fig), however this gene

is not a fossil, but it derives from *Talaromyces* by HGT (S7 Fig). The earliest diverged

species of section *Flavi* is supposed to be *A. avenaceus* [44, 45]. This is fully supported

by the position of the cluster-independent *pfdA* and *pfdC* genes in the phylogenetic tree

(S10 Fig). The cluster of this species, which includes *pfdB,* is similar to that of other

Flavi, except that *hxnP* is missing and neither of the two *hxnM* paralogues is included in

the cluster.

Reconciliation analysis of HxnT, HxnR and HxnM phylogeny restricted to

Eurotiomycetes confirmed the HGT between *Talaromyces* and section *Flavi* only in the

case of HxnR, despite the layout of the phylogenetic trees of HxnM and HxnT were

basically the same as in case of PfdB and HxnR (Fig 6). In spite of contradictory results,

the evidence strongly suggests the whole HG transfer of the cluster as detailed above.

Fig 6. HGT from *Talaromyces* to *Aspergillus* section *Flavi* is supported by the phylogenies of four different proteins based on *Eurotiomycetes* data set (see S6, S7, S9 and S10 Figs. for the complete phylogenies).

Cyan: *Talaromyces*; Blue: *Penicillium*; Red: *Aspergillus* section *Flavi*; Purple: *Aspergillus* section *Polypaecilum*; Light brown: *Aspergillus* section *Tannerorum*; Green: *Aspergillus* section *Terrei*; Gray: *Aspergillus* section *Cremei*; Black: *Aspergillus* section *Flavipedes*.

Disturbingly, in the *hxnR*, *hxnV* and *hxnT* phylogenies, *A. avenaceus* appears as out-species of the *Talaromyces*/*Penicillium* clade which transferred the cluster to other *Flavi* (Fig 6 and S6-S8 figs.). There is obviously a complex series of HGTs which may be solved when more genomes of closely related species become available.

A. sclerotialis (subgenus *Polypaecilum*) has an *hxn* cluster of complex origin. Supported

by phylogeny of HxnS, HxnR, HxnV and HxnT, the corresponding genes derived by

HGT from a *Fusarium* (*Nectria*) species (S6-S8 and S12 Figs). Its cluster includes a

pfdB gene, necessarily derived from a *Talaromyces/Penicillium* species. Its clustered

hxnM1 is sister to the *hxnM1* of *A. tanneri* (see below) (Fig 6, S9 Figs). The most

parsimonious hypothesis is that the complex *hxn* cluster of this species originated in the

confluence of two HGT events, one from a *Fusarium*/*Nectria* species and another from

a *Talaromyces/Penicillium* species, together with an extensive rearrangement of the

cluster, leading to a unique pattern of gene organisation.

Aspergillus tanneri belongs to section *Tannerorum*, a sister clade to section *Circumdati*

[32, 43]. In this species two clusters and an isolated *hxnM-hxnN* pair are extant (S5 Fig).

Five enzyme-encoding genes are present in two copies, *hxnV*, *hxnX* and *hxnM*, *hxnS* and

hxnW. No *hxnZ, hxnY* and *hxnP* genes are extant in *A. tanneri*. The *hxnS* gene present in

both clusters is a product of a recent duplication (S12 Fig). In the larger cluster (7

genes), *pfdB* originates from an HGT from *Talaromyces* or *Penicillium* (Fig 6), most

likely independently from the HGT to section *Flavi*, *hxnV1, hxnM1* and *hxnT* genes,

sister to those extant in *A. avenaceus,* are likely of the same origin, while *hxnR* has not

originated from an HGT event (S6-S10 Figs). This is a composite cluster, where HGT

events are coalesced with vertically inherited genes. The smaller cluster includes an

hxnV2 paralogue beside an *hxnS* paralogue that also originated from a recent duplication

together with an *hxnW* pseudogene (S8 and S12 Figs).

Concluding remarks

Experimental work has shown that three gene clusters in *A. nidulans* constitute a

nicotinate (actually a nicotinate derivative) inducible regulon, under the control of a

specific Zn-finger transcription factor, HxnR. Deletion of HxnR have shown that

expression of some or all of the genes in this regulon are necessary for NA, 6-NA and

the putative intermediate 2,5-dihydroxypyridine utilisation as nitrogen sources [11]. The

specific metabolic function of each encoded protein will be reported separately. This

regulon is extant only in the *Ascomycetes*. The variable organisation seen in different

species includes instances of complete clustering of all 11 genes, which may suggest an

evolutionary pressure towards the global integration of the *hxn* genes, together with

instances of de-clustering such as the separation of clusters 1(VI) and 2(VI) in section

Nidulantes of the Aspergilli. Several instances of HGT were detected (Fig 7), most

notably the origin of the cluster of *Aspergillus* section *Flavi* from

Talaromyces/Penicillia. The events of HGT, together with the recruitment of genes after

duplication, including *hxnS* and *hxnM*, and additional genes such as *pfdB*, underlies

both the dynamic nature and the reticulate character of metabolic cluster evolution.

Fig 7. Summary of supported HGT events on collapsed species tree related to *pfdB*,

***hxnM* and *hxnS* genes.**

F: *Aspergillus* section *Flavi*; U: *Aspergillus* section *Usti*; P: Penicillia*;* T: early

diverging species of *Talaromyces*; 1: HGT of *pfdB* gene found between Penicillia and

species of *Talaromyces* and between species of *Talaromyces* and *Aspergillus* section

Flavi; 2: HGTs of *hxnM* gene between *Xylonomycetes* and *Dothideomycetes*, Fusaria

(*Sordariomycetes*) and *Basidiomycota* and between a group of Penicillia containing *P.*

567   *brasilianum* and *Aspergillus* section *Usti*; 3: HGT of *hxnS* gene between *Aspergillus*

568   section *Usti* and Penicillia ; red asterisk: transfer of the whole *hxn* cluster composed of

569   nine *hxn* genes and including the *pfdB* gene from Penicillia to species of *Talaromyces*

570   and from *Talaromyces* to *Aspergillus* section *Flavi*. Solid lines mark confirmed HGTs.

Materials and Methods

Strains and growth conditions

The *A. nidulans* strains used in this work are listed in S1 Table. Standard genetic

markers are described in http://www.fgsc.net/Aspergillus/gene_list/. Minimal media

(MM) contained glucose as the carbon source; the nitrogen source varied according to

the experimental condition [11]. The media were supplemented according to the

requirements of each auxotrophic strain (www.fgsc.net). Nitrogen sources, inducers and

repressors were used at the following concentrations: 10 mM acetamide, 10 mM

nicotinic acid (1:100 dilution from 1 M nicotinic acid dissolved in 1 M sodium

hydroxide) and 5 mM L-(+)di-ammonium-tartrate as sole N-sources; 1 mM 6-

hydroxynicotinic acid sodium salt as inducer and 5 mM L-(+)di-ammonium-tartrate as

repressor. Growth conditions are detailed in the figure legends of corresponding

experiments.

RNA manipulation

Total RNA was isolated using a NucleoSpin RNA Plant Kit (Macherey-Nagel) and

RNase-Free DNase (Qiagen) according to the manufacturer's instructions. cDNA

synthesis was carried out with a mixture of oligo-dT and random primers using a

RevertAid First Strand cDNA Synthesis Kit (Fermentas). Quantitative RT-PCR (RT-

qPCR) were carried out in a CFX96 Real Time PCR System (BioRad) with SYBR

Green/Fluorescein qPCR Master Mix (Fermentas) reaction mixture (94 °C 3 min

followed by 40 cycles of 94 °C 15 s and 60 °C 1 min). Data processing was done by the

standard curve method [24]. DNA sequencing was done by the Sanger sequencing

service of LGC (http://www.lgcgroup.com). Primers used are listed in the S2 Table.

Data mining

The coding sequences of fungal *hxn* genes (ATG-STOP) were mined by TBLASTN

screening of DNA databases at the NCBI servers, mainly the Whole Genome Shotgun

contigs (WGS) database, using the available on-line tools [46]. For a few species (*N.*

crassa, *P. anserina*, *P. chrysogenum*, *A. oryzae*, *A. niger* ATCC 1015, *Leptosphaeria*

maculans and some *Saccharomycotina*), the sequence contings of the published genome

are located in the nr/nt database or the Refseq genome database. Additional *Eurotiales*

genomes (outside *Aspergillaceae*) are publicly accessible at the website of the Centre

for Structural and Functional Genomics (Concordia University Montreal, Canada;

https://gb.fungalgenomics.ca/portal/). We also included some species from the 1000

Fungal Genomes Project (http://1000.fungalgenomes.org) exclusively available at the

Mycocosm database (Joint Genome Institute, US Department of Energy)

(https://mycocosm.jgi.doe.gov/mycocosm/home). For the two classes of *Pezizomycotina*

for which few genome sequences are public (*Xylonomycetes*, *Pezizomycetes*), we have

obtained permission to use the *hxn* complement in the genome sequences of five species

lodged at JGI in our current work: *Symbiotaphrina kochii* (Project ID: 404190);

Trinosporium guianense (Project ID: 1040180); *Gyromitra esculenta* (Project ID:

1051239); *Plectania melastoma* (Project ID: 1040543); and *Sarcoscypha coccinea*

(Project ID: 1042915). TBLASTN query sequences for the 11 *hxn* genes were the full-

length proteins deduced from the cDNA sequences we experimentally determined for

each of the *A. nidulans hxn* genes (see Table 1 for GenBank Accession numbers).

Where necessary, to confirm gene orthology amongst multiple homologous sequences,

the TBLASTN hits and their surrounding sequences were further inspected for the

conservation of occupied intron positions between species and for synteny with other

hxn genes in the sequence contig identified (gene clustering). We did not use the results

of automated annotation ("Models" or "mRNA" at nr/nt) nor did we use deduced

protein databases for the eukaryotic (Hxn) proteins. We used a selection of

autoannotated proteins for the prokaryote HxnM outgroup extracted from the nr/nt

database, using the *Pseudomonas putida* cyclic imide hydrolase (GenBank AAY98498:

[33]) as the BLASTP query. We manually predicted the intron-exon structure of each

(*hxn*) gene, guided by comparative genomics and after (*in silico*) intron removal,

deduced the encoded proteins subsequently used in phylogenetic analyses (see below).

Alternative yeast nuclear codes were utilised where appropriate (*Pachysolen*:

CUG=Ala, *Priceomyces*: CUG=Ser). For some species in under-represented taxa, we

could use the Transcriptome Shotgun Assembly (TSA) database to obtain intronless

sequences coding for full-length protein.

Construction or Maximum likelihood trees.

Criteria for identification of orthologues/paralogues is detailed for each tree.

Alignments were done with MAFFT G-INS-i unless otherwise indicated, with default

parameters [47, 48] (https://mafft.cbrc.jp/alignment/server/). Alignments were trimmed

with BMGE with default parameters unless otherwise indicated

(https://ngphylogeny.fr/workflows/wkmake/42f42d079b0a46e9, [49]. Maximum

likelihood trees were constructed with PhyML 3.0. Automatic model selection by SMS

(http://www.atgc-montpellier.fr/phyml [50, 51]) and drawn with FigTree v 1.4.4.

Values at nodes of all trees are aLRTs (Approximate Likelihood ratio test, [52]). All

trees are shown in a circular cartooned form. Trees are rooted in the specified out group.

Reconciliation was done by GeneRax v1.2.3, a Maximum Likelihood based method

[53] with default settings in 500 replicates. Only those transfers were considered, which

were present in at least 70% of the replicates. Species tree for the reconciliation was

drawn after [54, 55].

Statements

Statements

Data accessibility

The datasets supporting this article are included in the paper and detailed in the

electronic supplementary material tables. Sequences determined by us are available on

GenBank accession numbers: MT707473, MT707472, MN718567, MN718568,

MN718569, MN718566, MN718565, KX585439, MT707474, MT707475.

Authors' contributions

ZH and CS conceived the project. EB, ZH and JA contributed to various aspects of the

wet laboratory work; ZH and CS wrote the manuscript. MF discovered the additional

two clusters in A. nidulans, manually curated gene models of hxnV, hxnP and hxnZ

orthologs and constructed the schemes of hxn clusters for hundreds of species. CV

contributed to in silico promoter analysis. CS and SK did the phylogenetic analysis. All

authors analysed the results and gave final approval for publication.

665 Competing interests

666 We have no competing interests.

668 Funding

669 Work was supported by the Hungarian National Research, Development and Innovation

670 Office (NKFIH K16-119516) and by the Hungarian Government (GINOP-2.3.2-15-

671 2016-00012).

673 **Acknowledgement**

674 JGI sequences used in the construction of the phylogenetic tree were from the US

675 Department of Energy Joint Genome Institute (http://www.jgi.doe.gov/) in collaboration

676 with the user community. We thank I. V. Grigoriev for permitting the use of genome

677 sequences included in the 1000 Fungal Genomes project and we thank J. Spatafora, J.

678 Magnuson and R. Gazis for allowing access to the genomes of some individual species

679 prior to publication (Project IDs: 404190, 1040180, 1051239, 1040543 and 1042915).

680 We thank Prof. Joan M. Kelly for allowing us to cite her early genetics work. We thank

681 László G. Nagy for critical comments on the manuscript.

683 **References**

684 1 Behrman, E. J., Stanier, R. Y. 1957 The bacterial oxidation of nicotinic acid. *The*

685 *Journal of biological chemistry*. **228**, 923-945.

686 2 Jimenez, J. I., Canales, A., Jimenez-Barbero, J., Ginalski, K., Rychlewski, L., Garcia,

687 J. L., Diaz, E. 2008 Deciphering the genetic determinants for aerobic nicotinic acid

degradation: the nic cluster from *Pseudomonas putida* KT2440. *Proceedings of the*

National Academy of Sciences of the United States of America. **105**, 11329-11334.

(10.1073/pnas.0802273105

0802273105 [pii])

3 Ensign, J. C., Rittenberg, S. C. 1964 The Pathway of Nicotinic Acid Oxidation by a

Bacillus Species. *The Journal of biological chemistry*. **239**, 2285-2291.

4 Alhapel, A., Darley, D. J., Wagener, N., Eckel, E., Elsner, N., Pierik, A. J. 2006

Molecular and functional analysis of nicotinate catabolism in *Eubacterium barkeri*.

Proceedings of the National Academy of Sciences of the United States of America. **103**,

12341-12346. (0601635103 [pii]

10.1073/pnas.0601635103)

5 Andreesen, J. R., Fetzner, S. 2002 The molybdenum-containing hydroxylases of

nicotinate, isonicotinate, and nicotine. *Met Ions Biol Syst*. **39**, 405-430.

6 Scazzocchio, C. 1973 The genetic control of molybdoflavoproteins in Aspergillus

nidulans. II. Use of NADH dehydrogenase activity associated with xanthine

dehydrogenase to investigate substrate and product inductions. *Mol Gen Genet*. **125**,

147-155.

7 Lewis, N. J., Hurt, P., Sealy-Lewis, H. M., Scazzocchio, C. 1978 The genetic control

of the molybdoflavoproteins in *Aspergillus nidulans*. IV. A comparison between purine

hydroxylase I and II. *Eur J Biochem*. **91**, 311-316.

8 Mehra, R. K., Coughlan, M. P. 1984 Purification and properties of purine hydroxylase

II from *Aspergillus nidulans*. *Arch Biochem Biophys*. **229**, 585-595. (0003-

9861(84)90191-7 [pii])

9 Scazzocchio, C., Holl, F. B., Foguelman, A. I. 1973 The genetic control of

molybdoflavoproteins in *Aspergillus nidulans*. Allopurinol-resistant mutants

constitutive for xanthine-dehydrogenase. *Eur J Biochem*. **36**, 428-445.

10 Glatigny, A., Scazzocchio, C. 1995 Cloning and molecular characterization of *hxA*,

the gene coding for the xanthine dehydrogenase (purine hydroxylase I) of *Aspergillus*

nidulans. *The Journal of biological chemistry*. **270**, 3534-3550.

11 Amon, J., Fernandez-Martin, R., Bokor, E., Cultrone, A., Kelly, J. M., Flipphi, M.,

Scazzocchio, C., Hamari, Z. 2017 A eukaryotic nicotinate-inducible gene cluster:

convergent evolution in fungi and bacteria. *Open Biol*. **7**, 170199. (170199 [pii]

10.1098/rsob.170199

rsob.170199 [pii])

12 Scazzocchio, C., Darlington, A. J. 1968 The induction and repression of the enzymes

of purine breakdown in *Aspergillus nidulans*. *Biochimica et biophysica acta*. **166**, 557-

568. (0005-2787(68)90243-8 [pii]

10.1016/0005-2787(68)90243-8)

13 Gournas, C., Oestreicher, N., Amillis, S., Diallinas, G., Scazzocchio, C. 2011

Completing the purine utilisation pathway of *Aspergillus nidulans*. *Fungal genetics and*

biology : FG & B. **48**, 840-848. (10.1016/j.fgb.2011.03.004

S1087-1845(11)00066-1 [pii])

14 Sibthorp, C., Wu, H., Cowley, G., Wong, P. W., Palaima, P., Morozov, I. Y.,

Weedall, G. D., Caddick, M. X. 2013 Transcriptome analysis of the filamentous fungus

Aspergillus nidulans directed to the global identification of promoters. *BMC Genomics*.

14, 847. (10.1186/1471-2164-14-847

1471-2164-14-847 [pii])

15 Arst, H. N., Jr., Cove, D. J. 1973 Nitrogen metabolite repression in *Aspergillus*

nidulans. *Mol Gen Genet*. **126**, 111-141. (10.1007/BF00330988)

16 Wilson, R. A., Arst, H. N., Jr. 1998 Mutational analysis of AREA, a transcriptional

activator mediating nitrogen metabolite repression in *Aspergillus nidulans* and a

member of the "streetwise" GATA family of transcription factors. *Microbiol Mol Biol*

Rev. **62**, 586-596.

17 Scazzocchio, C. 2000 The fungal GATA factors. *Curr Opin Microbiol*. **3**, 126-131.

(S1369-5274(00)00063-1 [pii])

18 Nutzmann, H. W., Scazzocchio, C., Osbourn, A. 2018 Metabolic Gene Clusters in

Eukaryotes. *Annu Rev Genet*. **52**, 159-183. (10.1146/annurev-genet-120417-031237)

19 McGary, K. L., Slot, J. C., Rokas, A. 2013 Physical linkage of metabolic genes in

fungi is an adaptation against the accumulation of toxic intermediate compounds.

Proceedings of the National Academy of Sciences of the United States of America. **110**,

11481-11486. (10.1073/pnas.1304461110

1304461110 [pii])

20 Rokas, A., Wisecaver, J. H., Lind, A. L. 2018 The birth, evolution and death of

metabolic gene clusters in fungi. *Nat Rev Microbiol*. **16**, 731-744. (10.1038/s41579-

018-0075-3

10.1038/s41579-018-0075-3 [pii])

21 Wortman, J. R., Gilsenan, J. M., Joardar, V., Deegan, J., Clutterbuck, J., Andersen,

M. R., Archer, D., Bencina, M., Braus, G., Coutinho, P.*, et al.* 2009 The 2008 update of

the *Aspergillus nidulans* genome annotation: A community effort. *Fungal Genetics and*

Biology. **46**, S2-S13. (10.1016/j.fgb.2008.12.003)

22 Martins, T. M., Martins, C., Guedes, P., Silva Pereira, C. 2021 Twists and Turns in

the Salicylate Catabolism of *Aspergillus terreus*, Revealing New Roles of the 3-

Hydroxyanthranilate Pathway. *mSystems*. **6**, (e00230-20 [pii]

10.1128/mSystems.00230-20

6/1/e00230-20 [pii])

23 Osherov, N., May, G. 2000 Conidial germination in *Aspergillus nidulans* requires

RAS signaling and protein synthesis. *Genetics*. **155**, 647-656.

24 Larionov, A., Krause, A., Miller, W. 2005 A standard curve based method for

relative real time PCR data processing. *BMC Bioinformatics*. **6**, 62. (1471-2105-6-62

[pii]

10.1186/1471-2105-6-62)

25 Platt, A., Langdon, T., Arst, H. N., Jr., Kirk, D., Tollervey, D., Sanchez, J. M.,

Caddick, M. X. 1996 Nitrogen metabolite signalling involves the C-terminus and the

GATA domain of the *Aspergillus* transcription factor AREA and the 3' untranslated

region of its mRNA. *The EMBO journal*. **15**, 2791-2801.

26 Abreu, C., Sanguinetti, M., Amillis, S., Ramon, A. 2010 UreA, the major urea/H+

symporter in *Aspergillus nidulans*. *Fungal genetics and biology : FG & B*. **47**, 1023-

1033. (10.1016/j.fgb.2010.07.004

S1087-1845(10)00130-1 [pii])

27 Kudla, B., Caddick, M. X., Langdon, T., Martinez-Rossi, N. M., Bennett, C. F.,

Sibley, S., Davies, R. W., Arst, H. N., Jr. 1990 The regulatory gene *areA* mediating

nitrogen metabolite repression in *Aspergillus nidulans*. Mutations affecting specificity

of gene activation alter a loop residue of a putative zinc finger. *The EMBO journal*. **9**,

1355-1364.

28 Ravagnani, A., Gorfinkiel, L., Langdon, T., Diallinas, G., Adjadj, E., Demais, S.,

Gorton, D., Arst, H. N., Jr., Scazzocchio, C. 1997 Subtle hydrophobic interactions

between the seventh residue of the zinc finger loop and the first base of an HGATAR

sequence determine promoter-specific recognition by the *Aspergillus nidulans* GATA

factor AreA. *The EMBO journal*. **16**, 3974-3986. (10.1093/emboj/16.13.3974)

29 Persikov, A. V., Singh, M. 2014 De novo prediction of DNA-binding specificities

for Cys2His2 zinc finger proteins. *Nucleic acids research*. **42**, 97-108.

(10.1093/nar/gkt890

gkt890 [pii])

30 Amrani, L., Cecchetto, G., Scazzocchio, C., Glatigny, A. 1999 The *hxB* gene,

necessary for the post-translational activation of purine hydroxylases in *Aspergillus*

nidulans, is independently controlled by the purine utilization and the nicotinate

utilization transcriptional activating systems. *Molecular microbiology*. **31**, 1065-1073.

31 Schwarz, G., Mendel, R. R., Ribbe, M. W. 2009 Molybdenum cofactors, enzymes

and pathways. *Nature*. **460**, 839-847. (10.1038/nature08302

nature08302 [pii])

32 Houbraken, J., Kocsube, S., Visagie, C. M., Yilmaz, N., Wang, X. C., Meijer, M.,

Kraak, B., Hubka, V., Bensch, K., Samson, R. A*., et al.* 2020 Classification of

Aspergillus, *Penicillium*, *Talaromyces* and related genera (*Eurotiales*): An overview of

families, genera, subgenera, sections, series and species. *Stud Mycol*. **95**, 5-169.

(10.1016/j.simyco.2020.05.002

S0166-0616(20)30012-9 [pii])

33 Shi, Y. W., Cui, L. F., Yuan, J. M. 2007 Gene cloning, expression, and substrate

specificity of an imidase from the strain *Pseudomonas putida* YZ-26. *Curr Microbiol*.

55, 61-64. (10.1007/s00284-005-0455-6)

34 Naseeb, S., Delneri, D. 2012 Impact of chromosomal inversions on the yeast DAL

cluster. *PLoS One*. **7**, e42022. (10.1371/journal.pone.0042022

PONE-D-12-15144 [pii])

35 Magliano, P., Flipphi, M., Sanglard, D., Poirier, Y. 2011 Characterization of the

Aspergillus nidulans biotin biosynthetic gene cluster and use of the *bioDA* gene as a

new transformation marker. *Fungal genetics and biology : FG & B*. **48**, 208-215.

(10.1016/j.fgb.2010.08.004

S1087-1845(10)00150-7 [pii])

36 Johnston, M., Davis, R. W. 1984 Sequences that regulate the divergent GAL1-

GAL10 promoter in *Saccharomyces cerevisiae*. *Molecular and cellular biology*. **4**,

1440-1448. (10.1128/mcb.4.8.1440)

37 Slot, J. C., Rokas, A. 2010 Multiple GAL pathway gene clusters evolved

independently and by different mechanisms in fungi. *Proceedings of the National*

Academy of Sciences of the United States of America. **107**, 10136-10141.

(10.1073/pnas.0914418107

0914418107 [pii])

38 Punt, P. J., Strauss, J., Smit, R., Kinghorn, J. R., van den Hondel, C. A.,

Scazzocchio, C. 1995 The intergenic region between the divergently transcribed *niiA*

and *niaD* genes of *Aspergillus nidulans* contains multiple NirA binding sites which act

bidirectionally. *Molecular and cellular biology*. **15**, 5688-5699.

(10.1128/mcb.15.10.5688)

39 Muro-Pastor, M. I., Gonzalez, R., Strauss, J., Narendja, F., Scazzocchio, C. 1999

The GATA factor AreA is essential for chromatin remodelling in a eukaryotic

bidirectional promoter. *The EMBO journal*. **18**, 1584-1597. (10.1093/emboj/18.6.1584)

40 Shen, X. X., Opulente, D. A., Kominek, J., Zhou, X., Steenwyk, J. L., Buh, K. V.,

Haase, M. A. B., Wisecaver, J. H., Wang, M., Doering, D. T.*, et al.* 2018 Tempo and

Mode of Genome Evolution in the Budding Yeast Subphylum. *Cell*. **175**, 1533-1545

e1520. (S0092-8674(18)31332-1 [pii]

10.1016/j.cell.2018.10.023)

41 Varga, J., Frisvad, J. C., Kocsube, S., Brankovics, B., Toth, B., Szigeti, G., Samson,

R. A. 2011 New and revisited species in *Aspergillus* section *Nigri*. *Stud Mycol*. **69**, 1-

17. (10.3114/sim.2011.69.01)

42 Gould, S. J., Keller, G. A., Hosken, N., Wilkinson, J., Subramani, S. 1989 A

conserved tripeptide sorts proteins to peroxisomes. *The Journal of cell biology*. **108**,

1657-1664. (10.1083/jcb.108.5.1657)

43 Steenwyk, J. L., Shen, X. X., Lind, A. L., Goldman, G. H., Rokas, A. 2019 A Robust

Phylogenomic Time Tree for Biotechnologically and Medically Important Fungi in the

Genera *Aspergillus* and *Penicillium*. *mBio*. **10**, (e00925-19 [pii]

10.1128/mBio.00925-19

mBio.00925-19 [pii])

44 Frisvad, J. C., Hubka, V., Ezekiel, C. N., Hong, S.-B., Novakova, A., Chen, A. J.,

Arzanlou, M., Larsen, T. O., Sklenar, F., Mahakarnchanakul, W.*, et al.* 2019 Taxonomy

of *Aspergillus* section *Flavi* and their production of aflatoxins, ochratoxins and other

mycotoxins. *Stud Mycol*. **93**, 1-63. (10.1016/j.simyco.2018.06.001)

45 Kjaerbolling, I., Vesth, T., Frisvad, J. C., Nybo, J. L., Theobald, S., Kildgaard, S.,

Petersen, T. I., Kuo, A., Sato, A., Lyhne, E. K.*, et al.* 2020 A comparative genomics

study of 23 *Aspergillus* species from section *Flavi*. *Nat Commun*. **11**, 1106.

(10.1038/s41467-019-14051-y

10.1038/s41467-019-14051-y [pii])

46 Altschul, S. F., Gish, W., Miller, W., Myers, E. W., Lipman, D. J. 1990 Basic local

alignment search tool. *Journal of molecular biology*. **215**, 403-410. (10.1016/S0022-

2836(05)80360-2

S0022-2836(05)80360-2 [pii])

47 Kuraku, S., Zmasek, C. M., Nishimura, O., Katoh, K. 2013 aLeaves facilitates on-

demand exploration of metazoan gene family trees on MAFFT sequence alignment

server with enhanced interactivity. *Nucleic acids research*. **41**, W22-28.

(10.1093/nar/gkt389

gkt389 [pii])

48 Katoh, K., Rozewicki, J., Yamada, K. D. 2019 MAFFT online service: multiple

sequence alignment, interactive sequence choice and visualization. *Brief Bioinform*. **20**,

1160-1166. (10.1093/bib/bbx108

4106928 [pii])

49 Criscuolo, A., Gribaldo, S. 2010 BMGE (Block Mapping and Gathering with

Entropy): a new software for selection of phylogenetic informative regions from

multiple sequence alignments. *BMC Evol Biol*. **10**, 210. (10.1186/1471-2148-10-210

1471-2148-10-210 [pii])

50 Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O.

2010 New algorithms and methods to estimate maximum-likelihood phylogenies:

assessing the performance of PhyML 3.0. *Syst Biol*. **59**, 307-321.

(10.1093/sysbio/syq010

syq010 [pii])

51 Lefort, V., Longueville, J. E., Gascuel, O. 2017 SMS: Smart Model Selection in

PhyML. *Mol Biol Evol*. **34**, 2422-2424. (10.1093/molbev/msx149

3788860 [pii])

52 Anisimova, M., Gascuel, O. 2006 Approximate likelihood-ratio test for branches: A

fast, accurate, and powerful alternative. *Syst Biol*. **55**, 539-552. (T808388N86673K61

[pii]

10.1080/10635150600755453)

53 Morel, B., Kozlov, A. M., Stamatakis, A., Szollosi, G. J. 2020 GeneRax: A Tool for

Species-Tree-Aware Maximum Likelihood-Based Gene Family Tree Inference under

Gene Duplication, Transfer, and Loss. *Mol Biol Evol*. **37**, 2763-2774.

(10.1093/molbev/msaa141

5851843 [pii])

54 Shen, X. X., Steenwyk, J. L., LaBella, A. L., Opulente, D. A., Zhou, X., Kominek,

J., Li, Y., Groenewald, M., Hittinger, C. T., Rokas, A. 2020 Genome-scale phylogeny

and contrasting modes of genome evolution in the fungal phylum *Ascomycota*. *Sci Adv*.

6, (eabd0079 [pii]

10.1126/sciadv.abd0079

6/45/eabd0079 [pii])

55 Prasanna, A. N., Gerber, D., Kijpornyongpan, T., Aime, M. C., Doyle, V. P., Nagy,

L. G. 2020 Model Choice, Missing Data, and Taxon Sampling Impact Phylogenomic

Inference of Deep *Basidiomycota* Relationships. *Syst Biol*. **69**, 17-37.

(10.1093/sysbio/syz029

5486399 [pii])

56 Neuberger, G., Maurer-Stroh, S., Eisenhaber, B., Hartig, A., Eisenhaber, F. 2003

Prediction of peroxisomal targeting signal 1 containing proteins from amino acid

sequence. *Journal of molecular biology*. **328**, 581-592. (S002228360300319X [pii]

10.1016/s0022-2836(03)00319-x)

Supporting information

S1 Fig. Exon-intron organisation of the *hxnV* coding region based on cDNA sequencing and deduced HxnV protein sequence.

(A) Exon-intron organisation of the coding region deduced through comparison of the sequenced cDNA with the genomic DNA sequence. Introns (lower case letters) are highlighted in blue (splice site consensus sequences: 5'-donor, lariat branch point sequence, and 3'-acceptor) and grey (other intronic sequences). cDNA (GenBank: MN718569) confirms the gene model. (B) Peptidic HxnV sequence deduced from the cDNA sequence.

S1 Table. *A. nidulans* strains used in this work. (All strains are *veA1* mutant)

S2 Fig. Analysis of the *hxn7* mutation by Southern blot.

(A) Location of the second *hxn* cluster on chromosome VI (cluster 2/VI, comprising the *hxnX*, *hxnW* and *hxnV genes* shown as blue, purple and yellow arrows, respectively) within a 50 kb genomic sequence (part of TPA Accession number BN001306). The figure shows the cleavage sites of DraI (D),NdeI (Nd), SalI (S), XbaI (X), NcoI (Nc), PvuII (P) and BamHI (B) restriction endonucleases that were used in Southern blot analysis (see Panel C). The gene probe used in Southern blots (labelled as a yellow box above the *hxnV* gene) was a 486 bp fragment of *hxnV* (probe), obtained by using "hxnV AS frw" (cagcgtcaagtctcatatctatactg) and "hxnV AS rev" (cagagcacgggtacaaagaaggtg) as PCR primers. Yellow arrows above the 50 kb genomic region show, for each enzyme used, the endonuclease-cleaved fragments that hybridize with the probe. (B) Predicted

fragments obtained by restriction of the 50 kb genomic region described above. (C)

Southern blots result with the HZS.145 control (hxn^+) and HZS.697 $hxn7$ mutant ($hxn7$)

strains. Southern hybridization was carried out with the DIG-DNA labelling- and

detection kit (Roche) on restriction endonuclease (listed in Panels A and B) digested

total DNA of the hxn^+ and $hxn7$ strains. M: DIG-labelled DNA Molecular Weight

Marker (Fermentas). The DraI digest excludes that $hxn7$ could be a sizeable deletion.

An inversion internal to the ≈15 kb DraI fragment is excluded by several digests such as

BamHI and NdeI. A translocation or more likely an insertion is compatible with the

results shown.

S2 Table. Primers used in this work.

S3 Fig. Correction of the gene model for the *A. nidulans hxnP* gene, locus

AN11189.

(A) The manually predicted exon-intron organisation of the coding region: introns

(lower case letters) highlighted in grey; donor-, lariat- and acceptor sequences

highlighted in blue. cDNA (accession KX585439) confirms this gene model. (B)

Peptidic sequence of HxnP. The 4th intron splits a Met codon highlighted in purple in

both panels, and this feature is conserved throughout the *Aspergillaceae* as well as in

Talaromyces. Both the *Aspergillus* database (http://www.aspergillusgenome.org/cgi-

bin/locus.pl?locus=AN11189&organism=A_nidulans_FGSC_A4) and the NCBI nr/nt

database (gb|AACD01000170.1|:21152-23003 and gb|BN001306.1|:193501-195352;

locus identifiers AN9176.2 and ANIA_11189, respectively) feature an erroneous gene

model resulting in an incorrect HxnP protein sequence (accession numbers XP_682445

and EAA61467; TPA accession number CBF82384).

S4 Fig. Exon-intron structure of the *hxnZ* coding region based on cDNA sequencing, and deduced HxnZ protein sequence.

(A) Exon-intron organisation of the coding region deduced through the comparison of the sequenced cDNA with the genomic DNA sequence. Introns (lower case letters) are highlighted in grey; the donor-, lariat- and acceptor sequences are highlighted in blue. cDNA (GenBank: MT707474) confirms this gene model. (B) Peptidic sequence of HxnZ deduced from the cDNA sequence.

S5 Fig. Distribution of hxn genes in gene clusters in selected *Eurotiomycetes* (Panel A) and in other classes of *Pezizomycotina* as well as in *Saccharomycotina* (Panel B).

Colour coded arrows indicate specific hxn genes and relative gene orientation, as detailed: *hxnN* (dark green), *hxnM* (pink), *hxnS* (grey), *hxnT* (khaki), *hxnR* (red), *hxnP* (light green), *hxnY* (mid blue), *hxnZ* (orange), *hxnX* (ice blue), *hxnW* (purple) and *hxnV* (yellow). A single vertical line symbolises physical separation of *hxn* genes on different contigs, while a double vertical line symbolises location of genes on different chromosomes (*A. nidulans*). Grey bar: classes of *Pezizomycotina* subphylum. Purple bar: class of *Saccharomycotina* subphylum.

S6 Fig. Phylogeny of the HxnR transcription factor.

All putative orthologues have the same protein domain organisation as the *A. nidulans* HxnR protein [11]. HxnR orthologues are > 30 % identical to the *A. nidulans* regulatory protein outside the N-terminal DNA-binding, zinc finger domain. The "HxnR-like" proteins are Cys2His2 proteins that appear exclusively in the early divergent class of the *Pezizomycetes,* and which also are > 30 % identical to *A. nidulans* HxnR (beyond the

zinc finger domain). Both *Pezizomycetes* genes have three centrally positioned introns,

the first two of which positions are conserved always flank an exon with 132-138 nt

length. In contrast to the orthologous *hxnR* gene, the "HxnR-like" gene is never

clustered with enzyme or transporter encoding *hxn* genes and occurs in all sequenced

Pezizomycetes species, i.e., including those that do not have *hxn* genes (except for *hxnM*

paralogues). Colour code: Purple: *Pezizomycetes*, including "HxnR-like proteins"

serving as an out group; Magenta: *Leotiomycetes*; Brown: *Sordariomycetes*; Orange:

Dothideomycetes; Green: *Aspergillus;* Olive Green: *Aspergillus Section Flavi;* Cyan:

Eurotiomycetes except *Aspergillus* and *Penicillium;* Darker Cyan: *Penicillium;* Red:

Saccharomycotina. Species names in red: those which map outside its cognate

phylogenetic clade, suggesting HGT events.

S7 Fig. Phylogeny of the HxnT putative FMN oxidoreductase.

Orthologous HxnT proteins are at least 40-45 % identical to the *A. nidulans* protein and

can be distinguish from other homologue sequences in the genome by the synteny of

hxn genes and the conservation of intron positions. The three-exon model of *hxnT* is

broadly conserved. In some taxa, like *Monascus* [11], *hxnT* is duplicated. In such cases,

we have labelled the protein from the cluster-associated *hxnT* gene, "1". The tree is

rooted with the *Saccharomycotina* clade, shown in blue, which also contains the HxnT-

like protein from the early divergent yeast, *Saitoella complicata* (*Taphrinomycotina*).

This species lacks the *hxnV* and *hxnX* genes, ubiquitous in all other *Ascomycota* with a

minimal *hxn* complement. Colour code: Brown: *Sordariomycetes*; Purple:

Leotiomycetes; Orange: *Dothideomycetes*. Other colours: Eurotiomycetes; Cyan: non-

Aspergillus, non-*Penicillium;* Darker Cyan: *Penicillium;* Magenta: *Aspergillus* sections

Nidulantes/Versicolores; Red: section *Flavi;* Green: section *Terrei. A. tanneri* (section

Circumdati) and *A. wentii* (Section *Cremei*) are indicated with black lines. Grey: section

Nigri. Species names marked in red indicate an anomalous phylogenetic position,

suggesting HGT events. *A. bertholletius* is in blue, to distinguish it from the other

members of section *Flavi*: *hxnT* is in this organism the only gene intact in an *hxn* cluster

where all other are pseudo genes.

1012

S8 Fig. Phylogeny of the HxnV putative monooxygenase.

Orthologous HxnV proteins are 40-45 % identical to the *A. nidulans* protein and can be

distinguish from other homologue sequences in the genome by the synteny of *hxn* genes

and the conservation of intron positions. Certain taxa of *Sordariomycetes* have

duplicated *hxnV* genes: in the smaller clade, the genes that encode the paralogues

labelled "1" are associated with other *hxn* genes. In the larger clade, the copy of the

hxnV gene for the paralogues labelled "2" are unlinked to resident *hxn* clusters. This

larger clade also includes several species, token species like *Neurospora crassa*,

Magnaporthe oryzae and *Trichoderma reesei*, that lack the *hxn* system and harbour only

a lone copy of the *hxnV* gene, suggesting a loss of the nicotinate assimilation pathway

and probably a novel function for *hxnV* in these *Sordariomycetes*. The tree is rooted in

the monophyletic HxnV clade of the *Saccharomycotina*. Colour code: Blue:

Saccharomycotina; Purple: *Pezizomycetes*; Magenta: *Leotiomycetes*; Brown:

Sordariomycetes; Black lines *Xylonomycetes*; Orange: *Dothideomycetes*; Cyan: non-

Aspergillus, non-*Penicillium Eurotiomycetes*; Darker Cyan: *Penicillium;* Green:

Aspergillus, except Olive green: section *Flavi;* Red: sections *Nidulantes/Versicolores*.

Species names in red, those mapping outside their cognate phylogenetic clade

suggesting HGT events. Species names in blue: those showing a duplication of the *hxnV*

gene.

1032

S9 Fig. Maximum Likelihood Phylogeny of the HxnM putative imino-hydrolase

protein.

All putative HxnM orthologues are >50% identical to the *A. nidulans* HxnM protein and

>50% identical to the biochemically characterised cyclic imide hydrolase from

Pseudomonas putida (strain YZ-26) (GenBank AAY98498). Methods used in the

construction of the tree are detailed in the Materials and Methods section. Numbers in

nodes are aLRTs (approximate Likelihood Ratio Tests). Names of species in black:

Putative HxnM proteins encoded by genes (*hxnM*) not clustered with any other *hxn*

gene. Names of species in blue: *hxnM* genes clustered only with *hxnN* genes (e.g., like

in *A. nidulans*, see Fig 1 or S5 Fig, or in *Lipomyces*). Names of species in red: *hxnM*

included in a large cluster, encoding ≥ 4 genes. Highlighted in yellow: a monophyletic

clade and putatively iso-functional clade comprising HxnMs from both clustered and

un-clustered *hxnM* genes. Highlighted in light green: a clade comprising only proteins

from un-clustered HxnM encoding genes, which appears as the outgroup to all clades

with HxnMs from clustered genes detailed above. Colour code: Grey: prokaryotic

outgroup (comprising both Bacterial and Archeal proteins); Blue: *Basidiomycota*

(phylum); Red: *Saccharomycotina* (subphylum); Other colours: Pezizomycotina taxa;

namely: Green: Eurotiomycetes; Olive green: *Aspergillus* section *Flavi* to highlight its

anomalous position (HGT); Brown: *Sordariomycetes*; Magenta: *Dothideomycetes*;

Orange: *Lecanoromycetes*; Cyan: *Leotiomycetes*; Purple: *Pezizomycetes*.

1053

S10 Fig. Phylogeny of the PfdA, B, and C paralogues in the *Eurotiales*.

The *pfdA* gene is ubiquitous in *Pezizomycotina* and encodes a well conserved protein of

~ 500 amino acids with a canonical peroxisome targeting sequence PTS-1 at its C-

terminus. Generally, *Eurotiales* PfdAs are >72 % identical to the *A. flavus* protein. The

pfdB and *pfdC* paralogues are restricted to specific taxa of the *Aspergillaceae* or

Trichocomaceae families of the *Eurotiales* order; some species have two- while others

have all three genes. The *pfdB* gene is regularly but not always, associated with the *hxn*

gene cluster. The PfdB and C proteins are respectively, >60 % and >50 % identical to *A.*

flavus PfdA. Typically, both paralogues are shorter than ubiquitous PfdA; multiple

sequence alignments of the three paralogues (from one species) show a distinct gap in

the middle of the alignment. All PfdAs and PfdBs feature a canonical PTS-1 [56] but

some PdfCs appear to have lost the canonical signal sequence for peroxisome entry. All

Eurotiales Pfd paralogue proteins included show the same domain organisation and

their encoding genes have a conserved exon/intron structure with five exons (see Sup.

Figure S12). In *Aspergillus*, the four introns in *pfdA* are confirmed by non-overlapping

EST clones accessions DR703303 (introns 1 and 2 near the ATG) and CO136618

(introns 3 and 4 near the STOP). The exon 2 is always 59 nt long and exon 4 is always

74 nt long in nearly all Pfd paralogues. The size of the large central exon 3 distinguishes

pfdA from its *B* and *C* paralogues. Tree construction was as detailed in Material and

Methods, except that a Blosum 30 matrix was used for the BMGE alignment trimming.

The species involved in the *hxn* cluster transfer (*Penicillium*, *Talaromyces*, *Aspergillus*

section *Flavi*) are in red. Species names in magenta are *Aspergillu*s of sections other

than *Flavi* which also include a *pfdB* gene in their *hxn* gene clusters. Colour code:

Green: Outgroups *Pezizomycetes* and *Lecanoromycetes* (all PfdA; these taxa do not

feature the *pfdBC* paralogues); Cyan: *Eurotiales* other than *Aspergillus* and *Penicillium*;

Darker Cyan: *Penicillium;* Other colours: sections of the genus *Aspergillus.* Magenta:

sections *Nidulantes*/*Versicolores*; Red: section *Flavi*; Orange: sections

Fumigati/*Clavati*; Brown: sections *Terrei*/*Candidi*; Yellow: section *Aspergillus*; Grey:

section *Nigri*; Purple: section *Circumdati*.

1083

S11 Fig. Intron positions in *pfdA*, *pfdB*, and *pfdC* paralogues in *Aspergillus nomius*.

According to the data of *A. nomius* whole genome sequence project (JNOM00000000,

https://www.ncbi.nlm.nih.gov/nuccore/JNOM00000000), all three *pfd* paralogues have

four introns in conserved positions, two near the 5'end of the CDS and two near the 3'.

Introns (lower case letters) are highlighted in blue (splice site consensus sequences: 5'-

donor, lariat branch point sequence, and 3'-acceptor) and grey (other intronic

sequences). The sizes of the small exons 2 and 4 are absolutely conserved throughout

the orthologues. The size of the large central exon 3 distinguishes the three paralogues.

In the ubiquitous *pfdA* gene exon 3 is 30 codons longer than the exon 3 of *pfdB* and 20

codons longer than exon 3 of *pfdC*. The canonical PTS-1 [56] has been lost in some but

not all PfdC paralogues that supports a scenario of gene duplication of *pfdA* with

simultaneous or subsequent cluster integration (mean similarity between *A* and *B*

paralogues 65% compared with 88% of *A* orthologues among themselves). The third

paralogue *pfdC* (mean identity between *A* and *C* paralogues 57%) present in section

Flavi, and in a number of *Talaromyces* and *Penicilium* species and in a few species of

other clades (supplementary fig. S11), which is consistent with a basal duplication

followed by several episodes of loss completely unrelated to the evolution of the *hxn*

cluster.

1102

S12 Fig. Phylogeny of HxnS in the *Pezizomycotina*.

The tree is shown in a circular, cartoon form. Tree was constructed as indicated in

Materials and Methods, except that MAFFT E-INS-i was used for the alignment. The

HxnS sequences included are from Ámon et al. [11] to which orthologue proteins from

some more recently published *Aspergillus*, *Penicillium* and *Talaromyces* species were

added: *A. sclerotiales*, *A. tanneri*, *P. steckii*, *T. wortmannii*, and *T. picheae*. Colour

code: Blue: Outgroups, including non-fungal xanthine hydroxylases and selected HxA

orthologues; Purple: *Pezizomycetes*; Magenta: *Leotiomycetes*; Brown: *Sordariomycetes*;

Black (lines): *Xylonomycetes*; Orange: *Dothideomycetes*; Cyan: non-*Aspergillus*, non-

Penicillium Eurotiales (such as *Talaromyces*); Darker Cyan: *Penicillium* genus*;* Red:

Sections *Nidulantes/Versicolores* (also called subgenus *Nidulantes* by Houbraken and

Samson, 2011); Grey: Section *Nigri;* Green: all other *Aspergilli*. The three *Penicillium*

species and *Aspergillus sclerotialis*, which cluster together within the *Sordariomycetes*

are shown in red. The duplicated HxnS sequences of *A. tanneri* are shown in blue. The

three *Talaromyces* species which conserve HxnS are shown in magenta.

1118