1

Origin and diversification of fibroblasts from the sclerotome in zebrafish

Roger C. Ma[1], Katrinka M. Kocha[1], Emilio E. Méndez-Olivos[1], Tyler D. Ruel[1], and Peng Huang[1]*

[1] Department of Biochemistry and Molecular Biology, Cumming School of Medicine, Alberta Children's

Hospital Research Institute, University of Calgary, 3330 Hospital Drive, Calgary, Alberta, Canada T2N

4N1

* Correspondence should be addressed to PH.

Email: huangp@ucalgary.ca

Tel: 403-220-4612

Running title: Origin and diversification of fibroblasts

Keywords: Sclerotome, Fibroblasts, Collagen, Axial skeleton, Fin mesenchymal cells, Zebrafish

ABSTRACT

Fibroblasts play an important role in maintaining tissue integrity by secreting components of the

extracellular matrix and initiating response to injury. Although the function of fibroblasts has been

extensively studied in adults, the embryonic origin and diversification of different fibroblast subtypes

during development remain largely unexplored. Using zebrafish as a model, we show that the

sclerotome, a sub-compartment of the somite, is the embryonic source of multiple fibroblast subtypes

including tenocytes (tendon fibroblasts), blood vessel associated fibroblasts, fin mesenchymal cells,

and interstitial fibroblasts. High-resolution imaging shows that different fibroblast subtypes occupy

unique anatomical locations with distinct morphologies. Long-term Cre-mediated lineage tracing

reveals that the sclerotome also contributes to cells closely associated with the axial skeleton.

Ablation of sclerotome progenitors results in extensive skeletal defects. Using photoconversion-based

cell lineage analysis, we find that sclerotome progenitors at different dorsal-ventral and anterior-

posterior positions display distinct differentiation potentials. Single-cell clonal analysis combined with

in vivo imaging suggests that the sclerotome mostly contains unipotent and bipotent progenitors prior

to cell migration, and the fate of their daughter cells is biased by their migration paths and relative

positions. Together, our work demonstrates that the sclerotome is the embryonic source of trunk

fibroblasts as well as the axial skeleton, and local signals likely contribute to the diversification of

distinct fibroblast subtypes.

INTRODUCTION

Fibroblasts are connective tissue cells present in most organs of animals. They function as tissue support cells by synthesizing and remodeling the extracellular matrix (ECM). Recent studies have shown that tissue-resident fibroblasts also play important regulatory roles in wound healing, inflammation, tumor microenvironment, and tissue fibrosis (Kalluri, 2016; Kendall and Feghali-Bostwick, 2014; Plikus et al., 2021). Although fibroblasts have historically been viewed as a homogenous cell population, recent work using single-cell RNA sequencing (scRNA-seq) has revealed a high level of heterogeneity in fibroblast populations from different tissues as well as within the same tissue (Buechler et al., 2021; Forte et al., 2022; Muhl et al., 2020; Tabib et al., 2018; Xie et al., 2018). This raises the question of how the different fibroblast subtypes are generated during embryonic development.

One mechanism to diversify fibroblast populations is through contributions from multiple embryonic sources. Previous studies suggest that fibroblasts in the same tissue may have heterogeneous embryonic origins. For example, in avian and mouse embryos, dermal fibroblasts in the head are of neural crest origin, whereas those in the dorsal and ventral trunk originate from the somite and lateral plate mesoderm, respectively (Thulabandu et al., 2018). Similarly, during heart development in chick and mouse embryos, most cardiac fibroblasts originate from lateral plate mesoderm derived epicardium, while a small population arises from an endothelial/endocardial lineage (Tallquist, 2020).

Somites are the embryonic source of many tissue support cells, including fibroblasts (Christ et al., 2007). During development, somite forms three separate compartments: the dermatome, myotome, and sclerotome. The sclerotome contributes to the axial skeleton of animals (Christ et al., 2004). Work in mouse and chick has shown that the sclerotome is also the developmental origin of tendon fibroblasts (tenocytes) (Brent et al., 2003; Schweitzer et al., 2001) and vascular smooth muscle cells (Pouget et al., 2008; Wiegreffe et al., 2007). In zebrafish, the sclerotome has a unique bipartite organization and contributes to tenocytes, perivascular fibroblasts, and pericytes in a stereotypical manner (Ma et al., 2018; Rajan et al., 2020). It raises the question of the cellular mechanisms governing the diversification of fibroblast subtypes from sclerotome progenitors.

In fish and amphibians, fin mesenchymal cells are a population of fibroblasts present in the larval fin fold. They express many ECM proteins to provide structural support for the developing fin fold (Durán et al., 2011; Feitosa et al., 2012). However, the developmental origin of fin mesenchymal cells remains controversial. Dye labeling experiments in *Xenopus* and zebrafish suggest that the neural crest contributes to the fin mesenchyme (Smith and Hall, 1990; Smith et al., 1994). However, later

experiments in *Xenopus* suggest that the mesoderm also contributes to fin mesenchymal cells in both

the dorsal and ventral fins (Garriock and Krieg, 2007; Tucker and Slack, 2004). Similarly, cell

transplantation experiments in axolotls reveal dual origin of the fin mesenchyme from both the neural

crest and somites (Sobkow et al., 2006). More recent work using genetic lineage tracing in fish

(zebrafish) and amphibians (*Xenopus* and axolotl) supports a model where fin mesenchymal cells

originate exclusively from the mesoderm with no contribution from the neural crest lineage (Lee et al.,

2013; Taniguchi et al., 2015). In particular, the dermomyotome compartment of the somite has been

suggested as the source of fin mesenchymal cells in zebrafish (Lee et al., 2013).

Here, we show that the sclerotome in zebrafish is a major source of multiple fibroblast populations

as well as cells associated with the axial skeleton. Different sclerotome-derived fibroblast subtypes

display unique morphologies and occupy distinct anatomical locations. In vivo time-lapse imaging and

single-cell clonal analysis reveal that sclerotome progenitors can generate multiple fibroblast

subtypes, and their differentiation potential is biased by their positions as well as their migratory

patterns. Together, our results suggest that distinct fibroblast subtypes are specified from sclerotome

progenitors by local cellular environments.

RESULTS

Characterization of a sclerotome reporter in zebrafish

To explore the lineage potential of the sclerotome in zebrafish, we recently developed a

sclerotome reporter, *nkx3.1:Gal4; UAS:NTR-mCherry* (*nkx3.1^{NTR-mCherry}*) (Ma et al., 2018). The

nkx3.1:Gal4 driver accurately recapitulated the expression of the endogenous *nkx3.1* gene (Fig S1).

The specificity of this *nkx3.1^{NTR-mCherry}* line was further validated by co-labeling with known sclerotome

markers, including *vcanb* (*versican b*) (Landolt et al., 1995), *foxc1b* (Topczewska et al., 2001), and

twist1b (previously known as *twist1a*) (Germanguz et al., 2007; Yeo et al., 2009) (Fig 1A). Due to the

perdurance of the mCherry protein, we used *nkx3.1^{NTR-mCherry}* to label the initial sclerotome domains

and all their descendants. We crossed *nkx3.1^{NTR-mCherry}* with the endothelial reporter *kdrl:EGFP* to

examine sclerotome-derived cells at 2 dpf (days post-fertilization). Based on the anatomical location

and morphology, we broadly defined five groups of mCherry⁺ cells from the *nkx3.1* lineage (Fig 1B-C):

1) cells closely associated with blood vessels in the trunk, including the dorsal longitudinal

anastomotic vessel (DLAV), intersegmental vessels (ISV), dorsal aorta (DA), and caudal vein plexus

(CVP); 2) tenocytes along somite boundaries as previously described (Ma et al., 2018; Subramanian

et al., 2018); 3) interstitial cells located in the space between the notochord, spinal cord, and muscles;

4) cells located in both dorsal and ventral fin folds; and 5) some muscle cells located at the dorsal and

ventral edges of each somite (Ma et al., 2018). The contribution of muscles from the *nkx3.1* lineage

suggests that *nkx3.1* labels some early bipotent somitic cells that generate both sclerotome

progenitors and muscles, which is consistent with previous single-cell lineage analysis of the ventral

somite (Morin-Kensicki and Eisen, 1997). In addition to the five groups of cells from the *nkx3.1*

lineage, the *nkx3.1:Gal4* line drove ectopic expression in some cuboidal cells at the edge of the fin

fold (Fig 1C and S2), which were not derived from the sclerotome. It is worth noting that although the

nkx3.1:Gal4 line nicely matched the endogenous *nkx3.1* expression, the amplification of the

GAL4/UAS system in *nkx3.1^{Kaede}* (*nkx3.1:Gal4; UAS:Kaede*) and *nkx3.1^{NTR-mCherry}* lines resulted in

various degrees of ectopic expression outside the *nkx3.1* domains (Fig S1). We have previously

shown that both tenocytes and ISV-associated perivascular fibroblasts express *col1a2* and *col5a1*

(Ma et al., 2018; Rajan et al., 2020), two pan-fibroblast markers defined by scRNA-seq experiments

(Metikala et al., 2021; Muhl et al., 2020). Indeed, all *nkx3.1^{NTR-mCherry}*-positive non-muscle cells (groups

1-4) were labeled by *col1a2:GFP* at 2 dpf (Fig 1D), confirming their fibroblast identity. Our results

suggest that the sclerotome is the embryonic source of multiple types of fibroblasts in zebrafish.

The sclerotome generates different types of trunk fibroblasts

We performed mosaic labeling and marker analysis to characterize fibroblasts derived from the

nkx3.1 lineage (Fig 2A-C). *nkx3.1*$^{NTR-mCherry}$-expressing fibroblasts in the trunk, including tenocytes,

blood vessel associated fibroblasts, and interstitial fibroblasts, were marked by pan-fibroblast markers

such as *tgfbi* and *col5a1* (Metikala et al., 2021; Muhl et al., 2020) (Fig 2B). A subset of these

sclerotome-derived fibroblasts was associated with different vascular beds (Fig 1C and 2A). We

previously described ISV-associated perivascular fibroblasts, characterized by a globular cell body

with short processes extending around the ISV (Rajan et al., 2020). By contrast, labeling with a

mosaic *col1a2:Gal4; UAS:Kaede* line (*col1a2*^{Kaede}) (Sharma et al., 2019) revealed that fibroblasts

associated with DLAV or DA (DLAV and DA fibroblasts, respectively) were more elongated along the

long axis of the blood vessel (Fig 1C and 2A). CVP is a transient venous network with a stereotypical

"honeycomb-like" structure. Fibroblasts marked by *nkx3.1*$^{NTR-mCherry}$ occupied the space in the

honeycomb pockets of the CVP (Fig 1C and 2A). Mosaic labeling showed that individual Kaede$^+$ cells

had a small and elongated cell body with long cellular processes along the underlying endothelium

and sometimes across the honeycomb pocket (Fig 2A). The location and morphology of these CVP-

associated fibroblasts suggest that they are stromal reticular cells (SRCs) (Murayama et al., 2015).

Tenocytes and interstitial fibroblasts were the two fibroblast subtypes that populated the trunk

region but were not closely associated with blood vessels (Fig 1C). We and others have previously

shown that tenocytes are located along the myotendinous junction (MTJ), extending long cellular

processes into the intersomitic space (Fig 2A) (Ma et al., 2018; Subramanian et al., 2018). Marker

analysis revealed that *nkx3.1*$^{NTR-mCherry}$-positive cells along the MTJ were specifically labeled by known

tendon markers such as *scxa*, *cilp*, and *tnmd* (Chen and Galloway, 2014; Ralliere et al., 2015) (Fig

2C). By contrast, interstitial fibroblasts were scattered throughout the interstitial space and displayed

varied morphologies (Fig 1C and 2A).

The sclerotome contributes to fin mesenchymal cells

The *nkx3.1*$^{NTR-mCherry}$ line labeled cells in the fin fold (Fig 1B-C). At 2 dpf, the fin fold is subdivided

into two parts: the major lobe wrapping around the trunk and tail, and the minor lobe beneath the yolk

extension (Parichy et al., 2009) (Fig S2). Cells from the *nkx3.1* lineage populated both the dorsal and

ventral regions in the major lobe (Fig 1C and S2). Using the mosaic *col1a2*^{Kaede} line, we found that

Kaede$^+$ fin cells displayed extensive "tree-like" cellular processes projecting towards the periphery of

the fin fold (Fig 2A). The location and morphology suggest that these sclerotome-derived fin fold

residing cells correspond to fin mesenchymal cells (Carney et al., 2010; Feitosa et al., 2012; Lee et

al., 2013). Previous studies have shown that fin mesenchymal cells are marked by *hemicentin2*

(*hmcn2*) and *fibulin1* (*fbln1*), while the neighboring cell layer of apical epidermal cells is labelled by

hmcn2 and *fras1* (Carney et al., 2010; Feitosa et al., 2012; Lee et al., 2013) (Fig 3A). Double staining

revealed two juxtaposing cell layers in the fin fold: $hmcn2^+fbln1^+$ fin mesenchymal cells in the proximal

layer, and $hmcn2^+fras1^+$ apical epidermal cells in the distal layer (Fig 3B). As expected, double

staining in $nkx3.1^{NTR-mCherry}$ embryos showed that mCherry$^+$ cells in the fin fold co-expressed *fbln1* but

not *fras1* (Fig 3C), confirming that the sclerotome contributes to fin mesenchymal cells, a specific cell

population in the fin fold.

Interestingly, $nkx3.1^{Kaede}$-expressing fin mesenchymal cells were absent in the caudal fin fold and

minor lobe (Fig S2). Cre-mediated lineage tracing in *nkx3.1:Gal4; UAS:Cre-ERT2; ubi:Switch* fish

showed similar results (Fig 3D). Switched mCherry$^+$ cells derived from the *nkx3.1* lineage gave rise to

fin mesenchymal cells in both the dorsal and ventral regions of the major lobe at 4 dpf, but cells in the

minor lobe and caudal fin remained GFP$^+$mCherry$^-$ (un-switched) (Fig 3D). Our results suggest that

sclerotome is not the only lineage that gives rise to fin mesenchymal cells. To determine whether the

dermomyotome (also known as the external cell layer) contributes to fin mesenchymal cells, we

imaged *pax7b:Gal4FF; UAS:EGFP* ($pax7b^{EGFP}$) fish at 50 hpf. Interestingly, despite extensive labeling

of dermomyotome cells in all somites, most fin mesenchymal cells were not marked by $pax7b^{EGFP}$ (Fig

3E). EGFP$^+$ fin cells were not observed in the minor lobe or caudal fin, while on average, only 4.6%

and 0.2% of fin mesenchymal cells were $pax7b^{EGFP}$-positive in the dorsal and ventral regions of the

major lobe, respectively (Fig 3F). This is in striking contrast to 95% (dorsal) and 94% (ventral)

contributions from the $nkx3.1^{NTR-mCherry}$ reporter (Fig 3F). Therefore, our results suggest that the

sclerotome, but not the dermomyotome, is the main contributor of fin mesenchymal cells in the dorsal

and ventral fin folds.

Together, our data show that the sclerotome generates multiple types of fibroblasts in the

zebrafish trunk, supporting different tissues such as muscles, blood vessels, and the fin fold. These

fibroblast subtypes reside in distinct anatomical locations with unique morphologies, some of which

can be defined by specific marker expression.

The sclerotome contributes to the axial skeleton in zebrafish

Since the sclerotome is known to contribute to the axial skeleton (Christ et al., 2004), we

performed Cre-mediated lineage tracing on cells from the *nkx3.1* lineage in zebrafish. Sclerotome-

derived cells in *nkx3.1:Gal4; UAS:Cre-ERT2; ubi:Switch* fish were switched to express mCherry with

4-hydroxytamoxifen (4OHT) treatment at 1-2 dpf. Co-labeling of calcein, a vital dye for calcified

structures, in switched fish at 17 dpf revealed that mCherry$^+$ cells were closely associated with the

vertebrae, including the centrum, neural arch, and hemal arch (Fig 4A). Interestingly, mCherry$^+$ cells

formed a thin layer external to the calcein-positive centrum, with enrichment in the region between

adjacent centra (Fig 4A). By contrast, calcein staining in neural and hemal arches appeared to be

embedded within mCherry$^+$ cells with overlapping staining (Fig 4A). Long-term tracing of *nkx3.1$^+$* cells

resulted in substantial labeling of tenocytes at the MTJ and cells associated with the vertebral column

in fish at 6 weeks post-fertilization (wpf) and adult stages (Fig 4B). To determine the function of

sclerotome-derived cells in skeletal development, we performed genetic ablation using the

nitroreductase (NTR) system (Curado et al., 2008). *nkx3.1$^{NTR-mCherry}$* embryos or their mCherry$^-$ siblings

were treated with metronidazole (MTZ) at 1-2 dpf and stained for bone and cartilage at 3 wpf (Fig 4C-

E). Since many ablated fish developed heart edema, likely caused by the depletion of perivascular

fibroblasts (Rajan et al., 2020), we can only analyze survivors from the ablated group, which likely

represents fish with more mosaic expression. MTZ-treated mCherry$^+$ fish were significantly smaller

than mCherry$^-$ controls (Fig 4D), suggesting impaired growth with partial ablation of the *nkx3.1*

lineage. Interestingly, while vertebral bodies formed normally, both neural and hemal arches displayed

variable defects, with an average of 35% of vertebrae missing neural arches and 3% missing hemal

arches (Fig 4C-E). Together, our results suggest that sclerotome-derived cells in zebrafish contribute

to the normal development of the vertebral column, especially skeletal elements attached to the

vertebral body.

Both dorsal and ventral sclerotome domains give rise to multiple types of fibroblasts

We have previously shown that the zebrafish sclerotome consists of a dorsal and ventral domain,

both of which generate tenocytes and perivascular fibroblasts (Ma et al., 2018; Rajan et al., 2020). We

asked how each sclerotome domain contributes to different fibroblast subtypes. Taking advantage of

the photoconvertible Kaede protein, we performed lineage tracing of sclerotome domains in

nkx3.1^{Kaede} fish (Fig 5A). Either the dorsal or ventral sclerotome domain of a single somite was

photoconverted from Kaede^{green} to Kaede^{red} at 24 hpf (hours post-fertilization). We then determined

the fate of Kaede^{red} cells after 24 hours based on their location and morphology.

When the dorsal sclerotome was labeled at 24 hpf, we observed small projections from these

Kaede^{red} cells, suggesting the initiation of cell migration at this stage (Fig 5B). At 48 hpf, all dorsal

sclerotome-derived Kaede^{red} cells were restricted to the upper half of the trunk dorsal to the notochord

(Fig 5B). Based on the morphology and location relative to the vasculature, we determined that the

dorsal sclerotome generated dorsal fin mesenchymal cells, DLAV fibroblasts, tenocytes, ISV-

associated perivascular fibroblasts, and interstitial fibroblasts.

Next, we performed photoconversion experiments to mark the ventral sclerotome with Kaede^{red} at

24 hpf. Similar to the dorsal domain, small cellular projections were visible from Kaede^{red} cells,

suggesting that both dorsal and ventral sclerotome progenitors initiate their migration at the same

stage (Fig 5C). Interestingly, in anteriorly converted somites (somites 16-18), cellular projections

pointed exclusively in the dorsal direction, while in posteriorly converted somites (somites 23-25), both

dorsal and ventral projections were observed (Fig 5C). This result suggests that the migration pattern

is different in ventral sclerotome progenitors along the anterior-posterior axis, which likely accounts for

the cells generated in the CVP and ventral fin fold region. All the cells generated by the ventral

sclerotome were restricted to approximately the ventral two-thirds of the trunk. In anterior somites, the

ventral sclerotome generated cells in the trunk, including tenocytes, perivascular fibroblasts, and

interstitial fibroblasts at 48 hpf (Fig 5C). In contrast, the ventral sclerotome of posterior somites gave

rise to cells in the trunk and cells ventral to the photoconverted region in the periphery, including CVP-

associated SRCs and fin mesenchymal cells in the ventral fin fold (Fig 5C).

Together, our results suggest that both the dorsal and ventral sclerotome domains can generate

multiple fibroblast subtypes, but the progenitors in each domain generate an anatomically restricted

population of cells according to their dorsal-ventral and anterior-posterior positions.

Fate of dorsal sclerotome progenitors is biased by the migration direction

Both dorsal and ventral sclerotome contribute to multiple types of fibroblasts. It is possible that

sclerotome progenitors are multipotent, or alternatively, individual progenitors are "fate-restricted" to

differentiate into a specific fibroblast subtype. To distinguish these possibilities, we performed single-

cell clonal analysis of sclerotome progenitors as previously described (Sharma et al., 2019). We

focused on the smaller dorsal domain to ensure precise single-cell photoconversion and accurate cell

tracing. We selected $nkx3.1^{Kaede}$ fish with more mosaic Kaede expression to minimize the

photoconversion of multiple cells. At 24 hpf, a single Kaede[green] cell in the dorsal domain between

somites 16 and 18 was photoconverted to Kaede[red] (Fig 6A). We then imaged the same fish after 24

hours to determine the identity of their Kaede[red] descendants. Most clones (35/42, 83%) contained at

least 2 daughter cells, suggesting at least one cell division between 24 and 48 hpf (Fig 6B). From 42

dorsal sclerotome progenitors, 38% (16/42) produced clones with at least two fibroblast subtypes, and

2% (1/42) generated three subtypes (Fig 6C). This result suggests that some dorsal sclerotome

progenitors are bipotent and possibly multipotent at 24 hpf. Consistent with domain tracing, we found

that dorsal sclerotome progenitors gave rise to all four types of fibroblasts, including dorsal fin

mesenchymal cells, tenocytes, blood vessel associated fibroblasts, and interstitial fibroblasts (Fig 6D).

As observed earlier, 33 of the 42 photoconverted Kaede[red] cells at 24 hpf showed visibly biased

projections pointing either dorsally (14 cells) or ventrally (19 cells), likely corresponding to their

migration direction. To correlate progenitor morphology with their lineage potential, we categorized

different fibroblasts into two types based on their final positions (Fig 6A). In the type I group,

fibroblasts are located dorsal to the somite, including dorsal fin mesenchymal cells and DLAV

fibroblasts. Type II cells, such as tenocytes, ISV-associated perivascular fibroblasts, and interstitial

fibroblasts, are located medial to the somite. Interestingly, progenitors with initial dorsal projections

predominantly generated type I cells (13/14, 93%), whereas cells projecting ventrally were more likely

to give rise to type II cells (11/19, 58%) than type I cells (3/19, 16%) and sometimes generated cells of

both types (5/19, 26%) (Fig 6E-F). It is important to note that ventral projecting progenitors at 24 hpf

never generated dorsal fin mesenchymal cells and all type I cells generated by this group were DLAV

fibroblasts found along the ventral side of the DLAV (Fig 6F). Together, our results suggest that the

dorsal sclerotome domain contains a mixture of unipotent and bipotent progenitors at 24 hpf, and their

migration direction likely restricts the types of fibroblasts they differentiate into.

Fate of ventral sclerotome progenitors is biased by the position along the migratory path

In posterior somites, some ventral sclerotome progenitors migrate ventrally towards the CVP

region to generate SRCs and ventral fin mesenchymal cells. To visualize this dynamic process, we

performed time-lapse imaging of *kdrl:EGFP; nkx3.1$^{NTR-mCherry}$* embryos from 25 to 50 hpf (Video S1

and Fig S3). At 25 hpf, some migrating mCherry$^+$ progenitors were found ventral to the somites.

These cells migrated ventrally towards the fin fold as more mCherry$^+$ cells emerged from the ventral

somite and joined the migration. To better describe the position of migrating cells, we referred to cells

at the front edge of the migration as "leading cells", while all cells behind were called "lagging cells".

We observed a stereotypical migration pattern whereby mCherry$^+$ cells "marched" ventrally in an

orderly fashion, and individual cells did not pass other cells that were ahead. This orderly migration

was largely maintained, even during cell division. As a result, descendants of leading cells remained

at the front of the migration throughout the movie and formed ventral fin mesenchymal cells, whereas

descendants of lagging cells mostly generated SRCs associated with the CVP (Video S1 and Fig S3).

Interestingly, the cardinal vein underwent similar ventral expansion to form the CVP during the same

time window. Lagging cells appeared to constantly interact with the forming CVP, while leading cells

remained ahead of the vascular sprouts (Video S1 and Fig S3). Together, our results suggest that the

fate of ventral sclerotome progenitors is biased by their position along the migratory path, potentially

due to their interactions with the local environment.

To further compare the lineage potential between the leading and lagging cells, we performed

single-cell clonal analysis in *nkx3.1^{Kaede}* fish (Fig 7A). Single ventrally migrating Kaede^{green} progenitors

between somites 21-25 were photoconverted to Kaede^{red} at 24 hpf, and their descendants were

imaged after 24 hours. Similar to our time-lapse movies, the majority of photoconverted leading cells

generated only ventral fin mesenchymal cells (17/20, 85%), but a small number gave rise to both

SRCs and ventral fin mesenchymal cells (3/20, 15%) (Fig 7B-C). Interestingly, leading cells never
generated SRCs only. By contrast, 41% of lagging cells generated only SRCs (9/22), 36% generated
both ventral fin mesenchymal cells and SRCs (8/22), and 23% produced only ventral fin mesenchymal
cells (5/22) (Fig 7B-C). Combining time-lapse imaging and single-cell tracing, our results suggest that
the relative position along the migratory path biases the fate of these sclerotome-derived cells.

DISCUSSION

Our work establishes the sclerotome as the embryonic source of different connective tissue cells in zebrafish. First, the sclerotome gives rise to distinct fibroblasts in the trunk. Second, the sclerotome contributes to proper development of the axial skeleton. Third, the sclerotome contains a mixture of unipotent and bipotent progenitors and their differentiation is biased by their axial positions and migratory paths.

The sclerotome is the embryonic source of trunk fibroblasts

The developmental potential of the sclerotome in zebrafish has been first explored with vital dye injections (Morin-Kensicki and Eisen, 1997). Using a classic sclerotome marker *nkx3.1* (Schneider et al., 2000), we have previously developed the *nkx3.1:Gal4* line in zebrafish to genetically label sclerotome progenitors and their descendants (Ma et al., 2018). Here, we provide additional evidence that *nkx3.1:Gal4* labels cells of the sclerotome lineage. First, the *nkx3.1^{NTR-mCherry}* reporter co-expresses known sclerotome markers such as *versican b* (Landolt et al., 1995), *foxc1b* (Topczewska et al., 2001), and *twist1b* (Germanguz et al., 2007; Yeo et al., 2009). Second, lineage tracing of *nkx3.1*-derived cells shows extensive labeling of cells associated with the axial skeleton, consistent with previous cell tracing experiments (Morin-Kensicki et al., 2002). Interestingly, the *nkx3.1^{NTR-mCherry}* reporter also labels some muscles adjacent to the dorsal and ventral sclerotome domains. This observation is in agreement with previous single-cell lineage analysis of the presumptive sclerotome in the ventral somite where a sub-population of these cells gives rise to both sclerotome derivatives and muscles (Morin-Kensicki and Eisen, 1997). We speculate that some bipotent somitic cells might transiently express *nkx3.1*, before committing to either the myotome or the sclerotome lineage. This idea is supported by the observation that progenitors at developmental branch points sometimes express markers characteristic of multiple fates (Farrell et al., 2018).

Our previous studies (Ma et al., 2018; Rajan et al., 2020) and current work provide strong evidence that the sclerotome is the embryonic source of multiple types of fibroblasts in the trunk of zebrafish. Based on marker expression, anatomical location, cell morphology, and function, we define at least four fibroblast subtypes derived from the sclerotome: tenocytes, fin mesenchymal cells, blood vessel associated fibroblasts, and interstitial fibroblasts. Tenocytes express classic tendon markers *scxa* and *tnmd* (Chen and Galloway, 2014; Ma et al., 2018). Their cell bodies are positioned medially along the myotendinous junction, while extending long cellular processes into the intersomitic space (Ma et al., 2018; Subramanian et al., 2018). Tenocytes stabilize muscle attachment likely by secreting a number of ECM proteins (Ma et al., 2018; Subramanian and Schilling, 2014). Fin mesenchymal

cells, on the other hand, are marked by the expression of ECM genes *hmcn2* and *fbln1*. They are located adjacent to apical epidermal cells in the fin fold with distally extended tree-like processes. Recent work has shown that fin mesenchymal cells guide fin formation by aligning collagen fibers of actinotrichia (Kuroda et al., 2020). Genetic ablation of fin mesenchymal cells in the pectoral and median fin folds results in collapse of the respective fins (Lalonde and Akimenko, 2018). Although blood vessel associated fibroblasts are labeled by pan-fibroblast markers similar as other fibroblast subtypes in the trunk, they are characterized by their close association with the vasculature. Interestingly, their morphology differs depending on the vascular bed. For example, ISV-associated perivascular fibroblasts are globular, whereas DLAV fibroblasts are more elongated. We have previously shown that perivascular fibroblasts play dual roles in stabilizing nascent ISVs and functioning as pericyte precursors (Rajan et al., 2020). It is likely that other blood vessel associated fibroblasts play similar roles in stabilizing the corresponding blood vessels. Interestingly, stromal reticular cells closely interact with CVP, which functions as an early hematopoietic niche during development (Wattrus and Zon, 2018). Indeed, recent work has shown that defects in SRC maturation result in a compromised CVP niche that fails to support hematopoietic stem cell (HSC) maintenance and expansion (Murayama et al., 2015). Therefore, SRCs might not only provide structural support to the CVP, but also modulate the microenvironment required for HSC differentiation. Despite extensive labeling of blood vessel associated fibroblasts, our lineage analysis shows that sclerotome progenitors do not contribute to the vasculature. This result suggests that the sclerotome domain defined by *nkx3.1* is distinct from the endotome, a sub-compartment of the somite that has been shown to contain endothelial precursors (Nguyen et al., 2014; Sahai-Hernandez et al., 2020). Compared to other fibroblast subtypes, interstitial fibroblasts are the least well-characterized. Based on their distribution around the notochord and spinal cord, it is conceivable that interstitial fibroblasts give rise to osteoblasts required for vertebral column development. This is consistent with our observation that long-term cell tracing of the *nkx3.1* lineage results in extensive labeling of cells associated with the vertebral bodies, neural and hemal arches. Together, our results suggest that the sclerotome generates distinct fibroblast subtypes to support a variety of tissues, including muscles, fin folds and blood vessels.

The sclerotome contributes to axial skeleton development in zebrafish

In amniotes, the sclerotome is the embryonic source of the vertebrae, structural units of the vertebral column (Christ et al., 2004). In teleosts, each vertebra is composed of three elements: vertebral body (centrum), neural, and hemal arches. In contrast to amniotes, studies in zebrafish and salmon have suggested that the centrum is formed by notochord sheath cells instead of the

sclerotome (Fleming et al., 2004; Grotmol et al., 2003; Lleras Forero et al., 2018; Pogoda et al., 2018;

Wang et al., 2013; Wopat et al., 2018). However, how the sclerotome contributes to the development

of the axial skeleton in teleosts remains unclear. Our long-term lineage tracing experiments

demonstrate that cells derived from the sclerotome lineage are closely associated with all three

skeletal elements of the vertebrae. Partial depletion of sclerotome-derived cells results in variable

defects in neural and hemal arches, while centra are largely normal. This result is consistent with the

current model, in which the centrum forms independently of the somite (Lleras Forero et al., 2018;

Wopat et al., 2018). However, it is not clear why sclerotome ablation leads to a stronger defect in

neural arches than hemal arches, despite similar labeling of sclerotome-derived cells on both

structures. There are three non-mutually exclusive explanations. First, severe loss of hemal arches

might be detrimental to survival; therefore, only fish with mild phenotypes can survive to 3 wpf (time of

analysis). Second, the precursors of hemal arches might have a greater capacity to regenerate after

ablation than those of neural arches. Finally, there might be an alternative non-sclerotome source that

contributes to the formation of hemal arches redundantly. Interestingly, our results in zebrafish show

remarkable parallel to previous observations in medaka. In medaka, sclerotome-derived cells marked

by *twist:GFP* are similarly enriched in the intervertebral regions, and are thought to be the source of

osteoblasts to generate different skeletal elements (Inohaya et al., 2007). Morpholino knockdown of

key sclerotome genes, including *pax1*, *pax9*, and *twist*, results in the loss of neural arches with largely

normal centra (Mise et al., 2008; Yasutake et al., 2004). Together, these results suggest that the

sclerotome contributes to proper development of the axial skeleton in teleosts.

The developmental origin of fin mesenchymal cells

There has been considerable debate regarding where fin mesenchymal cells originate during fish

and amphibian development. Lineage tracing studies in zebrafish suggest that the paraxial

mesoderm, specifically the dermomyotome, but not the sclerotome, is the embryonic source of fin

mesenchymal cells (Lee et al., 2013). Their conclusions contrast with our findings in two aspects.

First, using the *pax3a:EGFP* line as a marker for the dermomyotome, Lee et al. (2013) show

extensive contribution of EGFP[+] fin mesenchymal cells throughout the entire major lobe of the fin fold,

including the caudal fin. By contrast, we find that a different dermomyotome reporter, *pax7b[EGFP]*

(Pipalia et al., 2016), contributes minimally to fin mesenchymal cells in the fin fold. This discrepancy is

likely due to the different expression dynamics of *pax3a* and *pax7b*, despite both being

dermomyotome markers. It has been shown that *pax3a* is initially broadly expressed in the entire

somite before becoming restricted to the anterior somites (precursors of the dermomyotome) (Feng et

al., 2006; Hammond et al., 2007). However, *pax7b* is expressed several hours after *pax3a* and is

restricted to the anterior somites (Feng et al., 2006; Hammond et al., 2007; Minchin and Hughes,

2008). Therefore, the *pax3a:EGFP* line may not be as specific as *pax7b^{EGFP}* in labeling the

dermomyotome. Second, Lee et al. (2013) use the *ola-twist1:Gal4* transgenic line, in which Gal4

expression is driven by the medaka (*Oryzias latipes*) *twist1* promoter, to rule out sclerotome

contribution. However, our results using the *nkx3.1$^{NTR-mCherry}$* reporter indicate that the sclerotome

contributes to most fin mesenchymal cells in both the dorsal and ventral regions of the major lobe.

Interestingly, we have previously shown that the expression of a similar *ola-twist1:EGFP* reporter is

restricted to sclerotome-derived cells around the notochord, but is absent in both dorsal and ventral

sclerotome domains (Ma et al., 2018). This observation explains why the *ola-twist1:Gal4* line fails to

label any fin mesenchymal cells (Lee et al., 2013). Together, our work suggests an updated model

where fin mesenchymal cells are generated by multiple lineages, depending on their locations. Fin

mesenchymal cells in the dorsal and ventral fin fold have a sclerotome origin, those in the caudal fin

originate from a non-dermomyotome and non-sclerotome compartment of the somite, while fin

mesenchymal cells in the minor lobe of the fin fold have an unidentified embryonic source.

Mechanisms of fibroblast diversification

A fundamental question in developmental biology is how progenitors in the embryo diversify into

distinct differentiated cell types. Using *nkx3.1*-based transgenic reporters, our work reveals three key

features of cell fate diversification from the sclerotome lineage (Fig 7D). First, sclerotome progenitors

display regional differences in their differentiation potential along the dorsal-ventral and anterior-

posterior positions. Dorsally located fibroblasts, such as dorsal fin mesenchymal cells and DLAV

fibroblasts, are derived from the dorsal sclerotome, whereas ventrally located fibroblasts, such as

stromal reticular cells and ventral fin mesenchymal cells, are generated by the ventral sclerotome.

Similarly, the ventral sclerotome in anterior somites (before the end of the yolk extension) generates

only fibroblasts in the trunk, whereas the ventral sclerotome in posterior somites contributes to stromal

reticular cells and fin mesenchymal cells in addition to trunk fibroblasts. Second, fate of sclerotome

progenitors is biased by their migratory path. For dorsal sclerotome progenitors, dorsal migration

limits the cell to become either dorsal fin mesenchymal cells or DLAV fibroblasts, while ventral

migration leads to the differentiation of predominately tenocytes or perivascular fibroblasts. Third, the

relative order of migrating sclerotome progenitors highly correlates with their differentiation potential.

Leading cells are more likely to become ventral fin mesenchymal cells, while lagging cells tend to

differentiate into stromal reticular cells. The combination of these three mechanisms (spatial

regulation, migration direction, and migration order) (Fig 7D) likely contributes to the generation of

different fibroblast subtypes in a stereotypical manner (Ma et al., 2018; Rajan et al., 2020).

How does a sclerotome progenitor decide which cell type to differentiate into? Photoconversion-

based clonal analysis reveals that single sclerotome progenitors predominantly generate 1-2 different

fibroblast subtypes. This result suggests that the sclerotome contains a mixture of unipotent and

bipotent progenitors prior to migration. The presence of bipotent progenitors suggests that individual

sclerotome cells are plastic and are not fate-restricted to become a specific fibroblast subtype. We

speculate that the local microenvironment plays a key role in regulating the migration and subsequent

differentiation of sclerotome-derived cells. In the trunk region, the migration of sclerotome-derived

cells depends on Sonic Hedgehog (Shh) from the notochord and floor plate (Ma et al., 2018).

Inhibition of Shh signaling results in the loss of tenocytes between muscle boundaries (Ma et al.,

2018). Similarly, Slit-Robo signaling has been shown to mediate the polarization and migration of fin

mesenchymal cells in the peripheral fin fold (Mahabaleshwar et al., 2022). Cell type diversification of

the sclerotome is remarkably similar to that of the neural crest lineage. Previous studies have shown

that migrating neural crest cells are multipotent (Baggiolini et al., 2015), and the ultimate cell fate is

determined by their axial positions as well as the migration timing and direction (Rocha et al., 2020).

In summary, we demonstrate that the sclerotome is the embryonic source of multiple fibroblast

subtypes and contributes to axial skeletal development in zebrafish. Our work shows that the

sclerotome is an excellent model for studying cell type diversification during embryonic development.

The combination of positional and migratory information as well as the local microenvironment likely

contributes to the differentiation trajectory of sclerotome progenitors. Future work using scRNA-seq

may further reveal the heterogeneity of sclerotome-derived cells and identify key gene regulatory

networks underlying the lineage progression of sclerotome progenitors.

MATERIALS AND METHODS

Ethics statement

All animal experiments were conducted in accordance with the principles outlined in the current Guidelines of the Canadian Council on Animal Care. All protocols were approved by the Animal Care Committee of the University of Calgary (#AC17-0128 and #AC21-0102).

Zebrafish strains

Zebrafish strains used in this study were maintained and raised under standard conditions. The following transgenic strains were used in this study: *Gt(:gSAIz-GAL4FFD)nkgsaizGFFD164A* (*pax7b:Gal4FF*) (Pipalia et al., 2016), *TgBAC(col1a2:Gal4)ca102* (Ma et al., 2018; Sharma et al., 2019), *TgBAC(col1a2:GFP)ca103* (Ma et al., 2018), *Tg(kdrl:EGFP)la163* (Choi et al., 2007), *Tg(kdrl:mCherry)ci5* (Proulx et al., 2010), *TgBAC(nkx3.1:Gal4)ca101* (Ma et al., 2018), *Tg(UAS:EGFP)nkuasgfp1a* (Asakawa et al., 2008), *Tg(UAS:Cre-ERT2)ca105* (Sharma et al., 2019), *Tg(-3.5ubb:LOXP-EGFP-LOXP-mCherry)cz1701* (*ubi:Switch*) (Mosimann et al., 2011), *Tg(UAS:Kaede)s1999t* (Davison et al., 2007), and *Tg(UAS:NTR-mCherry)c264* (Davison et al., 2007). The mosaic *col1a2:Gal4; UAS:Kaede* line was maintained by selecting embryos with more mosaic Kaede expression.

In situ hybridization and immunohistochemistry

Whole-mount in situ hybridization and immunohistochemistry were performed according to previously established protocols. We used the following RNA probes in this study: *cilp*, *col5a1*, *fbln1*, *foxc1b*, *fras1*, *gal4*, *hmcn2*, *kaede*, *ntr-mCherry*, *scxa*, *tgfbi*, *tnmd*, *twist1b*, *vcanb* (*versican b*). Double fluorescent in situ hybridizations were performed using different combinations of digoxigenin (DIG) or dinitrophenyl (DNP) labeled probes. For immunohistochemistry, rabbit polyclonal antibody to RFP (1:1000, MBL, PM005) and chicken antibody to GFP (1:250, Aves Labs, GFP-1020) were used. For fluorescent detection of antibody labeling, the appropriate Alexa Fluor-568 or Alexa Fluor-647 conjugated secondary antibodies (1:500, Thermo Fisher) were used.

Kaede photoconversion

Photoconversion was carried out using a 405 nm laser and a 20x objective on an Olympus FV1200 confocal microscope. At the appropriate stage, Kaede-expressing embryos were anesthetized using tricaine and mounted in 0.8% low-melting-point agarose. The duration and intensity of laser required for complete green-to-red conversion depended on the Kaede expression

level and the size of the target region. For cell tracing of the sclerotome domains, a circular region of

20x20 pixels was converted using 1.5% 405 nm laser for 4 seconds. For single-cell clonal analysis of

sclerotome progenitors, typically a circular region of 10x10 pixels was converted using 1.5% 405 nm

laser for 4 seconds. After photoconversion, embryos were confirmed for the expression of Kaede[red]

and recovered in E3 fish water until the necessary stages.

Cre-mediated lineage tracing

nkx3.1:Gal4; UAS:Cre-ERT2; ubi:Switch embryos were treated with 10 μM 4-hydroxytamoxifen

(4OHT) from 1 to 2 dpf. After treatment, embryos were washed three times and recovered in fish

water for analysis at the appropriate stages.

Cell ablation

NTR-mCherry[+] embryos and NTR-mCherry[-] sibling controls at the appropriate stage were treated

with metronidazole (MTZ, Sigma Millipore) at a concentration of 5 mM in fish water. Embryos were

incubated in MTZ solution for at least 24 hours to achieve complete ablation. Fish were then washed

and recovered in fish water until the desired stage for analysis.

Bone and cartilage staining

Bone and cartilage staining was performed following a modified version of a previously described

protocol (Walker and Kimmel, 2007). Briefly, larvae at 3 wpf were euthanized in 0.4% tricaine and

fixed in 4% formaldehyde in PBS overnight at 4 °C. Fixed larvae were washed in a solution containing

100 mM Tris pH 7.5 and 10 mM $MgCl_2$, then stained overnight in 0.04% alcian blue solution. The

larvae were then washed with decreasing concentrations of ethanol diluted in 100 mM Tris pH 7.5

(80%, 50%, 25%) and bleached in a 3% H_2O_2/0.5% KOH solution until all pigments were lost. They

were washed twice with 25% glycerol/0.1% KOH and stained in the same solution with 0.01% alizarin

red for 1 hour. Finally, larvae were washed twice in 50% glycerol/0.1% KOH and switched to 100%

glycerol for imaging.

Calcein staining

Bones can be fluorescently labelled in live larvae using the vital dye, calcein. Larvae at 17 dpf

were placed in a 100 mm petri dish in a 0.2% calcein solution for 15 – 30 minutes at room

temperature. They were then washed at least six times with water from zebrafish vivarium. The larvae

were left for 2 – 3 hours prior to imaging to allow excess calcein to pass through their digestive tract.

Fish were anesthetized using tricaine and mounted in 0.8% low-melting-point agarose for live

imaging. To obtain transverse views, fish were fixed in 4% formaldehyde and sectioned for imaging.

Vibratome sectioning

Juvenile and adult zebrafish were fixed in 4% paraformaldehyde (PFA) with 1% dimethyl sulfoxide

(DMSO) overnight at 4 °C. Fixed fish were washed in PBS and decalcified in 0.5 M EDTA at room

temperature for 3 days. Following decalcification, the dissected trunks were embedded in 20% gelatin

and sectioned using a Leica VT1000S vibratome to obtain slices at 150 µm thick. The sections were

mounted on slides in 50% glycerol for confocal imaging. Confocal tile scans were stitched together

using Olympus Fluoview software to generate a full view of the transverse section.

Time-lapse imaging and cell tracking

Time-lapse imaging of zebrafish embryos was carried out using an Olympus FV1200 confocal

microscope. At the appropriate stage, fish were anesthetized using tricaine and mounted in 0.8% low-

melting-point agarose. To keep the agarose hydrated and minimize the development of pigments, as

well as minimize movement of the fish, a small volume of E3 water with phenylthiourea and tricaine

was carefully flooded around the agarose. Z-stack images of the region of interest were collected at

regular intervals (6-9 mins) for up to 25 hours. Images were then compiled into movies using Olympus

Fluoview software. Cells from each movie were manually tracked using the Fiji software (Schindelin et

al., 2012).

Statistical analysis

All graphs were generated using the GraphPad Prism software. Data were plotted as mean±SEM.

The significance between two samples was calculated using Mann-Whitney *U* test. p values: $p > 0.05$

(not significant); $p < 0.05$ (*); $p < 0.01$ (**); $p < 0.001$ (***); $p < 0.0001$ (****).

ACKNOWLEDGMENTS

We thank the zebrafish community for providing probes and reagents; Simon Hughes for sharing

the *pax7b^{EGFP}* line; Sarah Childs for sharing reagents, transgenic lines and providing critical input on

this project; members of the Childs and Huang laboratories for discussions; Arsheen Rajan for critical

comments on the manuscript.

COMPETING INTERESTS

The authors declare that no competing interests exist.

FUNDING

This study was supported by grants to P.H. from the Canadian Institute of Health Research (MOP-

136926 and PJT-169113), Canada Foundation for Innovation John R. Evans Leaders Fund (Project

32920), and Startup Fund from the Alberta Children's Hospital Research Institute (ACHRI).

    FIGURE LEGENDS

Fig 1. The sclerotome gives rise to different types of fibroblasts. (A) Double fluorescent in situ

staining of *nkx3.1^NTR-mCherry* embryos at 24 hpf showed co-expression of the sclerotome markers

versican b, *foxc1b*, and *twist1b* (cyan) with *ntr-mcherry* (red) in the dorsal sclerotome (short arrows),

ventral sclerotome (long arrows), and sclerotome-derived notochord-associated cells (arrowheads).

n=15 per staining. (B) Schematic of sclerotome-derived fibroblasts: blood vessel associated

fibroblasts (cyan), fin mesenchymal cells (black), tenocytes (magenta), and interstitial fibroblasts

(yellow). DLAV, dorsal longitudinal anastomotic vessel; ISV, intersegmental vessel; DA, dorsal aorta;

CVP, caudal vein plexus; n, notochord. (C) At 2 dpf, *nkx3.1^NTR-mCherry*-positive cells (red) were

observed throughout the trunk showing close association with the vasculature labeled by *kdrl:EGFP*

(green) (top panels). Zoomed-in views (bottom panels) show different mCherry⁺ fibroblast subtypes

(arrows) in four boxed regions of the trunk (i-iv). mCherry expression in muscles (asterisks) and skin

cells along the fin fold (arrowheads) is shown. *n*=13. (D) Most *nkx3.1^NTR-mCherry*-positive sclerotome-

derived cells (red) co-expressed *col1a2:GFP* (green) at 2 dpf. The upper panel shows the dorsal

region with dorsal fin mesenchymal cells (white arrows), DLAV fibroblasts (long cyan arrows), ISV-

associated perivascular fibroblasts (cyan arrows), tenocytes (magenta arrows), and interstitial

fibroblasts (yellow arrows). The middle panel shows DA fibroblasts (cyan arrows) associated with the

dorsal aorta. The bottom panel shows stromal reticular cells (cyan arrows) and ventral fin

mesenchymal cells (white arrows). *n*=12. Scale bars: 50 μm.

Fig 2. Characterization of sclerotome-derived fibroblasts. (A) The mosaic *col1a2^Kaede* line (green)

was crossed with *kdrl:mCherry* (red) to visualize the morphology of different fibroblasts at a high

resolution at 2 dpf: blood vessel associated fibroblasts (cyan arrows), fin mesenchymal cells (white

arrows), tenocytes (magenta arrow), and interstitial fibroblasts (yellow arrow). Cell processes are

indicated by arrowheads. The associated vascular bed and fin fold are indicated for blood vessel

associated fibroblasts and fin mesenchymal cells, respectively. *n*=20. (B) *nkx3.1^NTR-mCherry*; *kdrl:EGFP*

embryos at 53 hpf were co-stained with RFP antibody (red), GFP antibody (yellow), and probes for

fibroblast markers *tgfbi* or *col5a1* (cyan). mCherry⁺ blood vessel associated fibroblasts (cyan arrows),

tenocytes (magenta arrows), and interstitial fibroblasts (yellow arrows) co-expressed *tgfbi* and *col5a1*.

n=12 per staining. (C) *nkx3.1^NTR-mCherry* embryos at 53 hpf were co-stained with RFP antibody (red)

and probes for tenocyte markers *scxa, cilp,* or *tnmd* (cyan). mCherry⁺ tenocytes (magenta arrows)

along MTJ co-expressed tenocyte markers. *n*=12 per staining. Scale bars: (A) 20 μm; (B, C) 50 μm.

Fig 3. The sclerotome contributes to fin mesenchymal cells. (A) Schematic of the zebrafish fin

fold. Both dorsal and ventral fin folds are populated by two adjacent cell layers: the distal

hmcn2⁺fras1⁺ apical epidermal cells and proximal *hmcn2⁺fbln1⁺* fin mesenchymal cells. (B) Double

fluorescent in situ staining of *fbln1* (red) with either *fras1* or *hmcn2* (green) at 48 hpf. *fras1* labeled

apical epidermal cells (orange brackets), *fbln1* marked fin mesenchymal cells (white brackets), while

hmcn2 was expressed in both layers. n=15 per staining. (C) *nkx3.1^{NTR-mCherry}* embryos were costained

with *fbln1* or *fras1* probes (green) and anti-RFP antibody (red) at 52 hpf. Apical epidermal cells and fin

mesenchymal cells are indicated by arrowheads and arrows, respectively. n=10 per staining. (D)

nkx3.1^{Cre-ERT2}; ubi:Switch embryos treated with 4OHT at 1-2 dpf and imaged at 4 dpf. Switched

mCherry⁺ fin mesenchymal cells were present in the dorsal and ventral regions of the major lobe

(arrows) but absent in the minor lobe or caudal fin (arrowheads). n=10. (E) *pax7b^{EGFP}* embryos at 50

hpf showed extensive EGFP expression in the dermomyotome (arrow), while most fin mesenchymal

cells (arrowheads) in both the major and minor lobes of the fin fold were EGFP⁻. n=10. (F)

Quantification of lineage contribution of fin mesenchymal cells. The percentage of fin mesenchymal

cells labeled by each lineage reporter at 2 dpf was plotted as mean±SEM. n=22 (*pax7b^{EGFP}*) and 13

(*nkx3.1^{NTR-mCherry}*). Statistics: Mann-Whitney U test; p<0.0001 (****). Scale bars: (B, D, E) 50 μm; (C)

20 μm.

Fig 4. The sclerotome contributes to the axial skeleton. (A) *nkx3.1^{Cre-ERT2}; ubi:Switch* embryos

were treated with 4OHT at 1-2 dpf to switch on mCherry expression (red) in sclerotome-derived cells.

Switched embryos were stained with calcein (cyan) at 17 dpf to visualize calcified structures, including

the centra (white arrowheads), neural arches (yellow arrowheads), and hemal arches (magenta

arrowheads). Magnified views of the neural (i) and hemal (ii) arches as well as a cross-sectional view

of the centrum (dotted line, iii) are shown. Switched mCherry⁺ cells associated with different skeletal

structures are denoted by arrows with the corresponding colors. Enrichment of mCherry⁺ cells in the

regions between the vertebral bodies is indicated by asterisks. Since the calcein signal is very strong,

un-switched GFP expression is undetectable under the imaging setting. n=5. (B) *nkx3.1^{Cre-ERT2};*

ubi:Switch embryos treated with 4OHT at 1-2 dpf were raised and imaged at 6 wpf (transverse

section) and 1-year post-fertilization (ypf) (sagittal section). mCherry⁺ cells were associated with the

MTJ (magenta arrowheads), centrum (white arrowheads), neural arch (long arrows), and hemal arch

(short arrows). The dorsal aorta (asterisks) showed strong autofluorescence in both channels.

Standard lengths of imaged fish are indicated. n=2 per stage. (C) *nkx3.1^{NTR-mCherry}* embryos and their

mCherry⁻ siblings were treated with MTZ at 1-2 dpf and stained with alizarin red and alcian blue at 3

wpf. mCherry⁺ fish showed normal vertebral bodies (arrowheads) with missing neural arches (long

arrow) and some misconnected hemal arches (short arrows). n=19 (mCherry⁻) and 15 (mCherry⁺). (D,

E) Quantification of standard length and skeletal defects in MTZ-treated fish (C). The percentage of

vertebrae with missing neural or hemal arches was plotted as mean±SEM. Statistics: Mann-Whitney

U test. Asterisk representation: p<0.01 (**), p<0.0001 (****). Scale bars: (A) 50 µm; (B, C) 200 µm.

Fig 5. Lineage analysis of sclerotome domains. (A) Schematic of lineage analysis. The entire

dorsal or ventral sclerotome domain in *nkx3.1^Kaede* embryos was photoconverted at 24 hpf and

analyzed for cell composition at 48 hpf. (B, C) Examples of cell tracing of the dorsal (B) and ventral

(C) sclerotome domains in *nkx3.1^Kaede* embryos. At 48 hpf, full view of the trunk and close-up views of

the boxed regions are shown. Cell projections at 24 hpf (arrowheads) and the outline of blood vessels

(dotted lines) are denoted. Pigment cells (asterisks) showed strong autofluorescence in the red

channel. (B) Dorsal sclerotome domain generated dorsal fin mesenchymal cells (white arrows), DLAV

fibroblasts (long cyan arrows), perivascular fibroblasts (cyan arrows), tenocytes (magenta arrows),

and interstitial fibroblasts (yellow arrows). n=20. (C) In the anterior somites above the yolk extension

(upper panel), the ventral sclerotome domain generated perivascular fibroblasts (cyan arrows),

tenocytes (magenta arrows), and interstitial fibroblasts (yellow arrows). In the posterior somites above

the CVP region (bottom panel), the ventral sclerotome domain generated SRCs (cyan arrowheads),

ventral fin mesenchymal cells (white arrows), tenocytes (magenta arrows), and interstitial fibroblasts

(yellow arrows). The ventral boundary of the CVP is indicated by dotted lines. n=20. Scale bars: 50

µm.

Fig 6. Clonal analysis of the dorsal sclerotome. (A) Schematic of clonal analysis. A single dorsal

sclerotome progenitor in *nkx3.1^Kaede* embryos was photoconverted at 24 hpf and imaged at 48 hpf to

determine clone composition. Dorsal sclerotome-derived cells were subdivided into two groups based

on their dorsal-ventral positions: the more dorsal type I cells include dorsal fin mesenchymal cells and

DLAV fibroblasts (above the dotted line), while the more ventral type II cells include tenocytes,

perivascular fibroblasts, and interstitial fibroblasts (below the dotted line). (B) Quantification of the

clone size. (C) Quantification of fibroblast subtypes in each clone. (D) Distribution of different

fibroblast subtypes from all decedents of traced cells. F, fin mesenchymal cells; T, tenocytes; V, blood

vessel associated fibroblasts; I, interstitial fibroblasts. (E) Quantification of the clone composition. Only

photoconverted cells showing polarized cell projections at 24 hpf were graphed. n=14 (with dorsal

projection) and 19 (with ventral projection) cells. (F) Examples of clonal analysis of dorsal sclerotome

progenitors in *nkx3.1^Kaede* embryos at 24 and 48 hpf. In the top panel (type I cell lineage), the

photoconverted cell showing dorsal projections (arrowhead) generated two dorsal fin mesenchymal

cells (white arrows) and one DLAV fibroblast (long cyan arrows). In the bottom panel (type II cell

lineage), the photoconverted cell with ventral cell projections (arrowhead) generated one DLAV

fibroblast (long cyan arrows), one perivascular fibroblast (cyan arrows), and one tenocyte (magenta

arrows). The outline of the vasculature is denoted by dotted lines. *n*=42 photoconverted cells from 40

embryos. Scale bars: 50 µm.

Fig 7. Clonal analysis of the ventral sclerotome. (A) Schematic of clonal analysis. A single ventral

sclerotome progenitor in *nkx3.1^Kaede* embryos was photoconverted at 24 hpf and imaged after 24 hours

to determine clone composition. Cells were classified as leading or lagging cells based on their initial

position at 24 hpf, where cells at the front of the migration path were defined as leading cells. (B)

Quantification of the clone composition. (C) Examples of cell tracing at 24 and 48 hpf. A lagging cell

(top panel) generated four SRCs (arrowheads) dorsal to the ventral edge of the CVP (dotted lines). A

leading cell (bottom panel) generated two fin mesenchymal cells (arrows). *n*=42 photoconverted cells

from 38 embryos. (D) Three mechanisms contribute to the diversification of fibroblast subtypes from

sclerotome progenitors in zebrafish. The fate of sclerotome progenitors is biased by their dorsal-

ventral or anterior-posterior positions (a, spatial regulation), migration direction (b), and migration

order (c). n, notochord. Scale bars: 50 µm.

 SUPPLEMENTAL FIGURES

Fig S1. Characterization of the sclerotome reporter. Double fluorescent in situ staining of *nkx3.1*

(red) together with *gal4*, *kaede*, or *ntr-mcherry* (cyan) was performed in *nkx3.1^NTR-mCherry^*, *nkx3.1^Kaede^*,

or *UAS:NTR-mCherry* embryos at 20 hpf. *gal4* expression in *nkx3.1^NTR-mCherry^* embryos accurately

recapitulated endogenous *nkx3.1* expression in both dorsal and ventral sclerotome domains (short

and long arrows, respectively). *kaede* expression in *nkx3.1^Kaede^* embryos was largely consistent with

nkx3.1 expression, whereas *nkx3.1^NTR-mCherry^* embryos showed ectopic *ntr-mcherry* expression in

somites (arrowheads). Note that *UAS:NTR-mCherry* but not *UAS:Kaede* embryos also showed low

Gal4-independent expression in the tail bud (asterisks). Representative images are shown with *n*

numbers for each staining. Both the merged and individual channels are shown. Scale bar: 50 μm.

Fig S2. The sclerotome generates fin mesenchymal cells in the dorsal and ventral fin folds. A

tiled confocal image of *nkx3.1^Kaede^; kdrl:mCherry* embryo at 2 dpf showing Kaede⁺ sclerotome-derived

cells (cyan) throughout the trunk of the embryo. Kaede⁺ fin mesenchymal cells were present in both

dorsal and ventral fin folds (arrows), but absent from either the caudal fin or the minor lobe (asterisks).

Arrowheads indicate mCherry expression in skin cells at the edge of the fin fold. The major and minor

lobes of the fin fold are outlined by dashed lines. *n*=5. Scale bar: 200 μm.

Fig S3. Time-lapse imaging of migration of ventral sclerotome progenitors. Snapshots of a

*kdrl:EGFP; nkx3.1^NTR-mCherry^* embryo between 25 and 50 hpf are shown with timestamps (hh:mm). The

migration and division of two representative lagging cells (cyan arrows and arrowheads) and two

leading cells (white arrows and arrowheads) were traced throughout the movie. Daughter cells are

indicated by arrows or arrowheads in the same color. The lagging cell never migrated ahead of the

leading cell and generated 2 SRCs. The leading cell remained at the migration front ahead of the

lagging cell and generated 2 ventral fin mesenchymal cells. *n*=7. Scale bar: 50 μm.

Video S1. Time-lapse imaging of migration of ventral sclerotome progenitors. *kdrl:EGFP;*

*nkx3.1^NTR-mCherry^* embryos were imaged from 25 to 50 hpf with timestamps (hh:mm). The migration and

division of representative lagging (cyan arrows) and leading (white arrows) cells were traced

throughout the movie. Daughter cells are indicated by arrows in the same color. The lagging cell

never migrated ahead of the leading cell and generated 2 SRCs. The leading cell remained at the

migration front ahead of the lagging cell and generated 2 ventral fin mesenchymal cells. Snapshots of

this video are shown in Fig S3. *n*=7. Scale bar: 50 μm.

REFERENCES

Asakawa, K., Suster, M. L., Mizusawa, K., Nagayoshi, S., Kotani, T., Urasaki, A., Kishimoto, Y., Hibi, M. and Kawakami, K. (2008). Genetic dissection of neural circuits by Tol2 transposon-mediated Gal4 gene and enhancer trapping in zebrafish. *Proc. Natl. Acad. Sci. U. S. A.* **105**,.

Baggiolini, A., Varum, S., Mateos, J. M., Bettosini, D., John, N., Bonalli, M., Ziegler, U., Dimou, L., Clevers, H., Furrer, R., et al. (2015). Premigratory and migratory neural crest cells are multipotent in vivo. *Cell Stem Cell* **16**, 314–322.

Brent, A. E., Schweitzer, R. and Tabin, C. J. (2003). A somitic compartment of tendon progenitors. *Cell* **113**, 235–248.

Buechler, M. B., Pradhan, R. N., Krishnamurty, A. T., Cox, C., Calviello, A. K., Wang, A. W., Yang, Y. A., Tam, L., Caothien, R., Roose-Girma, M., et al. (2021). Cross-tissue organization of the fibroblast lineage. *Nature* **593**,.

Carney, T. J., Feitosa, N. M., Sonntag, C., Slanchev, K., Kluger, J., Kiyozumi, D., Gebauer, J. M., Talbot, J. C., Kimmel, C. B., Sekiguchi, K., et al. (2010). Genetic analysis of fin development in zebrafish identifies furin and Hemicentin1 as potential novel fraser syndrome disease genes. *PLoS Genet.* **6**,.

Chen, J. W. and Galloway, J. L. (2014). The development of zebrafish tendon and ligament progenitors. *Development* **141**, 2035–2045.

Choi, J., Dong, L., Ahn, J., Dao, D., Hammerschmidt, M. and Chen, J. N. (2007). FoxH1 negatively modulates flk1 gene expression and vascular formation in zebrafish. *Dev. Biol.* **304**, 735–744.

Christ, B., Huang, R. and Scaal, M. (2004). Formation and differentiation of the avian sclerotome. *Anat. Embryol. (Berl).* **208**,.

Christ, B., Huang, R. and Scaal, M. (2007). Amniote somite derivatives. *Dev. Dyn.* **236**,.

Curado, S., Stainier, D. Y. and Anderson, R. M. (2008). Nitroreductase-mediated cell/tissue ablation in zebrafish: a spatially and temporally controlled ablation method with applications in developmental and regeneration studies. *Nat Protoc* **3**, 948–954.

Davison, J. M., Akitake, C. M., Goll, M. G., Rhee, J. M., Gosse, N., Baier, H., Halpern, M. E., Leach, S. D. and Parsons, M. J. (2007). Transactivation from Gal4-VP16 transgenic insertions for tissue-specific cell labeling and ablation in zebrafish. *Dev Biol* **304**, 811–824.

Durán, I., Marí-Beffa, M., Santamaría, J. A., Becerra, J. and Santos-Ruiz, L. (2011). Actinotrichia collagens and their role in fin formation. *Dev. Biol.* **354**, 160–172.

Farrell, J. A., Wang, Y., Riesenfeld, S. J., Shekhar, K., Regev, A. and Schier, A. F. (2018). Single-cell reconstruction of developmental trajectories during zebrafish embryogenesis. *Science (80-.).*

360, eaar3131.

Feitosa, N. M., Zhang, J., Carney, T. J., Metzger, M., Korzh, V., Bloch, W. and Hammerschmidt, M. (2012). Hemicentin 2 and Fibulin 1 are required for epidermal-dermal junction formation and fin mesenchymal cell migration during zebrafish development. *Dev. Biol.* **369**, 235–248.

Feng, X., Adiarte, E. G. and Devoto, S. H. (2006). Hedgehog acts directly on the zebrafish dermomyotome to promote myogenic differentiation. *Dev Biol* **300**, 736–746.

Fleming, A., Keynes, R. and Tannahill, D. (2004). A central role for the notochord in vertebral patterning. *Development* **131**, 873–880.

Forte, E., Ramialison, M., Nim, H. T., Mara, M., Li, J. Y., Cohn, R., Daigle, S. L., Boyd, S., Stanley, E. G., Elefanty, A. G., et al. (2022). Adult mouse fibroblasts retain organ-specific transcriptomic identity. *Elife* **11**,.

Garriock, R. J. and Krieg, P. A. (2007). Wnt11-R signaling regulates a calcium sensitive EMT event essential for dorsal fin development of Xenopus. *Dev. Biol.* **304**, 127–140.

Germanguz, I., Lev, D., Waisman, T., Kim, C.-H. and Gitelman, I. (2007). Four *twist* genes in zebrafish, four expression patterns. *Dev. Dyn.* **236**, 2615–2626.

Grotmol, S., Kryvi, H., Nordvik, K. and Totland, G. K. (2003). Notochord segmentation may lay down the pathway for the development of the vertebral bodies in the Atlantic salmon. *Anat. Embryol. (Berl).* **207**,.

Hammond, C. L., Hinits, Y., Osborn, D. P., Minchin, J. E., Tettamanti, G. and Hughes, S. M. (2007). Signals and myogenic regulatory factors restrict pax3 and pax7 expression to dermomyotome-like tissue in zebrafish. *Dev Biol* **302**, 504–521.

Inohaya, K., Takano, Y. and Kudo, A. (2007). The teleost intervertebral region acts as a growth center of the centrum: In vivo visualization of osteoblasts and their progenitors in transgenic fish. *Dev. Dyn.* **236**, 3031–3046.

Kalluri, R. (2016). The biology and function of fibroblasts in cancer. *Nat. Rev. Cancer* **16**, 582–598.

Kendall, R. T. and Feghali-Bostwick, C. A. (2014). Fibroblasts in fibrosis: Novel roles and mediators. *Front. Pharmacol.* **5 MAY**, 123.

Kuroda, J., Itabashi, T., Iwane, A. H., Aramaki, T. and Kondo, S. (2020). The Physical Role of Mesenchymal Cells Driven by the Actin Cytoskeleton Is Essential for the Orientation of Collagen Fibrils in Zebrafish Fins. *Front. Cell Dev. Biol.* **8**, 580520.

Lalonde, R. L. and Akimenko, M. A. (2018). Effects of fin fold mesenchyme ablation on fin development in zebrafish. *PLoS One* **13**, e0192500.

Landolt, R. M., Vaughan, L., Winterhalter, K. H. and Zimmermann, D. R. (1995). Versican is selectively expressed in embryonic tissues that act as barriers to neural crest cell migration and

axon outgrowth. *Development* **121**, 2303–2312.

Lee, R. T. H., Knapik, E. W., Thiery, J. P. and Carney, T. J. (2013). An exclusively mesodermal

origin of fin mesenchyme demonstrates that zebrafish trunk neural crest does not generate

ectomesenchyme. *Dev.* **140**, 2923–2932.

Lleras Forero, L., Narayanan, R., Huitema, L. F. A., VanBergen, M., Apschner, A., Peterson-

Maduro, J., Logister, I., Valentin, G., Morelli, L. G., Oates, A. C., et al. (2018). Segmentation

of the zebrafish axial skeleton relies on notochord sheath cells and not on the segmentation

clock. *Elife* **7**,.

Ma, R. C., Jacobs, C. T., Sharma, P., Kocha, K. M. and Huang, P. (2018). Stereotypic generation of

axial tenocytes from bipartite sclerotome domains in zebrafish. *PLoS Genet.* **14**, e1007775.

Mahabaleshwar, H., Asharani, P. V, Loo, T. Y., Koh, S. Y., Pitman, M. R., Kwok, S., Ma, J., Hu,

B., Lin, F., Li Lok, X., et al. (2022). Slit-Robo signalling establishes a Sphingosine-1-phosphate

gradient to polarise fin mesenchyme. *EMBO Rep.* **23**, e54464.

Metikala, S., Chetty, S. C. and Sumanas, S. (2021). Single-cell transcriptome analysis of the

zebrafish embryonic trunk. *PLoS One* **16**,.

Minchin, J. E. N. and Hughes, S. M. (2008). Sequential actions of Pax3 and Pax7 drive xanthophore

development in zebrafish neural crest. *Dev. Biol.* **317**,.

Mise, T., Iijima, M., Inohaya, K., Kudo, A. and Wada, H. (2008). Function of Pax1 and Pax9 in the

sclerotome of medaka fish. *Genesis* **46**,.

Morin-Kensicki, E. M. and Eisen, J. S. (1997). Sclerotome development and peripheral nervous

system segmentation in embryonic zebrafish. *Development* **124**, 159–167.

Morin-Kensicki, E. M., Melancon, E. and Eisen, J. S. (2002). Segmental relationship between

somites and vertebral column in zebrafish. *Development* **129**, 3851–3860.

Mosimann, C., Kaufman, C. K., Li, P., Pugach, E. K., Tamplin, O. J. and Zon, L. I. (2011).

Ubiquitous transgene expression and Cre-based recombination driven by the ubiquitin promoter

in zebrafish. *Development* **138**, 169–177.

Muhl, L., Genové, G., Leptidis, S., Liu, J., He, L., Mocci, G., Sun, Y., Gustafsson, S.,

Buyandelger, B., Chivukula, I. V, et al. (2020). Single-cell analysis uncovers fibroblast

heterogeneity and criteria for fibroblast and mural cell identification and discrimination. *Nat.*

Commun. **11**, 3953.

Murayama, E., Sarris, M., Redd, M., Le Guyader, D., Vivier, C., Horsley, W., Trede, N. and

Herbomel, P. (2015). NACA deficiency reveals the crucial role of somite-derived stromal cells in

haematopoietic niche formation. *Nat. Commun.* **6**, 8375.

Nguyen, P. D., Hollway, G. E., Sonntag, C., Miles, L. B., Hall, T. E., Berger, S., Fernandez, K. J.,

Gurevich, D. B., Cole, N. J., Alaei, S., et al. (2014). Haematopoietic stem cell induction by

somite-derived endothelial cells controlled by meox1. *Nature* **512**,.

Parichy, D. M., Elizondo, M. R., Mills, M. G., Gordon, T. N. and Engeszer, R. E. (2009). Normal

table of postembryonic zebrafish development: Staging by externally visible anatomy of the living

fish. *Dev. Dyn.* **238**, 2975–3015.

Pipalia, T. G., Koth, J., Roy, S. D., Hammond, C. L., Kawakami, K. and Hughes, S. M. (2016).

Cellular dynamics of regeneration reveals role of two distinct Pax7 stem cell populations in larval

zebrafish muscle repair. *Dis Model Mech* **9**, 671–684.

Plikus, M. V., Wang, X., Sinha, S., Forte, E., Thompson, S. M., Herzog, E. L., Driskell, R. R.,

Rosenthal, N., Biernaskie, J. and Horsley, V. (2021). Fibroblasts: Origins, definitions, and

functions in health and disease. *Cell* **184**,.

Pogoda, H. M., Riedl-Quinkertz, I., Löhr, H., Waxman, J. S., Dale, R. M., Topczewski, J., Schulte-

Merker, S. and Hammerschmidt, M. (2018). Direct activation of chordoblasts by retinoic acid is

required for segmented centra mineralization during zebrafish spine development. *Dev.* **145**,.

Pouget, C., Pottin, K. and Jaffredo, T. (2008). Sclerotomal origin of vascular smooth muscle cells

and pericytes in the embryo. *Dev. Biol.* **315**, 437–47.

Proulx, K., Lu, A. and Sumanas, S. (2010). Cranial vasculature in zebrafish forms by angioblast

cluster-derived angiogenesis. *Dev. Biol.* **348**,.

Rajan, A. M., Ma, R. C., Kocha, K. M., Zhang, D. J. and Huang, P. (2020). Dual function of

perivascular fibroblasts in vascular stabilization in zebrafish. *PLoS Genet.* **16**, e1008800.

Ralliere, C., Fretaud, M., Thermes, V. and Rescan, P. Y. (2015). CILP1 is dynamically expressed in

the developing musculoskeletal system of the trout. *Int J Dev Biol* **59**, 505–509.

Rocha, M., Beiriger, A., Kushkowski, E. E., Miyashita, T., Singh, N., Venkataraman, V. and

Prince, V. E. (2020). From head to tail: regionalization of the neural crest. *Development* **147**,.

Sahai-Hernandez, P., Pouget, C., Svoboda, O. and Traver, D. (2020). Dermomyotome-derived

endothelial cells migrate to the dorsal aorta to support hematopoietic stem cell emergence.

bioRxiv 2020.05.14.096305.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,

S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for

biological-image analysis. *Nat. Methods* **9**, 676–82.

Schneider, A., Brand, T., Zweigerdt, R. and Arnold, H. (2000). Targeted disruption of the Nkx3.1

gene in mice results in morphogenetic defects of minor salivary glands: parallels to glandular

duct morphogenesis in prostate. *Mech. Dev.* **95**, 163–174.

Schweitzer, R., Chyung, J. H., Murtaugh, L. C., Brent, A. E., Rosen, V., Olson, E. N., Lassar, A.

and Tabin, C. J. (2001). Analysis of the tendon cell fate using Scleraxis, a specific marker for

tendons and ligaments. *Development* **128**, 3855–3866.

Sharma, P., Ruel, T. D., Kocha, K. M., Liao, S. and Huang, P. (2019). Single cell dynamics of

embryonic muscle progenitor cells in zebrafish. *Development* **146**, dev178400.

Smith, M. M. and Hall, B. K. (1990). Development and evolutionary origins of vertebrate skeletogenic

and odontogenic tissues. *Biol. Rev. Camb. Philos. Soc.* **65**, 277–373.

Smith, M., Hickman, A., Amanze, D., Lumsden, A. and Thorogood, P. (1994). Trunk neural crest

origin of caudal fin mesenchyme in the zebrafish Brachydanio rerio. *Proc. R. Soc. B Biol. Sci.*

256, 137–145.

Sobkow, L., Epperlein, H. H., Herklotz, S., Straube, W. L. and Tanaka, E. M. (2006). A germline

GFP transgenic axolotl and its use to track cell fate: Dual origin of the fin mesenchyme during

development and the fate of blood cells during regeneration. *Dev. Biol.* **290**, 386–397.

Subramanian, A. and Schilling, T. F. (2014). Thrombospondin-4 controls matrix assembly during

development and repair of myotendinous junctions. *Elife* **3**,.

Subramanian, A., Kanzaki, L. F., Galloway, J. L. and Schilling, T. F. (2018). Mechanical force

regulates tendon extracellular matrix organization and tenocyte morphogenesis through TGFbeta

signaling. *Elife* **7**,.

Tabib, T., Morse, C., Wang, T., Chen, W. and Lafyatis, R. (2018). SFRP2/DPP4 and FMO1/LSP1

Define Major Fibroblast Populations in Human Skin. *J. Invest. Dermatol.* **138**, 802–810.

Tallquist, M. D. (2020). Developmental pathways of cardiac fibroblasts. *Cold Spring Harb. Perspect.*

Biol. **12**,.

Taniguchi, Y., Kurth, T., Medeiros, D. M., Tazaki, A., Ramm, R. and Epperlein, H. H. (2015).

Mesodermal origin of median fin mesenchyme and tail muscle in amphibian larvae. *Sci. Rep.* **5**,

11428.

Thulabandu, V., Chen, D. and Atit, R. P. (2018). Dermal fibroblast in cutaneous development and

healing. *Wiley Interdiscip. Rev. Dev. Biol.* **7**,.

Topczewska, J. M., Topczewski, J., Solnica-Krezel, L. and Hogan, B. L. M. (2001). Sequence and

expression of zebrafish foxc1a and foxc1b, encoding conserved forkhead/winged helix

transcription factors. *Mech. Dev.* **100**,.

Tucker, A. S. and Slack, J. M. W. (2004). Independent induction and formation of the dorsal and

ventral fins in Xenopus laevis. *Dev. Dyn.* **230**, 461–467.

Walker, M. B. and Kimmel, C. B. (2007). A two-color acid-free cartilage and bone stain for zebrafish

larvae. *Biotech. Histochem.* **82**,.

Wang, S., Kryvi, H., Grotmol, S., Wargelius, A., Krossøy, C., Epple, M., Neues, F., Furmanek, T.

and Totland, G. K. (2013). Mineralization of the vertebral bodies in Atlantic salmon (Salmo salar
L.) is initiated segmentally in the form of hydroxyapatite crystal accretions in the notochord
sheath. *J. Anat.* **223**,.

Wattrus, S. J. and Zon, L. I. (2018). Stem cell safe harbor: The hematopoietic stem cell niche in
zebrafish. *Blood Adv.* **2**, 3063–3069.

Wiegreffe, C., Christ, B., Huang, R. and Scaal, M. (2007). Sclerotomal origin of smooth muscle cells
in the wall of the avian dorsal aorta. *Dev. Dyn.* **236**, 2578–2585.

Wopat, S., Bagwell, J., Sumigray, K. D., Dickson, A. L., Huitema, L. F. A., Poss, K. D., Schulte-
Merker, S. and Bagnat, M. (2018). Spine Patterning Is Guided by Segmentation of the
Notochord Sheath. *Cell Rep.* **22**, 2026–2038.

Xie, T., Wang, Y., Deng, N., Huang, G., Taghavifar, F., Geng, Y., Liu, N., Kulur, V., Yao, C., Chen,
P., et al. (2018). Single-Cell Deconvolution of Fibroblast Heterogeneity in Mouse Pulmonary
Fibrosis. *Cell Rep.* **22**, 3625–3640.

Yasutake, J., Inohaya, K. and Kudo, A. (2004). Twist functions in vertebral column formation in
medaka, Oryzias latipes. *Mech. Dev.* **121**, 883–894.

Yeo, G. H., Cheah, F. S. H., Winkler, C., Jabs, E. W., Venkatesh, B. and Chong, S. S. (2009).
Phylogenetic and evolutionary relationships and developmental expression patterns of the
zebrafish twist gene family. *Dev. Genes Evol.* **219**,.

Figure 1

Figure 2

Figure 3

A Fin fold — Fin mesenchymal cells (*hmcn2+; fbln1+*); Apical epidermal cells (*hmcn2+; fras1+*)

B *fras1 fbln1* | *hmcn2 fbln1*

C *nkx3.1:Gal4; UAS:NTR-mCherry* 52 hpf
fbln1 mCherry | *fras1 mCherry*

D *nkx3.1:Gal4; UAS:Cre-ERT2; ubi:Switch* +4OHT at 1-2 dpf, imaged at 4 dpf
GFP (un-switched) mCherry (Switched)
Mid-trunk | Posterior trunk | Caudal fin

E *pax7b:Gal4FF; UAS:EGFP* 50 hpf

F Sclerotome: *nkx3.1^NTR-mCherry*
Dermomyotome: *pax7b^EGFP*
% of fin cells
**** ****
nkx3.1 *pax7b* | *nkx3.1* *pax7b*
dorsal fin | ventral fin

Figure 4

A

nkx3.1:Gal4; UAS:Cre-ERT2; ubi:Switch +4OHT 1-2 dpf, imaged at 17 dpf, **Calcein** **mCherry (Switched)**

B

nkx3.1:Gal4; UAS:Cre-ERT2; ubi:Switch +4OHT 1-2 dpf

GFP (un-switched) **mCherry (Switched)**

transverse section

6 wpf
19 mm

sagittal section

1 ypf
30 mm

C

nkx3.1:Gal4; UAS:NTR-mCherry +MTZ 1-2 dpf, stained at 3 wpf
Alcian blue **Alizarin red**

mCherry⁻   mCherry⁺

D

Standard Length (mm)
****

E

% of vertebrae
**** **
● mCherry-
○ mCherry+

Dorsal defects   Ventral defects

Figure 5

Figure 6

A Conversion of single dorsal sclerotome cell

Photoconversion 405nm

24 hpf

24 h

48 hpf

Type I cells
Type II cells

nkx3.1:Gal4; UAS:Kaede

B

% of clones

n=7, n=23, n=5, n=4, n=3

Cells in clone
1 2 3 4 5

C

% of clones

n=25, n=16, n=1

Cell types in clone
1 2 3

D

% of total cells

n=35, n=30, n=23, n=11

F T V I

E

% of cells

n=14, n=19

Dorsal Ventral

Type I
Type I and II
Type II

F

nkx3.1:Gal4; UAS:Kaede

| 24 hpf | 48 hpf |

Dorsal projecting cell

Ventral projecting cell

Figure 7

A Conversion of single ventral sclerotome cell

nkx3.1:Gal4; UAS:Kaede

24 hpf
48 hpf
24 h

Photoconversion
405nm
● Lagging cell
● Leading cell

B

% of cells

- SRC only
- SRC + Fin
- Fin only

n=22   n=20

Lagging   Leading

C

nkx3.1:Gal4; UAS:Kaede

| 24 hpf | 48 hpf |

Lagging cell

Leading cell

D

fin fold

a) Spatial regulation
b) Migration direction
c) Migration order

n

a

b

c

yolk extension

caudal vein plexus

fin fold

Figure S1

Figure S2

nkx3.1:Gal4; UAS:Kaede; kdrl:mCherry 2 dpf

Major Lobe

Minor Lobe

Figure S3

kdrl:EGFP; nkx3.1:Gal4; UAS:NTR-mCherry, 25 - 50 hpf