occAssess: An R package for assessing potential biases in species occurrence data

[1]Boyd, R. J., [1,2]Powney, G., [1]Carvell, C., [1]Pescott, O. L.

[1]UK Centre for Ecology and Hydrology, MacLean Bldg, Benson Ln, Crowmarsh Gifford, Wallingford OX10 8BB

[2]Oxford Martin School & School of Geography and Environment, University of Oxford, Oxford, OX1 3BD, UK

Abstract

Species occurrence records from a variety of sources are increasingly aggregated into heterogeneous databases and made available to ecologists for immediate analytical use. However, these data are typically biased, i.e. they are not a probability sample of the target population of interest, meaning that the information they provide may not be an accurate reflection of reality. It is therefore crucial that species occurrence data are properly scrutinised before they are used for research. In this article, we introduce occAssess, an R package that enables straightforward screening of species occurrence data for potential biases. The package contains a number of discrete functions, each of which returns a measure of the potential for bias in one or more of the taxonomic, temporal, spatial and environmental dimensions. Users can opt to provide a set of time periods into which the data will be split; in this case separate outputs will be provided for each period, making the package particularly useful for assessing the suitability of a dataset for estimating temporal trends in species' distributions. The outputs are provided visually (as ggplot2 objects) and do not include a formal recommendation as to whether data are of sufficient quality for any given inferential use. Instead, they should be used as ancillary information and viewed in the context of the question that is being asked, and the methods that are being used to answer it. We demonstrate the utility of occAssess by applying it to data on two key pollinator taxa in South America: leaf-nosed bats (Phyllostomidae) and hoverflies (Syrphidae). In this worked example, we briefly assess the degree to which various aspects of data coverage appear to have changed over time. We then discuss additional applications of the package, highlight its limitations, and point to future development opportunities.

Key words:

Bias; biological records; R; species distributions; species occurrence data; convenience samples; nonprobability samples

Introduction

Species occurrence records comprise information in three basic dimensions: taxonomic, geographic, and temporal; that is to say, what was seen, where was it seen, and when. Humans have been accumulating species occurrence data for centuries: historically as preserved specimens in museums and herbaria (Newbold, 2010; Spear et al., 2017), and in written accounts (e.g. Oswald and Preston, 2011); and more recently through recording for distribution atlases (Preston, 2013), and various other structured and unstructured monitoring and citizen science initiatives (Boakes et al., 2010;

Pescott et al., 2015; Petersen et al., 2021). Taken together, these data provide an immense resource
documenting species' geographical distributions and opportunities to investigate how they may have
changed over time. Over the last two decades, species occurrence data have become increasingly
accessible, thanks to the digitisation of historic records and the launch of online data portals such as
the Global Biodiversity Information Facility [GBIF; Nelson and Ellis (2019)]. A corollary of this increase
in accessibility has been a surge in the use of species occurrence data for research in biodiversity
conservation and other fields (Ball-Damerow et al., 2019).

Whilst clearly an increasingly important resource for ecologists, species occurrence data should be
used with caution when drawing inferences about species' distributions and how they have changed
over time. Straightforward inference in statistics is predicated on the assumption that the data have
been sampled randomly from the population of interest [probability sampling; e.g., Krzanowski
(2010)]. In many, if not most, cases, species occurrence data available through aggregated databases
do not satisfy this assumption. For example, data collected through citizen science initiatives tend to
be collected opportunistically (sometimes called convenience sampling); that is, without a structured
sampling plan. In this case, recorders are free to decide what to record, where and when. This
generally leads to preferential sampling of attractive and accessible locations, and to documentation
of interesting (e.g., rare) species (Isaac and Pocock, 2015). These "sampling biases" give rise to
nonprobability samples which are not representative of the spatial, temporal and taxonomic
domains of interest. Structured monitoring data tend to more closely resemble probability samples
(although issues like sample dropout and patchy uptake may still create issues). However, when
multiple structured datasets, with different aims, extents and sampling protocols, are aggregated
(e.g. as on GBIF), the ultimate target population sampled by these activities is unlikely to be formally
identified for inferential purposes. It may be possible to mitigate for biases by modifying the data
[e.g., spatial thinning; Beck et al. (2014)] or through the use of statistical correction procedures (e.g.,
by modelling the data generation process; Turner et al., 2009). In order to decide on what mitigating
action might be required, or if the data are simply too unrepresentative for a given use, it would be
helpful to have a set of heuristics that can indicate the degree to which a dataset might suffer from
various forms of bias.

There is a growing literature of studies which take species occurrence datasets and screen them for
biases (Barends et al., 2020; Boakes et al., 2010; Meyer et al., 2016; Pescott et al., 2019a; Petersen
et al., 2021; Ruete, 2015; Speed et al., 2018; Sumner et al., 2019; Troudet et al., 2018); we also note
that various approaches to visualising the spatial and temporal coverage of occurrence records
across large areas have been commonplace in national species atlases for some time (e.g. Preston et
al., 2002). Studies of these types provide a template for how to conduct such assessments, and a
suite of heuristics which can be deployed in similar situations. For example, one could assess data for
spatial bias by comparing the nearest neighbour distances of the occurrence data with those from a
simulated random distribution (Sumner et al., 2019). The proportion of records identified to species
level can be used as a measure of how taxonomic uncertainty has changed over time (Troudet et al.,
2018). Multidimensional environmental space can be summarised using principal component
analyses (PCAs), or other ordination techniques, allowing one to map the distribution of records in
environmental space and scrutinise it for bias relative to the total domain of interest (Pescott et al.,
2019b). Whilst these metrics are often presented in studies whose primary aim is to assess datasets
for their limitations, we find that they are rarely presented in studies which use such aggregated
species occurrence data to investigate actual patterns of species' distributions and how they have

changed over time [see Ball-Damerow et al. (2019) for a sobering review of the lack of scrutiny
where species occurrence data are used across research fields more generally].

One way to encourage the proper use of species occurrence data is to develop software that can
facilitate the various tasks involved, thereby easing the burden on researchers' time. Indeed, a suite
of packages have been developed in the R statistical programming environment (R Core Team, 2019)
to facilitate the acquisition, cleaning and proper acknowledgement of species occurrence data
(Chamberlain et al., 2021; Owens et al., 2021; Zizka et al., 2019). Recently, Zizka et al. (2021)
developed what is, to our knowledge, the first R package dedicated to quantifying sampling biases in
species occurrence data. The package, called sampbias, quantifies the relative strengths of various
geographical biasing factors, such as roads, cities and airports, in a given dataset. While sampbias
provides useful information on a set of possible geographical biases in species occurrence data, it is
not designed to screen data for biases in other dimensions (e.g. taxonomic, temporal and
environmental), and is limited to a specific set of data-biasing mechanisms and the assumption that
data point locations are accurate (rather than, for example, grid-based summaries). It would be
useful, therefore, to build on the functionality provided by sampbias and develop additional
software that can screen species occurrence data for more general biases in a range of possible
dimensions. Note that we do not think that bias screening can ever be a completely automatic or
easy task: assessing the great number of things that could go wrong, or be misinterpreted, between
the numerous data collection, collation, digitisation, and interpretation tasks embodied by the use of
any slice of any aggregated database, for any given inferential purpose, should humble any scientist
(e.g. Pescott et al., 2018). Nevertheless, making some basic "risk of bias" assessments more
straightforward, and raising their profile, is a step in the right direction for ecology.

Here then, we present occAssess: an R package for assessing potential biases in species occurrence
data. The package takes a user-supplied dataset and returns a suite of metrics that have been used
in the literature to assess species occurrence data for common issues when broad-scale inferences
relating to distributions and their changes may be desired. occAssess is designed primarily to assess
the suitability of species occurrence data for estimating temporal trends in species' distributions.
Nevertheless, the package should also be useful for those who would like to screen their data for
biases of potential importance when estimating spatial variation in species' occurrences with no
explicit reference to time (e.g. using static species distribution models). The aim is to enable quick
and easy screening of data for common limitations, thereby enabling researchers to properly
scrutinise their data before using it in further analyses, whatever their inferential goal may be. We
start by providing an overview of the package, what data it requires, and what outputs it returns. We
then provide a worked example using data on the occurrences of leaf-nosed bats and hoverflies in
South America over the period 1950 to 2019, and refer the reader to the supporting information
where additional vignettes and tutorials can be found. Finally, we discuss different ways in which the
package can be used, highlight its limitations, and suggest how it could be improved in the future.

Package

Package specifications
occAssess is an open-source R (version >= 4.0.0) package (R Core Team, 2019), built around the
existing packages ggplot2 (Wickham, 2016), spatstat (Baddeley et al., 2015), raster (Hijmans, 2019),
and stats (R Core Team, 2019). A stable version (1.3.0) can be found at

and the development version can be found at
.. We provide three vignettes with the package: 1) a tutorial
using the data presented in this article; 2) a second example using data that are simulated to be
unbiased for the purpose of estimating trends in species' distributions; and 3) a fully-reproducible
example for which all required data are available within the package. Note that all required data is
not provided with vignettes one and two; they are provided for instruction, rather than reproducible
examples.

Package structure
occAssess comprises seven discrete functions (Table 1), each of which is designed to assess a
common form of potential bias in species occurrence data. The functions each assess species
occurrence data in at least one of the spatial, temporal, taxonomic, and environmental dimensions.
The user can provide a set of time periods into which the data will be split, meaning that all
functions are to some extent temporally explicit. For example, one function assesses spatial bias in
the data, but, if multiple periods are specified, then the function provides information on *temporal*
variation in spatial bias. We provide the option to split the data into periods to facilitate assessments
of the suitability of data for estimating *changes* in species distributions over time. However, in some
cases it may be preferable to specify one time period, perhaps covering the entire temporal extent
of the data. This may be useful for static species distribution modelling where one simply requires
information on, e.g., spatial or environmental bias in the dataset as a whole. At present the time
periods must be specified in units of years, and the minimum permitted length for a time period is
one year (see Discussion below).

Table 1. Summary of the functions provided in occAssess. Note that users can opt to split the data
into multiple time periods; in this case all functions are temporally explicit and hence provide
information on temporal variation in some characteristic of the data. See the worked example in the
main text for more details of each function.

Function	Type of bias assessed	Method (in brief)	Output	Dataset type	Additional data required
assessEnvBias	Environmental bias. Indicates whether the input data are likely to be sampled from the same portion of environmental space over time, or whether the data are sampled from a representative portion of	Reduces the dimensionality of environmental space using principal component analyses and maps the distribution of the data in this reduced environmental space	Maps of the distribution of the data on two user-selected principal components of environmental space. Displayed as ellipses or points. One ellipse or set of points per period	Single or multispecies. Highly likely to indicate strong bias with one or a small number of species	Environmental data corresponding to each occurrence data point and, optionally, a background sample (e.g. a random sample from the domain of interest for inference)

	environmental space in the spatial domain of interest				
assessRarityBias	Indicates whether rare species are overrepresented in the data and whether the degree to which they are overrepresented changes over time	Measures the congruence of the number of times species have been recorded and their estimated commonness (range sizes). Drops records not identified to species level	Time series showing congruence in each period (correlation or r^2 from regression of the number of records on commonness)	Multispecies only	None
assessRecordNumber	Identifies temporal variation in sampling intensity in the domain of interest	Sums the number of records in the dataset in each time period	Time series of counts	Single or multispecies	None
assessSpatialBias	Indicates whether the data resemble a random distribution in the geographic space of interest for inference, and whether the extent to which the data resemble a random distribution changes over time	Compares the average nearest neighbour distance of the data with the average nearest neighbour distances of simulated random distributions of the same density	Time series showing nearest neighbour index in each period	Single or multispecies. Highly likely to indicate strong bias with one or a small number of species	Raster layer indicating which areas fall inside the study extent
assessSpatialCov	Indicates whether a representative portion of the spatial domain of interest has been sampled,	Maps the data in geographical space	Either multiple (gridded) maps showing the distribution of the data in	Single or multispecies	If the input data are not on WGS84 coordinate reference system, then any country

			each period, or one map showing the number of periods in which grid cells have been sampled		or political boundaries that the user requires to be superimposed on the resultant plots must be supplied
assessSpeciesID	Taxonomic resolution and whether it changes over time	Calculates proportions or counts of records identified to species level	Time series of proportions or counts	Multispecies only	None
assessSpeciesNumber	Taxonomic coverage and how it changes over time	Sums the number of species recorded in each time period. Drops records not identified to species level	Time series of counts	Multispecies only	None

Input data
For all functions, users must provide their occurrence data and a list of time periods into which the
data should be split. The occurrence data must be provided as a dataframe object with six fields:
species (species name; note that whilst we use the word "species" here for convenience, essentially
any set of taxonomic levels could be used), x (x coordinate), y (y coordinate), spatialUncertainty
(uncertainty associated with the x and y coordinates; any units are permitted), year and identifier.
The column names in the input data need not match the names of the fields above; rather, the user
must pass arguments to each function indicating what columns in their data correspond to which
field. This ensures compatibility with data standards such as Darwin Core (https://dwc.tdwg.org/).
For example, in Darwin Core, the spatialUncertainty field would be called
coordinateUncertaintyInMetres, and the user can provide a mapping by specifying
spatialUncertainty = "coordinateUncertaintyInMetres". We would expect that
information on all six required fields would be provided by any typical species occurrence data
aggregator, e.g. GBIF. Note that users may specify a threshold spatial uncertainty above which data
are dropped before the heuristics are calculated. This allows users to ask the question "how do the
biases in my data change if I retain only the more precise records?". Any coordinate reference
system (CRS) may be used. In the spatialUncertainty field, any units are permitted (e.g. metres for
eastings/northings, or decimal degrees for lon/lat) but they must be consistent. The identifier field is
used to group the data; for example, it may denote specific taxonomic groups, countries, datasets

etc. Where there is no information available for a field, its values should be set to NA. See Table 2 for
an example set of input data.

Table 2. The first six rows of an example dataset as required by occAssess. Note that any units are
permitted in the spatialUncertainty field (here metres) but they must be consistent. Also note that
the column names in the input data need not match those in this example: users can provide a
mapping between their data and the fields presented here using arguments to each function.

species	x	y	year	spatialUncertainty	identifier
Anoura caudifer	-65.4	-17.0667	1993	11839	Phyllostomidae
Carollia perspicillata	-65.5497	-17.1072	1993	1043	Phyllostomidae
Carollia perspicillata	-65.4	-17.0667	1993	11839	Phyllostomidae
Sturnira erythromos	-65.8692	-17.2119	1993	1043	Phyllostomidae
Platyrrhinus dorsalis	-65.5497	-17.1072	1993	1043	Phyllostomidae
Artibeus lituratus	-56	-25.4667	1995	11010	Phyllostomidae

Outputs
Each function returns a list with two elements: a ggplot2 (Wickham, 2016) object, and the data that
underpin that plot. The ggplot2 objects generally display the various potential bias metrics for each
level of the identifier field (Table 2) and for each time period specified. We provide the outputs as
ggplot2 objects because these can be subsequently modified by the user for presentation in e.g.
published articles or supplementary material. The functions do not provide any formal
recommendation as to whether the data are too biased for any given inferential use; instead, we
expect that the heuristics will be used in combination with researchers' expert judgement to decide
on whether mitigating action must be taken, and how this might be done (if indeed it is possible at
all). In supplementary material 2 we provide the outputs of occAssess as applied to a simulated
dataset that has a random distribution in space and time, and is resolved to species level in all cases;
this is taken as an example of a dataset that is unbiased relative to the inferential use case of
assessing all species' distributions in a region over time. These outputs can be used as a point of
comparison in that they are likely to provide examples of how the heuristics would appear if a
dataset is unbiased.

Worked example
In this section, we provide a worked example of the functionality of occAssess. We use the package
to assess data on the occurrences of leaf-nosed bats and hoverflies in South America over the period
1950 to 2019. The data were downloaded from GBIF (GBIF, 2021; DOI in reference list) and were
cleaned for spatial issues (e.g. coordinates matching country centroids, capital cities, biodiversity
institutes, etc.) using the CoordinateCleaner package (Zizka et al., 2019). We specify seven time
periods, each one decade in duration. We use the identifier field to distinguish between the leaf-
nosed bats (Phyllostomidae) and hoverflies (Syrphidae). We do not provide the code in the main
text; instead, we refer the reader to the vignette in supplementary material 1 which provides the
code for this example. As we introduce each function, and where where applicable, we 1) outline
what form of bias it relates to and in what dimension(s); 2) provide the theory behind the metric; 3)
indicate where additional inputs – beyond the fields in Table 2 – are required; 4) present the ggplot2
object returned for this case study (noting that the data underpinning these plots are also returned
by each function); and 5) give guidance on how to interpret the outputs. We reiterate here that

these heuristics are designed to be used alongside expert judgement and careful thought relative to
the inferences desired by the analyst – we do not intend any function to provide a simple binary
answer to the question "are these data biased for answering my question?". Biases are challenging!

assessRecordNumber()

The simplest function in occAssess, assessRecordNumber, provides a measure of sampling intensity
in the domain of interest and how it changes over time (Fig. 1A). Although simple, it is important to
understand the extent to which the quantity of data varies over time, because a change in the
number of records could reflect a change in recording intensity, which is itself likely to affect the
prevalence of particular species in the dataset through time in a non-random fashion (Pescott et al.,
2019a).

One problem that may arise when using assessRecordNumber is that the counts may differ widely
between levels of identifier. This can make it difficult to assess temporal variation in record counts
for the level(s) with fewer records. To circumvent this problem, we include the option to normalize
the counts for each level of identifier. In this case, the indices for each level of identifier fall on
comparable scales.

Figure 1. ggplot2 objects returned by A) assessRecordNumber; B) assessSpeciesNumber; C)
assessSpeciesID; D) assessRarityBias; and E) assessEnvBias. Note that the ggplot2 objects can be
modified by the user (e.g. colours, axis labels, etc.).

assessSpeciesNumber()
The function assessSpeciesNumber returns a measure of taxonomic coverage and how it changes
over time. The function sums the number of species recorded in each time period and for each level
of identifier and displays the results as time series (Fig. 1B). Of course, changes in the numbers of
species recorded could also reflect true extinction/colonisation events in a dataset, but, for
heterogeneous, aggregated, data, issues of uneven representativeness across time are considerably
more likely. As with assessRecordNumber, users can choose to normalize the species counts for each
level of identifier for ease of interpretation.

assessSpeciesID()
The function assessSpeciesID provides a measure of taxonomic uncertainty and how it changes over
time. By default the function displays the proportion of records identified to species level each year
[Fig. 1C, as in Troudet et al. (2018) and Zattara and Aizen (2021)]. Records are considered not
identified to species level if they take the value NA. The user has the option to substitute
proportions for counts which may be preferable in some circumstances. For example, it is feasible
that, due to the increase in the number of records submitted by volunteer citizen scientists over
time, the *proportion* of records identified to species level may decrease, but the overall *quantity* may
show a different trend.

assessRarityBias()
The function assessRarityBias can be used to assess the degree to which rare species are
oversampled relative to commoner species, and whether this changes over time. The idea is that,
was there no sampling bias, species would be recorded in proportion to their commonness.
Commonness can be defined as local abundance or regional occupancy (Gaston, 2011). Following
Speed et al. (2018), we define a species' commonness as the number of grid cells on which it has
been recorded – a proxy for regional occupancy. The user may decide on the spatial resolution of the
grid cells, and whether commonness is calculated over the entire temporal extent of the data, or
separately for each time period (which could have important implications for the interpretation of
discovered patterns, given other biases in the dataset).

Once the numbers of times species' have been recorded, and their commonness, have been
calculated, assessRarityBias measures the congruence between these two quantities. The user may
decide on one of two methods that the function will use to do this. The first option is to regress the
number of records on commonness and use the r^2 (coefficient of variation) from the fitted model as
an indicator of to what extent the number of records are explained by range size. This method is an
extension of that used by Barends et al. (2020) and Speed et al. (2018) who fitted analogous
regression models and treated each species' residual as an index of whether they are over- or under-
sampled relative to some wider assemblage. This measure ranges from 0, indicating high bias, to 1,
indicating low bias. The second option is to use the Pearson's correlation coefficient between the
number of times species' have been recorded and their commonness as the measure congruence.
This measure ranges from -1 – 1, with values closer to 1 indicating smaller bias. Whichever method is
chosen, occAssess displays the index for each time period and level of identifier (Fig. 1D). Note that

both metrics produced assessRarityBias indicate the strength of the *linear* relationship between
range size and the number of records; users may wish to inspect the data for curvillinearity.

assessSpatialCov()

The function assessSpatialCov can be used to assess the extent to which the data are spatio-
temporally biased; that is, the extent to which the same portion of the geographic domain has been
sampled over time — note that this is likely to be crucial for robust estimates of temporal
distributional change. The function provides this information in one of two ways (selected by the
user). Both methods begin by gridding the data at a user-specified spatial resolution. The first
method then returns n ggplot2 objects, where n is the number of levels in the identifier field. Each
ggplot2 object contains N maps showing the density of records in each grid cell, where N is the
number of time periods. The second method returns one map showing the number of time periods
in which each grid cell has been sampled (Figs 2B and 2C; see supplementay materials 2 and 3 for
examples using the first method).It is worth pointing out that data originally provided on a grid are
often converted to point format by online data aggregators (e.g., using cell centroids). For these
data, it is possible that the mismatch between the original grids and the user-specified grid produced
by assessSpatialCov could result in some unexpected biases.

In some circumstances users will need to pass additional data to assessSpatialCov to superimpose
political/ geographical boundaries on the resultant plots. This is not required where the data are on
the WGS84 coordinate reference system; in this case, the user must simply specify the relevant
countries, otherwise, a shapefile is required.

Figure 2. ggplot2 objects returned by assessSpatialBias (A) and assessSpatialCov (B and C). Note that the user can modify these plots (e.g., by changing colours, axis labels, etc.).

assessSpatialBias()

The function assessSpatialBias screens data for geographical bias, i.e. the degree to which a sample deviates from a random distribution within the spatial domain of interest. The function is based on the widely-used nearest neighbour index (NNI; Clark and Evans, 1954). The NNI is given as the ratio of the average observed nearest neighbour distances (the Euclidean distance of each data point to its nearest neighbouring point) to the expected average nearest neighbour distance if the data were randomly distributed. In the standard NNI, the average expected nearest neighbour distance for a random distribution is given by $1/2 \sqrt{study\ area/number\ of\ points}$. However, in the case of irregularly shaped study boundaries (e.g., political or geographical boundaries), the above formula does not equal the expected average nearest neighbour distances for a random distribution. To circumvent this problem, assessSpatialBias simulates n datasets randomly across the study area in equal number to the occurrence data. The NNI can then be given as the ratio of the average observed nearest neighbour distances to the average of the simulated nearest neighbour distances (Fig. 2A). Another advantage of this approach is that, by simulating n (chosen by the user) random datasets, assessSpatialBias can provide uncertainty associated with the index (the function will display 90% confidence intervals by default). The NNI produced by assessSpatialBias can be interpreted as how far the observed distribution deviates from a random distribution of the same density. Values between 0 and 1 are more clustered than a random distribution, and values between 1 and 2.15 are more widely dispersed (i.e., over-dispersed). See Sumner et al. (2019) for a somewhat similar approach.

It is worth pointing out that the NNI produced by assessSpatialBias is a function of both sampling biases in the data and the true distributions of the focal taxa. If the function is used to assess data for one or a small number of species, the NNI will likely indicate a strong departure from a random distribution. This is to be expected because the geographical distribution of records will reflect e.g., the environmental niche of the target taxa. The function is therefore most appropriate for use with data spanning many species, in which case a more accurate picture of the distribution of sampling is likely to be obtained.

assessEnvBias()

The function assessEnvBias can be used to assess species occurrence data for two types of environmental bias: unrepresentative sampling in the environmental space of the domain of interest, and uneven sampling of environmental space over time. The function maps the data in environmental space in each user-specified time period. To do so, additional environmental data are required. As a minimum users must supply environmental data (this can be many variables) at the coordinates of the occurrence data. Users may optionally supply a "background" sample of the same environmental variables; this may be, for example, the environment at random locations across the domain of interest. Whether or not background data are supplied impacts interpretation of the assessEnvBias outputs. If background data are supplied, then the function maps the distribution of the occurrence data in the environmental space of the domain of interest. Otherwise, the data are mapped in the *sampled* environmental space across all periods. In this example we use the standard suite of 19 bioclimatic variables from worldclim (Fick and Hijmans, 2017). These data can be

downloaded at https://www.worldclim.org/data/worldclim21.html or through R using the getData
function in the raster package (Hijmans et al., 2016).

assessEnvBias reduces the dimensionality of the environmental data using PCAs. It then maps the
data in two dimensional environmental space (Fig. 1G), enabling the user to assess whether their
data are sampled from the same portion of environmental space across periods or, if background
data are supplied, whether the data are sampled from a representative portion of environmental
space in the domain of interest. By default, the data are displayed as ellipses delimiting 95% of the
occurrence data. Strictly speaking, PCAs assume multivariate normality in the environmental data,
and the ellipses displayed by assessEnvBias assume multivariate normality among the principal
component scores. Users may wish to assess their data, and the resultant PC scores (which are
returned by the function), for normality. If the data are non-normal, then transformations can be
applied. If the PC scores are non-normal, it is simple to substitute the ellipses for the actual data
points (see s1 for more details). For similar approaches see Pescott et al. (2019b) and Barends et al.
(2020). Note that this assessment assumes that the spatial resolution of the environmental data are
relevant to the responses of the target organism(s) at the spatial scale of the analysis desired.

Discussion
In this paper, we have introduced a new R package, occAssess, which enables rapid screening of
species occurrence data for biases of potential importance for drawing inferences about species'
distributions and how they have changed over time. The package takes a species occurrence dataset
as input, and returns a number of metrics relating to common forms of bias in one or more of the
taxonomic, temporal, spatial, and environmental dimensions. None of the metrics provided in the
package are new (although some are extended and/or modified). However, we hope that in
assembling these metrics in an easy-to-use R package, we will ease the burden on researchers who
would like to scrutinise their data. In turn, we hope to promote the proper assessment of species
occurrence data *before* they are used in attempts to answer important research questions regarding
ecological change. The heuristics returned by occAssess could be provided as, for example,
supplementary material to published articles to provide evidence of the fact that a proper
assessment has been conducted. In general, we would expect such evidence of assessment to be
accompanied by written commentary interpreting the patterns seen and considering their
implications for any analyses presented.

We have presented a single example of how occAssess may be used, but it is easy to imagine
additional use cases. In our example, we used the identifier field (Table 2) to split the data by
taxonomic group (Phyllostomidae and Syrphidae). One might instead use the identifier field to
denote specific datasets. For example, one level of identifier could denote a dataset before some
newly-digitized data were added, and a second could denote the same data with the addition of the
newly-digitized records. It would then be possible to make an assessment of to what extent the data
have improved as a result of digitization efforts. occAssess could also be used for model-based data
integration (Isaac et al., 2020), where the aim is to exploit the strengths of multiple datasets, each of
which could be specified in the identifier field. Another possibility is that occAssess could be used to
screen data for single species as opposed to whole taxonomic groups as presented in our worked
example. In this case note that some heuristics would require different interpretations; for example,
one would expect the data to be biased in environmental space relative to the domain of interest

because it would reflect a species' environmental niche. In summary, we feel that occAssess has the
potential to be useful for many applications where species occurrence data are used.

A key feature of occAssess is the periods argument in each function, which enables assessment of
how the limitations of a dataset may change over time. We include this feature because a common
application of species occurrence data is the estimation of temporal trends in species' distributions
(e.g. Outhwaite et al., 2019; Pescott et al., 2019a; Powney et al., 2019). For some applications,
however, it may be more appropriate to consider an entire dataset as comprising one time period,
thereby removing the temporal dimension. An obvious example is where data are to be used for
species distribution modelling (SDM). In this case the objective is typically estimation of spatial
variation in species' occurrences with no explicit reference to time (Guisan, 2017). Where occAssess
is used to screen data for use in SDMs, we suggest that the functions relating to spatial and
environmental bias will be of most importance, namely assessSpatialBias, assessSpatialCov and
assessEnvBias (although the other functions could still provide important context on the temporal
dynamics of the dataset).

The functions in occAssess provide heuristics relating to the quality of species occurrence data, but
stop short of making a formal recommendation as to whether the data are of sufficient quality for
any given use. It would not be appropriate to provide such recommendations, because the utility of
species occurrence data depend not only their biases, but also on the question being asked and the
methods used to answer it. For example, it may be possible to obtain relatively unbiased predictions
of species' geographical distributions using SDMs, even when the data themselves are spatially and
environmentally biased. Phillips et al. (2009) developed the "target group" approach whereby
background data are generated with similar sampling biases to the occurrence data. This approach
helps SDMs to distinguish between suitable and unsuitable habitats as opposed to popular and
unpopular sampling locations. There have also been attempts to correct for changes in recorder
effort statistically, thereby enabling estimation of how species' distributions have changed over time
from biased data (Franklin, 1999; Hill, 2012; Isaac et al., 2014; Szabo et al., 2010; Telfer et al., 2002;
Van Strien et al., 2013). While it is not always clear to what extent the above-mentioned methods
achieve the goal of mitigating for sampling biases, the point remains that relatively informative
inferences may be possible from biased data where the biases can either be modelled, or reduced
through appropriate resolution-based aggregation (Pescott et al., 2019a), or through more complex
methods designed to leverage unbiased estimates of model parameters from additional probability
samples (e.g. Ahmad Suhaimi et al., 2021). It is for this reason that we suggest the metrics provided
by occAssess be consulted in combination with other relevant information in order to decide
whether or not a dataset is of sufficient quality for use for a given inferential purpose.

The version of occAssess presented here has two key limitations. First, the temporal unit is the year,
meaning that the package can say nothing about intra-annual biases (e.g. phenological patterns in
the data). In future versions it might be feasible to increase the temporal resolution of the package.
Second, it will not always be possible to tease apart biases from true biological phenomena using the
package alone. For example, assessSpatialBias indicates whether the data deviate from a random
distribution but, particularly where there are few species in the dataset, it might not be clear
whether this reflects sampling biases or species' true distributions. To disentangle sampling biases
and the biological truth it will always be preferable to solicit advice from experts who are familiar

with the biology of the focal taxa – we stress again that our package is a compliment to, not
substitute for, expert knowledge.

Acknowledgements

We would like to thank Francesca Mancini, Nick Isaac and Robert Cooke for their helpful comments
on the heuristics presented in this article. RB, GP and CC were funded by the SURPASS2 project
under the Newton Fund Latin America Biodiversity Programme: Biodiversity - Ecosystem services for
sustainable development, awarded by the UKRI Natural Environment Research Council (NERC)
NE/S011870/2. We thank SURPASS2 partners Francisco Fonturbel, Marcelo Aizen, Eduardo Zattara,
Antonio Saraiva and Jeff Ollerton for comments on the outputs presented. The contribution of OLP
was supported by the Natural Environment Research Council award number NE/R016429/1 as part
of the UK Status, Change and Projections of the Environment (UK-SCAPE) programme delivering
National Capability.

Supporting information

We provide three vignettes as supporting information: S1) a vignette showing the code for the
worked example presented in the manuscript; S2) a second example using data that have been
simulated to unbiased relative to the inferential goal of estimating temporal trends in species'
distributions; and 3) a fully-reproducible example for which all data are available in the package.

References

Ahmad Suhaimi, S.S., Blair, G.S., Jarvis, S.G., 2021. Integrated species distribution models: A
comparison of approaches under different data quality scenarios. Divers. Distrib. 1–10.
doi:10.1111/ddi.13255

Baddeley, A., Rubak, E., Turner, R., 2015. Spatial point patterns: Methodology and applications with
R. Chapman & Hall, London.

Ball-Damerow, J.E., Brenskelle, L., Barve, N., Soltis, P.S., Sierwald, P., Bieler, R., LaFrance, R., Ariño,
A.H., Guralnick, R., 2019. Research applications of primary biodiversity databases in the digital
age. PLoS One 1–26. doi:10.1101/605071

Barends, J.M., Pietersen, D.W., Zambatis, G., Tye, D.R.C., Maritz, B., 2020. Sampling bias in reptile
occurrence data for the Kruger National Park. Koedoe 62, 1–9. doi:10.4102/koedoe.v62i1.1579

Beck, J., Böller, M., Erhardt, A., Schwanghart, W., 2014. Spatial bias in the GBIF database and its
effect on modeling species' geographic distributions. Ecol. Inform. 19, 10–15.
doi:10.1016/j.ecoinf.2013.11.002

Boakes, E.H., McGowan, P.J.K., Fuller, R.A., Chang-Qing, D., Clark, N.E., O'Connor, K., Mace, G.M.,
2010. Distorted views of biodiversity: Spatial and temporal bias in species occurrence data.
PLoS Biol. 8. doi:10.1371/journal.pbio.1000385

Chamberlain, S., Barve, V., Mcglinn, D., Oldoni, D., Desmet, P., Geffert, L., Ram, K., 2021. rgbif:
Interface to the 3.3.0, Global Biodiversity Information Facility API_. R package version.

Clark, P., Evans, F., 1954. Distance to Nearest Neighbour as a Measure of Spatial Relationships in

Populations. Ecology 35, 445–453. doi:10.1007/BF02315373

Fick, S.E., Hijmans, R.J., 2017. WorldClim 2 : new 1-km spatial resolution climate surfaces for global
land areas. Int. J. Climatol. doi:10.1002/joc.5086

Franklin, D.C., 1999. Evidence of disarray amongst granivorous bird assemblages in the savannas of
northern Australia, a region of sparse human settlement. Biol. Conserv. 90, 53–68.
doi:10.1016/S0006-3207(99)00010-5

Gaston, K.J., 2011. Common ecology. Bioscience 61, 354–362. doi:10.1525/bio.2011.61.5.4

GBIF, 2021. (7 April 2021) GBIF Occurrence Download: hoverflies and leaf-nosed bats.
doi:https://doi.org/10.15468/dl.wqz6z3

Guisan, A., 2017. Habitat Suitability and Distribution Models: With Applications in R, 1st ed.
Cambridge University Press.

Hijmans, R., van Etten, J., Cheng, J., Mattiuzzi, M., Sumner, M., Greenberg, J. a, Lamigueiro, O.,
Bevan, A., Racine, E., Shortridge, A., 2016. Package " raster ". CRAN -R.2.5-8.

Hijmans, R.J., 2019. raster: Geographic Data Analysis and Modeling. R package version 2.9-23.

Hill, M.O., 2012. Local frequency as a key to interpreting species occurrence data when recording
effort is not known 195–205. doi:10.1111/j.2041-210X.2011.00146.x

Isaac, N.J.B., Jarzyna, M.A., Keil, P., Dambly, L.I., Boersch-Supan, P.H., Browning, E., Freeman, S.N.,
Golding, N., Guillera-Arroita, G., Henrys, P.A., Jarvis, S., Lahoz-Monfort, J., Pagel, J., Pescott,
O.L., Schmucki, R., Simmonds, E.G., O'Hara, R.B., 2020. Data Integration for Large-Scale Models
of Species Distributions. Trends Ecol. Evol. 35, 56–67. doi:10.1016/j.tree.2019.08.006

Isaac, N.J.B., Pocock, M.J.O., 2015. Bias and information in biological records. Biol. J. Linn. Soc. 115,
522–531. doi:10.1111/bij.12532

Isaac, N.J.B., van Strien, A.J., August, T.A., de Zeeuw, M.P., Roy, D.B., 2014. Statistics for citizen
science: Extracting signals of change from noisy ecological data. Methods Ecol. Evol. 5, 1052–
1060. doi:10.1111/2041-210X.12254

Krzanowski, W., 2010. An introduction to statistical modelling, 1st ed. Wiley.

Meyer, C., Weigelt, P., Kreft, H., 2016. Multidimensional biases, gaps and uncertainties in global
plant occurrence information. Ecol. Lett. 19, 992–1006. doi:10.1111/ele.12624

Nelson, G., Ellis, S., 2019. The history and impact of digitization and digital data mobilization on
biodiversity research. Philos. Trans. R. Soc. B Biol. Sci. 374, 2–10. doi:10.1098/rstb.2017.0391

Newbold, T., 2010. Applications and limitations of museum data for conservation and ecology, with
particular attention to species distribution models. Prog. Phys. Geogr. 34, 3–22.
doi:10.1177/0309133309355630

Oswald, P.., Preston, C.D., 2011. John Ray's Cambridge Catalogue (1660)., (Eds). ed. Cambridge
University Press, London.

Outhwaite, C.L., Powney, G.D., August, T.A., Chandler, R.E., Rorke, S., Pescott, O.L., Harvey, M., Roy,
H.E., Fox, R., Roy, D.B., Alexander, K., Ball, S., Bantock, T., Barber, T., Beckmann, B.C., Cook, T.,
Flanagan, J., Fowles, A., Hammond, P., Harvey, P., Hepper, D., Hubble, D., Kramer, J., Lee, P.,
MacAdam, C., Morris, R., Norris, A., Palmer, S., Plant, C.W., Simkin, J., Stubbs, A., Sutton, P.,

Telfer, M., Wallace, I., Isaac, N.J.B., 2019. Annual estimates of occupancy for bryophytes,
lichens and invertebrates in the UK, 1970-2015. Sci. data 6, 259. doi:10.1038/s41597-019-0269-
1

Owens, H., Merow, C., Maitner, B., Kass, J., Barve, V., Guralnick, R., 2021. occCite: Querying and
Managing Large Biodiversity Occurrence Datasets. R package version 0.4.6. [WWW Document].
URL https://cran.r-project.org/package=occCite

Pescott, O.L., Humphrey, T.A., Stroh, P.A., Walker, K.J., 2019a. Temporal changes in distributions and
the species atlas: How can British and Irish plant data shoulder the inferential burden? Br. Irish
Bot. 1, 250–282. doi:10.33928/bib.2019.01.250

Pescott, O.L., Humphrey, T.A., Walker, K.J., 2018. A short guide to using British and Irish plant
occurrence data for. Wallingford. doi:10.13140/RG.2.2.33746.86720

Pescott, O.L., Walker, K.J., Harris, F., New, H., Cheffings, C.M., Newton, N., Jitlal, M., Redhead, J.,
Smart, S.M., Roy, D.B., 2019b. The design, launch and assessment of a new volunteer-based
plant monitoring scheme for the United Kingdom. PLoS One 14, 1–30.
doi:10.1371/journal.pone.0215891

Pescott, O.L., Walker, K.J., Pocock, M.J.O., Jitlal, M., Outhwaite, C.L., Cheffings, C.M., Harris, F., Roy,
D.B., 2015. Ecological monitoring with citizen science: The design and implementation of
schemes for recording plants in Britain and Ireland. Biol. J. Linn. Soc. 115, 505–521.
doi:10.1111/bij.12581

Petersen, T.K., Austrheim, G., Speed, J.D.M., Grøtan, V., 2021. Species data for understanding
biodiversity dynamics : The what , where and when of species occurrence data collection 1–17.
doi:10.1002/2688-8319.12048

Phillips, S.J., Dudík, M., Elith, J., Graham, C.H., Lehmann, A., Leathwick, J., Ferrier, S., 2009. Sample
selection bias and presence-only distribution models: Implications for background and pseudo-
absence data. Ecol. Appl. 19, 181–197. doi:10.1890/07-2153.1

Powney, G.D., Carvell, C., Edwards, M., Morris, R.K.A., Roy, H.E., Woodcock, B.A., Isaac, N.J.B., 2019.
Widespread losses of pollinating insects in Britain. Nat. Commun. 1–6. doi:10.1038/s41467-
019-08974-9

Preston, C.D., Pearman, D.A. & Dines, T.D., 2002. New Atlas of the British and Irish Flora., eds. ed.
Oxford University Press, Oxford.

Preston, C.D., 2013. Following the BSBI's lead: the influence of the Atlas of the British flora, 1962-
2012. New J. Bot. 3, 2–14.

R Core Team, 2019. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria.

Ruete, A., 2015. Displaying bias in sampling effort of data accessed from biodiversity databases using
ignorance maps. Biodivers. Data J. 3. doi:10.3897/BDJ.3.e5361

Spear, D.M., Pauly, G.B., Kaiser, K., 2017. Citizen science as a tool for augmenting museum collection
data from urban areas. Front. Ecol. Evol. 5, 1–12. doi:10.3389/fevo.2017.00086

Speed, J.D.M., Bendiksby, M., Finstad, A.G., Hassel, K., Kolstad, A.L., Prestø, T., 2018. Contrasting
spatial, temporal and environmental patterns in observation and specimen based species
occurrence data. PLoS One 13, 1–17. doi:10.1371/journal.pone.0196417

Sumner, S., Bevan, P., Hart, A.G., Isaac, N.J.B., 2019. Mapping species distributions in 2 weeks using
citizen science. Insect Conserv. Divers. 12, 382–388. doi:10.1111/icad.12345

Szabo, J.U.K.S., Vesk, P.E.A. V, Baxter, P.E.W.J.B., Possingham, H., 2010. Regional avian species
declines estimated from volunteer-collected long-term data using List Length Analysis. Ecol.
Appl. 20, 2157–2169.

Telfer, M.G., Preston, C.D., Rothery, P., 2002. A general method for measuring relative change in
range size from biological atlas data. Biol. Conserv. 107, 99–109. doi:10.1016/S0006-
3207(02)00050-2

Troudet, J., Vignes-Lebbe, R., Grandcolas, P., Legendre, F., 2018. The increasing disconnection of
primary biodiversity data from specimens: How does it happen and how to handle it? Syst. Biol.
67, 1110–1119. doi:10.1093/sysbio/syy044

Turner, R.M., Spiegelhalter, D.J., Smith, G.C.S., Thompson, S.G., 2009. Bias modelling in evidence
synthesis. J. R. Stat. Soc. Ser. A Stat. Soc. 172, 21–47. doi:10.1111/j.1467-985X.2008.00547.x

Van Strien, A.J., Van Swaay, C.A.M., Termaat, T., 2013. Opportunistic citizen science data of animal
species produce reliable estimates of distribution trends if analysed with occupancy models. J.
Appl. Ecol. 50, 1450–1458. doi:10.1111/1365-2664.12158

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis.

Zattara, E.E., Aizen, M.A., 2021. Worldwide occurrence records suggest a global decline in bee
species richness. One Earth 4, 114–123. doi:10.1016/j.oneear.2020.12.005

Zizka, A., Antonelli, A., Silvestro, D., 2021. Sampbias, a Method for Quantifying Geographic Sampling
Biases in Species Distribution Data. Ecography (Cop.). 44, 25–32. doi:10.1111/ecog.05102

Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Duarte Ritter, C., Edler, D., Farooq, H., Herdean,
A., Ariza, M., Scharn, R., Svantesson, S., Wengström, N., Zizka, V., Antonelli, A., 2019.
CoordinateCleaner: Standardized cleaning of occurrence records from biological collection
databases. Methods Ecol. Evol. 10, 744–751. doi:10.1111/2041-210X.13152

554