Deep Mutagenesis of a Transporter for Uptake of a Non-Native Substrate Identifies Conformationally Dynamic Regions

Heather J. Ellis,[1,□] Matthew Chan,[2,□] Balaji Selvam,[2,□] Evan Walter,[3] Christine A. Devlin,[1]

Steven K. Szymanski,[1] L. Keith Henry,[3] Diwakar Shukla,[2,4] and Erik Procko,[1,4]

1 Department of Biochemistry, University of Illinois, Urbana, IL, USA

2 Department of Chemical and Biomolecular Engineering, University of Illinois, Urbana, IL, USA

3 Department of Biomedical Sciences, University of North Dakota School of Medicine and Health Sciences, Grand Forks, ND, USA

4 Cancer Center at Illinois and Center for Biophysics and Quantitative Biology, University of Illinois, Urbana, IL, USA

□ These authors contributed equally.

Correspondence: procko@illinois.edu; diwakar@illinois.edu

Title Running Head: Deep mutagenesis of human SERT

Abstract

The serotonin transporter, SERT, catalyzes serotonin reuptake at the synapse to terminate neurotransmission via an alternating access mechanism, and SERT inhibitors are the most widely prescribed antidepressants. Here, deep mutagenesis is used to determine the effects of nearly all amino acid substitutions on human SERT surface expression and transport of the fluorescent substrate APP+, identifying many mutations that enhance APP+ import. Comprehensive simulations of the entire ion-coupled import process reveal that while binding of the native substrate, serotonin, reduces free energy barriers between conformational states to promote SERT dynamics, the conformational free energy landscape in the presence of APP+ instead resembles Na^+ bound-SERT, with a higher free energy barrier for transitioning to an inward-facing state. The deep mutational scan for SERT-catalyzed import of APP+ finds mutations that promote the necessary conformational changes that would otherwise be facilitated by the native substrate. Indeed, hundreds of gain-of-function mutations for APP+ import are found along the permeation pathway, most notably mutations that favor the formation of a solvent-exposed intracellular vestibule. The mutagenesis data support the simulated mechanism in which the neurotransmitter and a symported sodium share a common cytosolic exit pathway to achieve coupling. Furthermore, the mutational landscape for SERT surface expression, which likely filters out misfolded sequences, reveals that residues along the permeation pathway are mutationally tolerant, providing plausible evolutionary pathways for changes in transporter properties while maintaining folded structure.

Keywords: deep mutagenesis, transporter, SERT, alternating access mechanism, molecular dynamics, conformational free energy landscape

INTRODUCTION

The serotonin transporter (SERT/SLC6A4) catalyzes the reuptake of serotonin from the synapse to terminate synaptic transmission (*1*, *2*). As evidenced by the effects of pharmaceuticals and illicit drugs that inhibit SERT, and genetic studies of knockout mice (*3*) and natural human variants (*4–7*), dysregulation of serotonergic signaling is associated with broad psychopathological symptoms, including major depression, anxiety, obsessive compulsive disorder, post-traumatic stress disorder, and attention-deficit/hyperactivity disorder, amongst others (*8*). Similar to other closely related monoamine neurotransmitter transporters for dopamine (DAT/SLC6A3) (*9*, *10*) and norepinephrine (NET/SLC6A2) (*11*), SERT mediated serotonin (5-hydroxytryptamine; 5-HT) translocation is coupled to the favorable dissipation of a Na^+ electrochemical gradient. While the stoichiometry for coupled transport remains unclear (*12*), it most likely involves the Cl^--dependent import of one Na^+ and one positively charged 5-HT into the cell by SERT (*13–15*) using an alternating access mechanism (*16*, *17*), in which the transporter in an outward-facing (OF) conformation captures extracellular substrate, transitions through an occluded (OC) state where the central substrate binding sites become closed and inaccessible to the outside, and then an intracellular exit pathway opens in an inward-facing (IF) conformation for cytosolic release (*18*, *19*). SERT then returns to an outward-facing conformation through the antiport of one K^+, together with possible movement of a proton, for an overall electroneutral cycle (*12*, *13*). Alternative stoichiometries and even channel-like conduction states may occur under different conditions (*15*, *20–23*).

Monoamine transporters and other neurotransmitter:sodium symporters (NSS) share a common architecture (*24*) known as the LeuT fold, named after the bacterial LeuT transporter, which was the first member structurally characterized at atomic resolution (*25*). Static crystal

structures of human SERT and *Drosophila* DAT have revealed atomic details of OF and OC

conformations bound to drugs and substrates (*26–31*). These structures have resolved two Na^+

sites (Na1 and Na2) and the location of Cl^- binding, as well as specific contacts to the

neurotransmitter aromatic ring and amine moieties within subsites (*32*) of the central orthosteric

(or S1) binding site. In addition, structural, biochemical and pharmacological studies have shown

that substrates entering the outward-facing extracellular vestibule engage an allosteric (or S2)

site immediately adjacent to critical gating residues, before moving past the open gate to the

orthosteric site (*26*, *33–35*). Cryo-EM analysis of detergent-solubilized SERT in the presence of

ibogaine, a psychoactive plant product that stabilizes IF conformations, allowed for the

characterization of OF, OC and IF-like states at moderate resolution (*19*), showing that regions

already implicated in bacterial and mammalian NSS homologues through multiple methods (*18*,

36–41) move to open a solvent-accessible intracellular vestibule. In particular, transmembrane

helix (TM) 1a moves away from the helical bundle. These motions are supported by

comprehensive HDX-MS analysis of LeuT (*42*) and SERT (*43*). However, the overall molecular

mechanism for ion-coupled neurotransmitter transport remains incompletely understood. Key

questions that remain unresolved are how stoichiometry is achieved and how does the

neurotransmitter move from the orthosteric site to the cytosol. Cryo-EM structures of ibogaine-

bound SERT only confirmed the general region for neurotransmitter exit, and the authors were

uncertain as to the exact exit pathway amongst multiple possibilities (*19*). Understanding the full

mechanism of transport in atomic detail will assist in understanding how genetic variants are

associated with disease and may aid the development of drugs targeting discrete conformational

states.

We recently described molecular dynamics (MD) simulations of the entire SERT-

catalyzed import process of 5-HT (*44*), accomplished using adaptive sampling to efficiently

explore conformational space. The calculated highest flux pathway for substrate binding events

and conformational changes lead to several important conclusions, including the identification of

key residues involved in substrate recognition at the extracellular vestibule and the prediction

that the neurotransmitter exits along a pathway beginning at the Na2 site that is surrounded by

TM1a, TM5, TM6b, and TM8.

Here, we use deep mutational scanning (*45*) to define the mutational landscape of SERT

for surface expression, which indicates how mutations effect folding and escape from

intracellular protein quality control (*46*), and substrate transport. Deep mutagenesis involves (i)

creating a library of protein variants, (ii) using *in vitro* selection to enrich for the variants with

high activity, and (iii) measuring the enrichment or depletion of variants in the mutagenesis

library using next generation sequencing. Deep mutational scanning has proven insightful for

indirectly interrogating protein conformational states in living cells (*46–49*), and datasets that

include higher order mutations can even assist in modeling small protein structures with atomic

resolution (*50*, *51*). We build on this body of work by now applying deep mutagenesis to a

human transporter that undergoes substantial conformational change. By using a non-native

substrate that is unable to facilitate isomerization of SERT to an inwards-facing conformation

when compared with the native substrate, we find extensive mutational information that provide

insight on dynamic regions in close agreement with simulation.

RESULTS

Mutational landscapes of SERT for plasma membrane localization and substrate import

We sought to understand how SERT sequence relates to conformational dynamics and

mechanism using deep mutagenesis, which demands an effective selection strategy to distinguish

low and high activity SERT sequence variants. This is not possible using transport assays with

radioactive 5-HT. Instead, *in vitro* selection was based on the uptake of the fluorescent substrate

and monoamine neurotransmitter analogue APP+ (4-(4-dimethylamino)phenyl-1-

methylpyridinium) (Figure S1), permitting cells expressing active and inactive SERT variants to

be readily distinguished and separated by fluorescence. We used a fluorescence-based transport

assay to assess SERT activity by flow cytometry, which could then be seamlessly integrated with

fluorescence-activated cell sorting (FACS) to screen libraries of SERT variants. To detect

surface localized transporter, a synthetic gene encoding human SERT with a c-myc epitope tag

(flanked by gly/ser-rich linkers) inserted between residues Asp216 and Asn217 of EL2 was

constructed; the homologous site in *Drosophila* DAT features a large domain insertion. When

expressed from an episomal plasmid in Expi293F cells (a suspension culture derivative of human

HEK293 that has advantages for FACS-based selections of large libraries), surface localized

transporter was detected with a fluorescent anti-myc antibody, while uptake of APP+ was

simultaneously observed (Figures S2A and S2B). Validation of our construct showed that myc-

tagged SERT has ion dependence consistent with untagged human SERT (*14*, *52–54*) (Figure

7E), appropriate neurotransmitter selectivity (Figure 7D), and exhibits a similar K_M for APP+ as

for the endogenous substrate 5-HT (Figure 7E and Table 1). However, APP+ and 5-HT transport

velocity of myc-tagged SERT are reduced 4-fold and 2-fold compared to the native protein,

respectively, while insertion of a shorter gly/ser-rich sequence at the same site had no significant

effect (Figure S2C and Table 1). We speculate that the bulkier myc tag with a longer gly/ser-rich

linker (a 24 residue insertion in total) may partially obstruct substrate access to the extracellular

vestibule, a limitation that we considered offset by the capability of measuring transport and

surface localization simultaneously. Furthermore, a subset of mutations displaying enhanced

APP+ transport discovered through deep mutagenesis were confirmed to also enhance activity in

native untagged SERT (Figure S2D), and the presence of the myc tag is therefore very unlikely

to have impacted scientific conclusions.

Tagged SERT was mutagenized to generate nearly all single amino acid substitutions. To

increase sampling during FACS-based selection, substitutions were split across two libraries

spanning the N- (residues 2-325) and C- (residues 326-630) termini, each library containing

~6,000 mutants. The libraries were diluted with carrier DNA encoding a viral factor for

enhanced replication of the episomal plasmid (*46*) and transfected into Expi293F cells. A large

excess of carrier DNA ensures that typically only one SERT coding variant is expressed in any

cell (*48*), providing a tight link between genotype and phenotype. Cell libraries were incubated

with APP+ at a concentration below the K_M and stained with anti-myc antibody conjugated to an

orthogonal fluorochrome to simultaneously assess surface expression. Cells expressing SERT at

the plasma membrane, or expressing SERT and transporting the fluorescent substrate, were

collected via FACS (Figures S2E and S2F). After deep sequencing, mutation enrichment ratios

were calculated by comparing variant frequencies from RNA transcripts in the sorted cells to the

naive DNA library. These data experimentally define the mutational landscapes for SERT

surface expression and APP+ transport (Figure 1).

Figure 1. SERT mutational landscapes for surface expression and APP+ transport.
(A, B) SSM libraries of SERT spanning N- and C-terminal residues (the red diamond in the secondary structure schematic at top indicates the dividing point between the libraries and the grey triangle at positions 216-217 indicates the placement of the myc tag) were sorted for (A) surface expression and (B) cell uptake of APP+. Log$_2$ enrichment ratios are plotted from $\leq$ -3

(depleted mutations, orange) to 0 (neutral, white) to $\geq$ +4 (enriched, dark blue). Data are
averaged from two independent experiments. Missing mutations ($<$ 10 reads in the naïve
libraries) are grey. *, stop codons.
(C-F) Agreement between $\log_2$ enrichment ratios from replicated selections for surface
expression of the SERT (C) N-terminal and (D) C-terminal SSM libraries, or for APP+ transport
of SERT (E) N-terminal and (F) C-terminal SSM libraries. Agreement between independent
experiments generally increases for more frequent sequence variants (orange, 10-99 reads in the
naïve library; green, 100-199 reads; light blue, 200-399 reads; dark blue, 400+ reads). R^2 values
are correspondingly colored in the upper-left corners.
(G-J) Agreement between conservation scores (defined as the mean of the $\log_2$ enrichment ratios
at a single residue position) from replicated selections for surface expression of SERT (G) N-
terminal and (H) C-terminal SSM libraries, or APP+ transport of (I) N-terminal and (J) C-
terminal libraries.

Enrichment ratios are correlated between independently replicated selection experiments

(Figures 1C-1F). Agreement between replicate experiments tends to increase for highly abundant

sequence variants, presumably due to increased sampling. Metrics for mutational tolerance such

as sequence Shannon entropy do not distinguish between positions that tolerate many neutral

mutations versus positions that are hot spots for gain-of-function mutations. We therefore

calculate a simple, unambiguous average of the $\log_2$ enrichment ratios for all 20 amino acid

possibilities at each position, which we call a residue conservation score. Conservation scores are

highly correlated between replicates (Figures 1G-1J) and define regions of sequence that are

tightly conserved for function, or alternatively are under positive selection to change.

Mutational scans of transmembrane proteins for plasma membrane localization indicate

whether mutations adversely impact folding and structural stability, as folded proteins are

anticipated to preferentially escape intracellular quality control machinery (*46*). In the mutational

landscapes for both SERT surface expression and APP+ transport, mutations within secondary

structural elements are generally deleterious, especially polar mutations within transmembrane

segments (Figures 1A and 1B). In mutational scans of G protein-coupled receptors (*48*), proline

substitutions, which are disruptive to helical conformations, were depleted in regions (including

termini) that have ordered structure, yet no such depletion of proline substitutions is observed in

the cytosolic termini of SERT. Instead, the SERT cytosolic tails (N-terminal residues 1-75 and

C-terminal residues 614-630 beyond the C-terminal helix immediately following TM12) tolerate

most mutations, strongly indicating that these regions lack folded structure based on expression

and substrate import alone, despite previous modeling to the contrary (*55*). However, we cannot

exclude that these regions may have folded structure pertinent to regulatory mechanisms, such as

structural elements induced by protein-protein interactions or post-translational modifications,

that do not participate in our cell-based selection. In comparison, mapping conservation scores

for each residue position to the crystal structure of SERT (*26*, *27*) highlights regions of folded

structure under sequence constraints (Figure 2). Residues considered important for SERT

trafficking, namely disulfide bonded cysteines (*56*) and a putative cholesterol interaction site

(*57*), are moderately to highly conserved for high surface expression, whereas N-glycosylation

site N208 is partially conserved and N217 is not (Figure S3). Moreover, co-evolved residues also

tend to be well conserved in the mutational scan for surface expression, highlighting

evolutionary constraints to maintain proper transporter folding and stability (Figure S4). The

selection of SERT mutants for surface expression reveals conservation within packed

hydrophobic cores 'above' and 'below' the central orthosteric binding site (Figure 2B), while the

protein sequence along the permeation pathway is tolerant of many substitutions. This suggests

mutations in the permeation pathway, which may be relevant in the evolution of new transporter

specificities and coupled ion stoichiometries, may frequently occur without major disruption of

the basic protein architecture and fold. Finally, contrary to predicted free energy values for

membrane helix insertion (*58*), transmembrane helices surrounding the periphery of the

transporter are only moderately conserved for expression, while helices in the core are highly

conserved (Figure S5). The effects of mutations on SERT surface expression are therefore most

closely related to the protein's folded structure and not to the bioenergetics for membrane

insertion of individual helices.

In the deep mutational scan for APP+ import, the SERT sequence is more conserved than

was observed for surface expression alone, with sequence conservation expanded from core

regions important for folding towards residues lining the entrance vestibule, the neurotransmitter

and ion binding sites, and sites of structural rearrangement near the orthosteric binding site

(Figure 2C and 3). For example, a hydrogen bond interaction between Tyr176 and Asp98, which

lies 'above' the orthosteric binding site, is proposed to serve as a key molecular switch for

conformational transitions (*59*). Most mutations to either residue are deleterious for substrate

import in the deep mutational scan, which concurs with previous mutational studies of Tyr176 in

SERT (*60*), as well as equivalent mutations in DAT and LeuT that exhibit decreased transport

activity of their respective substrates (*59*, *61*). Furthermore, the hydrogen bonding network

between Glu136, Gly340, and Glu508 (located at the central break in TM6 and forming part of

the orthosteric binding site) is critical for mediating the coupling of outward and inward facing

states (*62–65*). Previous computational modeling has suggested that the buried Glu136 and

Glu508 coordinate a single proton (*66*), while Glu136 also interacts with the Gly340 backbone of

TM6. Nearly all mutations to these residues are deleterious for transport function in the deep

mutational scan.

Figure 2. SERT mutations that increase APP+ import are enriched in the intracellular exit pathway.

(**A**) Overlaid MD snapshots of APP+ translocation, from when APP+ enters the extracellular vestibule (blue) to its cytosolic exit (red). The positions of ions in the OC state are shown with spheres. SERT is shown as a semi-transparent ribbon and colored by experimental sequence conservation for APP+ transport, showing regions where mutations are depleted in orange to regions where mutations are enriched in dark blue. Also see structure file in Supporting Information online.

(**B**) A cross-section through SERT in a partial OF state (PDB 5I73; bound to (*S*)-citalopram in magenta) colored by conservation for folding and surface expression. Tightly conserved residues in the core are orange.

(**C**) Cross-sections through SERT in a partial OF state at left and in an ibogaine-bound IF-like state at right (PDB 6DZZ), highlighting in orange residues within the substrate-binding sites and extracellular cavity that become conserved for APP+ transport, while residues that are hot spots for gain-of-transport mutations are dark blue and map to the intracellular exit pathway. Drugs are magenta.

(**D, E**) Enrichment ratios for APP+ transport were converted to amino acid probabilities, weighted by hydropathy (yellow, hydrophobic; blue, polar), and averages are mapped onto the (**D**) surface and (**E**) core of SERT (PDB 5I73). Residues lining the entrance cavity and putative exit pathway tolerate polar substitutions.

(**F**) Natural conservation based on a sequence alignment of 150 SERT homologs (blue, conserved; to red, variable). Core, entrance, and exit pathways are conserved in natural history.

The mutational scan identifies hundreds of gain-of-function mutations based on positive

enrichment after sorting for APP+ import. We further tested by targeted mutagenesis 22

predicted gain-of-function mutations that are spatially dispersed across the structure and

validated 21 as trending toward increased APP+ uptake, with 11 having statistical significance

(Figure S6). Gain-of-function phenotypes persist when the mutations are introduced into the

untagged native cDNA (Figure S2D). We also tested previously reported allelic variants G56A,

I425V, and K605N, which have increased 5-HT transport due to altered PKG signaling (*6*, *7*, *67*,

68), and found these did not have highly elevated APP+ uptake when expressed individually,

consistent with their lack of positive enrichment in the mutational scan (Figures S2D and S6A).

Cyclic GMP-dependent signaling is therefore unlikely to be an important pathway for

modulating SERT activity in the experimental system. Expression of the mutants was generally

similar to or less than wild type and cannot explain the large increases in transport activity

(Figures S6B and S6C).

261
Targeted mutagenesis therefore validates predictions from the mutational scan that many amino

acid substitutions increase APP+ transport kinetics. Futher validation of the strength of this

approach is found in agreement between identified mutants and previous focused studies

characterizing the same residues. For instance, our previous study investigating the molecular

determinants necessary for transport of 3,4 methylendioxymethamphetamine (MDMA) and 5-HT

by SERT (*69*) show that outer gate residue Arg104 is intolerant of substitutions (as also observed

in our mutational scan for APP+ import, in which only substitution R104K had a $\log_2$ enrichment

value > -1) and that only acidic residues at Glu493 are permissive for MDMA transport but other

substitutions of Glu493 are tolerated for 5-HT (*69*) and APP$^+$ transport based on our findings

here. Furthermore, residues Ile179 and Asp400 are highly sensitive to substitution in regards to

APP$^+$ transport, V489 can only tolerate substitution by hydrophobics Leu, Met, Ile, and Phe, and

K490 can accept multiple substitutions, which is consistent with the findings of Anderson et al.

(*60*).

Figure 3. The Na1 site, Cl$^-$ binding site, and sites of structural rearrangement adjacent to
the orthosteric binding site are conserved for transport function.
Heat maps (colored as in Figure 1) show depleted mutations for SERT surface expression (left)
or APP+ transport (right) in orange and enriched mutations in dark blue. Accompanying
snapshots show SERT (PDB: 5I73) as a cartoon representation with residues colored according
to conservation score for transport (as described in Figure 2A).
(**A**) The Na1 site.
(**B**) The Na2 site.
(**C**) The Cl$^-$ ion site (modeled based on structural alignment with PDB 7LIA).
(**D**) The hydrogen bonding interaction between Tyr176 and Asp98.
(**E**) The hydrogen bonding network between Glu136, Gly340, and Glu508. Based on previous
computational modeling, Glu136 and Glu508 are predicted to coordinate a proton between the
two residues (*66*). For this study, Glu136 was modeled in the protonated form in MD
simulations.
291
Conformational transition to the inward-facing state is rate-limiting for APP+ import

Strikingly, the bulk of mutations that enhance APP+ uptake are clustered in the intracellular half

of the transporter (Figure 2C). This clustering of gain-of-function mutations is most noticeable at

TM1a, TM6b, and the intracellular segment of TM8, loosely focused near the axis of

pseudosymmetry and tightly aligned to the simulated neurotransmitter exit pathway (*70*). After

weighting the enrichment ratios by hydropathy and averaging, we find that this region of SERT

is highly tolerant of diverse mutations including polar substitutions (Figures 2D and 2E).

Substituted cysteine accessability analysis of TMs 1 and 6 by us and others have found 5-HT

transport is largely unaffected by cysteine substitutions in intracellular sections of the TMs and

accessibility of some of the cysteine substitutions to modifying agents, consistent with their

exposure in the transport pathway (*71*, *72*). We propose mutations tolerated in this region will

disrupt buried interactions along the exit pathway and thereby facilitate opening of a solvent-

exposed intracellular vestibule. That these mutations enhance APP+ uptake is exactly what one

would expect if formation of the IF state is rate-limiting.

As previously reported (*44*), the entire Na^+-coupled import process of the physiological

substrate 5-HT has been simulated using Markov state model (MSM) based adaptive sampling

(*73–75*). MD with adaptive sampling has been successfully used to simulate complex transport

processes in other systems (*74*, *76*). MSM-weighted SERT simulation data are projected onto a

coordinate system defined by distances between extracellular and intracellular gating residues

(*38*, *44*), thereby visually displaying the relative free energies and barriers between kinetically

relevant intermediate states (Figure 4). SERT free of neurotransmitter substrate (i.e. Na^+-SERT)

readily transitions between OF and OC states (Figure 4A) and only in the presence of the cognate

substrate 5-HT does the IF state become substantially occupied (Figure 4B), with a reduction in

the free energy barrier for the OC-IF transition. Here, the simulations are replicated, but with 5-

HT replaced by APP+ in the simulation box as the substrate. Na$^+$-bound SERT in an OF

conformation obtained from Na$^+$-SERT simulations (*44*) was used to seed MD simulations of

APP+ import totaling ~370 μs (referred to as APP-SERT simulations; Figures 4C and S4).

Figure 4. Simulated conformational free energy landscapes of SERT.
(**A, B**) Relative free energies from MSM-weighted simulation data plotted against the distances
between extracellular and intracellular gating residues for (A) Na$^+$-SERT and (B) 5-HT-SERT.
These simulation data were previously reported (*44*) and are reproduced here for comparison.
The outward-facing (OF) crystal structure (PDB 5I73) was used as the starting structure for MD
simulation and transitioned to occluded (OC) and inward-facing (IF) states.
(**C**) Calculated free energy landscape of SERT during the APP+ translocation process.
(**D**) MD snapshot of membrane-embedded SERT with APP+ bound in subsite B and ions bound
in respective sites. Extracellular gating residues Arg104 and Glu493 shown in orange sticks.
Intracellular distance was measured based on distances between TM1a, TM6b and the
cytoplasmic base of TM5 and TM8, colored in teal, magenta, green, and yellow, respectively.

The simulations reveal that the free energy basin corresponding to IF conformations

remains at a higher free energy value in the presence of APP+ (Figure 3C), resembling the

conformational free energy landscape for Na^+-SERT (Figure 3A). The free energy barriers for

OC-IF transitions in Na^+-SERT and APP-SERT are calculated to be 3.43 ± 0.06 kcal/mol and

2.89 ± 0.08 kcal/mol, respectively, which are higher than the estimated 1.66 ± 0.04 kcal/mol

barrier for the OC to IF transition when 5-HT is bound and permeating along the transport

pathway. Indeed, formation of the IF state is predicted to be rate-limiting for APP+ import,

whereas 5-HT interactions facilitate dynamic motions necessary for transport by reducing free

energy barriers and stabilizing the IF state (compare Figures 4B and 4C). Similar conclusions

regarding differences between substrates have been drawn from experimental observations of

bacterial LeuT; only a suitable LeuT substrate such as alanine readily facilitates transition from

the OC to IF state (*38, 77*). The conformational free energy landscape of APP+ import also

shows OF states to be further stabilized as the bulkier substrate occupies the extracellular

vestibule and the allosteric site. Partial OF-IF-like conformations are also observed, which we

previously referred to as an hourglass (HG) state where both gates are open but there is a central

constriction (*44*). We speculate that HG conformations may be associated with transient water-

conducting states observed in simulations of other membrane transporters (*78, 79*).

Natural evolution can be considered akin to a deep mutagenesis experiment for protein

function as it relates to organism fitness, and we note that residues of SLC6 family members

along the permeation pathway are conserved between species (Figure 2F). Furthermore, an

exome database shows little human variation (*80*). SERT has therefore reached an optimum (or

close to optimum) sequence for its physiological function of ion-coupled 5-HT transport within

the constraints of natural evolution. In contrast, our deep mutational scan identifies mutations

that increase import of the unnatural substrate APP+ for which the protein sequence is

suboptimal. In essence, mutations are found that impose on the APP-SERT conformational free

energy landscape the necessary changes that would ordinarily be facilitated by interactions with

the native substrate 5-HT.

Based on crystal structures and computational modeling, the N-terminal cytosolic tail of

LeuT, in particular LeuT residue Arg5 (equivalent to Arg79 of SERT), forms an interaction

network that stabilizes a closed conformation of the cytoplasmic exit cavity in the OF and OC

states (*25*, *38*, *41*). The simulations show dynamic motions of the equivalent residues in SERT

(a.a. 78-82), with Arg79 forming transient salt bridges to Glu444 and Asp452 to restrict opening

of the cytoplasmic exit. Consistent with transitions to the IF state being rate-limiting, mutations

to Arg79 and surrounding cytoplasmic gating residues increase APP+ import by destabilizing

closure of the intracellular vestibule. We note that alanine substitution of Arg79 (*81*) is

deleterious for SERT-catalyzed transport of 5-HT and likely imbalances the conformational

equilibrium. Such mutations introduced in the gating residues of GABA transporter GAT-1 have

also been shown to alter the equilibria of states and promote inward-facing conformations (*82*).

Likewise, mutations in the Na2 site, which are discussed further below, also increase APP+

import but reduce 5-HT transport (*52*). It is only for APP+, which is unable to facilitate

appropriate conformational dynamics, that these mutations impart a gain-of-function phenotype,

providing a unique mutational data set that functionally maps the substrate exit pathway.

Deep mutagenesis supports the simulated binding mechanism of Na$^+$ and neurotransmitter

The highest flux pathway for ion/APP+ binding events and conformational changes was

determined by transition path theory (*83*) and predicts an ordered process for APP+ import

identical to that previously simulated for 5-HT (Figure S7). Na$^+$ first binds an *apo*-SERT at the

Na1 site, followed by a second Na^+ occupying the Na2 site. APP+ then enters the transporter

through the extracellular vestibule and moves to the orthosteric site, with localized dynamics

around extracellular loop 4 (EL4) and the extracellular gate, consistent with static structures (*18*,

19), HDX data (*42*, *43*), and voltage-clamp fluorometry (*84*). Unlike the endogenous substrate 5-

HT, the increased structural rigidity of APP+ (Figure S1) results in its translocation 'down' the

extracellular vestibule to be more constrained, with decreased rotational freedom. In the

mutational landscape, hydrophobic substitutions are tolerated or weakly enriched at positions

Phe335 and Glu494, where they pack against the APP+ aromatic ring moieties (Figure 5A).

Cl^-, accompanied by an extracellular Na^+ ion, then engages the extracellular gating

residues Arg104 and Glu493, disrupting their electrostatic interactions to allow Cl^- to enter the

orthosteric cavity. The additional Na^+ ion does not further progress towards the orthosteric cavity

and diffuses back to the extracellular space. Once the neurotransmitter and ions are all bound to

their respective sites, the extracellular gates close to form OC conformations. The stability of the

OC state promotes the solvation of the intracellular vestibule, thereby weakening the hydrogen

bonding network involving various intracellular gating residues (Arg79-Glu452, Glu80-Lys275,

Asp87-Trp282, Tyr350-Glu444). Opening of the intracellular permeation pathway is associated

with displacement of Na^+ from the Na2 site into the cytoplasm (consistent with extensive

experimental evidence (*36*, *40*, *85*)), followed by cytosolic exit of APP+.

Figure 5. The simulated mechanism for SERT-catalyzed substrate import is supported by the mutational landscape.

Heat maps (colored as in Figure 1) show depleted mutations for SERT surface expression (left) or APP+ transport (right) in orange and enriched mutations in dark blue. Accompanying MD snapshots show SERT as a cartoon representation with residues colored according to conservation score for transport (as described in Figure 2A).
(**A**) APP+ entering the transporter via the allosteric site.
(**B**) The neurotransmitter substrate bound to the orthosteric site, straddling subsites A and B.
(**C**) APP+ exiting the orthosteric site through the Na2 site.
(**D**) APP+ permeation along the exit pathway.

Simulations show contacts made by the neurotransmitter within specific subsites (*32*) inside the orthosteric binding pocket change as import progresses. The aromatic ring of 5-HT shifts from subsite C to B, while flexibility in the aliphatic chain accommodates the amine moiety remaining bound to subsite A (*44*). APP+ never fully straddles subsites A to C, most likely due to steric limitations from having a rigid structure, yet it does interact with subsite B in a binding mode supported by crystallography of dopamine-bound DAT (*30*) and biochemical analyses of inhibitor binding (*32*, *86*, *87*). While residues surrounding the subsites are generally conserved for APP+ import, A169D within subsite B enhances transport (Figure 5B). This mutation is known to also increase the potency of inhibitors (*86*). A169D introduces a

carboxylate ~12 Å distal to where the substrate's amine group is positioned at subsite A. This

spacing is ideal for accommodating the length of APP+, which through resonance has positive

charge distributed on both ends for favorable electrostatics in the A169D mutant; this provides

further evidence that monoamine substrates are indeed accommodated in subsite B during the

transport process. In contrast, Asp in the 169 position has been shown to decrease paroxetine

potency via the electrostatic repulsion of the fluorophenyl moiety (*88*).

In the deep mutational scan, mutations to residues coordinating ions in the Cl$^-$ and Na1

binding sites are highly deleterious for transport activity (Figure 3A and 3C) and further supports

the specific ion dependence for proper function (*52, 85, 89*). In contrast, mutations to the Na2

site increase APP+ import, likely due to weakening the interactions between the gating helix

TM1 and the scaffold TM8 that would stabilize outward-facing conformations (Figure 3B).

Similar mutations introduced to the Na2 site in LeuT biases the conformational equilibria

towards the inward-facing state and is demonstrated by its crystal structure (*18, 85*). In SERT,

Na2 site mutations reduce 5-HT binding and transport and underscore the importance of

maintaining proper substrate-ion coupling (*52, 90*).

Gain-of-function mutations that enhance APP+ uptake begin with Leu343 and Asp437 in

the Na2 site and continue along the entire exit pathway to the cytoplasm (Figures 3B, 5C, and

5D). There is a near perfect alignment of gain-of-function mutations for APP+ import on TM1a,

TM5, TM6b, and TM8 with the simulated permeation trajectory (Figure 4D and structure

visualization file in Supporting Information). The Na2 site therefore forms the apex for the

cluster of the gain-of-function mutations that delineate the intracellular vestibule. The data are

consistent with the simulated release of Na$^+$ from the Na2 site, followed by the movement of 5-

HT or APP+ into the exit pathway by first passing through the vacated Na2 site. The sharing of a

common exit pathway by neurotransmitter and Na^+ beginning at the Na2 site elegantly explains

the 1:1 sodium to neurotransmitter stoichiometry of the transport cycle. Tyr95 forms cation-pi

and pi-pi interactions with the substrate and is highly conserved in the mutational scan, tolerating

only aromatic substitutions (Figure 5B). The release of 5-HT from the orthosteric site requires a

rotameric shift of Tyr95 that is coupled to Na^+ dissociation at the Na2 site (*44*, *91*). Compared to

5-HT, the structural rigidity of APP+ does not enable Tyr95 to freely rotate and the conformation

of Tyr95 is less closely coupled to the release of the substrate (Table S1). Whereas both Na1 and

Na2 are important in propagating conformational changes (*66*, *92–95*), the hypothesis that Na^+ in

the Na2 site but not the Na1 site is imported during translocation is further strengthened when

considering that members from the SLC5 (eg. galactose transporter) and SLC6 (eg. SERT)

families lack sequence homology but share almost superimposable core structures with

structurally conserved Na2 sites, whereas only SLC6 members have an Na1 site (*96*) that has

been shown to be important for substrate coordination and propagation of conformational

changes for translocation (*94*). These distinct roles for Na1 and Na2 are supported by our

previous mutagenesis analysis of the Na1 and Na2 sites, where some Na1 site mutants but not

Na2 site mutants allowed Ca^{2+} to functionally substitute for Na^+ in 5-HT transport.

Electrophysiological reversal potentials revealed that Ca^{2+} was not transported. These findings

along with molecular dynamic simulations supported Ca^{2+} bound to Na1, while not transported,

contributes to the coordinated conformational changes for 5-HT transport (*94*).

Gain-of-function mutations for APP+ uptake at positions outside the permeation pathway are

predicted to enhance SERT conformational dynamics.

We clustered the SERT mutational data for surface expression and APP+ transport

activity using an unsupervised learning algorithm, Uniform Manifold Approximation and

Projection (UMAP) (*97*). UMAP presents an approach to effectively reduce the high dimensional

mutational landscape and systematically derive quantitative relationships between a residue's

mutational response and its biophysical and biochemical properties (*98*, *99*). Visualization of the

mutational landscape on the two-dimensional UMAP space shows that residues cluster based on

various biochemical and biophysical properties (Figures 6 and S5).

Projection of the mutational landscape for APP+ transport shows that residues naturally

cluster based on residue site location (Figure 6B) and hydrophobicity (Figure 6C). When

compared to biophysical properties obtained from MD simulations, we observed that a residue's

conservation score in the deep mutational scan is related to its dynamic fluctuations (Figure 6D),

solvent accessible surface area (SASA) (Figure 6E), and local structural packing (Figure 6F).

Residues that enhance transport activity are associated with larger fluctuations, while residues

with low mutational tolerance exhibit low SASA and RMSF, suggesting these residues are

buried in the protein core and are important for stabilizing contacts to maintain proper folding.

Furthermore, the deep mutational landscape for SERT expression correlates with mutational

effects predicted by EVmutation (*100*) (Figure S8G), DeepSequence (*101*) (Figure S8H), and

FoldX (*102*) (Figure S8I), in addition to correlating with local structural packing (Figure S8F),

underscoring that sequence and structural constraints are necessary for proper expression and

trafficking to the plasma membrane (Table S1). By comparison, the mutational landscape for

APP+ transport is less correlated with the variant effect predictions, further emphasizing that

important protein features are only partially shared between the transport mechanisms for the

endogenous and nonendogenous substrates.

Figure 6. UMAP projection of the mutational landscape for SERT-catalyzed APP+ transport.
Each residue was clustered using the UMAP clustering algorithm based on the activities of all amino acid substitutions at that position, and are colored based on different biochemical and biophysical properties: (**A**) APP+ uptake enrichment ratio, (**B**) residue location, (**C**) Kyte-Doolittle hydrophobicity, (**D**) root mean square fluctuation (RMSF) (**E**) solvent accessible surface area (SASA), and (**F**) average number of contacts. The mutational landscape was also compared to scores from variant effect predictors (**G**) EVmutation, (**H**) DeepSequence, and (**I**) FoldX. Residues in which data is unavailable are colored gray. The first UMAP dimension strongly correlates with enrichment ratio (transport ρ = 0.966). The properties from MD simulations (RMSF, SASA, and average number of contacts) were calculated on 50,000 structures drawn from an MSM-weighted distribution. Projection of the SERT expression deep mutational landscape in 2D UMAP space is available in Supplemental Information Figure S8. Correlations between biochemical and biophysical properties with the UMAP dimensions and DMS enrichment ratios are listed in Table S2.

While less abundant, some mutations outside the simulated permeation pathway are also enriched for APP+ import, including mutations at helical hinges that flex during structural transitions (e.g. A173G, P499K, S585P), or on neighboring helices immediately adjacent (e.g. F263C, P499K, G582K); these substitutions may stabilize unwound structural features within the membrane phase, facilitate dynamic motion, alter packing against surrounding helices, or stabilize conformational states. The mutational tolerance of Ala173 at a helix kink is particularly striking; nearly all substitutions are deleterious for transport except for the small amino acid glycine that can increase flexibility (Figure 7B). DAT and NET both have glycine at this position. We also find a cluster of gain-of-function mutations for APP+ import within the hydrophobic core where TM1b packs against EL4 (Figure 7C); substantial EL4 motion in bacterial transporters can occlude the extracellular vestibule (*103*), while more limited EL4/TM1b motions are observed in cryo-EM structures of ibogaine-bound SERT (*19*) and in the simulations. In general, mutations in this region that increase APP+ import substitute large hydrophobics for smaller side chains that reduce packing and are therefore anticipated to increase dynamics, possibly creating localized molten globule-like behavior.

We further investigated four SERT mutations that increase APP+ import, chosen to represent different structural regions: R79V in the cytoplasmic gating loop (Figure 7A), F88H on TM1a proximal to the exit pathway (Figure 7A), A173G at a helix kink in TM3 (Figure 6B), and L406I in EL4b where it packs against TM1b (Figure 7C). All mutants retained selectivity for 5-HT over dopamine and norepinephrine (Figure 7D). Na^+ and Cl^- dependency of APP+ import is partially lost in two of the mutants, R79V and F88H (Figure 7E), which both lie on the exit pathway and are predicted to destabilize closure of the intracellular vestibule. Correct ion coupling for symport therefore demands appropriate free energy differences between

523 conformational states, ensuring that all ions and substrate must bind to energize structural

524 transitions.

Figure 7. Exit pathway mutations can cause a partial loss of Na$^+$ and Cl$^-$ dependency.
(**A-C**) Superimposed MD snapshots (blue, OF state; green, OC state; and pale red, IF state) showing dynamic regions. Experimental enrichment ratios for select residues are plotted alongside, colored orange (deleterious mutations) to dark blue (enriched mutations) as in Figure 1.
(**A**) Motion around the intracellular gate.
(**B**) A hinge in TM3.
(**C**) The hydrophobic core below EL4, adjacent to the extracellular gate.
(**D**) APP+ uptake in SERT-expressing cells was more potently inhibited by 5-HT (grey) than by norepinephrine (NE, green) or dopamine (DA, blue). (Mean; WT n = 5, R79V n =2, F88H n =2, A173G n = 3, L406I n = 2. For WT, error bars show SD.)
(**E**) APP+ uptake by SERT-expressing cells was measured in Ringer's solution (black), with replacement of Na$^+$ (red) or Cl$^-$ (grey) ions. Transport was Na$^+$ and Cl$^-$ dependent for WT, A173G, and L406I SERT, while showing a partial loss of ion dependence in R79V and F88H mutants. (WT n = 4, R79V n = 4, F88H n = 3, A173G n = 4, L406I n = 3, mean ± SD)

To further support the conclusion that SERT gain-of-function mutations for APP+ import

in the deep mutational scan are substrate specific, we evaluated the set of four SERT mutants for

transport of tritiated 5-HT in transfected HEK-293 cells (Table 1). SERT A173G and L406I

have similar kinetic parameters for 5-HT transport as the wild type, albeit with slightly reduced

maximum velocity (V_{max}). However, SERT mutants R79V and F88H have large decreases in

V_{max} and K_M, possibly due to loss of interactions with 5-HT in the intracellular exit pathway that

would ordinarily facilitate translocation out of the orthosteric site and down a narrow

intracellular vestibule toward bulk solvent (*44*). Others have also found that mutations that favor

IF conformations lower both V_{max} and K_M (*57*).

Table 1. 5-HT transport: K_M and V_{max} of SERT and mutants.			
Transporter	K_M (μM) [1]	V_{max} (amol cell^{-1} min^{-1})	n
native SERT	2.1 ± 0.7**	0.66 ± 0.15	3
opt myc-SERT [2]	1.3 ± 0.6	0.48 ± 0.15	4
R79V	0.25 ± 0.07*	0.05 ± 0.01**	3
F88H	0.14 ± 0.05*	0.07 ± 0.04*	3
N173G	1.2 ± 0.5	0.38 ± 0.13	4
L406I	1.1 ± 0.5	0.23 ± 0.01	4

[1] Kinetic parameters are mean ± SE. Data were evaluated using a 1-way ANOVA with post-hoc Tukey's comparison.
[2] Mutations are introduced into codon-optimized SERT with a c-myc tag in EL2.

DISCUSSION

Protein sequences are highly optimized over natural history for their physiological

functions. Here, we employed a fluorescent-based assay to determine the effects of nearly all

amino acid substitutions on human SERT surface expression and transport. The selection of

SERT mutants for surface expression reveals conservation and evolutionary constraints for

plasma membrane trafficking and demonstrates the necessity for maintaining the proper three-

dimensional fold, rather than just membrane helix insertion and biogenesis per se. Furthermore,

SERT variants were selected for enhanced transport of an unnatural substrate for which the

protein sequence is suboptimal, and which simulations indicate is unable to promote the

conformational changes that drive transport. As a consequence, hundreds of gain-of-function

mutations were discovered that richly inform molecular mechanism and provide insight into

conformational changes that limit transport kinetics. We conclude that in the absence of the

cognate substrate, there is a high free energy barrier for transition to the IF state, which will

maintain ion-neurotransmitter coupling and prevent ion leakage, and the symported Na^+ and

neurotransmitter share a common exit pathway that begins with the Na2 site at the apex of the

intracellular vestibule. This proposed import mechanism is likely shared with related

neurotransmitter:sodium symporters.

The technical requirements for *in vitro* selection restricted investigations to the

fluorescent substrate and neurotransmitter analogue APP+ (*104*), which is larger than and has

reduced configurational entropy relative to 5-HT (Figure S1), and its import was therefore

sensitive to mutations along the permeation pathway. The concept of discovering mechanism by

studying an alternative substrate is widely practiced in the field of enzyme directed evolution

(*105*), and engineering of major facilitator superfamily transporters to redirect substrate

preference has illuminated important motifs and sequence features (*106–109*). Notably amongst

studies of NSS homologues, rational mutagenesis switched LeuT from being inhibited by to now

transporting tryptophan, demonstrating that substrate complementarity is necessary for driving

the OF-OC structural transition (*110*). Our work here similarly uses an unnatural substrate to

inform the mechanism of monoamine NSS transporters through the discovery of gain-of-function

mutations.

SERT self-associates into poorly defined oligomers containing up to 8 subunits (*111*,

112), consistent with aggregation or lipid microdomain localization as opposed to distinct

protein-protein interactions. The structures of human SERT and *Drosophila* DAT are both

monomeric (*19*, *26*, *30*), and while detergent extraction can dissociate protein-protein interfaces,

it nonetheless suggests against a specific, tight oligomer. Crystal structures of LeuT obtained in a

dimer suggested that the two monomers may form an interface at TM12 (*18*). Periole et al.

conducted coarse-grained SERT simulations and observed the probability of various dimer

interfaces involving different transmembrane helices, of which symmetric TM12-TM12 and

asymmetric TM7-TM12 dimers were most favorable (*113*). In light of these studies, it remains

unknown whether SERT forms specific oligomers that are biologically relevant. In investigations

of a dimeric G protein-coupled receptor that is closely related to metabotropic neurotransmitter

receptors, deep mutagenesis identified membrane-exposed surface patches that were relatively

highly conserved during *in vitro* selection; the data guided the integrative structure determination

for the protein's dimeric architecture that closely agreed with cryo-electron microscopy (*46*).

Searching for surface patches on SERT that are conserved in the deep mutational scans, the

landscapes for SERT expression and APP+ transport identify the membrane-exposed surface

formed by TM2 and TM7 as a candidate oligomerization or lipid-interaction site. However, these

helices also form critical contacts to Cl^- and Na^+ within the protein core, and their higher

conservation may simply reflect their importance for proper folding with bound ions. The

mechanisms by which SERT proteins self-associate requires further investigation.

The mechanistic details described here may provide insights into how SERT kinetics are

regulated *in vivo*. For example, syntaxin 1A interacts with the SERT N-terminus to dampen

excess Na^+ flux during 5-HT transport (*23*, *114*), and it may be that syntaxin 1A stabilizes the

OF state with a bound N-terminal gating loop to ensure free energy barriers are sufficiently high

for strict coupled transport. By comparison, phosphorylation of Thr276 by PKG when SERT is

in the IF conformation increases transport velocity, possibly by preferentially stabilizing the IF

state (*68*). Hence under certain physiological conditions, nature may have exploited features of

the conformational free energy landscape to increase kinetics or change ion coupling and flux.

METHODS

Cell Culture. Expi293F cells (ThermoFisher) were cultured in Expi293F expression medium

(ThermoFisher) at 37ºC, 8% CO_2, 125 rpm. GripTite HEK-293 (ThermoFisher) cells were grown

in Dulbecco's Modified Eagle Medium supplemented with 10% fetal bovine serum and 150

ng/ml G418.

Reagents. APP+ was from Aobious, prepared as a 100 mM solution in DMSO, and stored at -

80°C. Neurotransmitters 5-HT, norepinephrine, and dopamine were obtained from Sigma-

Aldrich and prepared as 100 mM (dopamine and norepinephrine) or 25 mM (5-HT) solutions in

water. Neurotransmitter stocks were stored at -80°C until use.

Plasmid Construction. Codon optimized human SERT was assembled from synthetic DNA

fragments (IDT) and inserted into the NheI-XhoI sites of pCEP4 (Invitrogen). The use of a

synthetic gene ensures there are no sequence features, such as GC-rich tracts, resistant to PCR-

based mutagenesis, and therefore aids in library construction. A c-myc epitope tag flanked by a

glycine / serine rich linker was inserted into EL2 between residues 216 and 217. A version

without the tag was created by PCR-based mutagenesis. For protein expression from the native

human cDNA, SERT was subcloned from pcDNA3-hSERT (Addgene 15483)[68] into Kpn1-Xho1

sites of pCEP4. A c-myc epitope tag was inserted into EL2 using overlap extension PCR. To

measure surface expression of SERT normalized by total expression, superfolder GFP was fused

directly to the myc-SERT N-terminus. All constructs featured a consensus Kozak sequence.

Important plasmids are deposited with Addgene.

Library Construction. Two separate SSM libraries were created from plasmid pCEP4 encoding codon optimized myc-SERT. The SERT N-terminal library covered amino acids 2-325, and the C-terminal library covered amino acids 326-630. For each library, overlap extension PCR using primers containing degenerate NNK codons was applied (*115*), and the pooled PCR products were cut and ligated into NheI-NotI or HindIII-XhoI sites for the N-terminal and C-terminal libraries, respectively. Ligation products were electroporated into NEB5α cells (New England Biolabs), with the number of transformants at least 50 times the possible DNA sequence diversity. Plasmid libraries were purified using GeneJet Maxi Prep kit (ThermoFisher).

Library Selection. Libraries were expressed transiently using Expifectamine (ThermoFisher) in Expi293F cells. To reduce the likelihood of two SERT sequences being acquired by a single cell (*48*), 1ng library DNA was diluted with 1.5 μg pCEP4ΔCMV (*46*) carrier plasmid per milliliter of culture at 2×10^6 cells/ml. Cells were incubated at 37 °C, 8% CO_2, with continuous shaking at 125 rpm for 2 h, followed by media replacement, and then incubated for a further 22 h. Transport activity was assessed by washing with PBS-BSA (phosphate-buffered saline supplemented with 0.2% bovine serum albumin), and incubating in 0.5 μM APP+ (Aobious) in PBS-BSA for 15 min at room temperature with gentle agitation. Cells were then washed with ice cold PBS-BSA, incubated on ice for 20 m with 1/200 anti-myc-Alexa 647 (clone 9B11; Cell Signaling Technology), washed twice, and then analyzed on an Accuri C6 or BD LSRII cytometer, or sorted on a BD FACS ARIA II. During sorting, the main cell population was gated by forward scatter-A versus -W to exclude doublets and debris. Propidium iodide-negative events were then gated to remove dead cells. The top 50 % of SERT-expressing cells (based on Alexa 647 fluorescence) were collected and frozen at -80 °C. For analysis of transport, the top 10-15 % of APP+ fluorescent cells were collected within the SERT-expression gate. To adequately sample

each mutation in the libraries during sorts, at least 100,000 cells were collected, pooling frozen

cell pellets from separate days as necessary to create one sample from which cDNA was

prepared. Typically, ~8 h of sorting was required for each condition, and transfected cultures

were not sorted for longer than 4 hours to maintain viability. Each deep mutational scan

experiment consisted of at least two separately sorted and prepared transfected cell samples.

Library Sequencing and Analysis. Total RNA was extracted from the sorted cells (GeneJet kit

from ThermoFisher), followed by first strand cDNA synthesis primed with gene-specific

oligonucleotides using high-fidelity Accuscript (Agilent). DNA fragments were amplified in two

rounds of PCR as previously described (*46*, *115*), adding Illumina adapter sequences and

barcodes. Samples were sequenced on an Illumina MiSeq or HiSeq2500 using 2×250 nt paired

end read protocol. Data were processed using Enrich (*116*), and commands for analysis are

included in the GEO submission. Enrichment ratios for each variant were calculated by dividing

the transcript frequency from the selected population by the plasmid DNA frequency in the naive

plasmid library. Sequence variants with < 10 reads (equivalent to a frequency of ~3.5×10^{-6}) in

the naïve plasmid library were considered missing. Log_2 enrichment scores are displayed as heat

maps using gnuplot (http://www.gnuplot.info/), or displayed on the crystal structure of (s)-

citalopram-bound SERT (PDB 5I73) using PyMOL (Shrodinger, LLC).

Variant Selection, Targeted Mutagenesis, and Analysis. Variants for validation by targeted

mutagenesis were selected from the deep mutational scan with a log_2 enrichment ratio of >3 for

APP+ import and <1 for expression, or were reported human alleles. Primers containing the

specific codon change were used for site-directed mutagenesis using overlap extension PCR.

SERT variants (500 ng plasmid DNA per ml culture) were transiently transfected into Expi293F

cells (2×10^6 cells/ml) using Expifectamine (ThermoFisher). Transfection enhancers were added

16-18 h post-transfection, and cells were harvested for analysis 36 h post-transfection. APP+

transport activity was assessed as described for library preparation. Samples were analyzed on an

Accuri C6 cytometer. Total and surface protein expression were measured using the BD

Fixation/Permeabilization Solution Kit (BD Biosciences) in conjunction with anti-myc-Alexa

647 staining (clone 9B11; Cell Signaling Technology). For testing neurotransmitter selectivity,

assays were performed on Expi293F cells incubated with 0.5 µM APP+ and varying

concentrations of 5-HT, dopamine, or norepinephrine in Ringer's Solution (140 mM NaCl, 5

mM KCl, 1.2 mM $MgCl_2$, 5 mM Glucose, 2 mM $CaCl_2$, 10 mM HEPES, pH 7.5) supplemented

with 0.2% bovine serum albumin. Ion dependency was measured by replacing NaCl in Ringer's-

BSA solution with Choline-Cl (140 mM) and adjusting the pH to 7.5 with choline-hydroxide

solution, or by replacing chloride salts with salts of gluconate, and adjusting pH with KOH.

The mean fluorescence (F_{APP+}) of cells incubated with APP+ was measured by flow cytometry.

Background autofluorescence (F_0) was subtracted using SERT-transfected cells without APP+ to

calculate $\Delta F = F_{APP+} - F_0$. To assess transport activity (T), ΔF of mock transfected control cells

incubated with APP+ was subtracted from ΔF of SERT-expressing cells: $T = \Delta F_{SERT} - \Delta F_{negative}$.

For quantifying surface expression, autofluorescence of untransfected cells incubated with Alexa

647-conjugated anti-myc antibody was subtracted from the mean fluorescence units of the

SERT-transfected population. All replicates are independent samples and statistical tests as

described in legends were calculated using Prism 10.

[^{3}H] 5-HT Single-Point Uptake Assay. GripTite HEK-293 cells were transiently transfected

with 100 ng of plasmid DNA and grown in 24-well CulturPlate-24 plates (PerkinElmer) to 80-

90% confluence. Cells were washed twice with 500 µL of 37 °C Modified Krebs-Ringer HEPES

+ glucose (MKRGH: 5 mM Tris, 7.5 mM HEPES, 120 mM NaCl, 5.4 mM KCl, 1.2 mM $CaCl_2$,

1.2 mM MgSO$_4$, 10 mM glucose, pH 7.4). Uptake was performed in a final volume of 250 μL

with 20 nM hydroxytryptamine creatinine sulfate, 5-[1,2-^{3}H(N)] ([^{3}H]5-HT) at 37.5 Ci/mmole

(PerkinElmer, Cat# NET498001MC). Uptake was allowed to proceed for 10 min at 37 °C

followed by three washes with ice-cold MKRH, aspirated dry, and solubilized with 250 μL of

MicroScint-20 (PerkinElmer). [^{3}H]5-HT uptake was measured with TopCountNXT Microplate

Scintillation and Luminescence Counter (PerkinElmer) and nonspecific uptake determined with

10 μM (*S*)-citalopram treated and parental cells.

Saturation Uptake Analysis. GripTite HEK-293 cells were grown, transfected, and washed as

above. 200 μl of MKRHG was as added to each well followed by 2-fold serial dilutions of non-

radiolabled 5-HT prepared in MKRHG with ascorbic acid and iproniazid. [^{3}H]5-HT was added

to each well to a final concentration of 30 nM. After 2.5 minutes, the cells were washed three

times with ice-cold MKRH and radioactivity quantified (CPM) as above. CPMs were converted

to amoles based on specific activity of 5-HT in each well and counter efficiency. The rate of

uptake was plotted versus 5-HT concentration to determine K_M and V_{max}.

All Atom MD Simulations. Simulations were initiated from an OF Na$^+$-bound SERT structure,

with Na$^+$ bound in the Na1 and Na2 sites, obtained from our previous study (*44*). The protein

was embedded in a 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC) bilayer using the

Membrane Builder plugin in CHARMM-GUI (*117*). The terminal chains were capped with

acetyl and methyl amide groups. 1 APP+ molecule was added to the simulation box. 150 mM

NaCl was added to neutralize the system and mimic physiological conditions. Glu508 was

modeled in its protonated form. The disulfide bond between Cys200 and Cys209 of EL2 was

present. The system was solvated with TIP3P water molecules (*118*) in a box volume of 75 x 75

x 112 Å^3. Simulations were implemented with the Amber18 package (*119*) using Amber

ff14SB(*120*) and GAFF2 force fields(*121*). Parameters for the APP+ molecule were generated

from the Antechamber package (*122*). The MD system was minimized for 20,000 steps using the

conjugate gradient method, heated from 0 to 300 K at NVT, and equilibrated for 40 ns under

NPT conditions. All production runs were conducted at constant NPT conditions (300K, 1 atm),

periodic boundary conditions, and hydrogen mass repartitioning (*123*). An integration timestep

of 4 femtosecond (fs) was used. Temperature (300 K) was maintained using Langevian dynamics

with a 1.0 picosecond^{-1} damping coefficient. Pressure (1 atm) was maintained using Monte Carlo

barostat with an update interval every 100 simulation steps. The nonbonded distance cutoff was

set at 10 Å. Electrostatic interactions were treated with the Particle Mesh Ewald method (*124*),

and hydrogen bonds were constrained using SHAKE algorithm (*125*). Simulation snapshots were

saved for every 100 ps during production simulations.

As conducted in our previous study, we implemented a Markov state model (MSM)-based

adaptive sampling approach (*126*, *127*) to thoroughly sample the conformational landscape of

SERT during the APP+ import process. This approach involved an iterative sampling procedure

in which multiple simulations were conducted in parallel. Data were then clustered based on a

designated metric of the z-position of the symported Na^+, Cl^-, and APP+ using a K-means

algorithm. States from the least populated clusters were chosen for the next iteration of

simulations. A total of ~370 μs of simulation time was collected and thoroughly sampled the

conformational landscape (Figure S9A).

MSM Construction. Markov state modeling is a statistical approach for discretizing the

simulation data into kinetically relevant clusters and calculating the transition probability

between each cluster (*73*). MSM were constructed with the pyEMMA 2.5.6 Python package

(*128*). To compare the APP+ import process with Na^+ and 5-HT-SERT simulations, we

constructed the MSM using the same 16 interhelical distances surrounding the channel pore

radius and z-components of substrates (APP+, Cl⁻, and the symported Na$^+$) as featurization

metrics (Figure S10A). The number of clusters was optimized based on the VAMP scoring

function implemented in pyEMMA. 4 time independent components (tICs) and 600 clusters were

chosen to construct the MSM (Figure S10B). The lag time of 12 ns was determined to be a

Markovian lag time based on implied timescale plots (Figure S10C). To validate the model, we

perform a Chapman-Kolmogorov test (*129*) on a 4 macrostates MSM (Figure S11).

Trajectory Processing and Analysis. The CPPTRAJ module (*130*) implemented in

AmberTools and MDTraj (*131*) was used for post processing of trajectory data. All simulation

data were projected on the coordinate landscape defined by the extracellular gating distance as

the closest heavy atom between Arg104 and Glu493 and the intracellular distance as the closest

heavy atom between the groups of residues Val86-Ser91, Tyr350 and Val199-Trp282, Glu444.

To estimate the error of the free energy landscape, a bootstrapping method was implemented in

which 500 independent MSM were constructed using a random selection of 80% of the trajectory

dataset (Figure S9B). Trajectories and MSM states were visualized with Visual Molecular

Dynamics (VMD) (*132*) and PyMOL (Shrodinger, LLC). In-house scripts and the matplotlib

python package were used to generate plots.

Clustering the mutational landscape using UMAP. The activities of all amino acid

substitutions per SERT position were clustered using UMAP python package

(github.com/lmcinnes/umap) using the following hyperparameters: n_neighbors = 10, min_dist =

0.2, n_components = 2, metric = euclidian. The mutational data were then projected on the two

UMAP components and colored based on different biochemical and biophysical properties. The

hydrophobicity of residues was calculated using the Kyte-Doolittle scale using a window size of

9. Features from MD simulations, root mean square fluctuations (RMSF), solvent accessible

surface area (SASA), and average number of residue-residue contacts, were calculated on 50,000

structures drawn from an MSM-weighted distribution. The outward-facing SERT structure

(PDB: 5I73) was used as the reference structure for RMSF calculation. SASA calculation was

implemented using the MDtraj package and a 1.4 Å probe radius. An 8Å Cα-Cα distance cutoff

was used to determine residue-residue contacts. Variant effect predictions were generated by

EVmutation (github.com/debbiemarkslab/EVcouplings) and DeepSequence

(github.com/debbiemarkslab/DeepSequence) packages and used the EVcouplings package to

construct the input sequence alignment. *In silico* free energy changes as a result of a mutation

were predicted using the FoldX suite (foldxsuite.crg.eu).

Estimating Evolutionary Conservation. Natural sequence conservation was mapped to the

SERT structure (PDB 5I73) using the ConSurf server (*133*). Homologs were searched using CS-

BLAST from UNIREF-90 with an E-value cutoff of 0.0001 and 3 iterations. Maximum and

minimum identity were set at 95 % and 35 %, respectively. 500 NSS transporter sequences were

aligned with MAFFT and conservation scores calculated using Bayesian algorithm of

substitution.

Data and Software Availability. Raw deep sequencing data and calculated enrichment ratios

are deposited with NCBI's Gene Expression Omnibus under series accession number

GSE109499. Command lines for analysis with Enrich are also included. Representative

structures from the Markov state model and MD simulations are available on Zenodo at

https://doi.org/10.5281/zenodo.4695032.

SUPPORTING INFORMATION

Online Supporting Information includes:

PDF of Supplementary Figures S1 to S8

PyMOL file of SERT colored by conservation score for APP+ transport with overlaid APP+

permeation pathway

Movie S1 – morph of SERT conformational transitions colored by APP+ transport conservation

score

Excel file of enrichment ratios from the SERT deep mutational scans

AUTHOR INFORMATION

Corresponding Authors

* (E.P.) Email: procko@illinois.edu

* (D.S.) Email: diwakar@illinois.edu

Author Contributions

H.J.E. and S.K.S. did experimental characterization and deep mutagenesis. M.C.C. and B.S. did

MD simulations. E.W. and L.K.H. tested uptake of 5-HT. E.P., L.K.H., and D.S. supervised and

directed research.

Notes

The authors declare they have no conflict of interest. E.P. receives stock and compensation from

Cyrus Biotechnology for working on drug programs unrelated to this study.

ACKNOWLEDGEMENTS

Within the UIUC Roy J. Carver Biotechnology Center, Barbara Pilas and Angela Kouris assisted

with flow cytometry, and Alvaro Hernandez and Chris Wright assisted with deep sequencing.

We thank Jesse R. Horne for technical assistance on DeepSequence and FoldX calculations. This

work was funded by R21 MH113155 from NIMH to E.P., and D.S. is supported by NSF Early

Career Award by NSF MCB 18-45606. The research is part of the Blue Waters sustained-

petascale computing project, which is supported by the National Science Foundation (awards

OCI-0725070 and ACI-1238993) and the state of Illinois, and as of December, 2019, the

National Geospatial-Intelligence Agency. Blue Waters is a joint effort of the University of

Illinois at Urbana-Champaign and its National Center for Supercomputing Applications.

REFERENCES

1. R. D. Blakely, H. E. Berson, R. T. Fremeau, M. G. Caron, M. M. Peek, H. K. Prince, C. C.
Bradley, Cloning and expression of a functional serotonin transporter from rat brain.
Nature. **354**, 66–70 (1991).

2. B. J. Hoffman, E. Mezey, M. J. Brownstein, Cloning of a serotonin transporter affected by
antidepressants. *Science*. **254**, 579–580 (1991).

3. D. L. Murphy, G. R. Uhl, A. Holmes, R. Ren☐Patterson, F. S. Hall, I. Sora, S.
Detera☐Wadleigh, K.-P. Lesch, Experimental gene interaction studies with SERT mutant
mice as models for human polygenic and epistatic traits and disorders. *Genes, Brain and*
Behavior. **2**, 350–364 (2003).

4. F. Kilic, D. L. Murphy, G. Rudnick, A Human Serotonin Transporter Mutation Causes
Constitutive Activation of Transport Activity. *Mol Pharmacol*. **64**, 440–446 (2003).

5. N. Ozaki, D. Goldman, W. H. Kaye, K. Plotnicov, B. D. Greenberg, J. Lappalainen, G.
Rudnick, D. L. Murphy, Serotonin transporter missense mutation associated with a
complex neuropsychiatric phenotype. *Molecular Psychiatry*. **8**, 933–936 (2003).

6. H. C. Prasad, J. A. Steiner, J. S. Sutcliffe, R. D. Blakely, Enhanced activity of human
serotonin transporter variants associated with autism. *Philosophical Transactions of the*
Royal Society B: Biological Sciences. **364**, 163–173 (2009).

7. J. S. Sutcliffe, R. J. Delahanty, H. C. Prasad, J. L. McCauley, Q. Han, L. Jiang, C. Li, S. E.
Folstein, R. D. Blakely, Allelic Heterogeneity at the Serotonin Transporter Locus
(SLC6A4) Confers Susceptibility to Autism and Rigid-Compulsive Behaviors. *The*
American Journal of Human Genetics. **77**, 265–279 (2005).

8. M. K. Hahn, R. D. Blakely, Monoamine transporter gene structure and polymorphisms in
relation to psychiatric and other complex disorders. *The Pharmacogenomics Journal*. **2**,
217–235 (2002).

9. T. B. Usdin, E. Mezey, C. Chen, M. J. Brownstein, B. J. Hoffman, Cloning of the cocaine-
sensitive bovine dopamine transporter. *PNAS*. **88**, 11168–11171 (1991).

845 10. B. Giros, S. el Mestikawy, N. Godinot, K. Zheng, H. Han, T. Yang-Feng, M. G. Caron,
846    Cloning, pharmacological characterization, and chromosome assignment of the human
847    dopamine transporter. *Mol Pharmacol*. **42**, 383–390 (1992).

848 11. T. Pacholczyk, R. D. Blakely, S. G. Amara, Expression cloning of a cocaine-and
849    antidepressant-sensitive human noradrenaline transporter. *Nature*. **350**, 350–354 (1991).

850 12. G. Rudnick, W. Sandtner, Serotonin transport in the 21st century. *J Gen Physiol*. **151**,
851    1248–1264 (2019).

852 13. G. Rudnick, P. J. Nelson, Platelet 5-hydroxytryptamine transport, an electroneutral
853    mechanism coupled to potassium. *Biochemistry*. **17**, 4739–4742 (1978).

854 14. P. J. Nelson, G. Rudnick, The role of chloride ion in platelet serotonin transport. *J. Biol.*
855    *Chem.* **257**, 6151–6155 (1982).

856 15. K. Schicker, Z. Uzelac, J. Gesmonde, S. Bulling, T. Stockner, M. Freissmuth, S. Boehm, G.
857    Rudnick, H. H. Sitte, W. Sandtner, Unifying Concept of Serotonin Transporter-associated
858    Currents. *J. Biol. Chem.* **287**, 438–445 (2012).

859 16. P. Mitchell, A General Theory of Membrane Transport From Studies of Bacteria. *Nature*.
860    **180**, 134–136 (1957).

861 17. O. Jardetzky, Simple Allosteric Model for Membrane Pumps. *Nature*. **211**, 969 (1966).

862 18. H. Krishnamurthy, E. Gouaux, X-ray structures of LeuT in substrate-free outward-open and
863    apo inward-open states. *Nature*. **481**, 469–474 (2012).

864 19. J. A. Coleman, D. Yang, Z. Zhao, P.-C. Wen, C. Yoshioka, E. Tajkhorshid, E. Gouaux,
865    Serotonin transporter–ibogaine complexes illuminate mechanisms of inhibition and
866    transport. *Nature*. **569**, 141–145 (2019).

867 20. L. D. Jayanthi, S. Ramamoorthy, Regulation of monoamine transporters: Influence of
868    psychostimulants and therapeutic antidepressants. *AAPS J*. **7**, E728–E738 (2005).

869 21. F. Lin, H. A. Lester, S. Mager, Single-channel currents produced by the serotonin
870    transporter and analysis of a mutation affecting ion permeation. *Biophysical Journal*. **71**,
871    3126–3135 (1996).

872 22. S. Mager, C. Min, D. J. Henry, C. Chavkintt, B. J. Hoffman, N. Davidson, H. A. Lester,
873    Conducting states of a mammalian serotonin transporter. *Neuron*. **12**, 845–859 (1994).

874 23. M. W. Quick, Regulating the Conducting States of a Mammalian Serotonin Transporter.
875    *Neuron*. **40**, 537–549 (2003).

876 24. A. S. Kristensen, J. Andersen, T. N. Jørgensen, L. Sørensen, J. Eriksen, C. J. Loland, K.
877    Strømgaard, U. Gether, SLC6 Neurotransmitter Transporters: Structure, Function, and
878    Regulation. *Pharmacol Rev*. **63**, 585–640 (2011).

25. A. Yamashita, S. K. Singh, T. Kawate, Y. Jin, E. Gouaux, Crystal structure of a bacterial
homologue of Na + /Cl - -dependent neurotransmitter transporters. *Nature*. **437**, 215
(2005).

26. J. A. Coleman, E. M. Green, E. Gouaux, X-ray structures and mechanism of the human
serotonin transporter. *Nature*. **532**, 334–339 (2016).

27. J. A. Coleman, E. Gouaux, Structural basis for recognition of diverse antidepressants by the
human serotonin transporter. *Nature Structural & Molecular Biology*. **25**, 170 (2018).

28. A. Penmatsa, K. H. Wang, E. Gouaux, X-ray structures of Drosophila dopamine transporter
in complex with nisoxetine and reboxetine. *Nature Structural & Molecular Biology*. **22**,
506–508 (2015).

29. A. Penmatsa, K. H. Wang, E. Gouaux, X-ray structure of dopamine transporter elucidates
antidepressant mechanism. *Nature*. **503**, 85–90 (2013).

30. K. H. Wang, A. Penmatsa, E. Gouaux, Neurotransmitter and psychostimulant recognition
by the dopamine transporter. *Nature*. **521**, 322–327 (2015).

31. H. Wang, A. Goehring, K. H. Wang, A. Penmatsa, R. Ressler, E. Gouaux, Structural basis
for action by diverse antidepressants on biogenic amine transporters. *Nature*. **503**, 141–
145 (2013).

32. L. Sørensen, J. Andersen, M. Thomsen, S. M. R. Hansen, X. Zhao, A. Sandelin, K.
Strømgaard, A. S. Kristensen, Interaction of antidepressants with the serotonin and
norepinephrine transporters: mutational studies of the S1 substrate binding pocket. *J. Biol.
Chem.* **287**, 43694–43707 (2012).

33. Z. Zhou, J. Zhen, N. K. Karpowich, R. M. Goetz, C. J. Law, M. E. A. Reith, D.-N. Wang,
LeuT-Desipramine Structure Reveals How Antidepressants Block Neurotransmitter
Reuptake. *Science*. **317**, 1390–1393 (2007).

34. S. K. Singh, A. Yamashita, E. Gouaux, Antidepressant binding site in a bacterial
homologue of neurotransmitter transporters. *Nature*. **448**, 952–956 (2007).

35. H. A. Neubauer, C. G. Hansen, O. Wiborg, Dissection of an Allosteric Mechanism on the
Serotonin Transporter: A Cross-Species Study. *Mol Pharmacol*. **69**, 1242–1250 (2006).

36. L. Malinauskaite, M. Quick, L. Reinhard, J. A. Lyons, H. Yano, J. A. Javitch, P. Nissen, A
mechanism for intracellular release of Na+ by neurotransmitter/sodium symporters. *Nat.
Struct. Mol. Biol.* **21**, 1006–1012 (2014).

37. Y. Zhao, D. Terry, L. Shi, H. Weinstein, S. C. Blanchard, J. A. Javitch, Single-molecule
dynamics of gating in a neurotransmitter transporter homologue. *Nature*. **465**, 188–193
(2010).

38.   Y. Zhao, D. S. Terry, L. Shi, M. Quick, H. Weinstein, S. C. Blanchard, J. A. Javitch,
914   Substrate-modulated gating dynamics in a Na$^+$-coupled neurotransmitter transporter
915   homologue. *Nature*. **474**, 109–113 (2011).

916   39.   L. R. Forrest, Y.-W. Zhang, M. T. Jacobs, J. Gesmonde, L. Xie, B. H. Honig, G. Rudnick,
917   Mechanism for alternating access in neurotransmitter transporters. *PNAS*. **105**, 10338–
918   10343 (2008).

919   40.   A. M. Razavi, G. Khelashvili, H. Weinstein, A Markov State-based Quantitative Kinetic
920   Model of Sodium Release from the Dopamine Transporter. *Scientific Reports*. **7**, 40076
921   (2017).

922   41.   S. A. Shaikh, E. Tajkhorshid, Modeling and Dynamics of the Inward-Facing State of a
923   Na+/Cl− Dependent Neurotransmitter Transporter Homologue. *PLOS Computational
924   Biology*. **6**, e1000905 (2010).

925   42.   S. Adhikary, D. J. Deredge, A. Nagarajan, L. R. Forrest, P. L. Wintrode, S. K. Singh,
926   Conformational dynamics of a neurotransmitter:sodium symporter in a lipid bilayer.
927   *PNAS*. **114**, E1786–E1795 (2017).

928   43.   I. R. Möller, M. Slivacka, A. K. Nielsen, S. G. F. Rasmussen, U. Gether, C. J. Loland, K. D.
929   Rand, Conformational dynamics of the human serotonin transporter during substrate and
930   drug binding. *Nature Communications*. **10**, 1687 (2019).

931   44.   M. C. Chan, B. Selvam, H. J. Young, E. Procko, D. Shukla, The substrate import
932   mechanism of the human serotonin transporter. *Biophysical Journal*. **121**, 715–730 (2022).

933   45.   D. M. Fowler, S. Fields, Deep mutational scanning: a new style of protein science. *Nature
934   Methods*. **11**, 801–807 (2014).

935   46.   J. Park, B. Selvam, K. Sanematsu, N. Shigemura, D. Shukla, E. Procko, Structural
936   architecture of a dimeric class C GPCR based on co-trafficking of sweet taste receptor
937   subunits. *J. Biol. Chem.* **294**, 4759–4774 (2019).

938   47.   J. D. Heredia, J. Park, H. Choi, K. S. Gill, E. Procko, Conformational Engineering of HIV-1
939   Env Based on Mutational Tolerance in the CD4 and PG16 Bound States. *Journal of
940   Virology*. **93** (2019), doi:10.1128/JVI.00219-19.

941   48.   J. D. Heredia, J. Park, R. J. Brubaker, S. K. Szymanski, K. S. Gill, E. Procko, Mapping
942   Interaction Sites on Human Chemokine Receptors by Deep Mutational Scanning. *The
943   Journal of Immunology*. **200**, 3825–3839 (2018).

944   49.   A. C. McShan, C. A. Devlin, S. A. Overall, J. Park, J. S. Toor, D. Moschidi, D. Flores-
945   Solis, H. Choi, S. Tripathi, E. Procko, N. G. Sgourakis, Molecular determinants of
946   chaperone interactions on MHC-I for folding and antigen repertoire selection. *PNAS*. **116**,
947   25602–25613 (2019).

50. N. J. Rollins, K. P. Brock, F. J. Poelwijk, M. A. Stiffler, N. P. Gauthier, C. Sander, D. S.
Marks, Inferring protein 3D structure from deep mutation scans. *Nat Genet*. **51**, 1170–
1176 (2019).

51. J. M. Schmiedel, B. Lehner, Determining protein structures using deep mutagenesis. *Nature
Genetics*. **51**, 1177–1186 (2019).

52. B. Felts, A. B. Pramod, W. Sandtner, N. Burbach, S. Bulling, H. H. Sitte, L. K. Henry, The
Two Na+ Sites in the Human Serotonin Transporter Play Distinct Roles in the Ion
Coupling and Electrogenicity of Transport. *J Biol Chem*. **289**, 1825–1840 (2014).

53. S. Ramamoorthy, A. L. Bauman, K. R. Moore, H. Han, T. Yang-Feng, A. S. Chang, V.
Ganapathy, R. D. Blakely, Antidepressant- and cocaine-sensitive human serotonin
transporter: molecular cloning, expression, and chromosomal localization. *Proc Natl Acad
Sci U S A*. **90**, 2542–2546 (1993).

54. G. Rudnick, Active transport of 5-hydroxytryptamine by plasma membrane vesicles
isolated from human blood platelets. *J. Biol. Chem.* **252**, 2170–2174 (1977).

55. C. Fenollar-Ferrer, T. Stockner, T. C. Schwarz, A. Pal, J. Gotovina, T. Hofmaier, K.
Jayaraman, S. Adhikary, O. Kudlacek, A. R. Mehdipour, S. Tavoulari, G. Rudnick, S. K.
Singh, R. Konrat, H. H. Sitte, L. R. Forrest, Structure and Regulatory Interactions of the
Cytoplasmic Terminal Domains of Serotonin Transporter. *Biochemistry*. **53**, 5444–5460
(2014).

56. J.-G. Chen, S. Liu-Chen, G. Rudnick, External Cysteine Residues in the Serotonin
Transporter. *Biochemistry*. **36**, 1479–1486 (1997).

57. L. Laursen, K. Severinsen, K. B. Kristensen, X. Periole, M. Overby, H. K. Müller, B.
Schiøtt, S. Sinning, Cholesterol binding to a conserved site modulates the conformation,
pharmacology, and transport kinetics of the human serotonin transporter. *Journal of
Biological Chemistry*. **293**, 3510–3523 (2018).

58. T. Hessa, N. M. Meindl-Beinker, A. Bernsel, H. Kim, Y. Sato, M. Lerch-Bader, I. Nilsson,
S. H. White, G. von Heijne, Molecular code for transmembrane-helix recognition by the
Sec61 translocon. *Nature*. **450**, 1026–1030 (2007).

59. Y.-W. Zhang, S. Tavoulari, S. Sinning, A. A. Aleksandrova, L. R. Forrest, G. Rudnick,
Structural elements required for coupling ion and substrate transport in the
neurotransmitter transporter homolog LeuT. *PNAS*. **115**, E8854–E8862 (2018).

60. J. Andersen, L. Olsen, K. B. Hansen, O. Taboureau, F. S. Jørgensen, A. M. Jørgensen, B.
Bang-Andersen, J. Egebjerg, K. Strømgaard, A. S. Kristensen, Mutational Mapping and
Modeling of the Binding Site for (S)-Citalopram in the Human Serotonin Transporter. *J.
Biol. Chem.* **285**, 2051–2063 (2010).

61. T. Beuming, J. Kniazeff, M. L. Bergmann, L. Shi, L. Gracia, K. Raniszewska, A. H.
Newman, J. A. Javitch, H. Weinstein, U. Gether, C. J. Loland, The binding sites for

cocaine and dopamine in the dopamine transporter overlap. *Nat Neurosci*. **11**, 780–789
(2008).

62. V. M. Korkhov, M. Holy, M. Freissmuth, H. H. Sitte, The Conserved Glutamate (Glu136)
in Transmembrane Domain 2 of the Serotonin Transporter Is Required for the
Conformational Switch in the Transport Cycle. *J. Biol. Chem.* **281**, 13439–13448 (2006).

63. N. Chen, R. A. Vaughan, M. E. A. Reith, The role of conserved tryptophan and acidic
residues in the human dopamine transporter as characterized by site-directed mutagenesis.
Journal of Neurochemistry. **77**, 1116–1127 (2001).

64. S. Sucic, F. A. Paczkowski, F. Runkel, H. Bönisch, L. J. Bryan-Lluka, Functional
significance of a highly conserved glutamate residue of the human noradrenaline
transporter. *Journal of Neurochemistry*. **81**, 344–354 (2002).

65. G. I. Keshet, A. Bendahan, H. Su, S. Mager, H. A. Lester, B. I. Kanner, Glutamate-101 is
critical for the function of the sodium and chloride-coupled GABA transporter GAT-1.
FEBS Letters. **371**, 39–42 (1995).

66. L. R. Forrest, S. Tavoulari, Y.-W. Zhang, G. Rudnick, B. Honig, Identification of a chloride
ion binding site in Na+/Cl−-dependent transporters. *PNAS*. **104**, 12761–12766 (2007).

67. H. C. Prasad, C.-B. Zhu, J. L. McCauley, D. J. Samuvel, S. Ramamoorthy, R. C. Shelton,
W. A. Hewlett, J. S. Sutcliffe, R. D. Blakely, Human serotonin transporter variants display
altered sensitivity to protein kinase G and p38 mitogen-activated protein kinase. *PNAS*.
102, 11545–11550 (2005).

68. Y.-W. Zhang, J. Gesmonde, S. Ramamoorthy, G. Rudnick, Serotonin Transporter
Phosphorylation by cGMP-Dependent Protein Kinase Is Altered by a Mutation Associated
with Obsessive–Compulsive Disorder. *J. Neurosci.* **27**, 10878–10886 (2007).

69. N. R. Sealover, B. Felts, C. P. Kuntz, R. E. Jarrard, G. H. Hockerman, E. L. Barker, L. K.
Henry, The external gate of the human and Drosophila serotonin transporters requires a
basic/acidic amino acid pair for 3,4-methylenedioxymethamphetamine (MDMA)
translocation and the induction of substrate efflux. *Biochemical Pharmacology*. **120**, 46–
55 (2016).

70. M. Chan, B. Selvam, H. Young, E. Procko, D. Shukla, The Substrate Import Mechanism of
the Human Serotonin Transporter (2020), doi:10.26434/chemrxiv.9922301.v2.

71. L. K. Henry, E. M. Adkins, Q. Han, R. D. Blakely, Serotonin and Cocaine-sensitive
Inactivation of Human Serotonin Transporters by Methanethiosulfonates Targeted to
Transmembrane Domain I*. *Journal of Biological Chemistry*. **278**, 37052–37063 (2003).

72. J. R. Field, L. K. Henry, R. D. Blakely, Transmembrane Domain 6 of the Human Serotonin
Transporter Contributes to an Aqueously Accessible Binding Pocket for Serotonin and the
Psychostimulant 3,4-Methylene Dioxymethamphetamine*. *Journal of Biological
Chemistry*. **285**, 11270–11280 (2010).

73. B. E. Husic, V. S. Pande, Markov State Models: From an Art to a Science. *J. Am. Chem. Soc.* **140**, 2386–2396 (2018).

74. B. Selvam, S. Mittal, D. Shukla, Free Energy Landscape of the Complete Transport Cycle in a Key Bacterial Transporter. *ACS Cent Sci*. **4**, 1146–1154 (2018).

75. D. Shukla, C. X. Hernández, J. K. Weber, V. S. Pande, Markov State Models Provide Insights into Dynamic Modulation of Protein Function. *Acc. Chem. Res.* **48**, 414–422 (2015).

76. B. Selvam, Y.-C. Yu, L.-Q. Chen, D. Shukla, Molecular Basis of the Glucose Transport Mechanism in Plants. *ACS Cent. Sci.* **5**, 1085–1096 (2019).

77. D. P. Claxton, M. Quick, L. Shi, F. D. de Carvalho, H. Weinstein, J. A. Javitch, H. S. Mchaourab, Ion/substrate-dependent conformational dynamics of a bacterial homolog of neurotransmitter:sodium symporters. *Nature Structural & Molecular Biology*. **17**, 822–829 (2010).

78. M. H. Cheng, C. Kaya, I. Bahar, Quantitative Assessment of the Energetics of Dopamine Translocation by Human Dopamine Transporter. *J. Phys. Chem. B*. **122**, 5336–5346 (2018).

79. J. Li, S. A. Shaikh, G. Enkavi, P.-C. Wen, Z. Huang, E. Tajkhorshid, Transient formation of water-conducting states in membrane transporters. *PNAS*. **110**, 7696–7701 (2013).

80. M. Lek, K. J. Karczewski, E. V. Minikel, K. E. Samocha, E. Banks, T. Fennell, A. H. O'Donnell-Luria, J. S. Ware, A. J. Hill, B. B. Cummings, T. Tukiainen, D. P. Birnbaum, J. A. Kosmicki, L. E. Duncan, K. Estrada, F. Zhao, J. Zou, E. Pierce-Hoffman, J. Berghout, D. N. Cooper, N. Deflaux, M. DePristo, R. Do, J. Flannick, M. Fromer, L. Gauthier, J. Goldstein, N. Gupta, D. Howrigan, A. Kiezun, M. I. Kurki, A. L. Moonshine, P. Natarajan, L. Orozco, G. M. Peloso, R. Poplin, M. A. Rivas, V. Ruano-Rubio, S. A. Rose, D. M. Ruderfer, K. Shakir, P. D. Stenson, C. Stevens, B. P. Thomas, G. Tiao, M. T. Tusie-Luna, B. Weisburd, H.-H. Won, D. Yu, D. M. Altshuler, D. Ardissino, M. Boehnke, J. Danesh, S. Donnelly, R. Elosua, J. C. Florez, S. B. Gabriel, G. Getz, S. J. Glatt, C. M. Hultman, S. Kathiresan, M. Laakso, S. McCarroll, M. I. McCarthy, D. McGovern, R. McPherson, B. M. Neale, A. Palotie, S. M. Purcell, D. Saleheen, J. M. Scharf, P. Sklar, P. F. Sullivan, J. Tuomilehto, M. T. Tsuang, H. C. Watkins, J. G. Wilson, M. J. Daly, D. G. MacArthur, Analysis of protein-coding genetic variation in 60,706 humans. *Nature*. **536**, 285–291 (2016).

81. S. Sucic, S. Dallinger, B. Zdrazil, R. Weissensteiner, T. N. Jørgensen, M. Holy, O. Kudlacek, S. Seidel, J. H. Cha, U. Gether, A. H. Newman, G. F. Ecker, M. Freissmuth, H. H. Sitte, The N Terminus of Monoamine Transporters Is a Lever Required for the Action of Amphetamines. *J. Biol. Chem.* **285**, 10924–10938 (2010).

82. O. Dayan-Alon, B. I. Kanner, Internal gate mutants of the GABA transporter GAT1 are capable of substrate exchange. *Neuropharmacology*. **161**, 107534 (2019).

83. P. Metzner, C. Schütte, E. Vanden-Eijnden, Transition Path Theory for Markov Jump
Processes. *Multiscale Model. Simul.* **7**, 1192–1219 (2009).

84. P. C. Söderhielm, J. Andersen, L. Munro, A. T. Nielsen, A. S. Kristensen, Substrate and
Inhibitor–Specific Conformational Changes in the Human Serotonin Transporter Revealed
by Voltage-Clamp Fluorometry. *Mol Pharmacol*. **88**, 676–688 (2015).

85. S. Tavoulari, E. Margheritis, A. Nagarajan, D. C. DeWitt, Y.-W. Zhang, E. Rosado, S.
Ravera, E. Rhoades, L. R. Forrest, G. Rudnick, Two Na+ Sites Control Conformational
Change in a Neurotransmitter Transporter Homolog. *J. Biol. Chem.* **291**, 1456–1471
(2016).

86. J. Andersen, L. Olsen, K. B. Hansen, O. Taboureau, F. S. Jørgensen, A. M. Jørgensen, B.
Bang-Andersen, J. Egebjerg, K. Strømgaard, A. S. Kristensen, Mutational mapping and
modeling of the binding site for (S)-citalopram in the human serotonin transporter. *J. Biol.*
Chem. **285**, 2051–2063 (2010).

87. H. Koldsø, K. Severinsen, T. T. Tran, L. Celik, H. H. Jensen, O. Wiborg, B. Schiøtt, S.
Sinning, The two enantiomers of citalopram bind to the human serotonin transporter in
reversed orientations. *J. Am. Chem. Soc.* **132**, 1311–1322 (2010).

88. B. A. Davis, A. Nagarajan, L. R. Forrest, S. K. Singh, Mechanism of Paroxetine (Paxil)
Inhibition of the Serotonin Transporter. *Sci Rep*. **6**, 23789 (2016).

89. L. K. Henry, H. Iwamoto, J. R. Field, K. Kaufmann, E. S. Dawson, M. T. Jacobs, C.
Adams, B. Felts, I. Zdravkovic, V. Armstrong, S. Combs, E. Solis, G. Rudnick, S. Y.
Noskov, L. J. DeFelice, J. Meiler, R. D. Blakely, A Conserved Asparagine Residue in
Transmembrane Segment 1 (TM1) of Serotonin Transporter Dictates Chloride-coupled
Neurotransmitter Transport. *J. Biol. Chem.* **286**, 30823–30836 (2011).

90. H. Koldsø, P. Noer, J. Grouleff, H. E. Autzen, S. Sinning, B. Schiøtt, Unbiased Simulations
Reveal the Inward-Facing Conformation of the Human Serotonin Transporter and Na+ Ion
Release. *PLOS Computational Biology*. **7**, e1002246 (2011).

91. M. H. Cheng, I. Bahar, Molecular Mechanism of Dopamine Transport by Human
Dopamine Transporter. *Structure*. **23**, 2171–2181 (2015).

92. S. Tavoulari, E. Margheritis, A. Nagarajan, D. C. DeWitt, Y.-W. Zhang, E. Rosado, S.
Ravera, E. Rhoades, L. R. Forrest, G. Rudnick, Two Na+ Sites Control Conformational
Change in a Neurotransmitter Transporter Homolog. *J. Biol. Chem.* **291**, 1456–1471
(2016).

93. A. K. Kantcheva, M. Quick, L. Shi, A.-M. L. Winther, S. Stolzenberg, H. Weinstein, J. A.
Javitch, P. Nissen, Chloride binding site of neurotransmitter sodium symporters.
Proceedings of the National Academy of Sciences. **110**, 8489–8494 (2013).

94.   B. Felts, A. B. Pramod, W. Sandtner, N. Burbach, S. Bulling, H. H. Sitte, L. K. Henry, The
1096   Two Na+ Sites in the Human Serotonin Transporter Play Distinct Roles in the Ion
1097   Coupling and Electrogenicity of Transport. *J. Biol. Chem.* **289**, 1825–1840 (2014).

1098   95.   D. A. Caplan, J. O. Subbotina, S. Y. Noskov, Molecular Mechanism of Ion-Ion and Ion-
1099   Substrate Coupling in the Na+-Dependent Leucine Transporter LeuT. *Biophysical*
1100   *Journal*. **95**, 4613–4621 (2008).

1101   96.   J. Abramson, E. M. Wright, Structure and Function of Na+-Symporters with Inverted
1102   Repeats. *Curr Opin Struct Biol*. **19**, 425–432 (2009).

1103   97.   L. McInnes, J. Healy, J. Melville, UMAP: Uniform Manifold Approximation and Projection
1104   for Dimension Reduction. *arXiv:1802.03426 [cs, stat]* (2020) (available at
1105   http://arxiv.org/abs/1802.03426).

1106   98.   M. W. Dorrity, L. M. Saunders, C. Queitsch, S. Fields, C. Trapnell, Dimensionality
1107   reduction by UMAP to visualize physical and genetic interactions. *Nature*
1108   *Communications*. **11**, 1537 (2020).

1109   99.   E. M. Jones, N. B. Lubock, A. Venkatakrishnan, J. Wang, A. M. Tseng, J. M. Paggi, N. R.
1110   Latorraca, D. Cancilla, M. Satyadi, J. E. Davis, M. M. Babu, R. O. Dror, S. Kosuri,
1111   Structural and functional characterization of G protein–coupled receptors with deep
1112   mutational scanning. *eLife*. **9**, e54895 (2020).

1113   100.   T. A. Hopf, J. B. Ingraham, F. J. Poelwijk, C. P. I. Schärfe, M. Springer, C. Sander, D. S.
1114   Marks, Mutation effects predicted from sequence co-variation. *Nat Biotechnol*. **35**, 128–
1115   135 (2017).

1116   101.   A. J. Riesselman, J. B. Ingraham, D. S. Marks, Deep generative models of genetic variation
1117   capture the effects of mutations. *Nature methods*. **15**, 816–822 (2018).

1118   102.   J. Schymkowitz, J. Borg, F. Stricher, R. Nys, F. Rousseau, L. Serrano, The FoldX web
1119   server: an online force field. *Nucleic Acids Research*. **33**, W382–W388 (2005).

1120   103.   K. Kazmier, S. Sharma, S. M. Islam, B. Roux, H. S. Mchaourab, Conformational cycle and
1121   ion-coupling mechanism of the Na+/hydantoin transporter Mhp1. *PNAS*. **111**, 14752–
1122   14757 (2014).

1123   104.   E. Solis, I. Zdravkovic, I. D. Tomlinson, S. Y. Noskov, S. J. Rosenthal, L. J. De Felice, 4-
1124   (4-(dimethylamino)phenyl)-1-methylpyridinium (APP+) is a fluorescent substrate for the
1125   human serotonin transporter. *J. Biol. Chem.* **287**, 8852–8863 (2012).

1126   105.   C. A. Tracewell, F. H. Arnold, Directed enzyme evolution: climbing fitness peaks one
1127   amino acid at a time. *Current Opinion in Chemical Biology*. **13**, 3–9 (2009).

1128   106.   A. Reinders, Y. Sun, K. L. Karvonen, J. M. Ward, Identification of Amino Acids Important
1129   for Substrate Specificity in Sucrose Transporters Using Gene Shuffling. *J. Biol. Chem.*
1130   **287**, 30296–30304 (2012).

107. A. Farwick, S. Bruder, V. Schadeweg, M. Oreb, E. Boles, Engineering of yeast hexose
transporters to transport d-xylose without inhibition by d-glucose. *PNAS*. **111**, 5159–5164
(2014).

108. J. G. Nijland, H. Y. Shin, R. M. de Jong, P. P. de Waal, P. Klaassen, A. J. Driessen,
Engineering of an endogenous hexose transporter into a specific D-xylose transporter
facilitates glucose-xylose co-consumption in Saccharomyces cerevisiae. *Biotechnology for*
Biofuels. **7**, 168 (2014).

109. E. M. Young, A. Tong, H. Bui, C. Spofford, H. S. Alper, Rewiring yeast sugar transporter
preference through modifying a conserved protein motif. *PNAS*. **111**, 131–136 (2014).

110. C. L. Piscitelli, E. Gouaux, Insights into transport mechanism from LeuT engineered to
transport tryptophan. *The EMBO Journal*. **31**, 228–235 (2012).

111. A. Anderluh, E. Klotzsch, A. W. A. F. Reismann, M. Brameshuber, O. Kudlacek, A. H.
Newman, H. H. Sitte, G. J. Schütz, Single molecule analysis reveals coexistence of stable
serotonin transporter monomers and oligomers in the live cell plasma membrane. *J. Biol.*
Chem. **289**, 4387–4394 (2014).

112. A. Anderluh, T. Hofmaier, E. Klotzsch, O. Kudlacek, T. Stockner, H. H. Sitte, G. J. Schütz,
Direct PIP$_2$ binding mediates stable oligomer formation of the serotonin transporter.
Nature Communications. **8**, 14089 (2017).

113. X. Periole, T. Zeppelin, B. Schiøtt, Dimer Interface of the Human Serotonin Transporter
and Effect of the Membrane Composition. *Scientific Reports*. **8**, 5080 (2018).

114. M. A. Ciccone, M. Timmons, A. Phillips, M. W. Quick, Calcium/Calmodulin-Dependent
Kinase II Regulates the Interaction between the Serotonin Transporter and Syntaxin 1A.
Neuropharmacology. **55**, 763–770 (2008).

115. E. Procko, R. Hedman, K. Hamilton, J. Seetharaman, S. J. Fleishman, M. Su, J. Aramini, G.
Kornhaber, J. F. Hunt, L. Tong, G. T. Montelione, D. Baker, Computational Design of a
Protein-Based Enzyme Inhibitor. *Journal of Molecular Biology*. **425**, 3563–3575 (2013).

116. D. M. Fowler, C. L. Araya, W. Gerard, S. Fields, Enrich: software for analysis of protein
function by enrichment and depletion of variants. *Bioinformatics*. **27**, 3430–3431 (2011).

117. S. Jo, T. Kim, V. G. Iyer, W. Im, CHARMM-GUI: a web-based graphical user interface for
CHARMM. *J Comput Chem*. **29**, 1859–1865 (2008).

118. W. L. Jorgensen, J. Chandrasekhar, J. D. Madura, R. W. Impey, M. L. Klein, Comparison
of simple potential functions for simulating liquid water. *J. Chem. Phys.* **79**, 926–935
(1983).

119. R. Salomon⬚Ferrer, D. A. Case, R. C. Walker, An overview of the Amber biomolecular
simulation package. *WIREs Computational Molecular Science*. **3**, 198–210 (2013).

120. J. A. Maier, C. Martinez, K. Kasavajhala, L. Wickstrom, K. E. Hauser, C. Simmerling, ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J Chem Theory Comput.* **11**, 3696–3713 (2015).

121. J. Wang, R. M. Wolf, J. W. Caldwell, P. A. Kollman, D. A. Case, Development and testing of a general amber force field. *J Comput Chem.* **25**, 1157–1174 (2004).

122. J. Wang, W. Wang, P. A. Kollman, D. A. Case, Automatic atom type and bond type perception in molecular mechanical calculations. *Journal of Molecular Graphics and Modelling.* **25**, 247–260 (2006).

123. C. W. Hopkins, S. Le Grand, R. C. Walker, A. E. Roitberg, Long-Time-Step Molecular Dynamics through Hydrogen Mass Repartitioning. *J. Chem. Theory Comput.* **11**, 1864–1874 (2015).

124. T. Darden, D. York, L. Pedersen, Particle mesh Ewald: An N☐log(N) method for Ewald sums in large systems. *J. Chem. Phys.* **98**, 10089–10092 (1993).

125. V. Kräutler, W. F. van Gunsteren, P. H. Hünenberger, A fast SHAKE algorithm to solve distance constraint equations for small molecules in molecular dynamics simulations. *Journal of Computational Chemistry.* **22**, 501–508 (2001).

126. G. R. Bowman, D. L. Ensign, V. S. Pande, Enhanced Modeling via Network Theory: Adaptive Sampling of Markov State Models. *J. Chem. Theory Comput.* **6**, 787–794 (2010).

127. M. I. Zimmerman, J. R. Porter, X. Sun, R. R. Silva, G. R. Bowman, Choice of Adaptive Sampling Strategy Impacts State Discovery, Transition Probabilities, and the Apparent Mechanism of Conformational Changes. *J. Chem. Theory Comput.* **14**, 5459–5475 (2018).

128. M. K. Scherer, B. Trendelkamp-Schroer, F. Paul, G. Pérez-Hernández, M. Hoffmann, N. Plattner, C. Wehmeyer, J.-H. Prinz, F. Noé, PyEMMA 2: A Software Package for Estimation, Validation, and Analysis of Markov Models. *J. Chem. Theory Comput.* **11**, 5525–5542 (2015).

129. J.-H. Prinz, H. Wu, M. Sarich, B. Keller, M. Senne, M. Held, J. D. Chodera, C. Schütte, F. Noé, Markov models of molecular kinetics: Generation and validation. *J. Chem. Phys.* **134**, 174105 (2011).

130. D. R. Roe, T. E. Cheatham, PTRAJ and CPPTRAJ: Software for Processing and Analysis of Molecular Dynamics Trajectory Data. *J. Chem. Theory Comput.* **9**, 3084–3095 (2013).

131. R. T. McGibbon, K. A. Beauchamp, M. P. Harrigan, C. Klein, J. M. Swails, C. X. Hernández, C. R. Schwantes, L.-P. Wang, T. J. Lane, V. S. Pande, MDTraj: A Modern Open Library for the Analysis of Molecular Dynamics Trajectories. *Biophysical Journal.* **109**, 1528–1532 (2015).

132. W. Humphrey, A. Dalke, K. Schulten, VMD: Visual molecular dynamics. *Journal of*
Molecular Graphics. **14**, 33–38 (1996).

133. H. Ashkenazy, E. Erez, E. Martz, T. Pupko, N. Ben-Tal, ConSurf 2010: calculating
evolutionary conservation in sequence and structure of proteins and nucleic acids. *Nucleic*
Acids Res. **38**, W529–W533 (2010).

134. T. Hessa, N. M. Meindl-Beinker, A. Bernsel, H. Kim, Y. Sato, M. Lerch-Bader, I. Nilsson,
S. H. White, G. von Heijne, Molecular code for transmembrane-helix recognition by the
Sec61 translocon. *Nature*. **450**, 1026–1030 (2007).

1209