Mutant alleles differentially shape cattle complex traits and fitness

Ruidong Xiang[1,2,*], Ed J. Breen[2], Sunduimijid Bolormaa[2], Christy J. Vander Jagt[2], Amanda J.

Chamberlain[2], Iona M. Macleod[2], Michael E. Goddard[1,2]

[1] *Faculty of Veterinary & Agricultural Science, The University of Melbourne, Parkville 3052,*

Victoria, Australia

[2] *Agriculture Victoria, AgriBio, Centre for AgriBiosciences, Bundoora, Victoria 3083, Australia.*

[*]Corresponding Author: ruidong.xiang@unimelb.edu.au

Abstract

Classical mutant alleles (MAs), with large effects on phenotype, tend to be deleterious to

traits and fitness. Is this the case for mutations with small effects? We infer MAs for 8

million sequence variants in 113k cattle and quantify the effects of MA on 37 complex traits.

Heterozygosity for variants at genomic sites conserved across 100 vertebrates increase

fertility, stature, and milk production, positively associating these traits with fitness. MAs

decrease stature and fat and protein concentration in milk, but increase gestation length and

somatic cell count in milk (the latter indicative of mastitis). However, the allele frequency of

MAs that decrease fat and protein concentration and stature and increase gestation length and

somatic cell count is lower than the allele frequency of MAs with the opposite effect. These

results suggest bias in the direction of effect of mutation (e.g. towards reduced protein in

milk), but selection operating to reduce the frequency of these MAs. Taken together, our

results imply two classes of genomic sites subject to long-term selection: sites conserved

across vertebrates show hybrid vigour while sites subject to less long-term selection show a

bias in mutation towards alleles that are selected against.

Introduction

Classical mutations, with a large effect on phenotype, tend to decrease fitness, decrease fitness-related traits and be partially recessive [1-3] (also see the 1st category of mutations defined in [3]). However, the majority of the genetic variance in complex traits is due to mutations of small effect. Do these small-effect mutations show the same characteristics as those classical large-effect mutations? A study in *E. coli* showd that mutations with small effect on fitness tend to be deleterious to protein function [4]. However, how mutations affect complex traits such as body size, health and fertility is unknown.

A better understanding of the consequence of mutations not only updates scientific knowledge but also has practical implications. Domestic cattle support humans with food, labour, clothing material and transportation. Today, there are over 4 billion cattle across the world and over ~900 million tonnes of dairy products have been produced annually for human consumption (http://www.fao.org/3/ca8341en/CA8341EN.pdf). When practicing genomic selection, which is widely used in animal breeding [5], it would be an advantage to know a priori whether mutations are more likely to increase or decrease traits of interest. In particular, if a trait is related to fitness, one might expect mutations to be deleterious [2,6]. Therefore the first objective of this study is to determine whether mutations, defined as the non-ancestral allele (also known as derived alleles) at segregating sites, tend to increase or decrease individual complex traits and whether this depends on the trait's association with fitness.

Traits that are related to fitness typically show inbreeding depression and heterosis caused by directional dominance. That is, fitness decreases with increased inbreeding due to increased homozygosity at loci with recessive deleterious alleles [7]. Conversely, fitness generally increases with heterozygosity [8]. Therefore, directional dominance can be used to link traits to fitness. Here, we introduce a method testing for directional dominance by estimating the

effect of heterozygosity at genomic sites on traits of cattle and use this method to identify

traits that are associated with fitness. Then, we classify traits showing directional dominance

as 'fitness-related traits'.

A likely cause of directional dominance is that mutations tend to be deleterious and partially

recessive. However, not all sites in the genome affecting a trait may show this pattern. Our

second objective is to test the hypothesis that sites, where the same allele has been conserved

across vertebrate evolution, are the most likely to show directional dominance. Therefore, we

consider conserved sites and other polymorphic sites in this analysis.

Cattle presents a unique opportunity for studying the effects of mutation. The cattle family

diverged from other artiodactyls up to 30 million years ago [9]. Modern cattle are derived from

at least two different subspecies of wild aurochs, i.e., *Bos primigenius primigenius* (Eurasian

aurochs) and *Bos primigenius namadicus* (Indian aurochs) which diverged up to 0.5 million

years ago [10-17]. Domestication of *Bos p. primigenius* led to the humpless *Bos taurus*

subspecies, which has evolved some highly productive breeds for agriculture, such as the

famous black-and-white Holstein breed with superior milk productivity. Besides natural

selection, dairy cattle breeds experienced very recent and intensive selection for milk

production traits [18,19] and stature [20]. Domestication of *Bos p. namadicus* gave rise to the

humped *Bos indicus* subspecies which evolved breeds with strong resistance to hot climates,

such as Brahman and Gir cattle.

In the present study, we use yak, sheep and camel as outgroup species to assign cattle

ancestral alleles for 8M sequence variants (at 8M genomic sites). For each of these variants,

the alternative to the ancestral allele is the mutant allele (MA). We estimate the effect of the

mutant allele at these 8M variable sites on 37 traits of 113k cattle from 4 breeds. We also

estimate the effect of heterozygosity on these traits using both conserved sites and all

genomic sites.

If mutant alleles decrease fitness we expect selection to reduce their allele

frequencycompared with mutant alleles that either have no effect or increase fitness.

Therefore, we compare the allele frequency of mutant alleles that increase and decrease each

trait. We expand the analysis of mutant allele frequency to additional breeds of ancient and

modern cattle from the 1000 Bull Genomes database [21,22], which provides validation of our

results. Additional analyses of MAs with strong effects on milk production traits [23,24] suggests

that the direction of phenotypic effects of these MAs correlates with their direction of effects

on the expression of genes in milk cells[4,25].

Results

Directional dominance at sites conserved across 100 vertebrates

To identify traits related to fitness, we have introduced a method to estimate the effect of

heterozygosity on 37 traits (described in Supplementary Table 1) recorded in over 100k

animals. In total, there were 16,035,443 imputed sequence variants (at 16,035,443 genomic

sites) with imputation accuracy $R^2 > 0.4$ and the minor allele frequency (MAF) > 0.005

available for variant-trait association analysis. A subset of these sequence variants that could

be assigned with ancestral alleles was used for analyses related to mutant alleles (described

later). For the analysis of the effect of heterozygosity, we fit the average heterozygosity of

sequence variants at 317,279 genomic sites conserved across 100 vertebrates (H'_{cons_j}) and

heterozygosity from variants at the other 15,718,164 sites ($H'_{non-cons_j}$) simultaneously (see

Methods). We observed a significant effect of heterozygosity at consserved sites for the yield

of protein (Prot), fat (Fat) and milk (Milk), survival (Surv), fertility performance (Fert),

stature (Stat) and angularity (related to slimness and milk yield) (Figure 1 and Supplementary

Figure 1). For all these traits, heterozygosity at other sites ($H'_{non-cons_j}$) was not significant

when fitted together with H'_{cons_j}. This directional dominance implies that milk production,

fertility, survival and stature show inbreeding depression and heterosis and therefore we

classify them as fitness-related traits and this directional dominance for these traits is

exclusively explained by genomic sites conserved across vertebrates. To be conserved across

vertebrates, mutations at these sites must be deleterious, implying extremely long-term

consistent selection for the ancestral allele at these sites.

Figure 1. Directional dominance at conserved sites (*H'*) for traits of 104k cows. The beta

values and standard errors for each trait were generated using a mixed linear model, fitting *H'*

from 317,279 conserved sites (left panel) and *H'* from the remaining 15,718,164 sites (right

panel) together with other fixed effects (e.g., breed). Blue dashed lines indicate t value of -

1.96 and 1.96 commonly used to indicate the significance.

Assignment of bovine ancestral and mutant alleles

To assign the mutant alleles in cattle, we first determined the alternative, ancestral alleles

using artiodactyls, including cattle as the focal species (98 global cattle breeds from the 1000

Bull Genomes Project [21,22], Supplementary Table 2) and yak, sheep and camel as outgroup

ancestor species (Ensembl 46-mammal sequence data). A probabilistic method [25] was used to

assign an ancestral allele for each site mappable between 4 artiodactyl species (see Methods).

Out of 42,573,455 equivalent sites between the 4 species, 39,998,084 sites had the ancestral

allele assigned with high confidence (probability > 0.8). We compared our results with a

previous study using different methods [26]. Of 1,925,328 sites that were assigned ancestral

alleles with high confidence in both studies, 1,904,598 (98.7%) sites agreed. However, we

have assigned ancestral alleles with high confidence to ~10 times more sites than the previous

study due to the use of large sample size and whole-genome sequence data. The full results

are publicly available at https://figshare.com/s/dd5985b76a413b56106b.

Biases in trait effects between ancestral and mutant alleles

We conducted GWAS of 37 traits using over 16 million imputed sequence variants in bulls

(N ~ 9k) and cows (N ~ 104k) separately (see Methods). For 7,910,190 variants where the

ancestral allele was assigned, we compared the direction (increase or decrease) of the effect

of the mutant alleles (MAs) on the trait (Supplementary Figure 2-3). The same comparison

was also performed for variants at the 202,530 out of 317,279 conserved sites where the

ancestral alleles could be assigned. Note that for a variant, the effect of a MA is identical to -

$1 \times$ the effect of the ancestral allele. We focus the description of effects on MAs, but a MA

increasing the trait is identical to an ancestral allele decreasing the trait.

Within all analysed variants and conserved variants, for each trait we considered the

following three variant categories for systematic comparison: 1) large-effect variants, i.e., p-

value of GWAS (p_{gwas}) < 5e-8 and the effect direction agreed in both sexes; 2) medium-

effect variants, i.e., 5e-8 <= p_{gwas} < 5e-5 and the effect direction agreed in both sexes, and 3)

small-effect variants, i.e., 5e-5 <= p_{gwas} < 0.05 and the effect direction agreed in both sexes.

Here the effect size refers to the amount of variance explained by variants which is inversely

related to the p-value. The use of different effect size is because mutations of small and large

effects may be different in their direction of effect. Selecting variants that have the same

effect direction between independent GWAS populations [27], such as bulls and cows, helps to

eliminate variants with spurious trait associations from the comparison. Based on a previous

method [27], the True Discovery Rate by Effect Direction (TDRed) of GWAS between two

sexes across 37 analysed traits for the small-, medium- and large-effect variants was 0.8, 0.98

and 0.99, respectively.

Based on GWAS results of each trait, we calculated the ratio of the number of variants where

the MA increased the trait (positive effect) to the number of variants where the MA decreased

the trait (negative effect). Across 37 traits and three effect-size groups, MAs showed diverse

trait effect patterns (Supplementary Figure 3). Results observed from GWAS were confirmed

by BayesR analysis [28], which jointly fits on average 4.3 million variants per trait (See

methods and Supplementary Figure 3). Based on jointly estimated effects for a given set of

variants, the significance of the effect direction bias was tested using Kolmogorov-Smirnov

to estimate the p-value (p_{ks}) of the difference in the effect distribution between ancestral and

mutant alleles (see Methods). We also tested the significance of bias using LD-clumped ($r^2 <$

0.3) [29] variants to calculate the standard error (Supplementary Figure 4).

In addition, we checked the direction of effects of MAs which had large positive effects and

large negative effects on protein yield, fat yield, milk yield, protein % and fat % on the

expression of genes within ±1Mb distance to these MAs (cis eQTL genes, see Methods) in

milk cells[23,24]. For 4 out of 5 sets of variants where the mutant allele decreased the trait, we

found the mutant allele tended to decrease the expression of cis eQTL genes. For another 4

out of 5 sets of variants where mutant alleles increased the trait, the mutant allele tended to

increase the expression of cis eQTL genes (Supplementary Table 3). These results suggest

correlated direction of effects of MAs on milk production traits and the expression of genes

in milk cells.

In the following text, we focus on 1) MAs within the large- and small-effect categories for

milk production traits as these two sets of MAs showed distinct effect direction patterns

(Figure 2), and 2) MAs associated with other traits, including those with medium or small

effects on somatic cell count (Scc, indicative of mastitis, medium-effect), survival (Surv,

small-effect), fertility (Fert, frequency of pregnancy, small-effect), gestation length (Gl,

medium effect), temperament (Temp, docility, small-effect) and stature (Stat, medium effect)

(Figure 2).

Figure 2. The ratio (y-axis) between the number of variants with mutant alleles increasing
the trait (+) and the number of variants with mutant alleles decreasing the trait (−). GWAS
effects of mutant alleles are shown for all variants (**a**). BayesR joint effects of mutant alleles
from the same variants in (**a**) are shown for all variants (**b**). Pink colour: the majority of
variants with mutant alleles tend to increase the trait (taller than the blue-dashed line). Dark
grey: the majority of variants with mutant alleles tend to decrease the trait (shorter than the
blue-dashed line). Numbers in bars: total number of variants significant at the given
threshold. Stars: p-value for the significance of the difference in the distribution of BayesR
effects between ancestral and mutant alleles, '*': p < 0.05, '**': p < 0.01, '***' p < 0.001.

For somatic cell count (Scc), gestation length (Gl) and stature (Stat), the results are from

medium-effect (M) variants and the full results are shown in Supplementary Figure 3.

The classical model [1-3] predicts that the majority of MAs, or mutations, are deleterious or

slightly deleterious. In our study, MAs consistently showed biases towards decreasing protein

and fat concentration (Figure 2 and Supplementary Figure 3,4), docility and stature, and

towards increasing somatic cell count (an indicator of mastitis) and gestation length. Among

these traits only stature showed a significant effect of heterozygosity. For milk yield and

protein yield, both of which were classified as fitness-related traits (Figure 1), the bias in the

direction of MA depends on the size of the MA effect. Large-effect MAs tended to decrease

milk and protein yield whereas small-effect MAs tended to increase them. A possible

explanation is that mutation seldom has a large positive effect on milk protein yield or

fertility but small positive effect mutations occur and are increased in frequency by natural or

artificial selection.

Also, there was a slight majority of small-effect MAs which tended to increase fertility and

survival, both of which were positively related to fitness (Figure 1). The effects of these sets

of MAs is partially due to pleiotropy, i.e., the effect of these MAs on multiple traits

(Supplementary Table 4). For instance, while small-effect MAs increasing milk yield

decreased fat yield, protein % and fat %, they also increased protein yield. Also, while small-

effect MAs increasing fertility increased gestation length, they also increased stature.

The simplest explanation for the bias in the direction of MA effects is that it is due to a bias

in the direction of mutation. For instance, that mutation more often leads to a decrease in

fat% rather than an increase. However, it is also possible that mutations that decrease fat%

are selected and therefore more likely to be discovered than mutations that increase fat%.

Below we exclude this possibility by comparing the allele frequency at variants where the

MA increases or decreases the trait.

Allele frequency of mutant alleles in modern and ancient cattle

Across all variable sites, the allele frequency of MAs was lower than the allele frequency of

ancestral alleles (Supplementary Figure 5). Also, the frequency of MAs at conserved sites

(0.27) was lower than the frequency of MAs across all sites (0.32). This is consistent with the

selection for the ancestral allele which is necessary to maintain conservation of the same

allele across vertebrates.

We grouped variants based on their mutant allele reducing (MAs−) or increasing the trait

(MAs+) and compared their allele frequency in over 110k Holstein, Jersey, crossbreds and

Australian Red bulls and cows (Figure 3a,b). To account for LD, we estimated the error of

MA frequency based on LD-clumped ($r^2 < 0.3$) [29] variants. As an external validation, we also

considered this analysis in a selection of 7 subspecies/breeds of 1,720 ancient and modern

cattle from the 1000 Bull Genomes Project [21,22] (Figure 3c,d).

Figure 3. The allele frequency of mutant alleles (MAs) in cattle. The average frequency of

variants associated with different traits is shown with standard error bars based on LD

clumped variants. All variants include the 7.9M variants where mutant alleles were assigned.

Red dashed line represents the frequency of 0.5. In the dairy cattle section (**a** and **b**), 90,627

Holstein, 13,465 Jersey, 3,358 Australian Red (AusRed) and 4,649 crossbreds were used. In

the ancient and modern cattle (**c** and **d**), 210 Brahman, 25 Tibetan, 10 Eurasian Aurochs, 242

Simmental, 95 Jersey, 840 Holstein and 287 Angus were used. For panels **b** and **d**, results for

survival (Surv), fertility (Fert) and temperament (Temp) were from small-effect MAs while

results for somatic cell count (Scc), gestation length (Gl) and stature (Stat) were from

medium-effect MAs.

For fat%, protein%, docility and stature MAs that increase the trait had higher allele

frequency than MAs that decrease the trait. For somatic cell count and gestation length, the

reverse is true. That is, MAs increasing somatic cell count and gestation length had lower

allele frequency than MAs that decreased the trait (Figure 3). Thus, although MAs more

commonly decreased fat% than increased it, the allele frequency was higher at sites where the

MAs increased fat%. This implies that selection acts against MAs that decrease fat% or

favours MAs that increase fat%. Consequently, the higher incidence of MAs that decrease

fat% cannot be due to selection favouring them but must be due to mutation more often

resulting in an allele that decreases fat% than increases it. Comparing results in Figure 2 and

3 shows that this is the usual pattern – the more common direction of effects of mutation

generates alleles that are selected against and hence have a reduced allele frequency.

For other traits, the results are less clear-cut. For milk yield, the majority of MAs of large

effect tended to decrease the trait (Figure 2). Interestingly, these large-effect milk-decreasing

MAs, which were deleterious, had a higher frequency than those MAs increasing milk yield

(Figure 3). On the other hand, the majority of MAs of small effect tended to increase the milk

yield (Figure 2). Yet, these small-effect MAs that increase milk were at a lower frequency

than MAs that decrease milk yield (Figure 3). Interpretation of these results is helped by

remembering that milk yield is negatively correlated with fat% and protein% (Supplementary

Table 4).

Selection of trait-associated mutant alleles in modern and ancient cattle

The above results for MA frequency at trait-associated variants imply selection. The selection

could be consistent across breeds which would limit the divergence of allele frequency

between breeds or it could be different between breeds leading to divergence in allele

frequency. We compared the average of Wright's fixation index ($\overline{F_{ST}}$), for MA+ variants and

MA− variants calculated using dairy cattle (Figure 4a,b) and ancient and modern cattle

(Figure 4c,d). To account for LD, we estimated the error of $\overline{F_{ST}}$ based on LD-clumped ($r^2 <$

0.3) [29] variants.

Figure 4. Selection (average Wright's fixation index $\overline{F_{ST}}$) of variants with mutant alleles that

increase or decrease the trait in dairy cattle (**a,b**) and ancient and modern cattle (**c,d**). The $\overline{F_{ST}}$

is shown as dots with its standard error bars estimated using LD clumped variants. The blue

line represents the $\overline{F_{ST}}$ for 7.9M variants analysed (0.1±4.3e-05) in dairy cattle in **a** and **b**; and

$\overline{F_{ST}} = 0.157±5e-05$ in ancient and modern cattle in **c** and **d**). For panels **b** and **d**, results for

survival (Surv), fertility (Fert) and temperament (Temp) were from small-effect MAs while

results for somatic cell count (Scc), gestation length (Gl) and stature (Stat) were from

medium-effect variants.

In general, variants associated with milk production traits (including somatic cell count,

Figure 4a) showed higher than average F_{ST} among dairy breeds implying divergent selection,

while variants associated with other traits, including survival and fertility, tended to have

below-average F_{ST} indicating convergent selection (Figure 4b). $\overline{F_{ST}}$ for gestation length was

below average especially for MA+, probably due to selection against mutations that increase

gestation length in all breeds (Figure 4d).

Among ancient and modern cattle, $\overline{F_{ST}}$ is high for both MA+ and MA− variants for stature

indicating divergent selection for height (Figure 4b). The allele frequency of MAs decreasing

height was the least frequent in Holstein cattle and was the most frequent in Tibetan cattle

living at high altitude and Angus cattle selected for beef production (Figure 3d). This

suggests that the direction of selection could vary across cattle breeds under different

environmental conditions and/or artificial selection.

Discussion

For some traits (e.g. survival, fertility) we expect that an increase in the trait leads to an

increase in fitness. It is these traits which typically show heterosis and inbreeding depression

due to directional dominance. The simplest explanation for these observations is that

mutations at sites affecting the trait tend to reduce the trait and be partially recessive.

However, our results show that it is not all sites affecting these traits that show directional

dominance but only those where the same allele is highly conserved across vertebrates. This

result explains why the mutations tend to lead to a decrease in the trait - long-term selection

has nearly fixed the favourable allele and so any mutation will cause a decrease in the trait

and in fitness. We partially confirm this explanation by finding that mutations for these traits

(milk and protein yield, stature but not fertility and survival) do tend to decrease the trait

although, for milk and protein yield, it is only mutations of large effect for which the effects

tend to be negative. This long term selection cannot be directly on traits involving lactation

since the same allele is conserved in vertebrates other than mammals.

For other traits we expect that an intermediate value leads to the highest fitness. For instance,

too high or too low a fat% in milk might be detrimental to the fitness of the mother or the

infant or both. These traits do not typically show inbreeding depression or heterosis. The

fittest allele might vary between species and environments. Therefore one might expect that

mutations are equally likely to increase or decrease the trait. However, that is not what we

found: for fat% and protein% mutations tend to decrease the trait whereas for SCC and

gestation length they tend to increase the trait. We hypothesise that at some of the genomic

sites affecting these traits selection has been consistent enough in mammals, or at least in

cattle, so that mutations cause a decrease in fat% and protein% and an increase in mastitis or

SCC and gestation length (leading to difficulty calving). This hypothesis is supported by our

finding that selection decreases the allele frequency of these mutations. This low allele

frequency is not only seen in dairy cattle but in beef breeds and *Bos indicus* breeds.

The findings on individual traits can be unexpected due to pleiotropy. That is, mutations

affect multiple traits. There are mutations at *DGAT1* and *GHR* loci that increase milk yield

but decrease fat% and protein% (Supplementary Figure 6). These are only at appreciable

frequency in domesticated cattle, especially breeds artificially selected for milk volume.

Their low allele frequency in other breeds and species suggest that natural selection acts

against the mutation thus increasing fat% and protein% but decreasing milk yield. Similarly,

there is a negative genetic correlation between milk yield and fertility so mutations that

increase milk yield might be favoured despite their negative effect on fertility. MAs

decreasing fertility tended to be most frequent in the Holstein breed (Figure 3d), perhaps

because these alleles tended to increase milk yield and stature.

For milk, fat and protein yield the results differ between mutations of large and small effects.

Mutations with a large effect on milk protein yield more often decrease protein yield than

increase it perhaps because the physiology supporting milk protein synthesis has been

optimised in part at least. Mutations with a small effect on protein yield are almost equally

likely to increase or decrease yield perhaps because natural selection favours an intermediate

level of milk protein yield because too high a yield drains the cow of nutrients needed for

maintenance and reproduction.

Effects of MAs on phenotypes might be mediated by their effects on gene expression. Based

on cis eQTL data [23], we found that MAs with large effects on milk production traits had

direction of effects that were correlated with their direction of effects on gene expression in

milk cells. This result shows that the effect direction of MAs on gene expression may also

show systematic biases and this may be related to their effects on phenotypic traits. Future

studies with larger sample size and more tissues for eQTL mapping may update our

understanding of the MA effects on molecular phenotypes.

The selection which we have observed affecting the frequency of mutations of positive and

negative effect could be both natural selection acting over a long period before and since the

domestication of cattle, and artificial selection acting over the last 10,000 years and, more

intensely, over the last ~100 years in dairy cattle. Artificial selection may differ between

breeds and generate high Fst between breeds. For fat%, protein% and stature at least one

class of mutation is more common than random mutations and the overall F_{ST} between breeds

tended to be high. Our analysis also highlighted some specific breeds. For example, the

selection of variants associated with somatic cell count led to high F_{ST} among dairy cattle but

low F_{ST} in our other breeds. Holstein cattle have been selected to be tall [20] and this is

reflected in the low frequency of MAs decreasing stature in Holstein. On the other hand, the

high frequency of MAs decreasing stature in Tibetan cattle (Figure 4d) may be due to its

adaptation to high altitude [30].

Although mutation is biased in its effect on some traits, the bias is small for most traits. That

is, mutations decreasing protein yield are only slightly more common than mutations that

increase protein yield. Also, although conserved sites explain directional dominance and are

enriched for polymorphisms affecting complex traits [31], they do not explain the majority of

the genetic variance. That is, there are many sites affecting traits, such as milk yield and

stature, at which the allele carried varies between species implying that the fittest allele varies

depending on the environment and the background genotype of the species.

The sequence variants associated with a complex trait are not necessarily causal but likely to

be in high LD with the causal variants. This tends to dilute the signal that might be

discovered if causal variants were used. However, variants in high LD may share a similar

evolutionary history and therefore show some of the same characteristics. We used BayesR

which jointly fits variants and LD-clumping to account for LD. However, we acknowledge

that we cannot completely remove the effects of LD on our results. Therefore, future studies

with even larger sample sizes, e.g., ~1 million, may update our results.

Genomic selection [32], used in the breeding of livestock and crops, estimates the genetic value

of individuals for traits of interest from the alleles they carry at genetic markers such as

SNPs. The equation predicting genetic value uses the effect of each SNP on the trait

estimated in a training population. The best methods treat the SNP effects as random

variables drawn from a prior distribution. To date it has been assumed that the effects of a

mutation are equally likely to be positive or negative on the trait but, if it was known that one

direction of effect was more likely, this could be built into the prior distribution resulting in

an increase in the accuracy with which genetic value is predicted.

In conclusion, our results support a new hypothesis which provides a new picture of the

effects of mutation and selection on mammalian complex traits. Directional dominance,

which causes heterosis and inbreeding depression, is characteristic of loci where mutations

decrease the trait and fitness and this pattern has been consistent over the evolution of

vertebrates. More recent selection, although not causing directional dominance, leads to a

bias in the direction of mutation because the mutation results in an allele which is less fit than

the ancestral allele and tends to affect a complex trait in a consistent direction. This

hypothesis, if supported by future research, adds to our understanding of the evolution of

complex traits and has practical value in the artificial selection of livestock and crops.

Methods

Data preparation for calling bovine ancestral alleles. The assignment of bovine ancestral

alleles was based on a model comparison of alleles from cattle with alleles from outgroups of

yak (*Bos grunniens*), sheep (*Ovis aries*) and camel (*Camelus dromedarius*). According to the

evolutionary relationships reported previously [9], among ruminants, yak is an outgroup species

closely related to cattle, while sheep is less closely related to cattle than yak. Goat is

equivalent to sheep in its relationship to cattle, but we chose sheep in the current study.

Camel without the rumen is distantly related to cattle as they are artiodactyls. For the cattle

species, we used whole-genome sequence data of 98 individuals from Run 7 of the 1000 Bull

Genomes Project [21,22]. Each of those above mentioned 98 individuals represents a breed

collected by the consortium. Only those whole-genome sequence samples with coverage >

10x were selected and if multiple individuals were found for a breed, the whole-genome

sequence sample with the highest coverage was chosen. Both *Bos taurus* and *Bos indicus*

subspecies were included (Supplementary Table 2). The pre-processing of sequence reads

and alignment of sequence data is done by project partners using the standard 1000 Bull

Genomes Project pipeline: http://www.1000bullgenomes.com/. Only BAM files from 1000

Bull Genomes partners are collected and processed by the consortium. The latest published

data from the 1000 Bull Genomes Project (1832 samples) can be found at

https://www.ebi.ac.uk/eva/?eva-study=PRJEB42783. The details of variant calling can be

found in [33]. Briefly, Genome Analysis Toolkit (GATK v.3.8) [34] was used for variant calling.

Variants from the GATK VQSR (Variant Quality Score Recalibration) 99.90 to 100.00

Tranche for SNP and INDEL were excluded, and Beagle v.4.0 [35] was used to impute sporadic

missing. Whole-genome sequence data in VCF format for these 98 cattle, as a subset from

the 1000 Bull Genomes Project database, was generated for further analysis.

For the outgroup species (to determine ancestral alleles), we used whole-genome sequence

data of 46 mammals stored in the Multiple Alignment File generated by Ensembl EPO

pipeline (http://asia.ensembl.org/info/genome/compara/multiple_genome_alignments.html). The

46-mammal EPO Multiple Alignment File was downloaded. Then, the software WGAbed

(https://henryjuho.github.io/WGAbed/) from python v2.7 was used to retrieve sequence data

for cattle, yak, sheep and camel in bed file format. Only sites with sequence data available in

at least one outgroup species were kept. Using the cattle coordinates in the 4-species

WGAbed files, the sequence data of the outgroup species were matched with the 98 cattle. As

a result, 42,573,455 sites found in the 98 cattle and in at least one outgroup species were

found. Sequence data on these 42,573,455 sites across 4 species were used to determine the

bovine ancestral alleles.

Probabilistic determination of bovine ancestral alleles. We used the method proposed by

Keightley et al [25] with the model choice of The Kimura two-parameter (K2) which accounts

for allele frequency of the focal species to determine the probability of an allele being

ancestral at each available site. The method was implemented in estsfs [25] and the K2 model

was chosen due to its equivalent accuracy to other models but better computation efficiency.

As described above, the sequence data of three outgroup species were used. The order of

phylogenetic tree topology was cattle → yak → sheep → camel. As requested by the

software, allele counts of A, C, G and T were determined for the focal species (cattle) and for

out species at each available site. For cattle, the total allele count for each site was 196 (98

×2). For each outgroup species, the total allele count for each site was up to 1. Missing

sequence data in the outgroup species were treated as 0 counts. For each site, estsfs produced

a probability (P_{ancs}) of the major allele in the focal species being ancestral. We then

determined alleles which were major at a site with $P_{ancs} > 0.8$ or those alleles which were

minor at a site with $P_{ancs} < 0.2$ to be ancestral. For those sites where the major or minor

alleles could not be determined but the $P_{ancs} > 0.8$ or < 0.2, the cattle allele with the highest

frequency in the 3 out species was assigned ancestral. The rest of the sites were determined as

ambiguous where no clear ancestral alleles could be determined. The detailed results of

ancestral alleles for those 42,573,455 sites across 4 species and the probability of the alleles

being ancestral or ambiguous is publicly available at:

https://figshare.com/s/dd5985b76a413b56106b.

Sequence variants under conserved sites across 100 vertebrate species. The variant

selection followed previous procedures [31]. Briefly, conservation was determined by the

criteria of PhastCon score [36] > 0.9 based on the sequence data of those 100 species. The

choice of 0.9 as the cutoff was arbitrary. However, since PhastCons score ranges from 0 to 1,

this cutoff kept relatively highly conserved sites. Also, in a previous study [31], cattle variants

from sites with PhastCon score > 0.9 were highly enriched for the heritability of cattle traits.

The conserved sites were primarily determined using the human genome coordinates (hg38)

and were lifted over to the bovine genome ARS-UCD1.2 using the LiftOver software

(https://genome.ucsc.edu/cgi-bin/hgLiftOver) with a lift-over rate $> 92\%$. In total, 317,279

variants in the current study were assigned as the conserved variants.

Animals and phenotypes for variant-trait association analysis. Data was collected by

farmers and processed by DataGene Australia (http://www.datagene.com.au/) for the official

May 2020 release of National breeding values. No live animal experimentation was required.

Phenotype data was based on trait deviations for cows and daughter trait deviations for bulls.

Daughter trait deviations were the average trait deviations of a bull's daughters and all

phenotypes were pre-corrected for known fixed effects, with processing done by DataGene.

Phenotype data used included a total of 8,949 bulls and 103,350 cows from DataGene.

Holstein (6,886♂ / 87,003♀), Jersey (1562♂ / 13,353♀), cross-breed (36♂ / 5,037♀) and

Australian Red dairy (265♂ / 3,379♀) breeds were included. In total, 37 traits related to milk

production, mastitis, fertility, temperament and body conformation were studied

(Supplementary Table 1). Larger trait values of fertility (Fert), ease of birth (Ease),

temperament (Temp), milking speed (MSpeed), likeability (Like) meant poor performances,

so to assist the interpretability of the study, we have corrected the trait direction so that larger

values of Fert, Ease, Temp, MSpeed and Like meant increased fertility performance (calving

frequency), labour ease, docility, milking speed and the overall preference as a dairy cow

(Supplementary Table 1). This correction only affected the reported effect direction of the

mutant allele.

Genotype data for association analysis. The genotypes used in the current study included a

total of 16,035,443 imputed bi-allelic sequence variants with Minimac3 [37,38] imputation

accuracy $R^2 > 0.4$ and the minor allele frequency (MAF) > 0.005 in both sexes. Most bulls

were genotyped with a medium-density SNP array (50K: BovineSNP50 Beadchip, Illumina

Inc) or a high-density SNP array (HD: BovineHD BeadChip, Illumina Inc) and most cows

were genotyped with a low-density panel of approximately 6.9k SNPs overlapping with the

standard-50K panel. The low-density genotypes were first imputed to the Standard-50K panel

and then all 50K genotypes were imputed to the HD panel using Fimpute v3 [31,39]. Prior to

sequence imputation, the HD genotypes were converted to forward sequence format. Then,

all HD genotypes were imputed to sequence using Minimac3 with Eagle (v2) to pre-phase

genotypes ([38,40]). The reference set for imputation included sequences of 3090 *Bos taurus*

animals from Run7 of the 1000 Bull Genomes Project [21] aligned to the ARS-UCD1.2

reference bovine genome (https://www.ncbi.nlm.nih.gov/assembly/GCF_002263795.1/) [22,41].

The accuracy of the sequence data for individual animals in the 1000 Bull Genomes Project is

routinely checked against their own high-density SNP array genotypes and the concordance

has been above 95% [33]. The empirical accuracy of imputation to sequence using the 1000

Bull Genomes project has been routinely tested for dairy breeds: for example, in Holsteins

the average correlation between imputed and real sequence variants was 0.92 to 0.95 using

Run5 of the 1000 Bull Genomes project (N= 1577)[42]. Therefore, we believe our imputed data

is more accurate: first because the number of reference animals has almost doubled and

second because in our study we impose a Minimac3 R^2 filter to remove poorly imputed

variants. A Minimac3 R^2 threshold of 0.4 was used because our in-house tests demonstrate

that this is approximately equivalent to an empirical imputation accuracy (correlation) of

0.85.

Genome-wide association studies. The above mentioned traits were analysed one trait at a

time independently in each sex with linear mixed models using GCTA [43]:

$$\mathbf{y} = \mathbf{mean} + \mathbf{breed} + \mathbf{bx} + \mathbf{a} + \mathbf{error} \quad (equation\ 1)$$

where $\mathbf{y}$ = vector of phenotypes for bulls or cows, $\mathbf{breed}$ = three breeds for bulls, Holstein,

Jersey and Australian Red and four breeds for cows (Holstein, Jersey, Australian Red and

MIX); $\mathbf{bx}$ = regression coefficient b on variant genotypes $\mathbf{x}$; $\mathbf{a}$ = random polygenic effects

$\sim N(0, \mathbf{G}\sigma_g^2)$ where $\mathbf{G}$ = genomic relatedness matrix based on all variants and σ_g^2 = random

polygenic variance; $\mathbf{error}$ = the vector of random residual effects $\sim N(0, \mathbf{I}\sigma_e^2)$, where I =

the identity matrix and σ_e^2 the residual variance.The purpose of fitting breeds as fixed

effects together with the GRM in the model was to have strong control of the population

structure which may cause spurious associations between variants and phenotype. The

construction of GRM followed the default setting (--make-grm) in GCTA[43]:

(https://cnsgenomics.com/software/gcta/#MakingaGRM).

Bayesian mixture model analysis. In the above-described GWAS, sequence variants, many

of which are in high LD, were analysed one at a time. In order to assess variant effects and

account for LD, we fitted selected variants jointly in BayesR [28]. For each trait, variants that

showed the same sign between bulls and cows (regardless of p-value) and could be assigned

with an ancestral allele were analysed with BayesR. Across 37 traits, the number of variants

analysed ranged from 3,961,180 to 4,737,492. To reduce the computational burden of

BayesR, we estimated the joint effects of these variants for each trait in bulls. BayesR models

the variant effects as mixture distribution of four normal distributions including a null

distribution, $N(0, 0.0\sigma^2{}_g)$, and three others: $N(0, 0.0001\sigma^2{}_g)$, $N(0, 0.001\sigma^2{}_g)$,

$N(0, 0.01\sigma^2{}_g)$, where $\sigma^2{}_g$ was the additive genetic variance for the trait. The starting value

of $\sigma^2{}_g$ for each trait was estimated using GREML implemented in MTG2 [44] with a single

genomic relationship matrix made of all 16M sequence variants. The statistical model used in

the single-trait BayesR was:

$$\mathbf{y} = \mathbf{Wv} + \mathbf{Xb} + \mathbf{e} \text{ (equation 2)}$$

where $\mathbf{y}$ was a vector of phenotypic records; $\mathbf{W}$ was the design matrix of marker genotypes;

centred and standardised to have a unit variance; $\mathbf{v}$ was the vector of variant effects,

distributed as a mixture of the four distributions as described above; $\mathbf{X}$ was the design matrix

allocating phenotypes to fixed effects; $\mathbf{b}$ was the vector of fixed effects of breeds; $\mathbf{e}$ = vector

of residual errors. As a result, the effect v for each variant jointly estimated with other

variants were obtained for further analysis.

The difference in effect distribution between ancestral and mutant alleles. For an

analysed variant, one allele is ancestral and then the other is mutant. If there is a bias in effect

direction in ancestral alleles or mutant alleles in a given set of variants, the effect distribution

of the ancestral and mutant alleles would be different. We tested if the distribution of the

effect of ancestral alleles estimated from BayesR was significantly different from that of

mutant alleles using the two-sample Kolmogorov-Smirnov test implemented by ks.test() in R

v3.6.1. The coding was ks.test(a,m) where a was the vector of variant effects based on the

ancestral alleles and m was a vector of variant effects based on the mutant alleles. To be more

conservative, we also tested the significance of biases using LD-clumped ($r^2 < 0.3$ within

1Mb windows) variants with small, medium and large effects using default settings in

plink1.9 [29].

Heterozygosity of individuals at conserved sites. It is widely accepted that higher genomic

heterozygosity is linked to gene diversity, therefore, fitness. However, it is not clear at which

set of genes or variants heterozygosity is more related to fitness. Also, the simple estimation

of heterozygosity, i.e., assigning allele counts of 0 or 2 as homozygous and 1 as

heterozygous, leads to biases as the estimation is not independent of additive effects

(illustrated later). Our previous work showed conserved sites across 100 vertebrate species

significantly contribute to trait variation [31,45] and it is also logical to assume mutations at

conserved sites tend to have strong effects on fitness. Therefore, we firstly partitioned the

genome into 317,279 conserved and 15,718,164 non-conserved variants. Then, we re-

parameterised the genotype allele count for each variant commonly used to model the

dominance deviation, so that the estimation of dominance deviation is independent of the

additive effects. We focused on cows because their traits were largely measured on

themselves, contrasting to bull traits which were based on their daughters' traits. We

estimated the variant-wise sum of the re-parameterised allele count value for dominance

deviation which was later termed as z'_{D_i} for each variant i in cows. The sum was averaged by

the number of variants and this average value based on re-parameterised dominance allele

count for the individual j was termed as H'_j to represent the individual heterozygosity. We

estimated the individual heterozygosity from conserved sites (H'_{cons_j}) and non-conserved

sites ($H'_{non-cons_j}$) and these computations are specified in the following text.

According to quantitative genetics theory [46-48], the genetic value (G') of an individual can be

partitioned into the mean (μ), additive genetic value (A) arising from additive effect (a) and

dominance genetic value (D) arising from dominance deviation (d). At a single locus, let the

allele frequency of the three genotype classes of AA, AB and BB be p^2, $2pq$ and q^2,

respectively. In a simple genetic model, the genetic value can be decomposed as:

$$G' = \mu + A + D + e = \mu + x_{A_i}a + z_{D_i}d + e \text{ (equation 3)}$$

Where x_{A_i} was the allele count for genotype AA, AB and BB for locus or variant i which

were usually coded as 0, 1, 2, respectively, to represent the additive component, and z_{D_i} was

usually coded as 0, 1, 0, for genotype AA, AB and BB for variant i, respectively, which

differentiates the homozygous and heterozygous to represent the dominance component.

Therefore, in the simplest form, the genome-wide heterozygosity of the individual j can be

calculated as:

$$H_j = \left.\sum_i^N z_{D_i} \middle/ N\right. \text{ (equation 4)}$$

where H_j is the simple genome-wide heterozygosity of individual j, N is the total number of

variants. Note that such calculation of H_j can also be used to derive inbreeding coefficient,

where $I_j = (\sum_i^N 2p_i q_i) \times H_j$. I_j was the inbreeding coefficient for the j$_{th}$ individual.

In equation 3, however, due to the non-zero correlation between x_A and z_D under Hardy-

Weinberg equilibrium (HWE), the estimation of a and d is not independent, i.e.,

$cov(x_A, z_D) = 2p(1-p)(1-2p) \neq 0$ under HWE. This then resulted in the estimation of

H_j not being independent of the additive components. Therefore, we proposed to re-

parameterise this model to estimate a and d independently.

According to Falconer [47] at this locus, the additive effects can be derived using the regression

of genetic value on the number of A alleles, where $A'_{AA} = 2q \times \alpha$, $A'_{AB} = (p-q) \times \alpha$ and

$A'_{BB} = -2p \times \alpha$. A' is the re-parameterised additive genetic value and α is the allele

substitution effect: $\alpha = a + (p-q)d$. Because the dominance deviation is the difference

between the genetic value and the mean plus the additive value, the dominance effects can be

derived as $D'_{AA} = -2p^2 \times d$, $D'_{AB} = 2pq \times d$ and $D'_{BB} = -2q^2 \times d$. D' is the re-

parameterised dominance genetic value. Therefore, equation 3 can be re-parameterised as:

$$G' = \mu + A' + D' + e = \mu + x'_{A_i}\alpha + z'_{D_i}d + e \text{ (equation 5)}$$

Where x'_A was coded as $2q$, $p - q$ and $-2p$ for genotype of AA, AB and BB of variant i,

respectively, to represent the additive component and z'_D was coded as $-2p^2$, $2pq$, $-2q^2$ for

genotype of AA, AB and BB of variant i, respectively, to represent the dominance

component. Such re-parametrisation has the following features: 1) The covariance between

the additive and dominance effects is zero; 2) the variance of the additive effects gives the

additive variance; and 3) The variance of the dominance deviations gives the dominance

variance. Equation 5 then leads to:

$$H'_j = {\sum_i^N z'_{D_i}} \Big/ N \text{ (equation 6)}$$

Where H'_j was the re-parameterised genome-wide heterozygosity for individual j, z'_D was

$-2p^2$, $2pq$, $-2q^2$ for genotype of AA, AB and BB of variant i and N was the total number of

variants. We then applied equation 6 to conserved and non-conserved variants to estimate

individual heterozygosity from conserved sites (H'_{cons_j}) and non-conserved sites

($H'_{non-cons_j}$). We then fitted H'_{cons_j} and $H'_{non-cons_j}$ as fixed effects together with the fixed

effects of breed jointly in GREML similar to equation 1. The difference was that there is no

fixed effect of variants but more fixed effects due to the fitting of H'_{cons_j} and $H'_{non-cons_j}$.

The GREML analysis used the implementation with MTG2 [44].

Mutant allele frequency and F_{ST} in different breeds/subspecies. Two sets of data were

used for this analysis. The first dataset was the Australian dairy cattle (8,949 bulls and

103,350 cows, Holstein, Jersey, Australian Red and crossbreds) used for GWAS as described

above. The second data set used for the analysis of mutant allele frequency and F_{ST} was the

curated whole-genome sequence data of 1,720 cattle from the 1000 Bull Genomes database

(Run 7) [21,22], which we refer to as modern and ancient cattle. Samples that met the quality

criteria of the 1000 Bull Genomes project were selected and they included 210 Brahman, 25

Tibetan, 10 Eurasian Aurochs, 242 Simmental, 95 Jersey, 843 Holstein and 295 Angus.

Genome sequences from 6 Gir and 12 Nellore cattle from the 1000 Bull Genomes database

were also analysed to support the results of mutant allele frequency of *Bos indicus*.

Additional information on these 1720 animals including related accession numbers (if

available) can be found in Supplementary Data 1. The ancient genome data were part of the

project of Verdugo et al 2019 [15] who processed and published the original data (PRJEB31621

at European Nucleotide Archive). These data were collected by Run 7 of the 1000 Bull

Genomes Project and processed by its standard pipeline

(http://www.1000bullgenomes.com/).

Sequence data at 7,910,190 variants assigned with mutant alleles were retrieved for these

animals to make a plink (v1.9) binary genotype file. The A1 allele of the plink genotypes was

set to the mutant allele and its frequency was calculated using the '--freq' function for

different selections of populations and variant sets. Average mutant allele frequency and the

standard error were calculated for different selections of variants, e.g., variants with mutant

alleles increasing or decreasing traits. Standard errors for frequency and F_{ST} (described

below) were all estimated using LD-clumped variants in the same procedure in plink [29] as

described above. For variants associated with milk production traits, i.e., the yield of milk

protein, fat and milk and percentage of protein and fat, we selected variants with large

(GWAS p-value < 5e-8 in both sexes) and small (GWAS p-value < 5e-2 and p-value > 5e-5

in both sexes) effects to focus on. For other trait-associated variants, the group with the

largest effects available were selected for this comparison. For example, for stature, there

were no variants with p-value < 5e-8 in both sexes, we then selected the medium-effect

variants (GWAS p-value < 5e-5 and p-value > 5e-8 in both sexes). For fertility, there was no

variants with p-value < 5e-5 in both sexes, we then selected the small-effect variants (GWAS

p-value < 5e-2 and p-value > 5e-5 in both sexes) for the comparison. Average mutant allele

frequency and the standard error were also calculated for all 7.9M variants analysed as the

baseline. The analysis procedure for allele frequency on the Australian dairy cattle was

applied to these 1000 Bull Genomes individuals.

With the same plink binary genotype file described above and the population structure for

dairy cattle (4 dairy breeds) and for ancient and modern cattle (7 breeds/subspecies), GCTA

[43] was used to calculate the F_{ST} value with the method described in Weir [49] with the option of

'--fst' and '--sub-pop'. The average F_{ST} value with standard errors was then calculated for

different selections of variants in the same fashion for selecting variant groups to compare the

mutant allele frequency as described above.

cis eQTL in milk cells. This analysis was based on 105 Holstein cattle who had RNA-seq

data in milk cells described and published previously (NCBI SRA SRP111067) [23,24]. The raw

reads of these data were aligned to the ARS-UCD1.2 reference bovine genome using STAR[50]

and the quality check followed what was described previously [23]. FeatureCount [51] was used to

extract gene counts and the voom [52] normalised counts were used in the following analyses.

The normalised gene expression was analysed as phenotypes in the same GWAS model as

equation 1 using GCTA, except that there were no breed effects (all animals are Holstein) but

were other fixed effects of Experiment, Days in Milk, 1st PC and 2nd PC extracted from the

expression count matrix. Variants analysed were those that had large positive effects and

large negative effects (p_{gwas}< 5e-8) on protein yield, fat yield, milk yield, protein % and fat

%. For these variants, the normalised expression of genes within ±1Mb distance to them were

analysed as phenotype. In other words, the analysis focused on cis eQTL genes for these

large-effect variants were analysed. When GWAS results of gene expression were obtained

(cis eQTL), the effect allele was mapped to the ancestral allele to determine the effects of

MAs. For quantifying the number of eQTL for each effect direction of MAs, only the SNPs

with the smallest p-value were considered.

Data availability

Our predictions of cattle ancestral alleles for those 42,573,455 sites have been made publicly

available at: https://figshare.com/s/dd5985b76a413b56106b. Multiple alignment data used to

determine cattle ancestral alleles are publicly available via Ensembl EPO pipeline

(http://asia.ensembl.org/info/genome/compara/multiple_genome_alignments.html).

Australian farmers and DataGene Australia (http://www.datagene.com.au/) are owners and

custodians of the raw phenotype and genotype data of Australian dairy animals. Access to

these data for research requires permission from DataGene under a Data Use Agreement. The

DNA sequence data as part of the 1000 Bull Genomes Consortium [20-22] are available to

consortium members and the membership is open. Sequence data of 1832 samples from the

1000 Bull Genome Project have been made publicly available at:

https://www.ebi.ac.uk/eva/?eva-study=PRJEB42783. The gene expression data is publically

available (NCBI SRA SRP111067). In addition: 1. The summary data of the effect direction

and effect category of those 7.9M sequence variants for which the ancestral alleles can be

assigned is published at https://figshare.com/s/ef020d948523c31c0e67; 2. The allele frequency

of mutant alleles of those 7.9M sequence variants for which the ancestral alleles can be

assigned for the Holstein and Jersey cattle from the 1000 Bull Genome Project is published at

https://figshare.com/s/20154b1d8e60e012e532; 3. The coordinates of conserved sites analysed in

the manuscript is published at: https://figshare.com/s/df9d3662f8f7fb8e72da. Other supporting

data are shown in the supplementary materials of the current manuscript.

Code availability

The probability of ancestral allele assignment used the software published by [25]. The linear

mixed model used GCTA [43] and MTG2 [44]. The Bayesian analysis used BayesR [53]. The R

code of estimating heterozygosity across conserved sites will be made public upon

publication.

References:

1   Ohta, T. Slightly deleterious mutant substitutions in evolution. *Nature* **246**, 96-98 (1973).

2   Eyre-Walker, A., Keightley, P. D., Smith, N. G. & Gaffney, D. Quantifying the slightly
deleterious mutation model of molecular evolution. *Molecular Biology and Evolution* **19**,
2142-2149 (2002).

3   Eyre-Walker, A. & Keightley, P. D. The distribution of fitness effects of new mutations.
Nature Reviews Genetics **8**, 610-618 (2007).

4   Mehlhoff, J. D. *et al.* Collateral fitness effects of mutations. *Proceedings of the National
Academy of Sciences* **117**, 11597-11607 (2020).

5   Meuwissen, T., Hayes, B. & Goddard, M. Genomic selection: A paradigm shift in animal
breeding. *Animal frontiers* **6**, 6-14 (2016).

6   Chen, J., Glémin, S. & Lascoux, M. From drift to draft: how much do beneficial mutations
actually contribute to predictions of Ohta's slightly deleterious model of molecular
evolution? *Genetics* **214**, 1005-1018 (2020).

7   Keller, L. F. & Waller, D. M. Inbreeding effects in wild populations. *Trends in Ecology &
Evolution* **17**, 230-241, doi:https://doi.org/10.1016/S0169-5347(02)02489-8 (2002).

8   Turelli, M. & Ginzburg, L. R. Should individual fitness increase with heterozygosity? *Genetics*
104, 191-209 (1983).

9   Jiang, Y. *et al.* The sheep genome illuminates biology of the rumen and lipid metabolism.
Science **344**, 1168-1173, doi:10.1126/science.1252806 (2014).

10   Loftus, R. T., MacHugh, D. E., Bradley, D. G., Sharp, P. M. & Cunningham, P. Evidence for two
independent domestications of cattle. *Proceedings of the National Academy of Sciences* **91**,
2757-2761 (1994).

11   Bradley, D. G., MacHugh, D. E., Cunningham, P. & Loftus, R. T. Mitochondrial diversity and
the origins of African and European cattle. *Proceedings of the National Academy of Sciences*
93, 5131-5135 (1996).

12   Troy, C. S. *et al.* Genetic evidence for Near-Eastern origins of European cattle. *Nature* **410**,
1088-1091 (2001).

13   Chen, S. *et al.* Zebu cattle are an exclusive legacy of the South Asia Neolithic. *Molecular
biology and evolution* **27**, 1-6 (2010).

14   Utsunomiya, Y. *et al.* Genomic clues of the evolutionary history of Bos indicus cattle. *Animal
genetics* **50**, 557-568 (2019).

15   Verdugo, M. P. *et al.* Ancient cattle genomics, origins, and rapid turnover in the Fertile
Crescent. *Science* **365**, 173-176 (2019).

16   Upadhyay, M. *et al.* Genetic origin, admixture and population history of aurochs (Bos
primigenius) and primitive European cattle. *Heredity* **118**, 169-176 (2017).

17   Park, S. D. *et al.* Genome sequencing of the extinct Eurasian wild aurochs, Bos primigenius,
illuminates the phylogeography and evolution of cattle. *Genome biology* **16**, 234 (2015).

18   García-Ruiz, A. *et al.* Changes in genetic selection differentials and generation intervals in US
Holstein dairy cattle as a result of genomic selection. *Proceedings of the National Academy*
of Sciences **113**, E3995-E4004 (2016).

19   Coffey, E., Horan, B., Evans, R. & Berry, D. Milk production and fertility performance of
Holstein, Friesian, and Jersey purebred cows and their respective crosses in seasonal-calving
commercial farms. *Journal of Dairy Science* **99**, 5681-5689 (2016).

20   Bouwman, A. C. *et al.* Meta-analysis of genome-wide association studies for cattle stature
identifies common genes that regulate body size in mammals. *Nature genetics* **50**, 362
(2018).

21   Daetwyler, H. D. *et al.* Whole-genome sequencing of 234 bulls facilitates mapping of
monogenic and complex traits in cattle. *Nature genetics* **46**, 858 (2014).

22   Hayes, B. J. & Daetwyler, H. D. 1000 Bull Genomes Project to Map Simple and Complex
Genetic Traits in Cattle: Applications and Outcomes. *Annual review of animal biosciences* **7**,
89-102, doi:10.1146/annurev-animal-020518-115024 (2019).

23   Xiang, R. *et al.* Genome variants associated with RNA splicing variations in bovine are
extensively shared between tissues. *BMC Genomics* **19**, 521, doi:10.1186/s12864-018-4902-
8 (2018).

24   Chamberlain, A. *et al.* in *11th world congress on genetics applied to livestock production*
(WCGALP). Auckland, New Zealand: Volume Molecular Genetics. 254.

25   Keightley, P. D. & Jackson, B. C. Inferring the probability of the derived vs. the ancestral
allelic state at a polymorphic site. *Genetics* **209**, 897-906 (2018).

26   Rocha, D., Billerey, C., Samson, F., Boichard, D. & Boussaha, M. Identification of the putative
ancestral allele of bovine single-nucleotide polymorphisms. *Journal of Animal Breeding and*
Genetics **131**, 483-486 (2014).

27   Xiang, R., van den Berg, I., MacLeod, I. M., Daetwyler, H. D. & Goddard, M. E. Effect direction
meta-analysis of GWAS identifies extreme, prevalent and shared pleiotropy in a large
mammal. *Commun Biol* **3**, 88, doi:10.1038/s42003-020-0823-6 (2020).

28   Erbe, M. *et al.* Improving accuracy of genomic predictions within and between dairy cattle
breeds with imputed high-density single nucleotide polymorphism panels. *Journal of dairy*
science **95**, 4114-4129 (2012).

29   Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer
datasets. *Gigascience* **4**, 7, doi:10.1186/s13742-015-0047-8 (2015).

30   Chen, N. *et al.* Whole-genome resequencing reveals world-wide ancestry and adaptive
introgression events of domesticated cattle in East Asia. *Nature Communications* **9**, 1-13
(2018).

31   Xiang, R. *et al.* Quantifying the contribution of sequence variants with regulatory and
evolutionary significance to 34 bovine complex traits. *Proceedings of the National Academy*
of Sciences **116**, 19398-19408, doi:10.1073/pnas.1904159116 (2019).

32   Meuwissen, T., Hayes, B. & Goddard, M. Prediction of total genetic value using genome-wide
dense marker maps. *Genetics* **157**, 1819-1829 (2001).

33   Daetwyler, H. *et al.* in *Proc Assoc Adv Anim Breed Genet.* 201-204.

34   DePristo, M. A. *et al.* A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nature genetics* **43**, 491-498 (2011).

35   Browning, S. R. & Browning, B. L. Rapid and accurate haplotype phasing and missing-data
inference for whole-genome association studies by use of localized haplotype clustering.
American journal of human genetics **81**, 1084-1097, doi:10.1086/521987 (2007).

36   Siepel, A. *et al.* Evolutionarily conserved elements in vertebrate, insect, worm, and yeast
genomes. *Genome research* **15**, 1034-1050 (2005).

37   Fuchsberger, C., Abecasis, G. R. & Hinds, D. A. minimac2: faster genotype imputation.
Bioinformatics **31**, 782-784 (2014).

38   Howie, B., Fuchsberger, C., Stephens, M., Marchini, J. & Abecasis, G. R. Fast and accurate
775   genotype imputation in genome-wide association studies through pre-phasing. *Nature*
776   *genetics* **44**, 955 (2012).

777   39   Sargolzaei, M., Chesnais, J. P. & Schenkel, F. S. A new approach for efficient genotype
778   imputation using information from relatives. *BMC Genomics* **15**, 478, doi:10.1186/1471-
779   2164-15-478 (2014).

780   40   Loh, P.-R. *et al.* Reference-based phasing using the Haplotype Reference Consortium panel.
781   *Nature genetics* **48**, 1443 (2016).

782   41   Rosen, B. D. *et al.* De novo assembly of the cattle reference genome with single-molecule
783   sequencing. *GigaScience* **9**, giaa021 (2020).

784   42   Pausch, H. *et al.* Evaluation of the accuracy of imputed sequence variant genotypes and their
785   utility for causal variant detection in cattle. *Genetics Selection Evolution* **49**, 1-14 (2017).

786   43   Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide complex
787   trait analysis. *The American Journal of Human Genetics* **88**, 76-82 (2011).

788   44   Lee, S. H. & Van der Werf, J. H. MTG2: an efficient algorithm for multivariate linear mixed
789   model analysis based on genomic information. *Bioinformatics* **32**, 1420-1422 (2016).

790   45   Xiang, R. *et al.* Genome-wide fine-mapping identifies pleiotropic and functional variants that
791   predict many traits across global cattle populations. *Nature Communications* **12**, 860,
792   doi:10.1038/s41467-021-21001-0 (2021).

793   46   Fisher, R. A. XV.—The correlation between relatives on the supposition of Mendelian
794   inheritance. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*
795   **52**, 399-433 (1919).

796   47   Falconer, D. S. & Mackay, T. F. C. *Introduction to quantitative genetics*. (Longman, 1996).

797   48   Lynch, M. & Walsh, B. *Genetics and Analysis of Quantitative Traits*. (Sinauer Associates,
798   1998).

799   49   Weir, B. S. & Ott, J. Genetic data analysis II. *Trends in genetics* **13**, 379 (1997).

800   50   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
801   doi:10.1093/bioinformatics/bts635 (2013).

802   51   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
803   assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).

804   52   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. Voom: precision weights unlock linear model
805   analysis tools for RNA-seq read counts. *Genome Biology* **15**, R29 (2014).

806   53   Moser, G. *et al.* Simultaneous discovery, estimation and prediction analysis of complex traits
807   using a Bayesian mixture model. *PLoS genetics* **11**, e1004969 (2015).

Acknowledgements

Australian Research Council's Discovery Projects (DP160101056 and DP200100499) supported R.X. and M.E.G. DairyBio, a joint venture project between Agriculture Victoria (Melbourne, Australia), Dairy Australia (Melbourne, Australia) and the Gardiner Foundation (Melbourne, Australia), funded computing resources used in the analysis. The authors also thank the University of Melbourne, Australia for supporting this research. No funding bodies participated in the design of the study nor analysis, or interpretation of data nor writing of the manuscript. DataGene provided access to the reference data used in this study and the 1000 Bull Genomes consortium provided access to cattle sequence data. We thank Gert Nieuwhof and Kon Konstantinov (DataGene) for the preparation and provision of data. We thank Prof. Hans D. Daetwyler for assisting the access to the 1000 Bull Genomes data. We thank Prof. Naomi Wray for a critical read of the manuscript.

Author contributions

M.E.G. and R.X conceived the study. R.X. performed all analyses. E.J.B. contributed to the BayesR analysis. I.M.M., assisted with data curation. S.B., C.J.J., and A.J.C. contributed to the imputation of sequence variants. R.X. and M.E.G. wrote the paper. R.X., M.E.G., E.J.B. and I.M.M. revised the paper. All authors read and approved the final manuscript.

Competing Interests

The authors declare no competing interests.