Effects of CO_2 and RuBisCO concentration on cyanobacterial growth and carbon isotope fractionation

Amanda Garcia[1], Mateusz Kedzior[1], Arnaud Taton[2], Meng Li[3], Jodi Young[3] and Betül Kaçar [1*]

[1]Department of Bacteriology, University of Wisconsin – Madison, Madison, WI

[2]Division of Biological Sciences, University of California San Diego, La Jolla, CA

[3]School of Oceanography, University of Washington, Seattle, WA

*For correspondence: bkacar@wisc.edu

KEYWORDS

carbon fixation, carbon isotope fractionation, RuBisCO, biosignatures, cyanobacteria

ABSTRACT

Carbon isotope biosignatures preserved in the Precambrian geologic record are primarily interpreted to reflect ancient cyanobacterial carbon fixation catalyzed by Form I RuBisCO enzymes. The average range of isotopic biosignatures generally follows that produced by extant cyanobacteria. However, this observation is difficult to reconcile with several environmental (e.g., temperature, pH, and CO_2 concentrations), molecular, and physiological factors that likely would have differed during the Precambrian and can produce fractionation variability in contemporary organisms that meets or exceeds that observed in the geologic record. To test a range of genetic and environmental factors that may have impacted ancient carbon isotope biosignatures, we

engineered a mutant strain of the model cyanobacterium *Synechococcus elongatus* PCC 7942 that

overexpresses RuBisCO and characterized the resultant physiological and isotope fractionation

effects. We specifically investigated how both increased atmospheric CO_2 concentrations and

RuBisCO regulation influence cell growth, oxygen evolution rate, and carbon isotope fractionation

in cyanobacteria. We found that elevated CO_2 increases the growth rate of wild-type and mutant

strains, and that the pool of active RuBisCO enzyme increases with increased expression.

RuBisCO overexpression in our engineered strain does not significantly affect isotopic

discrimination at all tested CO_2 concentrations, yielding cellular $^{13}C/^{12}C$ isotope discrimination

(ε_p) of ~24‰ for both wild-type and mutant strains at elevated CO_2. Understanding the

environmental factors that impact gene regulation, physiology, and evolution is crucial for

reconciling microbially driven carbon isotope fractionation with the geologic record carbon

biosignatures.

INTRODUCTION

The conserved microbial metabolic pathways that drive global biogeochemistry emerged on Earth billions of years ago, the evolution of which has both shaped and been shaped by large-scale environmental transitions (Falkowski, Fenchel, & Delong, 2008; Knoll & Nowak, 2017; Lyons, Fike, & Zerkle, 2015). These microbial processes have left distinct signatures that are evidence of biological activity billions of years in the past. The oldest and most extensive signature of biological activity on Earth is the deviation in stable carbon isotopic compositions ($^{13}C/^{12}C$, expressed as $\delta^{13}C$) between preserved inorganic and organic carbon, interpreted to reflect the isotopic discrimination of ancient biological carbon fixation (Des Marais, 2001; Krissansen-Totton, Buick, & Catling, 2015; Lloyd et al., 2020; Schidlowski, 2001). This deviation is primarily shaped by enzymes that preferentially assimilate the lighter ^{12}C isotope from inorganic carbon sources. Carbon biosignatures preserved in the geologic record therefore reflect the long-term evolution of these enzyme-mediated processes and their hosts' physiologies.

The RuBisCO enzyme (ribulose 1,5-bisphosphate (RuBP) carboxylase/oxygenase) catalyzes the reduction of inorganic CO_2 as the initial step of carbon assimilation into organic biomass via the Calvin-Benson-Bassham (CBB) cycle (Erb & Zarzycki, 2018; Nisbet et al., 2007; Tcherkez, Farquhar, & Andrews, 2006). RuBisCO is one of the most abundant proteins on Earth (Bar-On & Milo, 2019; Ellis, 1979; Raven, 2013), and its presence in photoautotrophic organisms, including early-evolved cyanobacteria, suggests that this enzyme has played a significant role in primary production for much of Earth history (Hamilton, Bryant, & Macalady, 2016; Schirrmeister, Sanchez-Baracaldo, & Wacey, 2016; Schopf, 2011). Thus, the isotopic fractionation behavior of

RuBisCO is also thought to have constrained Precambrian carbon isotope signatures preserved in

the geologic record (Garcia, Cavanaugh, & Kacar, 2021; Schidlowski, 1988). Though there are

multiple forms of RuBisCO, the known range of the Form I RuBisCO isotopic effect ($\varepsilon \approx$ 20-30‰)

(Guy, Fogel, & Berry, 1993; Scott et al., 2007; von Caemmerer, Tazoe, Evans, & Whitney, 2014)

— the form utilized by extant cyanobacteria and responsible for the bulk of modern primary

production (Field, 1998) — is largely consistent with the ~25‰ mean deviation between preserved

inorganic and organic carbon isotopic compositions across geologic time (Des Marais, 2001;

Havig, Hamilton, Bachan, & Kump, 2017; Kedzior et al., 2022; Krissansen-Totton et al., 2015;

Lloyd et al., 2020; Schidlowski, 2001). The isotopic effects of cyanobacterial Form IA and IB

RuBisCO in particular have been measured between ~22 to 24‰ (Guy et al., 1993; Scott et al.,

2007), whereas land plant Form IB RuBisCO typically has larger effects of up to ~29‰ (von

Caemmerer et al., 2014) and Form ID RuBisCO have exhibited effects as small as ~11‰ (Boller,

Thomas, Cavanaugh, & Scott, 2011).

RuBisCO is an important factor in the generation of distinct carbon isotopic biosignatures

associated with CBB-utilizing organisms (Hayes, 2001; Laws, Popp, Bidigare, Kennicutt, &

Macko, 1995; Wilkes & Pearson, 2019). Nonetheless the carbon isotopic composition of bulk

photoautotrophic biomass often deviates from values obtained for purified RuBisCO, a

discrepancy at least partially explained by the fact that the latter are measured at CO_2 saturation

(e.g., (Boller et al., 2011; Guy et al., 1993; Scott et al., 2007; von Caemmerer et al., 2014)). This

discrepancy might also be attributable to several intracellular physiological and metabolic features

that additionally shift the isotopic composition in organism biomass. These include the activation

of carbon-concentrating mechanisms (CCM) that elevate CO_2 in RuBisCO-containing cellular

compartments (Hurley, Wing, Jasper, Hill, & Cameron, 2021; Laws, Popp, Cassar, & Tanimoto,

2002; Price, Badger, Woodger, & Long, 2008; Raven, Cockell, & De La Rocha, 2008; Wilkes &

Pearson, 2019) and the diffusive transport of CO_2 (Hayes, 1993; Rau, Riebesell, & Wolf-Gladrow,

1996). Studies have shown that photosynthetic carbon isotope discrimination (ε_p) may vary due to

environmental factors and cellular physiological responses, including temperature (Deleens,

Treichel, & O'Leary, 1985; Wong & Sackett, 1978), pH (Hinga, Arthur, Pilson, & Whitaker, 1994;

Roeske & O'Leary, 1984), growth rate (Bidigare et al., 1997; Laws, Bidigare, & Popp, 1997), and

CO_2 concentration (Eichner, Thoms, Kranz, & Rost, 2015; Freeman & Hayes, 1992; Hinga et al.,

1994; Hurley et al., 2021; Schubert & Jahren, 2012; Wilkes, Lee, McClelland, Rickaby, & Pearson,

2018). These variations indicate a compelling need to comprehensively characterize both internal

and external factors that can affect isotope fractionation, though the study of each tend to be siloed

in biological and geobiological fields, respectively.

An aspect of photoautotrophic carbon isotopic discrimination that has not been thoroughly

investigated is how RuBisCO expression levels may have influenced ancient host organism growth

and biosignature generation when CO_2 leves were higher. RuBisCO expression has been shown to

be CO_2-sensitive (Gesch et al., 2003; Onizuka et al., 2002; Sengupta, Sunder, Sohoni, & Wangikar,

2019). This observation is particularly important, considering that atmospheric CO_2 concentrations

exceeded present day levels by more than an order of magnitude for much of Earth history

(between ~0.001 and 0.1 bar CO_2 through the Precambrian (Catling & Zahnle, 2020)) and that

RuBisCO catalytic efficiency itself is sensitive to atmospheric CO_2/O_2 levels (Erb & Zarzycki,

2018; Kacar, Hanson-Smith, Adam, & Boekelheide, 2017; Poudel et al., 2020; Riebesell, Revill,

Holdsworth, & Volkman, 2000; Schubert & Jahren, 2012; Scott et al., 2007; Tcherkez et al., 2006;

Wilkes et al., 2018). Though experimental work has demonstrated the positive correlation between

CO_2 levels and RuBisCO isotopic discrimination (Freeman & Hayes, 1992; Hinga et al., 1994;

Schubert & Jahren, 2012; Wilkes et al., 2018), the role of RuBisCO expression, given its CO_2

sensitivity, has not yet been explored. More broadly, the influences of genetic regulatory processes

on ancient biosignatures—otherwise highlighted in geobiological and paleobiological studies of

metazoan evolution (Erwin, 2020; Sumner, 2002; Valentine, 1994)—are not often examined in the

context of Precambrian microbial metabolisms like carbon fixation. Interpretation of the carbon

isotope record should consider the coupled influence of CO_2 concentration and RuBisCO

expression in generating isotopic biosignatures and shaping ancient primary productivity.

To investigate the interplay between RuBisCO expression and CO_2 concentrations on

cyanobacterial physiology and carbon isotopic discrimination, we generated a genetic system to

manipulate the expression levels of RuBisCO in the model organism *Synechococcus elongatus*

PCC 7942 (hereafter *S. elongatus*).(Bonfil et al., 1998; Gabay, Lieman-Hurwitz, Hassidim, Ronen-

Tarazi, & Kaplan, 1998; Maeda, Price, Badger, Enomoto, & Omata, 2000; Omata, Gohta,

Takahashi, Harano, & Maeda, 2001; Tchernov et al., 2001). Genetic engineering is facilitated by

this strain's ability to take up and integrate exogenous DNA into its genome by homologous

recombination (Taton et al., 2020). *S. elongatus* also possesses the β-carboxysome-based CCM

that likely impacted the production of Precambrian carbon biosignatures following the rise of

atmospheric oxygen (Hurley et al., 2021; Lyons, Reinhard, & Planavsky, 2014). Thus, *S. elongatus*

is a suitable model to study the genetic basis of ancient carbon biosignatures. We engineered its

genome with an additional copy of the RuBisCO operon in a chromosomal neutral site to permit

genetic manipulation of the RuBisCO operon and confirmed overexpression of RuBisCO under

varying CO_2 levels. Then, to assess whether physiological and subcellular parameters and different

CO_2 concentrations modulate carbon isotopic discrimination in cyanobacteria, we determined

growth rate, photosynthetic oxygen evolution rate, and $^{13}C/^{12}C$ discrimination of the engineered

strain in comparison to wild-type *S. elongatus* under different atmospheric conditions.

RESULTS

A second copy of the *rbc* operon results in increased amount of active RuBisCO

The RuBisCO Form IB enzyme in *S. elongatus* is encoded by an operon that includes a CO_2-

sensitive promoter region (Sengupta et al., 2019) as well as the structural *rbcL* (large subunit) and

rbcS (small subunit) genes (Vijayan, Jain, & O'Shea, 2011). We designed *S. elongatus* strain Syn02

that harbors the native *rbc* operon and a second copy inserted in the chromosome neutral site 2

(NS2), a site that permits genetic modification without additional indirect phenotypic impact

(Andersson et al., 2000; Clerico, Ditty, & Golden, 2007) (**Fig. 1**). Additionally, we generated a

control strain (Syn01) whereby RuBisCO is provided solely by an engineered *rbc* operon at its

NS2 site (**Table 1**). Attributes of these strains related to carbon fixation, including RuBisCO

transcription and protein expression, carboxylase activity, and carbon isotope fractionation, were

evaluated under different CO_2 conditions, including ambient (~0.04%), 2%, and 5% CO_2

concentrations. These CO_2 concentrations were selected to measure the impact of a Precambrian-

like atmosphere, which is generally constrained between ~0.1% to ~10% CO_2 for much of the

early evolution of cyanobacteria (assuming ~1-bar total paleopressure) (Catling & Zahnle, 2020).

We evaluated whether the additional copy of the *rbc* operon in strain Syn02 resulted in increased

RuBisCO *rbcL* and *purK* (located downstream of *rbcL* in the operon) transcription under varying

CO_2 levels (ambient, 2%, and 5% CO_2). Transcription was measured by quantitative reverse-

transcription PCR (RT-qPCR), normalized to *secA* and *ppc* reference genes (Hood, Higgins,

Flamholz, Nichols, & Savage, 2016; Luo et al., 2019; Szekeres, Sicora, Dragos, & Druga, 2014).

We observed that elevated CO_2 enhanced WT *rbcL* transcription relative by at least ~5-fold in

PCC7942 ($p < 0.001$; **Fig. 2B, Fig. S1B; see Materials and Methods**). *rbcL* transcription was

further increased by at least ~2-fold in Syn02 relative to WT across all tested CO_2 levels and

growth phases, and as high as ~14-fold in air ($p < 0.01$; **Fig. 2A, Fig. S1A**). Expression of *purK* in

Syn02 increased by >2-fold at 2% and 5% CO_2 relative to WT, but decreased by ~0.5-fold in air

(**Fig. S1C**).

To determine *rbcL* overexpression at the protein-level, we quantified RbcL protein from crude cell

lysates by Western blot using rabbit anti-RbcL antibody (see **Materials and Methods**). In

agreement with RuBisCO overexpression indicated by the RT-qPCR results, densitometric

analyses revealed a mean ~2 to 4-fold increase of RbcL protein in Syn02 relative to WT across all

tested CO_2 concentrations ($p < 0.01$; **Fig. 2C**). Finally, we confirmed proper assembly of the large

(L) and small (S) subunits into the RuBisCO L_8S_8 complex in Syn02 as well as the Syn01 control

strain by native protein electrophoresis and detection by anti-RbcL antibody. At 2% CO_2, we found

a ~3-fold increase in assembled RuBisCO protein for Syn02 and ~2-fold increase for Syn01

relative to WT for cultures grown in 2% CO_2 (**Fig. 2D**).

Finally, we tested whether RuBisCO overexpression in Syn02 resulted in an increased amount of

active RuBisCO by determining the total carboxylase activity of cell lysates. Because we detected

the smallest increase in transcript levels for Syn02 relative to WT under 2% CO_2, we selected this

growth condition to provide a reasonable lower bound on Syn02 RuBisCO activity. We measured

a mean ~1.8-fold ($p < 0.01$) increase in Syn02 lysate RuBisCO activity relative to WT (**Fig. 2E**).

Despite the modest increase in assembled RuBisCO detected for the control strain, Syn01, no

significant difference in activity was found between Syn01 and WT cultures grown in these same

conditions.

RuBisCO overexpression does not strongly influence growth rate or photosynthetic activity

Following confirmation that Syn02 overexpresses active RuBisCO, we compared growth rates of

WT and Syn02 *S. elongatus* strains at different CO_2 levels to identify potential downstream

physiological effects of increased RuBisCO protein. WT and Syn02 both exhibited ~2.5-fold faster

growth rates under 2% and 5% CO_2 compared to ambient air ($p < 0.001$; **Fig. 3; Table 2**). Syn02

exhibited a slight ~1.1-fold increase in growth rate in ambient air relative to WT ($p < 0.05$). No

difference in growth rate was observed between the two strains under 2% and 5% CO_2. The

carrying capacity (maximum cell density measured at OD_{750} across the total growth period) for

cultures varied between different atmospheric conditions, with a maximum carrying capacity of

$OD_{750} \approx 8.4$ reached under 2% CO_2. No significant difference in carrying capacity was found

between WT and Syn02 cultures grown under the same atmospheric conditions. These experiments

were replicated with unsparged cultures, yielding consistent results (**Table S1**).

To further test for differences in photosynthetic activities between the WT and Syn02 strains, we

measured their oxygen evolution rates in ambient air. After brief incubation in the dark, culture

samples were exposed to saturated light in an oxygen electrode chamber to detect levels of

molecular oxygen. Oxygen evolution rates were normalized to chlorophyll a concentrations,

following Zavřel et al. (2015). We did not observe a significant difference in mean oxygen

evolution rate between WT and Syn02 (340 ± 30 $O_2 \cdot h^{-1} \cdot \mu g^{-1}$ chlorophyll a and 360 ± 30 $O_2 \cdot h^{-1} \cdot \mu g^{-1}$

^{1} chlorophyll a, respectively) (**Table 2**).

***S. elongatus* ^{13}C/^{12}C fractionation is insensitive to RuBisCO overexpression**

We tested the combined influence of RuBisCO overexpression and CO_2 concentration on the

magnitude of whole-cell ^{13}C/^{12}C isotopic discrimination in *S. elongatus*. The isotopic

discrimination associated with photosynthetic CO_2 fixation (ε_p) was calculated from measured

$\delta^{13}C_{biomass}$ and headspace $\delta^{13}C_{CO2}$ in our experimental system, as described in **Materials and**

Methods. We observed that increased CO_2 concentration (2% CO_2) increases the degree of *S.*

elongatus ^{13}C/^{12}C discrimination relative to air. Mean ε_p values for WT and Syn02 strains grown

in ambient air were $4.8 \pm 0.3‰$ and $5.1 \pm 0.4‰$, respectively (**Fig. 4; see Supplementary**

Information, Table S2 and **S3** for δ^{13}C and pH measurement valuesvalues). At 2% CO_2, mean ε_p

values for WT and Syn02 strains were $23.75 \pm 0.07‰$ and $23.7 \pm 0.5‰$, respectively. The

difference between WT and Syn02 ε_p values at both tested CO_2 conditions were found to be

insignificant ($p > 0.05$). These results demonstrate that the increased RuBisCO transcriptional

levels and total carboxylasecarboxylase activity measured for Syn02 in our experiments does not

significantly impact whole-cell *S. elongatus* $^{13}C/^{12}C$ discrimination, both at ambient and elevated

CO_2 concentrations.

DISCUSSION

Evolutionary biologists have long debated the role of genetic regulation versus protein-level

variation in evolutionary selection (Carroll, 2005; Fay & Wittkopp, 2008; Olson-Manning,

Wagner, & Mitchell-Olds, 2012; Taylor, Shepherd, Jackson, & Silby, 2022; Wilson, Maxson, &

Sarich, 1974). How such molecular and genetic variation leads to evolutionary change is often

overlooked in geobiological evaluation of ancient microbial biosignatures due to the challenges

involved with disentangling their effects from other variables. Laboratory experimental systems,

though admittedly incapable of capturing the full complexity of ancient, natural systems, are

necessary to understand the genetic factors underpinning the oldest records of life. In this study,

we developed an engineered laboratory system to investigate the coupled impact of elevated CO_2

levels and RuBisCO expression on cyanobacterial physiology and carbon isotope discrimination.

We tested whether the overexpression of RuBisCO influences *S. elongatus* growth, photosynthetic

oxygen evolution rate, and carbon isotopic signatures at different CO_2 atmosphere concentrations

by generating an engineered *S. elongatus* strain Syn02 that harbors a second copy of the *rbc* operon

at NS2.

We confirmed overexpression of RuBisCO in Syn02 by combining transcriptional analyses,

protein quantification, and assays of total protein carboxylation activity. Our results show elevated

RuBisCO transcription in *S. elongatus* strain Syn02 relative to WT at all tested CO_2 concentrations

(ambient air, 2%, and 5% CO_2) (**Fig. 2A**), with a ~2-fold increase in active, assembled RuBisCO

protein measured under 2% CO_2 (**Fig. 2D and 2E**). We also found that higher levels of CO_2 result

in increased RuBisCO transcription for both WT and Syn02 strains relative to ambient air, with

the greatest fold change increase detected at 2% CO_2. This result is in agreement with a previous

study reporting a regulatory element in the *S. elongatus* PCC 7942 *rbcL* operon that regulates the

expression of RbcL with increasing CO_2 (Sengupta et al., 2019).

Our experimental approach also enabled us to detect potential upper bounds on the ability of *S.*

elongatus to overexpress RuBisCO under a variety of CO_2 conditions. First, increased *rbcL*

transcription in Syn02 relative to WT did not always result in a proportional increase in translated

RuBisCO protein. For example, even though Syn02 showed a ~14-fold increase in *rbcL* transcript

relative to WT in air (**Fig. 2A**), total Syn02 RbcL protein only increased by ~3-fold under these

same conditions (**Fig. 2C**). These results indicate that other physiological factors, possibly

including translation efficiency or expression of ancillary proteins in our experimental system,

may ultimately limit RuBisCO expression beyond transcription. Further, it is important to consider

whether all translated RuBisCO protein is catalytically active. Previous studies have shown that

overexpression of RuBisCO in cyanobacteria generally produces an increased pool of active

carboxylase (Atsumi, Higashide, & Liao, 2009; Iwaki et al., 2006; Lechno-Yossef et al., 2020;

Liang & Lindblad, 2017). Indeed, for cells grown in 2% CO_2, we found that nearly all

overexpressed RuBisCO protein was likely to be assembled and catalytically active (**Fig. 2C** and

2E). Future work may determine whether this proportion of active, overexpressed RuBisCO holds

for cells cultivated at other CO_2 conditions. Finally, we observed that the transcriptional impact of

a second *rbc* copy in Syn02 was dampened at elevated CO_2 relative to air (**Fig. 2A**), suggesting a

potential limit to the combined expression-level effects of increased *rbc* copy number and

increased CO_2. Altogether, our findings suggest the presence of physiological constraints that may

be limiting RuBisCO overexpression at both transcript and protein levels, the exact mechanism of

which awaits further characterization. Similar approaches investigating other model microbial

systems would clarify the degree to which these observations might extend to cyanobacteria gene

regulation in the past.

We investigated whether the overexpression of RuBisCO in strain Syn02 impacted the growth

characteristics of *S. elongatus* and, by that manner, might have the potential to influence isotopic

discrimination (Bidigare et al., 1997; Laws et al., 1997; Wilkes et al., 2018). Though various

culture conditions including pH, temperature, CO_2 concentration, nutrient availability, and light

intensity may substantially affect the growth characteristics of cyanobacteria (Kuan, Duff, Posarac,

& Bi, 2015; Rillema, MacCready, & Vecchiarelli, 2020; Ungerer et al., 2018; Yu et al., 2015), the

influence of genetic factors, including the regulation of RuBisCO expression, is less well known.

Previous studies targeting the correlation between RuBisCO overexpression and cyanobacterial

physiology have yielded mixed results. For instance, faster growth rates and oxygen evolution

rates were observed for an engineered *Synechocystis* PCC 6803 strain that overexpresses RuBisCO

(Liang & Lindblad, 2017), as well as in *S. elongatus* upon co-overexpression of its

phosphoribulokinase (Kanno, Carroll, & Atsumi, 2017). However, in *Synechococcus* sp. PCC

7002, the overexpression of RuBisCO did not alter growth rate (De Porcellinis et al., 2018). The

impact of RuBisCO upregulation on cyanobacterial growth appears to be species/strain-specific.

Our data shows that RuBisCO overexpression in *S. elongatus* PCC 7942 strain Syn02 results in a

minor increase in growth rate in ambient air relative to WT, and no significant difference in growth

or oxygen evolution rates was observed at elevated CO_2 levels. Our results indicate that further

study is needed to fully understand the coupled interaction of different environmental conditions

and genetic backgrounds in determining cyanobacterial growth parameters of relevance to carbon

biosignature generation.

We found that the degree of RuBisCO overexpression generated in Syn02 did not measurably

impact *S. elongatus* carbon isotope discrimination, both in ambient air and at elevated CO_2 (**Fig.**

4). Rather, the ε_p variability observed in our experiments stemmed solely from variable CO_2, an

effect that has previously been observed empirically for a variety of autotrophs (Freeman & Hayes,

1992; Hinga et al., 1994; Schubert & Jahren, 2012; Wilkes et al., 2018), including for

cyanobacteria in particular (Eichner et al., 2015; Hurley et al., 2021). Other studies have shown

that photoautotrophs grown under a lower pH, and thus a higher proportion of CO_2, also exhibit

higher carbon fractionation (Mizutani & Wada, 1982; Roeske & O'Leary, 1984; Wang, Yeager, &

Lu, 2016; Yoshioka, 1997). In our experiments, ε_p values were increased by ~19‰ for WT and

Syn02 strains in 2% CO_2 relative to air, the latter of which was carbon limiting in our experimental

system (see **Supplementary Information**). Theoretically, under high CO_2 concentrations,

RuBisCO carbon fixation is presumed to be rate-limiting and its intrinsic kinetic isotope effect is

expressed, maximizing ε_p (Bidigare et al., 1997; Hayes, 1993; Schubert & Jahren, 2012; Wilkes et

al., 2018). Under high CO_2 availability, CCM modules have been found to be downregulated in

cyanobacteria, including *S. elongatus* PCC 7942 (Burnap, Hagemann, & Kaplan, 2015; Schwarz

et al., 2011; Woodger, Badger, & Price, 2003). Indeed, the mean $\varepsilon_p \approx 24‰$ value we observed

under high inorganic carbon availability is within the $^{13}C/^{12}C$ isotopic effect range previously

measured for cyanobacterial Form IA and IB RuBisCO at saturating CO_2 (~22–24‰) (Guy et al.,

1993; Scott et al., 2007). Nevertheless, we cannot rule out the impact of upstream cellular modules

in significantly shaping the whole-cell, isotopic signatures observed in our experimental system,

which recent work has demonstrated can exert important isotopic effects under a variety of growth

conditions (Burnap et al., 2015; Eichner et al., 2015; Hurley et al., 2021; Wilkes & Pearson, 2019).

Though we did not quantify the isotopic contributions of these other intracellular factors in our

engineered system, it seems apparent that the increased RuBisCO expression and total protein

carboxylase activity in Syn02 neither changed the $^{13}C/^{12}C$ substrate pool in the enzyme's vicinity

nor measurably altered the potential impact of other cellular modules on carbon isotope

discrimination, even at ambient CO_2 levels.

The range of ε_p values measured for both *S. elongatus* WT and Syn02 strains under elevated CO_2

levels plausibly experienced by ancient cyanobacteria falls within the ~8 to 24‰ ε_p range inferred

from the Precambrian carbon isotope record following the rise of atmospheric oxygen (Hurley et

al., 2021; Krissansen-Totton et al., 2015). Our data suggest that potential overexpression of

RuBisCO by past cyanobacterial primary producers likely does not help explain the distribution

of Precambrian carbon biosignatures. Rather, it is more probable that other environmental and

intracellular factors, including but not limited to CO_2 levels, growth rates, temperature, and CCM

effects, are more significant determinants of the isotopic compositions of organic carbon. We note

in particular that less well-characterized RuBisCO forms (i.e., Forms IC, ID, and II) that are found

in proteobacteria and marine phytoplankton can exhibit significantly smaller fractionation factors

than that observed for forms associated with land plants and cyanobacteria, and perhaps were more

dominant in ancient ecosystems (Boller et al., 2011; Robinson et al., 2003; Thomas et al., 2019).

The full potential breadth of geochemical carbon fixation signatures requires further study (Garcia

et al., 2021), which can be significantly augmented by the integration of taxonomically

uncharacterized systems, the regulatory dynamics of the carbon fixation protein network

beyond RuBisCO, and laboratory reconstructions of ancient carbon-fixing enzymes (Kedzior et

al., 2022). Our results underscore that the integration of synthetic biology, metabolic engineering,

and geochemistry can offer new insights into the study and interpretation of biogeochemical

reservoirs at the global scale. Integrative laboratory approaches are crucial to establish the extent

to which the modern biological carbon isotope fractionation properties can be correlated with

evidence for microbial activity in the Earth's deep past.

MATERIALS AND METHODS

Cyanobacterial growth and maintenance

S. elongatus PCC 7942 strains were cultured in BG-11 medium (Rippka, Deruelles, Waterbury,

Herdman, & Stanier, 1979) as liquid cultures or on agar plates (1.5% (w/v) agar and 1 mM

$Na_2S_2O_3 \cdot 5H_2O$). For recombinant strains, liquid and solid media were supplemented with

appropriate antibiotics: 2 $\mu g \cdot ml^{-1}$ Spectinomycin (Sp) plus 2 $\mu g \cdot ml^{-1}$ Streptomycin (Sm), 5

$\mu g \cdot ml^{-1}$ Kanamycin (Km). The cyanobacterial growth was measured by optical density at 750 nm

(OD_{750}).

The strains were archived at -80 °C in 15% (v/v) glycerol. The 2-ml glycerol stocks were rapidly

thawed and inoculated in liquid BG-11 (supplemented with antibiotics as needed). Liquid cultures

were maintained in a Percival environment chamber (Cat. No. I36LLVLC8) fitted to a CO_2 gas

cylinder input, permitting monitoring and control of internal atmospheric CO_2. Cultures were

shaken at 120 rpm at 30 °C under continuous low illumination of 45 µmol photon·m^{-2}·s^{-1}, and

ambient air, until they reached an OD_{750} between 0.4 and 0.6. These cultures were then used to

inoculate fresh cultures that were grown using similar conditions but under moderate illumination

of 80 µmol photon·m^{-2}·s^{-1} (following (Smith & Williams, 2006)) and sparged at selected CO_2

concentrations (ambient air, 2%, or 5% CO_2; cultures were also grown without sparging for carbon

isotope fractionation experiments and to assess differences in growth rate; **Table S1**). Cultures

were sampled at an OD_{750} of ~7 to 7.5 for most subsequent experiments, as described below.

Cultures were sampled at an earlier OD_{750} of 0.5-1.5 for carbon isotope fractionation measurement

and for additional RT-qPCR experiments (**Fig. S1**).

Genetic engineering of cyanobacteria

A recombinant strain of *S. elongatus* was constructed by natural transformation using standard

protocols (Clerico et al., 2007) and the plasmids and methods described below (**Table 1**). To

construct the plasmid pSyn02, pAM4937 was digested with *Swa*I to release the *ccdB* toxic gene

and produce a plasmid backbone that contains the pBR322 *E. coli* origin of replication, the base

of mobilization site for conjugal transfer, the *aph1* gene conferring kanamycin resistance, and

sequences for homologous recombination into *S. elongatus* chromosome at NS2. The *rbc* operon

was amplified from *S. elongatus* PCC 7942 gDNA with primers F01 and R01 (**Table S4**)

containing 20-nucleotide sequences that overlap with pAM4937 backbone. The resulting DNA

fragments were assembled using the GeneArt™ Seamless Cloning and Assembly Kit (Invitrogen,

Cat. No. A13288). pSyn02 was used to insert the *rbc* operon into NS2 of the wild-type genome in

the strain PCC 7942 through homologous recombination to create the strain of *S. elongatus*, Syn02,

carrying two copies of the *rbc* operon. Plasmid pSyn-01 was constructed by using two primer pairs

F02/R02 and F03/R03 (**Table S4**), 1) to amplify a fragment of pAM4951 that contains the *E. coli*

origin of replication and the site for conjugal transfer, and 2) to amplify the *aadA* gene conferring

spectinomycin/streptomycin resistance. Native *rbc* operon flanking sequences were amplified

from the *S. elongatus* PCC 7942 gDNA with the primer pairs F04/R04 and F05/R05 (**Table S4**)

containing 20-nucleotide sequences that overlap with the pAM4951 fragments.

Transformation was carried out after growing the WT strain in liquid culture at 30 °C with shaking

(120 rpm) and a light intensity of 80 µmol photon$\cdot$m$^{-2}\cdot$s^{-1} until an OD$_{750}$ ~0.7. The cells were

prepared for transformation according to the protocol by Clerico et al. (2007) and plated on BG-

11 agar containing the appropriate antibiotic(s) for recovery. Subsequently, the colonies were

picked using sterile pipette tips, patched onto BG-11 agar containing the appropriate antibiotic(s),

and further incubated to ensure complete chromosome segregation (i.e., incorporation of the *trans-*

gene into all chromosomes). The patched transformants were screened using colony PCR using

the primers F06/R06 and F07/R07 and the genotypes of the engineered strains were confirmed by

Sanger sequencing using the primers F08/R08 (**Fig. S3, Table S4**)

Extraction of total RNA and proteins

Cultures were collected during the exponential growth phase. Cells were pelleted by centrifugation

at $4,700 \times$ g for 10 min at room temperature and resuspended in 10 mL of TE buffer (10 mM Tris,

pH 8.0, 1 mM EDTA). To prepare the crude cell lysate, 8 mL of the cell suspension were pelleted

by centrifugation at 4,700 × g for 10 min at room temperature, resuspended in 500 µL of hot (pre-

warmed to 95 °C) TE buffer supplemented with 1% (w/v) SDS and incubated at 95 °C for 10 min.

The mixture was sonicated at 40% amplitude for 3 × 10 sec with 10 sec intervals and the cell debris

was centrifuged at 17,000 × g for 10 min at room temperature. The supernatant was collected and

stored at -80 °C. Total RNA was extracted from the remaining 2 mL of the cell suspension using

the RNeasy® Protect Bacteria Mini Kit (QIAGEN, Cat. No. 74524) following the manufacturer

instructions.

Analysis of *rbc* operon expression by RT-qPCR

Total RNA was quantified with a NanoDrop spectrophotometer and 1 µg of RNA was treated with

the amplification grade deoxyribonuclease I (Invitrogen, Cat. No. 18068-015). 10 µL of DNase I-

treated RNA was then used for reverse transcription (RT) performed with the SuperScript™ IV

First-Strand Synthesis System (Invitrogen, Cat. No. 18091050). The four pairs of qPCR primers

(listed in **Table S4**) were designed with Primer3Plus (http://www.bioinformatics.nl/cgi-

bin/primer3plus/primer3plus.cgi). Both reference genes have previously been shown to be stably

expressed under diverse conditions in *S. elongatus* (Luo et al., 2019). The quality of cDNA and

primer specificity was assessed by PCR using cDNA templates (RT positive reactions), RT

negative controls, and the qPCR primers. The analysis of gene expression levels was performed in

a real-time thermal cycler qTOWER[3] G (Analytik Jena AG), equipped with qPCRsoft software,

using the cycles: 50 °C/2 min, 95 °C/2 min, 40 × (95 °C/15 sec, 60 °C/1 min). The relative

expression of the *rbc* operon genes (*rbcL* and *purK*) was calculated as the average fold change

normalized to *secA or ppc* reference genes using the delta-delta Ct method. The experiment was

carried out using three biological replicates and three technical replicates.

Analysis of RbcL protein by Western blot

Total protein concentration in the crude cell lysates was measured using the Pierce™ BCA Protein

Assay Kit (Thermo Scientific, Cat. No. 23225). The lysates were loaded in the amount of 5 µg of

total protein in Laemmli sample buffer onto a 6% (v/v) polyacrylamide stacking gel. Proteins were

electrophoresed in a 12% polyacrylamide resolving gel in TGS buffer and blotted in transfer buffer

onto a PVDF membrane. Total protein load in each sample was visualized by Revert™ 700 Total

Protein Stain (LI-COR Biosciences, Cat. No. 926-11011) and used for RbcL signal normalization.

Detection of RbcL was performed by overnight incubation of the membrane at 4 °C with rabbit

anti-RbcL antibody (Agrisera, Cat. No. AS03 037), 1:5000 in TBST with 5% non-fat milk,

followed by one-hour incubation at room temperature with IRDye® 800CW goat anti-rabbit IgG

secondary antibody (LI-COR Biosciences, Cat. No. 926-32211), 1:20,000 in Intercept® (TBS)

blocking buffer (LI-COR Biosciences, Cat. No. 927-60001) with 0.1% (v/v) Tween-20 and 0.01%

(w/v) SDS. Both the total protein load and the amount of RbcL in each sample were documented

with Odyssey® Fc Imaging System (LI-COR Biosciences, Cat. No. 2800-03) at the near-infrared

detection mode. The images were acquired using Image Studio™ software. The densitometric

analysis of RbcL signal intensity, normalized to total protein load, was performed with Quantity

One® software (Bio-Rad) for six biological replicates. The amount of RbcL produced by each

replicate of the Syn02 strain culture was compared to the averaged level of RbcL in the WT PCC

7942 replicate cultures and expressed as the averaged percent of RbcL synthesized by the WT

strain.

Assembly of RuBisCO subunits

Assembly of the RuBisCO large and small subunits into a hexadecameric complex in each strain

was evaluated by native gel electrophoresis and immunodetection. Samples were collected during

the exponential growh phase. Cells grown under 2% CO_2 were pelleted by centrifugation at 4,700

$\times$ g for 10 min at room temperature and resuspended in 400 µL of native lysis buffer (50 mM Tris,

pH 8.0, 150 mM NaCl, 1 mM EDTA, 10% (v/v) glycerol) supplemented with 5 mM DTT, 100

µg/mL lysozyme from chicken egg white, and 1% (v/v) Halt™ Protease Inhibitor Cocktail

(Thermo Scientific, Cat. No. 78430). The cell suspensions were incubated at 30 °C for 15 min and

subjected to five consecutive freeze-thaw cycles (10 min at -80 °C followed by 5 min at 30 °C),

then were sonicated on ice for 3 minutes at 30% amplitude (2-sec on/off intervals), centrifuged at

17,000 $\times$ g for 15 min at 4 °C. The concentration of total soluble proteins in the lysates was

determined with a Pierce™ BCA Protein Assay Kit. The lysates were adjusted to 5 µg of total

soluble proteins in native sample buffer and then loaded onto a 4-20% Mini-PROTEAN® TGX™

Precast Protein Gel (Bio-Rad, Cat. No. 4561094). Protein electrophoreses were performed in TG

buffer (60 mM Tris, 192 mM glycine) at 100 V for 4 h at 4 °C and blotted in transfer buffer (48

mM Tris, pH 9.2, 39 mM glycine, 0.04% (w/v) SDS) onto a nitrocellulose membrane. After three

10-min washes in wash buffer (48 mM Tris, pH 9.2, 39 mM glycine, 20% (v/v) methanol), total

protein load in each sample was visualized by Revert™ 700 Total Protein Stain and used for the

normalization of RuBisCO complex quantity. Immunodetection of the RuBisCO complex was

performed with the same primary and secondary antibodies that were used to analyze the level of

RbcL, as described above.

Catalytic activity of RuBisCO

The activity of RuBisCO in cyanobacterial lysates was measured using a spectrophotometric

coupled-enzyme assay that links this activity with the rate of NADH oxidation (Kubien, Brown,

& Kane, 2011). The cyanobacterial strains were cultured under 2% CO_2, collected, and pelleted

as described above. The pellets were resuspended in 1 mL of ice-cold lysis buffer (50 mM EPPS,

1 mM EDTA, 2 mM DTT, pH 8.0) and transferred into 2 mL screw-capped tubes with Lysing

Matrix B (MP Biomedical) for lysis by bead beating using FastPrep-24™ 5G bead beater (MP

Biomedical) with 4 m/sec for 10 sec, followed by 2-min incubation on ice, repeated six times. The

cell lysates were transferred to new Eppendorf tubes to remove the beads and unbroken cells and

to pellet the thylakoid membrane by centrifugation at $10,000 \times g$ for 1 min and at $20,000 \times g$ for

30 min at 4 °C, sequentially. The resulting clear supernatants containing cytosolic soluble proteins,

including phycobiliproteins and RuBisCO, were used to determine protein concentration by

Pierce™ BCA Protein Assay Kit and to measure RuBisCO activity by employing an assay adapted

from Kubien et al. (2011). The assay buffer (100 mM HEPES, 25 mM $MgCl_2$, 1 mM EDTA, pH

7.6) was used considering the high Michaelis constant for CO_2 (K_C) for cyanobacterial RuBisCO.

20 µL of cell lysates were preincubated in the assay mix (with 5 mM $NaHCO_3$) at 25 °C for

activation before initiating the reaction by adding synthesized ribulose 1,5-bisphosphate (RuBP)

according to Kane et al. (1998) . The absorbance at 340 nm was monitored using a Synergy H1

plate reader (BioTek). RuBisCO activity was reported as RuBP consumption rate normalized to

total soluble protein content. The assay was performed for three biological replicates.

Cyanobacterial growth measurements

OD_{750} values were plotted as a function of time and analyzed in R with the Growthcurver package.

Growth curve data was fitted to the standard form of the logistic equation to calculate growth

parameters including growth rate, doubling time, and carrying capacity (Sprouffske & Wagner,

2016). Each strain was grown in triplicate for every condition.

Photosynthetic oxygen evolution rate

S. elongatus strain photosynthetic activity was assayed using a Clark-type oxygen electrode

chamber to measure the level of molecular oxygen produced in cyanobacterial cultures. Cells were

grown in 50 mL of BG-11 at 30 °C, illumination of 80 µmol photon·m^{-2}·s^{-1}, shaking at 120 rpm,

in ambient air, and with culture sparging. The samples were collected from triplicate cultures

during the exponential growth phase, pelleted by centrifugation at $4,700 \times g$ for 10 min at room

temperature, and resuspended in fresh BG-11 to an OD_{750} of ~1.0. Oxygen evolution rates were

normalized to chlorophyll *a* concentration (following (Liang & Lindblad, 2017) using the protocol

by Zavřel et al. (2015). The remaining suspension was incubated in the dark for 20 min with gentle

agitation. Samples from each suspension, prepared in three technical replicates, were analyzed in

an oxygen electrode chamber under saturated light, using the Oxygraph+ System (Hansatech

Instruments) equipped with the OxyTrace+ software. Oxygen evolution rate was monitored for 10

min and expressed as nanomoles of molecular oxygen evolved per hour per microgram of

chlorophyll *a*.

Carbon isotope fractionation in bulk cyanobacterial biomass

Carbon isotope fractionation experiments were conducted under ambient and 2% CO_2 conditions.

Cyanobacteria were cultured in shaken flasks (all other conditions as described above) to an OD_{750}

of 0.5 to 1.5 (seven and four biological replicates per strain for experiments conducted in air and

2% CO_2, respectively). Cells were pelleted by centrifugation at $4,700 \times g$ for 10 min at room

temperature and washed in 10 mL of 10 mM NaCl. The bacteria were resuspended in 1 mL of 10

mM NaCl and transferred to Eppendorf tubes. After centrifugation at $4,700 \times g$ for 10 min at room

temperature, the supernatants were completely removed, the pellets were dried in opened tubes in

a laboratory oven at 50 °C for 2 days, and the resultant dried biomass samples were transferred

into tin capsules. The pH of growth medium aliquots was measured at the beginning (sampled

prior to inoculation) and end of cyanobacterial culture (sampled from the supernatant of pelleted

cells, see above). Aliquots sampled at the end of the culture period were then filter-sterilized (0.2

μm), transferred to Extetainer vials leaving no headspace, and stored at 4 °C for isotopic analysis

of dissolved inorganic carbon (DIC). Finally, internal incubator gas was sampled for ambient and

2% CO_2 experiments, as well as CO_2 gas cylinders used to mix 2% CO_2, following the sample

collection protocol provided by the UC Davis Stable Isotope Facility.

The carbon isotope composition of bulk biomass ($\delta^{13}C_{biomass}$), DIC ($\delta^{13}C_{DIC}$), and CO_2 ($\delta^{13}C_{CO2}$)

samples was determined at the UC Davis Stable Isotope Facility. $\delta^{13}C_{biomass}$ was analyzed using a

PDZ Europa ANCA-GSL elemental analyzer interfaced to a PDZ Europa 20-20 isotope ratio mass

spectrometer (Sercon Ltd.). $\delta^{13}C_{DIC}$ (prepared by CO_2 gas evolution using 85% phosphoric acid)

and $\delta^{13}C_{CO2}$ was measured by a GasBench II system interfaced to a Delta V Plus IRMS (Thermo

Scientific). All carbon isotopic composition values are reported relative to the Vienna PeeDee

Belemnite standard (V-PDB):

$$\delta^{13}C_{sample} = \left(\frac{^{13}C/^{12}C_{sample}}{^{13}C/^{12}C_{V-PDB}} - 1 \right) \times 1000$$

545 The carbon isotope fractionation associated with photosynthetic CO_2 fixation (ε_p) was calculated

546 from measured $\delta^{13}C_{biomass}$ and $\delta^{13}C_{CO2}$ according to Freeman and Hayes (1992):

$$\varepsilon_p = \frac{\delta^{13}C_{CO_2} - \delta^{13}C_{biomass}}{1 + \delta^{13}C_{biomass}/1000}$$

Statistical analyses

Results for experimental analyses were presented as the mean and the sample standard deviation

(SD) values of at least three independent experiments. Statistical significance was analyzed with

the two-tailed *t*-test. The unpaired sample *t*-test assuming equal variances was used to compare the

values obtained for different cyanobacterial strains and the paired sample *t*-test was used to

compare the values for the same strain at different experimental conditions.

REFERENCES

Andersson, C. R., Tsinoremas, N. F., Shelton, J., Lebedeva, N. V., Yarrow, J., Min, H., &

Golden, S. S. (2000). Application of bioluminescence to the study of circadian rhythms in

cyanobacteria. *Methods Enzymol, 305*, 527-542.

Atsumi, S., Higashide, W., & Liao, J. C. (2009). Direct photosynthetic recycling of carbon

dioxide to isobutyraldehyde. *Nat Biotechnol, 27*(12), 1177-1180. doi:10.1038/nbt.1586

Bar-On, Y. M., & Milo, R. (2019). The global mass and average rate of rubisco. *Proc Natl Acad*

Sci USA, 10, 4738-4743. doi:10.1073/pnas.1816654116

Bidigare, R. R., Fluegge, A., Freeman, K. H., Hanson, K. L., Hayes, J. M., Hollander, D., . . .

Wakeham, S. G. (1997). Consistent fractionation of ^{13}C in nature and in the laboratory:

growth-rate effects in some haptophyte algae. *Global Biogeochem Cycles, 11*(2), 279-

292. doi:10.1029/96gb03939

Boller, A. J., Thomas, P. J., Cavanaugh, C. M., & Scott, K. M. (2011). Low stable carbon isotope

fractionation by coccolithophore RubisCO. *Geochimica et Cosmochimica Acta, 75*(22),

7200-7207. doi:10.1016/j.gca.2011.08.031

Bonfil, D. J., Ronen-Tarazi, M., Sultemeyer, D., Lieman-Hurwitz, J., Schatz, D., & Kaplan, A.

(1998). A putative HCO_3^- transporter in the cyanobacterium Synechococcus sp. strain

PCC 7942. *FEBS Lett, 430*(3), 236-240.

Burnap, R., Hagemann, M., & Kaplan, A. (2015). Regulation of CO_2 Concentrating Mechanism

in Cyanobacteria. *Life, 5*(1), 348-371. doi:10.3390/life5010348

Carroll, S. B. (2005). Evolution at two levels: on genes and form. *PLoS Biol, 3*(7), e245.

doi:10.1371/journal.pbio.0030245

Catling, D. C., & Zahnle, K. J. (2020). The Archean atmosphere. *Science Advances, 6*(9),

eaax1420. doi:10.1126/sciadv.aax1420

Clerico, E. M., Ditty, J. L., & Golden, S. S. (2007). Specialized techniques for site-directed

mutagenesis in cyanobacteria. *Methods Mol Biol, 362*, 155-171. doi:10.1007/978-1-

59745-257-1_11

De Porcellinis, A. J., Norgaard, H., Brey, L. M. F., Erstad, S. M., Jones, P. R., Heazlewood, J. L.,

& Sakuragi, Y. (2018). Overexpression of bifunctional fructose-1,6-

bisphosphatase/sedoheptulose-1,7-bisphosphatase leads to enhanced photosynthesis and

global reprogramming of carbon metabolism in Synechococcus sp. PCC 7002. *Metab

Eng, 47*, 170-183. doi:10.1016/j.ymben.2018.03.001

Deleens, E., Treichel, I., & O'Leary, M. H. (1985). Temperature dependence of carbon isotope

fractionation in CAM plants. *Plant Physiology, 79*(1), 202-206. doi:10.1104/pp.79.1.202

Des Marais, D. J. (2001). Isotopic evolution of the biogeochemical carbon cycle during the

Precambrian. *Reviews in Mineralogy and Geochemistry, 43*(1), 555-578.

doi:10.2138/gsrmg.43.1.555

Eichner, M., Thoms, S., Kranz, S. A., & Rost, B. (2015). Cellular inorganic carbon fluxes in

Trichodesmium: a combined approach using measurements and modelling. *Journal of

Experimental Botany, 66*(3), 749-759. doi:10.1093/jxb/eru427

Ellis, R. J. (1979). The most abundant protein in the world. *Trends in Biochemical Sciences,

4*(11), 241-244.

Erb, T. J., & Zarzycki, J. (2018). A short history of RubisCO: the rise and fall (?) of Nature's

predominant CO_2 fixing enzyme. *Curr Opin Biotechnol, 49*, 100-107.

doi:10.1016/j.copbio.2017.07.017

Erwin, D. H. (2020). Evolutionary dynamics of gene regulation. *Curr Top Dev Biol, 139*, 407-

431. doi:10.1016/bs.ctdb.2020.02.006

Falkowski, P. G., Fenchel, T., & Delong, E. F. (2008). The microbial engines that drive Earth's

biogeochemical cycles. *Science, 320*(5879), 1034-1039. doi:10.1126/science.1153213

Fay, J. C., & Wittkopp, P. J. (2008). Evaluating the role of natural selection in the evolution of

gene regulation. *Heredity (Edinb), 100*(2), 191-199. doi:10.1038/sj.hdy.6801000

Field, C. B. (1998). Primary Production of the Biosphere: Integrating Terrestrial and Oceanic

Components. *Science, 281*(5374), 237-240. doi:10.1126/science.281.5374.237

Freeman, K. H., & Hayes, J. M. (1992). Fractionation of carbon isotopes by phytoplankton and

estimates of ancient CO_2 levels. *Global Biogeochemical Cycles, 6*(2), 185-198.

doi:10.1029/92gb00190

Gabay, C., Lieman-Hurwitz, J., Hassidim, M., Ronen-Tarazi, M., & Kaplan, A. (1998).

Modification of topA in Synechococcus sp. PCC 7942 resulted in mutants capable of

growing under low but not high concentration of CO2. *FEMS Microbiology Letters,*

159(2), 343-347. doi:10.1111/j.1574-6968.1998.tb12881.x

Garcia, A. K., Cavanaugh, C. M., & Kacar, B. (2021). The curious consistency of carbon

biosignatures over billions of years of Earth-life coevolution. *ISME J.*

doi:10.1038/s41396-021-00971-5

Gesch, R. W., Kang, I. H., Gallo-Meagher, M., Vu, J. C. V., Boote, K. J., H. Allen, L., & Bowes,

G. (2003). Rubisco expression in rice leaves is related to genotypic variation of

photosynthesis under elevated growth CO_2 and temperature. *Plant, Cell & Environment,*

26(12), 1941-1950. doi:10.1046/j.1365-3040.2003.01110.x

Guy, R. D., Fogel, M. L., & Berry, J. A. (1993). Photosynthetic fractionation of the stable

isotopes of oxygen and carbon. *Plant Physiol, 101*(1), 37-47. doi:10.1104/pp.101.1.37

Hamilton, T. L., Bryant, D. A., & Macalady, J. L. (2016). The role of biology in planetary

evolution: cyanobacterial primary production in low-oxygen Proterozoic oceans.

Environmental Microbiology, 18(2), 325-340. doi:10.1111/1462-2920.13118

Havig, J. R., Hamilton, T. L., Bachan, A., & Kump, L. R. (2017). Sulfur and carbon isotopic

evidence for metabolic pathway evolution and a four-stepped Earth system progression

across the Archean and Paleoproterozoic. *Earth-Science Reviews, 174*, 1-21.

doi:10.1016/j.earscirev.2017.06.014

Hayes, J. M. (1993). Factors controlling ^{13}C contents of sedimentary organic compounds:

Principles and evidence. *Marine Geology, 113*(1-2), 111-125. doi:10.1016/0025-

3227(93)90153-m

Hayes, J. M. (2001). Fractionation of carbon and hydrogen isotopes in biosynthetic processes.

Reviews in Mineralogy and Geochemistry, 43(1), 225-277. doi:10.2138/gsrmg.43.1.225

Hinga, K. R., Arthur, M. A., Pilson, M. E. Q., & Whitaker, D. (1994). Carbon isotope

fractionation by marine phytoplankton in culture: the effects of CO_2 concentration, pH,

temperature, and species. *Global Biogeochemical Cycles, 8*(1), 91-102.

doi:10.1029/93gb03393

Hood, R. D., Higgins, S. A., Flamholz, A., Nichols, R. J., & Savage, D. F. (2016). The stringent

response regulates adaptation to darkness in the cyanobacterium Synechococcus

elongatus. *Proc Natl Acad Sci U S A, 113*(33), E4867-4876.

doi:10.1073/pnas.1524915113

Hurley, S. J., Wing, B. A., Jasper, C. E., Hill, N. C., & Cameron, J. C. (2021). Carbon isotope

evidence for the global physiology of Proterozoic cyanobacteria. *Science Advances, 7*(2).

doi:10.1126/sciadv.abc8998

Iwaki, T., Haranoh, K., Inoue, N., Kojima, K., Satoh, R., Nishino, T., . . . Wadano, A. (2006).

Expression of foreign type I ribulose-1,5-bisphosphate carboxylase/ oxygenase (EC

4.1.1.39) stimulates photosynthesis in cyanobacterium Synechococcus PCC7942 cells.

Photosynth Res, 88(3), 287-297. doi:10.1007/s11120-006-9048-x

Kacar, B., Hanson-Smith, V., Adam, Z. R., & Boekelheide, N. (2017). Constraining the timing

of the Great Oxidation Event within the Rubisco phylogenetic tree. *Geobiology, 15*(5),

628-640. doi:10.1111/gbi.12243

Kane, H. J., Wilkin, J.-M., Portis, A. R., & John Andrews, T. (1998). Potent Inhibition of

Ribulose-Bisphosphate Carboxylase by an Oxidized Impurity in Ribulose-1,5-

Bisphosphate. *Plant Physiology, 117*(3), 1059-1069. doi:10.1104/pp.117.3.1059

Kanno, M., Carroll, A. L., & Atsumi, S. (2017). Global metabolic rewiring for improved CO2

fixation and chemical production in cyanobacteria. *Nature Communications, 8*(1), 14724.

doi:10.1038/ncomms14724

Kedzior, M., Garcia, A. K., Li, M., Taton, A., Adam, Z. R., Young, J. N., & Kacar, B. (2022).

Resurrected Rubisco suggests uniform carbon isotope signatures over geologic time. *Cell

Rep, 39*(4), 110726. doi:10.1016/j.celrep.2022.110726

Knoll, A. H., & Nowak, M. A. (2017). The timetable of evolution. *Sci Adv, 3*(5), e1603076.

doi:10.1126/sciadv.1603076

Krissansen-Totton, J., Buick, R., & Catling, D. C. (2015). A statistical analysis of the carbon

isotope record from the Archean to Phanerozoic and implications for the rise of oxygen.

American Journal of Science, 315(4), 275-316. doi:10.2475/04.2015.01

Kuan, D., Duff, S., Posarac, D., & Bi, X. (2015). Growth optimization ofSynechococcus

elongatusPCC7942 in lab flasks and a 2-D photobioreactor. *The Canadian Journal of*

Chemical Engineering, 93(4), 640-647. doi:10.1002/cjce.22154

Kubien, D. S., Brown, C. M., & Kane, H. J. (2011). Quantifying the amount and activity of

Rubisco in leaves. *Methods Mol Biol, 684*, 349-362. doi:10.1007/978-1-60761-925-3_27

Laws, E. A., Bidigare, R. R., & Popp, B. N. (1997). Effect of growth rate and CO_2 concentration

on carbon isotopic fractionation by the marine diatom *Phaeodactylum tricornutum*.

Limnology and Oceanography, 42(7), 1552-1560. doi:10.4319/lo.1997.42.7.1552

Laws, E. A., Popp, B. N., Bidigare, R. R., Kennicutt, M. C., & Macko, S. A. (1995). Dependence

of phytoplankton carbon isotopic composition on growth rate and $[CO_2]_{aq}$: Theoretical

considerations and experimental results. *Geochimica et Cosmochimica Acta, 59*(6), 1131-

1138. doi:10.1016/0016-7037(95)00030-4

Laws, E. A., Popp, B. N., Cassar, N., & Tanimoto, J. (2002). ^{13}C discrimination patterns in

oceanic phytoplankton: likely influence of CO_2 concentrating mechanisms, and

implications for palaeoreconstructions. *Functional Plant Biology, 29*(3).

doi:10.1071/pp01183

Lechno-Yossef, S., Rohnke, B. A., Belza, A. C. O., Melnicki, M. R., Montgomery, B. L., &

Kerfeld, C. A. (2020). Cyanobacterial carboxysomes contain an unique rubisco-activase-

like protein. *New Phytologist, 225*(2), 793-806. doi:10.1111/nph.16195

Liang, F., & Lindblad, P. (2017). Synechocystis PCC 6803 overexpressing RuBisCO grow faster

with increased photosynthesis. *Metab Eng Commun, 4*, 29-36.

doi:10.1016/j.meteno.2017.02.002

Lloyd, M. K., McClelland, H. L. O., Antler, G., Bradley, A. S., Halevy, I., Junium, C. K., . . .

Zerkle, A. L. (2020). The Isotopic Imprint of Life on an Evolving Planet. *Space Science*

Reviews, 216(7). doi:10.1007/s11214-020-00730-6

Luo, X., Li, J., Chang, T., He, H., Zhao, Y., Yang, X., . . . Xu, Y. (2019). Stable Reference Gene

Selection for RT-qPCR Analysis in Synechococcus elongatus PCC 7942 under Abiotic

Stresses. *Biomed Res Int, 2019*, 7630601. doi:10.1155/2019/7630601

Lyons, T. W., Fike, D. A., & Zerkle, A. (2015). Emerging Biogeochemical Views of Earth's

Ancient Microbial Worlds. *Elements, 11*(6), 415-421. doi:10.2113/gselements.11.6.415

Lyons, T. W., Reinhard, C. T., & Planavsky, N. J. (2014). The rise of oxygen in Earth's early

ocean and atmosphere. *Nature, 506*(7488), 307-315. doi:10.1038/nature13068

Maeda, S., Price, G. D., Badger, M. R., Enomoto, C., & Omata, T. (2000). Bicarbonate binding

activity of the CmpA protein of the cyanobacterium Synechococcus sp. strain PCC 7942

involved in active transport of bicarbonate. *J Biol Chem, 275*(27), 20551-20555.

doi:10.1074/jbc.M003034200

Mizutani, H., & Wada, E. (1982). Effect of high atmospheric CO2 concentration on δ13 of algae.

Origins of life, 12(4), 377-390. doi:10.1007/BF00927070

Nisbet, E. G., Grassineau, N. V., Howe, C. J., Abell, P. I., Regelous, M., & Nisbet, R. E. R.

(2007). The age of Rubisco: the evolution of oxygenic photosynthesis. *Geobiology, 5*,

311-335. doi:10.1111/j.1472-4669.2007.00127.x

Olson-Manning, C. F., Wagner, M. R., & Mitchell-Olds, T. (2012). Adaptive evolution:

evaluating empirical support for theoretical predictions. *Nat Rev Genet, 13*(12), 867-877.

doi:10.1038/nrg3322

Omata, T., Gohta, S., Takahashi, Y., Harano, Y., & Maeda, S. (2001). Involvement of a CbbR

homolog in low CO_2-induced activation of the bicarbonate transporter operon in

cyanobacteria. *J Bacteriol, 183*(6), 1891-1898. doi:10.1128/JB.183.6.1891-1898.2001

Onizuka, T., Akiyama, H., Endo, S., Kanai, S., Hirano, M., Tanaka, S., & Miyasaka, H. (2002).

CO_2 Response Element and Corresponding trans-acting Factor of the Promoter for

Ribulose-1,5-bisphosphate Carboxylase/oxygenase Genes in Synechococcus sp.

PCC7002 Found by an Improved Electrophoretic Mobility Shift Assay. *Plant and Cell*

Physiology, 43(6), 660-667. doi:10.1093/pcp/pcf082

Poudel, S., Pike, D. H., Raanan, H., Mancini, J. A., Nanda, V., Rickaby, R. E. M., & Falkowski,

P. G. (2020). Biophysical analysis of the structural evolution of substrate specificity in

RuBisCO. *Proceedings of the National Academy of Sciences, 117*(48), 30451-30457.

doi:10.1073/pnas.2018939117

Price, G. D., Badger, M. R., Woodger, F. J., & Long, B. M. (2008). Advances in understanding

the cyanobacterial CO_2-concentrating-mechanism (CCM): functional components, Ci

transporters, diversity, genetic regulation and prospects for engineering into plants. *J Exp*

Bot, 59(7), 1441-1461. doi:10.1093/jxb/erm112

Rau, G. H., Riebesell, U., & Wolf-Gladrow, D. (1996). A model of photosynthetic [13]C

fractionation by marine phytoplankton based on diffusive molecular CO_2 uptake. *Marine*

Ecology Progress Series, 133, 275-285. doi:10.3354/meps133275

Raven, J. A. (2013). Rubisco: still the most abundant protein of Earth? *New Phytologist, 198*(1),

1-3. doi:10.1111/nph.12197

Raven, J. A., Cockell, C. S., & De La Rocha, C. L. (2008). The evolution of inorganic carbon

concentrating mechanisms in photosynthesis. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, 363(1504), 2641-2650. doi:doi:10.1098/rstb.2008.0020

Riebesell, U., Revill, A. T., Holdsworth, D. G., & Volkman, J. K. (2000). The effects of varying

CO_2 concentration on lipid composition and carbon isotope fractionation in Emiliania

huxleyi. *Geochimica et Cosmochimica Acta, 64*(24), 4179-4192. doi:10.1016/s0016-

7037(00)00474-9

Rillema, R., MacCready, J. S., & Vecchiarelli, A. G. (2020). Cyanobacterial growth and

morphology are influenced by carboxysome positioning and temperature.

doi:10.1101/2020.06.01.127845

Rippka, R., Deruelles, J., Waterbury, J. B., Herdman, M., & Stanier, R. Y. (1979). Generic

Assignments, Strain Histories and Properties of Pure Cultures of Cyanobacteria.

Microbiology, 111(1), 1-61. doi:https://doi.org/10.1099/00221287-111-1-1

Robinson, J. J., Scott, K. M., Swanson, S. T., O'Leary, M. H., Horken, K., Tabita, F. R., &

Cavanaugh, C. M. (2003). Kinetic isotope effect and characterization of form II RubisCO

from the chemoautotrophic endosymbionts of the hydrothermal vent tubeworm *Riftia*

pachyptila. *Limnology and Oceanography, 1*, 48-54. doi:10.4319/lo.2003.48.1.0048

Roeske, C. A., & O'Leary, M. H. (1984). Carbon isotope effects on the enzyme-catalysed

carboxylation of ribulose bisphosphate. *Biochemistry, 23*, 6275-6284.

Schidlowski, M. (1988). A 3,800-Million-Year Isotopic Record of Life from Carbon in

Sedimentary-Rocks. *Nature, 333*(6171), 313-318. doi:10.1038/333313a0

Schidlowski, M. (2001). Carbon isotopes as biogeochemical recorders of life over 3.8 Ga of

Earth history: evolution of a concept. *Precambrian Research, 106*(1-2), 117-134. doi:Doi

10.1016/S0301-9268(00)00128-5

Schirrmeister, B. E., Sanchez-Baracaldo, P., & Wacey, D. (2016). Cyanobacterial evolution

during the Precambrian. *International Journal of Astrobiology, 15*(3), 187-204.

doi:10.1017/s1473550415000579

Schopf, J. W. (2011). The paleobiological record of photosynthesis. *Photosynth Res, 107*(1), 87-

101. doi:10.1007/s11120-010-9577-1

Schubert, B. A., & Jahren, A. H. (2012). The effect of atmospheric CO_2 concentration on carbon

isotope fractionation in C_3 land plants. *Geochimica et Cosmochimica Acta, 96*, 29-43.

doi:10.1016/j.gca.2012.08.003

Schwarz, D., Nodop, A., Huge, J., Purfurst, S., Forchhammer, K., Michel, K. P., . . . Hagemann,

M. (2011). Metabolic and transcriptomic phenotyping of inorganic carbon acclimation in

the Cyanobacterium Synechococcus elongatus PCC 7942. *Plant Physiol, 155*(4), 1640-

1655. doi:10.1104/pp.110.170225

Scott, K. M., Henn-Sax, M., Harmer, T. L., Longo, D. L., Frame, C. H., & Cavanaugh, C. M.

(2007). Kinetic isotope effect and biochemical characterization of form IA RubisCO from

the marine cyanobacterium *Prochlorococcus marinus* MIT9313. *Limnology and*

Oceanography, 52(5), 2199-2204.

Sengupta, A., Sunder, A. V., Sohoni, S. V., & Wangikar, P. P. (2019). Fine-Tuning Native

Promoters of Synechococcus elongatus PCC 7942 To Develop a Synthetic Toolbox for

Heterologous Protein Expression. *ACS Synth Biol, 8*(5), 1219-1223.

doi:10.1021/acssynbio.9b00066

Smith, R. M., & Williams, S. B. (2006). Circadian rhythms in gene transcription imparted by

chromosome compaction in the cyanobacterium *Synechococcus elongatus*. *Proceedings*

of the National Academy of Sciences, 103(22), 8564-8569. doi:10.1073/pnas.0508696103

Sprouffske, K., & Wagner, A. (2016). Growthcurver: an R package for obtaining interpretable

metrics from microbial growth curves. *BMC Bioinformatics, 17*, 172.

doi:10.1186/s12859-016-1016-7

Sumner, D. Y. (2002). Biology and Geology: A Necessary Symbiosis. *Palaios, 17*(4), 307-308.

doi:10.1669/0883-1351(2002)017<0307:Bagans>2.0.Co;2

Szekeres, E., Sicora, C., Dragos, N., & Druga, B. (2014). Selection of proper reference genes for

the cyanobacterium Synechococcus PCC 7002 using real-time quantitative PCR. *FEMS*

Microbiology Letters, 359(1), 102-109. doi:10.1111/1574-6968.12574

Taton, A., Erikson, C., Yang, Y., Rubin, B. E., Rifkin, S. A., Golden, J. W., & Golden, S. S.

(2020). The circadian clock and darkness control natural competence in cyanobacteria.

Nat Commun, 11(1), 1688. doi:10.1038/s41467-020-15384-9

Taton, A., Unglaub, F., Wright, N. E., Zeng, W. Y., Paz-Yepes, J., Brahamsha, B., . . . Golden, J.

W. (2014). Broad-host-range vector system for synthetic biology and biotechnology in

cyanobacteria. *Nucleic Acids Research, 42*(17), e136-e136. doi:10.1093/nar/gku673

Taylor, T. B., Shepherd, M. J., Jackson, R. W., & Silby, M. W. (2022). Natural selection on

crosstalk between gene regulatory networks facilitates bacterial adaptation to novel

environments. *Curr Opin Microbiol, 67*, 102140. doi:10.1016/j.mib.2022.02.002

Tcherkez, G. G., Farquhar, G. D., & Andrews, T. J. (2006). Despite slow catalysis and confused

substrate specificity, all ribulose bisphosphate carboxylases may be nearly perfectly

optimized. *Proc Natl Acad Sci U S A, 103*(19), 7246-7251. doi:10.1073/pnas.0600605103

Tchernov, D., Helman, Y., Keren, N., Luz, B., Ohad, I., Reinhold, L., . . . Kaplan, A. (2001).

Passive entry of CO2 and its energy-dependent intracellular conversion to HCO3- in

cyanobacteria are driven by a photosystem I-generated deltamuH+. *J Biol Chem, 276*(26),

23450-23455. doi:10.1074/jbc.M101973200

Thomas, P. J., Boller, A. J., Satagopan, S., Tabita, F. R., Cavanaugh, C. M., & Scott, K. M.

(2019). Isotope discrimination by form IC RubisCO from *Ralstonia eutropha* and

Rhodobacter sphaeroides, metabolically versatile members of '*Proteobacteria*' from

aquatic and soil habitats. *Environ Microbiol, 21*(1), 72-80. doi:10.1111/1462-2920.14423

Ungerer, J., Lin, P.-C., Chen, H.-Y., Pakrasi, H. B., Giovannoni, S. J., Burnap, R., & Sherman,

L. (2018). Adjustments to Photosystem Stoichiometry and Electron Transfer Proteins Are

Key to the Remarkably Fast Growth of the Cyanobacterium Synechococcus elongatus

UTEX 2973. *MBio, 9*(1). doi:10.1128/mBio.02327-17

Valentine, J. W. (1994). Late Precambrian bilaterians: grades and clades. *Proceedings of the*

National Academy of Sciences, 91(15), 6751-6757. doi:10.1073/pnas.91.15.6751

Vijayan, V., Jain, I. H., & O'Shea, E. K. (2011). A high resolution map of a cyanobacterial

transcriptome. *Genome Biology, 12*(5), R47. doi:10.1186/gb-2011-12-5-r47

von Caemmerer, S., Tazoe, Y., Evans, J. R., & Whitney, S. M. (2014). Exploiting

transplastomically modified Rubisco to rapidly measure natural diversity in its carbon

isotope discrimination using tuneable diode laser spectroscopy. *J Exp Bot, 65*(13), 3759-

3767. doi:10.1093/jxb/eru036

Wang, S., Yeager, K. M., & Lu, W. (2016). Carbon isotope fractionation in phytoplankton as a

potential proxy for pH rather than for [CO2(aq)]: Observations from a carbonate lake.

Limnology and Oceanography, 61(4), 1259-1270. doi:10.1002/lno.10289

Wilkes, E. B., Lee, R. B. Y., McClelland, H. L. O., Rickaby, R. E. M., & Pearson, A. (2018).

Carbon isotope ratios of coccolith–associated polysaccharides of *Emiliania huxleyi* as a

function of growth rate and CO_2 concentration. *Organic Geochemistry, 119*, 1-10.

doi:10.1016/j.orggeochem.2018.02.006

Wilkes, E. B., & Pearson, A. (2019). A general model for carbon isotopes in red-lineage

phytoplankton: Interplay between unidirectional processes and fractionation by RubisCO.

Geochimica et Cosmochimica Acta, 265, 163-181. doi:10.1016/j.gca.2019.08.043

Wilson, A. C., Maxson, L. R., & Sarich, V. M. (1974). Two Types of Molecular Evolution.

Evidence from Studies of Interspecific Hybridization. *Proceedings of the National*

Academy of Sciences, 71(7), 2843-2847. doi:10.1073/pnas.71.7.2843

Wong, W. W., & Sackett, W. M. (1978). Fractionation of stable carbon isotopes by marine

phytoplankton. *Geochimica et Cosmochimica Acta, 42*(12), 1809-1815.

doi:10.1016/0016-7037(78)90236-3

Woodger, F. J., Badger, M. R., & Price, G. D. (2003). Inorganic carbon limitation induces

transcripts encoding components of the CO(2)-concentrating mechanism in

Synechococcus sp. PCC7942 through a redox-independent pathway. *Plant Physiol,*

133(4), 2069-2080. doi:10.1104/pp.103.029728

Yoshioka, T. (1997). Phytoplanktonic carbon isotope fractionation: equations accounting for

CO2-concentrating mechanisms. *Journal of Plankton Research, 19*(10), 1455-1476.

doi:10.1093/plankt/19.10.1455

Yu, J., Liberton, M., Cliften, P. F., Head, R. D., Jacobs, J. M., Smith, R. D., . . . Pakrasi, H. B.

(2015). Synechococcus elongatus UTEX 2973, a fast growing cyanobacterial chassis for

biosynthesis using light and CO2. *Scientific Reports, 5*(1), 8132. doi:10.1038/srep08132

Zavřel, T., Sinetova, M., & Červený, J. (2015). Measurement of Chlorophyll *a* and Carotenoids

Concentration in Cyanobacteria. *Bio-Protocol, 5*(9). doi:10.21769/BioProtoc.1467

853 Table 1. Strains and plasmids used in this study.

Strain or plasmid	Description/Genotype	Antibiotic resistance	Source/Reference
WT	Wild-type strain of *S. elongatus* PCC 7942	-	Susan S. Golden (UC San Diego)
pAM4937	Expression vector for *S. elongatus* PCC 7942 neutral site 2 (NS2)	Km	(Taton et al., 2014)
pSyn02	pAM4937 carrying the *rbc* operon including *rbcL*, *rbcS*, *purK,* and flanking sequences from *S. elongatus* PCC 7942 (CP000100: 1479071-1484283)	Km	This study
Syn02	*S. elongatus* PCC 7942 carrying a second copy of the *rbc* operon and flanking sequences at NS2: NS2::*aphI-rbcL-rbcS-purK*- Synpcc7942_1429-Synpcc7942_1430	Km	This study
pAM4951	Expression vector for *S. elongatus* PCC 7942 neutral site 1 (NS1)	Sp+Sm	(Taton et al., 2014)
pSyn01	Plasmid to replace *S. elongatus'* native *rbc* operon (CP000100: 1479070-	Sp+Sm	This study

	1482595) with a Sp/Sm resistance gene: Δ(*rbcL-rbcS-purK*)::*aadA*.		
Syn01	*S. elongatus* strain Syn02 with the native *rbc* operon removed: Syn02 and Δ(*rbcL-rbcS-purK*)::*aadA*.	Km, Sp+Sm	This study

856   **Table 2. Growth parameters and oxygen evolution rates of *S. elongatus* strains under varying**

857   **CO_2 concentrations[a].**

Strain	Atmosphere	Growth rate (h^{-1})	Carrying capacity (OD_{750})	Oxygen evolution rate (nmol $O_2 \cdot h^{-1} \cdot \mu g^{-1}$ chlorophyll *a)*
WT	Air	0.017 ± 0.001	6.8 ± 0.7	340 ± 30
	2% CO_2	0.043 ± 0.002	8.4 ± 0.5	-
	5% CO_2	0.047 ± 0.005	5.4 ± 0.3	-
Syn02	Air	$0.020 \pm 0.001*$	7.6 ± 0.5	360 ± 30
	2% CO_2	0.046 ± 0.001	$8.33 \pm 0.$	-
	5% CO_2	0.040 ± 0.001	$5. \pm 0.4$	-

858   [a]Values are means of three biological replicates $\pm$ 1 SD. Asterisks indicate *t*-test result from

859   comparison with WT for the same atmospheric condition, $* - p < 0.05$.

Figure 1. Experimental strategy. A – Overall methodology. 1) Genetic engineering: An additional copy of the *rbc* operon was inserted at the chromosomal NS2 site of *S. elongatu*s PCC 7942 to create the Syn02 strain. 2) RuBisCO characterization was carried out by investigating (a) catalytic activity, (b) assembly into a hexadecameric complex, and (c) expression at both the transcript and protein levels. 3) Physiology: *S. elongatu*s WT and Syn02 strains were evaluated by monitoring microbial growth characteristics and rate of photosynthetic oxygen evolution. 4) Carbon isotope fractionation: The impact of RuBisCO overexpression on biomass $\delta^{13}C$ and ε_p under varying CO_2 concentrations was assessed. B – Map of the genetic insert in *S. elongatus*

872   **strain Syn02**. Plasmid pSyn02 was used to insert the *rbc* operon and the *aph1* gene conferring

873   kanamycin resistance in PCC 7942 WT NS2 to generate the Syn02 strain. Crosslines indicate

874   homologous recombination sites and scale bars show DNA fragment sizes (in bp).

Figure 2. RuBisCO expression, assembly, and activity in *S*. elongatus strains under varying

CO₂ concentrations. A – **Expression of Syn02 *rbcL* relative to WT *rbcL*.** Asterisks indicate *t*-

test results compared to WT *rbcL* expression at the same growth condition. **B – Expression of**

WT *rbcL* at 2 and 5% CO₂ relative to ambient air. Asterisks indicate *t*-test results compared

to WT *rbcL* expression in air. **A-B –** Expression data was measured by RT-qPCR, normalized to

reference genes *secA* and *ppc*. Columns represent mean fold expression for three biological

replicates. **C – Immunodetection of RbcL.** (*left*) Western blot showing RbcL protein detected

by anti-RuBisCO antibody and total protein stain from crude cell lysates. (*right*) RbcL

percentage signal intensities were normalized to that for the total soluble protein load. Columns

represent mean RbcL intensities for six biological replicates, relative to mean values for WT

cultures at the same growth condition. Asterisks indicate *t*-test results compared to WT RbcL. **D**

– Immunodetection of assembled RuBisCO. Western blot showing the proper assembly of

RbcL and RbcS into the L_8S_8 hexadecameric complex L_8S_8 (520 kDa), detected by anti-RbcL

antibody. Rbc L_8S_8 percentage signal intensity was normalized to that for the total soluble

protein load, and is shown relative to WT. **E – Lysate RuBisCO activity.** Activity was

measured by the RuBP consumption rate, normalized to total soluble protein content of cell

lysate. Columns represent the mean activity of three biological replicates. **A-C, E –** Asterisks

indicate *t*-test results for pairwise comparison indicated by horizontal line. Error bars on all

graphs indicate 1 SD. ns – not significant; ** – $p < 0.01$; *** – $p < 0.001$.

Figure 3. Growth of *S. elongatus* strains. A – Growth profiles. Cultures of each strain were

maintained in ambient air or under 2% or 5% CO_2. Smoothed profiles were generated from mean

optical density values (measured at 750 nm, OD_{750})for three replicate cultures per growth

condition. **B – Doubling times.** Columns represent the mean doubling time of three replicates.

Error bars on all graphs indicate 1 SD, and asterisks indicate *t*-test results for pairwise comparisons

indicated by horizontal lines. ns – not significant; * – $p < 0.05$; *** – $p < 0.001$.

Figure 4. $^{13}C/^{12}C$ discrimination associated with photosynthetic CO_2 fixation (ε_p) of *S.*

***elongatus* strains under ambient and 2% CO_2 concentrations.** ε_p values are calculated relative

to measured $\delta^{13}C_{CO2}$ (Table S3; see **Materials and Methods** for calculation). Mean (long

horizontal bars) and 1 SD (vertical error bars) are shown in red, calculated for seven (air) and four

(2% CO_2) biological replicates. Asterisks indicate *t*-test results for pairwise comparisons indicated

by horizontal lines. ns – not significant; *** – $p < 0.001$.