System-level effects of CO_2 and RuBisCO concentration on carbon isotope fractionation

Amanda K. Garcia[1‡], Mateusz Kedzior[1‡], Arnaud Taton[2], Meng Li[3], Jodi N. Young[3] and Betül Kaçar [1,4*]

[1]Department of Molecular and Cellular Biology, University of Arizona, Tucson, AZ

[2]Division of Biological Sciences, University of California San Diego, La Jolla, CA

[3]School of Oceanography, University of Washington, Seattle, WA

[4]Lunar and Planetary Laboratory, Department of Planetary Sciences, University of Arizona, Tucson, AZ

[‡]These authors contributed equally to this work.

*For correspondence: betul@arizona.edu

KEYWORDS

ABSTRACT

Carbon isotope biosignatures preserved in the Precambrian geologic record are primarily interpreted to reflect ancient cyanobacterial carbon fixation catalyzed by Form I RuBisCO enzymes. The average range of isotopic biosignatures generally follows that produced by extant cyanobacteria. However, this observation is difficult to reconcile with several environmental (e.g., temperature, pH, and CO_2 concentrations), molecular and physiological factors that likely would have differed during the Precambrian and can produce fractionation variability in contemporary organisms that meets or exceeds that observed in the geologic record. To test a range of genetic and environmental factors that may have impacted ancient carbon isotope biosignatures, we engineered a mutant strain of the model cyanobacterium *Synechococcus elongatus* PCC 7942 that overexpresses RuBisCO and characterized the resultant physiological and isotope fractionation effects. We specifically investigated how both increased atmospheric CO_2 concentrations and RuBisCO regulation influence cell growth, oxygen evolution rate, and carbon isotope fractionation in cyanobacteria. We found that >2% CO_2 increases the growth rate of wild-type and mutant strains, and that the pool of active RuBisCO enzyme increases with increased expression. At elevated CO_2, carbon isotope discrimination (ε_p) is increased by ~8‰, whereas RuBisCO overexpression does not significantly affect isotopic discrimination at all tested CO_2 concentrations. Our results show that understanding the environmental factors that impact RuBisCO regulation, physiology, and evolution is crucial for reconciling microbially driven carbon isotope fractionation with the geologic record of organic and inorganic carbon isotope signatures.

IMPORTANCE

Carbon isotope biosignatures preserved in the geologic record are interpreted to reflect the long-

term evolution of microbial carbon fixation and provide the earliest evidence of life on Earth.

RuBisCO enzymes, distinctive and early-evolved catalysts that fix atmospheric CO_2, have likely

been responsible for the bulk of primary productivity through Earth history. Thus, a comprehensive

understanding of the molecular, physiological, environmental, and evolutionary factors that

influence the isotopic discrimination of cyanobacteria that utilize RuBisCO is essential for the

interpretation of ancient isotopic biosignatures. For example, the vastly different atmospheric CO_2

levels that characterized the Precambrian may have influenced the expression and regulation of

the ancient RuBisCO protein complex. These observations underscore the need to consider how a

broader range of environmental conditions and subcellular processes may have shaped isotopic

discrimination over geologic time. In this study, we establish a cyanobacterial metabolic-

engineering strategy that can test such hypotheses and offer insights into the biogeochemical

record of life.

INTRODUCTION

The conserved microbial metabolic pathways that drive global biogeochemistry emerged on Earth billions of years ago, the evolution of which has both shaped and been shaped by large-scale environmental transitions (Falkowski, Fenchel, & Delong, 2008; Knoll, Bergmann, & Strauss, 2016; Lyons, Fike, & Zerkle, 2015). These microbial processes have left distinct signatures that evidence biological activity billions of years in the past. Of these, the oldest and most extensive signature of biological activity on Earth is the deviation in stable carbon isotopic compositions ($^{13}C/^{12}C$, expressed as $\delta^{13}C$) between preserved inorganic and organic carbon, interpreted to reflect the isotopic discrimination of ancient biological carbon fixation (Des Marais, 2001; Krissansen-Totton, Buick, & Catling, 2015; Lloyd et al., 2020; Schidlowski, 2001). These deviations are primarily shaped by enzymes that preferentially assimilate the lighter ^{12}C isotope from inorganic carbon sources. Carbon biosignatures preserved in the geologic record therefore reflect the long-term evolution of these enzyme-mediated processes and their hosts' physiologies.

The RuBisCO enzyme (ribulose 1,5-bisphosphate (RuBP) carboxylase/oxygenase) catalyzes the reduction of inorganic CO_2 as the initial step of carbon assimilation into organic biomass via the Calvin-Benson-Bassham (CBB) cycle (Erb & Zarzycki, 2018; Nisbet et al., 2007; Tcherkez, Farquhar, & Andrews, 2006). RuBisCO is one of the most abundant proteins on Earth (Bar-On & Milo, 2019; Ellis, 1979; Raven, 2013), and its presence in photoautotrophic organisms, including early-evolved cyanobacteria, suggests that this enzyme has played a significant role in primary production for much of Earth history (Hamilton, Bryant, & Macalady, 2016; Schirrmeister, Sanchez-Baracaldo, & Wacey, 2016; Schopf, 2011). Thus, the isotopic fractionation behavior of RuBisCO is also thought to have primarily constrained Precambrian carbon isotope signatures

preserved in the geologic record (Garcia, Cavanaugh, & Kacar, 2021; Schidlowski, 1988). Though

there exist multiple forms of RuBisCO, the known range of Form I RuBisCO isotopic fraction (ε

$\approx$ 20-30‰) (R. D. Guy, Fogel, & Berry, 1993; Scott et al., 2007; von Caemmerer, Tazoe, Evans,

& Whitney, 2014) — the form utilized by extant cyanobacteria and responsible for the bulk of

modern primary production (Field, 1998) — is largely consistent with the ~25‰ mean deviation

between preserved inorganic and organic carbon isotopic compositions across geologic time (Des

Marais, 2001; Garcia et al., 2021; Havig, Hamilton, Bachan, & Kump, 2017; Krissansen-Totton et

al., 2015; Lloyd et al., 2020; Schidlowski, 2001).

RuBisCO primarily shapes the generation of distinct carbon isotopic biosignatures associated with

CBB-utilizing organisms. Nonetheless the carbon isotopic composition of bulk photoautotrophic

biomass often deviates from values obtained for purified RuBisCO. This discrepancy might be

attributable to several intracellular physiological and metabolic features that additionally shift the

isotopic composition of organism biomass. These include the activation of carbon-concentrating

mechanisms such as the partitioning of RuBisCO into carboxysomes (Hurley, Wing, Jasper, Hill,

& Cameron, 2021; Laws, Popp, Cassar, & Tanimoto, 2002; Price, Badger, Woodger, & Long,

2008; Raven, Cockell, & De La Rocha, 2008), the diffusive transport of CO_2 (Hayes, 1993; Rau,

Riebesell, & Wolf-Gladrow, 1996), and intermediary carbon-fixing steps (Eungrasamee, Miao,

Incharoensakdi, Lindblad, & Jantaro, 2019; H. I. Guy & Evans, 1996; Hayes, 1993; Rothschild &

DesMarais, 1989). Further, studies show that photosynthetic carbon isotope discrimination (ε_p)

may vary due to environmental factors and cellular physiological responses, including temperature

(Deleens, Treichel, & O'Leary, 1985; Wong & Sackett, 1978), pH (Hinga, Arthur, Pilson, &

Whitaker, 1994; Roeske & O'Leary, 1984), growth rate (Bidigare et al., 1997; Laws, Bidigare, &

Popp, 1997), and CO_2 concentration (Eichner, Thoms, Kranz, & Rost, 2015; Freeman & Hayes,

1992; Hinga et al., 1994; Hurley et al., 2021; Schubert & Jahren, 2012; Wilkes, Lee, McClelland,

Rickaby, & Pearson, 2018). These variations indicate a compelling need to comprehensively

characterize both internal and external factors that can affect isotope fractionation, though the

study of each tend to be siloed in biological and geobiological fields, respectively.

An aspect of photoautotrophic isotopic discrimination that has not been thoroughly investigated is

how variable RuBisCO expression can influence host organism growth parameters and the isotopic

composition of assimilated biomass. RuBisCO expression has been shown to be CO_2-sensitive

(Gesch et al., 2003; Onizuka et al., 2002; Sengupta, Sunder, Sohoni, & Wangikar, 2019). This

observation is particularly important, considering that atmospheric CO_2 concentrations exceeded

present day levels by more than an order of magnitude for much of Earth history (between ~0.001

and 0.1 bar CO_2 through the Precambrian (Catling & Zahnle, 2020)) and that the RuBisCO

catalytic efficiency itself is sensitive to the atmospheric CO_2/O_2 levels (Erb & Zarzycki, 2018;

Kacar, Hanson-Smith, Adam, & Boekelheide, 2017; Poudel et al., 2020; Riebesell, Revill,

Holdsworth, & Volkman, 2000; Schubert & Jahren, 2012; Scott et al., 2007; Tcherkez et al., 2006;

Wilkes et al., 2018). Thus, interpretation of the carbon isotope record should take into account the

coupled influence of CO_2 concentration and RuBisCO expression in generating isotopic

biosignatures and shaping ancient primary productivity.

To investigate the interplay between RuBisCO expression and CO_2 concentrations on

cyanobacterial fitness and carbon isotopic discrimination, we generated a genetic system to

manipulate the expression levels of RuBisCO in the model organism *Synechococcus elongatus*

PCC 7942 (hereafter *S. elongatus*). *S. elongatus* is a naturally competent obligate photoautotroph

that utilizes Form I RuBisCO to fix carbon (Bonfil et al., 1998; Gabay, Lieman-Hurwitz, Hassidim,

Ronen-Tarazi, & Kaplan, 1998; Maeda, Price, Badger, Enomoto, & Omata, 2000; Omata, Gohta,

Takahashi, Harano, & Maeda, 2001; Taton et al., 2020; Tchernov et al., 2001). Specifically, we

engineered the *S. elongatus* genome with an additional copy of the RuBisCO operon in a

chromosomal neutral site to overexpress RuBisCO and permit genetic manipulation of the

RuBisCO operon for future studies. We confirmed proper assembly of the overexpressed RuBisCO

complex and its catalytic activity in the engineered strain. Then, to assess whether physiological

and subcellular parameters and varying CO_2 concentrations modulate carbon isotopic

discrimination in cyanobacteria, we determined growth rate, photosynthetic oxygen evolution rate,

and $^{13}C/^{12}C$ discrimination of the engineered strain in comparison to wild-type *S. elongatus*. Our

results suggest that increased CO_2 concentrations result in significantly faster growth in

cyanobacteria and increase the magnitude of isotopic discrimination. On the other hand, we show

that changing cellular RuBisCO levels do not significantly alter carbon isotope discrimination.

RESULTS

A second copy of the *rbc* operon results in increased amount of active RuBisCO

The RuBisCO Form I enzyme in *S. elongatus* is encoded by an operon that includes a CO_2-

sensitive promoter region (Sengupta et al., 2019) as well as the structural *rbcL* (large subunit) and

rbcS (small subunit) genes (Vijayan, Jain, & O'Shea, 2011). We designed *S. elongatus* strain Syn02

that harbors the native *rbc* operon and a second copy inserted in the chromosome neutral site 2

(NS2), a site that permits genetic modification without additional indirect phenotypic impact

(Andersson et al., 2000; Clerico, Ditty, & Golden, 2007) (**Fig. 1**). Syn02 was contructed by

transforming wild-type (WT) *S. elongatus* with plasmid pSyn02 carrying the *rbc* operon and

homologous regions directing recombination at NS2 (**Table 1**). Additionally, we generated a

control strain Syn01 whereby RuBisCO is provided solely by an engineered *rbc* operon at its NS2

site.

We evaluated whether the additional copy of the *rbc* operon in strain Syn02 resulted in RuBisCO

overexpression by quantifying transcription of the *rbcL* and *purK* (located downstream of *rbcL* in

the operon) genes by quantitative reverse-transcription PCR (RT-qPCR). Transcripts were

quantified and normalized to those of *secA* and *ppc* reference genes (Hood, Higgins, Flamholz,

Nichols, & Savage, 2016; Luo et al., 2019; Szekeres, Sicora, Dragoş, & Drugă, 2014). We found

that the level of *rbcL* transcript in Syn02 increased by at least 2-fold relative to WT across all

tested CO_2 concentrations and growth phases, and as high as ~14-fold in air ($p < 0.01$; **Fig. 2A,**

Fig. S1A). In addition, we found that *rbcL* expression increased by >5-fold at elevated CO_2

concentrations relative to ambient air ($p < 0.001$; **Fig. 2B, Fig. S1B).** Expression of *purK* in Syn02

increased by >2-fold at 2% and 5% CO_2 relative to WT, but decreased by ~0.5-fold in air (**Fig.**

S1C).

To determine *rbcL* overexpression at the protein-level, we quantified RbcL protein from crude cell

lysates by Western blot using rabbit anti-RbcL antibody (see **Materials and Methods**). In

agreement with RuBisCO overexpression indicated by the RT-qPCR results, densitometric

analyses revealed a mean ~2 to 4-fold increase of RbcL protein in Syn02 relative to WT across all

tested CO_2 concentrations ($p < 0.01$; **Fig. 2C**). Finally, we confirmed proper assembly of the large

(L) and small (S) subunits into the RuBisCO L_8S_8 complex in Syn02 as well as the Syn01 control

strain by native protein electrophoresis and detection by anti-RbcL antibody. We found a ~3-fold

increase in assembled RuBisCO protein for Syn02 and ~2-fold increase for Syn01 relative to WT

for cultures grown in 2% CO_2 (**Fig. 2D**).

Finally, we tested whether or not RuBisCO overexpression in Syn02 resulted in an increased

amount of active RuBisCO by determining the total carboxylase activity of cell lysates. Enzyme

activity was evaluated by an *in vitro* spectrophotometric coupled-enzyme assay that links

carboxylase activity to NADH oxidation, reported as the RuBP consumption rate normalized to

total soluble protein content (Kubien, Brown, & Kane, 2011). We measured a mean ~0.8-fold (p

< 0.01) increase in Syn02 lysate RuBisCO activity relative to WT (**Fig. 2E**). No significant

difference in activity was detected between Syn01 and WT cultures.

RuBisCO overexpression does not strongly influence growth rate or photosynthetic activity

We cultured WT and Syn02 *S. elongatus* strains in order to determine the influence of Syn02

RuBisCO overexpression on growth rate. Cultures were continuously sparged with ambient air, as

well as 2% and 5% CO_2, with daily optical density measurements at 750 nm (OD_{750}) as described

in **Materials and Methods**. WT and Syn02 both exhibited ~2.5-fold faster growth rates under 2%

and 5% CO_2 compared to ambient air ($p < 0.001$; **Fig. 2A; Table 2**). Syn02 exhibited a slight

~0.1-fold increase in growth rate in ambient air relative to WT ($p < 0.05$). No difference in growth

rate was observed between the two strains under 2% and 5% CO_2. The carrying capacity

(maximum cell density measured at OD_{750} across the total growth period) for cultures varied

between different atmospheric conditions, with a maximum carrying capacity of $OD_{750} \approx 8.4$

reached under 2% CO_2. No significant difference in carrying capacity was found between WT and

Syn02 cultures grown under the same atmospheric conditions. To determine the degree to which

sparging impacted growth, we repeated the growth experiments without sparging. Minimal

differences were again observed between WT and Syn02 across different CO_2 concentrations

(**Table S1**).

To further test for differences in photosynthetic activities between the WT and Syn02 strains, we

measured their oxygen evolution rates. After brief incubation in the dark, culture samples were

exposed to saturated light in an oxygen electrode chamber to detect increased levels of molecular

oxygen. Oxygen evolution rates were normalized to chlorophyll a concentrations, following

Zavrel et al. (Zavrel, Sinetova, & Červený, 2015). We did not observe a significant difference in

mean oxygen evolution rate between WT and Syn02 (342 ± 26 $O_2 \cdot h^{-1} \cdot \mu g^{-1}$ chlorophyll a and 362

± 26 $O_2 \cdot h^{-1} \cdot \mu g^{-1}$ chlorophyll a, respectively) (**Table 2**).

CO_2 concentration impacts $S.\ elongatus$ $^{13}C/^{12}C$ fractionation

We tested both the influence of RuBisCO overexpression and CO_2 concentration on the magnitude

of $^{13}C/^{12}C$ isotopic discrimination in $S.\ elongatus,$ which was assessed by measuring the $^{13}C/^{12}C$

composition of biomass ($\delta^{13}C_{biomass}$). The isotopic discrimination associated with photosynthetic

CO_2 fixation (ε_p) was then calculated from measured $\delta^{13}C_{biomass}$ and a reference $\delta^{13}C_{CO2}$ value (8.4

‰; (Keeling et al., 2017)), as described in **Materials and Methods**. We observed that increased

CO_2 concentration increases the magnitude of $S.\ elongatus$ $^{13}C/^{12}C$ discrimination. Mean ε_p values

for $S.\ elongatus$ strains grown in ambient air were ~10‰ (**Fig. 4A**). At 2% and 5% CO_2, ε_p values

were ~7-9‰ greater than those for cultures grown in air, with mean values ranging between ~17‰

to 20‰ ($p < 0.001$). Mean ε_p values were similar for WT and Syn02 strains most tested CO_2

conditions, with the exception of 5% CO2, where a significant but slight difference in ε_p values

was detected (18.95‰ and 18.62‰ for WT and Syn02, respectively; $p < 0.01$). These results

demonstrate that RuBisCO overexpression in Syn02 does not strongly impact cyanobacterial

$^{13}C/^{12}C$ discrimination.

DISCUSSION

In this study, we developed an integrative approach to investigate the coupled impact of CO_2 levels

and RuBisCO expression on cyanobacterial growth, fitness and carbon isotope discrimination. We

tested whether the overexpression of RuBisCO influences *S. elongatus* growth, physiology and

carbon isotope fractionation at different CO_2 atmosphere concentrations by generating an

engineered *S. elongatus* strain Syn02 that harbors a second copy of the *rbc* operon at NS2.

Our transcriptional analyses indicate that *S. elongatus* strain Syn02 overexpresses RuBisCO

relative to WT at all tested CO_2 concentrations (ambient air, 2%, and 5% CO_2) (**Fig. 2A, Fig. 2C**).

In addition, we found that *rbcL* transcript levels increase for all strains at elevated CO_2 levels

relative to air (**Fig. 2B**). This result is in agreement with a previous study reporting a regulatory

element in the *S. elongatus* PCC 7942 *rbcL* operon that regulates the expression of RbcL with

increasing CO_2 (Sengupta et al., 2019). We found that the magnitude of increased *rbcL*

transcription between Syn02 and WT was lower at elevated CO_2 (**Fig. 2A**). For example, in air,

rbcL transcript levels in Syn02 were ~14-fold greater than in WT, but in 2% CO_2, only a ~2-fold

increase was observed. These results indicate that, whereas RuBisCO expression is impacted by

CO_2 concentration, other physiological factors such as the translation efficiency or expression of

other ancillary crucial proteins may ~~may~~ be limiting RuBisCO expression and dampen the effects

of an additional copy of the *rbc* operon in Syn02.

Protein-level analyses indicate that increased *rbc* transcription in Syn02 relative to WT does result

in increased cellular RuBisCO concentration. The magnitude of this increase is relatively constant

(~2-3 fold) across different CO_2 concentrations (**Fig. 2C**). These results are supported by a ~2-fold

increase in assembled RuBisCO L_8S_8 protein in Syn02 relative to WT (**Fig. 3A**). However, the

~0.8-fold increase in carboxylase activity of Syn02 cell lysate relative to that of WT was not

directly proportional to the increase in protein amount observed by Western blot analyses,

suggesting that not all overexpressed RuBisCO was catalytically active (**Fig. 3B**). Previous studies

showed overexpression of RuBisCO in cyanobacteria generally produces an increased pool of

active carboxylase (Atsumi, Higashide, & Liao, 2009; Iwaki et al., 2006; Lechno-Yossef et al.,

2020; Liang & Lindblad, 2017). However, it is possible that not all overexpressed RuBisCO may

be properly assembled, e.g., due to limited chaperonin GroEL/ES, which is known to be involved

in RuBisCO folding (Hayer-Hartl, 2017; Liu et al., 2010). Altogether, our findings suggest that

additional physiological constraints may be limiting RuBisCO expression beyond transcription.

The exact mechanism of the regulation we observed in our experiments await further

characterization.

Various culture conditions including pH, temperature, CO_2 concentration, and light intensity may

substantially affect the growth characteristics of cyanobacteria, and thus have the potential to

influence isotopic discrimination (Kuan, Duff, Posarac, & Bi, 2015; Rillema, MacCready, &

Vecchiarelli, 2020; Ungerer, Lin, Chen, & Pakrasi, 2018; Yu et al., 2015). The influence of genetic

factors, including the regulation of RuBisCO expression on cyanobacterial fitness, is less well

known. Previous studies targeting the correlation between RuBisCO overexpression and fitness in

cyanobacteria have yielded mixed results. For instance, faster growth rates and oxygen evolution

rates were observed for an engineered *Synechocystis* PCC 6803 strain that overexpresses RuBisCO

(Liang & Lindblad, 2017), as well as in *S. elongatus* upon co-overexpression of its

phosphoribulokinase (Kanno, Carroll, & Atsumi, 2017). However, in *Synechococcus* sp. PCC

7002, the overexpression of RuBisCO did not alter growth rate (De Porcellinis et al., 2018). The

impact of RuBisCO upregulation on cyanobacterial growth appears to be species/strain-specific.

Our data shows that RuBisCO overexpression in *S. elongatus* PCC 7942 strain Syn02 results in a

minor increase in growth rate in ambient air relative to WT, and no significant difference was

observed at elevated CO_2 levels, nor for oxygen evolution rates. Under ambient environmental

conditions, RuBisCO has a relatively high baseline level of expression in cyanobacteria (V.

Vijayan, I. H. Jain, & E. K. O'Shea, 2011), perhaps explaining why increased expression does not

significantly improve fitness. Our results indicate that further study is needed to fully understand

the coupled interaction of different environmental conditions and genetic backgrounds in

determining cyanobacterial fitness.

Another important aspect to consider is the degree to which genetic regulation and background

may impact system-level isotope fractionation in cyanobacteria. Understanding the connection

between genetics and biogeochemistry, though often overlooked, can augment interpretation of

isotopic biosignatures in deep time. We measured the carbon isotopic composition of

cyanobacterial biomass ($\delta^{13}C_{biomass}$) and photosynthetic carbon isotopic discrimination (ε_p) in WT

and Syn02 cultured under a range of CO_2 concentrations. The increased expression and activity of

RuBisCO in Syn02 apparently neither changed the availability of carbon isotopes in the enzyme's

vicinity nor altered the performance of other cellular modules contributing to carbon isotope

discrimination. $\delta^{13}C_{biomass}$ and ε_p values were comparable for WT and Syn02 grown at all tested

CO_2 concentrations (**Fig. 4**).

Our results show that at both 2% and 5% CO_2, ε_p values were increased by ~7-9‰ for WT and

Syn02 strains relative to air. Theoretically, under high CO_2 concentrations, the intrinsic kinetic

isotope effect of RuBisCO is expressed and maximizes ε_p (Bidigare et al., 1997; Hayes, 1993;

Schubert & Jahren, 2012; Wilkes et al., 2018). Under these conditions, other cellular modules,

such as carbon concentrating mechanisms capable of discriminating carbon isotopes, produce

negligible impact on net carbon discrimination at the organismal level (Hurley et al., 2021; Laws

et al., 2002). The positive relationship between CO_2 concentration and carbon isotope fractionation

has been observed empirically for a variety of autotrophs (Freeman & Hayes, 1992; Hinga et al.,

1994; Schubert & Jahren, 2012; Wilkes et al., 2018), including for cyanobacteria in particular

(Eichner et al., 2015; Hurley et al., 2021). Other studies showed that photoautotrophs that grew

under a lower pH, and thus a higher proportion of CO_2, also exhibited higher carbon fractionation

(Mizutani & Wada, 1982; Roeske & O'Leary, 1984; Wang, Yeager, & Lu, 2016; Yoshioka, 1997).

The ε_p values observed under high inorganic carbon availability fall within the range of 17-20‰

(**Fig. 4B**), which follows the range known for Form I RuBisCO (R. D. Guy et al., 1993; McNevin,

Badger, Kane, & Farquhar, 2006; Scott et al., 2007).

Based on the data generated here, we speculate that, in addition to environmental factors such as

CO_2 concentrations, RuBisCO variation at the gene and enzymatic level, rather than its

overexpression and regulation, may be a more significant determinant of the isotopic composition

of produced biosignatures. Less well-characterized RuBisCO forms (i.e., Forms IC, ID, and II)

that are found in proteobacteria and marine phytoplankton can exhibit significantly smaller

fractionation factors than that observed for forms associated with land plants and cyanobacteria

(Boller, Thomas, Cavanaugh, & Scott, 2011; Robinson et al., 2003; Thomas et al., 2019). The full

diversity of RuBisCO isotopic fractionation requires further study,. Our results underscore that the

integration of synthetic biology, metabolic engineering and geochemistry can offer new insights

into the study and interpretation of biogeochemical reservoirs at the global scale. Further work

needs to be performed to elucidate how the intracellular factors impact metabolisms responsible

for carbon isotope fractionation under ancient transitions in environmental conditions. A deeper

understanding of the impact of the RuBisCO regulation and the environmental factors on the

resulting organismal behavior is essential to establish the extent to which the RuBisCO isotope

fractionation properties can be correlated with evidence for biological activity in the Earth's deep

past.

ACKNOWLEDGEMENTS

This work was supported by the National Science Foundation Emerging Frontiers Program Award

No. 1724090 (BK), NASA Early Career Faculty (ECF) Award No. 80NSSC19K1617 (BK.),

NASA Postdoctoral Fellowship (AKG.), Simons Foundation Early Career Award No. 561645

(JNY and ML), and National Institute of General Medical Sciences of the National Institutes of

Health award number R01GM118815 (AT, to James W. Golden at the University of California-

San Diego). We sincerely thank Bob Blankenship, Zach Adam, Katie McGrath and the members

MATERIALS AND METHODS

Cyanobacterial growth and maintenance

S. elongatus PCC 7942 strains were cultured in BG-11 medium (Rippka, Deruelles, Waterbury,

Herdman, & Stanier, 1979) as liquid cultures or on agar plates (1.5% (w/v) agar and 1 mM

$Na_2S_2O_3 \cdot 5H_2O$). For recombinant strains, liquid and solid media were supplemented with

appropriate antibiotics: 2 $\mu g \cdot ml^{-1}$ Spectinomycin (Sp) plus 2 $\mu g \cdot ml^{-1}$ Streptomycin (Sm), 5

$\mu g \cdot ml^{-1}$ Kanamycin (Km). The cyanobacterial growth was measured at optical density at 750 nm

(OD_{750}).

The strains were archived at -80 °C in 15% (v/v) glycerol. The 2-ml vials were rapidly thawed and

inoculated in liquid BG-11 (supplemented with antibiotics as needed). Cultures were shaken at

120 rpm at 30 °C under continuous low illumination of 45 $\mu mol\ photon \cdot m^{-2} \cdot s^{-1}$, and ambient air,

until they reached an OD_{750} between 0.4 and 0.6. These cultures were then used to inoculate fresh

cultures that were grown using similar conditions but under moderate illumination of 80 μmol

$photon \cdot m^{-2} \cdot s^{-1}$ and sparged at selected CO_2 concentrations (ambient air, 2%, or 5% CO_2; cultures

were also grown without sparging to assess differences in growth rate; **Table S1**). When cultures

reached an OD_{750} of ~7 to 7.5, they were sampled for subsequent experiments as described below

(cultures were also sampled at an earlier OD_{750} of 0.5-1.5 for additional RT-qPCR experiments,

see below; **Fig. S1**).

Genetic engineering of cyanobacteria

A recombinant strains of *S. elongatus* was constructed by natural transformation using standard protocols (Clerico et al., 2007) and the plasmids and methods described below (**Table 1**) . To construct the plasmid pSyn02, pAM4937 was digested with *Swa*I to release the *ccdB* toxic gene and produce a plasmid backbone that contains the pBR322 *E. coli* origin of replication, the base of mobilization site for conjugal transfer, the *aph1* gene conferring kanamycin resistance, and sequences for homologous recombination into *S. elongatus* chromosome at NS2. The *rbc* operon was amplified from *S. elongatus* PCC 7942 gDNA with primers F01 and R01 (**Table S2**) containing 20-nucleotide sequences that overlap with pAM4937 backbone. The resulting DNA fragments were assembled using the GeneArt™ Seamless Cloning and Assembly Kit (Invitrogen, Cat. No. A13288). pSyn02 was used to insert the *rbc* operon into NS2 of the wild type genome in the strain PCC 7942 through homologous recombination to create the strain of *S. elongatus*, Syn02, carrying two copies of the *rbc* operon. Plasmid pSyn-01 was constructed by using two primer pairs F02/R02 and F03/R03 (**Table S2**), 1) to amplify a fragment of pAM4951 that contains the *E. coli* origin of replication and the site for conjugal transfer, and 2) to amplify the *aadA* gene conferring spectinomycin/streptomycin resistance. Native *rbc* operon flanking sequences were amplified from the *S. elongatus* PCC 7942 gDNA with the primer pairs F04/R04 and F05/R05 (**Table S2**) containing 20-nucleotide sequences that overlap with the pAM4951 fragments.

Transformation was carried out after growing the WT strain in liquid culture at 30 °C with shaking (120 rpm) and a light intensity of 80 μmol photon·m^{-2}·s^{-1} until an OD_{750} ~0.7. The cells were prepared for transformation according to the protocol by Clerico et al. (2007) and plated on BG-11 agar containing the appropriate antibiotic(s) for recovery. Subsequently, the colonies were

377   picked using sterile pipette tips, patched onto BG-11 agar containing the appropriate antibiotic(s),

378   and further incubated to ensure complete chromosome segregation (i.e., incorporation of the *trans-*

379   gene into all chromosomes). The patched transformants were screened using colony PCR using

380   the primers F06/R06 and F07/R07 and the genotypes of the engineered strains were confirmed by

381   Sanger sequencing using the primers F08/R08 **(Fig. S2, Table S2**)

Extraction of total RNA and proteins

Cultures were collected during the exponential growth phase. Cells were pelleted by centrifugation

at 4,700 × g for 10 min at room temperature and resuspended in 10 mL of TE buffer (10 mM Tris,

pH 8.0, 1 mM EDTA). To prepare the crude cell lysate, 8 mL of the cell suspension were pelleted

by centrifugation at 4,700 × g for 10 min at room temperature, resuspended in 500 µL of hot (pre-

warmed to 95 °C) TE buffer supplemented with 1% (w/v) SDS and incubated at 95 °C for 10 min.

Then, the mixture was sonicated at 40% amplitude for 3 × 10 sec with 10 sec intervals, and the

cell debris centrifuged at 17,000 × g for 10 min at room temperature. The supernatant was collected

and stored at -80 °C. Total RNA was extracted from the remaining 2 mL of the cell suspension

using the RNeasy® Protect Bacteria Mini Kit (QIAGEN, Cat. No. 74524) following the

manufacturer instructions.

Analysis of *rbc* operon expression by RT-qPCR

Total RNA was quantified with the NanoDrop spectrophotometer and 1 µg of RNA was treated

with the amplification grade deoxyribonuclease I (Invitrogen, Cat. No. 18068-015). 10 µL of

DNase I-treated RNA was then used in reverse transcription (RT) performed with the

SuperScript™ IV First-Strand Synthesis System (Invitrogen, Cat. No. 18091050). The four pairs

of qPCR primers (listed in **Table S2**) were designed with Primer3Plus

(http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi). Both reference genes have

previously been shown to be stably expressed under diverse conditions in *S. elongatus* (Luo et al.,

2019). The quality of cDNA and primer specificity was assessed by PCR using cDNA templates

(RT positive reactions), RT negative controls, and the qPCR primers. The analysis of gene

expression levels was performed in the real-time thermal cycler qTOWER^3 G (Analytik Jena AG),

equipped with the software qPCRsoft, using the cycles: 50 °C/2 min, 95 °C/2 min, 40 × (95 °C/15

sec, 60 °C/1 min). The relative expression of the *rbc* operon genes (*rbcL* and *purK*) was calculated

as the average fold change normalized to reference genes using the delta-delta Ct method. The

experiment was carried out using three biological replicates and three technical replicates.

Analysis of RbcL protein by Western blot

Total protein concentration in the crude cell lysates was measured using the Pierce™ BCA Protein

Assay Kit (Thermo Scientific, Cat. No. 23225). The lysates were loaded in the amount of 5 µg of

total protein in Laemmli sample buffer onto a 6% (v/v) polyacrylamide stacking gel. Proteins were

electrophoresed in a 12% polyacrylamide resolving gel in TGS buffer and blotted in transfer buffer

onto a PVDF membrane. Total protein load in each sample was visualized by Revert™ 700 Total

Protein Stain (LI-COR Biosciences, Cat. No. 926-11011) and used for the RbcL signal

normalization. Detection of RbcL was performed by overnight incubation of the membrane at 4

°C with rabbit anti-RbcL antibody (Agrisera, Cat. No. AS03 037), 1:5000 in TBST with 5% non-

fat milk, followed by one-hour incubation at room temperature with IRDye® 800CW goat anti-

rabbit IgG secondary antibody (LI-COR Biosciences, Cat. No. 926-32211), 1:20,000 in Intercept®

(TBS) blocking buffer (LI-COR Biosciences, Cat. No. 927-60001) with 0.1% (v/v) Tween-20 and

0.01% (w/v) SDS. Both the total protein load and the amount of RbcL in each sample were

documented with Odyssey® Fc Imaging System (LI-COR Biosciences, Cat. No. 2800-03) at the

near-infrared detection mode. The images were acquired using Image Studio™ software. The

densitometric analysis of RbcL signal intensity, normalized to total protein load, was performed

with Quantity One® software (Bio-Rad) for six biological replicates. The amount of RbcL

produced by each replicate of the Syn02 strain culture was compared to the averaged level of RbcL

in the WT PCC 7942 replicate cultures and expressed as the averaged percent of RbcL synthesized

by the WT strain.

Assembly of RuBisCO subunits

Assembly of the RuBisCO large and small subunits into a hexadecameric complex in each strain

was evaluated by native gel electrophoresis and immunodetection of the RuBisCO complexes.

Samples were collected during the exponential growh phase. Cells were pelleted by centrifugation

at 4,700 × g for 10 min at room temperature and resuspended in 400 µL of native lysis buffer (50

mM Tris, pH 8.0, 150 mM NaCl, 1 mM EDTA, 10% (v/v) glycerol) supplemented with 5 mM

DTT, 100 µg/mL lysozyme from chicken egg white, and 1% (v/v) Halt™ Protease Inhibitor

Cocktail (Thermo Scientific, Cat. No. 78430). The cell suspensions were incubated at 30 °C for

15 min and subjected to five consecutive freeze-thaw cycles (10 min at -80 °C followed by 5 min

at 30 °C), then were sonicated on ice for 3 minutes at 30% amplitude (2-sec on/off intervals),

centrifuged at 17,000 × g for 15 min at 4 °C. The concentration of total soluble proteins in the

lysates was determined with Pierce™ BCA Protein Assay Kit. The lysates were adjusted to 5 µg

of total soluble proteins in native sample buffer and then loaded onto a 4-20% Mini-PROTEAN®

TGX™ Precast Protein Gel (Bio-Rad, Cat. No. 4561094). Protein electrophoreses were performed

in TG buffer (60 mM Tris, 192 mM glycine) at 100 V for 4 h at 4 °C and blotted in transfer buffer

(48 mM Tris, pH 9.2, 39 mM glycine, 0.04% (w/v) SDS) onto a nitrocellulose membrane. After

three 10-min washes in wash buffer (48 mM Tris, pH 9.2, 39 mM glycine, 20% (v/v) methanol),

total protein load in each sample was visualized by Revert™ 700 Total Protein Stain and used for

the normalization of RuBisCO complex quantity. Immunodetection of the RuBisCO complex was

performed with the same primary and secondary antibodies that were used to analyze the level of

RbcL, as described above.

Catalytic activity of RuBisCO

The activity of RuBisCO in cyanobacterial lysates was measured using a spectrophotometric

coupled-enzyme assay that links this activity with the rate of NADH oxidation (Kubien et al.,

2011). The cyanobacterial strains were cultured, collected, and pelleted as described above. The

pellets were resuspended in 1 mL of ice-cold lysis buffer (50 mM EPPS, 1 mM EDTA, 2 mM

DTT, pH 8.0) and transferred into 2 mL screw-capped tubes with Lysing Matrix B (MP

Biomedical) for lysis by bead beating using FastPrep-24™ 5G bead beater (MP Biomedical) with

4 m/sec for 10 sec, followed by 2-min incubation on ice, repeated six times. The cell lysates were

transferred to new Eppendorf tubes to remove the beads and unbroken cells and to pellet the

thylakoid membrane by centrifugation at 10,000 × g for 1 min and at 20,000 × g for 30 min at 4

°C, sequentially. The resulting clear supernatants containing cytosolic soluble proteins, including

phycobiliproteins and RuBisCO, were used to determine protein concentration by Pierce™ BCA

Protein Assay Kit and to measure RuBisCO activity by employing an assay adapted from Kubien

et al. (2011). The assay buffer (100 mM HEPES, 25 mM $MgCl_2$, 1 mM EDTA, pH 7.6) was used

considering the high Michaelis constant for CO_2 (K_C) for cyanobacterial RuBisCO. 20 μL of cell

lysates were preincubated in the assay mix (with 5 mM $NaHCO_3$) at 25 °C for activation before

initiating the reaction by adding synthesized ribulose 1,5-bisphosphate (RuBP) according to Kane

et al. (Kane, Wilkin, Portis, & John Andrews, 1998). The absorbance at 340 nm was monitored

using the Synergy H1 plate reader (BioTek). RuBisCO activity was reported as RuBP consumption

rate normalized to total soluble protein content. The assay was performed for three biological

replicates.

Cyanobacterial growth measurements

OD_{750} values were plotted as a function of time and analyzed in R with the Growthcurver package.

Growth curve data was fitted to the standard form of the logistic equation to calculate growth

parameters including growth rate, doubling time, and carrying capacity (Sprouffske & Wagner,

2016). Each strain was grown in triplicate for every condition.

Photosynthetic oxygen evolution rate

S. elongatus strain photosynthetic activity was assayed using a Clark-type oxygen electrode

chamber to measure the level of molecular oxygen produced in cyanobacterial cultures. Cells were

grown in 50 mL of BG-11 at 30 °C, illumination of 80 μmol photon·m^{-2}·s^{-1}, shaking at 120 rpm,

in ambient air, and with culture sparging. The samples were collected from triplicate cultures

during the exponential growth phase, pelleted by centrifugation at 4,700 × g for 10 min at room

temperature, and resuspended in fresh BG-11 to an OD_{750} of ~1.0. Concentration of chlorophyll *a*

(for normalization) was measured following the protocol by Zavrel et al. (Zavrel et al., 2015). The

remaining suspension was incubated in the dark for 20 min with gentle agitation. Samples from

each suspension, prepared in three technical replicates, were analyzed in an oxygen electrode

chamber under saturated light, using the Oxygraph+ System (Hansatech Instruments) equipped

with the OxyTrace+ software. Oxygen evolution rate was monitored for 10 min and expressed as

nanomoles of molecular oxygen evolved per hour per microgram of chlorophyll *a*.

Carbon isotope fractionation in bulk cyanobacterial biomass

The bacteria were pelleted by centrifugation at 4,700 × g for 10 min at room temperature, washed

in 10 mL of 10 mM NaCl. The bacteria were resuspended in 1 mL of 10 mM NaCl and transferred

to Eppendorf tubes. After centrifugation at 4,700 × g for 10 min at room temperature, the

supernatants were completely removed, the pellets were dried in opened tubes in a laboratory oven

at 50 °C for 2 days, and the resultant dried biomass samples were transferred into tin capsules.

The carbon isotope composition of bulk biomass ($\delta^{13}C_{biomass}$) samples was determined at the UC

Davis Stable Isotope Facility. $\delta^{13}C_{biomass}$ was analyzed using a PDZ Europa ANCA-GSL elemental

analyzer interfaced to a PDZ Europa 20-20 isotope ratio mass spectrometer (Sercon Ltd.). The

carbon isotopic composition values were reported relative to the Vienna PeeDee Belemnite

standard (V-PDB):

$$\delta^{13}C_{sample} = \left(\frac{^{13}C/^{12}C_{sample}}{^{13}C/^{12}C_{V-PDB}} - 1 \right) \times 1000$$

509 The carbon isotope fractionation associated with photosynthetic CO_2 fixation (ε_p) was calculated

510 relative to a reference $\delta^{13}C_{CO2}$ value (-8.4‰; (Keeling et al., 2017)) according to Freeman and

511 Hayes (1992):

$$\varepsilon_p = \frac{\delta^{13}C_{CO_2} - \delta^{13}C_{biomass}}{1 + \delta^{13}C_{biomass}/1000}$$

Statistical analyses

Results for experimental analyses were presented as the mean and the sample standard deviation (SD) values of at least three independent experiments. Statistical significance was analyzed swith the two-tailed t-test. The unpaired sample t-test assuming equal variances was used to compare the values obtained for different cyanobacterial strains and the paired sample t-test was used to compare the values for the same strain at different experimental conditions.

REFERENCES

Andersson, C. R., Tsinoremas, N. F., Shelton, J., Lebedeva, N. V., Yarrow, J., Min, H., &
Golden, S. S. (2000). Application of bioluminescence to the study of circadian rhythms in
cyanobacteria. *Methods Enzymol, 305*, 527-542.

Atsumi, S., Higashide, W., & Liao, J. C. (2009). Direct photosynthetic recycling of carbon
dioxide to isobutyraldehyde. *Nat Biotechnol, 27*(12), 1177-1180. doi:10.1038/nbt.1586

Bar-On, Y. M., & Milo, R. (2019). The global mass and average rate of rubisco. *Proc Natl Acad
Sci USA, 10*, 4738-4743. doi:10.1073/pnas.1816654116

Bidigare, R. R., Fluegge, A., Freeman, K. H., Hanson, K. L., Hayes, J. M., Hollander, D., . . .
Wakeham, S. G. (1997). Consistent fractionation of ^{13}C in nature and in the laboratory:
growth-rate effects in some haptophyte algae. *Global Biogeochem Cycles, 11*(2), 279-
292. doi:10.1029/96gb03939

Boller, A. J., Thomas, P. J., Cavanaugh, C. M., & Scott, K. M. (2011). Low stable carbon isotope
fractionation by coccolithophore RubisCO. *Geochimica et Cosmochimica Acta, 75*(22),
7200-7207. doi:10.1016/j.gca.2011.08.031

Bonfil, D. J., Ronen-Tarazi, M., Sultemeyer, D., Lieman-Hurwitz, J., Schatz, D., & Kaplan, A.
(1998). A putative HCO3- transporter in the cyanobacterium Synechococcus sp. strain
PCC 7942. *FEBS Lett, 430*(3), 236-240.

Catling, D. C., & Zahnle, K. J. (2020). The Archean atmosphere. *Science Advances, 6*(9).
doi:10.1126/sciadv.aax1420

Clerico, E. M., Ditty, J. L., & Golden, S. S. (2007). Specialized techniques for site-directed
mutagenesis in cyanobacteria. *Methods Mol Biol, 362*, 155-171. doi:10.1007/978-1-
59745-257-1_11

De Porcellinis, A. J., Norgaard, H., Brey, L. M. F., Erstad, S. M., Jones, P. R., Heazlewood, J. L.,

& Sakuragi, Y. (2018). Overexpression of bifunctional fructose-1,6-

bisphosphatase/sedoheptulose-1,7-bisphosphatase leads to enhanced photosynthesis and

global reprogramming of carbon metabolism in Synechococcus sp. PCC 7002. *Metab*

Eng, 47, 170-183. doi:10.1016/j.ymben.2018.03.001

Deleens, E., Treichel, I., & O'Leary, M. H. (1985). Temperature dependence of carbon isotope

fractionation in CAM plants. *Plant Physiology, 79*(1), 202-206. doi:10.1104/pp.79.1.202

Des Marais, D. J. (2001). Isotopic evolution of the biogeochemical carbon cycle during the

Precambrian. *Reviews in Mineralogy and Geochemistry, 43*(1), 555-578.

doi:10.2138/gsrmg.43.1.555

Eichner, M., Thoms, S., Kranz, S. A., & Rost, B. (2015). Cellular inorganic carbon fluxes in

Trichodesmium: a combined approach using measurements and modelling. *J Exp Bot,*

66(3), 749-759. doi:10.1093/jxb/eru427

Ellis, R. J. (1979). The most abundant protein in the world. *Trends in Biochemical Sciences,*

4(11), 241-244.

Erb, T. J., & Zarzycki, J. (2018). A short history of RubisCO: the rise and fall (?) of Nature's

predominant CO_2 fixing enzyme. *Curr Opin Biotechnol, 49*, 100-107.

doi:10.1016/j.copbio.2017.07.017

Eungrasamee, K., Miao, R., Incharoensakdi, A., Lindblad, P., & Jantaro, S. (2019). Improved

lipid production via fatty acid biosynthesis and free fatty acid recycling in engineered

Synechocystis sp. PCC 6803. *Biotechnology for biofuels, 12*, 8-8. doi:10.1186/s13068-

018-1349-8

Falkowski, P. G., Fenchel, T., & Delong, E. F. (2008). The microbial engines that drive Earth's

biogeochemical cycles. *Science, 320*(5879), 1034-1039. doi:10.1126/science.1153213

Field, C. B. (1998). Primary Production of the Biosphere: Integrating Terrestrial and Oceanic

Components. *Science, 281*(5374), 237-240. doi:10.1126/science.281.5374.237

Freeman, K. H., & Hayes, J. M. (1992). Fractionation of carbon isotopes by phytoplankton and

estimates of ancient CO_2 levels. *Global Biogeochemical Cycles, 6*(2), 185-198.

doi:10.1029/92gb00190

Gabay, C., Lieman-Hurwitz, J., Hassidim, M., Ronen-Tarazi, M., & Kaplan, A. (1998).

Modification of topA in Synechococcus sp. PCC 7942 resulted in mutants capable of

growing under low but not high concentration of CO2. *FEMS Microbiology Letters,*

159(2), 343-347. doi:10.1111/j.1574-6968.1998.tb12881.x

Garcia, A. K., Cavanaugh, C. M., & Kacar, B. (2021). The curious consistency of carbon

biosignatures over billions of years of Earth-life coevolution. *ISME J.*

doi:10.1038/s41396-021-00971-5

Gesch, R. W., Kang, I. H., Gallo-Meagher, M., Vu, J. C. V., Boote, K. J., H. Allen, L., & Bowes,

G. (2003). Rubisco expression in rice leaves is related to genotypic variation of

photosynthesis under elevated growth CO2 and temperature. *Plant, Cell & Environment,*

26(12), 1941-1950. doi:10.1046/j.1365-3040.2003.01110.x

Guy, H. I., & Evans, D. R. (1996). Function of the major synthetase subdomains of carbamyl-

phosphate synthetase. *J Biol Chem, 271*(23), 13762-13769. doi:10.1074/jbc.271.23.13762

Guy, R. D., Fogel, M. L., & Berry, J. A. (1993). Photosynthetic fractionation of the stable

isotopes of oxygen and carbon. *Plant Physiol, 101*(1), 37-47. doi:10.1104/pp.101.1.37

Hamilton, T. L., Bryant, D. A., & Macalady, J. L. (2016). The role of biology in planetary

evolution: cyanobacterial primary production in low-oxygen Proterozoic oceans.

Environmental Microbiology, 18(2), 325-340. doi:10.1111/1462-2920.13118

Havig, J. R., Hamilton, T. L., Bachan, A., & Kump, L. R. (2017). Sulfur and carbon isotopic

evidence for metabolic pathway evolution and a four-stepped Earth system progression

across the Archean and Paleoproterozoic. *Earth-Science Reviews, 174*, 1-21.

doi:10.1016/j.earscirev.2017.06.014

Hayer-Hartl, M. (2017). From chaperonins to Rubisco assembly and metabolic repair. *Protein*

Science, 26(12), 2324-2333. doi:10.1002/pro.3309

Hayes, J. M. (1993). Factors controlling 13C contents of sedimentary organic compounds:

Principles and evidence. *Marine Geology, 113*(1-2), 111-125. doi:10.1016/0025-

3227(93)90153-m

Hinga, K. R., Arthur, M. A., Pilson, M. E. Q., & Whitaker, D. (1994). Carbon isotope

fractionation by marine phytoplankton in culture: the effects of CO_2 concentration, pH,

temperature, and species. *Global Biogeochemical Cycles, 8*(1), 91-102.

doi:10.1029/93gb03393

Hood, R. D., Higgins, S. A., Flamholz, A., Nichols, R. J., & Savage, D. F. (2016). The stringent

response regulates adaptation to darkness in the cyanobacterium Synechococcus

elongatus. *Proc Natl Acad Sci U S A, 113*(33), E4867-4876.

doi:10.1073/pnas.1524915113

Hurley, S. J., Wing, B. A., Jasper, C. E., Hill, N. C., & Cameron, J. C. (2021). Carbon isotope

evidence for the global physiology of Proterozoic cyanobacteria. *Science Advances, 7*(2).

doi:10.1126/sciadv.abc8998

Iwaki, T., Haranoh, K., Inoue, N., Kojima, K., Satoh, R., Nishino, T., . . . Wadano, A. (2006).

Expression of foreign type I ribulose-1,5-bisphosphate carboxylase/ oxygenase (EC

4.1.1.39) stimulates photosynthesis in cyanobacterium Synechococcus PCC7942 cells.

Photosynth Res, 88(3), 287-297. doi:10.1007/s11120-006-9048-x

Kacar, B., Hanson-Smith, V., Adam, Z. R., & Boekelheide, N. (2017). Constraining the timing

of the Great Oxidation Event within the Rubisco phylogenetic tree. *Geobiology, 15*(5),

628-640. doi:10.1111/gbi.12243

Kane, H. J., Wilkin, J.-M., Portis, A. R., & John Andrews, T. (1998). Potent Inhibition of

Ribulose-Bisphosphate Carboxylase by an Oxidized Impurity in Ribulose-1,5-

Bisphosphate. *Plant Physiology, 117*(3), 1059-1069. doi:10.1104/pp.117.3.1059

Kanno, M., Carroll, A. L., & Atsumi, S. (2017). Global metabolic rewiring for improved CO2

fixation and chemical production in cyanobacteria. *Nature Communications, 8*(1), 14724.

doi:10.1038/ncomms14724

Keeling, R. F., Graven, H. D., Welp, L. R., Resplandy, L., Bi, J., Piper, S. C., . . . Meijer, H. A. J.

(2017). Atmospheric evidence for a global secular increase in carbon isotopic

discrimination of land photosynthesis. *Proc Natl Acad Sci U S A, 114*(39), 10361-10366.

doi:10.1073/pnas.1619240114

Knoll, A.H., Bergmann, K.D. and Strauss, J.V., 2016. Life: the first two billion

years. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *371*(1707),

p.20150493.

Krissansen-Totton, J., Buick, R., & Catling, D. C. (2015). A statistical analysis of the carbon

isotope record from the Archean to Phanerozoic and implications for the rise of oxygen.

American Journal of Science, 315(4), 275-316. doi:10.2475/04.2015.01

Kuan, D., Duff, S., Posarac, D., & Bi, X. (2015). Growth optimization ofSynechococcus

636   elongatusPCC7942 in lab flasks and a 2-D photobioreactor. *The Canadian Journal of*

637   *Chemical Engineering, 93*(4), 640-647. doi:10.1002/cjce.22154

638   Kubien, D. S., Brown, C. M., & Kane, H. J. (2011). Quantifying the amount and activity of

639   Rubisco in leaves. *Methods Mol Biol, 684*, 349-362. doi:10.1007/978-1-60761-925-3_27

640   Laws, E. A., Bidigare, R. R., & Popp, B. N. (1997). Effect of growth rate and CO_2 concentration

641   on carbon isotopic fractionation by the marine diatom *Phaeodactylum tricornutum*.

642   *Limnology and Oceanography, 42*(7), 1552-1560. doi:10.4319/lo.1997.42.7.1552

643   Laws, E. A., Popp, B. N., Cassar, N., & Tanimoto, J. (2002). 13C discrimination patterns in

644   oceanic phytoplankton: likely influence of CO2 concentrating mechanisms, and

645   implications for palaeoreconstructions. *Functional Plant Biology, 29*(3).

646   doi:10.1071/pp01183

647   Lechno-Yossef, S., Rohnke, B. A., Belza, A. C. O., Melnicki, M. R., Montgomery, B. L., &

648   Kerfeld, C. A. (2020). Cyanobacterial carboxysomes contain an unique rubisco-activase-

649   like protein. *New Phytologist, 225*(2), 793-806. doi:10.1111/nph.16195

650   Liang, F., & Lindblad, P. (2017). Synechocystis PCC 6803 overexpressing RuBisCO grow faster

651   with increased photosynthesis. *Metab Eng Commun, 4*, 29-36.

652   doi:10.1016/j.meteno.2017.02.002

653   Liu, C., Young, A. L., Starling-Windhof, A., Bracher, A., Saschenbrecker, S., Rao, B. V., . . .

654   Hayer-Hartl, M. (2010). Coupled chaperone action in folding and assembly of

655   hexadecameric Rubisco. *Nature, 463*(7278), 197-202. doi:10.1038/nature08651

Lloyd, M. K., McClelland, H. L. O., Antler, G., Bradley, A. S., Halevy, I., Junium, C. K., . . .

Zerkle, A. L. (2020). The Isotopic Imprint of Life on an Evolving Planet. *Space Science*

Reviews, 216(7). doi:10.1007/s11214-020-00730-6

Luo, X., Li, J., Chang, T., He, H., Zhao, Y., Yang, X., . . . Xu, Y. (2019). Stable Reference Gene

Selection for RT-qPCR Analysis in Synechococcus elongatus PCC 7942 under Abiotic

Stresses. *Biomed Res Int, 2019*, 7630601. doi:10.1155/2019/7630601

Lyons, T. W., Fike, D. A., & Zerkle, A. (2015). Emerging Biogeochemical Views of Earth's

Ancient Microbial Worlds. *Elements, 11*(6), 415-421. doi:10.2113/gselements.11.6.415

Maeda, S., Price, G. D., Badger, M. R., Enomoto, C., & Omata, T. (2000). Bicarbonate binding

activity of the CmpA protein of the cyanobacterium Synechococcus sp. strain PCC 7942

involved in active transport of bicarbonate. *J Biol Chem, 275*(27), 20551-20555.

doi:10.1074/jbc.M003034200

McNevin, D. B., Badger, M. R., Kane, H. J., & Farquhar, G. D. (2006). Measurement of (carbon)

kinetic isotope effect by Rayleigh fractionation using membrane inlet mass spectrometry

for CO2-consuming reactions. *Functional plant biology : FPB, 33*(12), 1115-1128.

doi:10.1071/fp06201

Mizutani, H., & Wada, E. (1982). Effect of high atmospheric CO2 concentration on δ13 of algae.

Origins of life, 12(4), 377-390. doi:10.1007/BF00927070

Nisbet, E. G., Grassineau, N. V., Howe, C. J., Abell, P. I., Regelous, M., & Nisbet, R. E. R.

(2007). The age of Rubisco: the evolution of oxygenic photosynthesis. *Geobiology, 5*,

311-335. doi:10.1111/j.1472-4669.2007.00127.x

Omata, T., Gohta, S., Takahashi, Y., Harano, Y., & Maeda, S. (2001). Involvement of a CbbR

homolog in low CO2-induced activation of the bicarbonate transporter operon in

cyanobacteria. *J Bacteriol, 183*(6), 1891-1898. doi:10.1128/JB.183.6.1891-1898.2001

Onizuka, T., Akiyama, H., Endo, S., Kanai, S., Hirano, M., Tanaka, S., & Miyasaka, H. (2002).

CO2 Response Element and Corresponding trans-acting Factor of the Promoter for

Ribulose-1,5-bisphosphate Carboxylase/oxygenase Genes in Synechococcus sp.

PCC7002 Found by an Improved Electrophoretic Mobility Shift Assay. *Plant and Cell*

Physiology, 43(6), 660-667. doi:10.1093/pcp/pcf082

Poudel, S., Pike, D. H., Raanan, H., Mancini, J. A., Nanda, V., Rickaby, R. E. M., & Falkowski,

P. G. (2020). Biophysical analysis of the structural evolution of substrate specificity in

RuBisCO. *Proceedings of the National Academy of Sciences, 117*(48), 30451-30457.

doi:10.1073/pnas.2018939117

Price, G. D., Badger, M. R., Woodger, F. J., & Long, B. M. (2008). Advances in understanding

the cyanobacterial CO2-concentrating-mechanism (CCM): functional components, Ci

transporters, diversity, genetic regulation and prospects for engineering into plants. *J Exp*

Bot, 59(7), 1441-1461. doi:10.1093/jxb/erm112

Rau, G. H., Riebesell, U., & Wolf-Gladrow, D. (1996). A model of photosynthetic 13 C

fractionation by marine phytoplankton based on diffusive molecular CO2 uptake. *Marine*

Ecology Progress Series, 133(1/3), 275-285.

Raven, J. A. (2013). Rubisco: still the most abundant protein of Earth? *New Phytologist, 198*(1),

1-3. doi:10.1111/nph.12197

Raven, J. A., Cockell, C. S., & De La Rocha, C. L. (2008). The evolution of inorganic carbon

concentrating mechanisms in photosynthesis. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, 363(1504), 2641-2650. doi:doi:10.1098/rstb.2008.0020

Riebesell, U., Revill, A. T., Holdsworth, D. G., & Volkman, J. K. (2000). The effects of varying

CO2 concentration on lipid composition and carbon isotope fractionation in Emiliania

huxleyi. *Geochimica et Cosmochimica Acta, 64*(24), 4179-4192. doi:10.1016/s0016-

7037(00)00474-9

Rillema, R., MacCready, J. S., & Vecchiarelli, A. G. (2020). Cyanobacterial growth and

morphology are influenced by carboxysome positioning and temperature.

doi:10.1101/2020.06.01.127845

Rippka, R., Deruelles, J., Waterbury, J. B., Herdman, M., & Stanier, R. Y. (1979). Generic

Assignments, Strain Histories and Properties of Pure Cultures of Cyanobacteria.

Microbiology, 111(1), 1-61. doi:https://doi.org/10.1099/00221287-111-1-1

Robinson, J. J., Scott, K. M., Swanson, S. T., O'Leary, M. H., Horken, K., Tabita, F. R., &

Cavanaugh, C. M. (2003). Kinetic isotope effect and characterization of form II RubisCO

from the chemoautotrophic endosymbionts of the hydrothermal vent tubeworm *Riftia*

pachyptila. *Limnology and Oceanography, 1*, 48-54. doi:10.4319/lo.2003.48.1.0048

Roeske, C. A., & O'Leary, M. H. (1984). Carbon isotope effects on the enzyme-catalysed

carboxylation of ribulose bisphosphate. *Biochemistry, 23*, 6275-6284.

Rothschild, L. J., & DesMarais, D. (1989). Stable carbon isotope fractionation in the search for

life on early Mars. *Adv Space Res, 9*(6), 159-165.

Schidlowski, M. (1988). A 3,800-Million-Year Isotopic Record of Life from Carbon in

Sedimentary-Rocks. *Nature, 333*(6171), 313-318. doi:DOI 10.1038/333313a0

Schidlowski, M. (2001). Carbon isotopes as biogeochemical recorders of life over 3.8 Ga of

722   Earth history: evolution of a concept. *Precambrian Research, 106*(1-2), 117-134. doi:Doi

723   10.1016/S0301-9268(00)00128-5

724   Schirrmeister, B. E., Sanchez-Baracaldo, P., & Wacey, D. (2016). Cyanobacterial evolution

725   during the Precambrian. *International Journal of Astrobiology, 15*(3), 187-204.

726   doi:10.1017/s1473550415000579

727   Schopf, J. W. (2011). The paleobiological record of photosynthesis. *Photosynth Res, 107*(1), 87-

728   101. doi:10.1007/s11120-010-9577-1

729   Schubert, B. A., & Jahren, A. H. (2012). The effect of atmospheric CO_2 concentration on carbon

730   isotope fractionation in C3 land plants. *Geochimica et Cosmochimica Acta, 96*, 29-43.

731   doi:10.1016/j.gca.2012.08.003

732   Scott, K. M., Henn-Sax, M., Harmer, T. L., Longo, D. L., Frame, C. H., & Cavanaugh, C. M.

733   (2007). Kinetic isotope effect and biochemical characterization of form IA RubisCO from

734   the marine cyanobacterium *Prochlorococcus marinus* MIT9313. *Limnology and*

735   *Oceanography, 52*(5), 2199-2204.

736   Sengupta, A., Sunder, A. V., Sohoni, S. V., & Wangikar, P. P. (2019). Fine-Tuning Native

737   Promoters of Synechococcus elongatus PCC 7942 To Develop a Synthetic Toolbox for

738   Heterologous Protein Expression. *ACS Synth Biol, 8*(5), 1219-1223.

739   doi:10.1021/acssynbio.9b00066

740   Sprouffske, K., & Wagner, A. (2016). Growthcurver: an R package for obtaining interpretable

741   metrics from microbial growth curves. *BMC Bioinformatics, 17*, 172.

742   doi:10.1186/s12859-016-1016-7

Szekeres, E., Sicora, C., Dragoş, N., & Drugă, B. (2014). Selection of proper reference genes for

the cyanobacterium Synechococcus PCC 7002 using real-time quantitative PCR. *FEMS*

Microbiol Lett, 359(1), 102-109. doi:10.1111/1574-6968.12574

Taton, A., Erikson, C., Yang, Y., Rubin, B. E., Rifkin, S. A., Golden, J. W., & Golden, S. S.

(2020). The circadian clock and darkness control natural competence in cyanobacteria.

Nat Commun, 11(1), 1688. doi:10.1038/s41467-020-15384-9

Taton, A., Unglaub, F., Wright, N. E., Zeng, W. Y., Paz-Yepes, J., Brahamsha, B., . . . Golden, J.

W. (2014). Broad-host-range vector system for synthetic biology and biotechnology in

cyanobacteria. *Nucleic Acids Research, 42*(17), e136-e136. doi:10.1093/nar/gku673

Tcherkez, G. G., Farquhar, G. D., & Andrews, T. J. (2006). Despite slow catalysis and confused

substrate specificity, all ribulose bisphosphate carboxylases may be nearly perfectly

optimized. *Proc Natl Acad Sci U S A, 103*(19), 7246-7251. doi:10.1073/pnas.0600605103

Tchernov, D., Helman, Y., Keren, N., Luz, B., Ohad, I., Reinhold, L., . . . Kaplan, A. (2001).

Passive entry of CO2 and its energy-dependent intracellular conversion to HCO3- in

cyanobacteria are driven by a photosystem I-generated deltamuH+. *J Biol Chem, 276*(26),

23450-23455. doi:10.1074/jbc.M101973200

Thomas, P. J., Boller, A. J., Satagopan, S., Tabita, F. R., Cavanaugh, C. M., & Scott, K. M.

(2019). Isotope discrimination by form IC RubisCO from *Ralstonia eutropha* and

Rhodobacter sphaeroides, metabolically versatile members of '*Proteobacteria*' from

aquatic and soil habitats. *Environ Microbiol, 21*(1), 72-80. doi:10.1111/1462-2920.14423

Ungerer, J., Lin, P.-C., Chen, H.-Y., & Pakrasi, H. B. (2018). Adjustments to Photosystem

Stoichiometry and Electron Transfer Proteins Are Key to the Remarkably Fast Growth of

the Cyanobacterium Synechococcus elongatus UTEX 2973. *mBio, 9*(1),

e02327-02317. doi:10.1128/mBio.02327-17

Vijayan, V., Jain, I. H., & O'Shea, E. K. (2011). A high resolution map of a cyanobacterial

transcriptome. *Genome Biology, 12*(5), R47. doi:10.1186/gb-2011-12-5-r47

Vijayan, V., Jain, I. H., & O'Shea, E. K. (2011). A high resolution map of a cyanobacterial

transcriptome. *Genome Biol, 12*(5), R47. doi:10.1186/gb-2011-12-5-r47

von Caemmerer, S., Tazoe, Y., Evans, J. R., & Whitney, S. M. (2014). Exploiting

transplastomically modified Rubisco to rapidly measure natural diversity in its carbon

isotope discrimination using tuneable diode laser spectroscopy. *J Exp Bot, 65*(13), 3759-

3767. doi:10.1093/jxb/eru036

Wang, S., Yeager, K. M., & Lu, W. (2016). Carbon isotope fractionation in phytoplankton as a

potential proxy for pH rather than for [CO2(aq)]: Observations from a carbonate lake.

Limnology and Oceanography, 61(4), 1259-1270. doi:10.1002/lno.10289

Wilkes, E. B., Lee, R. B. Y., McClelland, H. L. O., Rickaby, R. E. M., & Pearson, A. (2018).

Carbon isotope ratios of coccolith–associated polysaccharides of *Emiliania huxleyi* as a

function of growth rate and CO_2 concentration. *Organic Geochemistry, 119*, 1-10.

doi:10.1016/j.orggeochem.2018.02.006

Wong, W. W., & Sackett, W. M. (1978). Fractionation of stable carbon isotopes by marine

phytoplankton. *Geochimica et Cosmochimica Acta, 42*(12), 1809-1815.

doi:10.1016/0016-7037(78)90236-3

Yoshioka, T. (1997). Phytoplanktonic carbon isotope fractionation: equations accounting for

CO2-concentrating mechanisms. *Journal of Plankton Research, 19*(10), 1455-1476.

doi:10.1093/plankt/19.10.1455

Yu, J., Liberton, M., Cliften, P. F., Head, R. D., Jacobs, J. M., Smith, R. D., . . . Pakrasi, H. B.

(2015). Synechococcus elongatus UTEX 2973, a fast growing cyanobacterial chassis for

biosynthesis using light and CO2. *Scientific Reports, 5*(1), 8132. doi:10.1038/srep08132

Zavrel, T., Sinetova, M. A., & Červený, J. (2015). Measurement of Chlorophyll a and

Carotenoids Concentration in Cyanobacteria. *Bio-protocol, 5*(9), e1467.

doi:10.21769/BioProtoc.1467

 TABLES

 Table 1. Strains and plasmids used in this study.

Strain or plasmid	Description/Genotype	Antibiotic resistance	Source/Reference
WT	Wild-type strain of *S. elongatus* PCC 7942	-	Susan S. Golden (UC San Diego)
pAM4937	Expression vector for *S. elongatus* PCC 7942 neutral site 2 (NS2)	Km	(Taton et al., 2014)
pSyn02	pAM4937 carrying the *rbc* operon including *rbcL*, *rbcS*, *purK*, and flanking sequences from *S. elongatus* PCC 7942 (CP000100: 1479071-1484283)	Km	This study
Syn02	*S. elongatus* PCC 7942 carrying a second copy of the *rbc* operon and flanking sequences at NS2: NS2::*aphI-rbcL-rbcS-purK*- Synpcc7942_1429-Synpcc7942_1430	Km	This study
pAM4951	Expression vector for *S. elongatus* PCC 7942 neutral site 1 (NS1)	Sp+Sm	(Taton et al., 2014)
pSyn01	Plasmid to replace *S. elongatus'* native *rbc* operon (CP000100: 1479070-	Sp+Sm	This study

	1482595) with a Sp/Sm resistance gene: Δ(*rbcL-rbcS-purK*)::*aadA*.		
Syn01	*S. elongatus* strain Syn02 with the native *rbc* operon removed: Syn02 and Δ(*rbcL-rbcS-purK*)::*aadA*.	Km, Sp+Sm	This study

800 **Table 2. Growth parameters and oxygen evolution rates of *S. elongatus* strains under varying**

801 **CO_2 concentrations[a].**

Strain	Atmosphere	Growth rate (h^{-1})	Carrying capacity (OD_{750})	Oxygen evolution rate nmol ($O_2 \cdot h^{-1} \cdot \mu g^{-1}$ chlorophyll *a*)
WT	Air	0.017 ± 0.001	6.79 ± 0.70	342 ± 26
	2% CO_2	0.043 ± 0.002	8.43 ± 0.53	-
	5% CO_2	0.047 ± 0.005	5.35 ± 0.28	-
Syn02	Air	0.020 ± 0.001*	7.63 ± 0.54	362 ± 26
	2% CO_2	0.046 ± 0.001	8.28 ± 0.22	-
	5% CO_2	0.040 ± 0.001	5.44 ± 0.36	-

[a]Values are means of three biological replicates $\pm$ 1 SD. Asterisks indicate *t*-test result from

comparison with WT for the same atmospheric condition, * – $p < 0.05$.

FIGURES

Figure 1. Experimental strategy. A – Overall methodology. 1) Genetic engineering: An

additional copy of the *rbc* operon was inserted at the chromosomal NS2 site of *S. elongatu*s PCC

7942 to create the Syn02 strain. **2)** RuBisCO characterization was carried out by investigating (a)

catalytic activity, (b) assembly into a hexadecameric complex, and (c) expression at both the

transcript and protein levels. **3)** Physiology: *S. elongatu*s WT and Syn02 strains were evaluated by

monitoring microbial growth characteristics, morphology, and rate of photosynthetic oxygen

evolution. **4)** Carbon isotope fractionation: The impact of RuBisCO overexpression on biomass

$\delta^{13}C$ and ε_p under varying CO_2 concentrations was assessed. **B – Map of the genetic insert in *S.***

elongatus **strain Syn02**. Plasmid pSyn02 was used to insert the *rbc* operon and the *aph1* gene

conferring kanamycin resistance in PCC 7942 WT NS2 to generate the Syn02 strain. Crosslines

indicate homologous recombination sites and scale bars show DNA fragment sizes (in bp).

Figure 2. RuBisCO expression, assembly, and activity in *S*. elongatus strains under varying

CO₂ concentrations. A – Expression of Syn02 *rbcL* relative to WT *rbcL*. Asterisks indicate *t*-

test results compared to WT *rbcL* expression at the same growth condition. **B – Expression of**

WT *rbcL* at 2-5% CO₂ relative to ambient air. Asterisks indicate *t*-test results compared to

WT *rbcL* expression in air. **A-B –** Expression data was measured by RT-qPCR, normalized to

reference genes *secA* and *ppc*. Columns represent mean fold expression for three biological

replicates. **C – Immunodetection of RbcL.** (*left*) Western blot showing RbcL protein detected

by anti-RuBisCO antibody and total protein stain from crude cell lysates. (*right*) RbcL

percentage signal intensities were normalized to that for the total soluble protein load. Columns

represent mean RbcL intensities for six biological replicates, relative to mean values for WT

cultures at the same growth condition. Asterisks indicate *t*-test results compared to WT RbcL. **D**

– Immunodetection of assembled RuBisCO. Western blot showing the proper assembly of

RbcL and RbcS into the L_8S_8 hexadecameric complex L_8S_8 (520 kDa), detected by anti-RbcL

antibody. Rbc L_8S_8 percentage signal intensity was normalized to that for the total soluble

protein load, and is shown relative to WT. **E – Lysate RuBisCO activity.** Activity was

measured by the RuBP consumption rate, normalized to total soluble protein content of cell

lysate. Columns represent the mean activity of three biological replicates. **A-C, E –** Asterisks

indicate *t*-test results for pairwise comparison indicated by horizontal line. Error bars on all

graphs indicate 1 SD. ns – not significant; ** – *p* <0.01; *** – *p* <0.001.

Figure 3. Growth of *S. elongatus* strains. A – Growth profiles. Cultures of each strain were maintained in ambient air or under 2% or 5% CO_2. Smoothed profiles were generated from mean optical density (measured at 750 nm, OD_{750}) values for three replicate cultures per growth condition. **B – Doubling times.** Columns represent the mean doubling time of three replicates. Error bars on all graphs indicate 1 SD, and asterisks indicate *t*-test results for pairwise comparisons indicated by horizontal lines. ns – not significant; * – $p < 0.05$; *** – $p < 0.001$.

Figure 4. $^{13}C/^{12}C$ discrimination of *S. elongatus* strains under varying CO₂ concentrations. A

– Biomass δ^{13}C. Reported δ^{13}C values are relative to the Vienna PeeDee Belemnite standard (V-

PDB). **B – $^{13}C/^{12}C$ discrimination associated with photosynthetic CO₂ fixation (ε_p).** ε_p values

are calculated relative to a reference value, $\delta^{13}C_{CO2}$ = -8.4‰ (Keeling et al., 2017) (see **Materials**

and Methods for calculation). **A-B –** Mean (middle horizontal bars) and 1 SD (vertical error bars)

are shown in red. Asterisks indicate *t*-test results for pairwise comparisons indicated by horizontal

lines. ns – not significant; ** – $p < 0.01$; *** – $p < 0.001$.