**Title**

A billion years arms-race between viruses, virophages and eukaryotes

**Author names and affiliations**

Jose Gabriel Nino Barreat[1] and Aris Katzourakis[1,*]

[1]Department of Biology, University of Oxford, Oxford, United Kingdom.

[*]Corresponding author: aris.katzourakis@biology.ox.ac.uk

# **Abstract**

Bamfordviruses are arguably the most diverse group of viruses infecting eukaryotes. They include the Nucleocytoplasmic Large DNA viruses (NCLDVs), virophages, adenoviruses, *Mavericks* and Polinton-like viruses. Two main hypotheses for their origins have been proposed: the "nuclear-escape" and "virophage-first" hypotheses. The "nuclear-escape" hypothesis proposes an endogenous, *Maverick*-like ancestor which escaped from the nucleus and gave rise to adenoviruses and NCLDVs. In contrast, the "virophage-first" hypothesis proposes that NCLDVs coevolved with protovirophages; *Mavericks* then evolved from virophages that became endogenous, with adenoviruses escaping from the nucleus at a later stage. Here, we test the predictions made by both models and consider alternative evolutionary scenarios. We use a data set of the four core virion proteins sampled across the diversity of the lineage, together with Bayesian and maximum-likelihood hypothesis-testing methods, and estimate rooted phylogenies. We find strong evidence that adenoviruses and NCLDVs are not sister groups, and that *Mavericks* and Mavirus acquired the rve-integrase independently. We also found strong support for a monophyletic group of virophages (family *Lavidaviridae*) and a most likely root placed between virophages and the other lineages. Our observations support alternatives to the nuclear-escape scenario and a billion years evolutionary arms-race between virophages and NCLDVs.

**Key words**

Evolutionary arms-race, virophages, eukaryotes, NCLDVs.

## Introduction

Viruses in the kingdom *Bamfordvirae* comprise one of the most diverse groups in terms of their genome complexity, ecology and morphology. These dsDNA viruses include the largest viruses characterised to date (Legendre et al., 2014; Scheid, 2016), virophages which are virus parasites of other viruses (Fischer and Suttle, 2011; la Scola et al., 2008), endogenous viruses that colonise the genomes of eukaryotes (Kapitonov and Jurka, 2006; Pritham et al., 2007), and a diverse set of viruses that have been identified from metagenomic data (Bellas and Sommaruga, 2021; Moniruzzaman et al., 2020; Paez-Espino et al., 2019; Roux et al., 2017; Schulz et al., 2020; Yutin et al., 2015). More formally, these viruses are classified into the Nucleocytoplasmic Large DNA Viruses (NCLDVs) which comprise families in the phylum *Nucleocytoviricota* (Aylward et al., 2021), the virophage parasites of NCLDVs (family *Lavidaviridae*) (Fischer, 2020), adenoviruses (family *Adenoviridae*) which infect vertebrates (Harrach et al., 2019), the *Maverick/Polinton* endogenous viruses that colonise the genomes of eukaryotes (Kapitonov and Jurka, 2006; Pritham et al., 2007), Polinton-like viruses (PLVs) which are abundant in aquatic metagenomes (Bellas and Sommaruga, 2021; Yutin et al., 2015), and capsidless elements such as transpovirons (Desnues et al., 2012), mitochondrial and cytoplasmic linear plasmids (Krupovič and Koonin, 2015; Meinhardt et al., 1997). The wide taxonomic distribution of NCLDV and *Maverick* hosts, which comprise all major eukaryotic lineages (Kapitonov and Jurka, 2006; Pritham et al., 2007; Schulz et al., 2020), seem to point to an ancient origin of these viral groups.

Despite their diversity, the elements share an ancestral gene module used for capsid morphogenesis, which is the feature that distinguishes viruses from other mobile genetic elements (Koonin et al., 2021). The ancestral module is formed by double and single jelly-roll

capsid proteins, a family C5 (adenoviral-like) protease and a family FtsK/HerA DNA

packaging ATPase (Krupovič and Koonin, 2016; Yutin et al., 2015). In fact, the jelly-roll

capsid proteins and DNA packaging ATPase, also occur in phages from the families

*Turriviridae*, *Tectiviridae*, *Corticoviridae* and *Autolykiviridae*, which suggests a common viral

ancestor for eukaryotic bamfordviruses and their prokaryotic virus relatives (Koonin et al.,

2020; Woo et al., 2021). These observations are at odds with the suggestion that NCLDVs

originated by reductive evolution, possibly from a fourth domain of cellular life (Colson et al.,

2018; Legendre et al., 2012; Patil and Kondabagil, 2021).

Currently, two hypotheses have been proposed for the origin of the eukaryotic *Bamfordvirae*:

the "nuclear-escape" and "virophage-first" scenarios (Figure 1). According to the nuclear-

escape scenario, the elements in this lineage evolved from an endogenous virus in the nucleus

of an early eukaryote, a *Maverick*-like ancestor (Koonin and Krupovič, 2017; Krupovič and

Koonin, 2015). Adenoviruses, cytoplasmic linear plasmids and NCLDVs evolved after their

ancestor escaped from the nucleus (Koonin et al., 2015; Krupovič and Koonin, 2015).

Therefore, the nuclear-escape hypothesis predicts that adenoviruses form a clade with

NCLDVs and cytoplasmic linear plasmids, and that the proteins shared by cytoplasmic linear

plasmids and NCLDVs have a single common origin (Koonin and Krupovič, 2017; Krupovič

and Koonin, 2015). The nuclear-escape hypothesis is based on phylogenetic trees of the

protein-primed DNA polymerase B, which show a paraphyletic group of *Mavericks* at the base

of the other lineages (Koonin and Krupovič, 2017; Krupovič, Yutin, et al., 2016; Krupovič and

Koonin, 2015). However, the trees are missing NCLDVs which encode a non-homologous

nucleotide-primed DNA polymerase, and could therefore not be included in the analyses, and

only show the position for Mavirus-like virophages which encode the protein-primed DNA

polymerase.

The "virophage-first" scenario is based on the close similarity in gene content between the

Mavirus virophage and *Mavericks*, and the ability of Mavirus to integrate into eukaryotic

genomes (Fischer and Hackl, 2016; Fischer and Suttle, 2011). Integrated Mavirus virophages

reactivate upon giant virus infection and confer protection against *Cafeteria roenbergensis*

*virus* in the flagellate *Cafeteria burkhardae* (Fischer and Hackl, 2016). The virophage-first

scenario therefore suggests that an ancestral virophage evolved shortly after the origin of

NCLDVs, and was able to parasitise NCLDVs by virtue of their shared promoter and poly-A

sequences (Campbell et al., 2017; Fischer and Suttle, 2011). One of these lineages of ancestral

virophages would have gained an integrase and endogenised into eukaryotic hosts providing

an immune defence system, which gave rise to *Mavericks* and other elements (Fischer and

Suttle, 2011; Katzourakis and Aswad, 2014). A nuclear-escape would have still occurred for

adenoviruses and cytoplasmic linear plasmids, but not NCLDVs, which would belong to a

different clade suggesting the immediate ancestor of NCLDVs was an exogenous virus.

**Figure 1.** The two main hypotheses for the origin of virophages and NCLDVs. In the virophage first hypothesis, NCLDVs diverge early with its sister lineage evolving into protovirophages. In the nuclear escape hypothesis, NCLDVs descend from endogenous elements (encoding an integrase) that became exogenous; virophages then evolved to become their parasites.

Phylogenetics can be used to test competing evolutionary models of virus origins in a rigorous statistical framework. Each tree represents an evolutionary hypothesis which is inferred from matrices of aligned homologous characters with different states (the data). In molecular phylogenetics, nucleotide or amino acid alignments are used as the data to obtain the topology of a tree (and also rates or branch lengths). In statistical terms, the data matrices are used to calculate the tree likelihood (probability of the data given the parameters) in both maximum-likelihood and Bayesian frameworks (Schmidt and von Haeseler, 2009; Nascimento et al., 2017). In maximum likelihood, the preferred tree is the one with the parameter values that maximise the tree likelihood (Schmidt and von Haeseler, 2009). In Bayesian inference, a posterior probability distribution for the parameters is estimated from their prior distributions and the tree likelihood (probability of the parameters given the data) (Nascimento et al., 2017).

Together, these approaches can be used to assess the plausibility of competing evolutionary models that make different predictions about tree topologies and the position of the root.

Understanding the diversification of these viruses remains a major open question in virus evolution. To address this problem, we have analysed a data set of virus representatives sampled across the diversity of the eukaryotic bamfordvirus lineage and focused on the 4 core virion proteins: major and minor capsid proteins, protease and ATPase. We use explicit hypothesis-testing methods and estimate the position of the root in our phylogenies, which allowed us to infer the direction of evolution followed by the proteins in the morphogenetic module. Our results suggest a new model for the evolutionary origins of these viruses, which is consistent with an exogenous, non-virophage ancestor.

**Materials and Methods**

We used maximum-likelihood and Bayesian hypothesis-testing methods to compare the plausibility of the nuclear-escape versus alternative evolutionary scenarios. We focused on the 4 core virion proteins shared by viruses in this lineage: major and minor capsid proteins, ATPase and protease. Rooted phylogenies were inferred from the data using a relaxed molecular-clock or an outgroup rooting method. We then assessed whether adenoviruses and NCLDVs formed a monophyletic group as predicted by the nuclear-escape scenario. Finally, we studied the origin of the proteins shared by cytoplasmic linear plasmids and NCLDVs. A more detailed description of methods is presented below.

**Selection of viral sequences**

We first compiled a data set of viruses belonging to the families *Phycodnaviridae*, *Mimiviridae*, *Ascoviridae*, *Iridoviridae*, *Marseilleviridae*, *Asfarviridae*, *Poxviridae*, *Adenoviridae* and *Lavidaviridae* that were obtained from the NCBI using accession numbers listed in the International Committee for the Taxonomy of Viruses (ICTV) master species list 2020.v1. Previous analyses have shown that *Mavericks/Polintons* fall into two major groups based on the protein-primed DNA polymerase: the group I *Mavericks* (that include elements from vertebrates) and the group II *Mavericks* (which include elements present in insects) (Kapitonov and Jurka, 2006). Group I *Mavericks* can be distinguished from group II elements by the presence of a ~140 aa insertion similar to a bacterial "very-short-patch" (VSR) repair endonuclease (Kapitonov and Jurka, 2006). To make sure both groups were represented in our analyses, we included the vertebrate *Mavericks* reported in Barreat and Katzourakis (Barreat and Katzourakis, 2021), and added 12 new intact *Maverick* elements that we discovered in the genomes of insects and which lack the VSR insertion (Table S1).

We also included the sequences for Polinton-like viruses, virophages and NCLDVs that have been reported in the metagenomic works of Bellas and Sommaruga (Bellas and Sommaruga, 2021), Moniruzzaman et al. (Moniruzzaman et al., 2020), Schultz et al. (Schulz et al., 2020), Paez-Espino et al. (Paez-Espino et al., 2019), Roux et al. (Roux et al., 2017) and Yutin et al. (Yutin et al., 2015). Additional genomes for medusavirus (Yoshikawa et al., 2019), cedratvirus (Jeudy et al., 2020), tupanviruses (Abrahão et al., 2018), pandoraviruses (Antwerpen et al., 2016; Legendre et al., 2013, 2018), pithovirus (Legendre et al., 2014), mollivirus (Legendre et al., 2015), insectomime virus (Boughalmi et al., 2013), tokyovirus (Takemura, 2016), kaumoebavirus (Bajrai et al., 2016), faustoviruses (Benamar et al., 2016), pacmanvirus,

181  orpheovirus and the Dishui lake virophages (Sayers et al., 2021), were also included. In total,

182  the data set consisted of 9,222 virus genomes.

**Protein prediction and subsampling approach**

We devised a subsampling approach to arrive at a representative subset of sequences to carry

out the phylogenetic analyses and tests of topology. First, open reading frames for all sequences

were predicted with getorf in EMBOSS (Rice et al., 2000). Next, we used a set of Hidden

Markov Models (hmmbuild/hmmsearch, HMMER version 3.3.2 using the default parameters)

for each of the 4 core proteins involved in virion morphogenesis (major and minor capsid

proteins, ATPase and protease) to extract homologous proteins in 9 groups:

genomic/metagenomic virophages, genomic/metagenomic NCLDVs, metagenomic PLVs,

PLVs described in Yutin et al. (Yutin et al., 2015), adenoviruses, and *Mavericks 1* and *2*

lineages. We used the major capsid protein to make multiple sequence alignments in MAFFT

(Katoh et al., 2002) and construct approximate phylogenies for each group in FastTree 2 (Price

et al., 2010). A total of 2,803 major capsid proteins could be identified in the genome data,

which comprised: adenoviruses (71 sequences), *Mavericks 1/2* (28 sequences), NCLDVs

NCBI/metagenomic (2,031 sequences), PLVs Yutin/metagenomic (327 sequences), and

virophages NCBI/metagenomic (346 sequences). The trees were used to subsample 6

representatives from each of the 9 groups maintaining the maximal diversity in Treemmer

(Menardo et al., 2018). The final data set contained representatives of 54 taxa (Table S1).

**Protein multiple sequence alignment**

Using the list of representative taxa for the major capsid proteins, we retrieved the corresponding sequences for the ATPase, protease and minor capsid proteins in each genome. We confirmed that each prediction corresponded to its homology group using HHpred (Söding et al., 2005). Sequences were aligned in MAFFT (Katoh et al., 2002) and trimmed using trimAl (Capella-Gutiérrez et al., 2009). We then built a concatenated alignment using a custom script written in Python (van Rossum et al., 2009).

**Bayesian phylogenetic inference**

We constructed single-protein and concatenated rooted trees in BEAST 2 (Bouckaert et al., 2019). Best models for amino acid substitution were selected for each partition in ModelTest-NG (Darriba et al., 2020). The model LG+G+I (with 4 categories for the gamma rate heterogeneity) was used for the major capsid, protease and ATPase, while the model LG+G (with 4 categories for the gamma rate heterogeneity) was used for the minor capsid protein. We imposed monophyletic constraints at the level of well-established viral clades/families: NCLDVs, *Adenoviridae*, vertebrate *Mavericks* and invertebrate *Mavericks* (however, the monophyly between both groups of *Mavericks* was not constrained). The monophyly of virophages and Polinton-like virophages were not constrained. We conducted a Bayesian MCMC with a chain length of 140 million generations for the concatenated data set and 200 million generations for the single proteins (sampling every 5,000th generation and using relaxed molecular clocks). Convergence was assessed by inspecting the runs for good-mixing and stationarity, and ensuring that Effective Sample Sizes were > 200 for all parameters in Tracer (Rambaut et al., 2018).

**Maximum-likelihood phylogenetic inference**

We estimated a maximum-likelihood tree in RAxML-NG (Kozlov et al., 2019) using the concatenated data set and the same monophyletic constraints on well-established groups as described above. Proteins of the *Enterobacteria phage PRD1* (ATPase, major and minor capsid) were included in the alignment for outgroup rooting. We chose the best substitution models in ModelTest-NG (Darriba et al., 2020). For the protease and ATPase, we used model LG+G+I+F (with 4 categories for the gamma rate heterogeneity), and for the major and minor capsid proteins we used model LG+G+F (with 4 categories for the gamma rate heterogeneity). The analysis was started from 200 random starting trees and run with 2,200 bootstrap replicates.

**Akaike information criterion**

To compare the plausibility of the maximum-likelihood trees consistent with each hypothesis, we inferred the best supported phylogenies in RAxML-NG and then performed model-selection based on the Akaike information criterion (Kozlov et al., 2019; Posada and Buckley, 2004). Specifically, we compared a constrained model consistent with nuclear-escape where adenoviruses and NCLDVs are sister groups, to a model where the sister relationship between adenoviruses and NCLDVs was not constrained (alternative model). We included a partitioned analysis of the concatenated data set and 3 core proteins individually (protease, major and minor capsid), using the amino acid substitution models described in the previous section. The analyses were run with 200 random starting trees and we kept the estimate for the log-likelihood of the best tree model for the calculation of the AICs. The AIC for each model was size-corrected given that the alignment (sample) sizes were small compared to the number of

free parameters in each model, or $n/k < 40$ (Posada and Buckley, 2004; Symonds and Moussalli,

2011). For the comparison we calculated the Akaike weights which can be interpreted as the

normalised relative likelihood of the model given the data (Posada and Buckley, 2004), or as

an analogue of the probability that a model is the best approximating model given the data

within a set of candidates (Symonds and Moussalli, 2011). The ATPase was not included in

the single-protein analyses since adenoviruses encode a non-homologous ABC family ATPase

(Burroughs et al., 2007), and therefore it cannot be used to test their evolutionary relationships

to other bamfordviruses.

**Posterior model odds**

The tree topologies obtained in the Bayesian MCMCs for the single proteins and the

concatenated data set, were loaded and filtered in PAUP (Swofford, 1998). Following the

posterior model odds method of Bergsten et al. (Bergsten et al., 2013), we counted the number

of trees which contained a sister grouping of adenoviruses and NCLDVs (nuclear-escape

hypothesis) and compared them to the number of trees inconsistent with this grouping

(alternative hypotheses). The posterior model odds were calculated as the ratio of these two

quantities (Bergsten et al., 2013). The ATPase was again excluded from the single-protein

analysis given that adenoviruses could not be included.

**Marginal likelihood and Bayes factors**

We estimated the marginal likelihood of each hypothesis using stepping-stone sampling. We

ran the analysis on the concatenated data set, for 20 million generations in MrBayes 3 (Ronquist

and Huelsenbeck, 2003). For both analyses, we used the monophyletic constraints described

above. In the marginal calculation for the nuclear-escape hypothesis, we used a topological

constraint on the sister grouping of adenoviruses and NCLDVs, while we imposed a negative

topological constraint on the monophyly of adenoviruses and NCLDVs for the alternatives

(adenoviruses and NCLDVs cannot be sister groups). We calculated the Bayes factors as the

number $e$ to the power of the difference in log-marginal likelihoods, and compared the resulting

value to the table of Kass and Raftery (Kass and Raftery, 1995).

**Cytoplasmic linear plasmids**

Cytoplasmic linear plasmids have lost the ancestral module of genes involved in formation of

the capsid, but they have gained three genes encoding a mRNA capping-enzyme, a helicase

and the Rpb2 subunit of the DNA-dependent RNA-polymerase II (Koonin, Krupovič, et al.,

2015; Krupovič and Koonin, 2015). We used the proteins in cytoplasmic linear plasmids (Table

S3), to search for homologous sequences in the non-redundant protein database using blastp

(Altschul et al., 1990) and restricting searches to 'Eukaryota (taxid:2759)' and 'Viruses

(taxid:10239)'. The identity of significant matches (evalue < 1e-10) was confirmed with

HHpred (Söding et al., 2005), and in the case of eukaryotes, only sequences mapping to

chromosome assemblies were used. Virus, eukaryotic and plasmid homologues were aligned

in MAFFT (Katoh et al., 2002) and conserved blocks recovered from trimAl (Capella-Gutiérrez

et al., 2009). We chose the best models for protein evolution in ModelTest-NG (Darriba et al.,

2020) and ran a maximum-likelihood tree search in RAxML-NG (Kozlov et al., 2019) with

1,000 non-parametric bootstrap replicates for each protein separately.

**Data availability**

The Hidden Markov Models used to find homologues of the four core proteins (n = 38), and

concatenated and single-protein alignments used for phylogenetic inference have been

deposited in Figshare (DOI: 10.6084/m9.figshare.19576117). The scripts used for sequence

concatenation and for the AIC analyses are available on GitHub

(https://github.com/josegabrielnb/virophage-origins).

**Results**

The rooted Bayesian and maximum-likelihood trees estimated from the concatenated data had

the best overall support and shared important topological similarities (Figures 2 and 3). In the

Bayesian tree, adenoviruses were placed in a clade with *Mavericks* and most PLVs with a high

posterior probability (0.94, Figure 2), and to the exclusion of NCLDVs. This arrangement was

also observed in the maximum likelihood tree topology shown in Figure 3, although with a low

bootstrap support (34%). The virophages (family *Lavidaviridae*) emerged as a strongly

supported monophyletic group on both trees (posterior probability = 1 in the Bayesian analysis,

95% bootstrap support in the maximum-likelihood analysis), while Polinton-like viruses were

clearly polyphyletic.

The position of the root inferred both through the Bayesian relaxed-clock analysis and by

outgroup rooting with the *Enterobacteria phage PRD1* (*Tectiviridae*), was between virophages

and all the other viral lineages. In the Bayesian analysis, the root was placed on this position

in 54% of trees (Figure 2), while in the maximum-likelihood analysis the clade of

(Adenoviruses + *Mavericks* + PLVs + NCLDVs), i.e. to the exclusion of virophages, received

a bootstrap support of 73% (Figure 3). We also examined alternative placements of the root

sampled during the Bayesian Markov-chain Monte Carlo (MCMC) to assess the frequency of

other hypotheses (Table S2). Rooting on the clade formed by NCLDVs + PLV BS_539

(ERX2821583_NODE_576) was observed in 27.4% of trees. The root positions at the next

highest frequencies were on the branch of PLV BS_539 (5.6%), NCLDVs (5.3%) and PLV

BS_539 + NCLDVs + virophages (5%). Roots on other branches were observed at frequencies

lower than 1% (Table S2).

The monophyly of virophages and the root on this branch were also observed in the single-

protein trees. The major and minor capsid single-protein trees recovered a monophyletic

*Lavidaviridae* (0.97 and 0.52 posterior probability), and also inferred the root to be placed

between virophages and the other viruses (29% and 28%, Figures S1-S2). Although the root

was observed at a different position for the ATPase and protease trees; these analyses also

failed to recover a monophyletic family of virophages (Figures S3-S4). However, when

virophages were constrained to be monophyletic the root was placed again between virophages

and the other lineages, and thus in agreement with our findings using the concatenated data,

major and minor capsid proteins (Figures S5-S6).

**Figure 2.** Rooted Bayesian maximum clade credibility tree of the major lineages of eukaryotic viruses in the kingdom *Bamfordvirae*. The tree is based on the concatenated alignment of 4 core proteins involved in virion morphogenesis (major and minor capsid proteins, ATPase and protease). Tree computed in BEAST 2 (Bouckaert et al., 2019) using a relaxed molecular clock and 140 million generations (relative burn-in of 25%). Black: reference viral genomes, Blue: endogenous elements, Orange: metagenomic sequences.

 **Figure 3.** Outgroup-rooted maximum likelihood tree of the major lineages of eukaryotic viruses in the

kingdom *Bamfordvirae*. The tree is based on the concatenated alignment of 4 core proteins involved in

virion morphogenesis (major and minor capsid proteins, ATPase and protease). *Enterobacteria phage*

*PRD1* (*Tectiviridae*) was used as the outgroup for rooting. Tree computed in RAxML-NG with 200

random starting trees and 2,200 bootstraps (Kozlov et al., 2019). Black: reference viral genomes, Blue:

endogenous elements, Orange: metagenomic sequences.

The Bayesian and maximum-likelihood tests of topology were also consistent with each other.

We used the Akaike information criterion (AIC) to compare the plausibility of the maximum-

likelihood topologies consistent with the nuclear-escape and alternative scenarios. All the

comparisons preferred a non-sister grouping of adenoviruses and NCLDVs (Table 1). Indeed,

the Akaike weights, which can be interpreted as the normalised relative likelihood of the

models given the data (Posada and Buckley, 2004), strongly favoured the alternative scenario

over the nuclear-escape hypothesis (Concatenated data = 99.7%, Protease = 99.7%, Major

capsid protein = 72%, Minor capsid protein = 94.3%). The approach using the posterior model

odds, calculated following the method of Bergsten et al. (Bergsten et al., 2013), also favoured

the alternative scenarios in all of the data sets: concatenated and single-protein (Table 2). This

was strongest for the concatenated data (posterior model odds, $H_0/H_1$= 0.000536), but the same

pattern held for the single-protein analyses (posterior model odds = {Protease: 0.00166, Major

capsid: 0.11485, Minor capsid: 0.000639}). The Bayesian stepping-stone analysis on the

concatenated data also strongly supported the alternative scenario (NCLDVs and adenoviruses

are not sister groups), over the nuclear-escape hypothesis (NCLDVs and adenoviruses are sister

groups) (Table 3).

As an independent test of the predictions made by the nuclear-escape hypothesis, we analysed

the origin of the transcriptional proteins shared between NCLDVs and cytoplasmic linear

plasmids. According to the nuclear-escape hypothesis, these genes were acquired by the most

recent common ancestor of cytoplasmic linear plasmids and NCLDVs, and should thus have a

single origin. However, our analyses indicate that these proteins were acquired independently

by cytoplasmic linear plasmids and NCLDVs. The maximum-likelihood trees for the Rbp2

subunit of the DNA-directed RNA polymerase, helicase and mRNA-capping proteins all

suggest they have independent origins in NCLDVs and cytoplasmic linear plasmids (Figure

S7). The homologues specific to cytoplasmic linear plasmids clustered with those of eukaryotes

and their distribution agrees with a monophyletic origin in this group. Interestingly, the

phylogenetic patterns for NCLDVs were consistent with a single-capture (monophyletic

origin) for the mRNA-capping enzyme and the helicase, while the distribution of the Rpb2 was

consistent with multiple-captures/exchanges (polyphyletic origin).

**Table 1.** Comparison between the nuclear-escape and alternative maximum-likelihood models based

on the Akaike information criterion (AIC). The log-likelihoods for the models were obtained from the

best maximum-likelihood tree consistent with each hypothesis found in RAxML-NG (Kozlov et al.,

2019). Results are shown for the concatenated data set of four core proteins (ATPase, protease, major

capsid and minor capsid) and for the protease, major and minor capsid proteins, respectively. The AIC

was size-corrected given that the alignment (sample) size was small relative to the number of free

parameters, i.e. n/k < 40 (Posada and Buckley, 2004; Symonds and Moussalli, 2011). The best model

is highlighted in boldface. The "alternatives" refer to non "nuclear-escape" scenarios, that is, models

which are not consistent with the predictions made by the nuclear-escape.

| Model | Characters | Log-likelihood | AIC[1] | AICc[2] | ΔAICc[3] | Weight[4] |
|---|---|---|---|---|---|---|
| Nuclear-escape ($M_0$) | Concatenated | -41,618.19 | 83,616.37 | 83,864.09 | 11.41 | 0.003 |
| **Alternatives ($M_1$)** | **Concatenated** | **-41,612.48** | **83,604.96** | **83,852.68** | **0.00** | **0.997** |
| Nuclear-escape ($M_0$) | Protease | -6,323.233 | 12,898.47 | 10,898.22 | 11.37 | 0.003 |
| **Alternatives ($M_1$)** | **Protease** | **-6,317.550** | **12,887.10** | **10,886.85** | **0.00** | **0.997** |
| Nuclear-escape ($M_0$) | Major capsid | -16,305.52 | 32,861.04 | 34,173.54 | 1.90 | 0.279 |
| **Alternatives ($M_1$)** | **Major capsid** | **-16,304.57** | **32,859.14** | **34,171.64** | **0.00** | **0.721** |
| Nuclear-escape ($M_0$) | Minor capsid | -10,234.56 | 20,719.11 | 17,855.48 | 5.62 | 0.057 |
| **Alternatives ($M_1$)** | **Minor capsid** | **-10,231.75** | **20,713.50** | **17,849.86** | **0.00** | **0.943** |

[1] $AIC_i = -2 \cdot LogL_i + 2 \cdot K$

[2] $AICc_i = AIC_i + \frac{2K(K+1)}{(n-K-1)}$

[3] $\Delta AICc_i = AICc_i - \min(AICc_i, AICc_j)$

[4] $Weight_i = \frac{e^{-\Delta AICc_i/2}}{\sum_i^j e^{-\Delta AICc_i/2}}$

**Table 2.** Posterior model odds of the nuclear-escape and alternative hypotheses using concatenated and

single-protein data sets. Tree topologies consistent with each hypothesis were filtered and counted from

a Bayesian MCMC following the method of Bergsten et al. (2013). All ratios favour the alternative

scenarios to the nuclear-escape hypothesis. The best model is highlighted in boldface. The

"alternatives" refer to non-"nuclear-escape" scenarios, that is, models which are not consistent with the

predictions made by the nuclear-escape.

| Model | Characters | MCMC tree frequency | Posterior model odds $P(M_0 \mid X)/P(M_1 \mid X)$ |
|---|---|---|---|
| Nuclear-escape ($M_0$) | Concatenated | 15/28,001 (0.0536%) | |
| **Alternatives ($M_1$)** | **Concatenated** | **27,986/28,001 (99.946%)** | $5.36 \cdot 10^{-4} < 1$ |
| Nuclear-escape ($M_0$) | Protease | 333/200,001 (0.166%) | |
| **Alternatives ($M_1$)** | **Protease** | **199,668/200,001 (99.998%)** | $1.66 \cdot 10^{-3} < 1$ |
| Nuclear-escape ($M_0$) | Major capsid | 20,605/200,001 (10.302%) | |
| **Alternatives ($M_1$)** | **Major capsid** | **179,396/200,001 (89.697%)** | $1.15 \cdot 10^{-1} < 1$ |
| Nuclear-escape ($M_0$) | Minor capsid | 128/200,001 (0.0639%) | |
| **Alternatives ($M_1$)** | **Minor capsid** | **199,873/200,001 (99.936%)** | $6.39 \cdot 10^{-4} < 1$ |

**Table 3.** Bayesian stepping-stone analysis of the nuclear-escape and alternative hypotheses. Each

scenario was run on the concatenated data set in MrBayes 3 (Ronquist and Huelsenbeck, 2003) for 20

million generations (average standard deviation of split frequencies < 0.01). The Bayes factor strongly

rejects a sister relationship between adenoviruses and NCLDVs (nuclear-escape hypothesis). The best

model is highlighted in boldface. The "alternatives" refer to non "nuclear-escape" scenarios, that is,

models which are not consistent with the predictions made by the nuclear-escape.

| Model | Likelihood of best state (cold chain) | Log-Marginal-likelihood (ln) | Mean Log-marginal-likelihood (ln) | Bayes factor $P(X \mid M_0)/P(X \mid M_1)$ |
|---|---|---|---|---|
| Nuclear-escape ($M_0$) | Run 1: -36,318.81 | Run 1: -36,376.42 | -36,353.44 | $3.5 \times 10^{-94} \ll 1$*** |
| | Run 2: -36,318.81 | Run 2: -36,352.75 | | |
| **Alternatives ($M_1$)** | **Run 1: -35,965.90** | **Run 1: -36,137.56** | **-36,138.25** | |
| | **Run 2: -35,981.46** | **Run 2: -36,450.19** | | |

446 ***Strong support against $M_0$

## Discussion

To gain a better understanding of the evolutionary history of the bamfordviruses of eukaryotes, we analysed a set of four core virion morphogenesis proteins (major and minor capsids, DNA-packaging ATPase and protease) sampled across the diversity of the lineage and using an explicit hypothesis-testing framework. In addition, we analysed the origins of 3 proteins shared by cytoplasmic linear plasmids and NCLDVs. We found strong evidence against the nuclear-escape hypothesis. We also found support for a position of the root between virophages (family *Lavidaviridae*) and the other viral lineages. This position of the root suggests a new evolutionary model for the origin of these viral lineages. However, we recognise that some alternative root positions were also observed at lower frequencies in the Bayesian analyses, which could be interpreted as "virophage-first"-like scenarios.

We have shown that adenoviruses and NCLDVs are not sister groups as proposed by the nuclear-escape hypothesis (Koonin and Krupovič, 2017; Krupovič and Koonin, 2015). Instead, adenoviruses belong to a clade with *Mavericks*, to the exclusion of NCLDVs. The phylogenies of the proteins shared by cytoplasmic linear plasmids and NCLDVs also disagree with a nuclear escape. Indeed, phylogenies of the protein-primed DNA polymerase place cytoplasmic linear plasmids as the sister-group to adenoviruses (Krupovič, Yutin, et al., 2016), implying that adenoviruses, cytoplasmic linear plasmids and NCLDVs form a clade. If a single nuclear-escape had occurred we would expect the Rbp2, helicase and mRNA capping enzymes shared by cytoplasmic linear plasmids and NCLDVs to have a monophyletic origin (since they descend from the same "escaped" ancestor). However, their distribution agrees with a more recent origin in cytoplasmic linear plasmids. These observations agree with the more restricted taxonomic distribution of cytoplasmic linear plasmids, which are known only to infect fungi in

the Order Saccharomycetales (Division Ascomycota) (Meinhardt et al., 1997), in contrast to

the taxonomically diverse eukaryotic hosts of NCLDVs.

Virophages emerged as a highly supported monophyletic group in our analyses. This supports

the validity of the family *Lavidaviridae* which has been based on shared gene contents of

virophages and their parasitic (or commensal) lifestyles (Fischer, 2020; Krupovič, Kuhn, et al.,

2016). By using two independent rooting methods, relaxed-molecular clocks and outgroup

rooting, we found that the root was placed most confidently between virophages and the other

lineages; making them the most basal lineage of eukaryotic bamfordviruses. The basal position

of virophages has also been found in an independent work that looked at phylogenies based on

the ATPase and a concatenation of the ATPase and the major capsid protein (Woo et al., 2021).

This basal position of virophages and their divergence prior to the origin of NCLDVs, suggests

that the protovirophages were not parasites of other viruses. Instead, they would have become

parasites of other viruses at a later point once the most recent common ancestor of NCLDVs

had evolved (see Ideas and speculation).

The tree topologies that we estimated (Figure 2 and Figure 3), also suggest that the most recent

common ancestor of *Mavericks*, adenoviruses and PLVs was not a virophage, which is at odds

with a virophage-first scenario. Indeed, the basal position of virophages is inconsistent with

the virophage-first hypothesis which proposes that protovirophages coevolved with NCLDVs,

acquired an integrase and then gave rise to *Mavericks* and other elements in the lineage

(Campbell et al., 2017; Fischer and Suttle, 2011). However, we found that the second-highest

frequency root sampled in the Bayesian MCMC was on the branch of PLV BS_539 + NCLDVs

(frequency = 27.4%). Rooting on this branch would be consistent with a "virophage-first"-like

scenario, where the lineage of NCLDVs and associated viruses evolved early and in parallel

with protovirophages. Therefore, we cannot definitely rule out a "virophage-first"-like

scenario. In contrast, root placements on the branch leading to *Mavericks* and consistent with

nuclear-escape were observed in 0% of the trees, which seems to rule out this possibility (Table

S2).

The relative timing of events suggests that the most recent common ancestor of eukaryotic

bamfordviruses existed more than a billion years ago. There is general agreement based on

paleontological and molecular evidence that the Last Eukaryotic Common Ancestor (LECA)

existed at least 1 billion years ago (Porter, 2020). Analyses of genetic exchanges between the

DNA-dependent RNA polymerases of NCLDVs and eukaryotes have determined that the two

super-clades of NCLDVs (PAM and MAPI) already existed before the origin of LECA

(Guglielmini et al., 2019). Moreover, discovery and phylogenetic analysis of actin and actin-

related proteins encoded in NCLDVs, also suggest that early gene transfers occurred between

these viruses and proto-eukaryotes before the emergence of LECA (da Cunha et al., 2022). The

basal placement that we observed for virophages, or alternatively, their early origin after the

divergence of the NCLDV lineage ("virophage-first"-like scenario), suggest that virophages

evolved after the origin of the first NCLDVs during the diversification of the early eukaryotes.

These observations, together with the existence of various phage relatives that infect bacteria

and archaea, point to a very early origin of the kingdom *Bamfordvirae*, perhaps extending to

the initial stages of cellular life. Studies that consider the time-dependent rate phenomenon in

reconstructing viral evolution may be able to shed light on these issues (Aiewsakun and

Katzourakis, 2016; Ghafari et al., 2021).

## Ideas and speculation

Our analyses favour a new model for the evolution of the major lineages of eukaryotic bamfordviruses; we call this model the "exogenous, non-virophage scenario". The model is presented in Figure 4, where we have mapped shared character states onto a diagram of the concatenated Bayesian phylogeny (which has the best overall support). According to this idea, the most recent common ancestor of the eukaryotic bamfordviruses was a small exogenous dsDNA virus with a linear genome, flanked by inverted repeats. The virus had an autonomous (i.e. non-virophage) lifestyle. It used a protein-primed DNA polymerase B for replication and its morphogenetic module was formed by the major and minor jelly-roll capsid proteins, an adenoviral-like protease and the FtsK/HerA family DNA-packaging ATPase. The gain of the adenoviral-like protease differentiated this eukaryotic lineage from their phage relatives (Krupovič and Koonin, 2015). The first divergence from this ancestor led to the emergence of the protovirophages, and another lineage that would give rise to *Mavericks*, adenoviruses, PLVs and NCLDVs (Figure 4).

**Figure 4.** Evolutionary model for the origin of the major lineages of eukaryotic viruses in the kingdom *Bamfordvirae*. The viral ancestor is inferred to have been an exogenous virus, while the rve-integrase was captured independently by the clade of *Mavericks* + Polinton-like virus BS_13 and Mavirus, possibly by horizontal gene transfer. Virophages evolved from an autonomous virus that became specialised to parasitise the ancestor of NCLDVs. The vertical cross-hatching indicates that the trait is found in some but not all members of the group. Acronyms refer to genes and genomic features present in the viral genomes: ***pPOLB*** (protein-primed DNA polymerase B), ***MCP*** (major capsid protein), ***mCP*** (minor capsid protein), ***int*** (rve-type integrase), ***pro*** (adenoviral-like protease), ***atp*** (FtsK/HerA DNA packaging ATPase), ***TIRs*** (terminal inverted repeats).

The second divergence would have given rise to the lineage from which NCLDVs evolved. The most recent common ancestor of NCLDVs is believed to have already had a complex genome (~40 genes), which involved the gain of numerous genes from their eukaryotic hosts, other viruses and bacteria (Iyer et al., 2006; Koonin and Yutin, 2010; Yutin and Koonin, 2012). Once this virus evolved the capacity to exploit significant cell resources, there would have been

a window of opportunity for protovirophages to evolve into specialised parasites of NCLDVs,

occupying this new ecological niche. The close match between virophage and NCLDV

regulatory sequences, which underlie the capacity of virophages to parasitise NCLDVs, may

have evolved by parallel evolutionary changes on the sequences of their shared virus ancestor.

It is plausible that functional regulatory sequences in virophages could evolve *de novo* by a

few changes in the ancestral promoters/terminators. For example, functional *lac* operon

promoters have been evolved successfully even from random sequences in the presence of

lactose in *E. coli* (Yona et al., 2018). A deletion mutant in the ORF 8 gene of the Guarani

virophage expanded its host range to previously non-permissive giant viruses, so the

acquisition and mutation of proteins involved in virus-virophage interactions may also be

critical for virophage adaptation (Mougari et al., 2020). Interestingly, the exchange of the

protein-primed DNA polymerase B by a nucleotide-primed DNA polymerase B (Mönttinen et

al., 2021; Yutin and Koonin, 2012), may have been an early counter-measure of NCLDVs to

decrease the parasitic burden imposed by virophages. The acquisition of this new polymerase,

may be in part responsible for the considerable increases in genome size that we see in

NCLDVs relative to other eukaryotic bamfordviruses. Therefore, we can hypothesise that the

large genomes of NCLDVs, which is one of their most distinctive features, could have evolved

as a result of their ancient interaction with virophages.

The distribution of the rve-integrase on the phylogeny suggests it was acquired independently

on two separate occasions. Indeed, the rve-integrase is a universal feature of *Mavericks* + PLV

BS_13, while it does not seem to be present in any of their closest relatives. Therefore, the rve-

integrase seems to be a unique derived feature of the clade of *Mavericks* + PLV BS_13.

Mavirus also encodes a rve-integrase but it appears to be the only virophage with this gene (no

other hits found in a blastp search of the nr database using the labels "*Lavidaviridae*" +

"unclassified *Lavidaviridae*"). Since Mavirus is firmly nested within the phylogenies for the

morphogenetic module, it seems likely that this is a unique derived feature of Mavirus. The

most parsimonious explanation for these observations is that the integrase gene was acquired

on two separate occasions and that the common ancestor of Mavirus and *Mavericks* was an

exogenous virus without an integrase (# character state changes = 2 gains). This is in line with

a previous proposal that Mavirus evolved by recombination of an ancestral virophage with a

*Maverick* (Yutin et al., 2013).

**Concluding remarks**

Our findings strengthen the view of a viral origin for NCLDVs and other eukaryotic elements

in the kingdom *Bamfordvirae*. We have decisively shown that adenoviruses and NCLDVs are

not sister groups, and that adenoviruses, virophages, PLVs and NCLDVs did not escape from

the nucleus. The rve-integrase appears to have been captured independently by *Mavericks* +

PLV BS_13 on the one hand, and by Mavirus on the other. These observations support a new

evolutionary model that is different from the nuclear-escape and virophage-first scenarios. This

new model proposes that the ancestor of the eukaryotic viruses in the kingdom *Bamfordvirae*

was an exogenous, non-virophage DNA virus with a small genome. The lifestyle of virophages

would have evolved at a later stage as these became specialised parasites of the ancestral

NCLDVs. Darwin's closing words in the *Origin of Species* are well-suited for this group of

viruses: "from so simple a beginning endless forms most beautiful and most wonderful have

been, and are being, evolved" (Darwin, 1859).

## Acknowledgements

We wish to thank Peter Simmonds and Alexander Suh for their critical reading and comments on the manuscript, which served to improve this work. We also thank the reviewers for their recommendations and feedback. This work was supported by a doctoral scholarship (Dr. Jose Gregorio Hernandez Award) to JGNB made by the National Academy of Medicine of Venezuela and Pembroke College, Oxford.

## References

Abrahão J, Silva L, Silva LS, Khalil JYB, Rodrigues R, Arante T, Assis F, Boratt P, Andrade M, Kroon, EG, Ribeiro B, Bergier I, Seligmann H, Ghigo E, Colson P, Levasseur A, Kroemer G, Raoult D and la Scola B. (2018). Tailed giant Tupanvirus possesses the most complete translational apparatus of the known virosphere. *Nature Communications* 9(1). https://doi.org/10.1038/s41467-018-03168-1

Aiewsakun P and Katzourakis A. (2016). Time-dependent rate phenomenon in viruses. *Journal of Virology*, 90(16), 7184–7195. doi: 10.1128/JVI.00593-16

Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, 215(3), 403–410. doi: 10.1016/S0022-2836(05)80360-2

Antwerpen MH, Georgi E, Zoeller L, Woelfel R, Stoecker K and Scheid P. (2016). Whole-genome sequencing of a Pandoravirus isolated from keratitis-inducing *Acanthamoeba*. *Genome Announcements*, 3(2). https://doi.org/10.1128/genomeA.00136-15

Aylward FO, Moniruzzaman M, Ha AD and Koonin, EV. (2021). A phylogenomic framework for charting the diversity and evolution of giant viruses. *PLoS Biology*. doi: 10.1371/JOURNAL.PBIO.3001430

Bajrai LH, Benamar S, Azhar EI, Robert C, Levasseur A, Raoult D, and la Scola B. (2016).

Kaumoebavirus, a new virus that clusters with faustoviruses and *Asfarviridae*. Viruses 8(11). doi:

10.3390/v8110278

Barreat JGN and Katzourakis A. (2021). Phylogenomics of the *Maverick* virus-like mobile genetic

elements of vertebrates. *Molecular Biology and Evolution.* doi: 10.1093/molbev/msaa291

Bellas, CM and Sommaruga R. (2021). Polinton-like viruses are abundant in aquatic ecosystems.

*Microbiome* 9(1). doi: 10.1186/s40168-020-00956-0

Benamar S, Reteno DGI, Bandaly V, Labas N, Raoult D and la Scola B. (2016). Faustoviruses:

comparative genomics of new megavirales family members. *Frontiers in Microbiology* 7(FEB).

doi: 10.3389/fmicb.2016.00003

Bergsten J, Nilsson AN and Ronquist F. (2013). Bayesian tests of topology hypotheses with an example

from diving beetles. *Systematic Biology* 62(5). doi: 10.1093/sysbio/syt029

Bouckaert R, Vaughan TG, Barido-Sottani J, Duchêne S, Fourment M, Gavryushkina A, Heled J, Jones

G, Kühnert D, de Maio N, Matschiner M, Mendes FK, Müller NF, Ogilvie HA, du Plessis L,

Popinga A, Rambaut A, Rasmussen D, Siveroni I, … Drummond AJ. (2019). BEAST 2.5: An

advanced software platform for Bayesian evolutionary analysis. *PLoS Computational Biology*

15(4). doi: 10.1371/journal.pcbi.1006650

Boughalmi M, Pagnier I, Aherfi S, Colson P, Raoult D and la Scola B. (2013). First isolation of a

marseillevirus in the diptera syrphidae eristalis tenax. *Intervirology* 56(6). doi:

10.1159/000354560

Burroughs A, Iyer L and Aravind L. (2007). Comparative genomics and evolutionary trajectories of

viral ATP dependent DNA-packaging systems. *Genome Dynamics* (Vol. 3). doi:

10.1159/000107603

Campbell S, Aswad A and Katzourakis A. (2017). Disentangling the origins of virophages and

*Polintons. Current Opinion in Virology.* doi: 10.1016/j.coviro.2017.07.011

Capella-Gutiérrez S, Silla-Martínez JM and Gabaldón T. (2009). trimAl: A tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25(15). doi:

10.1093/bioinformatics/btp348

Colson P, Levasseur A, la Scola B, Sharma V, Nasir A, Pontarotti P, Caetano-Anollés G and Raoult D. (2018). Ancestrality and mosaicism of giant viruses supporting the definition of the fourth TRUC of microbes. *Frontiers in Microbiology* 9. doi: 10.3389/fmicb.2018.02668

da Cunha V, Gaia M, Ogata H, Jaillon O, Delmont TO and Forterre P. (2022). Giant viruses encode actin-related proteins. *Molecular Biology and Evolution*, 39(2). doi: 10.1093/molbev/msac022

Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B and Flouri T. (2020). ModelTest-NG: a new and scalable tool for the selection of dna and protein evolutionary models. *Molecular Biology and Evolution*, 37(1). doi: 10.1093/molbev/msz189

Darwin, CR. (1859). On the origin of species by means of natural selection, or the preservation of favoured races in the struggle for life. (1st edition). John Murray.

Desnues C, la Scola B, Yutin N, Fournous G, Robert C, Azza S, Jardot P, Monteil S, Campocasso A, Koonin EV and Raoult D. (2012). Provirophages and transpovirons as the diverse mobilome of giant viruses. *Proc Natl Acad Sci U S A* 109(44). doi: 10.1073/pnas.1208835109

Fischer MG. (2020). The virophage family *Lavidaviridae*. *Current Issues in Molecular Biology*, 40. doi: 10.21775/cimb.040.001

Fischer MG and Hackl T. (2016). Host genome integration and giant virus-induced reactivation of the virophage Mavirus. *Nature* 540(7632). doi: 10.1038/nature20593

Fischer MG and Suttle CA. (2011). A virophage at the origin of large DNA transposons. *Science*. 10.1126/science.1199412

Ghafari M, Simmonds P, Pybus OG and Katzourakis A. (2021). A mechanistic evolutionary model explains the time-dependent pattern of substitution rates in viruses. *Current Biology* 31(21). doi: 10.1016/j.cub.2021.08.020

Guglielmini J, Woo AC, Krupovič M, Forterre P and Gaia M. (2019). Diversification of giant and large eukaryotic dsDNA viruses predated the origin of modern eukaryotes. *Proc Natl Acad Sci U S A*. doi: 10.1073/pnas.1912006116

Harrach B, Tarján ZL and Benkő M. (2019). Adenoviruses across the animal kingdom: a walk in the zoo. *FEBS Letters* (Vol. 593, Issue 24). doi: 10.1002/1873-3468.13687

Iyer LM, Balaji S, Koonin EV and Aravind L. (2006). Evolutionary genomics of nucleo-cytoplasmic

large DNA viruses. *Virus Research* 117(1). doi: 10.1016/j.virusres.2006.01.009

Jeudy S, Rigou S, Alempic JM, Claverie JM, Abergel C and Legendre M. (2020). The DNA methylation

landscape of giant viruses. *Nature Communications* 11(1). doi: 10.1038/s41467-020-16414-2

Kapitonov VV and Jurka, J. (2006). Self-synthesizing DNA transposons in eukaryotes. *Proc Natl Acad*

*Sci U S A* 103(12). doi: 10.1073/pnas.0600833103

Kass R. E and Raftery AE. (1995). Bayes factors. *Journal of the American Statistical Association*

90(430). doi: 2291091

Katoh K, Misawa K, Kuma KI and Miyata T. (2002). MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. *Nucleic Acids Research* 30(14). doi:

10.1093/nar/gkf436

Katzourakis A and Aswad A. (2014). The origins of giant viruses, virophages and their relatives in host

genomes. *BMC Biology*. doi: 10.1186/s12915-014-0051-y

Koonin EV, Dolja VV and Krupovič M. (2015). Origins and evolution of viruses of eukaryotes: the

ultimate modularity. *Virology*. doi: 10.1016/j.virol.2015.02.039

Koonin EV, Dolja VV, Krupovič M and Kuhn, JH. (2021). Viruses defined by the position of the

virosphere within the replicator space. *Microbiology and Molecular Biology Reviews* 85(4). doi:

10.1128/mmbr.00193-20

Koonin EV, Dolja VV, Krupovič M, Varsani A, Wolf YI, Yutin N, Zerbini FM and Kuhn JH. (2020).

Global organization and proposed megataxonomy of the virus world. *Microbiology and*

*Molecular Biology Reviews* 84(2). doi: 10.1128/mmbr.00061-19

Koonin EV and Krupovič M. (2017). *Polintons*, virophages and transpovirons: a tangled web linking

viruses, transposons and immunity. *Current Opinion in Virology* 25. doi:

10.1016/j.coviro.2017.06.008

Koonin EV, Krupovič M and Yutin N. (2015). Evolution of double-stranded DNA viruses of

eukaryotes: From bacteriophages to transposons to giant viruses. *Annals of the New York*

*Academy of Sciences* 1341(1). doi: 10.1111/nyas.12728

Koonin EV and Yutin N. (2010). Origin and evolution of eukaryotic large nucleo-cytoplasmic DNA

viruses. *Intervirology*. doi: 10.1159/000312913

Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A and Wren J. (2019). RAxML-NG: A fast,

scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics*

35(21). doi: 10.1093/bioinformatics/btz305

Krupovič M and Koonin, EV. (2015). *Polintons*: A hotbed of eukaryotic virus, transposon and plasmid

evolution. *Nature Reviews Microbiology* 13(2). doi: 10.1038/nrmicro3389

Krupovič M and Koonin EV. (2016). Self-synthesizing transposons: unexpected key players in the

evolution of viruses and defense systems. *Current Opinion in Microbiology* 31. doi:

10.1016/j.mib.2016.01.006

Krupovič M, Kuhn JH and Fischer MG. (2016). A classification system for virophages and satellite

viruses. *Archives of Virology* 161(1). doi: 10.1007/s00705-015-2622-9

Krupovič M, Yutin N and Koonin EV. (2016). Fusion of a superfamily 1 helicase and an inactivated

DNA polymerase is a signature of common evolutionary history of *Polintons*, Polinton-like

viruses, *Tlr1* transposons and transpovirons. *Virus Evolution* 2(1). doi: 10.1093/ve/vew019

la Scola B, Desnues C, Pagnier I, Robert C, Barrassi L, Fournous G, Merchat M, Suzan-Monti M,

Forterre P, Koonin E and Raoult D. (2008). The virophage as a unique parasite of the giant

mimivirus. *Nature* 455(7209). doi: 10.1038/nature07218

Legendre M, Arslan D, Abergel C and Claverie JM. (2012). Genomics of Megavirus and the elusive

fourth domain of life. *Communicative and Integrative Biology* 5(1). doi:  10.4161/cib.18624

Legendre M, Bartoli J, Shmakova L, Jeudy S, Labadie K, Adrait A, Lescot M, Poirot O, Bertaux L,

Bruley C, Couté Y, Rivkina E, Abergel C and Claverie JM. (2014). Thirty-thousand-year-old

distant relative of giant icosahedral DNA viruses with a pandoravirus morphology. *Proc Natl*

*Acad Sci U S A* 111(11). doi: 10.1073/pnas.1320670111

Legendre M, Doutre G, Poirot O, Lescot M, Arslan D, Seltzer V, Bertaux L, Bruley C and Claverie J.

(2013). Pandoraviruses: amoeba viruses with genomes up to 2.5 Mb reaching that of parasitic

eukaryotes. *Science* 341. doi: 10.1126/science.1239181

Legendre M, Fabre E, Poirot O, Jeudy S, Lartigue A, Alempic JM, Beucher L, Philippe N, Bertaux L,

Christo-Foroux E, Labadie K, Couté Y, Abergel C and Claverie JM. (2018). Diversity and

evolution of the emerging Pandoraviridae family. *Nature Communications* 9(1). doi:

10.1038/s41467-018-04698-4

Legendre M, Lartigue A, Bertaux L, Jeudy S, Bartoli J, Lescot M, Alempic JM, Ramus C, Bruley C,

Labadie K, Shmakova L, Rivkina E, Couté Y, Abergel C and Claverie JM. (2015). In-depth study

of *Mollivirus sibericum*, a new 30,000-yold giant virus infecting *Acanthamoeba*. *Proc Natl Acad*

*Sci U S A* 112(38). doi: 10.1073/pnas.1510795112

Meinhardt F, Schaffrath R and Larsen M. (1997). Microbial linear plasmids. *Applied Microbiology and*

*Biotechnology*. doi: 10.1007/s002530050936

Menardo F, Loiseau C, Brites D, Coscolla M, Gygli SM, Rutaihwa LK, Trauner A, Beisel C, Borrell S

and Gagneux S. (2018). Treemmer: a tool to reduce large phylogenetic datasets with minimal

loss of diversity. *BMC Bioinformatics* 19(1). doi: 10.1186/s12859-018-2164-8

Moniruzzaman M, Martinez-Gutierrez CA, Weinheimer AR and Aylward FO. (2020). Dynamic

genome evolution and complex virocell metabolism of globally-distributed giant viruses. *Nature*

*Communications* 11(1). doi: 10.1038/s41467-020-15507-2

Mönttinen HAM, Bicep C, Williams TA and Hirt RP. (2021). The genomes of nucleocytoplasmic large

DNA viruses: viral evolution writ large. *Microbial Genomics*. doi: 10.1099/MGEN.0.000649

Mougari S, Chelkha N, Sahmi-Bounsiar D, di Pinto F, Colson P, Abrahao J and la Scola B. (2020). A

virophage cross-species infection through mutant selection represses giant virus propagation,

promoting host cell survival. *Communications Biology*. doi: 10.1038/s42003-020-0970-9

Nascimento, F.F., Reis, M.d. & Yang, Z. (2017). A biologist's guide to Bayesian phylogenetic analysis.

Nature Ecology and Evolution 1. doi: 10.1038/s41559-017-0280-x

Paez-Espino D, Zhou J, Roux S, Nayfach S, Pavlopoulos GA, Schulz F, McMahon KD, Walsh D,

Woyke T, Ivanova NN, Eloe-Fadrosh EA, Tringe SG and Kyrpides NC. (2019). Diversity,

evolution, and classification of virophages uncovered through global metagenomics. *Microbiome*

7(1). doi: 10.1186/s40168-019-0768-5

Patil S and Kondabagil K. (2021). Coevolutionary and phylogenetic analysis of mimiviral replication

machinery suggest the cellular origin of mimiviruses. *Molecular Biology and Evolution* 38(5).

doi: 10.1093/molbev/msab003

Porter SM. (2020). Insights into eukaryogenesis from the fossil record. *Interface Focus* 10(4). doi:

10.1098/rsfs.2019.0105

Posada D and Buckley TR. (2004). Model selection and model averaging in phylogenetics: advantages

of Akaike information criterion and Bayesian approaches over likelihood ratio tests. *Systematic*

*Biology*. doi: 10.1080/10635150490522304

Price MN, Dehal PS and Arkin AP. (2010). FastTree 2 - Approximately maximum-likelihood trees for

large alignments. *PLoS ONE* 5(3). doi: 10.1371/journal.pone.0009490

Pritham EJ, Putliwala T and Feschotte C. (2007). *Mavericks*, a novel class of giant transposable

elements widespread in eukaryotes and related to DNA viruses. *Gene* 390. doi:

10.1016/j.gene.2006.08.008

Rambaut A, Drummond AJ, Xie D, Baele G and Suchard MA. (2018). Posterior summarization in

Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* 67(5). doi: 10.1093/sysbio/syy032

Rice P, Longden L and Bleasby A. (2000). EMBOSS: The european molecular biology open software

suite. *Trends in Genetics*. doi: 10.1016/S0168-9525(00)02024-2

Ronquist F and Huelsenbeck JP. (2003). MrBayes 3: Bayesian phylogenetic inference under mixed

models. *Bioinformatics* 19(12). doi: 10.1093/bioinformatics/btg180

Roux S, Chan LK, Egan R, Malmstrom RR, McMahon KD and Sullivan MB. (2017). Ecogenomics of

virophages and their giant virus hosts assessed through time series metagenomics. *Nature*

*Communications* 8(1). doi: 10.1038/s41467-017-01086-2

Sayers EW, Beck J, Bolton EE, Bourexis D, Brister JR, Canese K, Comeau DC, Funk K, Kim S, Klimke

W, Marchler-Bauer A, Landrum M, Lathrop S, Lu Z, Madden TL, O'Leary N, Phan L, Rangwala

SH, Schneider VA, … Sherry ST. (2021). Database resources of the National Center for

Biotechnology Information. *Nucleic Acids Res* 49(D1). doi: 10.1093/nar/gkaa892

Scheid P. (2016). A strange endocytobiont revealed as largest virus. *Current Opinion in Microbiology*.

doi: 10.1016/j.mib.2016.02.005

Schulz F, Roux S, Paez-Espino D, Jungbluth S, Walsh DA, Denef VJ, McMahon KD, Konstantinidis

KT, Eloe-Fadrosh EA, Kyrpides NC and Woyke T. (2020). Giant virus diversity and host

interactions through global metagenomics. *Nature* 578(7795). doi: 10.1038/s41586-020-1957-x

Schmidt, HA and von Haeseler A. (2009). Phylogenetic inference using maximum likelihood methods.

The Phylogenetic Handbook: a practical approach to phylogenetic analysis and hypothesis

testing. 2nd Edition. Cambridge University Press, Cambridge, 181-209.

Söding J, Biegert A and Lupas AN. (2005). The HHpred interactive server for protein homology

detection and structure prediction. *Nucleic Acids Res*. doi: 10.1093/nar/gki408

Swofford DL. (1998). PAUP*. Phylogenetic Analysis Using Parsimony (*and other methods).. Sinauer

Associates.

Symonds MRE and Moussalli A. (2011). A brief guide to model selection, multimodel inference and

model averaging in behavioural ecology using Akaike's information criterion. *Behavioral*

*Ecology and Sociobiology*. doi: 10.1007/s00265-010-1037-6

Takemura M. (2016). Draft genome sequence of Tokyovirus, a member of the family *Marseilleviridae*

isolated from the Arakawa River of Tokyo, Japan. *Genome Announcements* 4(3). doi:

10.1128/genomeA.00429-16

van Rossum G, Drake FL, Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P,

Cournapeau D, Wieser E, Taylor J, Berg S, Smith NJ, Kern R, Picus M, Hoyer S, van Kerkwijk

MH, Brett M, Haldane A, del Río JF, … Oliphant TE. (2009). Python 3 Reference Manual.

Woo AC, Gaia M, Guglielmini J, da Cunha V and Forterre P. (2021). Phylogeny of the Varidnaviria

morphogenesis module: congruence and incongruence with the tree of life and viral taxonomy.

*Frontiers in Microbiology* 12. doi: 10.3389/fmicb.2021.704052

Yona AH, Alm EJ and Gore J. (2018). Random sequences rapidly evolve into de novo promoters.

*Nature Communications* 9(1). doi: 10.1038/s41467-018-04026-w

Yoshikawa G, Blanc-Mathieu R, Song C, Kayama Y, Mochizuki T, Murata K, Ogata H and Takemura

M. (2019). Medusavirus, a novel large dna virus discovered from hot spring water. *Journal of*

*Virology* 93(8). doi: 10.1128/jvi.02130-18

Yutin N and Koonin EV. (2012). Hidden evolutionary complexity of Nucleo-Cytoplasmic Large DNA

viruses of eukaryotes. *Virology Journal* 9. doi: 10.1186/1743-422X-9-161

Yutin N, Raoult D and Koonin EV. (2013). Virophages, *Polintons*, and transpovirons: A complex

evolutionary network of diverse selfish genetic elements with different reproduction strategies.

*Virology Journal* 10. doi: 10.1186/1743-422X-10-158

Yutin N, Shevchenko S, Kapitonov V, Krupovič M and Koonin EV. (2015). A novel group of diverse

Polinton-like viruses discovered by metagenome analysis. *BMC Biology* 13(1). doi:

10.1186/s12915-015-0207-4

838