Reference genome-independent taxonomic profiling of microbiomes with mOTUs3

Hans-Joachim Ruscheweyh[1,*]   hansr@ethz.ch

Alessio Milanese[1,2,*]   alessio.milanese@biol.ethz.ch

Lucas Paoli[1]   lucas.paoli@biol.ethz.ch

Nicolai Karcher[2]   nicolai.karcher@embl.de

Quentin Clayssen[1]   quentin.clayssen@gmail.com

Marisa Isabell Metzger[2]   marisa.metzger@embl.de

Jakob Wirbel[2]   jakob.wirbel@embl.de

Peer Bork[2,3,4]   bork@embl.de

Daniel R. Mende[5]   danielrmende@gmail.com

Georg Zeller[2#]   zeller@embl.de

Shinichi Sunagawa[1,#]   ssunagawa@ethz.ch

Contributed equally to this work

#Corresponding authors

[1]Department of Biology, Institute of Microbiology and Swiss Institute of Bioinformatics, ETH Zürich,

Zürich 8093, Switzerland

[2]Structural and Computational Biology Unit, European Molecular Biology Laboratory, 69117

Heidelberg, Germany

[3]Max Delbrück Centre for Molecular Medicine, Robert-Rössle-Str. 10, 13092, Berlin, Germany

[4]Department of Bioinformatics, Biocenter, University of Würzburg, Am Hubland, 97074, Würzburg,

Germany

[5]Department of Medical Microbiology, Amsterdam UMC, University of Amsterdam, Amsterdam, the

Netherlands

Abstract

Background: Taxonomic profiling is a fundamental task in microbiome research that aims to detect and quantify the relative abundance of microorganisms in biological samples. Available methods using shotgun metagenomic data generally depend on the availability of sequenced and taxonomically annotated reference genomes. However, the majority of microorganisms have not been cultured yet and lack such reference genomes. Thus, a substantial fraction of microbial community members remains unaccounted for during taxonomic profiling of metagenomes, particularly in samples from underexplored environments. To address this issue, we have developed the mOTU profiler, a tool that enables reference genome-independent species-level profiling of metagenomes. As such, it supports the identification and quantification of both "known" and "unknown" species based on a set of select marker genes.

Results: Here, we present mOTUs3, a command line tool that enables the profiling of metagenomes for >33,000 species-level operational taxonomic units. To achieve this, we leveraged the reconstruction and analysis of >600,000 draft genomes, most of which are metagenome assembled genomes (MAGs), from diverse microbiomes, including soil, freshwater systems, and the gastrointestinal tract of ruminants and other animals, which we found to be greatly underrepresented by reference genomes. Overall, two-thirds of all species-level taxa lacked a reference genome. The cumulative relative abundance of these newly included taxa was low in well-studied microbiomes, such as the human body sites (6-11%). By contrast, they accounted for substantial proportions (ocean, freshwater, soil: 43-63%) or even the vast majority (pig, fish, cattle: 60-80%) of the relative abundance across diverse non-human-associated microbiomes. Using community-developed benchmarks and datasets, we found mOTUs3 to be more accurate than other methods and to be more congruent with 16S rRNA gene-based methods for taxonomic profiling. Furthermore, we demonstrate that mOTUs3 greatly increases the resolution of well-known microbial groups into species-level taxa and helps identify new differentially abundant taxa in comparative metagenomic studies.

Conclusions: We developed mOTUs3 to enable accurate species-level profiling of metagenomes. Compared to other methods, it provides a more comprehensive view of prokaryotic community diversity, in particular for currently underexplored microbiomes. To facilitate comparative analyses by the research community, it is released with >11,000 precomputed profiles for publicly available metagenomes and is freely available at: https://github.com/motu-tool/mOTUs.

Keywords: Metagenomics, microbial community, benchmarking, taxonomic profiling, marker gene, metagenome-assembled genome, single-cell genome, reference genome

Background

Identifying and quantifying the abundance of taxa (i.e., taxonomic profiling) is a critical step in linking the composition of microbial communities to environmental functions and host health-related phenotypes [1,2]. Metagenomic sequencing of DNA directly extracted from an environmental or host-derived sample has enabled researchers to taxonomically profile microbial communities in an unbiased and cultivation-independent manner. The development of tools to generate accurate taxonomic profiles from metagenomic data has therefore become important to our understanding of microbial communities [3]. However, existing tools rely on the availability of informative sequences (such as k-mers or marker genes [4,5]), which are predominantly extracted from taxonomically annotated reference genomes (RefGs).

In recent years, high-throughput culturing of microorganisms coupled with RefG sequencing (known as culturomics) [6] has substantially expanded the proportion of microbial taxa with whole genome sequences in data repositories (e.g., NCBI RefSeq) benefitting taxonomic profiling tools. However, there is a strong bias toward microorganisms from well-studied habitats (e.g., human body sites) and/or those that can be readily cultivated using standard laboratory methods. Thus, most microbes on Earth remain uncultivated and lack a representative RefG [7,8], although they can be both globally prevalent [9] and numerically dominant in many environments [10, 11, 12, 13]. As a result, the incorporation of RefGs from newly isolated microbes into taxonomic profiling tools can be slow and disproportional across environments. This poses an additional challenge for accurate taxonomic profiling, given that microorganisms that remain undetected bias the abundance estimates of those that are detected [14,15].

To close the gap between the detectable and actual diversity present in microbial community samples, we developed mOTUs [14,16], a software tool that uses universal, protein-coding, single-copy phylogenetic marker gene (MG) sequences to quantify the taxonomic composition of microbial communities from metagenomic sequence data (for further applications, see also Ruscheweyh et al. 2021 [17]). As these MGs are present in all organisms, they can be identified not only in RefGs, but also in metagenomic assemblies. Conceptually, mOTUs is based on clustering sets of MGs

representing individual organisms by sequence similarity into species-level units. In the absence of a

generalizable species concept for prokaryotes [18,19], we refer to these units as MG-based

operational taxonomic units (abbreviated as 'mOTUs').

As an alternative to RefG sequencing, draft genomes are increasingly reconstructed by computational

binning of metagenomic assemblies into metagenome-assembled genomes (MAGs [20]) or by

sequencing amplified DNA from individual cells, resulting in single cell genomes (SAGs [21]). These

cultivation-independent methods have provided genomic access to microbial diversity in previously

underexplored environments. Here, in addition to MGs found in RefG and metagenomic data, we now

incorporate those found in MAGs and SAGs to more than double the number of taxa represented,

adding >20,000 new mOTUs compared to the previous major release [14]. Our evaluations show that

mOTUs3 outperforms other methods as assessed using metrics for taxonomic tool benchmarking

developed independently from our study [3,22]. Furthermore, we found mOTUs3 to provide an

unprecedented view of the species-level diversity within the most dominant heterotrophic bacterial

clade in the ocean and to greatly extend the number of detected and differentially abundant species in

cross-sectional studies, as exemplified in a comparison between rumen microbiomes of high- and

low-level methane-emitting sheep.

Results

Taxonomic profiling of diverse environments with mOTUs3

We developed mOTUs3 to facilitate the metagenomic profiling of 33,570 mOTUs, which is a 4.3-fold

increase compared to mOTUs2 (Figure 1a). Among all mOTUs, 35% were represented by a RefG

(n=11,915; ref-mOTUs), while an additional 21,655 were derived using MGs from either

metagenomic contigs (n=2,297; meta-mOTUs) or extended sources, such as MAGs (*de novo*-

assembled or imported) and a smaller number of SAGs and isolate genomes (n=19,358; ext-mOTUs),

to substantially extend the database coverage for reference genome-independent taxonomic profiling

of diverse environments. MGs not assigned to any mOTU were additionally added to the database and

merged into a single 'unassigned' group to improve the quantification accuracy of taxonomic profiles,

as previously demonstrated [14].

The newly established database allowed us to determine and systematically compare the fraction of

taxa currently not represented by RefGs in various environments. These environments include

extensively studied human-associated ones, for which metagenomic studies are complemented by

several culturomics efforts (e.g., Lagier et al. [23]). Furthermore, we included data from >20

environmental and animal-associated microbiomes (Supplementary Tables 1 and 2) that have been

primarily studied by metagenomic approaches. Overall, we found that more than half (11,882) of all

meta/ext-mOTUs (i.e., mOTUs not represented by any RefG) could not be assigned to any known

family (Supplementary Table 3; Methods), illustrating the taxonomic novelty covered by mOTUs3.

The distribution of the newly included data into ref/meta/ext-mOTUs was highly variable across the

different environments (Supplementary Figure 1). As expected, 97% of the ~400,000 MAGs from

human microbiome samples (Supplementary Table 1) had already been represented by 2,360 pre-

existing (i.e., ref/meta-)mOTUs (Supplementary Table 4). Notably, the remaining 3% represented

2,750 new ext-mOTUs, showing that novel species can still be uncovered by studying

underrepresented populations, dietary habits and/or disease states [24,25]. By contrast, we found that

only ~25% of the 6,479 MAGs from mouse gut metagenomes (Supplementary Table 1) corresponded

to pre-existing mOTUs (n=72), despite ongoing cultivation efforts [6]; the remaining 75% were

grouped into 587 ext-mOTUs (Supplementary Table 4). However, the vast majority of ext-mOTUs

(n=16,021) resulted from the inclusion of other animal-associated (e.g., ruminants, fish, chicken, pig,

bee, dog, cat) and environmental (e.g., soil, freshwater, wastewater, ocean, air) microbiomes

(Supplementary Table 1) for which the generation of representative RefGs is lagging.

We used mOTUs3 to profile 10,541 available shotgun metagenomic data sets across the 23

environments covered by its database (Supplementary Table 1). For comparative analyses, we subset

the data to 5,756 high-quality samples (Methods; Supplementary Table 5) from 16 environments and

found the overall number of detected mOTUs to range from 247 (honey bee) to >6,000 (ocean,

wastewater and cattle microbiomes). To illustrate the proportion of quantifying taxa currently not

represented by RefGs (Figure 1b), we summarized the cumulative relative abundances of unassigned

taxa and the different types of mOTUs (ref-mOTUs, meta-mOTUs, ext-mOTUs). The fraction of

unassigned taxa was highest for soil samples (33%; s.d. 8%), which reflects the high microbial

diversity in soil as well as challenges in reconstructing genomes from this environment [26]. By

contrast, more than 87% (s.d. 0.7%) of the relative abundance was represented by ref-mOTUs in

human skin samples mainly due to the dominance of few taxa with cultivated representatives [27].

Similarly, the fraction of relative abundance assigned to ext-mOTUs varied considerably between

environments: on average, only ~6% of the bacterial abundance in human-associated samples was

assigned to newly added taxa, while this fraction was as high as ~80% in cattle rumen microbiomes.

Comparison with other taxonomic profilers

As in other fields of bioinformatics, there is broad consensus that the performance of analysis tools

needs to be carefully evaluated. However, best practices (e.g., balancing precision and recall,

selecting criteria for 'best' performance) are often debated [28,29], and in microbiome research, an

agreement on some fundamental concepts (e.g., sequence vs. taxonomic abundance, representation of

unknown taxa in ground truth data) is still lacking [30,31]. In an attempt to address some of these

issues in a community-driven effort, modeled after successful examples in other fields [32,33], the

Critical Assessment of Metagenome Interpretation (CAMI) has provided curated ground truth datasets

along with a tool (OPAL) to reproducibly evaluate metagenomic analysis tools [3,22].

Using the latest CAMI datasets with disclosed results [34], we compared mOTUs3 to its prior major

release version (mOTUs2) [14] and other selected metagenomic profiling tools (MetaPhlAn3 [5] and

Bracken [4,35], Methods) representing conceptually different, well-performing approaches to

taxonomic profiling [30]. Using the OPAL tool for scoring and evaluation, we first evaluated

presence/absence (F_1-score) and relative abundance predictions (L1 norm error) at the species level.

For the different datasets, which represented samples from five human body sites and the mouse gut

microbiome, mOTUs3 and MetaPhlAn3 performed generally better than Bracken and mOTUs2

(Figure 2a/b). At higher taxonomic ranks, mOTUs3 had similar or higher scores than the other tools.

For some datasets, taxonomic ranks and tools, there was little to no room for improvements of the F_1-

score or L1 norm error. This may be due to the simulated datasets being mainly based on taxa for

which RefGs are available and/or result from incongruencies of taxonomic annotations used by the

different profilers compared to the ground truth. In addition to the L1 norm error, OPAL computes

additional metrics for profiling quality (completeness, purity, weighted UniFrac error) and

summarizes them across taxonomic ranks into a composite score. Based on this evaluation criterion,

mOTUs3 outperformed the other tools (Figure 2c), as well as additional tools assessed in the CAMI

challenge (Methods; Supplementary Figure 2).

In the absence of independent ground truth data sets to benchmark taxonomic profiling tools for less

well-studied environments, we correlated taxonomic profiles obtained by mOTUs3 and other tools to

those obtained by analyzing 16S rRNA gene (16S) fragments. This approach leverages both the

availability of comprehensive 16S databases for taxonomic classification [36] and the possibility of

estimating taxonomic abundances based on 16S-based data from metagenomes [37]. Briefly, we

extracted 16S fragments from the same datasets we used for metagenomic profiling and generated

relative abundance profiles for them (Methods). To ensure comparability between 16S and

metagenomic profiles, the analysis was performed at the genus and higher taxonomic ranks (for

discussion, see Salazar et al. [37]). We found that mOTUs3 had consistently higher correlations with

16S profiles than the other tools across all environments, except for the human gut for which

MetaPhlAn3 showed correlation coefficients similar to those of mOTUs3 (Figure 3).

Resolving the diversity of Pelagibacterales with mOTUs3

In addition to the broader taxonomic coverage by mOTUs3 across environments, we sought to

investigate the capability of mOTUs3 to resolve microbial clades into more fine-grained taxonomic

units. To this end, we focused on Pelagibacterales (also referred to as the SAR11 clade), which is the

most abundant heterotrophic bacterial group in the global oceans [38]. Members of the

Pelagibacterales have previously been shown to display high genomic variability while maintaining

highly conserved 16S sequences [39]. This prompted us to evaluate the species-level resolution of

mOTUs3 and to compare the diversity represented by mOTUs to the diversity represented by

operational taxonomic units (OTUs) defined by 16S sequence similarity.

For this analysis, we selected from all mOTUs annotated as Pelagibacterales (n=1,029; 2,063

genomes) those that were represented by genomes with complete 16S sequences (n=602; 1,105

genomes). The number of mOTUs was comparable to the number resulting from a 95% average

nucleotide identity (ANI)-based clustering of the 1,105 genome sequences into species-level groups

(n=700; Figure 4a), which is common practice in the field of microbial phylogenomics [7,40].

Moreover, we found sequence identities of mOTUs-representing MGs to linearly correlate with those

of whole genomes across the whole range of observed values (r^2=0.71; Figure 4b). By contrast, 16S

sequence-based OTUs using a 97% or 99% sequence similarity cutoff resulted in a 31.7-fold (n=19)

or 5.8-fold (n=104) lower number of taxonomic units, respectively, compared to mOTUs (Figure 4a).

This discrepancy is also reflected by a weaker correlation (r^2=0.45; Figure 4b) of identities between

16S sequences and corresponding whole genome sequences. The minimum 16S identities were ca.

87% and started saturating at approximately 97% at which point genome identities were still as low as

~70-80% (Figure 4b). Similar findings were reported previously albeit on smaller datasets [39].

Finally, comparing the grouping of genomes by mOTUs and ANI into species-level clusters, we

found almost perfect congruence (Figure 4c, Methods).

Differential abundance of novel archaea in low/high methane-emitting sheep rumen metagenomes

High-resolution taxonomic profiling of metagenomes from underexplored environments can be

achieved by custom-made marker gene or genome databases selected for the microbial community

under study [12,41]. However, this approach is often labor- and resource-intensive and requires

specialized expertise, and its results cannot easily be compared across studies and communities. To

demonstrate the utility of mOTUs3 to address these challenges, we reanalyzed rumen metagenomes

from high- and low-methane emitting (HME and LME) sheep [41]. Importantly, these data were not

used for the database construction of mOTUs3.

Based on mOTUs3 taxonomic profiles, we identified 131 microbial species that differed significantly

in abundance between HME and LME samples and showed an at least tenfold increase or decrease in

relative abundance (corresponding to a generalized fold change of >= 1 [42]). Among these

differentially abundant species, 92% were represented by ext-mOTUs. These were therefore not

expected to be detectable by reference-based profilers. To test this, we applied the same workflow

using MetaPhlAn3 and Bracken (see Methods), which yielded only 10 and 30 differentially abundant

species for the respective tools (Figure 5a).

Given the metabolic importance of methanogenic archaea in ruminants as well as previous evidence

of uncharted archaeal diversity in the sheep rumen [12], we further investigated the species-level

diversity of known and unknown archaeal species. To this end, we reconstructed a phylogenetic tree

of the archaeal mOTUs detected in the sheep rumen metagenomes (n=15) and contextualized them

with reference genomes from members of the genera *Methanobrevibacter* and *Methanosphaera*

(Figure 5b). This analysis revealed that all six differentially abundant archaea in the sheep rumen

corresponded to ext-mOTUs. Two of them, which were significantly more abundant in high-methane

emitters, were most closely related to *Methanobrevibacter gottschalkii*, which itself was not detected.

Notably, the MG sequence similarity between these ext-mOTUs and *M. gottschalkii* was <85%

(Figure 5b), which is well below the species-level cutoff of 96.5% used by mOTUs [16] and therefore

suggests that these ext-mOTUs represent novel *Methanobrevibacter* spp.

Discussion

With mOTUs3, we have developed a taxonomic profiler that combines state-of-the-art accuracy, as

demonstrated in competitive benchmarks based on simulated datasets, with an innovative database

construction approach to detect and quantify underrepresented microbes from diverse environments at

high (i.e., species-level) taxonomic resolution. The ability to incorporate MG sequences from any

MAG and SAG to generate mOTUs *de novo* and independently from the availability of RefGs and/or

prior existence of taxonomic annotations (such as NCBI or GTDB species names) will allow users to

continuously extend the core database of mOTUs to represent microbial diversity from newly

explored microbiomes. Such future extensions could also target eukaryotic microorganisms, as these

are an integral part of many microbial communities, but are not well represented in databases of

existing taxonomic profiling tools.

However, the flexibility in defining operational taxonomic units *de novo* comes with a need for

taxonomic annotation, as is also the case for 16S rRNA-based *de novo* clustered OTUs. Despite the

calibration of MG sequence identity cutoffs to maximize congruence with the NCBI taxonomy [16],

this procedure can lead to conflicts with existing taxonomies. Irrespective of the ongoing debate on

whether prokaryotic species should be consistent with genomic similarity-based criteria, delineating

species by sequence identity puts mOTUs at a disadvantage in benchmarks, such as CAMI, which

rely on rigid matching of taxonomic labels. The high performance of mOTUs [34] despite this

disadvantage is likely due to the higher number of quantified taxa and the resulting reduction in

compositionality-related biases.

Conclusions

The present work introduces mOTUs3 as a reference-genome independent tool that allows for

charting the taxonomic landscape of many environments at species-level resolution. Its independence

from taxonomically annotated reference genomes, makes it generally applicable also beyond well-

studied environments to quantify and reveal yet uncharacterized microbial species of potential

biological relevance. To support the research community, mOTUs3 is documented and available as

open source software at https://github.com/motu-tool/mOTUs.

Methods

Collection and processing of data to compile the mOTUs3 database

To extend the taxonomic coverage of the mOTUs3 database, 4,531 publicly available metagenomic

datasets from 23 environments (Supplementary Table 1) were processed to generate 150,880 MAGs

as previously described [43]. Briefly, BBMap (v.38.71) was used to quality control sequencing reads

from all samples by removing adapters from the reads, removing reads that mapped to quality control

sequences (PhiX genome) and discarding low-quality reads (*trimq=14, maq=20, maxns=1* and

minlength=45). For metagenomic data of human origin, human genome-derived reads were removed

using the masked human reference genome provided by BBMap. Quality-controlled reads were

merged using bbmerge.sh with a minimum overlap of 16 bases, resulting in merged, unmerged paired

and single reads. The reads were assembled into scaffolded contigs (hereafter scaffolds) using the

SPAdes assembler (v3.14 or v3.12) [44] in metagenomic mode. Genes were predicted on length-

filtered (≥ 500 bp) scaffolded contigs (hereafter scaffolds) using Prodigal (v2.6.3) [45]. Universal

single-copy phylogenetic marker genes (MGs) were extracted using fetchMGs (v1.2; *-m extraction*)

[16].

Scaffolds were length-filtered (≥ 1000 bp) and within each study, quality-controlled reads from each

sample were mapped against the scaffolds of each sample. Mapping was performed using BWA

(v0.7.17-r1188; *-a*) [46]. Alignments were filtered to be at least 45 bp in length, with an identity of ≥

97% and a coverage of ≥ 80% of the read sequence. The resulting BAM files were processed using

the *jgi_summarize_bam_contig_depths* script of MetaBAT2 (v2.12.1) [20] to compute within- and

between-sample coverages for each scaffold. The scaffolds were binned by running MetaBAT2 on all

samples individually (*--minContig 2000* and *--maxEdges 500* for increased sensitivity). These

metagenomic bins were complemented with 454,773 external draft genomes (~96% MAGs; ~4%

isolate and single-cell genomes) from previous work (Supplementary Table 1). Complete genes in

external draft genomes and metagenomic bins were predicted using Prodigal (v2.6.3; *-c -m -g 11 -p*

single) and MGs were extracted using fetchMGs (v1.2) *(-m extraction -v -i)*.

Metagenomic bins and draft genomes were annotated with Anvio (v5.5.0) [47], quality controlled

using the CheckM (v1.0.13) [48] lineage workflow (completeness ≥ 50% and contamination < 10%)

and filtered for genomes containing at least six out of the 10 MGs used by mOTUs [16] to produce

the dataset of MGs from a total of 499,512 *de novo*-generated MAGs (i.e., quality-controlled

metagenomic bins) and external draft genomes used for the construction of the mOTUs3 database.

Construction of the mOTUs3 database

MGs from 499,512 genomes were mapped against the latest mOTUs database (v2.5.1), which was an

update of version 2.0 to account for a more recent release of the progenomes2 database [49] (Figure

1a) using vsearch [50] (v2.14.1; *--usearch_global --strand both --id 0.8 --maxaccepts 10000 --*

maxrejects 10000). MGs from a total of 283,250 and 136,429 genomes were assigned to existing ref-

mOTUs and meta-mOTUs, respectively. These genomes were removed since they were already

represented. The remaining 79,833 genomes resulted in an extension of the mOTUs database by

19,358 new mOTUs (ext-mOTUs). For consistency with the taxonomic annotation of ref-mOTUs,

ext-mOTUs were annotated using the STAG classifier (https://github.com/zellerlab/stag, version 0.7;

default parameters) trained on genomes in the proGenomes2 database [49] (NCBI taxonomy, version:

8 January 2019). MGs identified on scaffolds that were not binned into MAGs were used to update

the 'unassigned' mOTU, which contain unbinned MGs that are used to estimate the quantity of

unknown species, by aligning these MGs against the extended database using vsearch (v2.14.1;

usearch_global --maxaccepts 1000 --maxrejects 1000 --strand both). MGs that did not align within

MG-specific cutoffs [51] were clustered using vsearch (v2.14.1; *--cluster_fast*) using MG-specific

cutoffs and the representative sequence was added to the unassigned mOTU.

Computation of mOTUs3 profiles for comparative analyses

A total of 11,164 metagenomic and metatranscriptomic samples (Supplementary Table 1,

Supplementary Table 2) were quality controlled and merged as described above and profiled with

mOTUs3 using default parameters and the *-c* option to build a community resource of taxonomic

profiles. For comparative analyses across environments, 5,756 of these samples were used after

removing all (n=623) metatranscriptomic samples, metagenomic samples from environments with too

few samples (termite, panda, aerosols and bioreactor) or from studies comprising samples from

different environments and samples with less than 5,000 mapped inserts. To calculate the total

number of detected mOTUs for a given environment, we counted the number of mOTUs with a

prevalence greater than 0.1% (Supplementary Table 5). To compare the median number of detected

mOTUs across different environments, we downsampled the insert counts to 5,000 using the *rrarefy*

function of the vegan package [52].

Comparison of taxonomic profilers using the CAMI framework

The performance of mOTUs3 was evaluated and compared to mOTUs2 and other taxonomic profilers

by analyzing 113 publicly available samples (49 human-associated, 63 mouse gut metagenomes)

provided by the second CAMI challenge (https://cami-challenge.org/participate). The samples were

profiled with mOTUs3 (v3.0.1; *-C precision*), mOTUs2 (v2.1.1; *-C precision*), MetaPhlAn3 (v3.0.7; *-*

-CAMI_format_output --index mpa_v30_CHOCOPhlAn_201901) [5] and Kraken/Bracken (v2.1.2; *--*

db=k2_standard_20201202 --paired / v2.6.1; --db=k2_standard_20201202 -r 100 -l S|G|F|O|C|P|D)

[4,35]. Kraken/Bracken reports were further translated into the CAMI format ed files using the

tocami.py script provided at https://github.com/hzi-bifo/cami2_pipelines. For comparative analyses,

the OPAL framework (v1.0.9) [22] was used with default parameters providing the gold standard with

the parameter *--gold_standard_file,* the names of the tools with *--labels*, the description with *-d*, the

output with *--output_dir* and the taxonomic profiles files as positional arguments.

Comparison of metagenomic profiles with 16S rRNA gene-based profiles

The 16S rRNA-based taxonomic profiler mTAGs [37] (v1.0.1; *-ma 1000 -mr 1000*) was used to

generate relative abundance profiles for metagenomic samples (Supplementary Table 1). The output

of mTAGs was mapped to the NCBI taxonomy to facilitate comparative analysis. The same samples

were profiled with MetaPhlAn3 (v3.0.7; *--index mpa_v30_CHOCOPhlAn_201901*) and

Kraken/Bracken (v2.1.2; *--db=k2_standard_20201202 --paired / v2.6.1; --*

db=k2_standard_20201202 -r 100 -l S). Samples with small read/insert coverages (mTAGs<10,000,

mOTUs<1,000, Kraken/Bracken<10,000, no filtering was done on MetaPhlAn3 as profiles contain

relative abundances) were removed, leaving 6,119 samples for comparative analysis. Spearman

correlations were calculated for each taxonomic rank based on concatenated relative abundances

between mTAGs and the metagenomic profiling tools.

Comparison of Pelagibacterales genome clusters with marker gene and 16S rRNA gene sequences

Out of 2,063 genomes belonging to 1,029 mOTUs annotated as Pelagibacterales, 1,105 genomes

(from 602 mOTUs) that contained a complete copy of the 16S rRNA gene were selected. These

genomes were also clustered based on average nucleotide identity using dRep [53] (v2.5.4; *-comp 0 -*

con 1000 -sa 0.95 -nc 0.2) using a 95% cutoff as part of the OMD [43]. In addition, these genomes

were clustered based on their 16S rRNA gene identity (99% and 97%) using vsearch [50] (v2.14.1; *--*

cluster_smallmem --id 0.97 / 0.99). The consistency between the different clustering approaches was

evaluated using the V-measure, which combines both the homogeneity and completeness metrics

[54].

To correlate distances of the 1,105 genomes between the different clustering techniques we performed

exhaustive distance calculations at the whole-genome level, the 10 MGs used by mOTUs and the 16S

rRNA gene. Whole genome distances were computed using MASH [55] as implemented in dRep

(v2.5.4). MG- and 16S rRNA gene-based distances were computed using vsearch (v2.14.1; --

allpairs_global --id 0.0) and MG distances were averaged across the 10 genes prior to computing

correlations.

Differential abundance of mOTUs between low/high methane-emitting sheep

Samples from sheep rumen metagenomes (n=16) [41] were profiled with mOTUs3 (v3.0.1; *-c*),

MetaPhlAn3 (v3.0.7; *--index mpa_v30_CHOCOPhlAn_201901*) and Kraken/Bracken (v2.1.2; --

db=k2_standard_20201202 --paired / v2.6.1; *--db=k2_standard_20201202 -r 100 -l S*). To test for

differentially abundant species between low methane emitters (LMEs) and high methane emitters

(HMEs), the respective profiles were analyzed using SIAMCAT default workflows [42]. This

workflow includes filtering of species/mOTUs with a relative abundance of >0.1% in at least one

sample [42]. Wilcoxon test results were corrected for multiple testing using the Benjamini–Hochberg

method [56] at 5% FDR. The reported effect size measure is the generalized fold change (gFC),

calculated as the log10 of the geometric mean of quantile differences between groups as defined in

SIAMCAT [42].

A phylogeny was constructed for all archaeal mOTUs belonging to the *Methanobrevibacter* and

Methanosphaera genera or the *Thermoplasmata* class that passed the relative abundance filtering (14

ext-mOTUs, 1 ref-mOTU) together with ref-mOTUs from *Methanobrevibacter* and *Methanosphaera*

(n=15) and a randomly selected Thermoplasmata ref-mOTU as an outgroup. Representative genomes

from these 31 mOTUs were selected either by picking the centroid genome (for ext-mOTUs) or the

reference genome (for ref-mOTUs). Marker genes were individually aligned (*mafft* [57], v7.458), the

alignments were concatenated and a maximum-likelihood phylogeny was calculated using RAxML

[58] (v8.2.12; *raxmlHPC -p 12345 -m PROTGAMMAAUTO*). The distance between the 14 ext-

mOTUs and their closest ref-mOTU was calculated based on averaged marker gene distances across

the 10 genes (v2.14.1; *vsearch --allpairs_global --id 0.0*).

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The mOTUs3 software is documented and publicly available as open source software (GPL 3) at

https://github.com/motu-tool/mOTUs. The updated mOTUs3 database can be found at Zenodo

(https://doi.org/10.5281/zenodo.5140350) and contains all MGs used in this study and the public

profiles generated with mOTUs3. A complete list with all sequencing samples used for building the

database and/or for profiling can be found in Supplemental Tables 1 and 2.

Competing interests

none declared

Funding

We acknowledge funding from the Swiss National Foundation through project grants

205321_184955, the NCCR Microbiomes (51NF40_180575) and the Strategic Focal Area

"Personalized Health and Related Technologies (PHRT)" of the ETH Domain (#2018-521) to SS, and

the European Molecular Biology Laboratory and the German Federal Ministry of Education and

Research (BMBF, the de.NBI network, grant no. 031A537B and grant no. 031L0181A) to GZ.

Authors' contributions

GZ and SS conceived and supervised the work. HJR and AM developed code, generated the database with support from DRM, and performed the benchmark analysis. LP and NK performed the taxonomic diversity analysis of the SAR11 clade and the comparative metagenomic analysis, respectively. QC supported the collection and processing of data. MIM and JW contributed to the taxonomic annotation of mOTUs. HJR, AM, LP, NK, PB, DRM, GZ and SS wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgments

We would like to thank the ETH IT Services and HPC facilities for granting access to the EULER high performance cluster. We also thank Thea Van Rossum for her input on the taxonomic annotation of mOTUs and the users of mOTUs for their feedback and continuous support.

Authors' information

Hans-Joachim Ruscheweyh and Alessio Milanese contributed equally to this work.

a

mOTUs2
5,232 ref-mOTUs   **2,494** meta-mOTUs

87,920 Reference genomes

150,880 Newly constructed MAGs

454,773 External MAGs/SAGs/isolates

Quality control

mOTUs2.5.1
11,915 ref-mOTUs   **2,297** meta-mOTUs

499,512 Genomes

283,250

136,429

79,833

11,915 ref-mOTUs   **2,297** meta-mOTUs   **19,358** ext-mOTUs

mOTUs3 (33,570 mOTUs)

b

Human skin (1043 / 49)

Human vagina (644 / 56)

Human gut (4219 / 227)

Human oral (1439 / 212)

Cat (473 / 186)

Dog (463 / 116)

Bee (247 / 41)

Ocean (6970 / 742)

Wastewater (6290 / 505)

Freshwater (3515 / 258)

Pig (2740 / 555)

Soil (1979 / 470)

Chicken (542 / 210)

Mouse (974 / 164)

Fish (857 / 158)

Cattle (6108 / 733)

Rel. abundance
- ref-mOTUs
- meta-mOTUs
- unassigned
- ext-mOTUs

Figure 1. The mOTUs3 database enables species-level profiling across diverse environments.

(a) The database of the previous major release of mOTUs (version 2)[14] was updated to version 2.5

to account for the current release of the progenomes2 database[49]. Based on version 2.5, the

mOTUs3 database was constructed by adding universal, single-copy phylogenetic marker genes

(MGs) from 605,653 genomes (metagenome-assembled genomes (MAGs) and a smaller number of

isolate and single amplified genomes (SAGs)). This addition resulted in the extension of the database

by 19,358 new species-level, MG-based operational taxonomic units (ext-mOTUs). Genomes already

represented by ref- and meta-mOTUs in version 2.5 were not added (gray lines). **(b)** Breakdown by

the three types of mOTUs shows that mOTUs3 enables the reference genome-independent profiling

of a substantial fraction of microbial diversity across different environments. The numbers below the

ring charts represent the total number of mOTUs that were detected per environment (left)

considering only species with a prevalence of 0.1% and the median number of mOTUs per sample

that were detected after downsampling to 5,000 inserts (right).

Figure 2. Comparison of mOTUs to other taxonomic profilers.

The performance of mOTUs3 was compared to other taxonomic profiling tools based on the dataset

from the second Critical Assessment of Metagenome Interpretation (CAMI) challenge (see Methods).

The F1 score **(a)** and L1 norm error **(b)** are shown as reported by the OPAL tool[22] for each

taxonomic rank (x-axis). High L1 norm error values at the family and genus levels of GI samples

mostly derive from an updated taxonomy of the highly abundant Oscillospiraceae (previously

Ruminococcaceae)[59]. **(c)** Each method was ranked across all samples and for each taxonomic rank

using four measures (completeness, purity, L1 norm error and weighted UniFrac error), and the

OPAL sum of scores was calculated as a sum of these ranks (lower rank indicates better

performance). OR: oral cavity, SK: skin, AI: airways, UT: urogenital tract, GI: gastrointestinal tract,

MG: mouse gut.

Figure 3. Comparison of metagenomic profiling tools using 16S rRNA-based taxonomic profiles.

Spearman correlations between relative abundances generated by different metagenomic profiling

tools and 16S rRNA gene-based profiles from the same samples. The correlations were calculated at

different taxonomic ranks (x-axis; c: class, o: order, f: family, g: genus) and showed that mOTUs3

generally had the highest values for the different body sites tested, except for human gut samples with

similar values for mOTUs3 and MetaPhlAn3.

Figure 4. Species-level diversity of Pelagibacterales as resolved by mOTUs3.

(a) The number of taxonomic units within the Pelagibacterales order varies depending on the

clustering method used, which was based on using marker gene (MG) sequences (used by mOTUs),

Average Nucleotide Identity (ANI) of whole genomes, and full length 16S rRNA gene sequences. (b)

mOTUs marker gene distances better capture whole genome distances compared to full length 16S,

explaining the patterns observed in (a). In particular, 16S rRNA gene sequence identity saturates

while whole genome similarity can be as low as 70-80%. (c) The different clustering approaches vary

in their agreement with each other as determined by the V-measure, which captures both the

completeness and homogeneity of the clusterings. The highest agreement was found between mOTUs

and with whole genome clustering by ANI.

Figure 5. Detection of differentially abundant taxa in low/high-level methane-emiting sheep

rumen microbiomes.

(a) A comparison between metagenomic profilers shows that mOTUs3 detected 131 differentially

abundant species (*q*-value <0.05 and an absolute generalized fold change > 1; indicated by dotted

lines) between low- and high-level methane-emitting sheep, while MetaPhlAn3 and Bracken detected

nine and two species, respectively. Most of the species detected by mOTUs were represented by ext-

mOTUs only, demonstrating the added value of reference genome-independent profiling enabled by

mOTUs3. **(b)** Archaeal mOTUs present in the sheep rumen microbiome (highlighted in gray) were

phylogenetically contextualized with *Methanobrevibacter* spp. and *Methanosphaera* spp. represented

by ref-mOTUs. All differentially abundant ext-mOTUs (middle panel) correspond to distinct yet

undescribed *Methanobrevibacter* spp. as supported by MG sequence identities (right panel) to the

closest known species being below the species-level cutoff of 96.5% (dotted vertical line).

References

1. Fuhrman JA. Microbial community structure and its functional implications. Nature. Nature

Publishing Group; 2009;459:193–9.

2. Wang J, Jia H. Metagenome-wide association studies: fine-mining the microbiome. Nat Rev

Microbiol. 2016;14:508–22.

3. Sczyrba A, Hofmann P, Belmann P, Koslicki D, Janssen S, Dröge J, et al. Critical Assessment of

Metagenome Interpretation—a benchmark of metagenomics software. Nat Methods. Nature

Publishing Group; 2017;14:1063–71.

4. Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence classification using exact

alignments. Genome Biol. BioMed Central; 2014;15:1–12.

5. Beghini F, McIver LJ, Blanco-Míguez A, Dubois L, Asnicar F, Maharjan S, et al. Integrating

taxonomic, functional, and strain-level profiling of diverse microbial communities with bioBakery 3.

Elife [Internet]. 2021;10. Available from: http://dx.doi.org/10.7554/eLife.65088

6. Lagkouvardos I, Pukall R, Abt B, Foesel BU, Meier-Kolthoff JP, Kumar N, et al. The Mouse

Intestinal Bacterial Collection (miBC) provides host-specific insight into cultured diversity and

functional potential of the gut microbiota. Nat Microbiol. 2016;1:16131.

7. Konstantinidis KT, Rosselló-Móra R. Classifying the uncultivated microbial majority: A place for

metagenomic data in the Candidatus proposal. Syst Appl Microbiol. 2015;38:223–30.

8. Lloyd-Price J, Mahurkar A, Rahnavard G, Crabtree J, Orvis J, Hall AB, et al. Strains, functions and

dynamics in the expanded Human Microbiome Project. Nature. 2017;550:61–6.

9. Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, et al. A new view of the

tree of life. Nat Microbiol. 2016;1:16048.

10. Salazar G, Paoli L, Alberti A, Huerta-Cepas J, Ruscheweyh H-J, Cuenca M, et al. Gene

Expression Changes and Community Turnover Differentially Shape the Global Ocean

Metatranscriptome. Cell. 2019;179:1068–83.e21.

11. Lesker TR, Durairaj AC, Gálvez EJC, Lagkouvardos I, Baines JF, Clavel T, et al. An Integrated

Metagenome Catalog Reveals New Insights into the Murine Gut Microbiome. Cell Rep.

2020;30:2909–22.e6.

12. Stewart RD, Auffret MD, Warr A, Walker AW, Roehe R, Watson M. Compendium of 4,941

rumen metagenome-assembled genomes for rumen microbiome biology and enzyme discovery. Nat

Biotechnol. Nature Publishing Group; 2019;37:953–61.

13. Wilhelm RC, Cardenas E, Leung H, Maas K, Hartmann M, Hahn A, et al. A metagenomic survey

of forest soil microbial communities more than a decade after timber harvesting. Sci Data.

2017;4:170092.

14. Milanese A, Mende DR, Paoli L, Salazar G, Ruscheweyh H-J, Cuenca M, et al. Microbial

abundance, activity and population genomic profiling with mOTUs2. Nat Commun. 2019;10:1014.

15. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome Datasets Are

Compositional: And This Is Not Optional. Front Microbiol. 2017;8:2224.

16. Sunagawa S, Mende DR, Zeller G, Izquierdo-Carrasco F, Berger SA, Kultima JR, et al.

Metagenomic species profiling using universal phylogenetic marker genes. Nat Methods.

2013;10:1196–9.

17. Ruscheweyh H-J, Milanese A, Paoli L, Sintsova A, Mende DR, Zeller G, et al. mOTUs: Profiling

Taxonomic Composition, Transcriptional Activity and Strain Populations of Microbial Communities.

Curr Protoc. 2021;1:e218.

18. Rosselló-Mora R, Amann R. The species concept for prokaryotes. FEMS Microbiol Rev.

2001;25:39–67.

19. Staley JT. The bacterial species dilemma and the genomic-phylogenetic species concept. Philos

Trans R Soc Lond B Biol Sci. 2006;361:1899–909.

20. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive binning

algorithm for robust and efficient genome reconstruction from metagenome assemblies. PeerJ.

2019;7:e7359.

21. Woyke T, Doud DFR, Schulz F. The trajectory of microbial single-cell sequencing. Nat Methods.

2017;14:1045–54.

22. Meyer F, Bremges A, Belmann P, Janssen S, McHardy AC, Koslicki D. Assessing taxonomic

metagenome profilers with OPAL. Genome Biol. 2019;20:51.

23. Lagier J-C, Khelaifia S, Alou MT, Ndongo S, Dione N, Hugon P, et al. Culture of previously

uncultured members of the human gut microbiota by culturomics. Nat Microbiol. 2016;1:16203.

24. Wirbel J, Pyl PT, Kartal E, Zych K, Kashani A, Milanese A, et al. Meta-analysis of fecal

metagenomes reveals global microbial signatures that are specific for colorectal cancer. Nat Med.

2019;25:679–89.

25. Pasolli E, Asnicar F, Manara S, Zolfo M, Karcher N, Armanini F, et al. Extensive Unexplored

Human Microbiome Diversity Revealed by Over 150,000 Genomes from Metagenomes Spanning

Age, Geography, and Lifestyle. Cell. 2019;176:649–62.e20.

26. Alteio LV, Schulz F, Seshadri R, Varghese N, Rodriguez-Reillo W, Ryan E, et al. Complementary

Metagenomic Approaches Improve Reconstruction of Microbial Diversity in a Forest Soil. mSystems

[Internet]. 2020;5. Available from: http://dx.doi.org/10.1128/mSystems.00768-19

27. Byrd AL, Belkaid Y, Segre JA. The human skin microbiome. Nat Rev Microbiol. 2018;16:143–

55.

28. Buchka S, Hapfelmeier A, Gardner PP, Wilson R, Boulesteix A-L. On the optimistic performance

evaluation of newly introduced bioinformatic methods. Genome Biol. 2021;22:152.

29. Marx V. Bench pressing with genomics benchmarkers. Nat Methods. 2020;17:255–8.

30. Sun Z, Huang S, Zhang M, Zhu Q, Haiminen N, Carrieri AP, et al. Challenges in benchmarking

metagenomic profilers. Nat Methods. 2021;18:618–26.

31. Ye SH, Siddle KJ, Park DJ, Sabeti PC. Benchmarking Metagenomics Tools for Taxonomic

Classification. Cell. 2019;178:779–94.

32. Stolovitzky G, Monroe D, Califano A. Dialogue on reverse-engineering assessment and methods:

the DREAM of high-throughput pathway inference. Ann N Y Acad Sci. 2007;1115:1–22.

33. Moult J, Pedersen JT, Judson R, Fidelis K. A large-scale experiment to assess protein structure

prediction methods. Proteins. 1995;23:ii – v.

34. Meyer F, Lesker T-R, Koslicki D, Fritz A, Gurevich A, Darling AE, et al. Tutorial: assessing

metagenomics software with the CAMI benchmarking toolkit. Nat Protoc. 2021;16:1785–801.

35. Lu J, Breitwieser FP, Thielen P, Salzberg SL. Bracken: estimating species abundance in

metagenomics data. PeerJ Comput Sci. PeerJ Inc.; 2017;3:e104.

36. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA

gene database project: improved data processing and web-based tools. Nucleic Acids Res.

2013;41:D590–6.

37. Salazar G, Ruscheweyh H-J, Hildebrand F, Acinas SG, Sunagawa S. mTAGs: taxonomic profiling

using degenerate consensus reference sequences of ribosomal RNA genes. Bioinformatics [Internet].

2021; Available from: http://dx.doi.org/10.1093/bioinformatics/btab465

38. Giovannoni SJ. SAR11 Bacteria: The Most Abundant Plankton in the Oceans. Ann Rev Mar Sci.

2017;9:231–55.

39. Grote J, Thrash JC, Huggett MJ, Landry ZC, Carini P, Giovannoni SJ, et al. Streamlining and core

genome conservation among highly divergent members of the SAR11 clade. MBio [Internet]. 2012;3.

Available from: http://dx.doi.org/10.1128/mBio.00252-12

40. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes

with the Genome Taxonomy Database. Bioinformatics [Internet]. 2019; Available from:

http://dx.doi.org/10.1093/bioinformatics/btz848

41. Shi W, Moon CD, Leahy SC, Kang D, Froula J, Kittelmann S, et al. Methane yield phenotypes

linked to differential gene expression in the sheep rumen microbiome [Internet]. Genome Research.

2014. p. 1517–25. Available from: http://dx.doi.org/10.1101/gr.168245.113

42. Wirbel J, Zych K, Essex M, Karcher N, Kartal E, Salazar G, et al. Microbiome meta-analysis and

cross-disease comparison enabled by the SIAMCAT machine learning toolbox. Genome Biol.

2021;22:93.

43. Paoli L, Ruscheweyh H-J, Forneris C, Kautsar S, Clayssen Q, Salazar S, et al. Uncharted

biosynthetic potential of the ocean microbiome. submitted. 2021;

44. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile metagenomic

assembler. Genome Res. 2017;27:824–34.

45. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene

recognition and translation initiation site identification. BMC Bioinformatics. 2010;11:119.

46. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009;25:1754–60.

47. Eren AM, Esen ÖC, Quince C, Vineis JH, Morrison HG, Sogin ML, et al. Anvi'o: an advanced

analysis and visualization platform for 'omics data. PeerJ. 2015;3:e1319.

48. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality

of microbial genomes recovered from isolates, single cells, and metagenomes. Genome Res.

2015;25:1043–55.

49. Mende DR, Letunic I, Maistrenko OM, Schmidt TSB, Milanese A, Paoli L, et al. proGenomes2:

an improved database for accurate and consistent habitat, taxonomic and functional annotations of

prokaryotic genomes. Nucleic Acids Res. 2020;48:D621–5.

50. Rognes T, Flouri T, Nichols B, Quince C, Mahé F. VSEARCH: a versatile open source tool for

metagenomics. PeerJ. 2016;4:e2584.

51. Mende DR, Sunagawa S, Zeller G, Bork P. Accurate and universal delineation of prokaryotic

species. Nat Methods. Nature Publishing Group; 2013;10:881–4.

52. Oksanen J, Kindt R, Legendre P, O'Hara B, Stevens MHH, Oksanen MJ, et al. The vegan

package. Community ecology package. 2007;10:719.

53. Olm MR, Brown CT, Brooks B, Banfield JF. dRep: a tool for fast and accurate genomic

comparisons that enables improved genome recovery from metagenomes through de-replication.

ISME J. 2017;11:2864–8.

54. Hirschberg JB, Rosenberg A. V-Measure: A conditional entropy-based external cluster evaluation

[Internet]. Columbia University; 2007. Available from:

https://academiccommons.columbia.edu/doi/10.7916/D80V8N84

55. Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, et al. Mash: fast

genome and metagenome distance estimation using MinHash. Genome Biol. 2016;17:132.

56. Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful

approach to multiple testing. J R Stat Soc. Wiley; 1995;57:289–300.

57. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements

in performance and usability. Mol Biol Evol. 2013;30:772–80.

58. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. Bioinformatics. 2014;30:1312–3.

59. Zhang X, Tu B, Dai L-R, Lawson PA, Zheng Z-Z, Liu L-Y, et al. Petroclostridium xylanilyticum

gen. nov., sp. nov., a xylan-degrading bacterium isolated from an oilfield, and reclassification of

clostridial cluster III members into four novel genera in a new Hungateiclostridiaceae fam. nov. Int J

Syst Evol Microbiol. 2018;68:3197–211.

Supplementary Information

Supplementary Information for this manuscript includes:

- Legends for Supplementary Figures 1-2

- Legends for Supplementary Tables 1-5

Supplementary Figure 1. Environment-specific membership of genomes in ref-, meta- and ext-mOTUs.

A total of 499,512 genomes derived from 23 environments (environments with few genomes are grouped as 'Other', see Supplementary Tables 1 and 3) were used for the extension. The number of genomes was normalized by environments. The proportions of genomes per environment that are either associated with ref- and meta-mOTUs or were used to build ex-mOTUs are shown in the colors blue, green or orange, respectively. For example, the majority of genomes from the human gut match ref-mOTUs, whereas the vast majority of genomes from the fish environment are used to build ext-mOTUs.

Supplementary Figure 2. OPAL score broken down to individual metrics for the 63 mouse gut metagenomic samples.

The evaluation was performed using the OPAL tool [1] on 63 simulated mouse gut metagenomes [2], which also provided taxonomic profiles for seven different taxonomic profiling tools, and to which we have added mOTUs3 profiling results. The OPAL tool ranks the tools for each sample and for each taxonomic level. The measures considered are completeness, purity, L1 norm error and weighted UniFrac error, shown individually in the bottom 4 plots. Tools with a lower score perform better, as the OPAL score is a sum over rank. The top plot represents the OPAL sum of scores, which is the sum over the four individual measures. mOTUs3 scored best in all categories, including the OPAL sum of scores.

Supplementary Table Legends

Supplementary Table 1: Included studies and associated environments.

Data from 91 studies from 23 environments were included in the extension and/or profiling of the mOTUs database. Of these, 39 studies were selected for in-house MAG reconstruction and 11,164 sequencing samples from 67 studies were used for taxonomic profiling.

Supplementary Table 2: Sequencing samples included in the taxonomic profile.

A total of 11,164 samples were taxonomically profiled. Sample names are connected to public repositories by biosample and sequencing run ids. The project name column links the sample name to the study name used in Supplementary Table 1.

Supplementary Table 3: Breakdown of taxonomic novelty in ext-mOTUs.

Taxonomic novelty increases with higher ranks, i.e., more than 50% of ext-mOTUs were assigned to previously unknown families.

Supplementary Table 4: Contribution of genomes to ref-, meta- or ext-mOTUs.

Genomes/MAGs from different studies and environments contribute in varying proportions to the extension of the database.

Supplementary Table 5: Data for Figure 1.

For each sample that passed the filter (total 5,756), we reported the relative abundance for each mOTU type. Additionally, we added the total number of detected mOTUs and the habitat.

References

1. Meyer F, Bremges A, Belmann P, Janssen S, McHardy AC, Koslicki D. Assessing taxonomic metagenome profilers with OPAL. Genome Biol. 2019;20:51.

2. Meyer F, Lesker T-R, Koslicki D, Fritz A, Gurevich A, Darling AE, et al. Tutorial: assessing metagenomics software with the CAMI benchmarking toolkit. Nat Protoc. 2021;16:1785–801.