Title

Life on the leaf: Seasonal activities of the phyllosphere microbiome of perennial crops

Authors

Adina C. Howe[1,2,3], Nejc Stopnisek[,4,5,6], Shane K. Dooley[1], Fan Yang[1], Keara L. Grady[4,5], Ashley
Shade[,4,5,6,7,8]*

<u>**Affiliations**</u>

1. Department of Agricultural and Biosystems Engineering, Iowa State University,
10   Ames, IA 50011
11   2. Department of Bioinformatics and Computational Biology, Iowa State University,
12   Ames, IA 50011
13   3. Center for Advanced Bioenergy and Bioproducts Innovation, Ames IA 50011
14   4. The Great Lakes Bioenergy Research Center, Michigan State University, East Lansing,
15   MI 48824
16   5. Department of Microbiology and Molecular Genetics, Michigan State University, East
17   Lansing, MI 48824
18   6. The Plant Resilience Institute, Michigan State University, East Lansing MI 48824
19   7. Department of Plant, Soil and Microbial Sciences, Michigan State University, East
20   Lansing MI 48824
21   8. Program in Ecology, Evolution, and Behavior, Michigan State University, East Lansing
22   MI 48824

Abstract

Plants and microorganisms form beneficial associations. Understanding plant-microbe

interactions will inform microbiome management to enhance crop productivity and resilience

to stress. Here, we apply a genome-centric approach to identify key leaf microbiome members

on field-grown switchgrass and miscanthus, and quantify their activities for switchgrass over

two growing seasons. We integrate metagenome and metatranscriptome sequencing from 192

leaf samples collected over key time points in crop phenology. We curated 40 focal

metagenome-assembled-genomes (MAGs) and conservatively focus analysis on transcript

recruitment to medium and high-quality MAGs that were <10% contaminated and >50%

complete. Classes represented by these MAGs (Actinomycetia, Alpha- and Gamma-

Proteobacteria, and Bacteroidota) were active and had seasonal dynamics in key functions,

including enrichments in transcripts for of short chain dehydrogenase, molybdopterin

oxioreductase, and polyketide cyclase in the late season. The majority of MAGs had activated

stress-associated pathways, including trehalose metabolism, indole acetic acid degradation,

betaine biosynthesis, and reactive oxygen species degradation, suggesting direct engagement

with the host environment. We also detected seasonally activated biosynthetic pathways for

terpenes (carotenoid and isoprenoids), and for various non-ribosomal peptide pathways that

were poorly annotated. Overall, this study overcame laboratory and bioinformatic challenges

associated with field-based leaf metatranscriptome analysis to inform potential key activities of

these phyllosphere populations. These activities collectively support that leaf-associated

bacterial populations are seasonally dynamic, responsive to host cues and interactively engage

in feedbacks with the plant.

Keywords

Metagenome, metatranscriptome, leaf, bacteria, archaea, bioenergy, biofuel feedstocks,

phytobiome, switchgrass, miscanthus, time-series, ROS, IAA, isoprene

Introduction

Perennial plants are a crucial target for the sustainable development of biofuels

(Robertson *et al.*, 2017; Hestrin *et al.*, 2021; Ma *et al.*, 2021). In addition to yielding high

biomass that can be converted to biofuels and bioproducts, perennial crops offer a broad range

of ecosystem services that support efforts to mediate climate change, including greenhouse gas

mitigation and promotion of soil nutrient cycling (Heaton *et al.*, 2008; Langholtz *et al.*, 2016;

Robertson *et al.*, 2017; Roley *et al.*, 2018). Like all plants, perennials harbor diverse microbiota,

and many of these microbes are either known or expected to benefit their hosts. For example,

plant-associated microbes can increase productivity and protect against environmental

stressors. Because of the intimate engagement of many plant-associated microbiota with the

host, management of the plant microbiome is one tool proposed to promote crop vigor and

support crop resilience to global climate changes (Busby *et al.*, 2017; Toju *et al.*, 2018; Haskett

et al., 2020; Wang and Haney, 2020). Therefore, along with selective breeding and data-

informed field management, regulating the plant microbiome is expected to be key for the

sustainable production of perennial biofuel feedstocks.

Plants have anatomical compartments that each are inhabited by distinctive microbial

consortia. Generally, the diversity and composition of the plant microbiome narrows from

external compartments to internal, and the plant plays an active role in filtering the

microbiome composition inward (Hacquard *et al.*, 2015; Hardoim *et al.*, 2015; Gopal and Gupta,

2016). External plant compartments include the root zone, rhizosphere and rhizoplane below

ground, and the epiphytic phyllosphere above ground (Andrews and Harris, 2000). External

compartments have relatively higher representation of transient or commensal microbial taxa,

and these compartments engage with and recruit microbes from the immediate environment.

Internal compartments include the endosphere of above- and below-ground tissues, and these

have relatively low richness and harbor the most selected microbiota (Bulgarelli *et al.*, 2013;

Müller *et al.*, 2016). Of these compartments, the rhizosphere has received the most attention

as a key site of microbial-plant interactions that are important for nutrient and water

acquisition (e.g., Kuzyakov and Razavi, 2019). However, members of the microbiota that

inhabit the phyllosphere also provide important plant functions, such as pathogen exclusion

and immune priming (Bell *et al.*, 2019; Chen *et al.*, 2020). Phyllosphere microorganisms have

specialized adaptations to their exposed lifestyle (Lindow and Brandl, 2003; Vorholt, 2012;

Müller *et al.*, 2016; Koskella, 2020) and contribute to global carbon and other biogeochemical

cycling, including transformations relevant for climate change (Bringel and Couée, 2015;

Dorokhov *et al.*, 2018; Cavicchioli *et al.*, 2019). Because perennial biofuel feedstocks are

selected to maximize foliage surface area, understanding the phyllosphere microbiome is

expected to provide insights into key microbial engagements that benefit the plant to support

productivity and stress resilience.

There are two general challenges in regulating the microbiome to promote crop vigor

and resilience to environmental stress. The first challenge is to distinguish the beneficial

members of the plant microbiome from transient or commensal members, with recognition

that some members that provide host benefit likely change situationally, either over plant

development or given environmental stress (Edwards *et al.*, 2015; Xu *et al.*, 2018; Zhalnina *et*

al., 2018), while others are stable (Shade and Stopnisek, 2019; Stopnisek and Shade, 2021). The

second challenge is that plants and their agroecosystems are temporally dynamic over the

growing season, and their associated microbiota are also dynamic. It is currently unclear what

functions may be associated with phyllosphere microbial dynamics and their potential

interactions with plant hosts.

Previously, we used 16S rRNA gene amplicon analysis to identify a "core" cohort of

bacterial and archaeal taxa that were persistently associated with the phyllosphere

microbiomes' of two perennial biofuel feedstocks, miscanthus and switchgrass. Persistent

membership was established by collecting leaf samples over replicated field plots, over a

temperate seasonal cycle, and across two annual growing seasons for switchgrass (Grady *et al.*,

2019). Other studies in switchgrass have similarly reported that the leaf, and other plant

compartments, can be distinguished by microbiome compositions (e.g., Bulgarelli *et al.*, 2012;

Lundberg *et al.*, 2012; Bahulikar *et al.*, 2014; Roley *et al.*, 2019; Singer *et al.*, 2019), suggesting

selection to the leaf compartment. In the current study, we aimed to understand the functional

attributes and activities of persistent phyllosphere taxa, with an interest in specialized

adaptations to the leaf and interactions with the host plant that may inform the mechanisms

and nature of plant-microbe engagements. Therefore, we performed a seasonal analysis of

phyllosphere metagenomes for both miscanthus and switchgrass. We paired our metagenome

longitudinal series with metatranscriptome analyses at select time points in switchgrass

phenology to determine which functions were active seasonally. We performed genome-centric

analyses of the metagenome and focused on understanding the seasonal dynamics and

functions of a focal subset of medium- and high-quality metagenome-assembled-genomes

(MAGs) that we could bin from these data. Our results reveal functions supportive of a leaf-

associated lifestyle and seasonal activities of persistent phyllosphere members. Finally, we

provide evidence that these genomes were detected in various sites and years beyond our

original study plots, suggesting that they are general, consistent inhabitants of bioenergy

grasses.

Materials and Methods

Site description and sampling scheme

Switchgrass and miscanthus leaves and corresponding contextual data were collected at

the Great Lakes Bioenergy Research Center (GLBRC) located at the Kellogg Biological Station in

Hickory Corners, MI, USA (42°23'41.6" N, 85°22'23.1" W) (**Figure 1**). We sampled switchgrass

(*Panicum virgatum L.* cultivar Cave-in-rock) and miscanthus (*Miscanthus x giganteus*), from the

Biofuel Cropping System Experiment (BCSE) sites, plot replicates 1-4, as previously described

(Grady *et al.*, 2019). This included collecting leaves from switchgrass and miscanthus at 8 and 9

time points, respectively, in 2016 and switchgrass at 7 time points in 2017 seasons (**Table 1,**

Figure 1, Dataset 1, Dataset 2). We collected leaves for RNA isolation at three phenology-

informed switchgrass time points in 2016 (emergence, peak growth and senescence according

to GLBRC standard phenology methods https://data.sustainability.glbrc.org/protocols/165) to

assess the potential for sufficient mass and quality RNA extraction from the switchgrass leaf

surface, and then expanded to include leaves from all switchgrass sampling time points in 2017.

Leaves for RNA isolation were flash-frozen in liquid nitrogen immediately and stored at -80°C

until processing.

Phyllosphere RNA and DNA isolation

139   Phyllosphere epiphyte DNA was isolated and processed (Grady *et al.*, 2019). DNA

140 concentrations were normalized to 4 ng/ml. Phyllosphere epiphyte RNA was isolated using a

141 benzyl chloride liquid:liquid extraction, based on (Grady *et al.*, 2019) that was newly modified

142 for RNA isolation based on the published methods of (Suzuki *et al.*, 2001). Approximately 5 g of

143 intact, frozen leaf material was added to a 50 ml polypropylene conical tube (Corning #430290)

144 and kept frozen on liquid nitrogen while samples were weighed and transferred. Denaturing

145 Solution (DS) was prepared with 4.2 M guanidine thiocyanate, 25 mM sodium citrate dihydrate

146 pH 7.0, 0.5% (v/v) sodium n-laroyl sarcosine in Milli-q water and was filter sterilized through

147 0.22 mm filters. Immediately prior to extraction, a working stock of DS was prepared fresh by

148 adding 2-mercaptoethanol to a final concentration of 5% (v/v, DS/2-ME). To each leaf tube, 5

149 ml of benzyl chloride, 2.5 ml of 3M sodium acetate (pH 5.2), and 5 ml of the working stock of

150 DS/2-ME was added. The tube was incubated in a 60°C water bath for 20 minutes with

151 vortexing every 1 min. The leaves were removed from the conical tube using ethanol-sterilized

152 forceps and discarded.

153   Five ml of chloroform:isoamyl alcohol (24:1) were added to the remaining solution in

154 each conical tube, which were then shaken by hand for 15 seconds and incubated on ice for 15

155 minutes. The tubes were then centrifuged at 12,000 x g for 15 min at 4°C to separate aqueous

156 and organic phases. Up to 5 ml of the upper, aqueous phase was transferred to a clean 15 ml

157 polypropylene tube, without disrupting the white interface. Two and a half ml of sodium

158 citrate buffer (1.2 M sodium chloride, 0.8 M sodium citrate dihydrate in Milli-q water, filter

159 sterilized at 0.22 mm) and ice-cold isopropanol were added to achieve a final volume of 12.5

160 ml. Next, the tubes were centrifuged at 12,000 x g for 15 minutes to pellet the RNA, and the

remaining supernatant aspirated using a pipette. The RNA pellets were resuspended in 0.3 ml

of working DS/2-ME solution, and afterwards 0.3 ml of ice-cold isopropanol was added and

mixed by pipetting gently. The solution was incubated for 30 minutes at -20°C, and then the full

volume was transferred to a clean nuclease-free 1.7 ml tube and centrifuged at 16,000 x g for

15 minutes at 4 °C. The supernatant was removed, and the pellet washed in 1 ml of nuclease-

free 75% ethanol. Tubes were then centrifuged at 16,000 x g for 15 minutes at 4 °C, and

supernatant again removed using a pipette. The remaining pellet was air dried to completely

remove residual ethanol, then resuspended in 30 ml of nuclease-free Tris-EDTA buffer, pH 8.0.

Genomic DNA (gDNA) was removed using RNase-free DNase I (Thermo Fisher #AM2222)

per manufacturer's instructions. The RNA was then purified using the RNeasy MinElute Cleanup

Kit (Qiagen Germantown, MD, USA) according to manufacturer's instructions. The absence of

contaminating gDNA was confirmed by lack of amplification of the 16S rRNA gene V4 region by

PCR (Caporaso *et al.*, 2011) with positive and negative controls. This RNA isolation method was

developed to most closely align with our established phyllosphere epiphyte DNA isolation

(Grady *et al.*, 2019) in order to minimize potential bias introduced during biofilm disruption or

microbial cell lysis, as well as to minimize contamination from host RNA or genomic DNA by

leaving the plant tissue intact. Commercial RNA extraction kits are primarily based on grinding

or bead beating whole tissue samples, which would result in over-representation of host-

derived nucleic acids and potentially introduce bias in microbial cell lysis efficiencies.

Metagenome and metatrascriptome library preparation

The Joint Genome Institute (JGI) performed the library preparation and sequencing from

submitted RNA and DNA samples. Plate-based DNA library preparation for Illumina sequencing

was performed on the PerkinElmer Sciclone NGS robotic liquid handling system using Kapa

Biosystems library preparation kit. 1.82 ng of sample DNA was sheared to 436 bp using a

Covaris LE220 focused-ultrasonicator. The sheared DNA fragments were size selected by

double-SPRI and then the selected fragments were end-repaired, A-tailed, and ligated with

Illumina compatible sequencing adaptors from IDT containing a unique molecular index

barcode for each sample library. The prepared libraries were quantified using KAPA

Biosystems' next-generation sequencing library qPCR kit and run on a Roche LightCycler 480

real-time PCR instrument. Sequencing of the flowcell was performed on the Illumina HiSeq

sequencer following a 2x151 indexed run recipe.

At JGI, plate-based RNA sample prep was performed on the PerkinElmer Sciclone NGS

robotic liquid handling system using Illumina Ribo-Zero rRNA Removal Kit (Bacteria) and the

TruSeq Stranded Total RNA HT sample prep kit following the protocol outlined by Illumina in

their user guide: https://support.illumina.com/sequencing/sequencing_kits/truseqstranded-

total-rna.html, and with the following conditions: total RNA starting material of 100 ng per

sample and 10 cycles of PCR for library amplification. The prepared libraries were quantified

using KAPA Biosystems' next-generation sequencing library qPCR kit and run on a Roche

LightCycler 480 real-time PCR instrument. Sequencing of the flowcell was performed on the

Illumina NovaSeq sequencer using NovaSeq XP V1 reagent kits, S4 flow cell, following a 2x151

indexed run recipe.

Quality filtering of metagenomes and metatranscriptomes

We proceeded with bioinformatic analysis (**Figure 2**) of 192 metagenome (**Dataset 1**)

and 78 metranscriptome (**Dataset 2**) observations that met JGI standards for raw data quality

based on the Illumina proprietary software. We used Trimmomatic (v0.39) (Bolger *et al.*, 2014)

to remove adaptors and filter low-quality reads from fastq files using the following arguments:

PE -phred33 ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10:8:TRUE LEADING:3 TRAILING:3

SLIDINGWINDOW:4:15 MINLEN:36. After assembly, plant host reads were filtered out (for both

metagenomes and metatranscriptomes) by removing all reads that mapped to the switchgrass

genome (*Panicum virgatum* v1.0, DOE-JGI, http://phytozome.jgi.doe.gov/) or miscanthus

genome (*Miscanthus sinensis* V7.1 http://phytozome.jgi.doe.gov/) using bowtie2 (v 2.4.1),

samtools (v 1.13), and bedtools (v2.30.0) (Li *et al.*, 2009; Quinlan and Hall, 2010; Langmead and

Salzberg, 2012) (**Figure S1)**. To remove the fungal reads and improve the prokaryotic signal in

metagenome samples, we also filtered reads against 7 fungal genomes (Gostinčar *et al.*, 2014;

Druzhinina *et al.*, 2018; Gill *et al.*, 2019; Haridas *et al.*, 2020) that represent close relatives of

the most abundant fungal species in this system that we assessed and reported in our prior

work (Bowsher *et al.*, 2020) (**Table S1**). The genomes of these fungal species were retrieved

from the JGI Genome Portal.

Metagenome assemblies and metagenome-assembled genome binning, curation, refinement,

annotation

Two metagenome assemblies, one for switchgrass and one for miscanthus, were created based

on metagenomes collected in 2016. These filtered metagenome reads were combined and used

for co-assembly with MEGAHIT (v 1.2.9) using--kmin-1pass (low sequencing depth) and--presets

meta-large (complex metagenome) (Li *et al.*, 2015).   Additionally, we curated metagenome

assembled genomes (MAGs) from the 2016 switchgrass and miscanthus metagenome libraries

(n = 136 metagenomes) using Metabat (v2.2.15) (Kang *et al.*, 2019). MAG assemblies were

performed using filtered reads from switchgrass and miscanthus sampled from 2016,

separately, to maximize completeness and reliability, as also done in other studies (Nayfach *et*

al., 2019). To assess the MAGs quality and completeness, we used CheckM (v.1.13 with the

lineage_wf option) estimates of quality and completeness (Parks *et al.*, 2015). Among 238

MAGs assembled from the switchgrass or miscanthus phyllosphere metagenomes (**Dataset 3**),

we selected a subset of MAGs based on: completeness greater than 50% and contamination

less than 10%. We identified replicated bins associated with MAGs using dRep (v3.2.0,Olm et

al. 2017), resulting in the removal of a single MAG. An additional MAG was removed due to

insufficient read recruitment in metatranscriptomes (described below), for a final total of 40

focal MAGs, including 7 high-quality and 33 medium quality (Bowers *et al.*, 2017) (**Dataset 3).**

Read recruitment was performed with MAGs that meet either medium- and/or high-

quality standards (Swan *et al.*, 2013; Bowers *et al.*, 2017). The metagenome abundance of

contigs in MAGs in each sample was estimated based on the median coverage of filtered

metagenome reads associated to each MAG contig. Specifically, Bowtie2 (v2.2.2) was used to

align reads to all focal MAG contigs (using default setting and allowing for a single read to map

to only one reference). Bedtools (v2.28) was used to estimate the coverage of each basepair

within the contig. The estimated abundance of contig was based on the median basepair

coverage of all reads mapped to the contig, and the estimated abundance of a MAG was based

on the average median coverage of all contig within its bins. The metatranscriptome

abundance was estimated based on protein-encoding genes identified in MAGs. For each MAG

contig, open reading frames (ORFs) and functional genes were identified using Prodigal (v2.6.3,

default parameters). Transcripts were mapped to ORFs associated with each MAG to estimate

median base coverage of each ORF (Bowtie2, default parameters, no multiple mappings

allowed). To normalize varying sequencing depths, estimated abundances were normalized by

the sum of the median base pair coverage of housekeeping genes identified in each sample.

Housekeeping genes were identified based on full sequence alignment to the HMM models of

71 housekeeping single-copy genes with E-value of less than 1e-5 (Eren *et al.*, 2015; Lee, 2019).

If housekeeping genes were not identified in a metagenome or metatranscriptome, samples

were removed from further analysis. We also estimated the total reads associated with each

MAG for metagenomes and metatranscriptomes (**Dataset 4**). One MAG, M22, had an average

of 48 reads map to metatranscriptome and was removed for further analysis. For

metatranscriptome analyses, only ORFs with the top 75% of observed abundances and in at

least 10% of samples were considered.

Functional annotation of ORFs in focal MAGs was done with the DRAM tool (v1.1.1,

(Shaffer *et al.*, 2020) using UniRef90, MEROPS, PFAM, dbCAN-HMMdb (v8) databases (all

compiled with DRAM on February 12, 2021), and the KEGG database which we manually added

to the DRAM pipeline (release January 1, 2018).   To obtain functions related to terpene

metabolism, ORFs associated with any KEGG annotation that contained the phrase "terpen"

were selected.

MAGs were assigned taxonomy using GTDB-tk (v1.4.0, (Parks *et al.*, 2018)). Assembled

focal MAGs were aligned agatimst the chloroplast genome of *Panicum virgatum* (NC015990)

(BLAST, 2.10.1). We detected partial matches to eight bins and no full alignments, confirming

that those bins were not chloroplast genomes. Additionally, we compared the focal MAGs

taxonomy to our previously detected 16S rRNA gene core cohort (Grady *et al.*, 2019). This

cohort consists of 61 phylogenetically diverse bacterial OTUs (97% clustering). Taxonomy of

these 61 OTUs was compared with the 40 MAGs at the genus level only. Due to differences in

nomenclature between the GTDB and SILVA databases we removed extensions for MAGs

classified as *Pseudomonas_E* or *Aeromonas_A*.

Statistical analyses

Pairwise comparisons of functional roles (the cumulative sum of all associated ORFs) between

the early (May – June) and late (July – Sep) season were performed with the Kruskal-Wallis test

by ranks. Early and late season were delineated by plant phenology (flowering/senescence in

late season). The distribution and dynamics of MAGs' transcripts were compared with Ward's

method for hierchical clustering using euclidian distances of the estimated metatranscriptome

abundances using *pvclust* (version 2.20)(Suzuki and Shimodaira, 2006).

Assessment of biosynthetic gene clusters (BGC) on focal MAGs

287   Biosynthetic gene clusters (BGC) were predicted by antiSMASH (v6.0) (Blin *et al.*, 2021)

and further annotated by Big-SCAPE (v1.1.0) (Navarro-Muñoz *et al.*, 2020). While there are only

8 BGC classes used in Big-SCAPE (i.e. PKS I, PKS other, NRPS, RiPPs, Saccharides, Terpenes,

PKS/NRPS Hybrids, Others), antiSMASH provides more detailed classification. We leveraged

both outputs to investigate the diversity and expression of predicted BGCs in focal MAGs. From

each predicted gene cluster, we extracted the location of the biosynthetic gene (i.e.,

gene_kind="biosynthetic" and core_position). To evaluate the transcription of putative BGCs

within a MAG, we searched for any transcripts that are mapped to the same genomic region of

predicted biosynthetic genes.

Predicting functional potential of focal MAGs

We used gapseq (Zimmermann *et al.*, 2021) to predict completed metabolic pathways in

our focal MAGs. We used the 'find -p all –b 200' option to search for pathways against the

MetaCyc database. We filtered out incomplete pathways and remaining pathways were

grouped into broader categories using MetaCyc classification and manually curated to focus on

understanding the pathways relevant either to the plant environment or for microbial

interactions with plants. These categories were defined as potential involvement in: i) plant

(using plant metabolites/cell components), ii) phytohormone (known/potential involvement in

phytohormone homeostasis), iii) stress (e.g., drought, reactive oxygen species), and iv) general

(pathways that utilize potential plant derived products). Furthermore, we also manually

searched for genes/pathways that were missed by gapseq but are known to be related to

adaptation to plant-associated lifestyle, including secretion systems (Tseng *et al.*, 2009; Lucke *et*

al., 2020), oxidation of trace gases (H_2 and CO)(Bay *et al.*, 2021; Palmer *et al.*, 2021), oxalate

degradation, and phytohormone production/degradation. Similar to our biosynthetic gene

analysis, the activity of predicted pathways was estimated based on mapped transcripts to key

identified genes (**Table S2**). Contigs of focal MAGS were aligned (BLAST v2.7.1+) against nine

isoprenoid precursor biosynthesis genes previously reported (Julsing *et al.*, 2007). The top hit

of each query was considered and accepted as aligned if E-value was less than 1e-5.

Detection of MAGS in public metagenomes

To evaluate the presence of MAGs in other switchgrass and miscanthus metagenomes, 55

publicly available switchgrass, miscanthus, and corn metagenomes were used (**Dataset 5**).

Metagenome reads were mapped to focal MAGs, using the same methods for mapping and

abundance estimation described above for the metagenomes generated in this study.

Data and code availability

Metagenome and metatranscriptome data are available through the Joint Genome Institute

Genome Portal under proposal ID 503249. Code and links to data and metadata are available

on GitHub (https://github.com/ShadeLab/PAPER_Howe_2021_switchgrass_MetaT). MAGs are

available in NCBI under project IMG. All metadata, including metadata standards for

metagenomes, metatranscriptomes, and metagenome-assembled genomes are provided as

supporting datasets (**Datasets 1-6**).

<u>Results</u>

Key leaf bacterial populations and general dynamics

Among all assembled MAGs (n=238), we focused analysis on 40 that were high- and

medium- quality based on completeness and contamination standards (**Figure 3A**, **Dataset**

3)(Bowers *et al.*, 2017). The consistent detection of these 40 MAGs in both miscanthus and

switchgrass samples (**Figure 4**) suggests that their originating populations were not host-

specific but rather distributed among these perennial grasses; this supports our previous results

from an 16S rRNA gene amplicon survey that revealed substantial overlap in the major leaf

bacterial taxa across the two crops (Grady *et al.*, 2019). Furthermore, there were 38 "core"

OTUs from our prior survey that were also found in 25 focal MAGs here per their taxonomic

identifications (**Figure 3B**); our previous study prioritized OTUs based on abundance and

seasonal and field-replicated occupancy, and so this conservatively shows that at least half of

the focal MAGs represent the most abundant and consistently detected taxa in this ecosystem.

The 40 focal MAGs most represented the orders Rhizobiales (n=8/40), Actinomycetales

(n=6/40), Burkholderiales (n=6/40), and Sphingomonadales (n=6/40) (**Dataset 3**).

Focal MAGs exhibited varied seasonal patterns in their overall metagenome (**Figure 4A-**

C) and metatranscriptome (**Figure 4D-E**) read recruitment, with early, late, and consistent

seasonal detection observed. However, in the metatranscriptomes of both years for

switchgrass, there was a strong seasonal enrichment over time, with increases in the later

months of sampling. Phenologically, August samples corresponded with (late) peak biomass

and fruiting for switchgrass, and with a closed canopy for both switchgrass and miscanthus.

Senescence occurred as early as mid-September for switchgrass and as late as November for

miscanthus. Notably, transcript recruitment to the MAGs was robust even when there was

inconsistent detection in metagenome read recruitment, suggesting that these MAGs were

consistently active and had relatively high activity even when their genome detection was

obscured by the flanking community or host signal.

MAGs were clustered based on coherence in seasonal transcript dynamics (**Figure 4D-E,**

Figure S2). Five coherent and statistically supported groups (clusters) of 40 MAGs were

identified (**Figure S2**). Cluster 1 contained a single MAG S28, assigned to genus *Pseudomonas*

(S28, 98% complete, 1.9% contamination) and its genes were enriched early in the season

(2017) but the MAG was highly active throughout the season. The other four clusters contained

numerous MAGs and were relatively more dynamic, with trends towards late-season

enrichment in transcripts. Several clusters contained MAGs annotated to the same class,

suggestive of taxonomic coherence in seasonal activities.   For example, cluster 3 included all

but one (of eight) Actinomycetia MAGs.

MAG genes and activities support a leaf-associated lifestyle

Not surprisingly, the most abundant subsystems identified among the 40 focal MAGs were

associated with bacterial growth, such as carbohydrates, energy, amino acid, and nucleotide

metabolisms (**Figure 5**). Leaf metatranscriptomes also showed that most subsystem transcripts

were steadily enriched over the season. However, subsystem associated with colonization and

adaptation, such as cell motility, signal transduction, community-associated pathways, and

environmental adaptation trended down in their normalized transcripts over the season. A

closer investigation of these down-trending functions revealed that they could be attributed

almost exclusively to MAG S28 (*Pseudomodales* from cluster 1 in **Figure 4DE** and **Figure S3**).

Finally, a few KEGG classifications were relatively stable and lacked overall seasonal trends,

including transcripts for translation, transcription, and antimicrobial resistance.

We selected the most prominent phyllosphere transcripts based on their high detection

among the 40 focal MAGs, hypothesizing that some of these functions shared between

bacterial populations may represent functions for survival on the leaf. A total of 124 distinct

functional roles were identified in at least 39 of the 40 MAGs (**Dataset 6, Figure S4**). Of these,

most functions were associated to translation (n=48/124). Other broadly present functions

were associated with metabolism of energy (n=29/124) and carbohydrates (n=28/124). There

were also several functional roles associated with cofactors and vitamins (n=18/124). While

many functions were enriched in transcripts in the late season (July-Sept) relative to the early

season (May-June), three functional roles in particular stood out, with enrichments greater

than 20-fold in the late season transcripts and included proteins classified as short chain

dehydrogenase, molybdopterin oxidoreductase, and polyketide cyclase (**Table 2**, **Dataset 6,**

Figure S4).

We then performed metabolic, biosynthetic, and plant-associated gene pathway

analyses to understand the functions detected among focal MAGs in the phyllosphere and their

activities inferred by transcript recruitment to the pathways (**Figure 6**). Pathways for terpene

metabolism (34/40), betaine biosynthesis (30/40), trehalose metabolism (25/40), cyanide

degradation (21/40), ROS degradation (23/40), and indole acetic acid (IAA) degradation (40/40)

were the most common pathways shared and active across MAGs, and were found in lineages

from all four classes represented by the focal MAGs. Thus, these pathways likely are common

among phyllosphere members and highly supportive of a leaf-associated lifestyle. In addition,

there were some functional pathways and activities that may be more specialized functions

because they were detected in only a few MAGs (e.g., xylitol metabolism). However, there was

no clear phylogenetic pattern to the distributions of these putative specialized functions.

399   Because of the near-ubiquitous detection of terpene-related biosynthetic genes on the

focal MAGs, we examined their annotations and recruitment more closely using gene function

predictions (i.e., ORFs), which was expected to improve sensitivity of functional annotation

relative to metabolic pathway tools (e.g., BGC). We detected 278 ORFs associated with

terpenes annotations in the metatranscripts. Collectively, these terpene-related genes

followed the trend of gradual enrichment over the season (**Figure 7A**) and were relatively

abundant in both years. The highest enrichment subcategory was terpenoid backbone

biosynthesis, which included 29 ORFs. Among several isoprene biosynthesis genes detected, of

interest was relatively high, seasonal enrichment of the two terminal enzymes in the non-

mevalonate isoprene biosynthesis pathway that is employed by bacteria, the *gcpE* and *lytB*

(**Figure 7B, Figure S5**). *GcpE* and *lytB* gene transcripts were detected in more than a third of the

MAGs (13/40), and these included MAGs representing all three phyla detected. Six of the eight

Proteobacteria that recruited isoprene biosynthesis transcripts were annotated to

Methylobacterium (alpha). The four Actinomycetia were more distributed phylogenetically and

annotated as genera *Frigoribacterium* M1, *Microbacterium* M105, *Amnibacterium* M67, and

Pseudokineococcus M86. The single Bacteroidota that had isoprene biosynthesis transcripts

was a *Hymenobacter* M9. We then investigated the 40 MAGs for detection of any of the nine

genes previously reported to be involved in bacterial biosynthesis of isopentenyl diphosphate, a

precursor terpenoids like isoprene (Julsing *et al.*, 2007) and found that 29/40 MAGs had six or

more of the genes detected, and that all MAGs had at minimum 2 of the genes (**Table S2**). The

consistent detection of genes associated with isoprene pathways in these MAGs, which are

>50% complete, suggests that biosynthesis of isoprene-related molecules may be a prominent

leaf strategy by phyllosphere bacteria. *Psuedomonas* MAG S28, noted previously to be the

dominant population that colonized and activated early in the season (**Figure 4, Figure S2**), had

high isoprene biosynthesis transcript enrichment early in the season that then declined.

However, the other eleven MAGs harboring genes from isoprene biosynthesis pathways

compensated with increased activity in the late season.

Returning to the BGC annotations (by antiSMASH , **Figure 6**), most of the terpene

transcripts were associated to pigment biosynthesis (e.g., carotenoids). The second most

consistent class of BGC transcripts on the focal MAGs were non-ribosomal peptide synthase

(NRPK) genes, but the majority of these did not have additional annotations beyond the general

category. Therefore, the ORF analysis and BGC detection were complementary in results,

especially for terpene-related functions. Another notable BGC finding was that all but two (of

eight) Actinomycetia MAGs had transcripts for type III polyketide synthases, while this BGC was

less commonly detected among Proteobacteria and Bacteroidota.

MAGs were detected in metagenomes from different field sites, crops, and years

We next asked if these focal MAGs were detected more broadly in other crop

metagenomes. Because leaf metagenomes were not findable, we curated related soil

metagenomes from local cropping systems in Michigan and also from the publicly available

data within the Integrated Microbial Genomes (IMG) database, which serves as a repository for

the Joint Genome Institute's sequencing efforts through the US Department of Energy. We

were surprised to find that, for each metagenome investigated, reads could be mapped to this

set of MAGs. This included soil metagenomes from both switchgrass and miscanthus fields,

three locations (Iowa, Michigan, and Wisconsin) and sampled across five different years (**Figure**

8). These results suggest that the detected and analyzed MAGs can be widely distributed in

midwestern agroecosystems, and potentially of general importance for perennial crop

environments.

Discussion

Here, we report the first multi-year seasonal metagenome and metatranscriptome

assessment of plant phyllosphere in agricultural field conditions, focusing on the bacterial

functions associated with two promising biofuel feedstocks. We expect these findings to have

relevance for other grasses or systems with substantial aerial biomass, including native prairie.

Furthermore, our collection of MAGs included phyllosphere members previously reported as

abundant, persistent, or key for microbiome assembly in other plants, including the model

Arabidopsis (e.g., (Carlström *et al.*, 2019), specifically: Sphingomonadales, Pseudomonadales,

Actinomycetales, Burkholderiales, and Rhizobiales. Therefore, the patterns and consistently

detected functions among this curated collection offer general insights as well as seasonal

phyllosphere functions.

There are multiple lines of evidence that the focal MAGs discussed here represent key

lineages in the switchgrass and miscanthus phyllosphere. First, there is ample overlap among

these focal MAGs and the core taxa of high abundance and occupancy from our previous 16S

rRNA gene amplicon analysis of the switchgrass and miscanthus phyllosphere diversity (Grady

et al., 2019), including a MAG associated to *Hymenobacter* M9, which is a genus that had high

occupancy across fields and over time, as well as several OTUs assigned. This same time series

was investigated in our previous amplicon analysis and the core taxa were prioritized using

consistency across replicated fields at the same time point, persistence over time, and relative

abundance. This suggests that the populations represented by the MAGs are not rare taxa that

are transient to the system. Second, we were able to generate quality assemblies from the

complex metagenome data, which is a process that is generally biased towards abundant

members. These MAGs are highly represented in read abundance in metagenomes and

recruited relatively more metatranscriptome reads as well, suggesting that they are both

abundant and active in the phyllosphere. Though it is possible that there are key lineages

missing among our collection, we are confident that those discussed here are among the major

host- or environment-selected populations inhabiting the switchgrass and miscanthus

phyllosphere.

Stress responses: trehalose, betaine, reactive oxygen, IAA

Trehalose is a disaccharide that protects cells against salt, water, and osmolyte stress by

serving as a stabilizing chemical chaperone, either by displacing water from protein surfaces or

by vitrifying around protein structures to shield them (Laskowska *et al.*, 2020). Similarly,

betaine is another commonly biosynthesized osmoprotectant used by microorganisms to

contend with water, salt, and temperature stress (Zou *et al.*, 2016). Both have been

hypothesized to be important survival strategies of microorganisms in the phyllosphere, with

supporting evidence from isolate genome analyses (Rastogi *et al.*, 2013). Here, we show both

trehalose and betaine to be prominent among MAG populations and activated consistently on

the leaf surface, suggesting that they are not seasonally activated but rather necessary for the

leaf-associated lifestyle.

In bacteria, trehalose metabolism prevents cellular overflow metabolism and carbon

stress by redirecting glucose-6-phosphate from conversion to pyruvate (Laskowska *et al.*, 2020).

Trehalose biosynthesis is common among bacteria and archaea that live in arid, saline, thermal,

or seasonally dry environments (e.g., (Urrejola *et al.*, 2019), (Dai *et al.*, 2015; Lacerda-Júnior *et*

al., 2019), and it has also been reported to be induced in a pseudomonad by ethanol (Harty *et*

al., 2019), which can originate inside plant cells (Kimmerer and MacDonald, 1987) or more

generally in the roots (e.g., Ferner *et al.*, 2012), especially during stress, fruit ripening, or

senescence (Kimmerer and Kozlowski, 1982). In switchgrass and plants in general, trehalose

concentration is increased in response to drought conditions (Liu *et al.*, 2015), and its

precursor, trehalose-6-phosphate, induces senescence when carbon is readily available

(Wingler *et al.*, 2012). Furthermore, the *A. thaliana* phyllosphere member *Sphingomonas*

melonis was reported to regulate trehalose biosynthesis during growth conditions that

promoted mild stress (Gottschlich *et al.*, 2019). Given this, it makes sense that the majority of

MAGs had enrichment of transcripts related to trehalose metabolism, which would support a

plant-associated lifestyle during drought and host senescence (Gottschlich *et al.*, 2019).

Betaine biosynthesis in bacteria often begins with oxidation of choline, which is a part of

plant tissues and can be transported into the cell (Chen *et al.*, 2013). Indeed, choline

degradation was detected in 9/10 Gammaproteobacteria MAGs here (**Figure 6**). Microbial-

derived osmolytes such as betaine and trehalose have been suggested as targets for

biotechnology development to support crop stress tolerance. However, plants can

biosynthesize betaine and can be divided into groups of those that do and do not accumulate it

in concentrations that are supportive for stress tolerance (Valenzuela-Soto and Figueroa-Soto,

2019). For example, in switchgrass, the concentration of glycine betaine was not predictive of

differences in drought tolerance among different genotypes, while trehalose was, along with

abscisic acid, spermine, and fructose (Liu *et al.*, 2015). Furthermore, our data suggest no

notable microbial limitations in the genetic potential or activation of betaine biosynthesis in

phyllosphere.

Reactive oxygen species (ROS) serve as signals for various developmental and cellular

processes in plants, and here we detected active pathways for ROS degradation in the majority

of focal MAGs. Though the precise mechanisms are unclear, homeostasis of ROS are expected

to be involved in senescence (Kerchev *et al.*, 2015), which is relevant to our study given that at

least some to a majority of senescing plants were observed per plot in August and September

sampling dates, respectively. Additionally, ROS accumulate in plants that are exposed to abiotic

stress, to negative effects. ROS degradation is one of many functions phyllosphere

microorganisms employ to contend with expected fluctuations in ROS on the leaf surface

(though, these fluxes are difficult to measure (Kerchev *et al.*, 2015)). Previously, several genes

relevant for oxidative stress response were differentially regulated in a wild-type phyllosphere

bacteria *Sphingomonas melonis* strain Fr1 as compared to a knock-out mutant for regulation of

general stress response, when both were grown in a medium expected to induce low-levels of

stress. (Gottschlich *et al.*, 2019). Given that managing plant ROS is a target for reducing crop

stress and regulating plant development (Considine and Foyer, 2014; Kerchev *et al.*, 2015), it is

possible that manipulating microbial ROS degradation could be applied as one tool to achieve

such efforts, but much more research is needed to understand the microbial-host interaction

given ROS exposure or accumulation, and any possible ROS signaling between them.

IAA is a phytohormone produced by plants to regulate many processes in growth and

stress response (e.g., Spaepen *et al.*, 2007; Egamberdieva *et al.*, 2017). It is also made by many

microorganisms, including those shown to support plant grown promotion (e.g., Egamberdieva

et al., 2017). Therefore, the activity of IAA degradation pathways by focal MAGs is expected

given the redundancies between plants and microorganisms in synthesizing and responding to

IAA and demonstrates microbiome responsiveness to feedbacks in the host environment.

Biosynthesis of isoprene-related molecules

Most of the functions identified in our MAGs suggest general requirements for growth

and maintenance given a leaf-associated lifestyle (e.g., carbohydrate and amino acid

metabolism, pigment production to protect from radiation, similar to previous reports, e.g.,

Lajoie *et al.*, 2020). However, BGC analysis revealed surprising consistency in terpene

metabolism pathways, leading us to look more closely at transcript ORFs associated to

terpenes. This analysis revealed particular enrichment in pathways and key genes associated

with isoprenoid biosynthesis. Isoprenoids are a class of volatile terpenes that are generally

abundant and reactive, and they engage in indirect and complex feedbacks with methane and

nitrous oxide greenhouse gases (McGenity *et al.*, 2018). Isoprene is one of the simplest

isoprenoids. It is released by many plant species, and much of it is synthesized within the

methylerythritol phosphate pathway of the chloroplast (MEP, aka:non-mevalonate) (Sharkey *et*

al., 2008). Isoprene is thought to act as a signaling molecule in stress response (Zuo *et al.*,

2019). Studies have also found that isoprene emission protects leaf photosynthesis against

short episodes of high-temperature (Sharkey and Yeh, 2001). Plants emit isoprene from

matured, photosynthetically active leaves, and emissions are light responsive (Sharkey *et al.*,

2008). However, senescing leaves have been reported to decrease in their isoprene emissions

relative to leaves at peak growth (Sharkey *et al.*, 1991). Both switchgrass and miscanthus have

been reported to emit relatively low basal levels of isoprene (Eller *et al.*, 2011; Morrison *et al.*,

2016).

We hypothesize that that members of the biofuel feedstock phyllosphere bacterial

community may be either compensating for the loss of plant-derived isoprene, engaging in

interspecies isoprenoid signaling with the host, protecting plant photosynthesis from thermal

damage, quenching reactive oxygen species, or possibly producing isoprenoids as overflow

metabolites (as hypothesized for *Bacillus subtilus* (Sivy *et al.*, 2002)). Bacterial isoprene

degraders and synthesizers are widespread in nature (McGenity *et al.*, 2018) and have been

previously investigated in phyllosphere communities of the relatively high isoprene emitter

Populus spp (Crombie *et al.*, 2018), as well as in soils (El Khawand *et al.*, 2016), which can serve

as an isoprene sink. Stable isotope assays have been used to determine that a subset of

bacteria community members degrade isoprene, including several Actinobacteria (*Rhodococcus*

spp.) and *Variovorax* (Proteobacteria) (El Khawand *et al.*, 2016; Crombie *et al.*, 2018). Our MAG

collection contains several Actinomycetia, a *Hymenobacter*, several *Methylobacterium*, and

Pseudomonas MAG S28 that show activation of genes involved in isoprenoid biosynthesis, and

adding support for their involvement in related molecular feedbacks in the phyllosphere.

Furthermore, given that a third of our modest collection had activity of key isoprenoid

biosynthesis genes, there may be many more leaf bacterial contributors that we did not detect

here. In addition, these activities were commonly to three Bacteria phyla, it suggests that

biosynthesis of isoprene-related molecules may be a very common phyllosphere microbiome

function. As isoprene is a precursor to sidechains needed for several quinones (Nowicka and

Kruk, 2010), it could be speculated that leaf bacteria scavange isoprene emitted by the host

plant to supplement bacterial synthesis of these sidechains, but then compensate with de novo

biosynthesis if host decreases production. We observe that isoprenoid synthesis increases

seasonally in the majority of MAGs containing these pathways, and concurrently with when

plant isoprene emissions also is expected to decrease, directs future work to understand these

dynamics and potential isoprenoid-mediated bacterial-host engagement.

MAGs of interest

We highlight three MAG populations that were of interest because of their taxonomy,

functions, dynamics, or detection. All were shared with taxa in our prior 16S rRNA survey,

supporting their inclusion as part of the "core" set that was selected by abundance and

occupancy . First, high-quality MAG S28 (>97% complete, <2% contamination) was a prominent

pioneer and active colonizer of the leaf (Figure 4 Group 1). MAG S28 is related to *Pseudomonas*

cerasi, a species reported to have phytopathogen relatives (Kałużna *et al.*, 2016), but we did not

note any disease symptoms on the leaves analyzed. This population had expected traits of a

strong surface colonizer, including colonization, adaptation and motility subsystems. It also had

six pathways related to phytohormone responses (out of 7 total phytohormone pathways

observed in these data), including activated ethene biosynthesis, ACC deaminase, and

degradation of ethylene glycol, putrescine, salicylate and IAA. These data suggest that S28 has

several mechanisms to engage or respond to the host via phytohormones.

Next, MAG M9, identified as *Hymenobacter*, was of interest because it was associated to

the most numerous taxonomic group detected in our prior 16S rRNA gene survey (Grady *et al.*,

2019) and not among the most typically investigated phyllosphere lineages in the literature.

While M9 populations were first detected early in the season, its transcripts were enriched in

the late season (**Figure 4** Group 5). MAG M9 had detected and activated galactonate, N-

acetylglucosamime, and lactose metabolisms, which were not common among the focal MAGs.

It also had activated benzoate, curcumin, and putrescine degradation, as well as cyanide

detoxification, type VI secretion, and dihydrogen oxidation. While M9 also had some pathways

that were common among these MAGs (e.g., ROS and IAA degredation, terpene biosynthesis),

its suite of more sparsely detected pathways and functions suggest a specialized role in the

phyllosphere community. Notably, M9 has 65% completion and 0% detected contamination,

suggesting more functional potential remains to be discovered for this and similar

Hymenobacter lineages inhabiting the phyllosphere.

Finally, we selected a representative Actinomycetia MAG M60, a *Quadrisphaera* lineage

that had activated isoprenoid biosynthesis and had increased activity late in the season along

with the majority of focal MAGs (Figure 4, Group 3). Studies have found that members of

Actinomycetia are important part of phyllosphere that contribute to disease prevention and

plant growth [El-Tarabily 2009; Javed 2021; Anwar 2016]. MAG M60 had several

oligo/polysaccharide metabolisms that were infrequently detected in these data, including

glycogen, melibiose, and trehalose. Despite its high completeness and low contamination

(>95% and < 5%, respectively), M60 was sparsely annotated by the methods we applied.

However, *Quadrisphaera* have been reported to be highly abundant in the phyllosphere or

endosphere of various plants (Bao *et al.*, 2020).

Conclusions

Many recent review, perspective, and opinion pieces have urged integration of multi-

omics approaches to improve understanding of the microbiome and its relationship to the host

plant (Rastogi *et al.*, 2013; Levy *et al.*, 2018; Remus-Emsermann and Schlechter, 2018; Beilsmith

et al., 2019; Trivedi *et al.*, 2020). However, most integrative studies have focused almost

exclusively on the rhizosphere as the compartment of soil-plant feedbacks and nutrient and

water acquisition for the host. Though leaves are readily accessible for sampling, the

phyllosphere microbiome is challenging to investigate using throughput, cultivation-

independent approaches like metagenomics and metatranscriptomics. There are high levels of

host and chloroplast contamination in leaf samples, and relatively low microbial biomass per

leaf that that must be first dislodged from tightly-adhered biofilms. Signal from messenger RNA

in metatranscriptome analysis is masked by abundant ribosomal RNA signal, leading to further

challenge. Because of the combination of all of these challenges, much of our understanding of

the phyllosphere, as the largest surface area of microbial habitation on Earth (Peñuelas and

Terradas, 2014), has been learned from studies that employ model hosts and synthetic or

model microbial communities in controlled settings, or from description of the community

structure by sequencing of marker genes, amplified and bioinformatically depleted of

chloroplast genes to overcome the challenges of low signal and host contamination.

Here, we report optimized laboratory protocols (to minimize host and chloroplast

signals) combined with a genome-centric bioinformatic approach to perform focused functional

gene and transcript analysis of seasonally dynamic yet persistent phyllosphere microbiome

members. To our knowledge, this is the first untargeted bacterial metatranscriptomic work

performed on the leaf phyllosphere of field-grown crops. Other recent leaf metatranscriptome

studies have focused the viral communities of tomato and pepper (Choi *et al.*, 2020), soybean

(Marzano and Domier, 2016), and rice (Chao *et al.*, 2020 Preprint). A key strength of this work

is the challenging integration of phyllosphere metagenome and metatranscriptome data,

leveraging the higher coverage of the metagenomes with the activity information available

from the metatranscriptomes. Despite the relatively limited coverage of the MAGs (due to

substantial host and ribosomal DNA contamination), the analysis proved successful by

integrating both datasets and focusing on genome-centric interpretation. Thus, there are likely

many more prevalent and functionally active populations of the phyllosphere that were not

captured in this study, including those players previously known to be key in the phyllosphere

(e.g. Delmotte *et al.*, 2009; Vorholt, 2012). Substantial additional sequencing effort or an

enrichment strategy would improve signal for a cultivation-independent approach to target

those players. While the use of genome-centric approaches has the obvious shortcoming that

we have obviously not captured every microbiome member, our approach does allow us to link

actively transcribed functions to specific microbial membership. Furthermore, the functional

genes and activities documented here are logical given current understanding of microbial

adaptation to the host and phyllosphere environment.

Overall, this work provides evidence of a thriving, dynamic, functionally diverse, leaf-

specialized, and host-responsive microbiome on the phyllosphere of perennial grasses. It

provides evidence of specific phyllosphere functions that are seasonally activated in a

temperate agroecosystem and suggests several hypotheses of important host-microbe

interactions in the phyllosphere, for example via central metabolism, isoprenoid biosynthesis,

and stress response engagements. This research contributes to our broad understanding of the

dynamics and activities of phyllosphere microbial communities, and points to specific microbial

functions to target that could prove useful for plant-microbiome management.

Acknowledgements

Support for this research was provided by the Great Lakes Bioenergy Research Center, U.S.

Department of Energy, Office of Science, Office of Biological and Environmental Research

(Awards DE-SC0018409 and DE-FC02-07ER64494), by the National Science Foundation Long-

term Ecological Research Program (DEB 1637653 and 1832042) at the Kellogg Biological Station,

Michigan State University AgBioResearch, and by the DOE Center for Advanced Bioenergy and

Bioproducts Innovation (US Department of Energy, Office of Science, Office of Biological and

Environmental Research under award number DE-SC0018420). This work was also supported in

part by Michigan State University through computational resources provided by the Institute

for Cyber-Enabled Research and in part by the University of Wisconsin-Madison Wisconsin

Energy Institute as supported by GLBRC Information Services. The work

(proposal:10.46936/10.25585/60000818) conducted by the U.S. Department of Energy Joint

Genome Institute (https://ror.org/04xm1d337), a DOE Office of Science User Facility, is

supported by the Office of Science of the U.S. Department of Energy operated under Contract

No. DE-AC02-05CH11231. AS acknowledges support by the USDA National Institute of Food

and Agriculture and Michigan State University AgBioResearch. NS acknowledges support from

the Michigan State University Plant Resilience Institute. We thank three anonymous reviewers

for their thoughtful comments on a previous version of this work.

688
Author contributions

KLG and AS conceived and designed experiments; NS, KLG and AS performed the experiments;

ACH, NS, SKD, FY and AS analyzed the data; ACH, NS, SKD, FY, KLG and AS contributed

materials/analysis tools; and ACH, NS, SKD, FY, KLG and AS wrote the paper.

Competing Interests statement

The authors declare no competing interests.

References

Andrews, J.H. and Harris, R.F. (2000) The ecology and biogeography of microorganisms on plant surfaces. *Annu Rev Phytopathol* **38**: 145–180.

Bahulikar, R.A., Torres-Jerez, I., Worley, E., Craven, K., and Udvardi, M.K. (2014) Diversity of nitrogen-fixing bacteria associated with switchgrass in the native tallgrass prairie of Northern Oklahoma. *Appl Environ Microbiol* **80**: 5636–5643.

Bao, L., Cai, W., Cao, J., Zhang, X., Liu, J., Chen, H., et al. (2020) Microbial community overlap between the phyllosphere and rhizosphere of three plants from Yongxing Island, South China Sea. *Microbiologyopen* **9**: 1–18.

Bay, S.K., Dong, X., Bradley, J.A., Leung, P.M., Grinter, R., Jirapanjawat, T., et al. (2021) Trace gas oxidizers are widespread and active members of soil microbial communities. *Nat Microbiol* **6**: 246–256.

Beilsmith, K., Thoen, M.P.M., Brachi, B., Gloss, A.D., Khan, M.H., and Bergelson, J. (2019) Genome-wide association studies on the phyllosphere microbiome: Embracing complexity in host–microbe interactions. *Plant J* **97**: 164–181.

Bell, T.H., Hockett, K.L., Alcalá-Briseño, R.I., Barbercheck, M., Beattie, G.A., Bruns, M.A., et al. (2019) Manipulating wild and tamed phytobiomes: Challenges and opportunities. *Phytobiomes J* **3**: 3–21.

Blin, K., Shaw, S., Kloosterman, A.M., Charlop-Powers, Z., Van Wezel, G.P., Medema, M.H., and Weber, T. (2021) AntiSMASH 6.0: Improving cluster detection and comparison capabilities. *Nucleic Acids Res* **49**: W29–W35.

Bolger, A.M., Lohse, M., and Usadel, B. (2014) Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics* **30**: 2114–2120.

Bowers, R.M., Kyrpides, N.C., Stepanauskas, R., Harmon-Smith, M., Doud, D., Reddy, T.B.K., et al. (2017) Minimum information about a single amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat Biotechnol* **35**: 725–731.

Bowsher, A.W., Benucci, G.M.N., Bonito, G., and Shade, A. (2020) Seasonal Dynamics of Core Fungi in the Switchgrass Phyllosphere, and Co-Occurrence with Leaf Bacteria. *Phytobiomes J* PBIOMES-07-20-0.

Bringel, F. and Couée, I. (2015) Pivotal roles of phyllosphere microorganisms at the interface between plant functioning and atmospheric trace gas dynamics. *Front Microbiol* **6**:.

Bulgarelli, D., Rott, M., Schlaeppi, K., Ver Loren van Themaat, E., Ahmadinejad, N., Assenza, F., et al. (2012) Revealing structure and assembly cues for *Arabidopsis* root-inhabiting bacterial microbiota. *Nature* **488**: 91–95.

Bulgarelli, D., Schlaeppi, K., Spaepen, S., van Themaat, E.V.L., and Schulze-Lefert, P. (2013) Structure and functions of the bacterial microbiota of plants. *Annu Rev Plant Biol* **64**: 807–838.

Busby, P.E., Soman, C., Wagner, M.R., Friesen, M.L., Kremer, J., Bennett, A., et al. (2017) Research priorities for harnessing plant microbiomes in sustainable agriculture. *PLoS Biol* **15**: 1–14.

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J., et al. (2011) Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proc Natl Acad Sci* **108**: 4516–4522.

Carlström, C.I., Field, C.M., Bortfeld-Miller, M., Müller, B., Sunagawa, S., and Vorholt, J.A. (2019)
Synthetic microbiota reveal priority effects and keystone strains in the *Arabidopsis*
phyllosphere. *Nat Ecol Evol* **3**: 1445–1454.
Cavicchioli, R., Ripple, W.J., Timmis, K.N., Azam, F., Bakken, L.R., Baylis, M., et al. (2019)
Scientists' warning to humanity: microorganisms and climate change. *Nat Rev Microbiol*
17: 569–586.
Chao, S., Wang, H., Yan, Q., Chen, L., Chen, G., Wu, L., et al. (2020) Metatranscriptomics and
small RNA analysis revealed that viral covert coinfection resulted in disease symptoms
reminiscent of rice sterility. *Prepr available Res Sq*.
Chen, C., Li, S., McKeever, D.R., and Beattie, G.A. (2013) The widespread plant-colonizing
bacterial species Pseudomonas syringae detects and exploits an extracellular pool of
choline in hosts. *Plant J* **75**: 891–902.
Chen, T., Nomura, K., Wang, X., Sohrabi, R., Xu, J., Yao, L., et al. (2020) A plant genetic network
for preventing dysbiosis in the phyllosphere. *Nature* **580**: 653–657.
Choi, H., Jo, Y., Cho, W.K., Yu, J., Tran, P.T., Salaipeth, L., et al. (2020) Identification of viruses
and viroids infecting tomato and pepper plants in vietnam by metatranscriptomics. *Int J*
Mol Sci **21**: 1–16.
Considine, M.J. and Foyer, C.H. (2014) Redox regulation of plant development. *Antioxid Redox*
Signal **21**: 1305–1326.
Crombie, A.T., Larke-Mejia, N.L., Emery, H., Dawson, R., Pratscher, J., Murphy, G.P., et al. (2018)
Poplar phyllosphere harbors disparate isoprene-degrading bacteria. *Proc Natl Acad Sci U S*
A **115**: 13081–13086.
Dai, J., Dai, W., Qiu, C., Yang, Z., Zhang, Y., Zhou, M., et al. (2015) Unraveling adaptation of
Pontibacter korlensis to radiation and infertility in desert through complete genome and
comparative transcriptomic analysis. *Sci Rep* **5**: 1–9.
Delmotte, N., Knief, C., Chaffron, S., Innerebner, G., Roschitzki, B., Schlapbach, R., et al. (2009)
Community proteogenomics reveals insights into the physiology of phyllosphere bacteria.
Proc Natl Acad Sci **106**: 16428–16433.
Dorokhov, Y.L., Sheshukova, E. V., and Komarova, T. V. (2018) Methanol in plant life. *Front Plant*
Sci **871**: 1–6.
Druzhinina, I.S., Chenthamara, K., Zhang, J., Atanasova, L., Yang, D., Miao, Y., et al. (2018)
Massive lateral transfer of genes encoding plant cell wall-degrading enzymes to the
mycoparasitic fungus Trichoderma from its plant-associated hosts.
Edwards, J., Johnson, C., Santos-Medellín, C., Lurie, E., Podishetty, N.K., Bhatnagar, S., et al.
(2015) Structure, variation, and assembly of the root-associated microbiomes of rice. *Proc*
Natl Acad Sci U S A **112**: E911-20.
Egamberdieva, D., Wirth, S.J., Alqarawi, A.A., Abd-Allah, E.F., and Hashem, A. (2017)
Phytohormones and beneficial microbes: Essential components for plants to balance stress
and fitness. *Front Microbiol* **8**: 1–14.
Eller, A.S.D., Sekimoto, K., Gilman, J.B., Kuster, W.C., de Gouw, J.A., Monson, R.K., et al. (2011)
Volatile organic compound emissions from switchgrass cultivars used as biofuel crops.
Atmos Environ **45**: 3333–3337.
Eren, A.M., Esen, O.C., Quince, C., Vineis, J.H., Morrison, H.G., Sogin, M.L., and Delmont, T.O.
(2015) Anvi'o: An advanced analysis and visualization platform for 'omics data. *PeerJ* **2015**:

1–29.

Ferner, E., Rennenberg, H., and Kreuzwieser, J. (2012) Effect of flooding on C metabolism of
flood-tolerant (*Quercus robur*) and non-tolerant (*Fagus sylvatica*) tree species. *Tree Physiol*
32: 135–145.

Gill, U.S., Nandety, R.S., Krom, N., Dai, X., Zhuang, Z., Tang, Y., et al. (2019) Draft genome
sequence resource of switchgrass rust pathogen, puccinia novopanici isolate ard-01.
Phytopathology **109**: 1513–1515.

Gopal, M. and Gupta, A. (2016) Microbiome selection could spur next-generation plant
breeding strategies. *Front Microbiol* **7**: 1971.

Gostinčar, C., Ohm, R.A., Kogej, T., Sonjak, S., Turk, M., Zajc, J., et al. (2014) Genome sequencing
of four Aureobasidium pullulans varieties: Biotechnological potential, stress tolerance, and
description of new species. *BMC Genomics* **15**:.

Gottschlich, L., Geiser, P., Bortfeld-Miller, M., Field, C.M., and Vorholt, J.A. (2019) Complex
general stress response regulation in *Sphingomonas melonis* Fr1 revealed by
transcriptional analyses. *Sci Rep* **9**: 1–13.

Grady, K.L., Sorensen, J.W., Stopnisek, N., Guittar, J., and Shade, A. (2019) Assembly and
seasonality of core phyllosphere microbiota on perennial biofuel crops. *Nat Commun* **10**:.

Hacquard, S., Garrido-Oter, R., González, A., Spaepen, S., Ackermann, G., Lebeis, S., et al. (2015)
Microbiota and host nutrition across plant and animal kingdoms. *Cell Host Microbe* **17**:
603–616.

Hardoim, P.R., van Overbeek, L.S., Berg, G., Pirttilä, A.M., Compant, S., Campisano, A., et al.
(2015) The hidden world within plants: Ecological and evolutionary considerations for
defining functioning of microbial endophytes. *Microbiol Mol Biol Rev* **79**: 293–320.

Haridas, S., Albert, R., Binder, M., Bloem, J., LaButti, K., Salamov, A., et al. (2020) 101
Dothideomycetes genomes: A test case for predicting lifestyles and emergence of
pathogens. *Stud Mycol* **96**: 141–153.

Harty, C.E., Martins, D., Doing, G., Mould, D.L., Clay, M.E., Occhipinti, P., et al. (2019) Ethanol
stimulates trehalose production through a SpoT-DksA-AlgU-dependent pathway in
Pseudomonas aeruginosa. 1–21.

Haskett, T.L., Tkacz, A., and Poole, P.S. (2020) Engineering rhizobacteria for sustainable
agriculture. *ISME J* **15**: 949–964.

Heaton, E.A., Dohleman, F.G., and Long, S.P. (2008) Meeting US biofuel goals with less land: The
potential of Miscanthus. *Glob Chang Biol* **14**: 2000–2014.

Hestrin, R., Lee, M.R., Whitaker, B.K., and Pett-Ridge, J. (2021) The switchgrass microbiome: A
review of structure, function, and taxonomic distribution. *Phytobiomes J* **5**: 14–28.

Julsing, M.K., Rijpkema, M., Woerdenbag, H.J., Quax, W.J., and Kayser, O. (2007) Functional
analysis of genes involved in the biosynthesis of isoprene in *Bacillus subtilis*. *Appl Microbiol*
Biotechnol **75**: 1377–1384.

Kałużna, M., Willems, A., Pothier, J.F., Ruinelli, M., Sobiczewski, P., and Puławska, J. (2016)
Pseudomonas cerasi sp. nov. (non Griffin, 1911) isolated from diseased tissue of cherry.
Syst Appl Microbiol **39**: 370–377.

Kang, D.D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., and Wang, Z. (2019) MetaBAT 2: An
adaptive binning algorithm for robust and efficient genome reconstruction from
metagenome assemblies. *PeerJ* **2019**: 1–13.

Kerchev, P., De Smet, B., Waszczak, C., Messens, J., and Van Breusegem, F. (2015) Redox
strategies for crop improvement. *Antioxidants Redox Signal* **23**: 1186–1205.
El Khawand, M., Crombie, A.T., Johnston, A., Vavlline, D. V., McAuliffe, J.C., Latone, J.A., et al.
(2016) Isolation of isoprene degrading bacteria from soils, development of isoA gene
probes and identification of the active isoprene-degrading soil community using DNA-
stable isotope probing. *Environ Microbiol* **18**: 2743–2753.
Kimmerer, T.W. and Kozlowski, T.T. (1982) Ethylene, ethane, acetaldehyde, and ethanol
production by plants under stress. *Plant Physiol* **69**: 840–847.
Kimmerer, T.W. and MacDonald, R.C. (1987) Acetaldehyde and ethanol biosynthesis in leaves of
plants. *Plant Physiol* **84**: 1204–1209.
Koskella, B. (2020) The phyllosphere. *Curr Biol* **30**: R1143–R1146.
Kuzyakov, Y. and Razavi, B.S. (2019) Rhizosphere size and shape: Temporal dynamics and spatial
stationarity. *Soil Biol Biochem* **135**: 343–360.
Lacerda-Júnior, G. V., Noronha, M.F., Cabral, L., Delforno, T.P., De Sousa, S.T.P., Fernandes-
Júnior, P.I., et al. (2019) Land use and seasonal effects on the soil microbiome of a Brazilian
dry forest. *Front Microbiol* **10**: 1–14.
Lajoie, G., Maglione, R., and Kembel, S.W. (2020) Adaptive matching between phyllosphere
bacteria and their tree hosts in a neotropical forest. *Microbiome* **8**: 1–10.
Langholtz, M., Stokes, B., and Eaton, L. (2016) 2016 billion-ton report: Advancing domestic
resources for a thriving bioeconomy (Executive Summary). *Ind Biotechnol* **12**: 282–289.
Langmead, B. and Salzberg, S.L. (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9: 357–359.
Laskowska, E., Dorota Kuczyńska-Wiśnik, ·, and Kuczyńska-Wiśnik, D. (2020) New insight into
the mechanisms protecting bacteria during desiccation. *Curr Genet* **66**: 313–318.
Lee, M.D. (2019) GToTree: A user-friendly workflow for phylogenomics. *Bioinformatics* **35**:
4162–4164.
Levy, A., Conway, J.M., Dangl, J.L., and Woyke, T. (2018) Elucidating bacterial gene functions in
the plant microbiome. *Cell Host Microbe* **24**: 475–485.
Li, D., Liu, C.M., Luo, R., Sadakane, K., and Lam, T.W. (2015) MEGAHIT: An ultra-fast single-node
solution for large and complex metagenomics assembly via succinct de Bruijn graph.
Bioinformatics **31**: 1674–1676.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009) The Sequence
Alignment / Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.
Lindow, S.E. and Brandl, M.T. (2003) Microbiology of the phyllosphere. *Appl Environ Microbiol*
69: 1875–1883.
Liu, Y., Zhang, X., Tran, H., Shan, L., Kim, J., Childs, K., et al. (2015) Assessment of drought
tolerance of 49 switchgrass (Panicum virgatum) genotypes using physiological and
morphological parameters. *Biotechnol Biofuels* **8**: 1–18.
Lucke, M., Correa, M.G., and Levy, A. (2020) The role of secretion systems, effectors, and
secondary metabolites of beneficial rhizobia in interactions with plants and microbes.
Front Plant Sci **11**:.
Lundberg, D.S., Lebeis, S.L., Paredes, S.H., Yourstone, S., Gehring, J., Malfatti, S., et al. (2012)
Defining the core *Arabidopsis thaliana* root microbiome. *Nature* **488**: 86–90.
Ma, L., Rocha, F.I., Lee, J., Choi, J., Tejera, M., Sooksa-Nguan, T., et al. (2021) The impact of

stand age and fertilization on the soil microbiome of *Miscanthus × giganteus*. *Phytobiomes*
J **5**: 51–59.

Marzano, S.Y.L. and Domier, L.L. (2016) Novel mycoviruses discovered from
metatranscriptomics survey of soybean phyllosphere phytobiomes. *Virus Res* **213**: 332–
342.

McGenity, T.J., Crombie, A.T., and Murrell, J.C. (2018) Microbial cycling of isoprene, the most
abundantly produced biological volatile organic compound on Earth. *ISME J* **12**: 931–941.

Morrison, E.C., Drewer, J., and Heal, M.R. (2016) A comparison of isoprene and monoterpene
emission rates from the perennial bioenergy crops short-rotation coppice willow and
Miscanthus and the annual arable crops wheat and oilseed rape. *GCB Bioenergy* **8**: 211–
225.

Müller, D.B., Vogel, C., Bai, Y., and Vorholt, J.A. (2016) The Plant Microbiota: Systems-Level
Insights and Perspectives. *Annu Rev Genet* **50**: 120215–034952.

Navarro-Muñoz, J.C., Selem-Mojica, N., Mullowney, M.W., Kautsar, S.A., Tryon, J.H., Parkinson,
E.I., et al. (2020) A computational framework to explore large-scale biosynthetic diversity.
Nat Chem Biol **16**: 60–68.

Nayfach, S., Shi, Z.J., Seshadri, R., Pollard, K.S., and Kyrpides, N.C. (2019) New insights from
uncultivated genomes of the global human gut microbiome. *Nature* **568**: 505–510.

Nowicka, B. and Kruk, J. (2010) Occurrence, biosynthesis and function of isoprenoid quinones.
Biochim Biophys Acta - Bioenerg **1797**: 1587–1605.

Palmer, J.L., Hilton, S., Picot, E., Bending, G.D., and Schäfer, H. (2021) Tree phyllospheres are a
habitat for diverse populations of CO-oxidizing bacteria. *Environ Microbiol* **23**: 6309–6327.

Parks, D.H., Chuvochina, M., Waite, D.W., Rinke, C., Skarshewski, A., Chaumeil, P.A., and
Hugenholtz, P. (2018) A standardized bacterial taxonomy based on genome phylogeny
substantially revises the tree of life. *Nat Biotechnol* **36**: 996.

Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., and Tyson, G.W. (2015) CheckM:
Assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**: 1043–1055.

Peñuelas, J. and Terradas, J. (2014) The foliar microbiome. *Trends Plant Sci* **19**: 278–280.

Quinlan, A.R. and Hall, I.M. (2010) BEDTools : a flexible suite of utilities for comparing genomic
features. **26**: 841–842.

Rastogi, G., Coaker, G.L., and Leveau, J.H.J. (2013) New insights into the structure and function
of phyllosphere microbiota through high-throughput molecular approaches. *FEMS*
Microbiol Lett **348**: 1–10.

Remus-Emsermann, M.N.P. and Schlechter, R.O. (2018) Phyllosphere microbiology: at the
interface between microbial individuals and the plant host. *New Phytol* **218**: 1327–1333.

Robertson, G.P., Hamilton, S.K., Barham, B.L., Dale, B.E., Izaurralde, R.C., Jackson, R.D., et al.
(2017) Cellulosic biofuel contributions to a sustainable energy future: Choices and
outcomes. *Science (80-)* **356**: 1–9.

Roley, S.S., Duncan, D.S., Liang, D., Garoutte, A., Jackson, R.D., Tiedje, J.M., and Robertson, G.P.
(2018) Associative nitrogen fixation (ANF) across a nitrogen input gradient. *PLoS One* **13**:
1–19.

Roley, S.S., Xue, C., Hamilton, S.K., Tiedje, J.M., and Robertson, G.P. (2019) Isotopic evidence for
episodic nitrogen fixation in switchgrass (*Panicum virgatum* L.). *Soil Biol Biochem* **129**: 90–

98.

Shade, A. and Stopnisek, N. (2019) Abundance-occupancy distributions to prioritize plant core
microbiome membership. *Curr Opin Microbiol* **49**: 50–58.

Shaffer, M., Borton, M.A., McGivern, B.B., Zayed, A.A., La Rosa, S.L. 0003 3527 8101, Solden,
L.M., et al. (2020) DRAM for distilling microbial metabolism to automate the curation of
microbiome function. *Nucleic Acids Res* **48**: 8883–8900.

Sharkey, T.D., Loreto, F., and Delwiche, C. (1991) High carbon dioxide and sun/shade effects on
isoprene emission from oak and aspen tree leaves. *Plant, Cell Environ* **14**: 333–338.

Sharkey, T.D., Wiberley, A.E., and Donohue, A.R. (2008) Isoprene emission from plants: Why
and how. *Ann Bot* **101**: 5–18.

Sharkey, T.D. and Yeh, S. (2001) Isoprene emission from plants. *Plant Mol Biol* **52**: 407–436.

Singer, E., Bonnette, J., Kenaley, S.C., Woyke, T., and Juenger, T.E. (2019) Plant compartment
and genetic variation drive microbiome composition in switchgrass roots. *Environ*
Microbiol Rep **11**: 185–195.

Sivy, T.L., Shirk, M.C., and Fall, R. (2002) Isoprene synthase activity parallels fluctuations of
isoprene release during growth of *Bacillus subtilis*. *Biochem Biophys Res Commun* **294**: 71–
75.

Spaepen, S., Vanderleyden, J., and Remans, R. (2007) Indole-3-acetic acid in microbial and
microorganism-plant signaling. *FEMS Microbiol Rev* **31**: 425–448.

Stopnisek, N. and Shade, A. (2021) Persistent microbiome members in the common bean
rhizosphere: an integrated analysis of space, time, and plant genotype. *ISME J.*

Suzuki, R. and Shimodaira, H. (2006) Pvclust: An R package for assessing the uncertainty in
hierarchical clustering. *Bioinformatics* **22**: 1540–1542.

Suzuki, Y., Makino, A., and Mae, T. (2001) An efficient method for extraction of RNA from rice
leaves at different ages using benzyl chloride. *J Exp Bot* **52**: 1575–1579.

Swan, B.K., Tupper, B., Sczyrba, A., Lauro, F.M., Martinez-Garcia, M., Gonźalez, J.M., et al.
(2013) Prevalent genome streamlining and latitudinal divergence of planktonic bacteria in
the surface ocean. *Proc Natl Acad Sci U S A* **110**: 11463–11468.

Toju, H., Peay, K.G., Yamamichi, M., Narisawa, K., Hiruma, K., Naito, K., et al. (2018) Core
microbiomes for sustainable agroecosystems. *Nat Plants* **4**: 247–257.

Trivedi, P., Leach, J.E., Tringe, S.G., Sa, T., and Singh, B.K. (2020) Plant–microbiome interactions:
from community assembly to plant health. *Nat Rev Microbiol* **18**: 607–621.

Tseng, T.T., Tyler, B.M., and Setubal, J.C. (2009) Protein secretion systems in bacterial-host
associations, and their description in the Gene Ontology. *BMC Microbiol* **9**: 1–9.

Urrejola, C., Alcorta, J., Salas, L., Vásquez, M., Polz, M.F., Vicuña, R., and Díez, B. (2019)
Genomic features for desiccation tolerance and sugar biosynthesis in the extremophile
gloeocapsopsis sp. UTEX B3054. *Front Microbiol* **10**: 1–11.

Valenzuela-Soto, E.M. and Figueroa-Soto, C.G. (2019) Biosynthesis and Degradation of Glycine
Betaine and Its Potential to Control Plant Growth and Development. In *Osmoprotectant-*
Mediated Abiotic Stress Tolerance in Plants. Anwar Hossain, M., Kumar, Vi., Burritt, D.J.,
Fujita, M., and Makela, P.S.A. (eds). Springer, pp. 241–256.

Vorholt, J.A. (2012) Microbial life in the phyllosphere. *Nat Rev Microbiol* **10**: 828–840.

Wang, N.R. and Haney, C.H. (2020) Harnessing the genetic potential of the plant microbiome.
Biochem (Lond) **42**: 20–25.

Wingler, A., Delatte, T.L., O'Hara, L.E., Primavesi, L.F., Jhurreea, D., Paul, M.J., and
Schluepmann, H. (2012) Trehalose 6-phosphate is required for the onset of leaf
senescence associated with high carbon availability. *Plant Physiol* **158**: 1241–1251.
Xu, L., Naylor, D., Dong, Z., Simmons, T., Pierroz, G., Hixson, K.K., et al. (2018) Drought delays
development of the sorghum root microbiome and enriches for monoderm bacteria. *Proc*
Natl Acad Sci **115**: E4284–E4293.
Zhalnina, K., Louie, K.B., Hao, Z., Mansoori, N., da Rocha, U.N., Shi, S., et al. (2018) Dynamic root
exudate chemistry and microbial substrate preferences drive patterns in rhizosphere
microbial community assembly. *Nat Microbiol* **3**: 470.
Zimmermann, J., Kaleta, C., and Waschina, S. (2021) Gapseq: informed prediction of bacterial
metabolic pathways and reconstruction of accurate metabolic models. *Genome Biol* **22**: 1–
35.
Zou, H., Chen, N., Shi, M., Xian, M., Song, Y., and Liu, J. (2016) The metabolism and
biotechnological application of betaine in microorganism. *Appl Microbiol Biotechnol* **100**:
3865–3876.
Zuo, Z., Weraduwage, S.M., Lantz, A.T., Sanchez, L.M., Weise, S.E., Wang, J., et al. (2019)
Isoprene acts as a signaling molecule in gene networks important for stress responses and
plant growth. *Plant Physiol* **180**: 124–152.
978
979

Figures and Tables

Figures

Figure 1. Phyllosphere microbiome field sampling strategy at the Great Lakes Bioenergy
Research Center Bioenergy Cropping System Experiment (BCSE) in 2016 and 2017. (A) The
study site is at Kellogg Biological Station, a Long-Term Ecological Research site focused on
agroecosystems located in southwest Michigan. (B) Four replicate randomized cropping system
blocks from the BCSE were sampled at each time point for switchgrass and/or miscanthus, and
within each plot there was a fertilized main plot and unfertilized subplot sampled. (C) In 2016,
both switchgrass and miscanthus were sampled, and in 2017 only switchgrass was sampled.
Switchgrass leaves were flash frozen in liquid nitrogen for RNA extraction and
metatranscriptome analysis at a subset of time points in 2016 and at all points in 2017.

Figure 2. Overview of bioinformatic processing of the metagenome (green solid arrows) and
metatranscriptome (red solid arrows) datasets. Switchgrass reads are shown in light green and
miscanthus is dark green. The solid grey arrow represents the common analysis step for both
datasets. Figure was made with Lucidchart.

Figure 3. Summary of leaf-associated MAGs. **(A)** Abundance and occupancy of genomes
assembled and binned from switchgrass and miscanthus phyllosphere metagenomes. Quality
and contamination assessment were determined using checkM. 40 focal MAGs were selected
(B) Taxonomy of focal MAGs, as annotated with GTDB-tk, and taxonomic overlap with the
persistent taxa detected in our previous 16S rRNA gene amplicon survey.

1004

Figure 4. Seasonal patterns of the 40 focal MAG metagenome and metatranscriptome read
recruitment. MAG abundances for: (**A**) Miscanthus 2016 metagenomes (metaG); (**B**)
Switchgrass 2016 metagenomes; (**C**) Switchgrass 2017 metagenomes; (**D**) Switchgrass 2016
metatranscriptomes (metaT) and (E) Switchgrass 2017 metatranscriptomes. Abundances of
metagenome contigs were estimated with the median basepair of read recruitment divided by
the average median basepair coverage of housekeeping genes. Abundances of
metatranscriptome ORFs were estimated based on the median basepair coverage of all reads
mapped to ORFs and divided by median basepair coverage of housekeeping genes. The same
dendrogram is applied to all panel, and it is the result of hierarchical clustering (see Figure S2)
of metatranscriptome diversity and abundances in switchgrass.

1015

1016   **Figure 5**. 2016 (circle) and 2017 (triangle) switchgrass leaf transcript dynamics of KEGG
1017   metabolism classifications to the 40 focal MAGs. The y-axis is scaled for each classification.

1018

Figure 6. Key functional gene pathways detected in the 40 focal MAGs and their activities
(mapped transcripts) during the 2016-2017 switchgrass growing season. Functional gene
pathways were curated using antiSMASH for biosynthetic gene clusters, gapseq for general
metabolic pathways, and manual selection for plant-associative functions reported in the
literature. Pathways that were discovered in the MAGs but not detected in the transcripts are

represented by open circles, and pathways detected in the MAGs and mapped by transcripts
are represented by filled circles. Colors categorize different functional groups of pathways.
1026
Figure 7. **(A).** 2016 (circle) and 2017 (triangle) switchgrass leaf transcript dynamics of KEGG
metabolism classifications associated with terpene metabolism. **(B).** Transcripts in MAGs
associated with terminal enzymes in the non-mevalonate isoprene biosynthesis, *gcpE* and *lytB*.
MAG IDs include predicted taxonomy at the genus level: Methyl = *Methylobacterium*, Frigor =
Frigoribacterium, Pseudokineo = *Pseudokineococcus*, Microbac = *Microbacterium*, Amnibac =
Amnibacterium, Hymeno = *Hymenobacter,* Sphingo = *Sphingomonas,* and Pseudo =
Pseudomonas.
1034
1035   **Figure 8**. Detection of 36 focal MAGs (out of 40) in publicly available metagenomes and
1036   metatranscriptomes of bioenergy grasses.
1037
1038

1039   Tables
1040
1041   **Table 1**. Summary of RNA and DNA samples that returned reads and passed Illumina standard
1042   quality control at the Joint Genome Institute. Total collected samples submitted for sequencing
1043   is provided first, and the number of quality sequencing datasets returned for analysis is given in
1044   parentheses.

	2016		2017	
	Metagenome	Metatranscriptome	Metagenome	Metatranscriptome
Switchgrass	8 timepoints (64/64 successful)	3 (22/24 successful)	7 (56/56 successful)	7 (56/56 successful)
Miscanthus	9 (72/72 successful)	Not assessed	Not assessed	Not assessed

1045
1046
1047

1048 **Table 2.** Functional roles that exhibited strong seasonality with transcript enrichment by more
1049 than 20-fold in the late season than early. These roles were consistently detected among
1050 phyllosphere focal MAGs (39/40 detections).
1051

Functional role	Major classification(s)	No. features	Late:early ratio
short chain dehydrogenase [PF00106.26]; Enoyl-(Acyl carrier protein) reductase [PF13561.7]; KR domain [PF08659.11]	Carbohydrate metabolism, Lipid metabolism, nucleotide metabolism, metabolism of cofactors and vitamins	4	53.2
Molybdopterin oxidoreductase [PF00384.23], Molydopterin dinucleotide binding domain [PF01568.22]	Carbohydrate metabolism, metabolism of cofactors and vitamins, Energy metabolism, Signal transduction	4	29.7
Polyketide cyclase / dehydrase and lipid transport [PF10604.10]	Energy metabolism	1	23.6

1052

Datasets
Dataset 1. Metadata for metagenome samples according to MIMs standards.
Dataset 2. Metadata for metatranscriptome samples according to MIMs standards.
Dataset 3. Excel file. *Sheet 1.* Metadata for selected focal 40 metagenome-assembled genomes
(>50% complete, <10% contamination) according to MiMAG standards *Sheet 2.* Other MAGs
(non-focal) assembled from switchgrass or miscanthus metagenomes.
Dataset 4. Excel file. *Sheet 1.* Summary of metagenome reads mapped to the focal MAGs.
Sheet 2. Summary of metatranscriptome reads mapped to the focal MAGs. *Sheet 3*. Total reads
mapped from each metagenome. *Sheet 4.* Total reads mapped from each metatranscriptome.
Dataset 5. Public metagenomes used for comparison to the switchgrass and miscanthus focal
MAGs.
Dataset 6. Core KEGG functions identified in at least 39 focal MAGs and their estimated
abundances in early and late season.
1066
1067

Supplementary Figures
Figure S1. Contaminating plant or fungal sequences. Proportion of sequencing in metagenomes
originating from miscanthus and switchgrass phyllospheres associated with miscanthus and
switchgrass host genomes (A) and prevalent fungal genomes (B, Table S1). Proportion of
sequencing in metatreanscriptomes originating from and switchgrass phyllospheres associated
with switchgrass host genomes (C) and fungal genomes (D, Table S1).
1074
1075   **Figure S2.** Hierarchical clustering identified five clusters of MAGS to identify MAG populations
1076   with coherent seasonal activity dynamics. Clustering was based on metatranscriptome diversity
1077   and abundances.
1078
1079   **Figure S3.** Mean transcript dynamics by month and year for each MAG cluster. Filled circles
1080   indicate that identification of transcripts, and empty circles indicate no detection.
1081
1082   **Figure S4**. Summary of transcript seasonality of phyllosphere open reading frames (ORFs) that
1083   could be annotated as KEGG functional roles and were consistently detected among focal MAGs
1084   (at least 39/40 detections). Ratios are late-to-early normalized transcript abundances on
1085   MAGs.
1086
1087   **Figure S5**. 2016 (circle) and 2017 (triangle) switchgrass leaf transcript dynamics of KEGG
1088   metabolism classifications associated with terpenoid backbone biosynthesis.
1089
1090
1091
1092

1093

1094   **Supplementary Tables**

1095

1096   **Table S1**. Fungal genomes used for filtering metagenome reads to remove eukaryotic
1097   contamination. Genomes were selected to use for filtering based on the taxonomic identities
1098   of prevalent fungal taxa detected in our previous ITS2 amplicon survey that was conducted at
1099   the same location (Bowsher *et al.*, 2020).

1100

Fungal Genome	JGI Project ID (*) or GenBank accession (**)	Reference
Trichoderma harzianum	403727*	(Druzhinina et al., 2018)
Pleomassaria siparia	1011309*	(Haridas et al., 2020)
Aureobasidium pullulans	403628*	(Gostinčar et al., 2014)
Alternaria alternata	1103683*	NA
Didymella zeae-maydis	https://genome.jgi.doe.gov/portal/pages/dynamicOrganismDownload.jsf?organism=Didma1	NA
Sporobolomyces roseus	16892*	NA
Puccinia novopanici	GCA_004348175.1**	(Gill et al., 2019)

1101
1102
1103
1104

1105   **Table S2.**

1106

1107   Summary of the nine genes involved the *Bacillus subtilus* isoprene biosynthesis pathway that
1108   were detected among focal MAG contigs. Except for *ypgA*, these genes were directly linked to
1109   isoprene accumulation by (Julsing *et al.*, 2007).

1110

Gene	dxr	dxs	gcpE	idi	lytB	yacM	ychB	ygbB	yqiD
NCBI Acc. No.	NP_389537.2	NP_390307.1	NP_390386.1	BAB32625.1	NP_390395.2	NP_387971.1	NP_387927.1	NP_387972.1	NP_390308.2
M94	+	+	+	+	+		+	+	+
S30	+	+	+	+	+		+	+	+
M105	+	+	+		+		+	+	+
M12	+	+	+	+	+			+	+
M32	+	+	+		+		+	+	+
M52	+	+	+	+	+			+	+
M67	+	+	+		+	+	+		+
M77	+	+	+		+		+	+	+
S117		+	+	+	+		+	+	+
S28	+	+	+		+		+	+	+
S56	+	+	+	+			+	+	+
M100	+	+	+		+	+			+
M109	+	+	+		+			+	+
M1	+	+			+		+	+	+
M21	+	+	+		+		+		+
M44	+	+		+	+		+		+
M47		+	+		+	+		+	+
M60		+	+		+		+	+	+
M66	+	+	+				+	+	+
M86	+	+	+		+		+		+
M8		+	+		+		+	+	+
M99	+	+			+	+		+	+
M9	+	+	+		+			+	+
S27	+	+	+		+			+	+
S61	+	+	+		+			+	+
S71	+	+			+		+	+	+
S74	+	+	+		+	+			+
M102	+	+	+					+	+
M55	+				+		+	+	+
S80		+	+		+	+			+
M111			+	+	+				+
M87	+	+						+	+
S120		+	+		+		+		
S29		+					+	+	+
S50	+	+	+				+		
S36		+	+		+				
M17		+							+
M35		+							+
S8		+							+
S9		+							+

1111
1112
1113
1114

Figure 1

Figure 1. Phyllosphere microbiome field sampling strategy at the Great Lakes Bioenergy Research Center Bioenergy Cropping System Experiment (BCSE) in 2016 and 2017. (A) The study site is at Kellogg Biological Station, a Long-Term Ecological Research site focused on agroecosystems located in southwest Michigan. (B) Four replicate randomized cropping system blocks from the BCSE were sampled at each time point for switchgrass and/or miscanthus, and within each plot there was a fertilized main plot and unfertilized subplot sampled. (C) In 2016, both switchgrass and miscanthus were sampled, and in 2017 only switchgrass was sampled. Switchgrass leaves were flash frozen in liquid nitrogen for RNA extraction and metatranscriptome analysis at a subset of time points in 2016 and at all points in 2017.

Figure 2

Figure 2. Overview of bioinformatic processing of the metagenome (green solid arrows) and metatranscriptome (red solid arrows) datasets. Switchgrass reads are shown in light green and miscanthus is dark green. The solid grey arrow represents the common analysis step for both data types. Figure was made with Lucidchart.

Figure 3

Figure 3. Summary of leaf-associated MAGs. **(A)** Abundance and occupancy of genomes assembled and binned from switchgrass and miscanthus phyllosphere metagenomes. Quality and contamination assessment were determined using checkM. 40 focal MAGs were selected **(B)** Taxonomy of focal MAGs, as annotated with GTDB-tk, and taxonomic overlap with the persistent taxa detected in our previous 16S rRNA gene amplicon survey.

Figure 4

Figure 4. Seasonal patterns of the 40 focal MAG metagenome and metatranscriptome read recruitment. MAG abundances for: **(A)** Miscanthus 2016 metagenomes; **(B)** Switchgrass 2016 metagenomes (metaG); **(C)** Switchgrass 2017 metagenomes; **(D)** Switchgrass 2016 metatranscriptomes (metaT) and (E) Switchgrass 2017 metatranscriptomes. Abundances of metagenome contigs were estimated with the median basepair of read recruitment divided by the average median basepair coverage of housekeeping genes. Abundances of metatranscriptome ORFs were estimated based on the median basepair coverage of all reads mapped to ORFs and divided by median basepair coverage of housekeeping genes. The same dendrogram is applied to all panel, and it is the result of hierarchical clustering (see Figure S2) of metatranscriptome diversity and abundances in switchgrass.

Figure 5

Figure 5. 2016 (circle) and 2017 (triangle) switchgrass leaf transcript dynamics of KEGG metabolism classifications to the 40 focal MAGs. The y-axis is scaled for each classification. Coverage was estimated as the average median base pair coverage for each gene. All coverage estimates were normalized by estimated coverage of housekeeping genes in each sample.

Figure 6

Figure 6. Key functional gene pathways detected in the 40 focal MAGs and their activities (mapped transcripts) during the 2016-2017 switchgrass growing season. Functional gene pathways were curated using antiSMASH for biosynthetic gene clusters, gapseq

for general metabolic pathways, and manual selection for plant-associative functions reported in the literature. Pathways that were discovered in the MAGs but not detected in the transcripts are represented by open circles, and pathways detected in the MAGs and mapped by transcripts are represented by filled circles. Colors categorize different functional groups of pathways.

Figure 7

Figure 7. **(A).** 2016 (circle) and 2017 (triangle) switchgrass leaf transcript dynamics of KEGG metabolism classifications associated with terpene metabolism. **(B).** Transcripts in MAGs associated with terminal enzymes in the non-mevalonate isoprene biosynthesis, *gcpE* and *lytB*. MAG IDs include predicted taxonomy at the genus level: Methyl = *Methylobacterium*, Frigor = *Frigoribacterium*, Pseudokineo = *Pseudokineococcus*, Microbac = *Microbacterium*, Amnibac = *Amnibacterium*, Hymeno = *Hymenobacter*, Sphingo = *Sphingomonas,* and Pseudo = *Pseudomonas*.

Figure 8

Figure 8. Detection of MAGs in publicly available metagenomes and metatranscriptomes of bioenergy grasses.