Sox8 is sufficient to reprogram ectoderm into ear vesicles and associated neurons

Ailin Leticia Buzzi[1], Jingchen Chen[1,3,$], Alexandre Thiery[1,$], Julien Delile[2], Andrea Streit[1*]

[1] Centre for Craniofacial and Regenerative Biology, King's College London, London SE1 9RT, UK

[2] The Francis Crick Institute, London NW1 1AT, UK

[$] These authors contributed equally.

*Corresponding author: andrea.streit@kcl.ac.uk

[3] current address: Maijing Gene Diagnostics, Guangzhou 510030, China

The vertebrate inner ear arises from a pool of progenitors with the potential to give rise to all

the sense organs and cranial ganglia of the head[1-6]. Here we explore the molecular

mechanisms that control ear specification from these precursors. Using a multi-omics

approach combined with loss-of-function experiments we identify a core transcriptional circuit

that imparts ear identity, along with non-coding elements that integrate this information. This

analysis places the transcription factor Sox8 at the top of the ear determination network.

Introducing Sox8 into cranial ectoderm not only converts non-ear cells into ear progenitors,

but also activates the cellular programmes for ear morphogenesis and neurogenesis. Thus,

Sox8 emerges as a master regulator of ear identity and may be a key factor for sense organ

cell reprogramming.

Master regulators have the unique ability to reprogram cells from other cell types. They activate a set

of transcription factors that in turn control the expression of proteins required for cell-specific functions.

The classic example is the transcription factor MyoD, which can convert fibroblasts into myogenic

cells[7]. From insects to vertebrates, the transcription factor Pax6 induces ectopic eyes when mis-

expressed in non-eye cells[8], however master regulators for other sense organs have not yet been

discovered. Here we set out to identify such factors for the organ of hearing and balance, the

vertebrate inner ear.

During development, the inner ear arises from the posterior border of the cranial neural plate.

Here, precursors for neural, neural crest, ear and sensory ganglia are initially intermingled, but

segregate over time[9,10]. FGF signalling from adjacent tissues induces otic-epibranchial progenitors

(OEPs), which subsequently converge to distinct territories: the otic and epibranchial placodes[11-15]

(Fig. 1a, e). While epibranchial cells produce sensory ganglia, the otic placode invaginates to form a

vesicle, which is then transformed into the inner ear, containing many specialised cell types and

associated neurons. Using transcriptional profiling, we recently defined the first gene regulatory

network for early ear development[16]. However, the key factors that impart ear character remain

unknown as do the non-coding regions that integrate the process. Here we use single cell RNA-

sequencing (scRNAseq) and pseudo-time analysis to predict transcriptional changes as progenitors

undergo ear commitment. Using epigenomic profiling we also determine ear enhancers and their

upstream regulators. To ensure transparency and reproducibility, we have developed a fully

integrated workflow of the data analysis in Nextflow (available at https://alexthiery.github.io/otic-

reprogramming/). Together with functional experiments, we identify a minimal transcriptional circuit

that defines ear identity comprising Sox8, Pax2 and Lmx1a, with Sox8 at the top of the hierarchy.

Introduction of Sox8 alone into cranial ectoderm is sufficient to trigger the ear programme, to initiate

ear morphogenesis by forming ear vesicles and to activate neurogenesis. Thus, using a multi-omics

approach we have uncovered Sox8 as a master regulator of ear fate, which may in the future allow

reprograming of specific sensory cell types.

Dynamic changes in gene expression characterise the transition from progenitor to otic commitment

To unravel the genetic hierarchy that controls how otic and epibranchial cells diverge, we first characterised the transcriptional profile of cells committed to each fate[2] (Fig1. a-d, Extended Data Fig. 1a, b). To label each cell population we used two enhancers driving EGFP, the novel otic Lmx1aE1 enhancer (see below) and the epibranchial enhancer Sox3U3[17]. Reporter constructs were electroporated into the OEP territory of somite stage (ss) 3-5 embryos. At ss18-21, EGFP+ cells were isolated by fluorescence-activated cell sorting (FACS) and processed for RNA-seq. Differential expression analysis identifies 103 and 319 genes upregulated in Lmx1aE1-EGFP and Sox3U3-EGFP expressing cells, respectively (log2FC > 1.5; adjusted p-value < 0.05; Fig. 1d; Extended Data Fig. 1c, d and Supplementary Table1). This allows us to define the transcriptional states of definitive otic and epibranchial cells.

Next, we investigated the transcriptional changes which take place as OEPs transition from a common progenitor population to definitive otic and epibranchial cells. *Pax2* is expressed throughout this time-window in both cell populations (Extended Data Fig.1e, f)[2,9]. Using a Pax2E1-EGFP reporter, we isolated single cells by FACS from consecutive stages of ear specification: OEP (ss8-9), early-placode (ss11-12) and late-placode stage (ss14-15) and processed them for single-cell-RNA-seq (scRNAseq; Fig. 1e, f). To characterise cellular diversity, we first looked at groups of genes co-expressed across the dataset (gene modules). These gene modules were identified in an unbiased manner through hierarchical clustering of gene-gene Spearman correlation coefficient[18].Gene modules of interest were selected based on the presence of well-characterized makers for placodal, neural and neural-crest cells including the new otic or epibranchial genes identified above (Fig.1d). Initial cell clustering defines five major clusters (Extended Data Fig. 2a) to which we assigned identities using known markers (Fig. 1g-h, Extended Data Fig. 2c-f). Clusters C1 and C2 represent *Pax2+/Six1+* cells expressing high levels of OEP and placodal makers (Fig. 1h, i; Extended Data Fig. 2c). Surprisingly, we also find two clusters with relatively few *Pax2+* cells: one containing neural-like cells (C4; *Sox21+*) and another containing neural crest-like cells (C5; *Pax7+*) (Extended Data Fig. 2a, d, e). Since OEPs are mixed with neural and neural crest cells at early stages[9], this observation

suggests that while cells in clusters C4/5 initially activate Pax2E1-EGFP they subsequently

downregulate enhancer activity and *Pax2* expression.

To investigate the transcriptional dynamics accompanying otic and epibranchial fate

decisions, we subset the placodal clusters (C1/2 in Fig. 1; Extended Data Fig. 2a). Re-clustering these

cells using gene modules containing otic and epibranchial genes we obtained five placodal clusters

(PC1-5; Fig. 1m, j, k; Supplementary Table 2). PC1 largely consists of only OEPs (ss8-9 cells), while

the other clusters contain cells from both early and late placode stages (Fig.1m, j, k). Indeed, PC1 is

characterised by the expression of OEP genes (GM1-3; Fig. 1m), as well as sharing genes with the

otic module GM5/6 and the epibranchial module GM7-9. In contrast, PC2/3 and PC4/5 are

transcriptionally distinct from each other with profiles akin to otic (GM5/6) and epibranchial (GM7-9)

cells, respectively. To explore the relationship between different cell clusters we organised cells along

pseudo-time using Monocle2[19]. This analysis predicts that OEPs gradually split into one otic and one

epibranchial branch, each composed of early and late placodal cells (Fig.1n, o). Calculating RNA-

velocity independently[20] and embedding the corresponding vector field onto the Monocle2 trajectory

validates the directionality of predicted cell state transitions (Fig.1p).

To explore dynamic changes of gene expression accompanying these inferred trajectories,

we used Branch-Expression-Analysis-Modelling (BEAM)[19]. This identified groups of transcription

factors expressed in OEPs prior to the branching point, which subsequently segregate into either the

otic (e.g. *Sox8, Lmx1a*) or the epibranchial (e.g. *Foxi3, Tfap2e*) branch (Fig. 2a, Extended Data Fig.

3a). To quantify the changes in co-expression of otic and epibranchial genes, we assessed the

proportion of co-expressing cells before and after the branching point. A two-tailed Wilcoxon rank sum

test reveals significantly more cells co-expressing otic and epibranchial markers in OEPs than in

epibranchial (W=214, p=0.0013) and otic cells (W=235, p<0.0001) after the branching point (Fig. 2b,

c). Quantification of gene expression in the monocle trajectories and by *in situ* hybridization chain

reaction (HCR)[21] confirms that otic (*Sox8, Lmx1a*) and epibranchial (*Foxi3, Tfap2e*) transcripts overlap

at ss8-9, and their expression resolves as both placodes are firmly established (Fig. 2d-i, Extended

Data Fig. 3b-f).

Together, these results identify groups of transcription factors whose expression changes over

time as otic and epibranchial precursors segregate. Some of these may play a key role in imparting

ear identity.

109   Epigenomic profiling uncovers new regulatory elements and motifs in ear precursors

Transcription factors controlling cell fate choice regulate cell-specific downstream targets by

interacting with tissue specific cis-regulatory enhancer elements. Active enhancers are regions of

open chromatin flanked by nucleosomes enriched for histone 3 lysine-27 acetylation (H3K27ac), while

actively transcribed genes are marked by H3K4me3[22-25]. We therefore profiled ss8-9 OEPs by

ChIPseq for H3K27ac, H3K4me3 and the repressive mark H3K27me3 and determined chromatin

accessibility by ATACseq. Overlapping H3K27ac and ATACseq data identifies 10969 regions with

depleted H3K27me3 marks; average profiles show bimodal H3K27ac read distribution surrounding

ATACseq peaks (Extended Data Fig. 4a). Of these just over 70% are intergenic or intronic

representing putative enhancers (8316), while the remaining are close to transcription start sites (TSS;

Extended Data Fig. 3b). We associated each putative enhancer to the nearest TSS of protein coding

genes; GO term analysis of the corresponding genes returns MAP-kinase, Wnt and Notch signalling

known to mediate ear induction, development and neurogenesis (Extended Data Fig. 4c)[13,15]. We

selected putative enhancers in the vicinity of ear-enriched genes to assess their activity *in vivo*,

generated EGFP reporters and co-electroporated them with ubiquitously expressed mCherry into

head-fold-stage chick embryos. RT-PCR-based assays[26] (not shown) and fluorescence microscopy

confirm enhancer activity in ear progenitors and otic placodes (Fig. 3a, b; Extended Data Fig. 5-7).

Next, we performed motif enrichment analysis of all ear enhancers to identify upstream regulators

that may act as otic determinants. This reveals an over-representation of binding sites for Sox, TEAD

and Six family members and for Tfap2a (Fig. 3c; Extended Data Fig. 4d). Of these, Tfap2a and Six

proteins have previously been implicated in cranial placode development[27-31], confirming that this

strategy can identify relevant regulatory factors.

We also exploited the idea that cell identity genes may be regulated by super-enhancers

characterised by broad H3K27ac, while their gene bodies harbour broad H3K4me3[32-34]. Focusing on

OEP factors segregating to the ear lineage (Fig. 2a) that are also prominent in the otic GRN[16], we find

that the Sox8 locus is marked with H3K4me3, while enhancers close to Lmx1a, Zbtb16 and Sox13

show broad H3K27ac marks (Fig. 3a, Extended data 4e, 6a, 7a). Together, this analysis points to

Lmx1a, Zbtb16 and members of the Sox transcription factor family as potential key regulators of ear

identity.

Sox8, Lmx1a and Pax2 are core components of the ear determination network

Our transcriptome analysis identifies several Sox genes enriched in otic progenitors. Of these, *Sox3*

and *Sox13* are expressed prior to ear specification, while *Sox9* and *-10* are activated later[16]. These

genes are therefore unlikely to initiate the otic programme. In contrast, the SoxE group factor *Sox8* is

highly enriched in OEPs prior to segregating to otic cells (Fig. 3d). *In situ* hybridisation reveals that

Sox8 begins to be expressed at 3ss, followed shortly thereafter by *Pax2*, *Zbtb16*, and *Lmx1a*, while

Foxg1, *Soho1* and *Sox10* are activated later (Fig. 3d, Extended Data Fig. 5a).

146   To explore the regulatory interactions between the earliest OEP transcription factors, we

systematically knocked down each one in turn and assayed the expression of all others as well as

that of *Foxg1* and *Soho1* as a readout for otic identity using *in situ* hybridisation and RT-qPCR (Fig.

2e; Extended Data Fig. 5b, c). Control or antisense oligonucleotides targeting *Sox8, Pax2, Zbtb16* or

Lmx1a were electroporated into future OEPs of head-fold-stage chick embryos and gene expression

analysed at OEP stage (ss8-9). We find that *Sox8* is necessary for the expression of all assayed ear

transcription factors. *Pax2* is required for the expression of *Zbtb16* and *Lmx1a*, which in turn is

necessary for *Pax2,* suggesting that they act in a positive feedback loop. All gene expression changes

can be rescued by co-electroporation of the appropriate full-length constructs (Extended Data Fig. 9).

These experiments put Sox8 at the top of the ear determination network forming a regulatory circuit

with Pax2 and Lmx1a.

Sox8 induces ectopic otic vesicles and vesicle-derived neurons

If Sox8, Pax2 and Lmx1a indeed form a minimal circuit driving otic specification, they should be able

to convert non-ear cells into cells with ear identity. We tested this hypothesis by electroporating a

combination of Sox8/Pax2/Lmx1a-mCherry-tagged constructs into head-fold-stage ectoderm not destined to contribute to the ear. After 72hrs we observe many otic vesicles scattered across the head ectoderm. Some of these contain neurofilament-positive neurons, with processes projecting out from the ectopic vesicle (Fig. 4a).

We next asked whether Sox8 alone can initiate the otic programme using the Lmx1aE1-EGFP enhancer and otic markers as a readout. We find that misexpression of Sox8-mCherry not only activates Lmx1aE1-EGFP in ectopic vesicles, but also the expression of *Pax2* and *Lmx1a*, and the ear marker *Soho-1* (Fig. 4b-d; Extended Data Fig. 10a). To assess to what extent Sox8 can confer ear identity, we isolated double-positive Sox8-mCherry/Lmx1a-EGFP cells from ss11-12 by FACS and compared their transcriptome with control ectoderm labelled with constitutive mCherry/EGFP. Differential expression analysis shows upregulation of 399 transcripts in comparison to controls, while 112 genes are downregulated (log2FC > 1.5; adjusted p-value < 0.05) (Fig. 4e, Extended Data Fig. 9b-d, Supplementary Table 3). Seventeen of the twenty seven upregulated transcription factors are known to be expressed in the otic placode or vesicle, while for the remaining ten no expression data are available (Supplementary Table 3). In contrast, among the downregulated genes are typical neural crest and forebrain transcription factors. The first cell types specified during ear formation are neurons of the cochlear-vestibular ganglion, we therefore asked whether Sox8 can also trigger neurogenesis. We find that Sox8-induced ectopic vesicles are associated with neurofilament-positive neurites generated from Sox8 expressing cells themselves (Fig. 4f, g). Together, these results show that Sox8 can activate the transcriptional program for ear identity in non-ear cells by direct lineage reprograming *in vivo*.

Conclusion

We have established an integrated workflow for multi-omics data analysis to ensure accessibility and reproducibility. Using transcriptional profiling at single cell level, enhancer discovery and analysis, together with functional experiments, we have identified the critical components of the ear determination network: Sox8, Pax2 and Lmx1a (Fig. 4h). Sox8 lies at the top of the transcriptional hierarchy, and alone is sufficient to impart ear identity to cells otherwise destined to form head

epidermis, other sense organs or cranial ganglia. Previous findings have implicated Pax2 in otic

placode development and patterning[35-38] and Sox8 in regulating ear-specific enhancers[39-41]. Sox

proteins cooperate with a variety of transcription factors to exert their cell type specific function[42]. It is

therefore tempting to speculate that, in analogy to the eye, where Sox2 cooperates with Pax6 to

regulate lens-specific transcription[43], in the ear Sox8 might partner with Pax2.

Our results also show that OEPs initially express competing transcriptional programmes that

resolve over time as otic and epibranchial cell states are established. We capture previously unknown

gene modules that accompany this process as well as regulatory regions associated with otic-

epibranchial specification. In turn, this information is critical to unravel the underlying gene regulatory

networks and identify the transcription factor codes that determine cell identity in the cranial sensory

nervous system. In the long term, this will enhance our ability to engineer specific sensory cell types

for basic research and regenerative purposes.

References

1 Jacobson, A. G. The determination and positioning of the nose, lens, and ear. III. Effects of

204  reversing the antero-posterior axis of epidermis, neural plate and neural fold. *J Exp Zool*

205  **154**, 293-303 (1963).

206 2 Groves, A. K. & Bronner-Fraser, M. Competence, specification and commitment in otic

207  placode induction. *Development* **127**, 3489-3499 (2000).

208 3 Baker, C. V., Stark, M. R., Marcelle, C. & Bronner-Fraser, M. Competence, specification and

209  induction of Pax-3 in the trigeminal placode. *Development* **126**, 147-156 (1999).

210 4 Bhattacharyya, S. & Bronner-Fraser, M. Competence, specification and commitment to an

211  olfactory placode fate. *Development* **135**, 4165-4177 (2008).

212 5 Schlosser, G. Early embryonic specification of vertebrate cranial placodes. *Wiley Interdiscip*

213  *Rev Dev Biol* **3**, 349-363, doi:10.1002/wdev.142 (2014).

214 6 Thiery, A., Buzzi, A. L. & Streit, A. Cell fate decisions during the development of the

215  peripheral nervous system in the vertebrate head. *Curr Top Dev Biol* **139**, 127-167,

216  doi:10.1016/bs.ctdb.2020.04.002 (2020).

217   7   Davis, R. L., Weintraub, H. & Lassar, A. B. Expression of a single transfected cDNA converts

218   fibroblasts to myoblasts. *Cell* **51**, 987-1000, doi:10.1016/0092-8674(87)90585-x (1987).

219   8   Gehring, W. J. The master control gene for morphogenesis and evolution of the eye. *Genes*

220   *Cells* **1**, 11-15. (1996).

221   9   Streit, A. Extensive cell movements accompany formation of the otic placode. *Dev Biol* **249**,

222   237-254 (2002).

223   10   Pieper, M., Eagleson, G. W., Wosniok, W. & Schlosser, G. Origin and segregation of cranial

224   placodes in Xenopus laevis. *Dev Biol* **360**, 257-275, doi:S0012-1606(11)01268-1 [pii]

225   10.1016/j.ydbio.2011.09.024 (2011).

226   11   Ladher, R. K., Anakwe, K. U., Gurney, A. L., Schoenwolf, G. C. & Francis-West, P. H.

227   Identification of synergistic signals initiating inner ear development. *Science* **290**, 1965-1968.

228   (2000).

229   12   Wright, T. J. & Mansour, S. L. Fgf3 and Fgf10 are required for mouse otic placode induction.

230   *Development* **130**, 3379-3390 (2003).

231   13   Ohyama, T., Groves, A. K. & Martin, K. The first steps towards hearing: mechanisms of otic

232   placode induction. *Int J Dev Biol* **51**, 463-472 (2007).

233   14   Ladher, R. K., O'Neill, P. & Begbie, J. From shared lineage to distinct functions: the

234   development of the inner ear and epibranchial placodes. *Development* **137**, 1777-1785,

235   doi:137/11/1777 [pii] 10.1242/dev.040055 (2010).

236   15   Chen, J. & Streit, A. Induction of the inner ear: stepwise specification of otic fate from

237   multipotent progenitors. *Hear Res* **297**, 3-12, doi:10.1016/j.heares.2012.11.018 (2013).

238   16   Chen, J. *et al.* A systems-level approach reveals new gene regulatory modules in the

239   developing ear. *Development* **144**, 1531-1543, doi:10.1242/dev.148494 (2017).

240   17   Nishimura, N. *et al.* A Systematic Survey and Characterization of Enhancers that Regulate

241   Sox3 in Neuro-Sensory Development in Comparison with Sox2 Enhancers. *Biology (Basel)*

242   **1**, 714-735, doi:10.3390/biology1030714 (2012).

18   Delile, J. *et al.* Single cell transcriptomics reveals spatial and temporal dynamics of gene
expression in the developing mouse spinal cord. *Development* **146**, doi:10.1242/dev.173807
(2019).

19   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by
pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381-386, doi:10.1038/nbt.2859
(2014).

20   La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494-498, doi:10.1038/s41586-
018-0414-6 (2018).

21   Choi, H. M. T. *et al.* Third-generation in situ hybridization chain reaction: multiplexed,
quantitative, sensitive, versatile, robust. *Development* **145**, doi:10.1242/dev.165753 (2018).

22   Creyghton, M. P. *et al.* Histone H3K27ac separates active from poised enhancers and
predicts developmental state. *Proc Natl Acad Sci U S A* **107**, 21931-21936,
doi:10.1073/pnas.1016071107 (2010).

23   Kharchenko, P. V. *et al.* Comprehensive analysis of the chromatin landscape in Drosophila
melanogaster. *Nature* **471**, 480-485, doi:10.1038/nature09725 (2011).

24   Rada-Iglesias, A. *et al.* A unique chromatin signature uncovers early developmental
enhancers in humans. *Nature* **470**, 279-283, doi:nature09692 [pii] 10.1038/nature09692
(2011).

25   Zentner, G. E., Tesar, P. J. & Scacheri, P. C. Epigenetic signatures distinguish multiple
classes of enhancers with distinct cellular functions. *Genome Res* **21**, 1273-1283,
doi:10.1101/gr.122382.111 (2011).

26   Chen, J. & Streit, A. A medium-scale assay for enhancer validation in amniotes. *Dev Dyn*
244, 1291-1299, doi:10.1002/dvdy.24306 (2015).

27   Brugmann, S. A., Pandur, P. D., Kenyon, K. L., Pignoni, F. & Moody, S. A. Six1 promotes a
placodal fate within the lateral neurogenic ectoderm by functioning as both a transcriptional
activator and repressor. *Development* **131**, 5871-5881 (2004).

28   Ozaki, H. *et al.* Six1 controls patterning of the mouse otic vesicle. *Development* **131**, 551-
562 (2004).

29   Zheng, W. *et al.* The role of Six1 in mammalian auditory system development. *Development*
130, 3989-4000 (2003).

30   Maharana, S. K. & Schlosser, G. A gene regulatory network underlying the formation of pre-
placodal ectoderm in Xenopus laevis. *BMC Biol* **16**, 79, doi:10.1186/s12915-018-0540-5
(2018).

31   Kwon, H. J., Bhat, N., Sweet, E. M., Cornell, R. A. & Riley, B. B. Identification of early
requirements for preplacodal ectoderm and sensory organ development. *PLoS Genet* **6**,
doi:10.1371/journal.pgen.1001133 (2010).

32   Benayoun, B. A. *et al.* H3K4me3 breadth is linked to cell identity and transcriptional
consistency. *Cell* **158**, 673-688, doi:10.1016/j.cell.2014.06.027 (2014).

33   Hnisz, D. *et al.* Super-enhancers in the control of cell identity and disease. *Cell* **155**, 934-
947, doi:10.1016/j.cell.2013.09.053 (2013).

34   Whyte, W. A. *et al.* Master transcription factors and mediator establish super-enhancers at
key cell identity genes. *Cell* **153**, 307-319, doi:10.1016/j.cell.2013.03.035 (2013).

35   Torres, M., Gomez-Pardo, E. & Gruss, P. Pax2 contributes to inner ear patterning and optic
nerve trajectory. *Development* **122**, 3381-3391 (1996).

36   Burton, Q., Cole, L. K., Mulheisen, M., Chang, W. & Wu, D. K. The role of Pax2 in mouse
inner ear development. *Dev Biol* **272**, 161-175 (2004).

37   Christophorou, N. A., Mende, M., Lleras-Forero, L., Grocott, T. & Streit, A. Pax2 coordinates
epithelial morphogenesis and cell fate in the inner ear. *Dev Biol* **345**, 180-190, doi:S0012-
1606(10)00911-5 [pii] 10.1016/j.ydbio.2010.07.007 (2010).

38   Freter, S. *et al.* Pax2 modulates proliferation during specification of the otic and epibranchial
placodes. *Dev Dyn* **241**, 1716-1728, doi:10.1002/dvdy.23856 (2012).

39   Betancur, P., Sauka-Spengler, T. & Bronner, M. A Sox10 enhancer element common to the

otic placode and neural crest is activated by tissue-specific paralogs. *Development* **138**,

3689-3698, doi:dev.057836 [pii] 10.1242/dev.057836 (2011).

40   Murko, C. & Bronner, M. E. Tissue specific regulation of the chick Sox10E1 enhancer by

different Sox family members. *Dev Biol* **422**, 47-57, doi:10.1016/j.ydbio.2016.12.004 (2017).

41   Okamoto, Y. *et al.* Cooperation of Sall4 and Sox8 transcription factors in the regulation of the

chicken Sox3 gene during otic placode development. *Dev Growth Differ* **60**, 133-145,

doi:10.1111/dgd.12427 (2018).

42   Kondoh, H. & Kamachi, Y. SOX-partner code for cell specification: Regulatory target

selection and underlying molecular mechanisms. *The International Journal of Biochemistry &*

Cell Biology **42**, 391-399 (2010).

43   Kamachi, Y., Uchikawa, M., Tanouchi, A., Sekido, R. & Kondoh, H. Pax6 and SOX2 form a

co-DNA-binding partner complex that regulates initiation of lens development. *Genes Dev*

15, 1272-1286 (2001).

   Figures and Figure Legends

Figure 1. Transcriptomic characterisation of ear development. a-c, *In ovo* electroporation (**a**) was

used to label and collect otic (**b**; Lmx1aE1-EGFP⁺) and epibranchial (**c;** Sox3U3-EGFP⁺) cells for bulk

RNAseq. **d**, Volcano plot showing genes differentially expressed (absolute log2 fold-change >1.5 and

adjusted p-value < 0.05) in otic (orange) and epibranchial cells (green). **e**, **f**, Cells were collected for

scRNAseq after *ex ovo* electroporation of the Pax2E1-EGFP enhancer, which is active in OEPs, otic

and epibranchial placodes (**f**); cells were collected at the stages indicated. **g-l**, tSNE representation

of unsupervised hierarchical clustering of all cells (**g-i**) and of the placodal cell subset (**j-l**); cells were

colour-coded according to stage collected (**g**, **j**; OEP grey, ss11-12 blue, ss14-15 black), clusters (**h**,

k) and placodal markers *Six1* (**i**) and *Pax2* (**l**) expression. **m**, Heatmap showing partitioning of the

placodal subset (C1, C2 in h) into five clusters (PC1-5) based on gene modules (GM). Expression

profiles reveal that PC1 largely contains OEP-like cells, while PC2/ and PC4/5 are composed of otic-

like and epibranchial-like cells. **n-o**, Pseudo-time ordering using Monocle2 shows trajectories

between stages (**n**) and clusters (**o**). Note: otic- and epibranchial-like cells segregate into two

branches. **p**, RNAvelocity vector field verifies the directional trajectories predicted by Monocle2.

Figure 2. Dynamic gene expression during OEP segregation into otic and epibranchial lineages. a, BEAM identifies genes regulated in a branch-dependant manner. **b**, Histogram showing the proportion of cells co-expressing genes that are expressed before the branching point, but segregate to the otic or epibranchial branch. Significantly more cells co-express such genes before the branching point than thereafter. Error bars indicate +/- 1 standard deviation. ** p-value < 0.01; *** p-value < 0.001. **c**, Dot plot for OEP, otic and epibranchial markers based on scRNAseq data. Expression level is indicated by colour intensity and gene expression frequency by dot size. **d**, **e**, A proportion of cells co-expresses otic (*Sox8*, *Lmx1a*) and epibranchial (*Foxi3*, *Ap2e*) markers prior to the branching point. **f-i**, *In situ* HCR (**f**, **h**) and intensity measurements (**g**, **i**) along the medial (M) to lateral (L) axis of the OEP-domain confirms gradual segregation of OEP gene expression as distinct otic and epibranchial cell populations emerge.

Figure 3. Chromatin accessibility profile of OEPs identifies putative regulatory regions. a,

Genome browser view of ATAC, H3K27ac and H3K27me3 profiles in OEPs at the Lmx1a locus.

Cloned putative enhancer is shown in green. **b**, Coelectroporation of Lmx1aE1-EGFP reporter (green)

and constitutive mCherry (magenta) reveals *in vivo* enhancer activity in the otic placode. **c**, Motif

enrichment analysis of all identified enhancers. **d**, Diagram showing the activation of expression of

potential regulators of otic identity during specification and commitment stages. **e**, Knock-down of

selected transcription factors using fluorescein labelled antisense oligonucleotides (aON; green) lead

to downregulation of otic markers as identified by *in situ* hybridisation (blue).

Figure 4. Direct reprogramming of ectoderm by Sox8. a, Co-electroporation of mCherry-tagged

vectors containing the full-length sequence of Sox8/Pax2/Lmx1a (magenta) generate ectopic

vesicles across the cranial ectoderm after 72hrs culture. Neurofilament antibody staining (NF;

green) shows axonal projections from the vesicle. **b-d**, Overexpression of Sox8 (Sox8 OE) alone

activates Lmx1aE1-EGFP (green) and otic marker expression on the electroporated, but not on the

control side (Ctrl). **e**, RNAseq of Sox8 and control electroporated cranial ectoderm; volcano plot

shows enrichment of otic genes in the Sox8 OE (absolute log2 fold-change >1.5 and adjusted p-

value < 0.05). **f**, Sox8-mCherry overexpression (magenta) alone generates ectopic otic vesicles with

neuronal projections (green), while controls do not (**g**). **h**, BioTapestry model showing the minimal

transcriptional circuit for otic specification with Sox8 at the top of the hierarchy.

Methods

Expression and enhancer constructs

Putative enhancers were amplified from chick genomic DNA and cloned into pTK-EGFP reporter vectors after digestion with XcmI[26]. To generate expression constructs total RNA was isolated from HH8-12 chick embryos with RNAqueous-Micro Total RNA Isolation Kit (Thermo Fisher Scientific), reverse transcribed using Superscript II reverse transcriptase (Thermo Fisher Scientific, 180644-014) and oligo-dT primer. Specific primers were used to amplify the full-length coding sequence of Sox8, Pax2, Lmx1a and Zbtb16 and PCR products were cloned into pCAB-IRES-mCherry. All sequences were verified by Sanger sequencing.

Chick embryos, electroporation and culture

Fertilised hens' eggs (Stewart, Norfolk UK) were incubated at 38°C and staged according to Hamburger and Hamilton (HH)[44]. All experiments were performed on embryos younger than 12 days, thus were not regulated by the Animals Scientific Procedures Act 1986.

OEPs for bulk RNAseq were labelled using *in ovo* electroporation[45]; eggs were incubated until the 3-6 somite stage (ss), pTK-Lmx1aE1-EGFP or pTK-Sox3U3-EGFP plasmids (1µg/µl) were injected targeting the OEP territory. Electroporation using Ovodyne electroporator (TSS20, Intracel) was performed with five 50ms pulses of 8V at 100ms intervals. After incubation until ss18-21, embryos were collected in PBS and processed for bulk RNAseq.

For *ex ovo* culture, embryos were harvested using filter paper rings[46] and cultured on egg albumen; for long-term culture (72hrs) the 'modified Cornish pasty' method[47] was used. *Ex ovo* electroporation was performed to collect cells for scRNAseq, for knock-down, rescue and overexpression experiments. The posterior placodal region or the cranial surface ectoderm of HH6 embryos was targeted for electroporation by injecting plasmid DNA at 1µg/ul (pTK-Pax2E1-EGFP to label OEPs; pCES-Sox8-mCherry + pTK-Lmx1aE1-EGFP, pCES-Sox8-mCherry + pCES-Lmx1a-mCherry + pCES-Pax2-mCherry; pCES-mCherry + pCES-EGFP for misexpression), control or antisense-oligonucleotides (aON; GeneTools; 0.75mM with 0.3µg/µl carrier DNA), or a combination of aON and expression construct (0.75mM aON + 0.75µg/µl plasmid). Oligonucleotides used were Sox8: 5'-

400   CTCCTCGGTCATGTTGAGCATTTGG-3'[39], Pax2: 5'-GGTCTGCCTTGCAGTGCATATCCAT-3'[37],

401   Lmx1a: 5'-CCTCCATCTTCAAGCCGTCCAGCAT-3'[16],   Zbtb16:   5'-

402   GTCAAATCCATAGCACTCCCGAGGT-3', Sox13: 5'-CTCCTCATGGACATCCATTTCATTC-3' and

403   control 5'-ATGGCCTCGGAGCTGGAGAGCCTCA-3'. Electroporation was performed in a chamber

404   using five 5V pulses of 50ms in 100ms intervals[45].

Whole mount and hybridization chain reaction fluorescent *in situ* hybridisation

In situ hybridization was carried out following previously described protocols[48]. Whole-mount pictures

were taken using an Olympus SZX12 with a Retiga2000R camera and Q-Capture Pro7 software.

HCR v3 was performed using the Molecular Technologies protocol[21]. Briefly, embryos were

fixed in 4% PFA for one hour at room temperature, dehydrated in a series of methanol in PBT and

stored overnight at -20°C. After rehydration and proteinase-K treatment (20mg/ml; 3 min) embryos

were post-fixed in 4% PFA for 20 min. Embryos were then washed on ice in PBS, 1:1 PBT/5X SSC

(5X sodium chloride sodium citrate, 0.1% Tween-20) and 5X SSC for 5 min each. Pre-hybridization

in hybridisation buffer was performed for 5 min on ice, followed by 30 minutes at 37°C. Embryos were

hybridised overnight at 37°C with probes at 4pmol/ml in hybridization buffer. After four 15 min washes

with probe wash buffer at 37°C, preamplification was carried out in amplification buffer for 5 min at

room temperature. Hairpins were prepared individually at 30pmol final concentration; they were

incubated at 95°C for 90 seconds followed by cooling to room temperature for 30 min, protected from

light. Cooled hairpins were added to 500µl amplification buffer, embryos were incubated in hairpins

overnight at room temperature followed by two 5 min and two 30 min washes in 5X SSC. After a 5

min incubation in DAPI (10 mg/ml), they were washed three times for 10 min with 5X SSC, before

being imaged using Leica SP5 laser scanning confocal inverted microscope using the LAS AF

software.

For HCR image analysis, Z-stacks were collected for 50-70µm; figures show projections of all

stacks. Images were processed using ImageJ and the ImageJ Plot Profile tool was used to calculate

intensity plots. In brief, an optical section in the centre of the placode territory was selected using the

Pax2 channel as a reference. The Pax2 channel was added to the Sox8, Lmx1a, Foxi3 or Tfap2e

channels, respectively. Intensity values were then calculated across the area of interest and plotted.

Whole mount immunostaining

Embryos were collected in PBS, fixed for 25 minutes at room temperature in 4% PFA, washed in

PBS-Tx (PBS + 0.2% Triton X-100) and blocked in 5% goat serum in PBS-Tx for 3–5 h at room

temperature. Embryos were then incubated in primary antibody diluted in blocking buffer for 24-72

hours at 4°C. Primary antibodies were rabbit anti–mCherry (1:200; Abcam ab167453), mouse anti–

NF (1:100; Thermo Fisher Scientific 13-0700), rabbit anti-GFP (1:500; Abcam a11122) or mouse anti

GFP (1:1000; Molecular Probes A11120). After five 60 min washes and one overnight wash in PBS-

Tx, embryos were incubated in secondary antibodies (1:800) at 4°C overnight. Secondary antibodies

used were goat anti–rabbit IgG Alexa Fluor 488 (Thermo Fisher Scientific, A11036), donkey anti–

rabbit Alexa Fluor 568 (Thermo Fisher Scientific, A11001), goat anti mouse IgG Alexa Fluor 488

(Molecular Probes A11001) and goat anti mouse IgG Alexa Fluor 568 (Molecular Probes A11004).

Embryos were then were briefly incubated in PBS containing 10 mg/ml DAPI and washed at least five

times in PBS-Tx before being mounted on slides and imaged using Leica SP5 laser scanning confocal

inverted microscope using a 10x objective or an Olympus SZX12 with a Retiga2000R camera and Q-

Capture Pro7 software. Confocal whole mount images in the manuscript are maximum intensity

projections of embryo z-stacks. Sections were imaged using a 63x oil immersion objective and

maximum intensity projections are shown.

Cryosectioning

Embryos were embedded in gelatine as previously described[49] and cryo-sectioned at 15-20μm using

a Bright OTF5000 cryostat. Sections were mounted using Mowiol 4-88 (Sigma Aldrich, 81381) and

imaged using Leica SP5 laser scanning confocal inverted microscope (LAS AF software) or a Zeiss

Axiovert 200M microscope with a Hamamatsu C4742-95 camera and using OCULAR software.

Quantitative PCR (qPCR)

RNA from dissected otic tissue was isolated using the RNAqueous-Micro Kit (Ambion, AM1931) and

reverse transcribed. Primers for target genes were designed with PrimerQuest (IDT). qPCR was

performed using Aria Mx Real-Time System (Agilent Technologies) with SYBR green master mix

(Roche, 64913850001). RT-qPCR for antisense oligonucleotide experiments was carried out with a

minimum of three biological replicates; two sample Welch's t-tests with Benjamini-Hochberg multiple

test correction were used to determine statistical significance between control and antisense

oligonucleotide ΔCt values. Relative expression ($2^{-\Delta\Delta Ct}$)[50] was calculated using the NormqPCR

package in R. The geometric mean of Gapdh and Rplp1 expression, or Rplp1 expression alone, was

used to normalise gene expression.

Bulk RNA sequencing

To label otic and epibranchial cells embryos were electroporated with Lmx1aE1-EGFP and Sox3U3-

EGFP plasmids, respectively, and whole heads were used for cell collection. For overexpression

experiments embryos were electroporated with Sox8-mCherry+Lmx1aE1-EGFP or with pCAB-

mCherry+pCAB-EGFP, the endogenous otic placode and the trunk were removed before cell

dissociation. Cells were dissociated in FACSmax cell dissociation solution (Ambion, T200100)

containing papain (30U/mg, Sigma-Aldrich, 10108014001) for 20min at 37°C before being transferred

to Hanks Balanced Solution without calcium and magnesium (HBSS, Life Technologies, 14185045)

containing 5% heat-inactivated foetal-bovine-serum (FBS), rock inhibitor (10 µM, Stemcell

Technologies, Y-27632) and non-essential amino acids (Thermo Fisher Scientific, 11140035). Cells

were disaggregated by pipetting, sequentially filtered through 0.35µm and 0.20µm filters (Miltenyi

Biotech, 130-101-812). Pelleted cells were resuspended in 500µl HBSS and isolated by fluorescent

activated cell sorting (FACS) using a BD FACS-Aria Diva. 2000 cells per biological replicate were

collected, centrifuged at 200xg for 5 mins at 4°C, washed with PBS, and resuspended in lysis buffer.

RNA was extracted using Ambion RNAqueous Micro Total RNA isolation kit (AM1931, ThermoFisher

Scientific). RNA integrity was checked using Bioanalyser with Agilent RNA 6000 pico kit (Agilent

Technologies, 5067-1513); samples with RIN >7 were processed for library preparation. Sequencing

libraries were prepared using Nextera Sample low input kit (Illumina, 15028212) and sequenced using

75bp paired end reads on the Illumina Hiseq4000 platform. A minimum of three biological replicates

were used for analysis.

Single-cell RNA extraction, library preparation and sequencing

HH6-7 embryos were electroporated with Pax2E1-EGFP to label OEPs; cells were dissociated as

described above and 288 EGFP[+] cells from each ss8-9, ss11-12 and ss14-15 were collected by FACS

in 96-well plates. Sequencing libraries were prepared following the SmartSeq2 protocol[51]. Libraries

were sequenced on the Illumina NextSeq500 platform using single end 75bp sequencing (ss8-9) or

on the HiSeq4000 platform using paired end 75bp sequencing (ss11-12, ss14-15).

Nuclei isolation, ATAC library preparation and sequencing

The ATACseq library was prepared following published protocols[52]. Approximately 30 pieces of the

OEP territory were dissected from ss8-9 embryos, dissociated with Dounce homogenizer (tight pestle)

in lysis buffer (10mM Tris-HCl, pH7.4, 10mM NaCl, 3mM $MgCl_2$, 0.1% IGEPAL CA-630). Nuclei were

pelleted at 4000 rpm at 4°C for 10 min. After removing the lysis buffer, 1.25µl Tn5 transposase

(Illumina, FC-131-1024) were added to 25µl reaction volume and incubated at 37°C for 10 mins.

Tagmented DNA was then purified with Mini Elute PCR purification kit (Qiagen, 28004) followed by 9

cycles of PCR enrichment using NEB High fidelity PCR kit (NEB, M0541S). The quality of ATAC

libraries was assessed with Agilent Bioanalyzer with DNA High Sensitivity kit (Agilent Technologies,

5067-4627) and quantified with Kapa NGS library quantification kit (Kapa Biosystems, KK4824). The

libraries were sequenced with Illumina HiSeq 2000/2500 in 2x100 cycles.

H3K27ac, H3K27me3 and H3K4me4 ChIP library preparation and sequencing

Approximately 200 pieces of OEP ectoderm were dissected from ss8-9 embryos. The tissue was

dissociated in nuclei expulsion buffer (0.5% NP-40, 0.25% Triton X-100, 10 mM Tris-HCl, pH 7.5,

3mM $CaCl_2$, 0.25M sucrose, 1x protease inhibitor (Roche), 1mM DTT, and 0.2mM PMSF) using a

Dounce homogenizer (loose pestle); crosslinking was performed in 1% formaldehyde for 9 minutes,

followed by quenching with 1M glycine for 5 minutes. Cells were then pelleted and washed 3 times

with PBS containing protease inhibitor (Roche, 11873580001). The pellets were snap-frozen and

stored at -80°C for later use. Crosslinked chromatin was fragmented by sonication in an ice bath

(Misonix Q700 at 7 Amplitude, 5x40 seconds, 30 seconds on, 60 seconds off) and immunoprecipitated

following the Nano-ChIP protocol[53]. For chromatin immunoprecipitation (ChIP), the following

antibodies were used anti-H3K27Ac (Abcam, ab4729), anti-H3K4me3 (Diagenode, A5051-001P) and

anti-H3K27me3 (Millipore, 07449). Adaptors and primers from NEBNext library preparation kit

(Illumina, E6040S) were used to prepare the library following NanoChip protocol[53]. The library was

enriched using 14 PCR cycles and quantified with Kapa NGS library quantification kit (Kapa

Biosystems, KK4824) before pooling at a concentration of 20nM. Library quality was assessed with

Agilent Bioanalyzer with DNA High Sensitivity kit (Agilent Technologies, 5067-4627) and quantified

with Kapa NGS library quantification kit (Kapa Biosystems, KK4824). The libraries were sequenced

with Illumina HiSeq 2000/2500 in 2x100 cycles.

High-Throughput-Sequencing-Data (HTSD) Analysis

All data alignment and downstream analysis was carried out using NF-core and custom Nextflow

pipelines to allow full reproducibility. All code used, including Nextflow pipelines and downstream

analysis, can be found at https://github.com/alexthiery/otic-reprogramming. Full detailed

documentation for the pipeline is also available at https://alexthiery.github.io/otic-reprogramming/. A

custom Docker container used for the downstream analysis pipeline can be found at

https://hub.docker.com/repository/docker/alexthiery/otic-reprogramming-r_analysis. This also allows

for interactive exploration of the data.

Bulk-RNA-seq

Bulk-RNA-seq data were processed and aligned to GalGal6 using the default NF-core RNA-seq (v2.0)

pipeline[54] which uses the STAR aligner. Downstream differential expression analysis (Lmx1aE1-

EGFP vs Sox3U3-EGFP; Sox8OE vs ControlOE) was carried out with the DESeq2 package in R[55].

Adjusted p-values were calculated using the default DESeq2 multiple test correction (Benjamini-

Hochberg). Differentially expressed transcripts were determined by an absolute log2 fold-change >1.5

and adjusted p-value < 0.05.

Single-cell-RNA-seq alignment

SmartSeq2 single-cell-RNA-seq reads were aligned and processed using a custom Nextflow DSL2

pipeline. Adaptor sequences were trimmed using Cutadapt (v2.10). HISAT2 (v2.2.1) was then used

to build a genome index from GalGal6 (amended to include a GFP sequence), before extracting splice

sites from the GTF and aligning reads. Read counts were obtained using HTSeq (v0.12.4). BAM files

from HISAT2 were also passed to Velocyto (v0.17)[20] in order to get spliced and unspliced expression

matrices for further downstream analysis.

Single-cell-RNA-seq data analysis

Downstream data analysis was carried out primarily using the Antler R package (version:

Development2019)[18]. We excluded from the dataset: cells which expressed fewer than 1k genes or

fewer than 500k reads, cells with more than 6% reads from mitochondrial genes, genes which were

expressed in fewer than 3 cells and genes with CPM < 10.

Identification of gene modules

To identify clusters of genes with correlated expression unbiasedly (gene modules), we used the

Antler *identifyGeneModules* function. Genes which had a Spearman correlation greater than 0.3 with

at least 3 other genes were first removed. Genes were then hierarchically clustered iteratively into

gene modules and filtered. Gene modules were filtered based on the minimum expression level (5

CPM) and the proportion (0.4) of cells expressing a gene module. The number of final gene modules

was set to 40 to achieve reasonably large clusters, which broadly characterise cell type diversity

across the dataset. Gene modules were then filtered based on the presence of genes with known

expression profiles in ectodermal derivatives, before cells were re-clustered based on the remaining

gene modules. We used the expression of known markers to assign cell states. Placodal cells were

then subset and re-clustered using a new set of gene modules identified from the subset cell

population. Three main cell states were identified as otic, epibranchial and OEP populations.

Gene expression dynamics at the otic-epibranchial branching point

To model the transcriptional dynamics at the otic-epibranchial branching point, we ordered cells along

pseudotime using Monocle2[56]. The lineage tree was rooted to the earliest cell state (ss8-9) which was

comprised mostly of OEPs. The expression of genes along the bifurcation point was then modelled

using branch expression analysis modelling (BEAM).

To assess whether individual OEPs simultaneously express markers associated with the otic

or epibranchial state, we calculated the co-expression of key otic and epibranchial genes within each

of the Monocle branches. First, gene expression was binarized based on the presence or absence of

gene expression; then the proportion of cells co-expressing pairs of otic-epibranchial genes within

each of the branches was calculated. A Kruskal Wallis test was then used to compare the proportion

of cells co-expressing otic and epibranchial genes between the three branches. Post-hoc pairwise

comparisons between the OEP branch and the otic or epibranchial branch were carried out using two-

tailed Wilcoxon rank sum tests.

RNA velocity provides an alternative to Monocle2 for inferring future cell state of individual

cells. Spliced, un-spliced and spanning matrices obtained from Velocyto[20] were subset in R based on

genes and cells used to generate the Monocle trajectory. RNA velocity was then calculated using the

Velocyto.R package, where spanning reads were used to fit gene offsets. Single cell velocities as well

as vector fields were subsequently visualised on both tSNE and Monocle DDRTree embeddings.

ChIP-seq and ATAC-seq alignment and peak calling

ChIP-seq and ATAC-seq data were processed and aligned to GalGal6 using the default NF-core

ChIP-seq (v1.2.0) and NF-core ATAC-seq (v1.2.0) pipelines[54], respectively. Reads were aligned using

the Burrows-Wheeler aligner. MACS v2.2.7.1 was used to call broad peaks (FDR < 0.1) and narrow

peaks (FDR < 0.05) for ChIP-seq and ATAC-seq data, respectively.

Enhancer discovery

Putative enhancers were identified using a custom Nextflow DSL2 pipeline. First, bedtools (v2.29.2) was used to subset ATAC-seq narrow peaks, which overlap with a broad H3K27ac peak, whilst removing those overlapping with broad H3K27me3 peaks. Remaining ATAC-seq peaks were then annotated with Homer (v4.11), using a GalGal6 GTF which was filtered for protein coding genes. Promoter and exonic peaks were then removed. Motif enrichment and functional enrichment analysis were carried out using Homer and g:Profiler, respectively.

Data availability

All data have been deposited in Gene Expression Omnibus (GEO; Accession number GSE168089). Nextflow pipelines and downstream analysis, can be found at https://github.com/alexthiery/otic-reprogramming. Full detailed documentation is available at https://alexthiery.github.io/otic-reprogramming/. A custom Docker container used can be found at https://hub.docker.com/repository/docker/alexthiery/otic-reprogramming-r_analysis.

References

44   Hamburger, V. & Hamilton, H. L. A series of normal stages in the development of the chick embryo. *J Morphol* **88**, 49-92 (1951).

45   Norris, A. & Streit, A. Morpholinos: Studying gene function in the chick. *Methods* **66**, 454-465, doi:10.1016/j.ymeth.2013.10.009 (2014).

46   Chapman, S. C., Collignon, J., Schoenwolf, G. C. & Lumsden, A. Improved method for chick whole-embryo culture using a filter paper carrier. *Dev Dyn* **220**, 284-289, doi:10.1002/1097-0177(20010301)220:3<284::AID-DVDY1102>3.0.CO;2-5 (2001).

47   Nagai, H., Lin, M. C. & Sheng, G. A modified cornish pasty method for ex ovo culture of the chick embryo. *Genesis* **49**, 46-52, doi:10.1002/dvg.20690 (2011).

48   Streit, A. & Stern, C. D. Combined whole-mount in situ hybridization and

immunohistochemistry in avian embryos. *Methods* **23**, 339-344,

doi:10.1006/meth.2000.1146 (2001).

49   Stern, C. D. & Holland, P. W. H. *Essential Developmental Biology: A Practical Approach.*,

(1993).

50   Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time

quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402-408,

doi:10.1006/meth.2001.1262 (2001).

51   Picelli, S. *et al.* Full-length RNA-seq from single cells using Smart-seq2. *Nat Protoc* **9**, 171-

181, doi:10.1038/nprot.2014.006 (2014).

52   Buenrostro, J. D., Giresi, P. G., Zaba, L. C., Chang, H. Y. & Greenleaf, W. J. Transposition

of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-

binding proteins and nucleosome position. *Nat Methods* **10**, 1213-1218,

doi:10.1038/nmeth.2688 (2013).

53   Adli, M. & Bernstein, B. E. Whole-genome chromatin profiling from limited numbers of cells

using nano-ChIP-seq. *Nat Protoc* **6**, 1656-1668, doi:10.1038/nprot.2011.402 (2011).

54   Ewels, P. A. *et al.* The nf-core framework for community-curated bioinformatics pipelines.

Nat Biotechnol **38**, 276-278, doi:10.1038/s41587-020-0439-x (2020).

55   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8 (2014).

56   Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat*

Methods **14**, 979-982, doi:10.1038/nmeth.4402 (2017).

Acknowledgements. We thank Tatjana Sauka-Spengler and Ruth Williams for assistance with

establishing ChIPseq and scRNAseq protocols, Monica Tambalo for help with tissue collection,

Rosalinda Guerra and Ewa Kolano-Merlin for technical assistance, and James Briscoe for advice on

single cell transcriptomics. Thanks to Claudio D. Stern for comments on the manuscript and the

Streit group for discussions. This work was funded by grants to AS from the Biotechnology and

Biological Sciences Research Council (BB/S005536/1; BB/M006964/1) and the National Institute on

Deafness and Other Communication Disorders (DC011577).

Author contributions. A.L.B., J.C. and A.S. designed experiments; A.L.B. performed RNAseq for

Lmx1a-E1-EGFP, Sox3U3-EGFP, Sox8-mCherry/Lmx1aE1-EGFP, mCherry/EGFP populations,

scRNAseq, HCR analysis, Sox8/Pax2/Lmx1a OE and some Sox8 OE experiments; J.C. performed

ATACseq, ChIPseq, enhancer cloning and testing, in situ hybridisation, qPCR, knock-down and

Sox8 OE experiments; J.C. performed the initial analysis of ATACseq and ChIPseq data. A.L.B.

interpreted and supervised bioinformatics analysis; J.D. performed some scRNAseq analysis, A.T.

analysed all transcriptomic and epigenomic data and established the integrated pipeline; A.L.B.

generated all figures; A.L.B., A.T. and A.S. wrote the manuscript.

Competing interests. The authors declare no competing interests.

Additional information

Supplementary information is available for this paper.

Correspondence and requests for material should be addressed to Andrea Streit:

andrea.streit@kcl.ac.uk.