Pseudogene-mediated gene conversion after CRISPR-Cas9 editing demonstrated by partial *CD33* conversion with *SIGLEC22P*

Benjamin C. Shaw, MS[1] and Steven Estus, PhD[1*]

[1]Department of Physiology and Sanders-Brown Center on Aging, University of Kentucky

Correspondence to:

Steve Estus, PhD

Sanders-Brown Center on Aging, Room 330

800 S. Limestone St.,

Lexington, KY USA 40536

Phone: 859-218-2388

e-mail: steve.estus@uky.edu

Pseudogene-mediated gene conversion after CRISPR-Cas9 editing demonstrated by partial *CD33* conversion with *SIGLEC22P*

Running Head: Pseudogene conversion after Cas9 cleavage

Abstract

Although gene editing workflows typically consider the possibility of off-target editing, pseudogene-directed homology repair has not, to our knowledge, been reported previously. Here, we employed a CRISPR-Cas9 strategy for targeted excision of exon 2 in *CD33* in U937 human monocytes. Candidate clonal cell lines were screened by using a clinically relevant antibody known to label the IgV domain encoded by exon 2 (P67.6, gemtuzumab). In addition to the anticipated deletion of exon 2, we also found unexpected P67.6-negative cell lines which had apparently retained CD33 exon 2. Sequencing revealed that these lines underwent gene conversion from the nearby SIGLEC22P pseudogene during homology repair that resulted in three missense mutations relative to CD33. Ectopic expression studies confirmed that the P67.6 epitope is dependent upon these amino acids. In summation, we report that pseudogene-directed homology repair can lead to aberrant CRISPR gene editing.

Keywords: homology directed repair, CD33, gemtuzumab ozogamicin, D2-CD33

Introduction

The CRISPR–Cas9 system has revolutionized gene editing.[1] In this process, a single-stranded

guide RNA (sgRNA) directs Cas9 endonuclease to cleave DNA at a sequence-specific site. The

DNA cleavage results in a double stranded DNA break (DSB) that is repaired by either

homology-directed repair (HDR) or nonhomologous end-joining (NHEJ). The former results in

targeted integration of DNA sequence while the latter typically results in gene disruption through

the introduction of insertions or deletions (indels). To generate a targeted knock-in or knock-out,

an HDR template of exogenous DNA is often supplied as part of the process.[2] Alternatively,

endogenous HDR templates have been described, including *HBD* sequence being incorporated

into *HBB* and sequence from one allele of *HPRT* being incorporated into the other allele.[3, 4]

However, to our knowledge, HDR directed by a pseudogene has not been previously reported.

CD33 genetic variants, including rs124594919, have been associated with reduced risk of

Alzheimer's Disease (AD) in genome-wide studies.[5-7] We and others subsequently identified

rs12459419 as a functional SNP that increases the proportion of CD33 lacking exon 2 (D2-

CD33).[8-12] This exon encodes the ligand-binding IgV domain of this member of the sialic acid-

binding immunoglobulin-type lectin (SIGLEC) family.[13] Hence, while CD33 normally includes

an extracellular IgV and IgC_2 domain, D2-CD33 encodes a protein without the IgV domain.[14]

CD33 inhibits microglial activity through its immunomodulatory tyrosine inhibitory motif

(ITIM) and ITIM-like domains, which recruit protein tyrosine phosphatases, SHP1 and SHP2, to

impact intracellular calcium flux, phagocytosis, and microglial migration.[9, 11, 12, 15-20]

Given that the AD-protective rs12459419 increases D2-CD33 at the expense of CD33, the

prevailing theoretical mechanism has been that rs12459419 reduces AD risk through decreased

CD33 function. However, recent findings that a *bona fide* loss of function indel, rs201074739, is

not associated with AD risk, has led to this hypothesis being revised to suggest that rs12459419 and its related D2-CD33 isoform represent a gain of function.[13, 21, 22] The gain-of-function mechanism and localization of D2-CD33 protein remain heavily debated. [8, 9, 11, 13, 17, 20-25]

Here, we sought to generate a model of physiologic D2-CD33 expression by using CRISPR-Cas9 to excise *CD33* exon 2 in the human U937 cell line. During these experiments, we identified a subset of cells which apparently underwent HDR directed by the *SIGLEC22P* pseudogene, located 13.5 kb away from *CD33*. Although the *SIGLEC22P* pseudogene shares approximately 87% identity over 1800 bp with *CD33*, this gene conversion was detected because three nucleotides in SIGLEC22P differ from those within the targeted *CD33* exon 2 and result in three missense amino acids in CD33, including p.N20K, p.F21I, and p.W22R. Hence, we report pseudogene directed gene conversion as a mechanism for unanticipated CRISPR mutations.

Results

CRISPR-Cas9-mediated CD33 Exon 2 deletion leads to loss of P67.6 epitope

To generate an *in vitro* model of D2-CD33, we targeted exon 2 for deletion by using guide RNAs corresponding to sequences in the flanking introns as previously described.[25] Clonal cell lines were established and examined by flow cytometry with the antibodies P67.6 and HIM3-4, which target the IgV and IgC$_2$ epitopes encoded by exon 2 and exon 3, respectively. CD33 domains targeted by each antibody used in this study are shown in Table 1. While unedited cells showed robust labeling by both HIM3-4 and P67.6 (Figure 1A), edited cell lines showed strong labeling by HIM3-4 but not P67.6 (Figure 1B), or no labeling by either HIM3-4 or P67.6 (Figure 1C). Data are representative of three independently established cell lines for each phenotype. Since D2-CD33 protein is not readily apparent on the cell surface, we expected that the latter cells (Figure 1C) were candidates for exon 2 excision, which was confirmed by a PCR product of the appropriate size (Figure 1D, right) and by sequencing.[8, 23-25] However, cell lines with robust cell surface HIM3-4 but not P67.6 labeling were unexpected. Screening by the size of the PCR amplicon with primers corresponding to exon 1 and exon 3 suggested that exon 2 was still present (Figure 1D, middle). Sequencing of this PCR fragment revealed that the HIM3-4$^+$ P67.6$^-$ clones contained three apparent single nucleotide polymorphisms (SNPs) in exon 2 compared to the unedited, wild-type (WT-CD33) U937 cell line. These SNPs have an identical minor allele frequency (MAF) = 9.86 x 10^{-5} and are indexed as rs3987761, rs3987760, and rs35814802.[26] Introduction of these SNPs results in changes in three consecutive amino acids (p.N20K, p.F21I, p.W22R) which we refer to as KIR-CD33. The nonsynonymous amino acids are the 4-6th amino-terminal residues of the mature protein. EMBOSS and PSORT II predict cell surface localization for KIR-CD33 and no change in the signal peptide cleavage site.[27, 28] Consistent with typical

cell-surface localization, HIM3-4 labeled both CD33 and KIR-CD33 in a similar fashion (Figure 1A-B). Notably, total *CD33* gene expression and exon 2 splicing in KIR-CD33 cells does not differ from that of unedited cells (Figure 1E-F). CD33 expression was increased in D2-CD33 cells (Figure 1E) which exclusively express the *D2-CD33* isoform (Figure 1F).

Validation of the CRISPR-induced in-frame disruption of the P67.6 epitope

To rigorously demonstrate that the lack of P67.6 labeling was solely due to the KIR mutations, we introduced the KIR mutations into a previously described CD33 expression vector.[8] CD33 and KIR-CD33 vectors were transfected into HEK293 cells, which do not naturally express CD33, and the cells processed for flow cytometry (Figure 2) and immunofluorescent confocal microscopy (Figure 3). For flow cytometry, cells were labeled with P67.6 or WM53, each of which was conjugated to the same fluorochrome to facilitate direct comparisons (Figure 2). Importantly, these antibodies both target exon 2, are both mouse IgG1κ and have comparable degrees of labeling. In the CD33 HEK293 cells, labeling with both WM53 and P67.6 correlated well with HIM3-4 (Figure 2A-B). However, in the KIR-CD33 HEK293 cells, labeling with WM53 but not P67.6 correlated with HIM3-4 labeling as P67.6 labeling was not apparent (Figure 2C-D). Further gating on the HIM3-4[+] cells shows that WM53 labeling is not affected by the KIR-CD33 mutation (Figure 2E), while P67.6 labeling in KIR-CD33 cells is comparable to non-transfected control cells (Figure 2F). These results were confirmed with immunofluorescent confocal microscopy using an array of anti-CD33 antibodies (Figure 3). The *CD33*-transfected HEK293 cells showed consistent double-labeling between an antibody against a cytoplasmic epitope (H110) and either WM53, P67.6, or PWS44 (Figure 3A-C). For the KIR-CD33 cells, robust co-labeling was observed between H110 and WM53 or PWS44 (Figure 3D, F). However, P67.6 labeling of KIR-CD33 cells was not detected (Figure 3E). We thus conclude that the

residues identified—p.N20, p.F21, and p.W22—are necessary for P67.6 binding and that these changes are the reason that this antibody failed to label the CRISPR-edited cells. Although P67.6 has been humanized and used clinically, prior studies have not mapped its epitope at this resolution.[17, 29, 30]

Identification of SIGLEC22P as a homology-directed repair template for CD33

The three nucleotide changes (Figure 4, red) are 23-27 bp away from the putative Cas9 cleavage site (Figure 4, scissors), were not consecutive, and were present and identical in each of the clones with the KIR-CD33 phenotype. Since this was unlikely due to chance, we hypothesized that this was due to HDR from elsewhere in the genome. A search of a 50 bp sequence centered on the KIR mutations revealed that this sequence occurs in the *SIGLEC22P* pseudogene which is located 13.5 kb away from *CD33*. Further investigation found an extended region of homology between *CD33* and *SIGLEC22P* that flanked the Cas9 cleavage site; indeed, the first 500 bp of the genes share 97% identity, including a 143 bp region of otherwise complete identity centered on the KIR mutations (Figure 4, underline). The *SIGLEC22P* pseudogene exon 2 contains 11 additional mutations as well as two intronic mutations relative to CD33. None of these mutations were detected in the CRISPR-edited cell lines, indicating the region used for repair was limited to, at most, the 143 bp region surrounding the KIR mutations near the Cas9-induced double-stranded break. We interpret these results as indicating that, upon double-strand breakage at the beginning of exon 2 in *CD33* (Figure 4, blue), the *SIGLEC22P* pseudogene was used as a repair template because of the strong sequence homology to *CD33* (Figure 4, underlined). In this process, three *SIGLEC22P*-specific mutations were introduced into *CD33*, resulting in missense mutations in three adjacent codons and thus KIR-CD33. This indicates an in-frame, ectopic gene conversion using pseudogene sequence.

Discussion

We show here that the DSB repair pathways initiated after S. pyogenes Cas9 cleavage can lead to ectopic gene conversion from a pseudogene in a mitotic human cell line. This gene conversion resulted in an in-frame chimeric protein, wherein a fragment smaller than 150 bp of a nearby pseudogene replaced the targeted gene sequence (Figure 4). Indeed, this distance is consistent with meiotic gene conversion events observed by Jeffreys et al., who showed that gene conversion occurs through relatively short tracts with a mean length between 55-290 bp.[31] Exogenous double-stranded regions of homology as short as 58 bp have been used *in vitro* to introduce mutations through CRISPR-Cas9/HDR mechanisms.[2] Coincidentally, this ectopic gene conversion disrupted the epitope of a well-validated antibody, P67.6 (Figure 5). This murine antibody is clinically relevant as gemtuzumab. Using transiently transfected HEK293 cells expressing the chimeric protein, we demonstrated that these mutations are sufficient to abrogate P67.6 binding, providing the most precise epitope mapping to date of this clinically relevant antibody.

The KIR-CD33 mutations—rs3987761, rs3987760, rs35814802—are indexed in dbSNP and gnomAD with minor allele frequencies $< 10^{-4}$, and roughly equivalent across populations.[26] It is possible that these mutations, while rare, occur naturally and are clinically relevant. The SIGLEC family of genes are undergoing rapid evolution in many species, including humans.[32] This rapid evolution has resulted in the pseudogenization of many SIGLECs, including the *CD33* pseudogene *SIGLEC22P*. These same bases are indexed as SNPs in *SIGLEC22P* (rs997169007, rs1049597792, and rs1005338799), also have identical frequencies (MAF = 2.1×10^{-4}) in the gnomAD database, and the major and minor alleles are the inverse of the KIR-CD33 SNPs.[26] The region of homology identified here between *CD33* and *SIGLEC22P* is 143 bp, substantially

shorter than the 50 bp reads upon which the gnomAD and dbSNP databases are built.[33] While it is possible these are *bona fide* variants in *CD33*, the most likely explanation is that the *SIGLEC22* pseudogene sequences have been mapped incorrectly to *CD33* by algorithms. These missense SNPs are also not recorded in the BeatAML variant database, which records known AML-associated functional variants such as missense SNPs, further underscoring the low probability that these are true *CD33* variants.[34]

Off-target editing is a frequent concern in gene editing workflows, including CRISPR-Cas9. However, gene conversion is often overlooked as a potential confound. For instance, within *CD33*, off-target editing of *SIGLEC22P* resulting in a 14 kb deletion and subsequent *SIGLEC22P*/*CD33* gene fusion has been previously reported, which may have been the result of a mitotic crossover.[35, 36] Neither report noted a gene conversion between *SIGLEC22P* and *CD33*. Mitotic crossover initiated by CRISPR-Cas9 cleavage has also been reported in the *HPRT* locus, resulting in a 36 kb crossover.[4] Gene conversion via CRISPR-Cas9 between a 101 bp homologous region in *HBD* and *HBB* has also been reported; notably, these genes were also the first reported gene conversion event in humans.[3, 37] To our knowledge, this is the first report of pseudogene-mediated gene conversion during the CRISPR-Cas9 editing process. Our results demonstrate the need for rigorous screening in studies which rely on gene editing, and analysis of the region flanking the expected cut site for homology and possible gene conversion. Since the human genome contains 8000-12000 pseudogenes and approximately 3400 genes in the human genome have known pseudogenes, pseudogene directed homology repair is a potentially considerable confound.[38, 39] As pseudogenes are by definition nonfunctional, they may be overlooked as irrelevant during the design process. Given the pseudogene conversion described here, there is a clear need to analyze the sequence of the edit, rather than relying on gene or

protein expression. While off-target editing of a nonfunctional pseudogene may have limited impact on downstream results, incorporating elements from a nonfunctional pseudogene into a target gene may lead to deleterious mutations that abrogate function of the target gene entirely. This is especially important to consider for designs which incorporate unedited cells which have undergone sorting and single cell cloning as a control. By coincidence, our initial screen for editing included an antibody which overlapped the KIR sequence. This gene conversion, at the protein level, is masked when using an alternative antibody (WM53) targeting the same domain and is not apparent by PCR alone. Combining the data from Figures 1D-F and 2C, one could reason that these cells were unedited as there are no apparent differences in PCR fragment size, gene expression, splicing, or cell surface protein expression, and thus incorrectly assume that sequencing is unnecessary. We conclude that, in addition to off-target Cas9-editing confounds, researchers should be aware of the potential for pseudogene-directed homology repair.

Methods

Cell Lines and Antibodies

U937 and HEK293 cell lines were obtained from American Type Culture Collection (ATCC).

U937 cells were cultured in RPMI 1640 with HEPES (Gibco 22400-089) supplemented with

10% fetal bovine serum, defined (HyClone, GE Healthcare SH30070.03); 50 U/mL penicillin, 50

µg/mL streptomycin (Gibco 15070-063); and 2 µM L-glutamine (Gibco A2916801). HEK293

cells were cultured in EMEM, ATCC formulation (ATCC 30-2003) supplemented with 10%

fetal bovine serum, defined (HyClone, GE Healthcare SH30070.03); 50 U/mL penicillin, 50

µg/mL streptomycin (Gibco 22400-089). Cells were maintained at 37°C in a 5% CO_2 in air

atmosphere. Antibodies, concentrations, and CD33 domains targeted are shown in Table 1.

CRISPR-Cas9 Gene Editing

CRISPR reagents were purchased from Integrated DNA Technologies (IDT). Single-guide

RNAs (sgRNAs) and S. pyogenes Cas9 protein (IDT 1081059) were incubated at a 1:1 molar

ratio (0.5 nmol each) at room temperature for 10 minutes to form ribonucleotide-protein

complexes (RNPs). The sgRNA sequences targeting CD33 exon 2 were 5'-

TCCATAGCCAGGGCCCCTGT and 5'-GCATGTGACAGGTGAGGCAC.[25] U937 cells were

washed three times in PBS (Gibco 10010-023) and resuspended in complete Nucleofector Kit C

(Lonza Biosciences VCA-1004) media (10^6 cells per transfection) with 5 µL electroporation

enhancer (IDT 1075916) and RNPs. Cells were electroporated using a Nucleofector IIb device

(Lonza Biosciences) under protocol V-001 and immediately added to a 12 well plate with 1.5 mL

complete media and cultured for two weeks.

Cell Sorting and Flow Cytometry

Edited U937 cells were washed in PBS with 5% heat-inactivated fetal bovine serum (Gibco

10082-147), resuspended at 10^6 cells/mL and then treated with Human TruStain FcX blocker (BioLegend 422302). Cell sorting was carried out in azide-free buffers; for flow cytometry, 0.02% sodium azide was included in all buffers. Cells were stained with HIM3-4-FITC and P67.6-BV711 for one hour on ice, washed twice with HBSS, then stained with Fixable Viability Dye eFluor780 (Invitrogen 65-0865-18). Cells were resuspended in HBSS (Gibco 24020-117) with 5% heat-inactivated fetal bovine serum (Gibco 10082-147) for sorting. Viable cells were gated using scatter and viability exclusion stain, sorted as either HIM3-4[+] P67.6[+], HIM3-4[+] P67.6[-], or HIM3-4[-] P67.6[-] and collected in complete media. At 48 hours post-sort, cells were split using limiting dilution into a 96 well plate at an average density of 0.5 cells/well and expanded until sufficient cell numbers for analysis were achieved, approximately 8 weeks. Clones were screened by flow cytometry again prior to PCR and sequence analysis.

PCR Screening and Cloning

Genomic DNA from CRISPR-edited U937 clones was isolated with a DNeasy kit (Qiagen 69506) per manufacturer instructions and amplified with Q5 High-Fidelity DNA Polymerase (New England BioLabs M0439L) using forward primer 5'-CACAGGAAGCCCTGGAAGCT and reverse primer 5'-GAGCAGGTCAGGTTTTTGGA (Invitrogen). Thermocycling parameters were as follows: 98°C 1 min; 98°C 15 s, 66°C 15 s, 72°C 45 s, 32 cycles; 72°C 2 min, 25°C hold. PCR products were separated on a 0.8% agarose-TBE gel, purified using a Monarch gel extraction kit (New England BioLabs T1020L), and sequenced by a commercial company (ACGT, Wheeling, IL). The three missense mutations identified were introduced into a previously described pcDNA3.1-CD33-V5/HIS vector using a QuikChange XL kit (Agilent 200517) with forward primer 5'-GCACTTGCAGCCGGATTTTTGGATCCATAGCCAGGGCC and reverse primer 5'- GGCCCTGGCTATGGATCCAAAAATCCGGCTGCAAGTGC

(Invitrogen) to generate the pcDNA3.1-KIR-CD33-V5/HIS vector, transformed into TOP10 E. coli (Invitrogen C404003), isolated using a Plasmid Plus Midiprep Kit (Qiagen 12945) and verified by sequencing (ACGT).[8]

Gene expression by qPCR

Quantitative PCR (qPCR) was used to quantify expression of total CD33 and D2-CD33 as previously described.[14] Briefly, primers corresponding to sequences within exons 4 and 5 were used to quantify total *CD33* expression (forward, 5'-TGTTCCACAGAACCCAACAA-3'; reverse, 5'-GGCTGTAACACCAGCTCCTC-3'), as well as primers corresponding to sequences at the exon 1–3 junction and exon 3 to quantify the D2-CD33 isoform (forward, 5'-CCCTGCTGTGGGCAGACTTG-3'; reverse, 5'-GCACCGAGGAGTGAGTAGTCC-3'). PCR was conducted using an initial 2 min incubation at 95°, followed by cycles of 10 s at 95°C, 20 s at 60°C, and 20 s at 72°C. The 20 µl reactions contained 1 µM each primer, 1X PerfeCTa SYBR Green Super Mix (Quanta Biosciences), and 20 ng of cDNA. Experimental samples were amplified in parallel with serially diluted standards that were generated by PCR of cDNA using the indicated primers, followed by purification and quantitation by UV absorbance. Results from samples were compared relative to the standard curve to calculate copy number in each sample. Assays were performed in triplicate and normalized to expression of ribosomal protein L32 (*RPL32*) as the housekeeping gene.[14, 40]

HEK293 Transfection

HEK293 cells were seeded at approximately 70% confluency 24 hours before transfection. Cells were then transfected with Lipofectamine 3000 with Plus Reagent (Invitrogen L3000001) per manufacturer instructions, 250 ng plasmid per well in 8 well glass chamber slides (MatTek CCS-8) for immunofluorescence or 1000 ng per well in 12 well plates (Corning 3513) for flow

cytometry. Cells were transfected with either the previously described wild-type CD33 vector (pcDNA3.1-CD33-V5/HIS), pcDNA3.1-KIRCD33-V5/HIS, or no vector control. Cells were incubated for 24 hours before analysis by flow cytometry or immunofluorescence and confocal microscopy.

Confocal Immunofluorescence Microscopy

Transfected HEK293 cells were fixed with 10% neutral buffered formalin (Fisher Scientific SF100-4) for 30 minutes then blocked and permeabilized for 30 minutes with 10% goat serum (Sigma S26-LITER), 0.1% Triton X-100 (Fisher Scientific BP151-500) in PBS (Fisher BioReagents BP665-1). Primary and secondary antibodies were diluted in the same blocking and permeabilization buffer and incubated at room temperature for 90 minutes. Cells were washed three times in blocking and permeabilization buffer between primary and secondary antibodies, and three times in PBS prior to coverslip mounting with Prolong Glass with NucBlue mounting media (Invitrogen P36981) and high-tolerance No. 1.5 coverglass (ThorLabs CG15KH1). Images were acquired using a Nikon A1R HD inverted confocal microscope with a 60X oil objective and NIS Elements AR software.

Statistical analyses

Analyses were performed using GraphPad Prism 8.4.2. Gene expression data were analyzed by one-way ANOVA followed by Dunnett's multiple comparisons to unedited U937 cells.

Acknowledgements

This work was supported by RF1AG05971701S1 and R21AG068370 (SE). We would like to thank Ann M. Stowe for flow cytometry support, the University of Kentucky Light Microscopy Core Facility for help with image acquisition and processing, and the University of Kentucky Flow Cytometry and Immune Monitoring Core Facility, supported in part by the Office of the Vice President for Research, the Markey Cancer Center and an NCI Center Core Support Grant (P30 CA177558).

Authorship Contributions

BCS conducted all experiments. BCS and SE designed experiments, analyzed data, and drafted the manuscript.

Disclosure of Conflicts of Interest

Authors declare no conflicts of interest.

References

1.   Jinek M, Chylinski K, Fonfara I et al. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science.* 2012;337:816-821. DOI: 10.1126/science.1225829

2.   Renaud J-B, Boix C, Charpentier M et al. Improved Genome Editing Efficiency and Flexibility Using Modified Oligonucleotides with TALEN and CRISPR-Cas9 Nucleases. *Cell Reports.* 2016;14:2263-2272. DOI: 10.1016/j.celrep.2016.02.018

3.   Javidi-Parsijani P, Lyu P, Makani V et al. CRISPR/Cas9 increases mitotic gene conversion in human cells. *Gene Ther.* 2020;27:281-296. DOI: 10.1038/s41434-020-0126-z

4.   Susani L, Castelli A, Lizier M et al. Correction of a Recessive Genetic Defect by CRISPR-Cas9-Mediated Endogenous Repair. *Crispr j.* 2018;1:230-238. DOI: 10.1089/crispr.2018.0004

5.   Hollingworth P, Harold D, Sims R et al. Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease. *Nat Genet.* 2011;43:429-435. DOI: 10.1038/ng.803

6.   Jansen IE, Savage JE, Watanabe K et al. Genome-wide meta-analysis identifies new loci and functional pathways influencing Alzheimer's disease risk. *Nat Genet.* 2019. DOI: 10.1038/s41588-018-0311-9

7.   Naj AC, Jun G, Beecham GW et al. Common variants at MS4A4/MS4A6E, CD2AP, CD33 and EPHA1 are associated with late-onset Alzheimer's disease. *Nat Genet.* 2011;43:436-441. DOI: 10.1038/ng.801

8.   Malik M, Chiles IIIJ, Xi HS et al. Genetics of CD33 in Alzheimer's disease and acute myeloid leukemia. *Human Molecular Genetics.* 2015;24:3557-3570. DOI: 10.1093/hmg/ddv092

9.   Raj T, Ryan KJ, Replogle JM et al. CD33: increased inclusion of exon 2 implicates the Ig V-set domain in Alzheimer's disease susceptibility. *Human Molecular Genetics.* 2014;23:2729-2736. DOI: 10.1093/hmg/ddt666

10.   Schwarz F, Springer SA, Altheide TK et al. Human-specific derived alleles of CD33 and other genes protect against postreproductive cognitive decline. *Proc Natl Acad Sci U S A.* 2016;113:74-79. DOI: 10.1073/pnas.1517951112

11.   Griciuc A, Serrano-Pozo A, Parrado Antonio R et al. Alzheimer's Disease Risk Gene CD33 Inhibits Microglial Uptake of Amyloid Beta. *Neuron.* 2013;78:631-643. DOI: https://doi.org/10.1016/j.neuron.2013.04.014

12.   Bradshaw EM, Chibnik LB, Keenan BT et al. CD33 Alzheimer's disease locus: altered monocyte function and amyloid biology. *Nat Neurosci.* 2013;16:848-850. DOI: 10.1038/nn.3435

13.   Estus S, Shaw BC, Devanney N et al. Evaluation of CD33 as a genetic risk factor for Alzheimer's disease. *Acta Neuropathol.* 2019. DOI: 10.1007/s00401-019-02000-4

14.   Malik M, Simpson JF, Parikh I et al. CD33 Alzheimer's Risk-Altering Polymorphism, CD33 Expression, and Exon 2 Splicing. 2013;33:13320-13325. DOI: 10.1523/JNEUROSCI.1224-13.2013 %J The Journal of Neuroscience

15.   Balaian L, Zhong RK, Ball ED. The inhibitory effect of anti-CD33 monoclonal antibodies on AML cell growth correlates with Syk and/or ZAP-70 expression. *Exp Hematol.* 2003;31:363-371. DOI: 10.1016/s0301-472x(03)00044-4

16.   Hernández-Caselles T, Martínez-Esparza M, Pérez-Oliva AB et al. A study of CD33 (SIGLEC-3) antigen expression and function on activated human T and NK cells: two isoforms of CD33 are generated by alternative splicing. *Journal of Leukocyte Biology.* 2006;79:46-58. DOI: https://doi.org/10.1189/jlb.0205096

17.   Perez-Oliva AB, Martinez-Esparza M, Vicente-Fernandez JJ et al. Epitope mapping, expression and post-translational modifications of two isoforms of CD33 (CD33M and CD33m) on lymphoid and myeloid human cells. *Glycobiology.* 2011;21:757-770. DOI: 10.1093/glycob/cwq220

18.   Walter RB, Raden BW, Zeng R et al. ITIM-dependent endocytosis of CD33-related Siglecs: role of intracellular domain, tyrosine phosphorylation, and the tyrosine phosphatases, Shp1 and Shp2. *J Leukoc Biol.* 2008;83:200-211. DOI: 10.1189/jlb.0607388

19.   Paul SP, Taylor LS, Stansbury EK et al. Myeloid specific human CD33 is an inhibitory receptor with differential ITIM function in recruiting the phosphatases SHP-1 and SHP-2. *Blood.* 2000;96:483-490.

20.   Bhattacherjee A, Rodrigues E, Jung J et al. Repression of phagocytosis by human CD33 is not conserved with mouse CD33. *Commun Biol.* 2019;2:450. DOI: 10.1038/s42003-019-0698-6

21.   Bhattacherjee A, Jung SJ, Ho M et al. The CD33 short isoform is a gain-of-function variant that enhances Aβ1-42 phagocytosis in microglia. *Molecular Neurodegeneration.* 2021.

22.   Wissfeld J, Nozaki I, Mathews M et al. Deletion of Alzheimer's disease-associated CD33 results in an inflammatory human microglia phenotype. *Glia.* 2021. DOI: 10.1002/glia.23968

23.   Siddiqui SS, Springer SA, Verhagen A et al. The Alzheimer's disease-protective CD33 splice variant mediates adaptive loss of function via diversion to an intracellular pool. *J Biol Chem.* 2017;292:15312-15320. DOI: 10.1074/jbc.M117.799346

24.   Godwin CD, Laszlo GS, Wood BL et al. The CD33 splice isoform lacking exon 2 as therapeutic target in human acute myeloid leukemia. *Leukemia.* 2020;34:2479-2483. DOI: 10.1038/s41375-020-0755-7

25.   Humbert O, Laszlo GS, Sichel S et al. Engineering resistance to CD33-targeted immunotherapy in normal hematopoiesis by CRISPR/Cas9-deletion of CD33 exon 2. *Leukemia.* 2019;33:762-808. DOI: 10.1038/s41375-018-0277-8

26.   Karczewski KJ, Francioli LC, Tiao G et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature.* 2020;581:434-443. DOI: 10.1038/s41586-020-2308-7

27.   Nakai K, Horton P. PSORT: a program for detecting sorting signals in proteins and predicting their subcellular localization. *Trends Biochem Sci.* 1999;24:34-36. DOI: 10.1016/s0968-0004(98)01336-x

28.   Rice P, Longden I, Bleasby A. EMBOSS: the European Molecular Biology Open Software Suite. *Trends Genet.* 2000;16:276-277. DOI: 10.1016/s0168-9525(00)02024-2

29.   Mortland L, Alonzo TA, Walter RB et al. Clinical significance of CD33 nonsynonymous single-nucleotide polymorphisms in pediatric patients with acute myeloid leukemia treated with gemtuzumab-ozogamicin-containing chemotherapy. *Clin Cancer Res.* 2013;19:1620-1627. DOI: 10.1158/1078-0432.CCR-12-3115

30.   Chauhan L, Shin M, Wang YC et al. CD33_PGx6_Score Predicts Gemtuzumab Ozogamicin Response in Childhood Acute Myeloid Leukemia: A Report From the Children's Oncology Group. *JCO Precis Oncol.* 2019;3. DOI: 10.1200/po.18.00387

31.   Jeffreys AJ, May CA. Intense and highly localized gene conversion activity in human meiotic crossover hot spots. *Nature Genetics.* 2004;36:151-156. DOI: 10.1038/ng1287

32.   Padler-Karavani V, Hurtado-Ziola N, Chang YC et al. Rapid evolution of binding specificities and expression patterns of inhibitory CD33-related Siglecs in primates. *The FASEB Journal.* 2014;28:1280-1293. DOI: 10.1096/fj.13-241497

33.   Lonsdale J, Thomas J, Salvatore M et al. The Genotype-Tissue Expression (GTEx) project. *Nature Genetics.* 2013;45:580-585. DOI: 10.1038/ng.2653

34.   Tyner JW, Tognon CE, Bottomly D et al. Functional genomic landscape of acute myeloid leukaemia. *Nature.* 2018;562:526-531. DOI: 10.1038/s41586-018-0623-z

35.   Borot F, Wang H, Ma Y et al. Gene-edited stem cells enable CD33-directed immune therapy for myeloid malignancies. *Proceedings of the National Academy of Sciences.* 2019:201819992. DOI: 10.1073/pnas.1819992116

36.   Kim MY, Yu K-R, Kenderian SS et al. Genetic Inactivation of CD33 in Hematopoietic Stem Cells to Enable CAR T Cell Immunotherapy for Acute Myeloid Leukemia. *Cell.* 2018;173:1439-1453.e1419. DOI: 10.1016/j.cell.2018.05.013

37.   Slightom JL, Blechl AE, Smithies O. Human fetal G gamma- and A gamma-globin genes: complete nucleotide sequences suggest that DNA can be exchanged between these duplicated genes. *Cell.* 1980;21:627-638. DOI: 10.1016/0092-8674(80)90426-2

38.   Zhang Z, Carriero N, Zheng D et al. PseudoPipe: an automated pseudogene identification pipeline. *Bioinformatics.* 2006;22:1437-1439. DOI: 10.1093/bioinformatics/btl116

39.   Pei B, Sisu C, Frankish A et al. The GENCODE pseudogene resource. *Genome Biology.* 2012;13:R51. DOI: 10.1186/gb-2012-13-9-r51

40.   Grear KE, Ling IF, Simpson JF et al. Expression of SORL1 and a novel SORL1 splice variant in normal and Alzheimers disease brain. *Mol Neurodegener.* 2009;4:46. DOI: 10.1186/1750-1326-4-46

Tables

Clone	Conjugate	Vendor	Catalog #	Lot	Dilution/Use	CD33 domain targeted
P67.6	BV711	BioLegend	366624	B302694	2.5 µg/mL FC	IgV
P67.6	Unconjugated	BioLegend	825601	B258818	5 µg/mL IF	IgV
WM53	BV711	BioLegend	303424	B253729	5 µg/mL FC	IgV
WM53	Unconjugated	BioLegend	96281	B274701	5 µg/mL IF	IgV
HIM3-4	FITC	BioLegend	303304	B284834	20 µg/mL FC	IgC$_2$
PWS44	Unconjugated	Leica	NCL-L-CD33	6024275	2 µg/mL IF	IgC$_2$
H110	Unconjugated	Santa Cruz	sc-28811	K2211	1 µg/mL IF	Intracellular
Goat anti-Mouse	AlexaFluorPlus 488	Invitrogen	A32723	TF266577	10 µg/mL IF	
Goat anti-Rabbit F(ab)$_2$	AlexaFluor 568	Invitrogen	A21069	2087701	10 µg/mL IF	

Table 1. Antibodies used in this study. IF: immunofluorescence, FC: flow cytometry

Figure Legends

Figure 1. CRISPR-Cas9 editing of *CD33* exon 2 leads to loss of P67.6 epitope. (A) Unedited U937 cells display robust P67.6 and HIM3-4 labeling. (B) Edited U937 clone that is robustly labeled by HIM3-4 but not P67.6. (C) Edited U937 clone that is not labeled by either HIM3-4 or P67.6. The depicted results in B-C are representative of at least three clonal cell populations established for each phenotype. (D) Genomic DNA PCR of *CD33* exon 1 to exon 3 of the above cell lines. PCR products at 789 bp and 428 bp correspond to the expected sizes for the presence and absence of exon 2, respectively. (E) *KIR-CD33* mutations do not affect total *CD33* gene expression, but removal of exon 2 increases total *CD33* gene expression by 39.8%. (F) *KIR-CD33* mutations do not affect splicing efficiency of exon 2, but removal of exon 2 at the genomic level increases exon 1-exon 3 junction to 100% of total *CD33* expression. Data from E-F analyzed by one-way ANOVA followed by Dunnett's multiple comparisons test to unedited control. ** $p < 0.01$; **** $p < 0.0001$.

Figure 2. Loss of P67.6 epitope is in-frame and preserves cell surface expression. WT-CD33 (A, B) or KIR-CD33 (C, D) expressing HEK293 cells were labeled with HIM3-4 and either WM53 (A, C) or P67.6 (B, D). HIM3-4[+] cells identified in the "Transfected" gate (A-D) were gated to show WM53 (E) or P67.6 (F) binding in transfected cells.

Figure 3. Loss of P67.6 epitope does not alter other common CD33 epitopes. WT-CD33 (A-C) or KIR-CD33 (D-F) HEK293 cells were labeled with H110 (red) and either WM53 (A, D; green), P67.6 (B, E; green), or PWS44 (C, F; green) and DAPI (blue).

Figure 4. Alignment of cell line sequencing data. Unedited U937 cells at top, followed by KIR-CD33 sequence, D2-CD33 sequence, and reference *SIGLEC22P* sequence. Cas9 cleavage

site marked by scissor icons. Mismatches from the unedited *CD33* sequence are denoted by either the differing base (KIR-CD33) or dash in the case of a gap (D2-CD33). Intron 1 in black, exon 2 in blue. The *SIGLEC22P* region used as a repair template is underlined. Mutations introduced into *CD33* in red.

Figure 5. Model of pseudogene repair mechanism and anti-CD33 antibody binding sites. CD33 is normally a transmembrane, cell surface receptor with one IgV domain and one IgC_2 domain. The KIR-CD33 mutation introduced by pseudogene directed-repair abrogates P67.6, but not WM53, binding. IgC_2 domain antibodies HIM3-4 and PWS44, and the intracellular domain H110 antibody bind both CD33 and KIR-CD33. D2-CD33 is not readily apparent on the cell surface in CRISPR-Cas9 edited U937 cells, implying that under physiologic expression, the D2-CD33 protein is retained in an intracellular vesicle. Overall, pseudogene repair of a CRISPR-Cas9-targeted gene can disrupt the binding of well-validated antibodies without introducing a frameshift, and protein-level expression alone is not sufficient for knockout confirmation. Created with Biorender.

Figures

Figure 1.

Figure 2.

Figure 3.

```
WT-CD33          TGAGTGGCTG TGGGGAGAGG GGTTGTCGGG CTGGGCCGAG CTGACCCTCG TTTCCCCACA
KIR-CD33         .......... .......... .......... .......... .......... ..........
D2-CD33          .......... .......... .......... .......... .......... ..........
SIGLEC22P pseudogene ...A...... C......... .......... .......... .......... ..........

WT-CD33          GGGGCCCTGG CTATGGATCC AAATTTCTGG CTGCAAGTGC AGGAGTCAGT GACGGTACAG
KIR-CD33         .......... .......... ...AA..C.. .......... .......... ..........
D2-CD33          .--------- ---------- ---------- ---------- ---------- ----------
SIGLEC22P pseudogene .......... .......... ...AA..C.. .......... .......... ..........

WT-CD33          GAGGGTTTGT GCGTCCTCGT GCCCTGCACT TTCTTCCATC CCATACCCTA CTACGACAAG
KIR-CD33         .......... .......... .......... .......... .......... ..........
D2-CD33          ---------- ---------- ---------- ---------- ---------- ----------
SIGLEC22P pseudogene .......... .......... .......... ...C..... .......... ...A...G.
```

Figure 4.

Figure 5.