Compartimentalized control of Cdk1 drives mitotic spindle assembly.

Angela Flavia Serpico[1,2], Francesco Febbraro[1,2], Caterina Pisauro[1,2] & Domenico Grieco[1,3*].

[1]CEINGE Biotecnologie Avanzate, Naples, Italy; [2]DMMBM, University of Naples "Federico II",

Naples, Italy; [3]Department of Pharmacy, University of Naples "Federico II", Naples, Italy.

*Correspondence and requests for materials should be addressed to D. G. (email:

domenico.grieco@unina.it).

Abstract

During cell division, dramatic microtubular rearrangements driven by cyclin B-cdk1 (Cdk1)

kinase activity mark mitosis onset leading to interphase cytoskeleton dissolution and mitotic

spindle assembly. Once activated by Cdc25, that reverses inhibitory phosphorylation operated

by Wee1/Myt1, Cdk1 clears the cytoplasm from microtubules by inhibiting microtubule

associated proteins (MAPs) with microtubule growth-promoting properties. Nevertheless, some

of these MAPs are required for spindle assembly, creating quite a conundrum. We show here

that a Cdk1 fraction bound to spindle structures escaped Cdc25 action and remained inhibited

by phosphorylation (i-Cdk1) in mitotic human cells. Loss or restoration of i-Cdk1 inhibited or

promoted spindle assembly, respectively. Furthermore, polymerizing spindle microtubules

fostered i-Cdk1 by aggregating with Wee1 and excluding Cdc25. Our data reveal that spindle

assembly relies on compartimentalized control of Cdk1 activity.

To prevent chromosome damage during spinlde assembly, the microtubular interphase cytoskeleton

must be dismantled at the onset of mitosis and, as many mitotic features, this is an event downstream

Cdk1 activation. Indeed, Cdk1 activity clears cytoplasmic microtubules by directly or indirectly

affecting several microtubule-stabilizing and -destabilizing MAPs, inhibiting, for instance, the

microtubule growth-promoting properties of ch-Tog and Map4 (*1, 2, 3, 4, 5*). Nevertheless, a number

of these MAPs are required for spindle microtubule growth (*2, 6, 7*). Thus, how spindle assembles

despite the potent microtubule-destabilizing action of Cdk1 is unknown. During interphase, Cdk1,

accumulates in an inactive state, phosphorylated at inhibitory sites by Wee1/Myt1 kinases (*8*). At

mitosis onset, initial reversal of these phosphorylation by Cdc25 phosphatase is rapidly followed by

full Cdk1 activation via positive feedback loops in which Cdk1 inhibits Wee1/Myt1 and stimulates

Cdc25 (*8*). Thus, inhibitory phopshorylation of Cdk1 has a crucial role in ensuring completion of

interphase tasks before mitosis onset, but once Cdk1 activation begins this control appears to be

rapidly suppressed by autoactivatory feedback loops. Cdk1 is believed to be only inactivated upon

spindle assembly by cyclin B degradation (*8*). Nevertheless, early evidence indicated that cells

expressing a Cdk1 version non-phosphorylatable at the inhibitory sites threonine 14 and tyrosine 15

(Cdk1AF), rapidly entered mitosis but failed to assemble spindles (*9*). These mitotic defects were,

however, ascribed to premature mitosis onset in presence of incompletely replicated DNA and not as

a requirement for inhibitory phosphorylation of Cdk1 in spindle assembly (*9*). In addition, evidence

that i-Cdk1 is present in meiotic and mitotic extracts from Xenopus eggs and mouse eggs was later

reported (*10, 11*). Here we examined if and how inhibitory phosphorylation of Cdk1 had a role for

spindle assembly.

To this end, we begun analysing spindle assembly upon downregulation of Wee1 expression by small

interfering RNAs (siRNAs) in hTERT-RPE1 cells. To prevent premature mitosis onset upon Wee1

downregulation, 6 hours post Wee1-siRNA treatment cells were incubated with the selective and

reversible Cdk1 inhibitor RO-3306 at 9 μM (high-RO) for further 16 hours, to allow completion of

DNA replication and arrest cell cycle at G2 (*12*). As control, part of Wee1-siRNA-treated cells were

previously complemented with siRNA-resistant Wee1 expression vector (fig. 1A; Fig. S1) (*13*). Upon

release from G2-arrest into fresh medium, containing the proteasome inhibitor MG-132 to block

mitosis exit and the protein synthesis inhibitor cycloheximide to prevent abnormal protein

accumulation (M/C medium), the majority of control cells built normal bipolar spindles within 60

min incubation, that remained stable for further 20 min incubation (fig. 1A). Conversely, spindle

assembly was markedly impaired in Wee1-downregulated cells, with defects that ranged from

monopolarity to poorly structured bipolar spindles with gross chromosome alignment abnormalities

(fig. 1A). To establish whether impaired spindle assembly was due to Wee1 loss *per se* or because it

affected Cdk1 activity, we asked if mild Cdk1 inhibition could restore spindle assembly in Wee1-

downregulated cells. Indeed, addition of RO-3306 at 0.5 μM (low-RO), from 60 to 80 min post G2

release, substantially restored spindle assembly (fig. 1A). Defects in spindle assembly were also

induced by cdk1AF overexpression under similar conditions of G2-arrest and release, paralleling

what described by Krek and Nigg in 1991, as well as in Wee1-siRNA-treated HeLa cells or by

chemical inhibition of Wee1, in all cases defects were substantially reversed by low-RO (figs. S2, S3

and S4A) (*9, 14*). In another set of experiments, non-targeting-siRNA-, as control, and Wee1-siRNA-

treated hTERT-RPE1 cells were arrested at prometaphase by the reversible microtubule inhibitor

nocodazole added shortly after siRNA treatments (as described in Materials and Method). Cells were,

then, released in M/C medium for 80 min and a portion of Wee1-siRNA-treated cells also received

low-RO at 60 min incubation. Within this period, the majority of control cells assembled spindles,

while assembly was strongly impaired in Wee1-siRNA-treated cells but restored by low-RO (fig.

1B). When cells were released just into fresh medium, mitosis exit was delayed in Wee1-siRNA-

treated cells, as shown by delayed cyclin B1 degradation, presumably by the action of the spindle

assembly checkpoint (SAC; fig. S5) (*15*). Together, these data suggest that inhibitory

phosphorylation-dependent control of Cdk1 activity is required for spindle assembly.

To directly test if i-Cdk1 was present in mitotic cells and where it was localised, we probed metaphase

cells with an antibody recognizing cdk1 phosphorylated at tyrosine 15 (p-Y15-cdk1; fig. 2A). The p-

Y15-cdk1 signal appeared decorating the spindle area paralleling total cdk1 localization (fig. 2A). To

biochemically dissect localization and function of mitotic i-Cdk1, we adapted a method described by

Nigg and coworkers to isolate mitotic spindles and tightly associated proteins by separating

microtubular, insoluble, pellet fraction (P) from soluble fraction (S) of mitotic cells (see Materials

and Methods) (*16*). Nocodazole-treated, prometaphase-arrested hTERT-RPE1 cells (Pro) were

collected and taken at time 0 or taken after release in M/C medium for 60 min to have metaphase-

arrested cells (Meta; fig. 2B). Cells were fractionated and proteins from S and P fractions analyzed

(fig. 2B; P fraction was enriched 4 folds relatively to S fraction). In Pro cell P fraction, centrosomes

were present as indicated by significant amounts of the central protein γ-tubulin and Nuclear and

Mitotic Apparatus protein 1 (NuMA1), a centrosome-associated protein (fig. 2B) (*3*). Conversely, α-

tubulin was substantially present in S fraction and much less in P fraction, as expected because of the

microtubule polymerization block, as well as the MAPs Map4 and ch-Tog (fig. 2B). In Meta cells,

instead, α-tubulin and significant amounts of Map4 and ch-Tog, the latter also involved in activating

centrosome-independent, intra-spindle, microtubule nucleation, were present in P fraction, reflecting

spindle assembly (fig. 2B) (*17, 18*). Distribution of NuMA1 and γ-tubulin in S and P fractions did not

substantially change between Pro and Meta cells (fig. 2B). P fraction also contained small amounts

of Wee1 that increased in Meta cells, while minimal amounts of Cdc25C were equally present in P

fraction of both Pro and Meta cells (fig. 2B; similar protein distribution was found in HeLa cells

under similar experimental conditions; see Fig. S6). Small amounts of cyclin B1 and cdk1 were

present in P fraction of Pro cells and more of Meta cells (fig. 2B). However, i-Cdk1 was found quite

exclusively in P fractions, as shown by p-Y15-cdk1 signal and by retarded cdk1 mobility on

SDS/PAGE, and more from Meta than from Pro cells (Fig. 2B) (*19*). Quantitation of the upshifted

cdk1 forms from Meta cell P fraction indicated that i-Cdk1 represents less than 10% of total Cdk1.

The selective presence of i-Cdk1 in P fraction was confirmed also by probing immunoprecipitates

(Ips) of comparable amounts of cyclin B1 from S and P fraction of Meta cells (Fig. 2C). In Meta cell

P fraction, Map4 and ch-Tog appeared to have increased mobility on SDS/PAGE compared to S

fraction, suggesting that these proteins were relatively dephosphorylated when microtubule-bound.

Indeed, when Meta cells were further treated with nocodazole, from 60 to 80 min, to revert to a

prometaphase condition, distribution of proteins between S and P fractions returned similar to Pro

cells; Map4 and ch-Tog regained slower migration on SDS/PAGE, presumably reflecting their re-

phosphorylation when returning into the cytoplasm, while Wee1 and i-Cdk1 were reduced in the P

fraction, indicating their dependence also on microtubule polymerization (fig. 2D).

To ask whether a cause/effect relationship existed between i-Cdk1 and the control of Map4 and ch-

Tog dephosphorylation and distribution, prometaphase-arrested control and Wee1-siRNAs were

released into M/C medium as described in Fig. 1b. Cells were further fractionated and the S and P

fractions analyzed (fig. 2E). Control cells showed a similar distribution as in Meta cells described in

Figs. 2b, d. In Wee1-siRNA-treated cells the distribution of γ-tubulin and NuMA1 between the S and

P fractions were similar to control cells, however, Map4, ch-Tog and even part of α-tubulin were

substantially less present in the P fraction relatively to control cells, a distribution pattern similar to

that of Pro cells described in figs 2A, D (fig. 2E). Addition of low-RO to Wee1-siRNA-treated cells

reversed protein distribution relocating Map4, ch-Tog and α-tubulin from S to P fraction similarly to

control cells (fig. 2E). As expected, i-Cdk1 was reduced in Wee1-dowregulated cell P fraction

compared to controls, however, addition of low-RO induced accumulation of part of residual Wee1,

that escaped siRNA-mediated downregulation, in P fraction and restored i-Cdk1 content (fig. 2E;

note Wee1 in short, SE, and long exposures, LE). No increase of p-Y15-cdk1 signal was detected in

S fraction upon low-RO addition, indicating that small fluctuation of Cdk1 activity are indeed

buffered in the cytosol by strong hysteresis and bistability of the Cdk1 autoactivatory loops, while

nucleating spindle microtubules appeared insulated from them (fig. 2E) (*8, 20, 21*). Localized

inhibition of endogenous Cdk1 by low-RO treatment, could possibly explain spindle assembly

restoration in cdk1AF-expressing cells (fig. S2). Immunofluorescence (IF) analysis confirmed that

the two MAPs poorly localised to defectively structured spindles in Wee1-downregulated cells in a

low-RO-reversible manner (fig. 2F). Similar mitotic defects and changes in protein distribution were

induced by the chemical Wee1 inhibitor AZD1775 upon release from prometaphase-arrest, also in

this case corrected by low-RO (fig. S4B). Together, these data indicate that i-Cdk1 is required for

proper mitotic spindle assembly and spindle microtubule association of Map4 and ch-Tog.

Microtubule growth- and stability-promoting properties of MAPs like ch-Tog and Map4 are

antagonized by direct Cdk1 phosphorylation (*2, 3, 4, 5, 6*). We were intrigued by the fact that changes

in migration on SDS/PAGE suggested that these MAPs were hypophosphorylated when associated

with spindle microtubules (fig. 2B, D). We developed an anti-phosphospecific antibody recognizing

phosphorylated Map4-serine 787 (p-S787-Map4), a target site for inhibition by Cdk1 *in vivo* (*22*).

Indeed, when the fusion proteins tGFP-Map4-wild type (WT) and tGFP-Map4-S787A (S787A), in

which Map4 serine 787 was mutated into non-phosphorylatable alanine, were expressed and isolated

from mitotic cells, the antibody recognized wild type but not the mutant version (fig. S7). Probing

Ips of comparable amounts of endogenous Map4 from S and P fractions of Meta cells for p-S787-

Map4 showed that the signal was readily detected from S fraction while hardly from P fraction (fig.

3A). In addition, treating P fraction Map4 with active Cdk1, *in vitro*, restored strong p-S787-Map4

signal and slower Map4 migration (fig. 3B). These data indicate that Map4 is dephosphorylated at

Cdk1-dependent sites when bound to microtubules. Map4 and ch-Tog have been shown to physically

interact with Cdk1, mediating association of the kinase with microtubules (*6, 23*). Our data from

Wee1 downregulation experiments suggested that ch-Tog and Map4 distribution in the P fraction

depended on i-Cdk1. Indeed, Ips of comparable amounts of Map4 or ch-Tog from the S and P fraction

of Meta cells showed that the MAPs selectively interacted with Wee1 and i-Cdk1 in P fraction (figs.

3C, D). Together, these observations suggest that the interaction of microtubule-associated MAPs

with i-Cdk1 was instrumental for their dephosphorylation (fig. 3A, C, D). In mitosis, major protein

phosphatases like PP1 and PP2A are directly or indirectly inhibited by Cdk1 (*1*). PP1 catalytic subunit

is directly inhibited by Cdk1 phosphorylation, in particular the alpha isoform, PP1α, at threonine 320

(p-T-320-PP1α) and phosphorylation in analogous sites inhibits other PP1 isoforms (*24*). PP1 can,

however, rapidly autoactivate by dephosphorylating itself, provided no Cdk1 activity in the proximity

(*24*). In addition, PP1 often interacts with its substrates through specific domains (*25*). Thus, first we

asked whether PP1 interacted with Map4 and ch-Tog and found that the MAPs interacted with PP1α

only in the P fraction (figs. 3C, D). Because of the presence of i-Cdk1, we hypothesised that PP1α in

P fraction was dephosphorylated at the Cdk1 inhibitory site. Indeed, when comparable amounts of

PP1α were analyzed from S and P fractions of Meta cells, the p-T-320-PP1 signal was substantially

detectable in S fraction and barely in P fraction (fig. 3E). In addition, probing comparable amount of

PP1α from total S fraction and from Map4 and ch-Tog Ips showed that MAP-associated PP1α was

dephosphorylated at the inhibitory Cdk1-dependent site (fig. 3F). Moreover, active PP1α was able to

dephosphorylate Map4-S787 *in vitro* (fig. 3G). Together, these data suggest that i-Cdk1 coopts active

PP1 to reverse inhibitory phosphorylation of Map4 and ch-Tog, and possibly other MAPs, for spindle

microtubule growth. When prometaphase-arrested control-siRNA- and Wee1-siRNA-treated cells

were released into M/C medium for 80 min, PP1α was readily detectable in Map4 and ch-Tog Ips

from total extracts of control but not of Wee1-siRNA-treated cells (fig. 3H, lane 6). Addition of low-

RO to Wee1-siRNA-treated cells, that restored i-Cdk1 and spindle assembly (see fig. 2E), also

restored efficient PP1α binding to Map4 and ch-Tog (fig. 3H, lane 9). Thus, i-Cdk1 allowed Map4

and ch-Tog interaction with active PP1 to promote their dephosphorylation. The Map4 sequence

KDVRW (from aa 318 to 322) matched a rather conserved PP1 binding motif (R/KxVxF/W; where

x is any aa) (*25*). We mutated it into KDARA in tGFP-Map4-WT expression vector to produce tGFP-

Map4-PP1-KO, a mutant impaired in PP1 binding. Indeed, when tGFP-Map4-WT and tGFP-Map4-

PP1-KO were expressed at comparable levels and isolated from Meta cells, PP1α substantially bound

tGFP-Map4-WT but not tGFP-Map4-PP1-KO (figs. 3J, K). When the distribution in S and P fractions

of tGFP-Map4-WT- and tGFP-Map4-PP1-KO-transfected Meta cells was analyzed, tGFP-Map4-WT

was readily detectable in the P fraction while tGFP-Map4-PP1-KO remained substantially in S

fraction (fig. 3I). We concluded that i-Cdk1 is required for spindle localized interaction of

microtubule-stabilizing MAPs with active PP1 and for PP1-dependent reversal of their inhibitory

phosphorylations, operated by Cdk1 and other mitotic kinases, in order to locally restore their

microtubule growth-promoting properties.

Probing the cyclin B1 Ips form S and P fractions of Meta cells, shown in fig. 2C, for γ-tubulin revealed

that it interacted with Cdk1 in P fraction (fig. 4A). Conversely, probing γ-tubulin Ips from S and P

fraction of Pro cells showed that it selectively interacted with i-Cdk1 and Wee1, in P fraction (fig.

4B). In Meta cells P fraction, i-Cdk1 and Wee1 increased relatively to the P fraction of Pro cells,

suggesting a mutual relationship between microtubule polymerization and i-Cdk1 formation (see figs.

2B, D and 3C, D). This may reflect further Wee1 recruitment and i-Cdk1 formation at intra-spindle

microtubule nucleation points, in a γ-tubulin-dependent manner, but also further i-Cdk1 formation by

growing spindle microtubules, in a mutual fashion, as suggested by the interaction of i-Cdk1 with

Map4 in Meta cell P fraction (see fig. 3C) (*17, 18*). When the repartition of the Cdk1 inhibitory

pathway players between S and P fractions in Pro and Meta cells was probed on blots where the P

fraction was enriched 10 folds over the S fraction, it appeared more evident that the increase of Wee1

in P fraction of Meta cells relatively to Pro cells was microtubule-dependent, Myt1 also increased

albeit to a lesser extent (fig. 4C). No changes in distribution between fractions in Meta and Pro cells

were, instead, detected for Cdc25 phosphatase family members (fig. 4C). Thus, polymerizing spindle

microtubules appeared to bind Wee1 but not with Cdc25, further promoting i-Cdk1 insulated from

the cytoplasmic Cdk1 autoactivatory loops (fig. 4C) (*8, 20, 21*). To corroborate the hypothesis that

nucleating microtubules compartimentalized the Cdk1 control fostering i-Cdk1 formation, we used

concentrated cytoplasmic cell extracts from prometaphase-arrested HeLa that were incubated with nocodazole or with the microtubule stabilizer taxol for 20 min at 23ºC, the pellets were then eluted and analyzed (*25*). The microtubular pellets aggregated with Wee1 and i-Cdk1 but not with Cdc25 family members (fig. 4d). The presence of Wee1 on spindles in cells was also confirmed by IF (fig. 4E). Thus, in mitosis, positive feedback loops keep Cdk1 active in the cytoplasm to clear peripheral cytoplasmic microtubules by phosphorylating MAPs for safe chromosome movements and possibly to destabilize spindle microtubules that do not, or weakly, bind chromosomes (fig. 4F). At the same time, centrosomes, and perhaps intra-spindle γ-tubulin nucleating centers, along with their nucleating microtubules foster localized i-Cdk1 for PP1-dependent reversal of inhibitory MAP phosphorylation and spindle microtubule growth (fig. 4f) (*3, 17, 18, 27*). Compartimentalized control of Cdk1 activity might also be contributed by other mechanisms in addition to inhibitory phosphorylation of Cdk1 and relevant to terminate the SAC-sustaining action of Cdk1 upon stable microtubule-kinetochore attachments (*15, 28, 29*).

References

1. F. Verde, J. C. Labbé, M. Dorée, E. Karsenti, Regulation of microtubule dynamics by cdc2 protein kinase in cell-free extracts of Xenopus eggs. *Nature. 343,* 233-238 (1990).

2. A. Lieuvin, J. C. Labbé, M. Dorée, D. Job, Intrinsic microtubule stability in interphase cells. *J Cell Biol* **124**, 985-996 (1994).

3. N. Mchedlishvili, H. K. Matthews, A. Corrigan, B. Baum, Two-step interphase microtubule disassembly aids spindle morphogenesis. *BMC Biol 16,* 14 (2018).

4. Chang, W. et al. Phosphorylation of MAP4 affects microtubule properties and cell cycle progression. J Cell Sci **114**, 2879-2887 (2001).

5. Vasquez ,R. J., Gard, D. L. & Cassimeris, L. Phosphorylation by CDK1 regulates XMAP215 function in vitro. *Cell Motil Cytoskeleton* **43**, 310-321 (1999).

6. Ookata, K. et al. Cyclin B interaction with microtubule-associated protein 4 (MAP4) targets p34cdc2 kinase to microtubules and is a potential regulator of M-phase microtubule dynamics. *J Cell Biol* **128**, 849-862 (1995).

7. Gergely, F., Draviam, V. M. & Raff, J. W. The ch-TOG/XMAP215 protein is essential for spindle pole organization in human somatic cells. *Genes Dev* **17**, 336-341 (2003).

8. Hunt, T. On the regulation of protein phosphatase 2A and its role in controlling entry into and exit from mitosis. *Adv Biol Regul* **53**, 173-178 (2013).

9.   Krek, W. & Nigg, E. A. Mutations of p34cdc2 phosphorylation sites induce premature mitotic
events in HeLa cells: evidence for a double block to p34cdc2 kinase activation in vertebrates.
EMBO J **10**, 3331-3341 (1991).

10. D'Angiolella, V., Palazzo, L., Santarpia, C., Costanzo, V. & Grieco, D. Role for non-
proteolytic control of M-phase-promoting factor activity at M-phase exit. *PLoS One* **2**, e247
(2007).

11. Oh, J. S., Susor, A. & Conti, M. Protein tyrosine kinase Wee1B is essential for metaphase II
exit in mouse oocytes. *Science* **332**, 462-465 (2011).

12. Vassilev, L. T. et al. Selective small-molecule inhibitor reveals critical mitotic functions of
human CDK1. *Proc Natl Acad Sci U S A* **103** (2006).

13. Visconti, R. et al. The Fcp1-Wee1-Cdk1 axis affects spindle assembly checkpoint robustness
and sensitivity to antimicrotubule cancer drugs. *Cell Death Differ*. **22**, 1551-1560 (2015).

14. Serpico, A. F., D'Alterio, G., Vetrei, C., Della Monica, R., Nardella, L., Visconti, R. & Grieco,
D. Wee1 rather than Plk1 is inhibited by AZD1775 at therapeutically relevant concentrations.
Cancers **11**, 819 (2019).

15. Serpico, A. F. & Grieco, D. Recent advances in understanding the role of Cdk1 in the Spindle
Assembly Checkpoint. *F1000Res* **9**, F1000 Faculty Rev-57 (2020).

16. Silljé, H. H. W. & Nigg, E. A. Purification of mitotic spindles from cultured human cells.
Methods **38**, 25-28 (2006).

17. David, A. F. et al. Augmin accumulation on long-lived microtubules drives amplification and
kinetochore-directed growth. *J Cell Biol* **218**, 2150-2168 (2019).

18. Thawani, A. et al. The transition state and regulation of γ-TuRC-mediated microtubule
nucleation revealed by single molecule microscopy. *Elife* **9**, e54253. (2020).

19. Solomon, M. J., Lee, T. & Kirschner, M. W. Role of phosphorylation in p34cdc2 activation:
identification of an activating kinase. *Mol Biol Cell* **3**, 13-27 (1992).

20. Novák, B. & Tyson, J. J. Mechanisms of signalling-memory governing progression through
the eukaryotic cell cycle. *Curr Opin Cell Biol*. **69**, 7-16 (2021).

21. Pomerening, J. R., Sontag, E. D. & Ferrell, J. E. Jr. Building a cell cycle oscillator: hysteresis
and bistability in the activation of Cdc2. *Nat Cell Biol*. **5**, 346-351 (2003).

22. Ookata, K. et al. MAP4 is the in vivo substrate for CDC2 kinase in HeLa cells: identification
of an M-phase specific and a cell cycle-independent phosphorylation site in MAP4.
Biochemistry **36**, 15873-15883 (1997).

23. Charrasse, S., Lorca, T., Dorée, M. & Larroque, C. The Xenopus XMAP215 and its human homologue TOG proteins interact with cyclin B1 to target p34cdc2 to microtubules during mitosis. *Exp Cell Res* **254**, 249-256 (2000).

24. Kwon, Y. G., Lee, S. Y., Choi, Y., Greengard, P. & Nairn, A. C. Cell cycle-dependent phosphorylation of mammalian protein phosphatase 1 by cdc2 kinase. *Proc Natl Acad Sci U S A* **94**, 2168-2173 (1997).

25. Meadows, J. C. et al. Spindle checkpoint silencing requires association of PP1 to both Spc7 and kinesin-8 motors. *Dev Cell* **20**, 739-750 (2011).

26. Visconti, R., Palazzo, L., Della Monica, R. & Grieco, D. Fcp1-dependent dephosphorylation is required for M-phase-promoting factor inactivation at mitosis exit. *Nat Commun* **3**, 894 (2012).

27. Woodruff, J. B., et al. The centrosome is a selective condensate that nucleates microtubules by concentrating tubulin. *Cell* **169**, 1066-1077 (2017).

28. Yim, H. & Erikson R. L. Cell division cycle 6, a mitotic substrate of polo-like kinase 1, regulates chromosomal segregation mediated by cyclin-dependent kinase 1 and separase. *Proc Natl Acad Sci U S A* **107**,19742-19747 (2010).

29. Schmidt, A. K. et al. The p53/p73 - p21[CIP1] tumor suppressor axis guards against chromosomal instability by restraining CDK1 in human cancer cells. *Oncogene* **40**, 436-451 (2021).

Acknowledgements

The authors acknowledge V. E. Avvedimento for suggestions, R. Visconti and A. Romano for technical advice. D.G. acknowledges AIRC, Associazione Italiana per la Ricerca sul Cancro, for support (IG grant 2017; Id. 19851 to D.G.).

Author contributions

A.F.S. and D.G. designed and performed experiments and analyzed data. C.P. and F.F. performed immunoblot and immunofluorescence experiments and analyzed data. D. G. wrote the manuscript. A.F.S. revised the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no conflict of interest.

303   **Figures**

Fig. 1. Wee1 downregulation affects spindle assembly. (**A**) hTERT-RPE1 cells were arrested at

G2 by high-RO treatment, released and fixed at 60 and 80 min of incubation in M/C medium (see

Methods section). Control: Wee1-siRNA-treated cells previously transfected with siRNA-resistant

Wee1 expression vector; Wee1-siRNAs: Wee1-siRNA-treated cells previously transfected with

empty vector; Wee1-siRNAs + low-RO: Wee1-siRNAs to which low-RO was added by 60 min from

G2-arrest release. (**B**) hTERT-RPE1 cells were arrested at prometaphase with nocodazole, released

and fixed after 80 min of incubation in M/C medium (see Materials and Methods). Control: cells

treated with non-targeting siRNAs; Wee1-siRNAs: just Wee1-siRNA-treated cells; Wee1-siRNAs +

low-RO: Wee1-siRNA-treated cells to which low-RO was added by 60 min from prometaphase-arrest

release. Cells were fixed and stained for CREST (centromere marker, green), α-tubulin (α-Tub, red)

and DNA (blue). Addition of vehicle (DMSO) had no effect on spindle assembly in Wee1-siRNA-

treated cells. Graphs: quantitation of cells containing normal aligned (blue bar) or abnormal

misaligned (orange bar) spindles at 80 min incubation; shown are data from two independent

experiments (1-4, experiment 1; 5-8, experiment 2), around 100 cells were scored in 4 independent

microscopy slide fields. Scale bar: 5 μm.

Fig. 2. i-Cdk1 localizes at spindle structures and is required for spindle assembly. (**A**) Growing
hTERT-RPE1 and HeLa cells were treated for 20 min with MG-132, fixed and stained for indicated
antigens (γ-Tub; γ-tubulin). (**B**) Prometaphase-arrested hTERT-RPE1 cells were released in M/C
medium for 60 min and either nocodazole was added back at time 0 of incubation (Pro), or just vehicle
(Meta). S and P fractions were probed for indicated antigens. Bottom: IFs of representative cells
probed for indicated antigens. (**C**) Cyclin B1 Ips from Meta cell S and P fractions probed for indicated
antigens. (**D**) Meta cells obtained as described in (B) were further treated with vehicle (Meta-) or
nocodazole (-Pro) from 60 min to 80 min of incubation. S and P fractions were probed for indicated
antigens. (**E**, **F**) hTERT-RPE1 cells were treated with non-targeting-siRNAs (Control) and Wee1-
siRNAs, prometaphase-arrested and released in M/C medium for 80 min incubation as described in
Fig. 1b and a portion of Wee1-siRNA-treated cells received low-RO at 60 min post release. (**E**) S and
P fractions probed for indicated antigens; (**F**) Cells were fixed and stained for indicated antigens.
Scale bar: 5 μm.

Fig. 3. i-Cdk1 assists spindle-localized MAP interaction with PP1 and dephosphorylation. From Meta hTERT-RPE1 cells: (**A**) Map4 Ips from S and P fractions were probed for indicated antigens. (**B**) Map4 Ips were incubated in kinase reaction buffer at 37° C for 15 min: lane 1, from S fraction; lanes 2, 3 and 4, from P fraction – or + active Cdk1 or + active Cdk1 + RO-3306, respectively. After incubation, Ips were probed for indicated antigens. (**C**) Map4 Ips and (**D**) ch-Tog Ips from S and P fractions were probed for indicated antigens. **e**, S and P fractions were probed for indicated antigens. (**F**) total S fraction and Map4 and ch-Tog Ips from P fraction were probed for indicated antigens. (**G**) Map4 Ips from S fraction were incubated in phosphatase reaction buffer at 37° C for 15 min in - or + active PP1α, then probed for indicated antigens. (**H**) and (**I**) Non-targeting-siRNA- (Control) and

Wee1-siRNA-treated (Wee1) hTERT-RPE1 cells released from prometaphase-arrest into M/C medium for 80 min incubation. A sample of control cells received nocodazole at time 0 (Noco) and a sample of Wee1-siRNAs cells received low-RO at 60 min of incubation (RO). Cells were lysed in high salt buffer and (**H**) Map4 or (**I**) ch-Tog Ips were probed for indicated antigens (lanes 1-4, total lysate; Mk Ip, mock Ips from Noco cell lysates). (**J**, **K**) hTERT-RPE1 cells were transfected with tGFP-Map4-wild type (WT) and tGFP-Map4-PP1-KO expression vectors, arrested at metaphase and either (J) lysed in high salt buffer, tGFP Ips were probed for indicated antigens or (K) S and P fraction isolated and probed for indicated antigens.

Fig. 4. Nucleating microtubules bind and foster i-Cdk1. (**A**) Cyclin B1 Ips described in fig. 2C were probed for γ-tubulin. (**B**) γ-tubulin Ips from S and P fraction of Pro hTERT-RPE1 cells were probed for indicated antigens (P fraction enriched 10 folds over S fraction). (**C**) S and P fraction from Pro, Meta, Meta- and -Pro, as described in Fig. 2b, d, were probed for indicated antigens on blots in which the P fraction was enriched 10 folds over S fraction. (**D**) Eluates from pellets of mitotic HeLa cell extracts (P) incubated with nocodazole (Noco) or taxol (Tax) were probed for indicated antigens; Tot: 1/20 of total extract. **E**, growing hTERT-RPE1 cells were treated for 20 min with MG-132, fixed and stained for indicated antigens. Scale bar: 5 μm. (**F**) a synoptic scheme depicting Cdk1 activity control in mitosis. At prophase, Cdk1 activity, sustained by positive feedback loops in the cytoplasm, destabilizes cytoplasmic microtubules by directly (continuous arrow) or indirectly (dotted arrow) phosphorylating MAPs. Centrosomes, however, maintain/promote inhibitory phosphorylation of a small Cdk1 fraction by selectively interacting with Wee1 and excluding Cdc25. From prometaphase to metaphase, further interaction of Wee1, inhibited Cdk1 and active PP-1 with microtubule-stabilizing MAPs expands from centrosomes, and possibly from intra-spindle γ-tubulin-dependent nucleating centers, to promote reversal of inhibitory MAP phosphorylation and spindle microtubule growth.

Supplementary Materials

Materials and Methods

Cell lines and cell culture

HeLa cells were grown in Roswell Park Memorial Institute Medium (RPMI-1640; Sigma-Aldrich), hTERT-RPE1 cells were grown in Dulbecco's Modified Eagle Medium: Nutrient Mixture F-12 (DMEM/F12; Gibco, Thermo Fisher Scientific), both supplemented with 10% fetal bovine serum (FBS; Gibco), 1% GlutaMAX-supplement (Gibco), 1% penicillin/streptomycin (Euroclone), at 37° C with 5% CO_2.

Cell treatments

Transfections were performed using linear polyethylenimine (PEI; Polysciences, Inc) or MegaTran 2.0 Transfection Reagent (OriGene Technologies, Inc.). The expression plasmids for turbo-GFP-Map4-wild type (tGFP-Map4-WT), turbo-GFP-Map4 rendered non-phosphorylatable at serine 787 (tGFP-Map4-S787A) and turbo-GFP-Map4 defective in PP1 binding Map4 (tGFP-Map4-PP1-KO), in which the Map4 peptide KDVRW (from aa 318 to 32; human Map4 isoform 1) was mutated into KDARA, were generated and purchased from OriGene Technologies Inc. Cdk1WT and cdk1AF expression plasmids were purchased from Addgene. 3XFlag-Wee1 expression construct was obtained as described[13]. SiRNAs targeting the 3′-UTR of human WEE1 (5′-GGGCUUU AUUACAGACAUAUU-3′, 5′-GUACAUAGCUGUUUGAAAUUU-3′ and 5′-UGUAAACUUGUAGCAUUAUU-3′) were purchased from Dharmacon Inc. For efficient knockdown cells were transfected with 25 nM of siRNAs duplex using DharmaFECT 1 siRNA Transfection Reagent (Dharmacon Inc.). For siRNA treatment and complementation experiments in synchronized cells, cells were mock- or 3XFlag-Wee1 expression construct-transfected 24 hours prior to treatment with non-targeting or specific siRNAs. For synchronization of siRNA-treated cells, 6 hours after siRNA-treatment, cells were incubated with RO-3306 at 9 μM (high-RO; Calbiochem) for further 16 hours of incubation, to have G2-arrsted cells, or with nocodazole (Abcam) at 1 μg/ml and 100 ng/ml for further 16 or 14 hours of incubation for hTERT-RPE1 and HeLa cells, respectively, to have prometaphase-arrested cells (Pro cells). To obtain metaphase-arrested cells (Meta cells), Pro cells were collected and washed twice with fresh medium and twice with phosphate buffer saline (PBS; Euroclone) solution before plating into fresh medium containing MG-132 (20 μM; Calbiochem) and cycloheximide (CHX; 60 μg/ml; Santa Cruz Biotechnology, Inc) for further 60 min incubation. Meta- and -Pro cells were obtained by adding either DMSO or nocodazole (1 μg/ml) to Meta cells at 60 min and prolonging incubation for further 20 min. Where indicated AZD1775 (Selleckchem) was added at 1 μM. Where indicated, partial reversal of Cdk1 activity was obtained by adding RO-3306 at 0.5 μM (low-RO).

Cell fractionation and extracts

To isolate spindle-bound protein we adapted a previously described method[15]. All the steps, except the final elution, were performed at room temperature (rt) to preserve microtubules. Cells were harvested by centrifugation and washed once with PBS containing taxol (1 μM; Sigma-Aldrich). 5 x 10^6 cells were resuspended in 800 μl of fractionation lysis buffer (FLB; 40 mM β-glycerophosphate, 15 mM MgCl$_2$, 20 mM EGTA, 0,25% Igepal, 20 mM Hepes, 5 μM taxol, 6 μg/ml latrunculin B; Sigma-Aldrich) supplemented with 300 μg/ml RNase A and 120 U/ml DNase I (Roche). After careful resuspension, samples were incubated in thermomixer (Eppendorf thermomixer comfort) at 34 °C for 20 min with constant shaking at 1,200 rpm. Lysates were spun at 6,800 rpm (Eppendorf centrifuge 5425) for 2 min: the supernatant fractions were collected and transferred into new tubes and supplemented with 250 mM NaCl (S fraction), the pellet fractions were resuspended in FLB + 300 μg/ml RNase A and 120 U/ml DNase I and again incubated in thermomixer at 34 °C for 20 min with constant shaking at 1,200 rpm. Lysates were collected again by centrifugation and the pellets were washed twice with washing buffer (WB; 5 mM Hepes, 5 μM taxol). Finally, the pellet fractions were resuspended in 200 μl of FLB minus taxol and supplemented with 250 mM NaCl (P fraction). Both supernatant and pellet fractions were incubated on ice, to help destabilize microtubules, for 30 min and then spun at 3,200 rpm at 4 °C for 10 min (Eppendorf centrifuge 5424 R) and the supernatants of S and P fraction further analyzed by Ibs or Ips. To induce microtubule polymerization and analyze microtubule-bound proteins in mitotic cell extracts, HeLa cell extracts were obtained as previously described[24]. Cell extracts were supplemented with 1 mM GTP (Sigma-Aldrich), 50 μl aliquots were treated with nocodazole (1μg/ml; Abcam), as control, or taxol (5 μM; Sigma-Aldrich) for 20 min at rt, diluted 10 times with FLB and spun at 6,800 rpm (Eppendorf centrifuge 5425) for 2 min. The pellets were washed once more in FLB and finally resuspended in 30 μl FLB supplemented with 250 mM NaCl, incubated on ice for 30 min and then spun 13,200 rpm at 4 °C for 10 min (Eppendorf centrifuge 5424 R). Supernatants were analyzed by Ibs; total samples = 5 μl of total mitotic extract.

Antibodies

Antibodies used for immunoblotting according to standard protocols: mouse anti-cdc2 (1:500; BD Biosciences), rabbit anti-phospho-tyrosine-15-cdc2 (P-Y15-cdc2; 1:1000; Boster Bio), rabbit anti-phospho-threonine-14-cdc2 (P-T14-cdc2; 1:1000; Cell Signaling Technology Inc.; CST), rabbit anti-Wee1 (1:1000; CST), rabbit anti-Myt1 (1:1000; CST), rabbit anti-Cdc25C (1:1000; CST), rabbit anti-phospho-threonine-320-PP1α (P-T320-PP1; 1:1000; CST), rabbit anti-cyclin B1 (1:2000; Bethyl Laboratories, Inc.), mouse anti-γ-tubulin (1:2000; Sigma-Aldrich), mouse anti- α-tubulin (1:2000;

Sigma-Aldrich), mouse anti-PP1γ (1:500; Santa Cruz Biotechnology, Inc.), rabbit anti-Map4 (1:2000;

Bethyl Laboratories Inc.), rabbit anti-NuMA (1:1000; Novus Biologicals), rabbit anti-ch-Tog

(1:1000; Abcam), mouse anti-turboGFP (1:2000; OriGene Technologies). Rabbit polyclonal antibody

against phosphorylate serine 787 of human Map4 (P-S787-Map4) was raised using C-

DAKAPEKRA[Sp]PSKPA-coNH2 peptide as immunogen; peptide synthesis, rabbit inoculation,

serum production and cross-affinity antibody purification were carried out by CovalAb S.A.S. Sheep

anti-mouse IgG peroxidase-(HRP) linked and donkey anti-rabbit IgG HRP-linked whole antibodies

(1:2000; GE Healthcare). Antibodies used for IF: rabbit anti-phospho-tyrosine-15-cdc2 (P-Y15-cdc2;

1:50; CST), anti-cdc2 (1:100; BD Biosciences), mouse anti-γ-tubulin (1:1000; Sigma-Aldrich),

mouse anti-α-tubulin (1:1000; Sigma-Aldrich), rabbit anti-γ-tubulin (1:1000; Sigma-Aldrich), human

anti-centromere (CREST; 1:50; Antibodies Incorporated), rabbit anti-Map4 (1:1000; Abcam), rabbit

anti-ch-Tog (1:400; Proteintech Group, Inc.), rabbit anti-Wee1 (1:100; CST), goat anti-human IgG

rhodamine conjugated (4 µg/ml; Santa Cruz Biotechnology, Inc.), rabbit anti-human IgG FITC

conjugated (4 µg/mL; Dako Agilent), Alexa Fluor 594 donkey anti-rabbit IgG (2 µg/mL; Invitrogen),

Alexa Fluor 594 donkey anti-mouse IgG (4 µg/ml; Invitrogen), Alexa Fluor 488 donkey anti-rabbit

IgG (2 µg/ml; Invitrogen), Alexa Fluor 488 goat anti-mouse IgG (4 µg/ml; Invitrogen). Antibodies

used for Ips: rabbit anti-Map4 (2µg/mL; Bethyl Laboratories Inc.), rabbit anti-chTOG (2µg/mL;

Abcam), mouse anti-γ-tubulin (2µg/mL; Sigma-Aldrich). Antibody-conjugated beads used for Ips:

mouse anti-cyclin B1 agarose (Santa Cruz Biotechnology), mouse anti-turboGFP agarose (OriGene),

protein A/G PLUS-agarose (Santa Cruz Biotechnology), mouse anti-γ-tubulin agarose conjugate

(Santa Cruz Biotechnology).

Immunoblotting and immunoprecipitation

Immunoblots (Ib) were performed as described[26]. For immunoprecipitations (Ips) from S fraction and

and P fraction eluates, samples were diluted 6 folds in lysis buffer (LB; 80 mM β-glycerophosphate,

15 mM $MgCl_2$, 20 mM EGTA, 20 mM Hepes pH 7.4, 100 mM NaCl, 0,1 % Igepal; Sigma-Aldrich)

supplemented with 1 tablet of phosphatase inhibitor cocktail (PhosSTOP; Roche) and incubated with

antibody and protein A/G PLUS-agarose beads or with agarose bead-conjugated antibody overnight

at 4 °C in constant rotation. Beads were washed twice in LB and beads-bound proteins were eluted

by boiling in SDS loading buffer. For total cell lysates, 5×10^6 cells were lysed in 180 µl high salt

lysis buffer (HSB; 16 mM β-glycerophosphate, 3 mM $MgCl_2$, 4 mM EGTA, 0,5 mM DTT, 20 mM

Hepes, 0,2 % Igepal, 250 mM NaCl; Sigma-Aldrich) supplemented with 1 tablet of phosphatase

inhibitor cocktail (PhosSTOP; Roche) to preserve phosphorylation. After 30 min incubation on ice,

lysates were spun at 13,200 rpm for 10 min at 4° C (Eppendorf centrifuge 5424 R). Cleared lysates

were incubated with antibody and protein A/G PLUS-agarose beads or with agarose bead-conjugated

antibody overnight at 4 °C in constant rotation. Beads were washed twice in LB and beads-bound

proteins were eluted by boiling in SDS loading buffer. For kinase reactions, after the two LB washes,

Ips were washed once in kinase reaction buffer (KRB; 80 mM β-glycerophosphate, 15 mM MgCl$_2$,

20 mM EGTA, 20 mM Hepes pH 7.4, 100 μM ATP; Sigma-Aldrich) and incubated for 20 min at 37°

C in KRB -/+ recombinant active Cdk1 (0.05 mg/ml; Sigma-Aldrich) and + active Cdk1 + RO-3306

(10 μM). For phosphatase reactions, after the two LB washes, Ips were washed once in phosphatase

reaction buffer (PRB; 20 mM Hepes pH 7.4, 150 mM NaCl, 1 mM DTT, 1 mM MnCl$_2$; Sigma-

Aldrich) and incubated in PRB -/+ 0.01 mg/ml recombinant human PP1α (Novus Biologicals) for 20

min at 30° C.

Immunofluorescence staining and microscopy

Cells were grown onto poly-D-lysine (0,1 mg/ml; Sigma-Aldrich) coated glass coverslips and treated

as described in the text. Before immunofluorescence (IF) procedure initiation, coverslips were

washed twice with PBS, then cells were fixed with 4 % paraformaldehyde (Sigma-Aldrich) in PBS

(Euroclone) for 10 min at rt and with a methanol (99,9 %; Exacta + Optech) additional step at -20° C

for 10 min, the latter except for α-tubulin staining. Cells were permeabilized with 0,25-0,5 % Triton

X-100 (Sigma-Aldrich) in PBS for 15-30 min, washed once with PBS and incubated with 3% bovine

serum albumin (BSA; Sigma-Aldrich) in PBS for 1 hour at rt. Coverslips were transferred into a

humidity chamber and incubated with primary antibodies in 1,5 % (w/v) BSA-PBS solution for 2-12

hour at 4° C. Samples were washed three times and incubated with fluorescently labelled secondary

antibodies, dissolved in 1,5 % BSA-PBS solution, for 1 hour at rt. DNA was stained with Hoechst

33258 (1 μg/ml; Invitrogen) by incubation for 10 min. Finally, samples were washed four times and

mounted with Mowiol 40-88 (Sigma-Aldrich) on a glass side. Fixed cells were observed using LSM

980 inverted confocal microscope (Zeiss) with a 63X/1,4 oil objective. Representative images were

obtained collecting 3 Z-stack series over 42 μm. The acquisitions were deconvoluted and projected

into one plane using the LAS-AF software.

Statistics and reproducibility

For quantitative analyses of immunofluorescence experiments investigators were blinded to sample

allocation. Metaphases were visually scored and those with more than three chromosomes falling

outside the two internal quarters of the interpolar distance were considered misaligned; around 100

cells were scored in 4 fields of each slide for each experimental condition; each experiment was

repeated 3-5 times with similar results. Experiments analyzed by immunoblotting were repeated 2-8

times with similar results.

Supplementary Figures

Figure S1

Fig. S1. siRNAs treatment of hTERT-RPE1 cells. Lysates of hTERT-RPE1 cells treated with non-

targeting siRNAs (NT) and with Wee1-siRNAs (Wee1) as described in fig.1 were probed for

indicated antigens. A portion of Wee1-siRNAs was previously transfected with a siRNAs-resistant,

Flag-tagged Wee1 expression vector (F-Wee1).

Fig. S2. Expression of an inhibitory-phosphorylation-resistant version of Cdk1 impairs spindle

assembly. hTERT-RPE1 cells were transfected with (**A**) wild type cdk1 (cdk1WT) expression vector

or (**B**) a mutant cdk1 expression vector in which cdk1-threonine 14 and tyrosine 15 are mutated into

non-phosphorylatable alanine and phenylalanine, respectively (cdk1AF). 6 hours post transfection,

cells were arrested at G2 by addition high-RO and for further 16 hours incubation. Cells were fixed

and stained for DNA (blue), α-tubulin (red) and CREST (green) at 60 and 80 min upon release from

G2 arrest. A portion of cdk1AF-transfected cells received low-RO at 60 min; as control vehicle was

added (DMSO). Graphs: quantitation of cells containing normal aligned (blue bar) or abnormal

misaligned (orange bar) spindles at 80 min incubation; shown are data from two independent

experiments, in which around 100 cells were scored in 4 independent microscopy slide fields (1-4,

experiment 1; 5-8, experiments 2). Scale bar: 5 μm. (**C**) samples of cells treated as in (A) and (B)

were lysed prior to release and probed for indicated antigens.

Fig. S3. Downregulation of Wee1 impairs spindle assembly in HeLa cells. HeLa cells were treated

and arrested at G2 as described for hTERT-RPE1 cells in Fig 1a. Cells were fixed and stained for

DNA (blue), α-tubulin (red) and CREST (green) at 60 and 80 min upon release from G2 arrest. (**A**)

Wee1-siRNA-treated cells complemented with siRNA-resistant expression vector (Wee1-siRNAs +

F-Wee1). (**B**) Wee1-siRNA-treated cells (Wee1-siRNAs). A portion of Wee1-siRNAs cells received

low-RO at 60 min; as control vehicle was added (DMSO). Graphs: quantitation of cells containing

normal aligned (blue bar) or abnormal misaligned (orange bar) spindles at 80 min incubation; shown

are data from two independent experiments, in which around 100 cells were scored in 4 independent

microscopy slide fields (1-4, experiment 1; 5-8, experiments 2). Scale bar: 5 μm. **c**, samples of cells

treated as in (A) and (B) were lysed prior to release and probed for indicated antigens.

Fig. S4. Chemical inhibition of Wee1 impairs spindle assembly. (**A**) hTERT-RPE1 cells were

released from a G2 arrest by high-RO treatment into M/C medium and either vehicle (Control) or

AZD1775 was added. Cells were fixed after further 80 min of incubation and stained for α-tubulin

(green), centromeres (red) and DNA (blue). A sample of the AZD1775-treated cells also received

low-RO 60 min post release. Scale bar: 5 μm. (**B**) prometaphase-arrested hTERT-RPE1 were released

into M/C medium and either vehicle (Control) or AZD1775 was added. S and P fraction were

prepared after further 80 min of incubation and probed for indicated antigens. A sample of the

AZD1775-treated cells also received low-RO 60 min post release.

Fig. S5. Downregulation of Wee1 delays mitosis exit. Non-targeting-siRNA- (Control) or Wee1-

siRNA-treated (Wee1) hTERT-RPE1 cells were arrested at prometaphase and released into fresh

medium for 60 min of incubation. Cells were lysed and probed for indicated antigens.

Figure S6

Fig. S6. Mitotic HeLa cell fractionation. Prometaphase-arrested HeLa cells were collected and taken immediately (Pro) or released in M/C medium for 60 min (Meta). Fractionated S and P fractions were probed for indicated antigens (P fraction enriched 4 folds over S fraction).

Figure S7

Fig. S7. Detection of S787-Map4 phosphorylation. hTERT-RPE1 cells were transfected with expression vectors of tGFP-Map4 wild type (WT) or tGFP-Map4 mutant version in which serine 787 of Map4 was mutated into alanine (S787A) and arrested at prometaphase. Total cell lysates (Tot) and relative anti-tGFP Ips were probed for indicated antigens (Mk: mock-transfected cells).

586