1

BET family members Bdf1/2 modulate global transcription initiation and elongation in *Saccharomyces*

cerevisiae

Rafal Donczew[1] and Steven Hahn[1,*]

[1]Fred Hutchinson Cancer Research Center, Division of Basic Sciences, 1100 Fairview Ave N. A1-162,

Seattle, WA 98109

*Correspondence: email: shahn@fredhutch.org, phone: 206 667 5261

Abstract

Human bromodomain-containing BET family members are promising targets for therapy of cancer and immunoinflammatory diseases, but their mechanisms of action and functional redundancies are poorly understood. Bdf1/2, yeast homologues of the human BET factors, were previously proposed to target transcription factor TFIID to acetylated histone H4, analogous to bromodomains that are present within the largest subunit of metazoan TFIID. We investigated the genome-wide roles of Bdf1/2 and found that their important contributions to transcription extend beyond TFIID function, as transcription of many genes is more sensitive to Bdf1/2 than to TFIID depletion. Bdf1/2 co-occupy the majority of yeast promoters and affect preinitiation complex formation through recruitment of TFIID, Mediator and basal transcription factors to chromatin. Surprisingly, we discovered that hypersensitivity of genes to Bdf1/2 depletion results from combined defects in transcription initiation and processive elongation, a striking functional similarity to human BET proteins, most notably Brd4. Our results establish Bdf1/2 as critical for yeast transcription and provide important mechanistic insights into the function of BET proteins in all eukaryotes.

Introduction

Bromodomains (BDs) are reader modules that allow protein targeting to chromatin via interactions with acetylated histone tails. BD containing factors are usually involved in gene transcription and their deregulation has been implicated in a spectrum of cancers and immunoinflammatory and neurological conditions (Fujisawa and Filippakopoulos, 2017; Wang et al., 2021). The bromo and extra-terminal (BET) family is characterized by the presence of a double BD, which has the highest affinity towards hyperacetylated histone H4, and an ET domain that interacts with non-histone proteins (Rahman et al., 2011; Slaughter et al., 2021). BET bromodomain inhibitors have shown promising results in treating both blood cancers and solid tumors (Stathis and Bertoni, 2018). Early studies implicated mammalian BET proteins in regulation of selected lineage-specific genes (Delmore et al., 2011; Lovén et al., 2013), however, it has become clear that their regulatory roles extend to the majority of genes transcribed by RNA Polymerase II (Pol II) (Muhar et al., 2018; Winter et al., 2017).

While most mammalian tissues express three BET factors (Brd2, Brd3, Brd4) (Uhlén et al., 2015), many studies have implicated Brd4 as most important for widespread changes in transcription after BET inactivation (Muhar et al., 2018; Zheng et al., 2021; Zuber et al., 2011). These broad genome-wide expression defects are due to the important roles of Brd4 in both transcription initiation and elongation. Brd4 contributes to recruitment of the transcription coactivator Mediator to enhancers and promoters and cooperates with Mediator in forming nuclear condensates at actively transcribed regions (Bhagwat

et al., 2016; Han et al., 2020; Sabari et al., 2018). Brd4 is also important for productive transcription

elongation but it is unclear what mechanisms are involved. Brd4 associates with the positive

transcription elongation factor b (P-TEFb), but no global defects in P-TEFb recruitment to chromatin

were observed following inactivation of Brd4 or all BET factors (Muhar et al., 2018; Winter et al., 2017).

Proposed alternative mechanisms of elongation control include release of P-TEFb inhibition, histone

chaperone activity, or direct kinase activity (Devaiah et al., 2012; Itzen et al., 2014; Kanno et al., 2014).

Finally, Brd4 was shown to affect preinitiation complex (PIC) formation at promoters, although its roles

in later stages of transcription are believed to be dominant (Kanno et al., 2014; Winter et al., 2017).

TFIID is a conserved Pol II factor that, in yeast, regulates transcription of almost all Pol II-transcribed

genes (Donczew et al., 2020; Huisinga and Pugh, 2004; Warfield et al., 2017). TFIID is comprised of

TBP (TATA binding protein) and 13-14 Tafs (TBP associated factors). TFIID acts as a TBP-DNA loading

factor, a promoter recognition factor and a molecular scaffold guiding PIC assembly (Patel et al., 2020).

Recruitment of metazoan TFIID to promoters is thought to be aided by interactions between two Taf

subunits and chromatin marks that are enriched at the +1 nucleosome. The human Taf1 double BD and

Taf3 PHD domain recognize acetylated histone H4 tails and trimethylated histone H3 lysine 4,

respectively (Jacobson et al., 2000; van Ingen et al., 2008). In contrast, Taf1 and Taf3 in budding yeast

are missing both of these chromatin readers as well as recognizable promoter DNA sequence motifs

such as the INR and DPE that are known to interact with metazoan Tafs (Patel et al., 2020).

Yeast BET family member Bdf1 and its paralogue Bdf2 are genetically redundant, with at least one

required for viability. Deletion of *BDF1* but not *BDF2* results in growth retardation, suggesting a

dominant role for Bdf1 (Matangkasombut et al., 2000). Bdf1 binds Taf7 and recognizes acetylated

histone H4. Both of these functions are required to support cell growth and transcription at several loci.

Based on these findings, it was proposed that Bdf1 double BD substitutes for the missing BDs of yeast

Taf1, linking yeast TFIID to promoter-enriched chromatin marks (Matangkasombut et al., 2000;

Matangkasombut and Buratowski, 2003). Gene deletion or targeted depletion of Bdf1 revealed a small

contribution of Bdf1 to Taf1 recruitment genome-wide with a bias towards a group of genes classified

as "TFIID-dominated" and led to a modest defect in transcription at a limited gene set (Durant and Pugh,

2007; Joo et al., 2017; Ladurner et al., 2003).

By applying degron-mediated protein depletion and monitoring nascent mRNA levels, we recently

showed that yeast protein-coding genes can be classified into two broad categories based on

transcription changes after rapid depletion of the coactivators TFIID and SAGA (Donczew et al., 2020).

We found that the majority of yeast genes are strictly TFIID-dependent while a subset, which we termed

coactivator-redundant (CR) genes, are co-regulated by TFIID and SAGA. This latter category overlaps
~50% with the gene set earlier termed "SAGA-dominated" (Huisinga and Pugh, 2004). In contrast, long-
term ablation of SAGA via gene deletions showed that nearly all yeast genes are dependent on
chromatin modifications directed by SAGA. However, the molecular basis for the TFIID and CR gene
classes remains unknown. For example, it's not known if there are other factors that preferentially
participate in the TFIID or SAGA-directed pathways.
94
Here, we have examined the functions of Bdf1 and Bdf2 in yeast transcription and we found broad
genome-wide roles for the Bdfs in both transcription initiation and elongation. Rapid Bdf depletion
strongly decreases transcription from the TFIID-dependent genes and, at many genes, the Bdfs are
more important than Tafs for normal levels of transcription. The Bdfs contribute to PIC formation and
TFIID and Mediator recruitment to gene regulatory regions. In addition, at many genes, the Bdfs also
regulate transcription elongation and this role contributes to the strong defects in mRNA synthesis upon
Bdf depletion. These striking functional similarities with mammalian BET factors suggest broad
conservation of BET function in eukaryotes.
103

Results
Transcription of most TFIID-dependent genes is more sensitive to Bdf1/2 than to TFIID depletion
A limitation of previous studies on Bdf1/2 function was that both proteins could not be simultaneously
eliminated since at least one Bdf is required for viability. In this study, we used the auxin-degron system
(Nishimura et al., 2009) to achieve rapid depletion of Bdf1, Bdf2, or simultaneous depletion of both
proteins, and used 4-thioU RNA-seq to monitor changes in newly-synthesized mRNA (Donczew et al.,
2020; Rabani et al., 2011). We first assessed protein degradation after a 30-minute treatment with the
auxin indole-3-acetic acid (IAA) (**Figure S1A**). Bdf1 was efficiently degraded with <10% of protein
remaining in IAA treated samples. Bdf2 degradation was less complete, with ~15% of protein remaining,
so we prepared a *bdf2* deletion strain combined with a Bdf1 degron (Bdf1, *Δbdf2*). Growth of all strains
with Bdf derivatives was similar to wild type (**Figure S1B**) and analysis of 4-thioU mRNA levels in control
DMSO-treated cultures showed that changes in transcription due to these genetic alterations were
minimal (**Figure S1C**).
117

Experiments were done in two or three biological replicates and the variation in nascent mRNA levels
between replicate samples was <30% for nearly all experiments (**Figure S1D and Table S2**). Bdf2
degradation or *bdf2* deletion showed minimal changes in transcription while Bdf1 degradation resulted
in modest defects, in agreement with the genetic phenotypes of *BDF1/2* mutants (**Figures 1A and S1E**

and Table S2). Strikingly, depletion of both Bdf1 and Bdf2, either through a double degron or by
combining a *bdf2* deletion with a Bdf1 degron, resulted in a global collapse of transcription with median
decreases of ~4-fold and ~5.4-fold, respectively (**Figures 1A and S1F and Table S2**). Transcriptional
changes in both Bdf1/2 depletion experiments are highly correlated (r = 0.95) (**Figure S1G**). The results
of the Bdf1 and Bdf1/2 depletion experiments also correlate (r = 0.82), showing redundant roles of Bdf1
and Bdf2 in transcription (**Figure S1H**). We compared results of the Bdf depletion experiment with
previously published results of rapid Taf1 depletion (Donczew et al., 2020). Surprisingly, we observed
that transcription changes upon Bdf or Taf1 depletion correlate but the relationship between them is not
linear (**Figure 1B**). Labeling the points based on the TFIID and CR gene categories revealed that genes
less sensitive to Bdf depletion (on the right of the diagonal line) include almost all coactivator-redundant
(CR) genes and one third of TFIID-dependent genes. Interestingly, most of the TFIID-dependent genes
(68%) are more sensitive to Bdf than Taf1 depletion and, for 12% of them, the difference is 4-fold or
higher (**Table S2**). This result shows that, while Bdfs likely act in conjunction with TFIID at many
promoters, Bdfs also have important TFIID-independent contributions to transcription of many genes.
Comparison of the Bdf results with depletion of Taf13, another TFIID subunit, leads to similar
conclusions (**Figure S1I**).
138
Ribosomal protein (RP) genes are frequently analyzed as a separate gene category due to high
expression levels, common regulatory mechanisms, and important roles in cell growth and stress
response (Zencir et al., 2020). About 95% of RP genes belong to the TFIID-dependent class (Donczew
et al., 2020; Huisinga and Pugh, 2004). Surprisingly, we found that nearly all RP genes are significantly
less dependent on Bdf1/2 compared with TFIID, an uncommon feature for TFIID-dependent genes
(**Figure S1J**).
145
With many TFIID-dependent genes losing >85% of detectable transcription after Bdf1/2 depletion, we
investigated how these changes compare to defects caused by depletion of basal components of the
preinitiation complex (PIC). We depleted basal factors with degron tags on either TFIIA subunit Toa1
or the TFIIH translocase subunit Ssl2 followed by 4-thioU RNA-seq. The results are summarized in a
boxplot divided into CR and TFIID-dependent genes (**Figure 1C and Table S2**). Depletion of Toa1 or
Ssl2 decreases all mRNA transcription independent of gene class with median changes of ~15-fold and
~23-fold, respectively. Interestingly, 38% and 19% of TFIID-dependent genes are similarly sensitive to
Toa1 or Ssl2 depletion as they are to Bdf depletion. Combined, our results show that Bdf dependence
is as good or even a better classifier for the TFIID and CR gene classes than Taf-dependence.
Transcription at many TFIID-dependent genes is severely compromised in the absence of Bdfs while
transcription of CR genes is only weakly affected (**Figure 1C**).

Bdf1 is also a subunit of the SWR1 complex that incorporates the histone variant H2A.Z into promoter-

proximal nucleosomes (Krogan et al., 2003; Zhang et al., 2005). We used the degron system to rapidly

deplete H2A.Z and check if the loss of transcription after Bdf depletion is mediated by defects in H2A.Z

deposition (**Figure S1K**). RNA-seq analysis did not reveal significant defects in gene transcription

following the rapid loss of H2A.Z (**Figure S1L and Table S2**), suggesting that the global transcriptional

defects upon Bdf depletion are not mediated via SWR1 and H2A.Z.

Figure 1. Many genes are more sensitive to Bdf1/2 than to TFIID depletion. (**A**) Transcriptional changes caused by degron depletion of Bdf1/2 for 30 min. Scatter plots comparing expression level (in log scale) with log$_2$ change in transcription per gene measured by 4-thioU RNA-seq. Mean values from replicate experiments are plotted for 5313 genes with detectable signals in all RNA-seq samples collected in this work. (**B**) Scatter plot comparing log$_2$ change in transcription after depleting Bdf1/2 (this work) or Taf1 (Donczew et al., 2020). Data for 4883 genes classified into TFIID-dependent and coactivator-redundant (CR) categories are shown (Donczew et al., 2020). (**C**) Boxplot showing log$_2$ change in transcription for 4883 genes measured by 4-thioU RNA-seq after depleting indicated factors. Genes are grouped into TFIID-dependent and CR categories. (**D**) Phylogenetic analysis of amino acid sequences of individual bromodomains from human Taf1, human Brd4 and yeast Bdf1. (**E**) BLAST global alignment of amino acid sequences within indicated domains of Bdf1 and Brd4. (**F**) Boxplot showing log$_2$ change in transcription after depleting BET factors in yeast (this work) or human cells (Winter et al., 2017). Genes are classified depending on the presence of a consensus TATA box (TATAWAW) in their promoter. A list of TATA-containing promoters in human cells was obtained from Eukaryotic Promoter Database (EPD). A single, most representative promoter per gene was used in this analysis. Results of the Welch's t-test are shown. The asterisks represent p-value with the following cut-off levels: **** - 0.0001, *** - 0.001, ** - 0.01, * - 0.05, ns - >0.05. See also Figures S1 and S2 and Tables S2 and S3.

Bdf1/2 share significant similarities with human BET factors

Bdf1 was initially proposed to constitute a missing part of yeast Taf1 (Matangkasombut et al., 2000), but later work classified Bdf1/2 as members of the BET family based on a conserved domain organization (Wu and Chiang, 2007) (**Figure S2A**). Since Bdfs have important TFIID-independent functions at many genes, we explored similarities between yeast and human BET proteins. Multiple sequence alignment and phylogenetic analysis of individual bromodomains (BDs) of Bdf1, Brd4 and human Taf1 (hTaf1) revealed that Bdf1 BDs, especially BD2, are more closely related to Brd4 than hTaf1 BDs (**Figures 1D and S2B**). Other conserved domains of Bdf1 and Brd4 show ≥50% similarity (**Figure 1E**). We next compared gene-specific responses to Bdf/BET depletion in yeast and human cells. We used a published RNA-seq dataset from an experiment where BET factors were targeted with the chemical degrader dBET6 in the MOLT4 leukemia cell line (Winter et al., 2017) (**Table S2**). Since the CR/TFIID-dependent classes are biased for TATA-containing/TATA-less categories (Donczew et al., 2020; Huisinga and Pugh, 2004) we divided yeast and human genes into two categories based on the presence or absence of a consensus TATA box. We found that TATA-less genes in both systems were significantly more affected by Bdf/BET depletion than TATA-containing genes (**Figure 1F**). Finally, in the 25% most affected genes in each dataset, we observed an extensive overlap in major gene ontology terms (**Figure S2C and Table S3**). Altogether, these results suggest a common biological role of BET factors in yeast and human cells, a conclusion supported by other results presented below.

Bdf1/2 and Taf1 have similar genome-wide binding patterns

We recently mapped Taf1 and other TFIID subunits using an improved ChEC-seq method, where DNA cleavage by protein-MNase fusions is applied to map genome-wide binding (Donczew et al., 2021, 2020). TFIID is detectable at over 3000 yeast promoters and is found at both TFIID-dependent and CR genes, which agrees with its general role in transcriptional regulation. We used ChEC-seq to map Bdf1 and Bdf2 binding genome-wide. Comparison of Bdf1/2 and Taf1 ChEC-seq showed similar binding patterns with broad peak spanning between -1 and +1 nucleosomes and signals extending into the gene body, especially for Bdf1 (**Figures 2A and S3A**). We identified 3700, 3466 and 3103 promoters bound by Bdf1, Bdf2 and Taf1, respectively. Data for Bdf1 and Bdf2 confirm redundancy between both factors. There is an extensive overlap of promoters bound by both Bdfs and the signals at common bound promoters are highly correlated, although Bdf1 signals are consistently stronger, in agreement with its dominant role (**Figures 2B and S3B and Table S4**). There is also extensive overlap between Bdf1 and Taf1 bound promoters (**Figure 2B**), although Bdf1 is found less frequently at CR genes (**Figure 2C**), consistent with the lesser role of Bdfs at this gene class. Importantly, for both classes of genes, the change in transcription after Bdf depletion significantly differs depending on the presence of a Bdf1 peak (**Figure S3C**). We validated Bdf1 binding using ChIP-seq as an orthogonal approach. The

200 results are similar to ChEC-seq as well as to previously published ChIP-exo data (**Figure S3D and**
201 **Table S4**) (Rhee and Pugh, 2012). The number of bound promoters detected by ChIP-seq is lower due
202 to worse signal to background ratio but shows a good overlap with ChEC-seq results (**Figure S3E**).

Figure 2. A significant overlap between Bdf1/2 and Taf1 bound promoters. (**A**) Average plot of Bdf1, Bdf2 and Taf1 ChEC-seq signals versus free MNase signal at 5888 yeast promoters. Published MNase-seq data are shown as a grey area plot (Oberbeckmann et al., 2019). Mean values from replicate experiments are plotted. (**B**) Venn diagrams showing the overlap of promoters bound by indicated factors. Results of the hypergeometric test are shown. The asterisks represent p-value with the following cut-off levels: **** - 0.0001, *** - 0.001, ** - 0.01, * - 0.05, ns - >0.05. (**C**) Bar plot showing percentage of promoters in each class (TFIID-dependent or CR) bound by a given factor. (**D**) Boxplot showing the H4K12ac signal at TSS normalized to H3 signal. Genes are grouped into TFIID-dependent and CR categories and data are plotted separately depending on the presence of a significant Bdf1 peak. Signals were calculated in a -100 to +200 bp window relative to TSS. Published H3 ChIP-seq dataset was used for this analysis (Bruzzone et al., 2018). Results of the Welch's t-test are shown. (**E**) Average plot comparing Bdf2 ChEC-seq signal before (DMSO, solid line) and after (IAA, dashed line) Bdf1 depletion at 3223 promoters bound by Bdf2 and classified into TFIID-dependent and CR categories. (**F**) Scatter plot comparing $\log_2$ change in transcription and $\log_2$ change in Bdf2 ChEC-seq occupancy after depleting Bdf1/2 or Bdf1, respectively. Spearman correlation coefficient (r) is shown. Bdf2 signal was calculated in a 200 bp window centered on a dominant peak assigned to 3223 promoters. See also Figure S3 and Table S4.

203
204

We also performed a Bdf2 ChIP-seq experiment, but the quality of the data was too low to call peaks.
Instead, we calculated Bdf2 signal at promoters bound by Bdf1 and found that the signals correlate well,
supporting conclusions obtained by ChEC-seq (**Figures S3B and S3F and Table S4**). However, a
major difference between ChEC-seq and ChIP-seq was observed at the RP genes. We detected Bdf1
binding at 15% of RP promoters using ChEC-seq, which is consistent with a limited role of Bdfs at RP
genes. Conversely, almost all (94%) RP promoters have a significant Bdf1 ChIP signal, but this result
is not predictive of Bdf1 function (**Figures S1J and S3G**).
212
Next, we investigated whether Bdf1 binding correlates to histone H4 acetylation levels. We measured
the genome-wide acetylation of H4 lysine 12 (H4K12ac), which was shown to be a preferred target for
Bdf1 (Slaughter et al., 2021) (**Figure S3H**). Importantly, for both classes of genes, the normalized
H4K12ac level at promoters significantly differs depending on the presence of a Bdf1 peak (**Figure 2D**
and Table S4). To conclude, Bdf1 occupancy at promoters predicts higher H4K12ac levels and
transcription dependence on Bdfs.
219
It was earlier proposed that Bdf2 redistributes to preferred Bdf1 sites in a *bdf1* deletion strain (Durant
and Pugh, 2007). We tested this hypothesis by measuring changes in Bdf1 or Bdf2 occupancy after
depleting Bdf2 or Bdf1, respectively. Following Bdf1 depletion, the Bdf2 signal increases at TFIID
promoters at the expense of CR promoters (**Figure 2E**). The changes at individual promoters are
relatively weak but they correlate well with gene dependence on Bdfs (r = -0.52) (**Figure 2F and Table**
S4). Conversely, Bdf1 does not redistribute in response to Bdf2 depletion (**Figure 3I and Table S4**).
This result provides more evidence of redundancy between Bdf1 and Bdf2, with Bdf1 being the
dominant factor and Bdf2 serving an auxiliary role.
228
Bdf1 can be targeted to promoters independently of histone H4 acetylation
Esa1, a catalytic subunit of the NuA4 complex, is almost exclusively responsible for acetylation of
lysines 5, 8 and 12 on histone H4 in yeast (Chang and Pillus, 2009; Suka et al., 2001). We compared
the roles of Esa1 and Bdfs in transcriptional regulation using an Esa1 degron strain. We used a 1-hour
treatment with IAA, which allowed for a substantial loss of both H4K12ac and Esa1 (**Figure S4A**). 4-
thioU RNA-seq analysis revealed a global loss of transcription after Esa1 depletion, which has a good
but non-linear correlation with the Bdf experiment (r = 0.73) (**Figures 3A, 3B and S4B and Table S2**).
Importantly, Bdf depletion results in stronger transcriptional defects, suggesting that a portion of Bdf
function is H4Ac independent. Our findings show that Bdf1/2, Esa1 and TFIID cooperate in regulation
of many genes but their contributions to transcription can be significantly different at individual genes
(**Figures 3C and S4C and Table S2**).

Figure 3. Esa1 regulates similar set of genes as Bdf1/2 but is not essential for Bdf1 targeting to chromatin. (**A**) Scatter plot comparing expression level (in log scale) with $\log_2$ change in transcription per gene after depleting Esa1 for 60 min. Mean values from replicate experiments are plotted for 5313 genes with detectable signals in all RNA-seq samples collected in this work. (**B**) Scatter plot comparing $\log_2$ change in transcription after depleting Bdf1/2 or Esa1. Spearman correlation coefficient (*r*) is shown. Data for 4883 genes classified into TFIID-dependent and CR categories are plotted. (**C**) Heatmap comparing results of indicated depletion experiments on transcription of 4883 genes. (**D**) Average plots comparing H4K12ac ChIP-seq and Bdf1 ChEC-seq signals before (DMSO, solid line) and after (IAA, dashed line) Esa1 depletion at 3426 promoters bound by Bdf1 and classified into TFIID-dependent and CR categories. Mean values from replicate experiments are plotted. (**E**) Scatter plot comparing $\log_2$ change in transcription and $\log_2$ change in Bdf1 occupancy after depleting Esa1 at 3426 promoters. Spearman correlation coefficient (*r*) is shown. Bdf1 signal was calculated in a 200 bp window centered on a dominant peak. See also Figure S4 and Table S4.

Bdf1 interactions with chromatin were previously shown to be modulated by Esa1 (Durant and Pugh,

2007; Koerber et al., 2009). We used ChEC-seq to measure changes in Bdf1 occupancy following Esa1

depletion. In parallel, we measured changes in H4K12ac level using ChIP-seq. After 1 hour of IAA

treatment, the H4K12ac chromatin signal decreased dramatically, but Bdf1 was still bound at substantial

levels, especially at promoters (**Figures 3D and S4D and Table S4**). Nevertheless, the quantified
changes in Bdf1 promoter signal and transcription following Esa1 depletion are correlated, suggesting
that the effect of Esa1 on transcription is largely mediated by Bdf1 (**Figure 3E**). Consistent with our
results, Brd4 was also detected at human promoters following bromodomain inhibition or mutation which
explains why BET degradation outperforms BET inhibition in preclinical cancer models (Bauer et al.,
2021; Kanno et al., 2014; Winter et al., 2017).

Bdf1/2 play a role in both TFIID and Mediator recruitment
Human BET factors are involved in recruitment of Mediator (Bhagwat et al., 2016) and Bdf1/2 were
proposed to assist in recruitment of TFIID to yeast promoters (Matangkasombut et al., 2000). We used
ChEC-seq to measure changes in chromatin occupancy of selected subunits of TFIID (Taf1, Taf11),
Mediator (Med8, Med17) and SAGA (Spt3) after Bdf depletion. We first defined promoters bound by
each factor and restricted the downstream analysis only to these locations. Different factors mapped in
this study had characteristic distributions between the two classes of yeast promoters. Bdf1/2 are
enriched at TFIID-dependent promoters, TFIID does not exhibit preference against either class, and
SAGA and Mediator are enriched at CR promoters (**Figure S5A and Table S4**). We observed modest
(~1.7-fold) decreases in TFIID and Mediator occupancy at all bound promoters following Bdf depletion
(**Figures 4A, 4B, S5B and S5C and Table S4**). In contrast, SAGA binding was relatively insensitive to
Bdf depletion with only small decreases observed, mostly at CR promoters. Importantly, changes
measured for different subunits of TFIID or Mediator correlate (**Figures S5D and S5E and Table S4**).
Using the ChIP-seq assay, similar results were observed for Bdf-dependent Taf1 binding (**Figure S5C**
and S5F and Table S4).

We next tested whether the decreases in TFIID and Mediator recruitment were due to cooperative
interactions between these two factors (Grünberg et al., 2016; Knoll et al., 2018) or via a more direct
role of Bdfs in modulating TFIID and Mediator binding. We used ChEC-seq to measure changes in Taf1
and Med8 promoter signals after depleting Med14 and Taf13, respectively. We confirmed modest
binding cooperativity between both factors but the decreases in TFIID and Mediator binding were
significantly stronger after depleting Bdfs (**Compare Figure 4A,B with 4C,D; Table S4**). This indicates
a role of Bdfs in both TFIID and Mediator binding beyond the cooperative interactions of these two
factors. Altogether, our results illustrate that Bdfs have independent contributions to the recruitment of
TFIID and Mediator although, in the absence of Bdfs, both factors are still recruited to promoters at
substantial levels. This suggests that the strong transcription defects upon Bdf depletion are not solely
mediated by defects in TFIID and Mediator binding.

Figure 4. Bdf1/2 participate in recruitment of TFIID and Mediator to chromatin. (**A**) Average plots comparing Taf1, Med8 and Spt3 ChEC-seq signals before (DMSO, solid line) and after (IAA, dashed line) Bdf1/2 depletion at promoters bound by each factor and classified into TFIID-dependent and CR categories (2879, 890 and 2526 promoters, respectively). Mean values from replicate experiments are plotted. (**B**) Boxplot showing $\log_2$ change in promoter occupancy of indicated factors after Bdf1/2 depletion. Signals were calculated in a 200 bp window centered on a dominant peak. (**C**) Average plots comparing Taf1 and Med8 ChEC-seq signals before (DMSO, solid line) and after (IAA, dashed line) Med14 or Taf13 depletion, respectively. Same set of promoters as in Figure 4A was used for this analysis. (**D**) Boxplot showing $\log_2$ change in promoter occupancy of indicated factors after Med14 or Taf13 depletion. Signals were calculated in a 200 bp window centered on a dominant peak. List of Spt3 bound promoters was used to calculate Spt7 occupancy. See also Figure S5 and Table S4.

PIC assembly, especially at TFIID-dependent genes, is strongly dependent on Bdf1/2

Since the strong decrease in transcription of TFIID-dependent genes could not be explained solely by decreases in TFIID or Mediator binding, we probed for defects in PIC assembly, using ChIP-seq to quantitate promoter binding of the basal factor TFIIB. Depletion of either Bdfs or Taf13 caused a global

loss of TFIIB, most pronounced at TFIID-dependent promoters (**Figures 5A and S6A and Table 4**).
Comparing the defects in TFIIB binding due to Bdf depletion vs TFIID depletion, we found that Bdf
depletion caused a significantly larger loss of TFIIB at TFIID-dependent promoters than does TFIID
depletion (**Figure 5B and Table S4**). Conversely, at CR genes, we found that TFIID depletion caused
a greater loss in TFIIB signal compared with Bdf depletion. These results are consistent with the above
RNA-seq experiments. However, comparison of transcriptional and TFIIB binding defects caused by
Bdf depletion shows that Bdfs have a broader role in transcription than solely regulating PIC assembly.
This larger role of Bdfs is especially apparent for the most Bdf-dependent genes where transcription
defects significantly exceed the defect in TFIIB binding (**Figure 5C**). Combined, our results suggest an
additional role of Bdfs in transcription that extend beyond organizing a platform for recruitment of PIC
components.

Figure. 5. Depletion of Bdf1/2 compromises TFIIB recruitment to promoters. (**A**) Average plots comparing TFIIB ChIP-seq signals before (DMSO, solid line) and after (IAA, dashed line) Bdf1/2 or Taf13 degradation at 4900 promoters classified into TFIID-dependent and CR categories. Mean values from replicate experiments are plotted. (**B**) Boxplot showing $\log_2$ change in promoter occupancy of TFIIB after Taf13 or Bdf1/2 degradation. Signals were calculated in a -200 to 100 bp window relative to TSS. Results of the Welch's t-test are shown. (**C**) Scatter plot comparing $\log_2$ change in transcription and $\log_2$ change in TFIIB occupancy after Bdf1/2 degradation at 4883 promoters/genes analyzed in RNA-seq experiments. Spearman correlation coefficient (*r*) is shown. See also Figure S6 and Table S4.

Bdf1/2 promote productive elongation at a subset of genes

At many metazoan promoters, Pol II pauses after transcribing ~20-100 nucleotides and release of
paused Pol II, mediated in part by the C-terminal repeat domain (CTD) Ser2 phosphorylation, is a critical
step in gene regulation (Core and Adelman, 2019). Interestingly, Brd4 was shown to be involved in Ser2
phosphorylation (Muhar et al., 2018; Winter et al., 2017). While metazoan-like Pol II pausing does not

occur at yeast promoters, Pol II stalling and shifts in Pol II distribution under specific conditions have
been observed in both budding and fission yeast (Badjatia et al., 2021; Shetty et al., 2017).

We tested whether Bdfs, like the mammalian BET factors, play a role in Pol II elongation. Elongating
Pol II was quantitated using ChIP-seq for Rpb1 before and after depleting Bdfs and, as a comparison,
Taf1. We first calculated the change in Pol II occupancy along the whole transcribed region. We found
that the change in transcription following Bdf depletion is not proportional to the loss of Pol II signal,
even though the two results show a good correlation (**Figure 6A and Table S5**). As we observed for
TFIIB, the decrease in Pol2 signal at the most Bdf-dependent genes is less severe than the loss of
transcription. Importantly, we did not detect such difference after depleting Taf1 (**Figure S7A and Table**
S5). Next, we divided genes into quintiles, based on Bdf dependence, and plotted the change in Pol II
occupancy up to 1 kb downstream from the TSS (**Figure 6**). With the exception of the first quintile (the
least Bdf-dependent genes), Bdf depletion caused a similar loss of Pol II at TSSs of all genes.
Interestingly, the Pol II loss increased further downstream of TSS at the most Bdf-dependent genes
until stabilizing at ~400 bp downstream of TSS. Conversely, at the least Bdf-dependent genes, the Pol
II signal partially recovered in the gene body relative to the loss at TSS. We calculated the Pol II
travelling ratio (TR) defined as the ratio of Rpb1 signal at 5' versus the 3' end of transcribed region (Rahl
et al., 2010). We observed a decrease of TR at the least Bdf-dependent genes and an increase of TR
at the most dependent genes following Bdf depletion (**Figure 6C and Table S5**). Analysis of individual
genes revealed shifts in Pol II distribution towards 3' or 5' ends of genes, respectively (**Figure S7B**).
We did not detect similar changes in Pol II distribution after depleting Taf1 (**Figures S7C and S7D and**
Table S5). Importantly, the change in TR after depleting Bdfs correlates with the change in transcription
at individual genes (**Figure 6D**). Combined with our earlier findings, this suggests that the most Bdf-
dependent genes experience transcription defects both at the initiation and elongation stages upon Bdf
depletion. This explains the disproportionally large decrease in the amount of nascent RNA. On the
other hand, more efficient elongation at the least Bdf-dependent genes seems to partially nullify the
initiation defects leading to little total change in transcription upon Bdf depletion (**Figure 1C**).

To further explore elongation defects, we performed ChIP-seq with antibodies specific to
phosphorylated CTD residues Ser5P and Ser2P. Sequential deposition of these marks coordinates the
assembly of multiple factors during transcription and they are involved in successful promoter escape
and elongation, respectively (Harlen and Churchman, 2017). We measured total Ser5P and Ser2P
signals along transcribed regions, normalized them by corresponding Rpb1 signals, and calculated the
change in CTD phosphorylation status following Bdf depletion (**Figure 6E and Table S5**). In agreement
with the above findings that the most Bdf-dependent genes have a defect in elongation, Pol II becomes

hypophosphorylated on Ser2 and, to a lesser extent, on Ser5 after depleting Bdfs. Conversely, at the

least Bdf-dependent genes, Pol II becomes weakly hyperphosphorylated, in agreement with the

suggestion that at this gene set elongation becomes more efficient. Importantly, we did not observe

similar changes in Pol II phosphorylation after Taf1 depletion, which validates that Bdfs have a TFIID-

independent role in regulating transcriptional elongation (**Figure S7E and Table S5**).

Figure. 6. Bdf1/2 are involved in transcriptional elongation. (**A**) Scatter plot comparing $\log_2$ change in transcription and occupancy of the largest Pol II subunit Rpb1 after Bdf1/2 degradation. Rpb1 occupancy was calculated along the whole transcribed region for 4615 genes longer than 300 bp, and with annotated TSS and PAS locations (Park et al., 2014). Spearman correlation coefficient (*r*) is shown. Mean values from replicate experiments are plotted. (**B**) Average plot showing $\log_2$ change in Rpb1 occupancy from 200 bp upstream to 1000 bp downstream of TSS after depleting Bdf1/2. Data in this and following panels are divided into five groups based on gene dependence on Bdf1/2 as measured by 4-thioU RNA-seq for 3438 genes longer than 1 kb and with annotated TSS and PAS locations (Park et al., 2014). Published MNase-seq dataset is shown as a grey area plot (Oberbeckmann et al., 2019). (**C**) Boxplot comparing Pol II traveling ratio (TR) in DMSO or IAA treated samples in the Bdf1/2 degron experiment. TR represents a ratio of Rpb1 occupancy calculated in 100 bp windows at the beginning and end of a transcribed region. The same set of 4615 genes as in Figure 6A was used. (**D**) Scatter plot comparing $\log_2$ change in transcription and $\log_2$ change in Pol II traveling ratio after depleting Bdf1/2. Spearman correlation coefficient (*r*) is shown. (**E**) Boxplot comparing $\log_2$ change in Rpb1 CTD phosphorylation status at Ser2 and Ser5 residues after depleting Bdf1/2. Data were calculated along the whole transcribed region for the same set of 4615 genes as in Figure 6A and signals were normalized to the total Rpb1 signal. (**F**) Boxplot comparing $\log_2$ change in Bur1, Ctk1 and Spt5 occupancy after depleting Bdf1/2. Data for DMSO and IAA treated samples were calculated along the whole transcribed region for the same set of 4615 genes as in Figure 6A and signals were normalized to the total Rpb1 signal. See also Figure S7 and Table S5.

Bur1 and Ctk1, homologues of metazoan Cdk9 and Cdk12/13, are responsible for CTD Ser2P marks
in yeast. We used ChIP-seq to measure Bur1 and Ctk1 levels along the transcribed regions, normalized
them by corresponding Rpb1 signals, and calculated the change in their relative occupancy following
Bdf depletion (**Figure 6F and Table S5**). We observed ~2-fold increases in occupancy of both kinases
relative to Rpb1 at the least Bdf-dependent genes, a result that agrees with observed CTD
hyperphosphorylation. Interestingly, at the rest of genes, the relative occupancy of both kinases further
increased, correlating with gene dependence on Bdfs. As a comparison, we also investigated defects
in the recruitment of Spt5, an essential elongation factor. We observed a uniform, small (~1.4-fold)
increase in Spt5 occupancy, normalized to Rpb1, at all tested genes (**Figure 6F and Table S5**). These
results show that changes in the recruitment of Spt5 and the CTD kinases cannot account for the
function of the Bdfs in transcription elongation.
356

Discussion

After development of specific inhibitors targeting BET bromodomains (BDs), BET factors, especially
Brd4, emerged as key regulators of transcription and as promising targets in the therapy of cancer and
immunoinflammatory diseases (Fujisawa and Filippakopoulos, 2017; Wang et al., 2021). In this work,
we investigated the roles of yeast BET family members Bdf1 and Bdf2. We explored transcription
dependence on Bdf1/2, their genome-wide binding patterns and, their interplay with key components of
the transcriptional machinery. We found that Bdf functions go beyond solely regulating TFIID, as
expression from most genes is more sensitive to Bdf vs TFIID depletion. Our results establish the Bdfs
as critical for normal expression of most yeast genes, with functions both during transcription initiation
and elongation. Our work reveals that the roles of yeast BET factors are surprisingly similar to their
human equivalents and suggests conserved mechanisms of BET function across eukaryotes.
368

Bdf1 was proposed to act as part of yeast TFIID, substituting for the metazoan Taf1 BDs
(Matangkasombut et al., 2000). This function might be especially important in yeast, as H4 acetylation
is enriched at most promoters and yeast promoters lack known DNA sequence motifs that are bound
by TFIID. However, we found that that the organization and amino acid sequence of Bdf1 is more similar
to human BET factors than to the human Taf1 (hTaf1) bromodomains. Our results showed that Bdfs
regulate a similar set of genes as TFIID, but that there are important differences. Transcription from
most genes defined earlier as coactivator-redundant (CR) is largely insensitive to Bdf depletion, while
a large subset of TFIID-dependent genes is significantly more affected by Bdf vs TFIID depletion.
377

Genome-wide mapping revealed that Bdf1/2 are found together with TFIID at many promoters and that
they contribute to TFIID recruitment. However, a substantial fraction of TFIID is still bound to promoters
in the absence of Bdfs, in agreement with the sub-stochiometric association between Bdf1 and TFIID
(Sanders et al., 2002). Interestingly, human BET proteins were shown to associate with TFIID in a
proteomic screen and both Brd4 and hTaf1 were proposed to have synergistic effects on gene
expression and cell growth (Lambert et al., 2019; Sdelci et al., 2016). Finally, hTaf1 BDs have the same
substrate specificity as BET BDs and both Brd4 and TFIID are bound to the majority of active promoters
in human cells (Bhagwat et al., 2016; Lauberth et al., 2013; Slaughter et al., 2021). Based on our
findings and the presented evidence from human cells, we propose that cooperation between BET
factors and TFIID is a conserved feature of eukaryotic gene regulation. The gain of bromodomains by
metazoan TFIID may result from the increased complexity of gene expression programs and allow for
BET-independent TFIID interactions with acetylated histone H4.

Figure. 7. Yeast BET proteins regulate transcription initiation and elongation. Yeast BET proteins bind
to regions of the genome highly acetylated at histone H4. Together with sequence-specific transcription
factors (TFs) they provide a nucleation center for dynamic recruitment of TFIID and Mediator to the
promoter-proximal nuclear territory. In turn, TFIID and Mediator create a platform which facilitates PIC
assembly and stimulates transcription initiation. Yeast BET proteins also affect processive elongation serving
as negative regulatory factors at CR genes and positive regulatory factors at TFIID-dependent genes.

Brd4 interacts with Mediator and participates in Mediator recruitment to enhancers and promoters
(Bhagwat et al., 2016; Wu et al., 2003). Our experiments revealed that Bdfs are important for normal
levels of Mediator occupancy. A direct role of Bdfs in Mediator targeting to chromatin was surprising
and we considered the possibility that other factors are involved in that process. Mediator was proposed
to be recruited cooperatively with TFIID (Grünberg et al., 2016; Johnson et al., 2002; Knoll et al., 2018).

We confirmed this modest cooperativity, but we found that the contribution of Bdfs to Mediator
recruitment surpasses that of TFIID. There is strong evidence that TFIID and Mediator act as molecular
scaffolds to organize the recruitment of other components of transcriptional machinery (Allen and
Taatjes, 2015; Patel et al., 2020). Human Mediator and Brd4 were also found to participate in the
formation of dynamic nuclear condensates at the sites of active transcription (Han et al., 2020; Sabari
et al., 2018). Finally, a recent study suggested cooperation of yeast TFIID and Mediator in restricting
the diffusion of PIC components to shared subnuclear territories in order to facilitate gene transcription
(Nguyen et al., 2020). We propose that chromatin tethered yeast BET factors serve as a nucleation
center for dynamic recruitment of Mediator and TFIID, that in turn create a platform for efficient PIC
assembly. This role of Bdf1/2 seems widespread and likely complementary to the action of sequence-
specific transcription factors (TFs) (**Figure 7**).

At the least Bdf-dependent genes, we observed a modest decrease in TFIID, Mediator and TFIIB
recruitment after Bdf depletion, which did not translate into appreciable defects in mRNA synthesis.
Conversely, at the most Bdf-dependent genes, the loss of transcription exceeded the loss of Mediator,
TFIID, TFIIB and Pol II recruitment. The unexpected role of the yeast BET factors in transcription
elongation provided an explanation for these seemingly conflicting results (**Figure 7**). At CR genes and,
at a small subset of TFIID-dependent genes, Bdf depletion results in a 3' shift in Pol II distribution and
increased CTD Ser2 phosphorylation. This suggests more efficient elongation of these genes in the
absence of Bdfs. In contrast, Bdf depletion at many TFIID-dependent genes causes Pol II accumulation
at 5' gene ends and a loss of CTD Ser2 phosphorylation. This change in Pol II profile is highly similar
to the defects in pause release caused by BET depletion in human cells (Muhar et al., 2018; Winter et
al., 2017). At these yeast genes, the combined initiation and elongation defects resulting from Bdf
depletion lead to a large decrease in mRNA synthesis.

CTD Ser2 residues in yeast are phosphorylated by kinases Bur1 and Ctk1, homologues of metazoan
Cdk9 and Cdk12/13, respectively. Ctk1 is the dominant CTD kinase in yeast while in metazoans
Cdk12/13 seem to be less important, although the interplay between Cdk9 and Cdk12/13 is poorly
understood (Harlen and Churchman, 2017). At Bdf-independent genes, depletion of Bdfs results in a
modest increase of Bur1 and Ctk1 occupancy normalized to Rpb1 level, and this agrees with a gain of
CTD Ser2P marks at these genes. Surprisingly, at Bdf-dependent genes, occupancy of both kinases
relative to Rpb1 level increased significantly after Bdf depletion, and in apparent conflict with observed
defects in CTD Ser2 phosphorylation. Interestingly, depletion of Brd4 or all BET factors in human cells
was shown to result in a small increase in Cdk9 and Cyclin T1 chromatin occupancy, even though it
causes a decrease in CTD Ser2P marks and a defect in pause release (Muhar et al., 2018; Winter et

al., 2017). It is not known if Brd4 affects Cdk12/13 recruitment. There are alternative models explaining

the role of Brd4 in transcription elongation. Brd4 was shown to release Hexim1-mediated inhibition of

Cdk9 *in vitro* thus acting as a positive regulator of P-TEFb (Itzen et al., 2014). It was also proposed that

Brd4 can act as a histone chaperone and facilitate Pol II elongation independently of its interaction with

P-TEFb (Kanno et al., 2014). Finally, it is possible that CTD hypophosphorylation at BET-dependent

genes may result from defects in recruitment and/or reduction in function of other elongation factors.

Since the role(s) of the BET factors in elongation are unresolved in both systems, it will be of great

interest to determine whether the yeast and metazoan factors use similar mechanisms to regulate

elongation.

Redundancy between three (Brd2, Brd3, Brd4) of the four mammalian BET factors was proposed based

on their ubiquitous expression in most tissues, similar binding patterns and, because they are all

essential for embryonic development (Gyuris et al., 2009; Khoueiry et al., 2019; Uhlén et al., 2015). The

extent of this redundancy is poorly understood because only a few context-specific examples of shared

functions of BET factors have been reported (Gilan et al., 2020; Rahman et al., 2011; Stonestrom et al.,

2015). In most studies the roles of BET factors in transcriptional regulation were attributed to Brd4 alone

(Muhar et al., 2018; Zheng et al., 2021; Zuber et al., 2011). Our data indicate that yeast BET proteins

are functionally redundant under normal growth conditions, with Bdf1 being the dominant factor. The

changes in transcription following depletion of Bdf1 or Bdf1/2 are highly correlated, while depletion of

Bdf2 does not have significant consequences for the cell. In addition, Bdf1 and Bdf2 have nearly

identical genome-wide binding patterns and Bdf2 redistributes from CR to TFIID-dependent genes upon

Bdf1 depletion. Interestingly, it was reported that Bdf2 overexpression can suppress temperature- and

salt-sensitive phenotypes in *bdf1* deletion strain, but at the same time Bdf1 was found to be a negative

regulator of Bdf2 expression (Fu et al., 2013; Matangkasombut et al., 2000). This latter finding is also

supported by our RNA-seq results where we observed ~2-fold upregulation of *BDF2* transcription

following Bdf1 depletion. Importantly, a similar interplay between human BET factors in regulating

expression of other family members was recently proposed (Lambert et al., 2019). It remains to be

investigated if and to what extent Brd2 and/or Brd3 can replace Brd4 in supporting transcription in

human cells.

Recent years have brought remarkable discoveries highlighting the essential nature of mammalian BET

factors. Still, many aspects of their biology remain poorly understood, including details of the relationship

with other components of the transcriptional machinery at enhancers and promoters and the extent of

redundancies. We uncovered striking similarities between yeast and mammalian BET family members.

Many of the most BET sensitive genes in both systems lack a TATA element in their promoters and

control highly similar cellular processes. BET proteins are also universally involved in the recruitment

of Mediator to chromatin and modulation of transcriptional elongation. Considering the relative simplicity

compared to metazoans, the yeast model system offers unique opportunities for a detailed investigation

of different aspects of BET biology and this work provides a foundation for these future studies.

Materials and Methods

Yeast cell growth

All *S. cerevisiae* and *S. pombe* strains used in this study are listed in **Table S1**. *S. cerevisiae* strains

were grown in YPD medium (1% yeast extract, 2% peptone, 2% glucose, 20 µg/ml adenine sulfate) at

30°C with shaking. *S. pombe* strains were grown in YE medium (0.5% yeast extract, 3% glucose) at

30°C with shaking. In experiments involving degron depletion of target proteins, *S. cerevisiae* strains

were treated with 500 µM indole-3-acetic acid (IAA) dissolved in DMSO or with DMSO alone for 30 min

(or 60 min for Esa1-degron strains) to induce protein degradation followed by protocol-specific steps.

In RNA-seq experiments, *S. cerevisiae* and *S. pombe* strains were grown to A600 ~ 1.0. In ChEC-seq

experiments *S. cerevisiae* strains were grown to A600 ~ 0.6. In ChIP-seq experiments *S. cerevisiae*

and *S. pombe* strains were grown to A600 ~ 0.8. The number of biological replicates collected in each

experiment is listed in **Table S6**.

Strain construction

S. cerevisiae strains (**Table S1**) were constructed using standard methods. Proteins were

chromosomally tagged by yeast transformation and homologous recombination of PCR-amplified DNA.

Plasmid pFA6a-3V5-IAA7-KanMX6 (Chan et al., 2018), or a derivative with the *KanMX* marker replaced

by *URA3* (pSH1855), were used as a template for generating the IAA7 degron tags. This C-terminal tag

contains three copies of the V5 epitope tag followed by the IAA7 degron. For ChEC-seq experiments,

proteins were tagged with 3xFLAG-MNase::TRP1 using pGZ110 (Zentner et al., 2015). Plasmids p2L-

3Flag-NAT and pFA6a-13Myc-Hyg (a gift from Toshio Tsukiyama, Fred Hutch) were used as templates

to generate PCR fragments for tagging proteins with 3xFlag and 13xMyc epitope tags. A strain

expressing free MNase under control of the native *BDF2* promoter was constructed the following way.

First, the *MED8* promoter in pSG79 (Grünberg et al., 2016) was exchanged with the *BDF2* promoter

(containing 500 bp upstream DNA from the *BDF2* start codon). A XhoI/SacI fragment containing the

pBDF2-MNase fusion DNA was inserted to the yeast integrating vector pRS303. The resulting pRD15

plasmid was linearized with BstEII and integrated into strain BY4705. *BDF2* deletion strain was

constructed by replacing the *BDF2* gene with the *HPH* marker amplified from plasmid pAG32 (Goldstein

and McCusker, 1999). *ESA1*-containing vectors were constructed by inserting a XhoI/SacI fragment

containing the *ESA1* gene with its native promoter and terminator into vectors pRS316 (pRD5) and

pRS315 (pRD21). A plasmid shuffle strain was constructed containing WT *ESA1* on plasmid pRD5 and

a chromosomal deletion of *ESA1* in strain SHY1036. pRD21 was modified by replacing nucleotides 301

– 354 in the *ESA1* ORF with the 3xV5-IAA7 fragment to create the Esa1-degron. Plasmid shuffling was

used to replace pRD5 with pRD21 by counter selection on 5-FOA plates.

Western blot analysis

1 ml cell culture was collected and pelleted from strains after treatment with IAA or DMSO, washed with

500 µl water, then resuspended in 100 µl yeast whole cell extract buffer (60 mM Tris, 6.8, 10% glycerol,

2% SDS, 5% 2-mercaptoethanol, 0.0025% bromophenol blue). After heating for 5 min at 95°C, samples

were centrifuged for 5 min at 21K x g and analyzed by SDS-PAGE and Western blot where protein

signals were visualized by using the Odyssey CLx scanner and quantified using Odyssey Image Studio

software (Li-Cor) by generating a standard curve using a titration from WT extract. Each protein

analyzed was normalized to the amount of the TFIIF subunit Tfg2.

Spot assay for yeast growth

S. cerevisiae strains were streaked from glycerol stocks on YPD plates and incubated for 3 days at

30°C. Single colonies from freshly grown plates were used to start overnight cultures for the spot assay.

After reaching saturation, cultures were diluted to A600 = 1.0 and 10x serial dilutions were prepared.

10 µl of appropriate dilution was spotted on YPD plates and the plates were incubated for 2-3 days at

30°C to compare growth rates.

Isolation of newly synthesized RNA

All steps were done essentially as described with minor modifications (Donczew et al., 2020). Two

(Bdf1, Bdf2, *Δbdf2*, Bdf1*Δbdf2*, H2A.Z) or three (Bdf1/2, Toa1, Ssl2) replicate samples were collected

for each experiment. 10 ml *S. cerevisiae* or 20 ml *S. pombe* (strain SHY1058) cultures were labeled

with 5 mM 4-thiouracil (4-thioU) (Sigma-Aldrich #440736) for 4 min, the cells were pelleted at 3000 x g

for 3 min, flash-frozen in liquid N_2, and then stored at −80°C for further use. *S. cerevisiae* and *S. pombe*

cells were mixed in an 8:1 ratio and total RNA was extracted using reagents from RiboPure yeast kit

(Thermo Fisher Scientific #AM1926) using the following volumes: 480 µl lysis buffer, 48 µl 10% SDS,

480 µl phenol/CHCl$_3$/isoamyl alcohol (25:24:1) per *S. cerevisiae* pellet + 50 µl *S. pombe* cell solution

(from a single *S. pombe* pellet resuspended in 850 µl lysis buffer). Cells were lysed using 1.25 ml

zirconia/silica beads (RPI #9834) in a Mini Beadbeater-96 (BioSpec Products) for 5 min. Lysates were

spun for 5 min at 16k x g, then the following volumes were combined in a 5 ml tube: 400 µl supernatant,

1400 µl binding buffer, 940 µl 100% ethanol. Samples were processed through Ambion filter cartridges

until all sample was loaded, then washed with 700 µl Wash Solution 1, and twice with 500 µl Wash

Solution 2/3. After a final spin to remove residual ethanol, RNA was eluted with 25 µl 95°C preheated

Elution Solution. The elution step was repeated, and eluates combined. RNA was then treated with

DNaseI using 5 µl DNaseI buffer and 4 µl DNaseI for 30 min at 37°C, then treated with Inactivation

Reagent for 5 min at RT. Total RNA samples were stored at -20°C for up to 2 months or at -80°C for

longer periods.

RNA was biotinylated using 40 µl (~40 µg) total RNA and 4 µg MTSEA biotin-XX (Biotium #90066-1) in

the following reaction: 40 µl total 4-thioU-labeled RNA, 20 mM HEPES, 1 mM EDTA, 4 µg MTSEA

biotin-XX (80 µl 50 µg/ml diluted stock) in a 400 µl final volume. Biotinylation reactions occurred for 30

min at RT with rotation and under foil. Unreacted MTS-biotin was removed by phenol/CHCl$_3$/isoamyl

alcohol (25:24:1) extraction. RNA was precipitated with isopropanol and resuspended in 100 µl

nuclease-free H2O. Biotinylated RNA was purified using 80 µl MyOne Streptavidin C1 Dynabeads

(Thermo Fisher Scientific #65002) + 100 µl biotinylated RNA for 15 min at RT with rotation and under

foil. Prior to use, MyOne Streptavidin beads were washed in a single batch with 3 × 3 ml H$_2$O, 3 × 3 ml

High Salt Wash Buffer (100 mM Tris, 7.4, 10 mM EDTA, 1 M NaCl, 0.05% Tween-20), blocked in 4 ml

High Salt Wash Buffer containing 40 ng/µl glycogen (Millipore Sigma #10901393001) for 1 hr at RT,

then resuspended to the original volume in High Salt Wash Buffer. After incubation with biotinylated

RNA, the beads were washed 3 × 0.8 ml High Salt Wash Buffer, then eluted into 25 µl streptavidin

elution buffer (100 mM DTT, 20 mM HEPES, 7.4, 1 mM EDTA, 100 mM NaCl, 0.05% Tween-20) at RT

with shaking, then the elution step was repeated, and eluates were combined for a total of 50 µl.

Samples were stored at -20°C until further processing.

10% input (not biotinylated) RNA (4 µl) was diluted into 50 µl streptavidin elution buffer and processed

the same as the biotinylated RNA samples to determine the extent of recovery. 50 µl of each input and

biotinylated RNA was adjusted to 100 µl with nuclease-free water and purified on RNeasy columns

(Qiagen #74104) using the modified protocol. To each 100 µl sample, 350 µl RLT lysis buffer (supplied

by the Qiagen kit and supplemented with 10 µl 1% βME per 1 ml RLT) and 250 µl 100% ethanol was

added, mixed well, and applied to columns. Columns were washed with 500 µl RPE wash buffer

(supplied by the Qiagen kit and supplemented with 35 µl 1% βME per 500 µl RPE), followed by a final

5 min spin at 21K xg. RNAs were eluted into 14 µl nuclease-free water and the RNA concentration was

measured using Qubit HS RNA assay (Thermo Fisher Scientific #Q32852). Samples were stored at -

20°C. One sample per batch prepared in a single day was tested for enrichment of labeled RNA by RT-

qPCR, probing both unlabeled and labeled RNA from at least three transcribed genes as previously

described (Donczew et al., 2020). The purified 4-thioU labeled RNA contained 2–11% contamination of

unlabeled RNA.

Preparation of RNA-seq libraries for NGS

Newly synthesized RNA isolated via 4-thioU labeling and purification was prepared for sequencing

using the Ovation Universal RNA-seq Library Preparation Kit with *S. cerevisiae* AnyDeplete reagent

(Tecan #0364-A01) according to the manufacturer's instructions and 50 ng input RNA. Libraries were

sequenced on the Illumina HiSeq2500 platform using 25 bp paired-end reads at the Fred Hutchinson

Genomics Shared Resources facility.

ChEC-seq experiments

ChEC-seq was performed as previously described (Donczew et al., 2021, 2020). In experiments

involving Bdf1, Bdf2 or Bdf1/2 depletion six replicate samples were collected. In Esa1 depletion

experiment two replicate samples were collected, as well as in experiments mapping SAGA subunits

(Spt3, Spt7) following Med14 or Taf13 depletion. In all other experiments three replicate samples were

collected. *S. cerevisiae* 50 ml cultures were pelleted at 2000 x g for 3 min. Cells were resuspended in

1 ml of Buffer A (15 mM Tris, 7.5, 80 mM KCl, 0.1 mM EGTA, 0.2 mM spermine (Millipore Sigma

#S3256), 0.3 mM spermidine (Millipore Sigma #85558), protease inhibitors (Millipore Sigma

#04693159001)), transferred to a 1.5 ml tube and pelleted at 1500 x g for 30 sec. Cell were washed

twice with 1 ml of Buffer A and finally resuspended in 570 μl of Buffer A. 30 μl 2% digitonin (Millipore

Sigma #300410) was added to a final concentration of 0.1% and cells were permeabilized for 5 min at

30°C with shaking (900 rpm). 0.2 mM CaCl$_2$ was added to the samples followed by incubation for

another 5 min at 30°C. 100 μl cell suspension was mixed with 100 μl Stop Solution (400 mM NaCl, 20

mM EDTA, 4 mM EGTA). Stop Solution was supplemented with 5 μg MNase digested *D. melanogaster*

chromatin. Samples were incubated with 0.4 mg/ml Proteinase K (Thermo Fisher Scientific #AM2548)

for 30 min at 55°C and the DNA was purified by phenol/CHCl$_3$/isoamyl alcohol (25:24:1) extraction and

ethanol precipitation. Pellets were resuspended in 30 μl 0.3 mg/ml RNase A (Thermo Fisher Scientific

#EN0531) (10 mM Tris, 7.5, 1 mM EDTA, 0.3 mg/ml RNase A) and incubated for 15 min at 37°C. 60 μl

of Mag-Bind reagent (Omega Biotek #M1378-01) was added and the samples were incubated for 10

min at RT. Supernatants were transferred to a new tube and the volume was adjusted to 200 μl (10 mM

Tris, 8.0, 100 mM NaCl). DNA was purified again by phenol/CHCl$_3$/isoamyl alcohol (25:24:1) extraction

and ethanol precipitation, and resuspended in 25 μl 10 mM Tris, 8.0.

ChIP-seq experiments

ChIP-seq experiments were performed similarly as described (Donczew et al., 2020). Two replicate samples were collected for all experiments except for Bur1 ChIP-seq where three replicates were collected. 100 ml *S. cerevisiae* or *S. pombe* cultures were crosslinked with 1% formaldehyde (Sigma-Aldrich #252549) for 20 min in the above growth conditions, followed by another 5 min treatment with 130 mM glycine. Cells were pelleted at 3000 x g for 5 min, washed with cold TBS buffer, pelleted at 2000 x g for 3 min, flash-frozen in liquid N_2, and then stored at −80°C for further use. Cell pellets were resuspended in 300 µl Breaking Buffer (100 mM Tris, 8.0, 20% glycerol, protease inhibitors (Millipore Sigma #04693159001)). Cells were lysed using 0.4 ml zirconia/silica beads (RPI #9834) in a Mini Beadbeater-96 (BioSpec Products) for 5 min. Lysates were spun at 21K x g for 2 min. Pellets were resuspended in 1 ml FA buffer (50 mM HEPES, 7.5, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate, protease inhibitors (Millipore Sigma #04693159001)) and transferred to 15 ml polystyrene tubes. In experiments with antibodies specific against phosphorylated Rpb1 CTD, Breaking Buffer and FA buffer were supplemented with phosphatase inhibitors (Thermo Fisher Scientific #A32957). Samples were sonicated in a cold Bioruptor sonicator bath (Diagenode #UCD-200) at a maximum output, cycling 30 sec on, 30 sec off, for a total of 45 min. Samples were spun twice in fresh tubes at 21K x g for 15 min. Prepared chromatin was flash-frozen in liquid N_2, and then stored at −80°C for further use.

20 µl of the chromatin sample was used to estimate DNA concentration. First, 20 µl Stop buffer (20 mM Tris, 8.0, 100 mM NaCl, 20 mM EDTA, 1% SDS) was added to samples followed by incubation at 70°C for 16-20 hrs. Samples were digested with 0.5 mg/ml RNase A (Thermo Fisher Scientific #EN0531) for 30 min at 55°C and 1 mg/ml Proteinase K for 90 min at 55°C. Sample volume was brought to 200 µl and DNA was purified by two phenol/CHCl$_3$/isoamyl alcohol (25:24:1) extractions and ethanol precipitation. DNA was resuspended in 20 µl 10 mM Tris, 8.0 and the concentration was measured using Qubit HS DNA assay (Thermo Fisher Scientific #Q32851).

20 µl Protein G Dynabeads (Thermo Fisher Scientific #10003D) was used for a single immunoprecipitation. Beads were first washed three times with 500 µl PBST buffer (PBS buffer supplemented with 0.1% Tween 20) for 3 min with gentle rotation. Beads were resuspended in a final volume of 20 µl containing PBST buffer and 5 – 8 µl of appropriate antibody (Key Resources Table). The following antibody volumes were used: H4K12ac, FLAG-Tag, Myc-Tag – 5 µl; Rpb1 CTD (total or phosphorylated) – 8.5 µl. The bead suspension was incubated for 60 min with shaking (1400 rpm) at RT, washed with 500 µl PBST buffer and 500 µl FA buffer. Beads were finally resuspended in 25 µl FA

buffer. 1.5 µg *S. cerevisiae* chromatin and 30 ng *S. pombe* chromatin (strain SHY1110) were combined

and samples were brought to a final volume of 500 µl. 25 µl of each sample was mixed with 25 µl Stop

buffer and set aside (input sample). 25 µl of beads was added to remaining 475 µl of samples followed

by incubation for 16-20 hrs at 4°C.

The beads were washed for 3 min with gentle rotation with the following: 3 times with 500 µl FA buffer,

2 times with FA-HS buffer (50 mM HEPES, 7.5, 500 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1%

sodium deoxycholate), once with 500 µl RIPA buffer (10 mM Tris, 8.0, 0.25 M LiCl, 0.5% NP-40, 1 mM

EDTA, 0.5% sodium deoxycholate). DNA was eluted from beads with 25 µl Stop buffer at 75°C for 10

min. Elution was repeated, eluates were combined and incubated at 70°C for 16-20 hrs together with

input samples collected earlier. Samples were digested with 0.5 mg/ml RNase A (Thermo Fisher

Scientific #EN0531) for 30 min at 55°C and 1 mg/ml Proteinase K for 2 hrs at 55°C. Sample volume

was brought to 200 µl and DNA was purified by two phenol/CHCl$_3$/isoamyl alcohol (25:24:1) extractions

and ethanol precipitation. DNA was resuspended in 15 µl 10 mM Tris, 8.0 and the concentration was

measured using Qubit HS DNA assay (Thermo Fisher Scientific #Q32851).

Preparation of NGS libraries for ChEC-seq and ChIP-seq samples

NGS libraries for ChEC-seq and ChIP-seq experiments were prepared similarly as described (Donczew

et al., 2020; Warfield et al., 2017). 12 µl of ChEC samples and 5 µl of ChIP samples was used as input

for library preparation. Samples were end-repaired, phosphorylated and adenylated in 50 µl reaction

volume using the following final concentrations: 1X T4 DNA ligase buffer (NEB #B0202S), 0.5 mM each

dNTP (Roche #KK1017), 0.25 mM ATP (NEB #P0756S), 2.5% PEG 4000, 2.5 U T4 PNK (NEB

#M0201S), 0.05 U T4 DNA polymerase (Invitrogen #18005025), and 0.05 U Taq DNA polymerase

(Thermo Fisher Scientific #EP0401). Reactions were incubated at 12°C 15 min, 37°C 15 min, 72°C 20

min, then put on ice and immediately used in adaptor ligation reactions. Adaptor ligation was performed

in a 115 µl volume containing 6.5 nM adaptor, 1X Rapid DNA ligase buffer (Enzymatics #B101L) and

3000 U DNA ligase (Enzymatics #L6030-HC-L) and reactions were incubated at 20 deg for 15 min.

Following ligation, a two-step cleanup was performed for ChEC-seq samples using 0.25x vol Mag-Bind

reagent (Omega Biotek # M1378-01) in the first step and 1.1x vol in the second step. In case of ChIP-

seq samples a single cleanup was performed using 0.4x vol Mag-Bind reagent. In both cases DNA was

eluted with 20 µl 10 mM Tris, 8.0. Library Enrichment was performed in a 30 µl reaction volume

containing 20 µl DNA from the previous step and the following final concentrations: 1X KAPA buffer

(Roche #KK2502), 0.3 mM each dNTP (Roche #KK1017), 2.0 µM each P5 and P7 PCR primer, and 1

U KAPA HS HIFI polymerase (#KK2502). DNA was amplified with the following program: 98°C 45 s,

[98°C 15 s, ramp to 60°C @ 3°C /s, 60°C 10 s, ramp to 98°C @ 3°C /s] 16-18x, 72°C 1 min. 18 cycles

were used for library amplification for ChEC-seq samples and 16 cycles for ChIP-samples. A post-PCR

cleanup was performed using 1.4x vol Mag-Bind reagent and DNA was eluted into 30 μl 10 mM Tris,

8.0. Libraries were sequenced on the Illumina HiSeq2500 platform using 25 bp paired-end reads at the

Fred Hutchinson Cancer Research Center Genomics Shared Resources facility.

Analysis of NGS data

Data analysis was performed similarly as described (Donczew et al., 2020). The majority of the data

analysis tasks except sequence alignment, read counting and peak calling (described below) were

performed through interactive work in the Jupyter Notebook (https://jupyter.org) using Python

programming language (https://www.python.org) and short Bash scripts. All figures were generated

using Matplotlib and Seaborn libraries for Python; (https://matplotlib.org; https://seaborn.pydata.org). All

code snippets and whole notebooks are available upon request.

Paired-end sequencing reads were aligned to *S. cerevisiae* reference genome (sacCer3), *S. pombe*

reference genome (ASM294v2.20) or *D. melanogaster* reference genome (release 6.06) with

Bowtie(Langmead and Salzberg, 2012) using optional arguments '-I 10 -X 700 --local --very-sensitive-

local --no-unal --no-mixed --no-discordant'. Details of th`e analysis pipeline depending on the

experimental technique used are described below.

Analysis of RNA-seq data

SAM files for *S. cerevisiae* data were used as an input for HTseq-count (Anders et al., 2015) with default

settings. The GFF file with *S. cerevisiae* genomic features was downloaded from the Ensembl website

(assembly R64-1-1). Signal per gene was normalized by the number of all *S. pombe* reads mapped for

the sample and multiplied by 10000 (arbitrarily chosen number). Genes classified as dubious,

pseudogenes or transposable elements were excluded leaving 5797 genes for the downstream

analysis. As a next filtering step, we excluded all the genes that had no measurable signal in at least

one out of 42 samples collected in this work. The remaining 5313 genes were used to calculate

coefficient of variation (CV) to validate reproducibility between replicate experiments (**Figures S1D and**

S4B and Table S2). This gene set was further compared to a list of 4900 genes we found previously to

provide the best reproducibility with a minimal loss of information(Donczew et al., 2020). The

overlapping set of 4883 genes was used in all plots where genes were divided into TFIID-dependent

and coactivator-redundant (CR) categories. The results of biological replicate experiments for each

sample were averaged. Corresponding samples were compared to calculate log$_2$ change in expression

per gene (IAA to DMSO samples for degron experiments, and *BDF2* deletion mutant to WT strain

(BY4705)) (**Table S2**).

Analysis of ChEC-seq data

SAM files for *S. cerevisiae* data were converted to tag directories with the HOMER

(http://homer.ucsd.edu (Heinz et al., 2010)) 'makeTagDirectory' tool. Data for six DMSO treated

replicate samples from experiments involving Bdf1, Bdf2 or Bdf1/2 depletion were used to define

promoters bound by Bdf1, Bdf2, Taf1, Taf11, Med8, Med17 and Spt3 as described below (**Table S4**).

Peaks were called using HOMER 'findPeaks' tool with optional arguments set to '-o auto -C 0 L 2 F 2',

with the free MNase dataset used as a control. These settings use a default false discovery rate (0.1%)

and require peaks to be enriched 2-fold over the control and 2-fold over the local background. Resulting

peak files were converted to BED files using 'pos2bed.pl' program. For each peak, the peak summit

was calculated as a mid-range between peak borders. For peak assignment to promoters the list of all

annotated ORF sequences (excluding sequences classified as 'dubious' or 'pseudogene') was

downloaded from the SGD website (https://www.yeastgenome.org). Data for 5888 genes were merged

with TSS positions (Park et al., 2014). If the TSS annotation was missing, TSS was manually assigned

at position −100 bp relative to the start codon. Peaks were assigned to promoters if their peak summit

was in the range from −300 to +100 bp relative to TSS. In a few cases, where more than one peak was

assigned to the particular promoter, the one closer to TSS was used. Promoters bound in at least four

out of six replicate experiments were included in a final list of promoters bound by a given factor and

were used to calculate promoter occupancy in all relevant experiments.

Coverage at each base pair of the *S. cerevisiae* genome was calculated as the number of reads that

mapped at that position divided by the number of all *D. melanogaster* reads mapped for the sample and

multiplied by 10000 (arbitrarily chosen number). To quantify log$_2$ change in factor promoter occupancy

signal per promoter was calculated as a sum of normalized reads per base in a 200 bp window around

the promoter peak summit. Peak summits were defined using Homer as described above. If no peak

was assigned to a promoter using Homer, the position of the strongest signal around TSS was used as

a peak summit. Manual inspection of selected cases confirmed the validity of this approach.

Corresponding IAA and DMSO treated samples were compared to calculate log$_2$ change in occupancy

and the results of biological replicate experiments for each sample were averaged (**Table S4**).

Analysis of ChIP-seq data

Data for Bdf1 was used to call peaks as described for ChEC-seq with input sample used as a control.

Promoters bound in at least one out of two replicate experiments were included in a final list of Bdf1

bound promoters as defined by ChIP-seq. For all samples coverage at each base pair of the *S.*

cerevisiae genome was calculated as the number of reads that mapped at that position divided by the

number of all *S. pombe* reads mapped in the sample, multiplied by the ratio of *S. pombe* to *S. cerevisiae*

reads in the corresponding input sample and multiplied by 10000 (arbitrarily chosen number). The list

of all annotated ORF sequences (excluding sequences classified as 'dubious' or 'pseudogene') was

downloaded from the SGD website (https://www.yeastgenome.org). Data for 5888 genes were merged

with TSS positions obtained from(Park et al., 2014). If the TSS annotation was missing the TSS was

manually assigned at position −100 bp relative to the start codon. Signal per gene was calculated as a

sum of normalized reads per base in a fixed window relative to TSS (defined in figure legends). The

results of biological duplicate experiments for each sample were averaged and the $\log_2$ change in signal

per gene was calculated by comparing corresponding IAA and DMSO treated samples (**Table S4**). In

experiments mapping Rpb1 CTD, Bur1, Ctk1 and Spt5, the signal per gene was calculated between

TSS and PAS(Park et al., 2014) (**Table S5**).

FASTQ files for the Bdf1 ChIP-exo (SRR397550) and H3 ChIP-seq (SRR6495880, SRR6495888,

SRR6495913, SRR6495921) experiments were obtained from the SRA. Data were processed as

described above except for the use of RPM normalization. The four datasets for H3 were averaged and

used to normalize H4K12ac data.

Phylogenetic analysis

Amino acid sequences of yeast and human BET proteins and human Taf1 were downloaded from NCBI

RefSeq database. Positions of individual bromodomains were obtained from (Wu and Chiang, 2007).

Multiple sequence alignment of individual bromodomains was performed using Clustal Omega (Sievers

et al., 2011). Neighbor joining tree was visualized with Jalview v 2.11.1.3 using PAM250 scoring matrix

(Waterhouse et al., 2009).

Gene ontology analysis

Gene ontology analysis for yeast and human BET regulated genes was done using PANTHER

overrepresentation test and PANTHER GO-Slim Biological Process annotation dataset with default

settings (http://www.pantherdb.org; v 16.0) (Mi et al., 2021). Only the most specific, enriched child

processes with FDR < 0.05 were used to compare BET functions between yeast and human model

systems.

Competing interests

No competing interests declared.

Acknowledgements

We thank Christine Cucinotta and Sarah Swygert for helpful comments about ChIP-seq experiments, Toshi Tsukiyama for plasmids, and Sandipan Brahma and all members of the Hahn lab for comments on manuscript. This work was supported by grants NIH GM053451 and GM075114 to SH and NIH P30 CA015704 to the Fred Hutch Genomics and Computational Shared Resources facility.

References

Allen BL, Taatjes DJ. 2015. The Mediator complex: a central integrator of transcription. *Nat Rev Mol Cell Biol* **16**:155–166. doi:10.1038/nrm3951

Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinforma Oxf Engl* **31**:166–169. doi:10.1093/bioinformatics/btu638

Badjatia N, Rossi MJ, Bataille AR, Mittal C, Lai WKM, Pugh BF. 2021. Acute stress drives global repression through two independent RNA polymerase II stalling events in Saccharomyces. *Cell Rep* **34**:108640. doi:10.1016/j.celrep.2020.108640

Bauer K, Berghoff AS, Preusser M, Heller G, Zielinski CC, Valent P, Grunt TW. 2021. Degradation of BRD4 - a promising treatment approach not only for hematologic but also for solid cancer. *Am J Cancer Res* **11**:530–545.

Bhagwat AS, Roe J-S, Mok BYL, Hohmann AF, Shi J, Vakoc CR. 2016. BET Bromodomain Inhibition Releases the Mediator Complex from Select cis-Regulatory Elements. *Cell Rep* **15**:519–530. doi:10.1016/j.celrep.2016.03.054

Bruzzone MJ, Grünberg S, Kubik S, Zentner GE, Shore D. 2018. Distinct patterns of histone acetyltransferase and Mediator deployment at yeast protein-coding genes. *Genes Dev* **32**:1252–1265. doi:10.1101/gad.312173.118

Chan LY, Mugler CF, Heinrich S, Vallotton P, Weis K. 2018. Non-invasive measurement of mRNA decay reveals translation initiation as the major determinant of mRNA stability. *eLife* **7**. doi:10.7554/eLife.32536

Chang CS, Pillus L. 2009. Collaboration between the essential Esa1 acetyltransferase and the Rpd3 deacetylase is mediated by H4K12 histone acetylation in Saccharomyces cerevisiae. *Genetics* **183**:149–160. doi:10.1534/genetics.109.103846

Core L, Adelman K. 2019. Promoter-proximal pausing of RNA polymerase II: a nexus of gene regulation.
Genes Dev **33**:960–982. doi:10.1101/gad.325142.119

Delmore JE, Issa GC, Lemieux ME, Rahl PB, Shi J, Jacobs HM, Kastritis E, Gilpatrick T, Paranal RM,
Qi J, Chesi M, Schinzel AC, McKeown MR, Heffernan TP, Vakoc CR, Bergsagel PL, Ghobrial
IM, Richardson PG, Young RA, Hahn WC, Anderson KC, Kung AL, Bradner JE, Mitsiades CS.
2011. BET bromodomain inhibition as a therapeutic strategy to target c-Myc. *Cell* **146**:904–917.
doi:10.1016/j.cell.2011.08.017

Devaiah BN, Lewis BA, Cherman N, Hewitt MC, Albrecht BK, Robey PG, Ozato K, Sims RJ, Singer DS.
2012. BRD4 is an atypical kinase that phosphorylates serine2 of the RNA polymerase II carboxy-
terminal domain. *Proc Natl Acad Sci U S A* **109**:6927–6932. doi:10.1073/pnas.1120422109

Donczew R, Lalou A, Devys D, Tora L, Hahn S. 2021. An improved ChEC-seq method accurately maps
the genome-wide binding of transcription coactivators and sequence-specific transcription
factors. *bioRxiv* 2021.02.12.430999. doi:10.1101/2021.02.12.430999

Donczew R, Warfield L, Pacheco D, Erijman A, Hahn S. 2020. Two roles for the yeast transcription
coactivator SAGA and a set of genes redundantly regulated by TFIID and SAGA. *eLife*
9:e50109. doi:10.7554/eLife.50109

Durant M, Pugh BF. 2007. NuA4-Directed Chromatin Transactions throughout the Saccharomyces
cerevisiae Genome. *Mol Cell Biol* **27**:5327–5335. doi:10.1128/MCB.00468-07

Fu J, Hou J, Liu L, Chen L, Wang M, Shen Y, Zhang Z, Bao X. 2013. Interplay between BDF1 and BDF2
and their roles in regulating the yeast salt stress response. *FEBS J* **280**:1991–2001.
doi:10.1111/febs.12219

Fujisawa T, Filippakopoulos P. 2017. Functions of bromodomain-containing proteins and their roles in
homeostasis and cancer. *Nat Rev Mol Cell Biol* **18**:246. doi:10.1038/nrm.2016.143

Gilan O, Rioja I, Knezevic K, Bell MJ, Yeung MM, Harker NR, Lam EYN, Chung C-W, Bamborough P,
Petretich M, Urh M, Atkinson SJ, Bassil AK, Roberts EJ, Vassiliadis D, Burr ML, Preston AGS,
Wellaway C, Werner T, Gray JR, Michon A-M, Gobbetti T, Kumar V, Soden PE, Haynes A,
Vappiani J, Tough DF, Taylor S, Dawson S-J, Bantscheff M, Lindon M, Drewes G, Demont EH,
Daniels DL, Grandi P, Prinjha RK, Dawson MA. 2020. Selective targeting of BD1 and BD2 of
the BET proteins in cancer and immunoinflammation. *Science* **368**:387–394.
doi:10.1126/science.aaz8455

Goldstein AL, McCusker JH. 1999. Three new dominant drug resistance cassettes for gene disruption
in Saccharomyces cerevisiae. *Yeast Chichester Engl* **15**:1541–1553. doi:10.1002/(SICI)1097-
0061(199910)15:14<1541::AID-YEA476>3.0.CO;2-K

Grünberg S, Henikoff S, Hahn S, Zentner GE. 2016. Mediator binding to UASs is broadly uncoupled
from transcription and cooperative with TFIID recruitment to promoters. *EMBO J* **35**:2435–2446.
doi:10.15252/embj.201695020

Gyuris A, Donovan DJ, Seymour KA, Lovasco LA, Smilowitz NR, Halperin ALP, Klysik JE, Freiman RN.
2009. The chromatin-targeting protein Brd2 is required for neural tube closure and
embryogenesis. *Biochim Biophys Acta* **1789**:413–421. doi:10.1016/j.bbagrm.2009.03.005

Han X, Yu D, Gu R, Jia Y, Wang Q, Jaganathan A, Yang X, Yu M, Babault N, Zhao C, Yi H, Zhang Q,
Zhou M-M, Zeng L. 2020. Roles of the BRD4 short isoform in phase separation and active gene
transcription. *Nat Struct Mol Biol* **27**:333–341. doi:10.1038/s41594-020-0394-8

Harlen KM, Churchman LS. 2017. The code and beyond: transcription regulation by the RNA
polymerase II carboxy-terminal domain. *Nat Rev Mol Cell Biol* **18**:263–273.
doi:10.1038/nrm.2017.10

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK.
2010. Simple combinations of lineage-determining transcription factors prime cis-regulatory
elements required for macrophage and B cell identities. *Mol Cell* **38**:576–589.
doi:10.1016/j.molcel.2010.05.004

Huisinga KL, Pugh BF. 2004. A genome-wide housekeeping role for TFIID and a highly regulated stress-
related role for SAGA in Saccharomyces cerevisiae. *Mol Cell* **13**:573–585.

Itzen F, Greifenberg AK, Bösken CA, Geyer M. 2014. Brd4 activates P-TEFb for RNA polymerase II
CTD phosphorylation. *Nucleic Acids Res* **42**:7577–7590. doi:10.1093/nar/gku449

Jacobson RH, Ladurner AG, King DS, Tjian R. 2000. Structure and function of a human TAFII250
double bromodomain module. *Science* **288**:1422–1425. doi:10.1126/science.288.5470.1422

Johnson KM, Wang J, Smallwood A, Arayata C, Carey M. 2002. TFIID and human mediator coactivator
complexes assemble cooperatively on promoter DNA. *Genes Dev* **16**:1852–1863.
doi:10.1101/gad.995702

Joo YJ, Ficarro SB, Soares LM, Chun Y, Marto JA, Buratowski S. 2017. Downstream promoter
interactions of TFIID TAFs facilitate transcription reinitiation. *Genes Dev* **31**:2162–2174.
doi:10.1101/gad.306324.117

Kanno T, Kanno Y, LeRoy G, Campos E, Sun H-W, Brooks SR, Vahedi G, Heightman TD, Garcia BA,
Reinberg D, Siebenlist U, O'Shea JJ, Ozato K. 2014. BRD4 assists elongation of both coding
and enhancer RNAs by interacting with acetylated histones. *Nat Struct Mol Biol* **21**:1047–1057.
doi:10.1038/nsmb.2912

Khoueiry P, Ward Gahlawat A, Petretich M, Michon AM, Simola D, Lam E, Furlong EE, Benes V,
Dawson MA, Prinjha RK, Drewes G, Grandi P. 2019. BRD4 bimodal binding at promoters and

drug-induced displacement at Pol II pause sites associates with I-BET sensitivity. *Epigenetics*

Chromatin **12**:39. doi:10.1186/s13072-019-0286-5

Knoll ER, Zhu ZI, Sarkar D, Landsman D, Morse RH. 2018. Role of the pre-initiation complex in Mediator

recruitment and dynamics. *eLife* **7**:e39633. doi:10.7554/eLife.39633

Koerber RT, Rhee HS, Jiang C, Pugh BF. 2009. Interaction of transcriptional regulators with specific

nucleosomes across the Saccharomyces genome. *Mol Cell* **35**:889–902.

doi:10.1016/j.molcel.2009.09.011

Krogan NJ, Keogh M-C, Datta N, Sawa C, Ryan OW, Ding H, Haw RA, Pootoolal J, Tong A, Canadien

V, Richards DP, Wu X, Emili A, Hughes TR, Buratowski S, Greenblatt JF. 2003. A Snf2 family

ATPase complex required for recruitment of the histone H2A variant Htz1. *Mol Cell* **12**:1565–

1576. doi:10.1016/s1097-2765(03)00497-0

Ladurner AG, Inouye C, Jain R, Tjian R. 2003. Bromodomains Mediate an Acetyl-Histone Encoded

Antisilencing Function at Heterochromatin Boundaries. *Mol Cell* **11**:365–376.

doi:10.1016/S1097-2765(03)00035-2

Lambert J-P, Picaud S, Fujisawa T, Hou H, Savitsky P, Uusküla-Reimand L, Gupta GD, Abdouni H, Lin

Z-Y, Tucholska M, Knight JDR, Gonzalez-Badillo B, St-Denis N, Newman JA, Stucki M, Pelletier

L, Bandeira N, Wilson MD, Filippakopoulos P, Gingras A-C. 2019. Interactome Rewiring

Following Pharmacological Targeting of BET Bromodomains. *Mol Cell* **73**:621-638.e17.

doi:10.1016/j.molcel.2018.11.006

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**:357–359.

doi:10.1038/nmeth.1923

Lauberth SM, Nakayama T, Wu X, Ferris A, Tang Z, Hughes SH, Roeder RG. 2013. H3K4me3

Interactions with TAF3 Regulate Preinitiation Complex Assembly and Selective Gene Activation.

Cell **152**:1021–1036. doi:10.1016/j.cell.2013.01.052

Lovén J, Hoke HA, Lin CY, Lau A, Orlando DA, Vakoc CR, Bradner JE, Lee TI, Young RA. 2013.

Selective inhibition of tumor oncogenes by disruption of super-enhancers. *Cell* **153**:320–334.

doi:10.1016/j.cell.2013.03.036

Matangkasombut O, Buratowski RM, Swilling NW, Buratowski S. 2000. Bromodomain factor 1

corresponds to a missing piece of yeast TFIID. *Genes Dev* **14**:951–962.

Matangkasombut O, Buratowski S. 2003. Different sensitivities of bromodomain factors 1 and 2 to

histone H4 acetylation. *Mol Cell* **11**:353–363.

Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, Thomas PD. 2021. PANTHER

version 16: a revised family classification, tree-based classification tool, enhancer regions and

extensive API. *Nucleic Acids Res* **49**:D394–D403. doi:10.1093/nar/gkaa1106

Muhar M, Ebert A, Neumann T, Umkehrer C, Jude J, Wieshofer C, Rescheneder P, Lipp JJ, Herzog
VA, Reichholf B, Cisneros DA, Hoffmann T, Schlapansky MF, Bhat P, von Haeseler A, Köcher
T, Obenauf AC, Popow J, Ameres SL, Zuber J. 2018. SLAM-seq defines direct gene-regulatory
functions of the BRD4-MYC axis. *Science* **360**:800–805. doi:10.1126/science.aao2793

Nguyen VQ, Ranjan A, Liu S, Tang X, Ling YH, Wisniewski J, Mizuguchi G, Li KY, Jou V, Zheng Q,
Lavis LD, Lionnet T, Wu C. 2020. Spatio-Temporal Coordination of Transcription Preinitiation
Complex Assembly in Live Cells. *bioRxiv* 2020.12.30.424853. doi:10.1101/2020.12.30.424853

Nishimura K, Fukagawa T, Takisawa H, Kakimoto T, Kanemaki M. 2009. An auxin-based degron system
for the rapid depletion of proteins in nonplant cells. *Nat Methods* **6**:917–922.
doi:10.1038/nmeth.1401

Oberbeckmann E, Wolff M, Krietenstein N, Heron M, Ellins JL, Schmid A, Krebs S, Blum H, Gerland U,
Korber P. 2019. Absolute nucleosome occupancy map for the Saccharomyces cerevisiae
genome. *Genome Res* **29**:1996–2009. doi:10.1101/gr.253419.119

Park D, Morris AR, Battenhouse A, Iyer VR. 2014. Simultaneous mapping of transcript ends at single-
nucleotide resolution and identification of widespread promoter-associated non-coding RNA
governed by TATA elements. *Nucleic Acids Res* **42**:3736–3749. doi:10.1093/nar/gkt1366

Patel AB, Greber BJ, Nogales E. 2020. Recent insights into the structure of TFIID, its assembly, and its
binding to core promoter. *Curr Opin Struct Biol* **61**:17–24. doi:10.1016/j.sbi.2019.10.001

Rabani M, Levin JZ, Fan L, Adiconis X, Raychowdhury R, Garber M, Gnirke A, Nusbaum C, Hacohen
N, Friedman N, Amit I, Regev A. 2011. Metabolic labeling of RNA uncovers principles of RNA
production and degradation dynamics in mammalian cells. *Nat Biotechnol* **29**:436–442.
doi:10.1038/nbt.1861

Rahl PB, Lin CY, Seila AC, Flynn RA, McCuine S, Burge CB, Sharp PA, Young RA. 2010. c-Myc
regulates transcriptional pause release. *Cell* **141**:432–445. doi:10.1016/j.cell.2010.03.030

Rahman S, Sowa ME, Ottinger M, Smith JA, Shi Y, Harper JW, Howley PM. 2011. The Brd4
extraterminal domain confers transcription activation independent of pTEFb by recruiting
multiple proteins, including NSD3. *Mol Cell Biol* **31**:2641–2652. doi:10.1128/MCB.01341-10

Rhee HS, Pugh BF. 2012. Genome-wide structure and organization of eukaryotic pre-initiation
complexes. *Nature* **483**:295–301. doi:10.1038/nature10799

Sabari BR, Dall'Agnese A, Boija A, Klein IA, Coffey EL, Shrinivas K, Abraham BJ, Hannett NM, Zamudio
AV, Manteiga JC, Li CH, Guo YE, Day DS, Schuijers J, Vasile E, Malik S, Hnisz D, Lee TI, Cisse
II, Roeder RG, Sharp PA, Chakraborty AK, Young RA. 2018. Coactivator condensation at super-
enhancers links phase separation and gene control. *Science* **361**. doi:10.1126/science.aar3958

Sanders SL, Jennings J, Canutescu A, Link AJ, Weil PA. 2002. Proteomics of the Eukaryotic
Transcription Machinery: Identification of Proteins Associated with Components of Yeast TFIID

by   Multidimensional   Mass   Spectrometry.   *Mol   Cell   Biol*   **22**:4723–4738.
doi:10.1128/MCB.22.13.4723-4738.2002

Sdelci S, Lardeau C-H, Tallant C, Klepsch F, Klaiber B, Bennett J, Rathert P, Schuster M, Penz T,
Fedorov O, Superti-Furga G, Bock C, Zuber J, Huber KVM, Knapp S, Müller S, Kubicek S. 2016.
Mapping the chemical chromatin reactivation landscape identifies BRD4-TAF1 cross-talk. *Nat*
Chem Biol **12**:504–510. doi:10.1038/nchembio.2080

Shetty A, Kallgren SP, Demel C, Maier KC, Spatt D, Alver BH, Cramer P, Park PJ, Winston F. 2017.
Spt5 plays vital roles in the control of sense and antisense transcription elongation. *Mol Cell*
66:77-88.e5. doi:10.1016/j.molcel.2017.02.023

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding
J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of high-quality protein multiple
sequence alignments using Clustal Omega. *Mol Syst Biol* **7**:539. doi:10.1038/msb.2011.75

Slaughter MJ, Shanle EK, Khan A, Chua KF, Hong T, Boxer LD, Allis CD, Josefowicz SZ, Garcia BA,
Rothbart SB, Strahl BD, Davis IJ. 2021. HDAC inhibition results in widespread alteration of the
histone acetylation landscape and BRD4 targeting to gene bodies. *Cell Rep* **34**:108638.
doi:10.1016/j.celrep.2020.108638

Stathis A, Bertoni F. 2018. BET Proteins as Targets for Anticancer Treatment. *Cancer Discov* **8**:24–36.
doi:10.1158/2159-8290.CD-17-0605

Stonestrom AJ, Hsu SC, Jahn KS, Huang P, Keller CA, Giardine BM, Kadauke S, Campbell AE, Evans
P, Hardison RC, Blobel GA. 2015. Functions of BET proteins in erythroid gene expression. *Blood*
125:2825–2834. doi:10.1182/blood-2014-10-607309

Suka N, Suka Y, Carmen AA, Wu J, Grunstein M. 2001. Highly specific antibodies determine histone
acetylation site usage in yeast heterochromatin and euchromatin. *Mol Cell* **8**:473–479.
doi:10.1016/s1097-2765(01)00301-x

Uhlén M, Fagerberg L, Hallström BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson Å, Kampf C,
Sjöstedt E, Asplund A, Olsson I, Edlund K, Lundberg E, Navani S, Szigyarto CA-K, Odeberg J,
Djureinovic D, Takanen JO, Hober S, Alm T, Edqvist P-H, Berling H, Tegel H, Mulder J,
Rockberg J, Nilsson P, Schwenk JM, Hamsten M, von Feilitzen K, Forsberg M, Persson L,
Johansson F, Zwahlen M, von Heijne G, Nielsen J, Pontén F. 2015. Proteomics. Tissue-based
map of the human proteome. *Science* **347**:1260419. doi:10.1126/science.1260419

van Ingen H, van Schaik FMA, Wienk H, Ballering J, Rehmann H, Dechesne AC, Kruijzer JAW, Liskamp
RMJ, Timmers HTM, Boelens R. 2008. Structural insight into the recognition of the H3K4me3
mark   by   the   TFIID   subunit   TAF3.   *Struct   Lond   Engl   1993*   **16**:1245–1256.
doi:10.1016/j.str.2008.04.015

Wang N, Wu R, Tang D, Kang R. 2021. The BET family in immunity and disease. *Signal Transduct Target Ther* **6**:1–22. doi:10.1038/s41392-020-00384-4

Warfield L, Ramachandran S, Baptista T, Devys D, Tora L, Hahn S. 2017. Transcription of Nearly All Yeast RNA Polymerase II-Transcribed Genes Is Dependent on Transcription Factor TFIID. *Mol Cell* **68**:118-129.e5. doi:10.1016/j.molcel.2017.08.014

Waterhouse AM, Procter JB, Martin DMA, Clamp M, Barton GJ. 2009. Jalview Version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**:1189–1191. doi:10.1093/bioinformatics/btp033

Winter GE, Mayer A, Buckley DL, Erb MA, Roderick JE, Vittori S, Reyes JM, di Iulio J, Souza A, Ott CJ, Roberts JM, Zeid R, Scott TG, Paulk J, Lachance K, Olson CM, Dastjerdi S, Bauer S, Lin CY, Gray NS, Kelliher MA, Churchman LS, Bradner JE. 2017. BET Bromodomain Proteins Function as Master Transcription Elongation Factors Independent of CDK9 Recruitment. *Mol Cell* **67**:5-18.e19. doi:10.1016/j.molcel.2017.06.004

Wu S-Y, Chiang C-M. 2007. The Double Bromodomain-containing Chromatin Adaptor Brd4 and Transcriptional Regulation. *J Biol Chem* **282**:13141–13145. doi:10.1074/jbc.R700001200

Wu S-Y, Zhou T, Chiang C-M. 2003. Human mediator enhances activator-facilitated recruitment of RNA polymerase II and promoter recognition by TATA-binding protein (TBP) independently of TBP-associated factors. *Mol Cell Biol* **23**:6229–6242. doi:10.1128/mcb.23.17.6229-6242.2003

Zencir S, Dilg D, Rueda MP, Shore D, Albert B. 2020. Mechanisms coordinating ribosomal protein gene transcription in response to stress. *Nucleic Acids Res* **48**:11408–11420. doi:10.1093/nar/gkaa852

Zentner GE, Kasinathan S, Xin B, Rohs R, Henikoff S. 2015. ChEC-seq kinetics discriminates transcription factor binding sites by DNA sequence and shape in vivo. *Nat Commun* **6**:8733. doi:10.1038/ncomms9733

Zhang H, Roberts DN, Cairns BR. 2005. Genome-wide dynamics of Htz1, a histone H2A variant that poises repressed/basal promoters for activation through histone loss. *Cell* **123**:219–231. doi:10.1016/j.cell.2005.08.036

Zheng B, Aoi Y, Shah AP, Iwanaszko M, Das S, Rendleman EJ, Zha D, Khan N, Smith ER, Shilatifard A. 2021. Acute perturbation strategies in interrogating RNA polymerase II elongation factor function in gene expression. *Genes Dev*. doi:10.1101/gad.346106.120

Zuber J, Shi J, Wang E, Rappaport AR, Herrmann H, Sison EA, Magoon D, Qi J, Blatt K, Wunderlich M, Taylor MJ, Johns C, Chicas A, Mulloy JC, Kogan SC, Brown P, Valent P, Bradner JE, Lowe SW, Vakoc CR. 2011. RNAi screen identifies Brd4 as a therapeutic target in acute myeloid leukaemia. *Nature* **478**:524–528. doi:10.1038/nature10334

1010