**Heterologous reporter expression in the planarian *Schmidtea mediterranea* through somatic mRNA transfection**

Richard Nelson Hall[1,12], Uri Weill[2,12], Margarita Khariton[1], Sergio Leal-Ortiz[3], Leonard Drees[2], Chew Chai[1], Yuan Xue[1], Benyamin Rosental[4], Stephen R. Quake[1,5,6], Alejandro Sánchez Alvarado[7,8], Nicholas A. Melosh[3], Andrew Fire[9,10], Jochen C. Rink[2,*], Bo Wang[1,11,*]

[1]Department of Bioengineering, Stanford University, Stanford, CA, USA

[2]Department of Tissue Dynamics and Regeneration, Max Planck Institute for Biophysical Chemistry, Göttingen, Germany

[3]Department of Materials Science and Engineering, Stanford University, Stanford, CA, USA

[4]The Shraga Segal Department of Microbiology, Immunology, and Genetics, Faculty of Health Sciences, Center for Regenerative Medicine and Stem Cells, Ben Gurion University of the Negev, Beer-Sheva, Israel

[5]Department of Applied Physics, Stanford University, Stanford, CA, USA.

[6]Chan Zuckerberg Biohub, San Francisco, CA, USA.

[7]Stowers Institute for Medical Research, Kansas City, MO, USA

[8]Howard Hughes Medical Institute, Kansas City, MO, USA.

[9]Department of Pathology, [10]Department of Genetics, [11]Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA, USA

[12]These authors contributed equally to this work.

*Correspondence should be addressed to J.C.R. (jochen.rink@mpibpc.mpg.de) and B.W. (wangbo@stanford.edu)

## Abstract

Planarians have long been studied for their regenerative abilities, but they possess limited genetic tools due to challenges in gene delivery, expression, and detection, despite decades of work. We developed a toolbox for heterologous protein expression in planarian cells and in live animals. Specifically, we identified and optimized nanotechnological and chemical transfection methods to efficiently deliver mRNA encoding nanoluciferase into somatic cells, including planarian adult stem cells (neoblasts). The use of a luminescent reporter allowed us to quantitatively measure protein expression through spectroscopy and microscopy, thus overcoming the strong autofluorescent background of planarian tissues. Using this platform, we investigated the use of endogenous untranslated region (UTR) sequences and codon usage bias to post-transcriptionally alter gene expression. Our work provides a strong foundation for advancing exogenous gene expression and for the rapid prototyping of genetic constructs to accelerate the development of transgenic techniques in planarians.

**Introduction**

Planarian flatworms have fascinated generations of scientists with their astounding regenerative abilities and have played a critical role in our efforts to understand stem cells and regeneration (Newmark & Sánchez Alvarado, 2002; Reddien, 2018; Rink, 2018). Planarians can regenerate their entire body from a small tissue fragment. During this process, they restore their body axes and rebuild organs with appropriate proportions using a population of adult pluripotent somatic stem cells called neoblasts (Baguñà et al., 1989; Wagner et al., 2011; Zeng et al., 2018). In sexual strains, neoblasts also regenerate germ cells, breaching the Weissman barrier that separates soma and germline (Wang et al., 2007). While gene knockdown by RNA mediated genetic interference (RNAi) and sequencing techniques have been widely used in planarian research (Sánchez Alvarado & Newmark, 1999; Reddien et al., 2005; Collins et al., 2010; Böser et al., 2013; Duncan et al., 2015; Lakshmanan et al., 2016; Molinaro & Pearson, 2016; Grohme et al., 2018; Fincher et al., 2018; Plass et al., 2018; Mihaylova et al., 2018; Pascual-Carreras et al., 2020), tools for transgene expression have yet to be developed. Earlier efforts have used electroporation in whole animals (González-Estévez et al., 2003), and more recently in purified neoblasts under improved culture conditions (Lei et al., 2019). However, the lack of orthogonal verification of transgene expression combined with the intense autofluorescence of planarian tissue has limited the practical utility of both studies. It is unclear whether the issue is technical, owing to the vast range of experimental conditions to be explored, or biological, as a result of efficient gene silencing in planarians (Kim et al., 2019). Therefore, establishing a robust and quantitative positive control of transgene expression is still a paramount challenge in the planarian field.

Although transgene expression has become commonplace in many model organisms, establishing transgenesis *de novo* requires solving three coupled challenges: gene delivery, expression, and detection. Construct delivery and expression are necessary prerequisites for transgene detection. Conversely, developing delivery and expression methods are impossible without sensitive and quantitative detection methods (e.g., evaluating promoter activity requires a validated transgenic reporter, but measuring reporter expression requires a well characterized promoter). Therefore, the initial demonstration of transgene expression is essential as it can

transform the task of method development from a blind pursuit into a well constrained problem

of parameter optimization.

For delivery, most transgenesis protocols begin by injecting a zygote; however, this is

challenging in sexual planarians given that their zygotes are dispersed among numerous yolk

cells (Cardona et al., 2006; Davies et al., 2017). Furthermore, the commonly used planarian

strains are exclusively asexual, reproducing through fission and regeneration (Vila-Farré & Rink,

2018; Arnold, Benham-Pyle et al., 2019), making somatic cell transfection the only option. Such

transfection methods, however, are primarily optimized for vertebrate tissues and immortalized

cell lines. Here, to address the problem of delivery, we used a direct nanoscale injection method

to first establish a reference in delivering nucleic acids into planarian cells. Based on this

reference, we tested many other methods and identified several chemical transfection reagents

that also showed high delivery efficacy.

A parallel challenge is that measuring functional gene delivery often relies on expression as the

ultimate readout for efficiency. Due to the distant relationship between planarians and other

model organisms, it is unclear whether utilizing ubiquitously expressed reporter constructs used

in other research organisms would be sufficient. Furthermore, transgenes can be epigenetically

modified and silenced through endogenous mechanisms, as has been noted in other systems like

mouse and *C. elegans* (Chen et al., 2003; Aljohani et al., 2020). Therefore, we chose to transfect

*in vitro* synthesized mRNA to circumvent the numerous variables associated with DNA based

methods, such as nuclear import, transcription, and splicing, as well as for rapid expression due

to the limitations of planarian cell culture (Lei et al., 2019).

Finally, planarian autofluorescence limits the utility of commonly used fluorescent reporters,

especially during the initial stages of reporter optimization when signals may be weak. To

overcome this detection challenge, we used an alternative strategy which relies on luminescence.

Nanoluciferase (Nluc), derived from the deep sea shrimp *Oplophorus gracilirostris,* is a compact

(19 kDa) and stable luciferase which is orders of magnitude brighter than other luciferases

(England et al., 2016). This allows for high signal-to-noise quantification of luminescence from

Nluc expression, so even weak signals can be detected to guide further optimization.

Combining all these technical advances, we have established a robust means for exogenous

expression in both cell cultures and live animals of the planarian model species *Schmidtea*

*mediterranea*. Taking advantage of the sensitivity of our Nluc expression assay, we quantified

the influence of adding endogenous planarian untranslated regions (UTRs) and changed codon

usage bias to manipulate and optimize transgene expression through a wide dynamic range. Our

results not only provide the first positive control for exogenous gene expression in planarians to

guide the future development of planarian transgenesis, but also offer a new route to manipulate

the functional transcriptome in planarian cells. We anticipate that our strategies may also prove

useful in other asexually reproducing animals which lack a germline and cannot otherwise be

transformed through embryonic manipulations.

**Results**

*Nanoluciferase mRNA delivered through nanostraws is expressed in planarian cells*

In order to establish mRNA expression in the asexual strain of *S. mediterranea*, we sought to identify an efficient and species-agnostic platform for delivering genetic material into primary planarian cells. We selected nanostraws, which combine the robustness of mechanical methods like microinjection with the throughput of electrical methods like electroporation (Tay & Melosh, 2019).

Nanostraws are approximately100-200 nm wide hollow aluminum oxide tubes protruding from a polycarbonate substrate situated above a buffer reservoir containing the genetic material to be delivered (**Figure 1A-B**). Cells are centrifuged against the straws which engage in close contact with the cell membrane. Electric pulses are then used to locally porate the membranes and electrophorese genetic material through the nanostraws and directly into the cellular cytoplasm (**Figure 1 – figure supplement 1**). By only permeabilizing membranes immediately in contact with nanostraws, they have been shown to improve both delivery efficiency and viability of transformed cells (Xie et al., 2013; Cao et al., 2018).

To prepare cells for nanostraw delivery, we flow-sorted a neoblast enriched population (so-called X1FS) from dissociated planarians (Wagner et al., 2011). The resulting cells were uniform in size and depleted of debris, which are important preconditions for nanostraw delivery. Our initial experiments used capped and polyadenylated *in vitro* transcribed mRNA encoding the red fluorescent protein, mScarlet, fused to the planarian histone H2B (**Supplementary Table 1**). We reasoned that red fluorescence might be more easily detectable against planarian autofluorescence, which is biased towards shorter wavelengths, and that a nuclear localization signal might further enhance signal to noise. Initial experiments used a pulse train protocol that was empirically optimized for delivery in human hematopoietic stem cells, which have similar morphological characteristics to neoblasts (i.e., small size, minimally adherent, high nucleus to cytoplasm ratio) (Schmiderer et al., 2020).

Microscopic examination of the transfected cells showed a weak and sometimes nuclear localized fluorescence. However, negative controls also contained similarly fluorescent cells

(**Figure 1C**). Therefore, we performed flow cytometry analysis to quantify fluorescence signal

and found, even in this relatively uniform cell population, a broad distribution of fluorescence

intensity, spanning three orders of magnitude in both experimental and negative conditions

(**Figure 1D**). This highly variable autofluorescence in our measurements highlights the

difficulties of using fluorescent reporters. Since some cells genuinely exhibit brighter

fluorescence than others, false positives may be common and true positives could be obscured by

the broad autofluorescent background such that H2B-mScarlet expression cannot be

unambiguously assessed. Together, these results compelled us to seek an alternative non-

fluorescent reporter to quantify gene expression in planarian cells.

Unlike fluorophores, luciferases produce light as the result of an oxidative chemical reaction

with an exogenously added substrate (luciferin), and most animal tissues are devoid of auto-

luminescence due to the enzymatic nature of luciferases (**Figure 1E**). To take advantage of the

high signal-to-noise ratios achievable with luminescence-based reporters, we delivered capped

and polyadenylated mRNA encoding a planarian codon optimized nanoluciferase (sNluc1)

(**Supplementary Table 1**) into X1FS cells and transfected them as was done previously with

H2B-mScarlet mRNA. Transfected cells were maintained for 24 hrs in Iso-L15, a nutrient rich

medium with reduced osmolarity. A similar medium has been shown previously to improve

planarian cell viability and maintain neoblast pluripotency (Lei et al., 2019). Strikingly, we saw

that transfected cells produced a clear and reproducible luminescence signal over 100-fold above

the background signal of the negative controls. Moreover, the physical dimensions of the

nanostraws strongly influenced the signal intensity, as has been observed in previous nanostraw

transfection experiments (Xie et al., 2013), implicating that we were in the regime where

expression was dependent on the amount of mRNA delivered (**Figure 1F**). Overall, our results

provide unambiguous proof-of-principle for heterologous protein expression in planarian cells.

*A screen identifies chemical reagents to efficiently transfect planarian cells*

With a validated reporter construct in hand, we next sought to identify a more broadly accessible

mRNA delivery method. Chemical transfection reagents have become increasingly popular

because of their ease of use, high efficiency, and scalability, but they are often developed and

optimized for specific cell types.

To identify suitable reagents for planarian cell transfection, we screened a panel of commercially

available reagents by transfecting *in vitro* transcribed sNluc1 mRNA containing 5' and 3' UTR

sequences from the highly expressed planarian gene, YB1 (**Supplementary Table 1**). Our screen

used total cells from freshly dissociated animals rather than X1FS cells due to the large number

of individual transfections needed. Initially, all reagents were utilized according to the

manufacturer's recommended protocols (**Supplemental Table 2**). While most transfections had

little to no effect, Viromer and Trans-IT (Mirus) reagents consistently achieved luminescent

signals 100 to 1,000-fold above the negative control (**Figure 2A**).

To confirm that luminescent signals in the chemical transfections indeed measured Nluc

expression in planarian cells, we exposed Trans-IT transfected total cells 24 hr post transfection

(hpt) to NanoGlo-Live furimazine substrate (Promega) and imaged the cultures on an LV200

bioluminescence imaging system (Olympus). Individual luminescent cells positive for the live

cell indicator Calcein AM were apparent in Trans-IT transfections but never observed in controls

(**Figure 2B**). High magnification luminescence imaging confirmed the cytoplasmic origin of the

luminescence signal. These observations establish chemical delivery as a viable method for

mRNA transfection of planarian cells, with Viromer and Trans-IT as promising leads.

To further optimize Nluc expression, we tested various ratios between Nluc mRNA and reagent

components (**Figure 2C-D**). For Viromer, increasing both the amount of reagent and/or mRNA

in the transfection mix doubled the baseline signal, with 0.8 µL reagent per 1 µg mRNA per

reaction as an economic compromise for further experiments. A similar two-fold boost over the

baseline signal was also achieved for Trans-IT, but maximal signal intensities in cell cultures

remained below those achieved by Viromer transfections. To investigate the kinetics of

exogenous mRNA expression in both regimens, we performed time course experiments.

Luminescence was apparent as early as 4 hpt for both reagents, consistent with the expected

kinetics of mRNA transfection and translation. Luminescence continued to increase past the last

assayed time point at 72 hpt, which was surprising given the short transcript half-life, suggesting

that transfected Nluc mRNA is stable in the transfection complex and/or inside planarian cells,

allowing for sustained transfection and expression (**Figure 2E-F**).

Finally, with the planarian's prominence as a model for studies of regeneration, we sought to

determine whether neoblasts were amenable to chemical transfection. In the absence of

detectable anti-Nluc immunostaining or live imaging compatible with neoblast markers, we

compared the luminescence signal produced between parallel Viromer transfections of neoblast

enriched (X1FS) cell populations and an equal number of total cells. Both conditions showed

comparable signal 24 hpt. Since X1FS is not a pure population of neoblasts, and luminescence

may be coming from differentiated cells in the sample, we developed a new sorting strategy to

isolate a cell population (CRNeoblasts) containing approximately 94% *piwi-1$^+$* cells (**Figure 2 –**

**figure supplement 1**). When transfected and assayed 24 hpt, CRNeoblasts showed

approximately 25% of the expression measured in the bulk population, suggesting that neoblasts

can be transfected by our method. Altogether, these results establish chemical transfection to

deliver mRNA into planarian cells including neoblasts, the ease and scalability of which provide

a strong basis for further optimization of planarian transgene expression.

***Untranslated regions and codon optimization can modulate expression of delivered mRNA***

We next examined the influence of post-transcriptional regulators of gene expression such as

UTRs and codon usage bias, which can be exploited to further increase Nluc expression. The

ability to transfect and express exogenous mRNA also opens new possibilities for the analysis of

gene regulation in planarians, which have so far been limited to observing phenotypic changes

and measuring net RNA depletion under the influence of RNAi.

First, we considered non-coding regions, such as the 5' and 3' UTRs, which can restrict gene

expression to specific cell types (Brun et al., 2003; Merritt et al., 2008) and can contain targets of

small RNA silencing pathways (Diag et al., 2018). To identify UTR elements that can

significantly alter Nluc expression, we utilized expression data and genomic mappings available

on PlanMine (Rozanski et al., 2019) to identify four endogenous genes with high expression

across both neoblasts and differentiated cell types, reasoning that they may enhance Nluc mRNA

stability and/or translation efficiency in total cell transfections. We also limited our search to

shorter genes which mapped uniquely to the genome, allowing them to be more easily isolated

via PCR. We cloned the 5' and 3' UTRs to flank sNluc1 (**Figure 3 – figure supplement 1**) and

produced capped and polyadenylated mRNA from these templates (**Figure 3 – figure**

**supplement 2**). We found that UTRs either increased (RPL15, YB1, RPL10) or decreased

(ENO) expression of Nluc relative to the construct lacking endogenous UTRs (No UTR

condition, see **Supplementary Table 3** for sequence details) demonstrating the significant

influence of UTRs on planarian transgene expression (**Figure 3A**). Further, the 3-fold signal

boost granted by RPL15 UTRs suggests that these sequences are promising starting points for

further expression optimization.

Second, we considered codon optimization, the commonplace practice of altering synonymous

codons of a transgene to better match the preferences of the host organism in order to increase

expression. Codon composition has been shown to modulate translation kinetics and mRNA

stability (Quax et al., 2015; Jeacock et al., 2018); moreover, the global A/T-bias in the genome

(70% A/T) and a strong preference for A/T at wobble positions make codon optimization an

important consideration in *S. mediterranea*. To investigate this effect, we generated another five

codon optimized variants in addition to the baseline Nluc construct that cover a range of codon

adaptation indices (CAI), a 0-1 bounded value measuring the genes codon bias compared to a

reference codon preference table (Puigbò et al., 2008) (**Supplementary Table 1**). We also

included a Nluc sequence optimized for mammalian expression as a reference for comparison.

Genes with higher CAI values tended to have lower GC content, reflecting the A/T bias in *S.*

*mediterranea* genome (**Supplementary Table 1**) (Grohme et al., 2018).

The luminescence quantifications in total live cell transfections did not reveal the expected

correlation between luminescence signal and CAI (Jeacock et al., 2018). Instead, the construct

with the highest CAI (sNluc0; 0.905) showed the weakest expression, while hNluc, optimized for

expression in human cells, with an accordingly poor planarian CAI of 0.509, was the second

most highly expressed (**Figure 3B**). The most highly expressed construct (sNluc2; CAI = 0.725)

was part of a series of constructs, with very similar CAIs and GC content, generated by randomly

sampling the planarian codon table (sNluc2-4; **Supplementary Tables 1, 3**). These results hint

at a complex relationship between codon usage bias and gene expression in planarians;

nevertheless, they provide us with a coding sequence (sNluc2) that provides about approximately

2 fold higher expression than the original baseline (sNluc1) construct.

After observing variations between constructs with different UTRs and codon optimized Nluc

sequences, we asked whether their effects might be additive. To this end, we constructed a range

of combined constructs mixing UTRs and codon optimized Nluc sequences and assayed their

expression in total cells (**Figure 3C**). Although the "best of both" construct (RPL15-sNluc2) had

the highest average expression, it did not show a significant difference compared to other

constructs. Moreover, the stark difference between the UTR effects of ENO (worst) and RPL15

(best) disappeared when combined with sNluc2. The "worst" codon-optimized sNluc0 in

combination with RPL15 UTRs was also expressed at a level comparable to the other constructs.

Overall, these results highlight the complex interactions between different construct design

variables. Regardless, they identify RPL15-sNluc2 as an improved reporter construct for future

experiments.

By testing the effects of UTR sequences, codon optimization, and combinations thereof, we have

demonstrated a framework for investigating post-transcriptional factors modulating the

expression of exogenously introduced mRNAs in planarians. Furthermore, we anticipate that the

optimizations made here will hold true for DNA-based transgenic methods in the future.

### *Heterologous reporter expression in vivo*

Although the ability to transfect planarian cells with mRNA *in vitro* represents a significant

advance in the genetic manipulation of planarian cells, the lack of neoblast proliferation in

culture limits the range of questions that can be addressed *in vitro*. We therefore explored

whether our transfection protocols might be sufficient for generating detectable reporter

expression *in vivo*.

We injected RPL15-sNluc2 mRNA complexed with Viromer or Trans-IT into the parenchymal

tissue along the tail midline, which reduces the risk of misinjections into the abundant gut

branches. We then measured luminescence on individually dissociated worms at multiple time

points post-transfection (**Figure 4A**). The luminescence background in tissue lysates from sham-

injected animals was universally very low, mirroring what was observed *in vitro*. In contrast,

mRNA injection with Viromer or Trans-IT led to luminescence signals up to 100-fold above

background (**Figure 4B**). For Viromer, the highest fraction of expressing animals was detected

already at 12 hpt (13/30 hpt) and tapered to 4/30 after 72 hpt (**Figure 4B**). Contrary to our

observations *in vitro*, Trans-IT transfections produced luminescence comparable to Viromer,

with similar kinetics and fraction of expressing animals (**Figure 4C**), suggesting that for *in vivo*

transfections, Trans-IT can be as effective as Viromer. The early peak and rapid decline in

luminescence intensity represents another difference compared to our previous *in vitro*

experiments and may be due to the reduced stability of transfection complexes *in vivo* or

efficient clearing of the transfection complexes by the planarian excretory system (Thi-Kim Vu

et al., 2015).

We next asked whether Nluc expression in live worms was sufficient for luminescence imaging.

Initial experiments showed that customary concentrations of the cell-permeable substrate

furimazine are well-tolerated by planarians, with no overt signs of toxicity even after overnight

incubations, which makes live imaging feasible. Injected worms were anaesthetized with

linalool, incubated for 15 min in furimazine solution, agarose-mounted on glass bottom dishes,

and imaged on a luminescence microscope. This experiment revealed bright luminescence in

both Viromer and Trans-IT-injected animals, while no luminescence was detected in negative

controls injected with mRNA alone (**Figure 4D**). The success rate observed through imaging

(~30%) matched our results from the plate reader assay. Moreover, luminescence was always

found within the vicinity of the tail injection site as expected if the signal were from Nluc mRNA

uptake and expression. Although the spatial resolution of *in vivo* luminescence imaging did not

allow for cellular resolution, the size of the injected region in the maximum projections was ~50

µm and consistent with the transfection of a small cluster of cells immediately adjacent to the

injection site. All together, these experiments demonstrate proof-of-principle for mRNA reporter

delivery and expression in live planarians.

**Discussion**

Transgene expression in planarian flatworms has been a persistent challenge in the field since the

molecular biology revival of the system around two decades ago (Agata & Watanabe, 1999;

Newmark & Sánchez Alvarado, 2002). Here, we accomplished proof-of-principle heterologous

protein expression in the planarian model species *S. mediterranea* by combining three

experimental approaches: (1) nanostraw electro-delivery to establish an initial positive control,

which enabled subsequent optimization of chemical delivery methods; (2) delivering mRNA

instead of DNA to bypass the complexities of nuclear import, transcription, and splicing, and

optimizing  post-transcriptional processes for enhanced expression; and (3) using a luminescent

reporter to circumvent the strong autofluorescence which complicates the use of fluorescent

reporters. By making these choices, we observed a clear signal from an exogenously supplied

Nluc mRNA in both planarian cells *in vitro* and *in vivo*.

Exogenous reporter expression in our results is supported by multiple independent lines of

evidence. First, we showed that planarians are minimally autoluminescent, and most

experimental conditions tested reproducibly reached luminescence intensities as high as 3-4

orders of magnitude above the background. Second, we observed that the luminescence showed

a dose-dependence on Nluc mRNA amounts and exhibited rapid expression kinetics

characteristic of mRNA expression. Third, luminescence intensity was modulated by biologically

relevant factors such as UTR sequences and codon usage. Fourth, imaging of transfected cell

cultures confirmed a cytoplasmic origin of the luminescence signal. Fifth, we succeeded in

imaging luminescence in injected live worms, with the signal consistently restricted to the

injection site. Finally, the results presented in this manuscript were gathered in two different

laboratories, which highlights the robustness and general applicability of the technique.

Collectively, our results represent the first robust demonstration of exogenous mRNA expression

in planarians.

In terms of applications, our current reporter assay already represents a significant expansion of

the planarian toolkit. Besides UTRs, our protocol allows for exploring other post-transcriptional

gene regulatory mechanisms in planarians. For example, incorporating trans-spliced 5' leader

sequences (Zayas et al., 2005; Rossi et al., 2014), adding target sites of known small RNAs (Kim

et al., 2019), including internal ribosomal entry sites (IRES), or manipulating secondary

structures of mRNA (Leppek et al., 2021) may enable the identification of modulatory elements

that have a stimulatory or suppressive effect on mRNA function and expression. Indeed, our

observation that CAI and expression were not well correlated, along with the non-additive effects

of UTRs and codon usage (**Figure 3B-C**), suggests that a complex set of sequence factors

influence the expression of exogenous mRNA. Our platform allows for the systematic

characterization and analysis of these factors to better predict how sequence informs expression.

Another important direction is to investigate the effect of modified nucleotides, which is gaining

traction as a means of increasing RNA expression and stability (Andries et al., 2015; Svitkin et

al., 2017). In fact, we have already compared transfections with RPL15-sNluc2 mRNA

incorporating the canonical uridine versus mRNA with the base analog 1-methyl pseudouridine

(m1Ψ). While we did not observe any significant changes in gene expression (**Figure 3 – figure**

**supplement 3**), our protocol provides a route to explore the effects of other modified nucleotides

on mRNA expression in planarian cells. Finally, while overexpression is desirable for labeling

and imaging purposes, many endogenous genes are expressed at much lower levels than most

constitutively active transgenic constructs might drive. In this sense, it will be important to

explore whether current expression levels are sufficient for achieving neomorphic phenotypes,

such as ectopic expression of Wnt or other signaling proteins with essential roles in planarian

body patterning (Petersen & Reddien, 2009; Stückemann et al., 2017).

A direct utility of our method is to provide a robust positive control and a baseline for further

optimization of gene delivery and reporter expression in planarians. First, utilizing RNA-based

synthetic biology tools like self-replicating RNA replicons including alphavirus (Beal et al.,

2015; Li et al., 2019) and nodavirus (Taning et al., 2018) may significantly enhance mRNA

driven expression, which is currently still transient and limited to small fractions of cells, as

observed in our *in vitro* imaging and *in vivo* transfection experiments (**Figure 2C**). Such tools

may enable increased expression to levels detectable through western blot or fluorescence

reporters. Second, specialized delivery protocol and expression constructs might be required to

optimize expression in specific cell types. Our study provides a simple procedure to quickly

isolate a highly pure population of neoblasts (**Figure 2 – figure supplement 1**) which opens the

possibility for identifying and optimizing transfection protocols specifically tailored for targeting

neoblasts. Third, with luminescence as a readout, DNA based transgene expression can be

quantitatively evaluated against an mRNA-based control. Constructs developed in this work can

also serve as the foundation for the design of DNA encoded reporters. Fourth, the ability to

deliver and express mRNA *in vivo* also opens the door to testing various genome editing

strategies such as CRISPR and transposon integration by co-transfecting Cas9 mRNA with a

repair template or transposase mRNA with an integration cassette, respectively. Finally, we

anticipate that further screens of transfection reagents using our constructs will allow for the

identification of other, potentially more effective, alternatives, and remove our current

dependence on a limited set of reagents. We note that Viromer reagents are no longer

commercially available since the supplier's (Lipocalyx) acquisition by BioNTech, one of the

SARS-CoV-2 mRNA vaccine producers. Although Viromer reagents are still available through

our labs upon request and Trans-IT (Mirus) reagent is equally effective at least for *in vivo*

transfection, there is an obvious need to enrich our options.

Altogether, our method proves conclusively that transgene expression is possible in the planarian

model species *S. mediterranea*. Our study provides a validated reporter construct, a quantitative

expression assay, and several transfection reagents, which collectively narrow the enormous

parameter space that make the *de novo* establishment of transgenesis in phylogenetically distant

model systems so challenging. As such, our results transform the quest for planarian transgenesis

from a blind trial-and-error process into a constrained parameter optimization process, which we

hope to proceed rapidly within our community.

**Materials and Methods**

**Planarian cell dissociation.** Planarian cells were prepared by finely mincing the worm with a razor blade and suspending the tissue in CMF (Ca/Mg-Free media: $NaH_2PO_4$ 480 mg/L, NaCl 960 mg/L, KCl 1.44 g/L, $NaHCO_3$ 960 mg/L, HEPES 3.57 g/L, D-glucose 0.24 g/L, BSA 1 g/L, pH 7.4 in MilliQ $H_2O$). The tissue was rocked for 5 min, followed by gentle pipetting for 10 min, or until the tissue was visibly homogenized. The cells were centrifuged at 250 g for 4 min, and the supernatant was removed and replaced with 1.5 mL of fresh CMF. The cell suspension was then serially filtered through 100, 70, 40, and 30-µm mesh strainers. The filtered cell suspension was centrifuged and transferred to Iso-L15 (1:1 Leibovitz's L-15 to MilliQ $H_2O$, $1\times$ MEM nonessential amino acids, $1\times$ antibiotic-antimycotic, $1\times$ MEM vitamin solution, 1 mM Sodium Pyruvate, 2.5 g/L HEPES, 5% FBS, buffer to pH 7.8).

**FACS.** To isolate neoblast-enriched X1FS population, an aliquot of sacrificial cells was stained with Hoechst 33342 (10 µg/mL) in CMF for 15 min, filtered, and sorted on a Sony SH800 with either the 100 or 130 µm sorting chip (LE-C3210, LE-C3213, Sony). Following the identification of the neoblast population using Hoechst fluorescence (Wagner et al., 2011), unstained planarian cells were loaded and sorted using the neoblast gate overlayed on the forward and side-scatter into Iso-L15 medium. After sorting, the cells were centrifuged at 250 g for 5 min and resuspended in fresh Iso-L15.

To isolate CRNeoblasts, dissociated planarian cells at a density of $1\text{-}5\times10^6$ cells/mL were stained with 1:500 (5 µM) CellRox Green (Invitrogen, C10444) and 1:7000 (143 nM) LysoTracker Deep Red (Invitrogen, L12492). The sample was rocked gently in the dark for 30 min at room temperature. After incubation, the cells were pipetted up and down before being strained through a 30 µm filter cap FACS tube. The cells were sorted on a Sony SH800 with a 100 µm sorting chip. Cells were first gated by forward and side-scatter, then a final gate around CellRox Green high and LysoTracker Deep Red low identified the CRNeoblast population (**Figure 2 – figure supplement 1**).

**Single-cell RNA-seq analysis of CRNeoblast.** Single cell SmartSeq2 protocol was carried out

as previously described (Li et al., 2021). Paired-end reads were mapped to dd_Smed_v6

reference transcriptome (Rozanski et al., 2019) using Salmon (v1.4.0) (Patro et al., 2017).

Downstream preprocessing and analysis were performed using estimated counts in the Salmon

output, for which we sum up counts from different isoforms of the same gene. Cells with fewer

than 2,400 genes detected were filtered out, passing 481 cells for downstream analysis. Raw

gene counts were then normalized for sequencing coverage such that each cell has a total read

count equal to that of the median library size for all cells. The resulting counts were added with a

pseudo count of 1 and log-2 transformed. 2D embedding was performed with the SAM algorithm

(version 0.8.1) (Tarashansky et al., 2019) using default parameters. *piwi-1$^+$* cells were defined as

those for which their log-2 transformed and normalized read counts were greater than zero. *piwi-*

*1$^+$* cells were clustered using the Leiden clustering algorithm, and each cluster was annotated

using progenitor marker genes previously identified (N_epidermal: *soxP3*,

dd_Smed_v6_5942_0_1; N_gut: *hnf*, dd_Smed_v6_1694_0_1; N_muscle: *pcdh11*,

dd_Smed_v6_9283_0_1, cNeoblast: *tgs1*, dd_Smed_v6_10988_0_1).

**Nanostraw electro-delivery.** Nanostraws were fabricated according to previously established

methods (Cao et al., 2018). Each nanostraw cartridge was loaded with 200,000 X1FS cells in 300

µL of Iso-L15 media and centrifuged at 300 g for 10 min to ensure close contact between straws

and cells. For each cartridge, 3 µg of *in vitro* synthesized mRNA was diluted in PBS to a total

volume of 35 µL and placed on the titanium anode, and a cartridge was carefully lowered onto

the mRNA solution. The titanium cathode was placed atop, and the electro-delivery assembly

was subjected to a 35 V, 200 µs, 40 Hz square wave pulse 3 times for 45 s each, with a 1 min

rest in between pulses. Transfected cells were incubated at 20°C in the dark for 24 hr before

being transferred from the nanostraw cartridge to a 96 well plate for assaying luminescence.

**Chemical transfection.** For *in vitro* experiments, dissociated planarian cells were suspended at a

concentration of 0.88×10$^6$ cells/mL in Iso-L15 medium supplemented with 10 µg/mL

ciprofloxacin. 225 µL of cell suspension was added to each well of a 96 well plate for a total of

approximately 200,000 cells per well. For the initial screen, each reagent was prepared as

specified in **Supplementary Table 2**. After adding transfection complexes to each well, the cells

were incubated at 20°C in the dark before assaying luminescence. For live animal injections,

transfection mixture was loaded on needles pulled from glass capillaries (Drummond). Needles

were loaded on a FemtoJet injection system (Eppendorf) and worms were injected with 900 nL

transfection mix. Worms were left to rest in the dark at 20°C until assaying.

**Nanoluciferase assay.** For dissociated cells, luminescence was measured using the NanoGlo

Live Cell Assay (Promega). Furimazine substrate was added to NanoGlo buffer at a ratio of 1:20,

and 25 µL of reagent was added to 250 µL of transfected cells. For live planarians, each injected

worm was individually dissociated by finely mincing the worm with a razor blade and

suspending the tissue in 250 µL Iso-L15 medium just prior to assaying. The resuspended tissue

was transferred to a 96-well white bottom plate. Nluc expression was measured using the

NanoGlo Assay Lysis (Promega) kit. Furimazine substrate was added to the NanoGlo lysis

buffer at a ratio of 1:50 and 100 µL of reagent was added to the cells. Cells were lysed by

pipetting up and down 10 times. Luminescence was measured on a plate reader (BioTek HTX, or

BioTek Synergy™ Neo2, or EnVision Microplate Reader). Integration time was set at 1 s and the

digital gain was kept consistent on each instrument for all experiments.

**Luminescence imaging.** For imaging planarian cells *in vitro,* Nluc expression was imaged using

the NanoGlo Live Cell Assay. 35 mm glass-bottom dishes were coated with 0.5 mg/mL

Concanavalin A (Sigma-Aldrich) dissolved in deionized water for 2 hrs, washed, and air dried.

Transfected cells were transferred to the coated dish in a total of 1 mL of Iso-L15, allowed to

adhere to the glass surface for 1 hr, and stained with Calcein AM (ThermoFisher) (4 ng/ µL) to

label live cells. 12.5 µL of NanoGlo live (Promega) substrate was added to 237.5 µL of NanoGlo

buffer, mixed, and then added to the media. Cells were imaged using an LV200 Bioluminescence

Imaging System (Olympus) with a 20× air objective (Olympus: UPLXAPO20X) or a 100× oil

immersion objective (Olympus: UPLXAPO100XO). For *in vivo* imaging, NanoGlo live

(Promega) substrate was added to planarian water with 1% DMSO at a ratio of 1:20. Worms

were incubated in 100 µL of reagent mixture for 15 min, and then mounted on glass bottom

dishes, embedded in agarose gel (1.5%) supplemented with 1:5000 linalool and 1:20 furimazine

substrate, which was covered with a cover slip. They were imaged for luminescence and

brightfield using an LV200 with a 20× air objective (Olympus: UPLXAPO20X).

**Cloning.** To generate a plasmid capable of *in vitro* transcription and for harvesting the

endogenous regulatory sequences (5' and 3' UTRs) of planarian genes, we first amplified the

backbone of pDONOR221 (ThermoFisher) using primers BW-NH-104-105 (all primer

sequences are provided in **Supplementary Table 4**), as well as the LacZ cassette from pUC19

(Addgene #50005) with a T7 promoter sequence followed by a BsaI restriction site for

subsequent cloning steps, all flanked between BbsI restriction sites and M13 forward and M13

reverse primer sites (for *in vitro* transcription template production) using primers BW-NH-106-

107. The amplified backbone was digested with BsaI-HF (NEB) and the LacZ insert was

digested with BbsI-HF (NEB). The digested backbones were purified using the Zymo Clean and

Concentrate kit (Zymo). The purified fragments were ligated together using T4 DNA Ligase

(NEB) to create pNHT7 (**Figure 3 – figure supplement 1A**).

We cloned the gene of interest (GOI) from a pool of planarian cDNA using primers (BW-NH-

108-125) containing BsaI restriction sites to produce overhangs compatible with pNHT7. The

amplicons were purified using the Zymo Clean and Concentrate kit and then inserted into

pNHT7 via a golden gate reaction containing 40 ng of backbone and 20 ng for each insert to be

cloned in a 20 µL reaction volume containing 2 µL T4 Ligase Buffer, 1 µL T4 DNA Ligase, and

1 µL BsaI-HF to produce pNHT7::GOI (**Figure 3 – figure supplement 1B**).

Finally, to insert a reporter between the 5' and 3' UTRs, we amplified pNHT7::GOI with

outward facing primers (BW-NH-128-139) which bind to the end and beginning of the 5' and 3'

UTRs respectively containing the BsaI restriction sites. The reporter was then amplified to

append compatible BsaI restriction sites using primers BW-NH-174-185 and inserted between

the two UTR sequences via a golden gate reaction. The resulting plasmid was amplified with

M13 forward and M13 reverse primers to produce linear template for *in vitro* transcription

reactions (**Figure 3 – figure supplement 1C**).

*In vitro* **transcription.** *In vitro* transcription was performed using the T7 mScript™ Standard

mRNA Production System (CELLSCRIPT, C-MSC100625) according to the manufacturer's

protocol, opting for a 1 hr incubation during T7 transcription, a 2 hr incubation for 5' capping,

and a 1 hr incubation for poly-A tailing. RNA purification was performed by adding 600 µL of

ethanol to 50 µL of 10 M ammonium acetate. A standard 60 µL reaction typically yields 60 µg

of mRNA. For expected results, see **Figure 3 – figure supplement 2**. For mRNA containing

m1Ψ, the rNTP mix provided in the CELLSCRIPT kit was substituted for a homemade mixture

of rNTPs containing 10 mM rGTP, 10 mM rCTP, 10 mM rATP, and 10 mM m1Ψ (TriLink).

**Acknowledgements**

We thank K. Lei, C. Kuhn, A. Voskoboynik for stimulating discussion, E. Liu and D. Burghardt

for experimental assistance, Y. Lim and T. Boothe for microscopy assistance. RNH and CC are

supported by NSF Graduate Research Fellowships. UW is supported by an EMBO Long-Term

Fellowship. BW is Beckman Young Investigator. UW, LD, and JCR are supported by the Max

Planck Society. AF is supported by a NIH grant (R35GM130366). BW and BR thank the support

from a HFSP grant (RGY0085/2019). This work was initiated through a Stanford Bio-X

Interdisciplinary Initiative seed grant (IIP9-64) to BW, AF, and NAM, and later supported by an

NSF EDGE grant (IOS-1923534) to BW, ASA, and NAM, and a Volkswagen Foundation grant

(Az.-99498) to JCR.

**References**

Agata, K., & Watanabe, K. (1999). Molecular and cellular aspects of planarian regeneration. *Seminars in Cell and Developmental Biology*, *10*(4), 377–383. https://doi.org/10.1006/scdb.1999.0324

Aljohani, M. D., El Mouridi, S., Priyadarshini, M., Vargas-Velazquez, A. M., & Frøkjær-Jensen, C. (2020). Engineering rules that minimize germline silencing of transgenes in simple extrachromosomal arrays in *C. elegans*. *Nature Communications*, *11*, 6300. https://doi.org/10.1038/s41467-020-19898-0

Andries, O., Mc Cafferty, S., De Smedt, S. C., Weiss, R., Sanders, N. N., & Kitada, T. (2015). N1-methylpseudouridine-incorporated mRNA outperforms pseudouridine-incorporated mRNA by providing enhanced protein expression and reduced immunogenicity in mammalian cell lines and mice. *Journal of Controlled Release*, *217*(2015), 337–344. https://doi.org/10.1016/j.jconrel.2015.08.051

Arnold, C. P., Benham-Pyle, B. W., Lange, J. J., Wood, C. J., & Sánchez Alvarado, A. (2019). Wnt and TGFβ coordinate growth and patterning to regulate size-dependent behaviour. *Nature*, *572*(7771), 655–659. https://doi.org/10.1038/s41586-019-1478-7

Baguñà, J., Saló, E., & Auladell, C. (1989). Regeneration and pattern formation in planarians. III. that neoblasts are totipotent stem cells and the cells. *Development*, *107*(1), 77–86.

Beal, J., Wagner, T. E., Kitada, T., Azizgolshani, O., Parker, J. M., Densmore, D., & Weiss, R. (2015). Model-driven engineering of gene expression from RNA replicons. *ACS Synthetic Biology*, *4*(1), 48–56. https://doi.org/10.1021/sb500173f

Böser, A., Drexler, H. C. A., Reuter, H., Schmitz, H., Wu, G., Schöler, H. R., Gentile, L., & Bartscherer, K. (2013) SILAC proteomics of planarians identifies Ncoa5 as a conserved component of pluripotent stem cells. *Cell Reports*, *5*(4), 1142–1155. https://doi: 10.1016/j.celrep.2013.10.035

Brun, S., Faucon-Biguet, N., & Mallet, J. (2003). Optimization of transgene expression at the posttranscriptional level in neural cells: Implications for gene therapy. *Molecular Therapy*, *7*(6), 782–789. https://doi.org/10.1016/S1525-0016(03)00097-2

Cao, Y., Chen, H., Qiu, R., Hanna, M., Ma, E., Hjort, M., Zhang, A., Lewis, R. S., Wu, J. C., & Melosh, N. A. (2018). Universal intracellular biomolecule delivery with precise dosage control. *Science Advances*, *4*(10), eaat8131. https://doi.org/10.1126/sciadv.aat8131

Cardona, A., Hartenstein, V., & Romero, R. (2006). Early embryogenesis of planaria: A cryptic

larva feeding on maternal resources. *Development Genes and Evolution*, *216*(11), 667–681.

https://doi.org/10.1007/s00427-006-0094-3

Chen, Z. Y., He, C. Y., Ehrhardt, A., & Kay, M. A. (2003). Minicircle DNA vectors devoid of

bacterial DNA result in persistent and high-level transgene expression in vivo. *Molecular*

*Therapy*, *8*(3), 495–500. https://doi.org/10.1016/S1525-0016(03)00168-0

Collins III, J. J., Hou, X., Romanova, E. V., Lambrus, B. G., Miller, C. M., Saberi, A., Sweedler,

J. V., & Newmark, P. A. (2010) Genome-wide analyses reveal a role for peptide hormones

in planarian germline development. *PLoS Biology*, *8*(10), e1000509. https://doi:

10.1371/journal.pbio.1000509.

Davies, E. L., Lei, K., Seidel, C. W., Kroesen, A. E., McKinney, S. A., Guo, L., Robb, S. M. C.,

Ross, E. J., Gotting, K., & Sánchez Alvarado, A. (2017). Embryonic origin of adult stem

cells required for tissue homeostasis and regeneration. *eLife*, *6*, e21052.

https://doi.org/10.7554/eLife.21052

Diag, A., Schilling, M., Klironomos, F., Ayoub, S., & Rajewsky, N. (2018). Spatiotemporal

m(i)RNA architecture and 3′ UTR regulation in the *C. elegans* germline. *Developmental*

*Cell*, *47*(6), 785-800.e8. https://doi.org/10.1016/j.devcel.2018.10.005

Duncan, E. M., Chitsazan, A. D., Seidel, C. W., & Alvarado, A. S. (2015). Set1 and MLL1/2

target distinct sets of functionally different genomic loci in vivo. *Cell Reports*, *13*(12),

2741–2755. https://doi.org/10.1016/j.celrep.2015.11.059

England, C. G., Ehlerding, E. B., & Cai, W. (2016). NanoLuc: A small luciferase is brightening

up the field of bioluminescence. *Bioconjugate Chemistry*, *27*(5), 1175–1187.

https://doi.org/10.1021/acs.bioconjchem.6b00112

Fincher, C. T., Wurtzel, O., de Hoog, T., Kravarik, K. M., & Reddien, P. W. (2018). Cell type

transcriptome atlas for the planarian *Schmidtea mediterranea*. *Science*, *360*(6391),

eaaq1736. https://doi.org/10.1126/science.aaq1736

González-Estévez, C., Momose, T., Gehring, W. J., & Saló, E. (2003). Transgenic planarian lines

obtained by electroporation using transposon-derived vectors and an eye-specific GFP

marker. *Proceedings of the National Academy of Sciences of the United States of America*,

*100*(24), 14046–14051. https://doi.org/10.1073/pnas.2335980100

Grohme, M. A., Schloissnig, S., Rozanski, A., Pippel, M., Young, G. R., Winkler, S., Brandl, H.,

Henry, I., Dahl, A., Powell, S., Hiller, M., Myers, E., & Rink, J. C. (2018). The genome of

*Schmidtea mediterranea* and the evolution of core cellular mechanisms. *Nature*, *554*(7690),

56–61. https://doi.org/10.1038/nature25473

Jeacock, L., Faria, J., & Horn, D. (2018). Codon usage bias controls mRNA and protein

abundance in trypanosomatids. *eLife*, *7*, e32496. https://doi.org/10.7554/eLife.32496

Kim, I. V., Duncan, E. M., Ross, E. J., Gorbovytska, V., Nowotarski, S. H., Elliott, S. A.,

Sánchez Alvarado, A., & Kuhn, C. D. (2019). Planarians recruit piRNAs for mRNA

turnover in adult stem cells. *Genes & Development*, *33*(21–22), 1575–1590.

https://doi.org/10.1101/gad.322776.118

Lakshmanan, V., Bansai, D., Kulkarni, J., Poduval, D., Krishna, S., Sasidharan, V., Anand, P.,

Seshasayee, A., & Palakodeti, D. (2016). Genome-wide analysis of polyadenylation events

in *Schmidtea mediterranea*. *G3: Genes, Genomes, Genetics*, *6*(10), 3035–3048.

https://doi.org/10.1534/g3.116.031120

Lei, K., McKinney, S. A., Ross, E. J., Lee, H. C., & Sánchez Alvarado, A. (2019). Cultured

pluripotent planarian stem cells retain potency and express proteins from exogenously

introduced mRNAs. *bioRxiv*. https://doi.org/10.1101/573725

Leppek, K., Woo Byeon, G., Kladwang, W., Wayment-Steele, H. K., Kerr, C. H., Xu, A. F.,

Soon Kim, D., Topkar, V. V, Choe, C., Rothschild, D., Tiu, G. C., Wellington-Oguri, R.,

Fujii, K., Sharma, E., Watkins, A. M., Nicol, J. J., Romano, J., Tunguz, B., Participants, E.,

… Das, R. (2021). Combinatorial optimization of mRNA structure, stability, and translation

for RNA-based therapeutics. *bioRxiv*. https://doi.org/10.1101/2021.03.29.437587

Li, P., Nanes Sarfati, D., Xue, Y., Yu, X., Tarashansky, A. J., Quake, S. R., & Wang, B. (2021).

Single-cell analysis of *Schistosoma mansoni* identifies a conserved genetic program

controlling germline stem cell fate. *Nature Communications*, *12*, 485.

https://doi.org/10.1038/s41467-020-20794-w

Li, Y., Teague, B., Zhang, Y., Su, Z., Porter, E., Dobosh, B., Wagner, T., Irvine, D. J., & Weiss,

R. (2019). *In vitro* evolution of enhanced RNA replicons for immunotherapy. *Scientific

Reports*, *9*, 6932. https://doi.org/10.1038/s41598-019-43422-0

Merritt, C., Rasoloson, D., Ko, D., & Seydoux, G. (2008). 3' UTRs are the primary regulators of

gene expression in the *C. elegans* germline. *Current Biology*, *18*(19), 1476–1482.

https://doi.org/10.1016/j.cub.2008.08.013

Mihaylova, Y., Abnave, P., Kao, D., Hughes, S., Lai, A., Jaber-Hijazi, F., Kosaka, N., &

Aboobaker, A. A. (2018). Conservation of epigenetic regulation by the MLL3/4 tumour

suppressor in planarian pluripotent stem cells. *Nature Communications*, *9*, 3633.

https://doi.org/10.1038/s41467-018-06092-6

Molinaro, A. M., & Pearson, B. J. (2016). In silico lineage tracing through single cell

transcriptomics identifies a neural stem cell population in planarians. *Genome Biology*,

*17*(1), 87. https://doi.org/10.1186/s13059-016-0937-9

Newmark, P. A., & Sánchez Alvarado, A. (2002). Not your father's planarian: A classic model

enters the era of functional genomics. *Nature Reviews Genetics*, *3*(3), 210–219.

https://doi.org/10.1038/nrg759

Pascual-Carreras, E., Marín-Barba, M., Castillo-Lara, S., Coronel-Córdoba, P., Magri, M. S.,

Wheeler, G. N., Abril, J. F., Gomez-Skarmeta, J. L., Saló, E., & Adell, T. (2020). Genomic

analyses reveal FoxG as an upstream regulator of wnt1 required for posterior identity

specification in planarians. *bioRxiv*,. https://doi.org/10.1101/2020.12.08.416008

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast

and bias-aware quantification of transcript expression. *Nature Methods*, *14*(4), 417–419.

https://doi.org/10.1038/nmeth.4197

Petersen, C. P., & Reddien, P. W. (2009). A wound-induced Wnt expression program controls

planarian regeneration polarity. *Proceedings of the National Academy of Sciences of the*

*United States of America*, *106*(40), 17061–17066. https://doi.org/10.1073/pnas.0906823106

Plass, M., Solana, J., Alexander Wolf, F., Ayoub, S., Misios, A., Glažar, P., Obermayer, B.,

Theis, F. J., Kocks, C., & Rajewsky, N. (2018). Cell type atlas and lineage tree of a whole

complex animal by single-cell transcriptomics. *Science*, *360*(6391), eaaq1723.

https://doi.org/10.1126/science.aaq1723

Puigbò, P., Bravo, I. G., & Garcia-Vallvé, S. (2008). E-CAI: A novel server to estimate an

expected value of Codon Adaptation Index (eCAI). *BMC Bioinformatics*, *9*(1), 65.

https://doi.org/10.1186/1471-2105-9-65

Quax, T. E. F., Claassens, N. J., Söll, D., & van der Oost, J. (2015). Codon bias as a means to

fine-tune gene expression. *Molecular Cell*, *59*(2), 149–161.

https://doi.org/10.1016/j.molcel.2015.05.035

Reddien, P. W. (2018). The cellular and molecular basis for planarian regeneration. *Cell*, *175*(2),

327–345. https://doi.org/10.1016/j.cell.2018.09.021

Reddien, P. W., Bermange, A. L., Murfitt, K. J., Jennings, J. R., & Sánchez Alvarado, A. (2005). Identification of genes needed for regeneration, stem cell function, and tissue homeostasis by systematic gene perturbation in planaria. *Developmental Cell*, *8*(5), 635–649. https://doi: 10.1016/j.devcel.2005.02.014

Rink, J. C. (2018). Stem cells, patterning and regeneration in planarians: Self-organization at the organismal scale. *Methods in Molecular Biology*, *1774*, 57–172. https://doi.org/10.1007/978-1-4939-7802-1_2

Rossi, A., Ross, E. J., Jack, A., & Sánchez Alvarado, A. (2014). Molecular cloning and characterization of SL3: A stem cell-specific SL RNA from the planarian Schmidtea mediterranea. *Gene*, *533*(1), 156–167. https://doi.org/10.1016/j.gene.2013.09.101

Rozanski, A., Moon, H. K., Brandl, H., Martín-Durán, J. M., Grohme, M. A., Hüttner, K., Bartscherer, K., Henry, I., & Rink, J. C. (2019). PlanMine 3.0 - improvements to a mineable resource of flatworm biology and biodiversity. *Nucleic Acids Research*, *47*(D1), D812–D820. https://doi.org/10.1093/nar/gky1070

Sánchez Alvarado, A., & Newmark, P. A. (1999). Double-stranded RNA specifically disrupts gene expression during planarian regeneration. *Proceedings of the National Academy of Sciences of the United States of America*, *96*(9), 5049–5054. https://doi.org/10.1073/pnas.96.9.5049

Schmiderer, L., Subramaniam, A., Žemaitis, K., Bäckström, A., Yudovich, D., Soboleva, S., Galeev, R., Prinz, C. N., Larsson, J., & Hjort, M. (2020). Efficient and nontoxic biomolecule delivery to primary human hematopoietic stem cells using nanostraws. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(35), 21267–21273. https://doi.org/10.1073/pnas.2001367117

Stückemann, T., Cleland, J. P., Werner, S., Thi-Kim Vu, H., Bayersdorf, R., Liu, S. Y., Friedrich, B., Jülicher, F., & Rink, J. C. (2017). Antagonistic self-organizing patterning systems control maintenance and regeneration of the anteroposterior axis in planarians. *Developmental Cell*, *40*(3), 248–263. https://doi.org/10.1016/j.devcel.2016.12.024

Svitkin, Y. V., Cheng, Y. M., Chakraborty, T., Presnyak, V., John, M., & Sonenberg, N. (2017). N1-methyl-pseudouridine in mRNA enhances translation through eIF2α-dependent and independent mechanisms by increasing ribosome density. *Nucleic Acids Research*, *45*(10),

6023–6036. https://doi.org/10.1093/nar/gkx135

Taning, C. N. T., Christiaens, O., Li, X., Swevers, L., Casteels, H., Maes, M., & Smagghe, G. (2018). Engineered flock house virus for targeted gene suppression through RNAi in fruit flies (*Drosophila melanogaster*) *in vitro* and *in vivo*. *Frontiers in Physiology*, *9*, 805. https://doi.org/10.3389/fphys.2018.00805

Tarashansky, A. J., Xue, Y., Li, P., Quake, S. R., & Wang, B. (2019). Self-assembling manifolds in single-cell RNA sequencing data. *eLife*, *8*, e48994. https://doi.org/10.7554/eLife.48994

Tay, A., & Melosh, N. (2019). Transfection with nanostructure electro-injection is minimally perturbative. *Advanced Therapeutics*, *2*(12), 1900133. https://doi.org/10.1002/adtp.201900133

Thi-Kim Vu, J., Rink, J. C., McKinney, S. A., McClain, M., Lakshmanaperumal, N., Alexander, R., & Sánchez Alvarado, A. (2015). Stem cells and fluid flow drive cyst formation in an invertebrate excretory organ. *eLife*, *4*, e07405. https://doi.org/10.7554/eLife.07405

Vila-Farré, M., & Rink, J. C. (2018). The ecology of freshwater planarians. *Methods in Molecular Biology*, *1774*, 173–205. https://doi.org/10.1007/978-1-4939-7802-1_3

Wagner, D. E., Wang, I. E., & Reddien, P. W. (2011). Clonogenic neoblasts are pluripotent adult stem cells that underlie planarian regeneration. *Science*, *332*(6031), 811–816. https://doi.org/10.1126/science.1203983

Wang, Y., Zayas, R. M., Guo, T., & Newmark, P. A. (2007). *nanos* function is essential for development and regeneration of planarian germ cells. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(14), 5901–5906. https://doi.org/10.1073/pnas.0609708104

Xie, X., Xu, A. M., Leal-Ortiz, S., Cao, Y., Garner, C. C., & Melosh, N. A. (2013). Nanostraw-electroporation system for highly efficient intracellular delivery and transfection. *ACS Nano*, *7*(5), 4351–4358. https://doi.org/10.1021/nn400874a

Zayas, R. M., Bold, T. D., & Newmark, P. A. (2005). Spliced-leader trans-splicing in freshwater planarians. *Molecular Biology and Evolution*, *22*(10), 2048–2054. https://doi.org/10.1093/molbev/msi200

Zeng, A., Li, H., Guo, L., Gao, X., McKinney, S., Wang, Y., Yu, Z., Park, J., Semerad, C., Ross, E., Cheng, L. C., Davies, E., Lei, K., Wang, W., Perera, A., Hall, K., Peak, A., Box, A., & Sánchez Alvarado, A. (2018). Prospectively isolated tetraspanin[+] neoblasts are adult

732      pluripotent stem cells underlying planaria regeneration. *Cell*, *173*(7), 1593–1608.

733      https://doi.org/10.1016/j.cell.2018.05.006

734  **Main Figures**

736  **Figure 1**

**Figure 1: Nanoluciferase mRNA delivered through nanostraws is expressed in planarian cells.** (**A**) Schematics showing the steps of nanostraw electro-delivery. In each experiment, 200,000 cells are placed into a nanostraw cartridge. (**B**) SEM images of nanostraws. Inset: a magnified view showing individual straws. (**C**) Epi-fluorescence images showing planarian cells after nanostraw delivery. Insets: magnified views showing cells with nuclear autofluorescence. Arrows: example red autofluorescent cells. (**D**) Histogram of red fluorescence measured by flow cytometry. Both samples show nearly identical distributions. Statistical insignificance ($p = 0.49$, t-test) was established by bootstrapping, which calculates the p-value for 10,000 random subsamples of 100 cells between both distributions. (**E**) Unlike autofluorescence, autoluminescence is absent in planarian tissues, making luminescent reporters easy to detect. (**F**) Luminescence measured from 200,000 cells on a plate reader at 24 hr post nanostraw delivery with sNluc1 mRNA via the addition of furimazine, the substrate utilized by Nluc. Cells were subjected to 45 s of a square wave pulse train of 35 V, 200 μs, and 40 Hz applied 3 times with 1 min between each cycle to inject the cargo. Each data point represents one biological replicate which was conducted on an independently isolated population of cells using nanostraws from the same batch and mRNA synthesized in the same reaction. Negative control: nanostraw delivery with PBS alone. Diffusion: mock experiments without applying electrical pulses. Error bars represent the standard deviation and p-values were calculated using two-sided Welch's t-test.

**Figure 2**

**Figure 2: A screen identified chemical reagents to efficiently transfect planarian cells.** (**A**) Luminescence from 200,000 cells transfected with 1 µg of YB1-sNluc1 mRNA across a range of chemical transfection reagents, using manufacturer's recommended protocols (**Supplementary Table 2**). p-values were calculated by comparing experimental groups with the negative control in which mRNA was delivered with no transfection reagent. (**B**) Luminescence images of planarian cells transfected with YB1-sNluc1 mRNA using Trans-IT mRNA. Cells were imaged on an Olympus LV200 luminescence microscope with either a 20× air objective (Olympus: UPLXAPO20X) for a large field of view or a 100× oil immersion objective (Olympus: UPLXAPO100XO) to observe subcellular localization. Cells were stained with 4 ng/µL Calcein AM (green) with luminescence shown in magenta. Optimization of (**C**) Viromer mRNA and (**D**) Trans-IT mRNA transfections identified conditions with increased expression that are used for all following experiments (asterisk), compared to the condition initially used in the screen (arrowheads). 200,000 cells were transfected in each experiment. (**E**) Viromer mRNA and (**F**) Trans-IT transfected cells showing gradual increase of luminescence over time. Luminescence was measured on 200,000 cells per replicate at each time point and normalized against the maximum luminescence across replicates. The dashed line marks the mean intensity of the negative controls (n = 3) using only furimazine substrate and mRNA in the absence of any transfection reagent. Shaded regions: standard deviation. (**G**) Luminescence from 40,000 sorted neoblast enriched cells or bulk sorted cells transfected with YB1-sNluc1 mRNA. In all experiments, cells were transfected with 1 µg of YB1-sNluc1 mRNA unless otherwise specified (**Supplementary Table 3**), assayed for luminescence at 24 hpt. Data point in panels **A** and **G** represents the mean luminescence of three parallel technical replicates, and the experiments were conducted on three biological replicates each using independently dissociated cells. Data points in panels **C-F** are from technical replicates. p-values were calculated using two-sided Welch's t-test.

**Figure 3**

**Figure 3: Expression of exogenously delivered mRNA is regulated post-transcriptionally.**

(**A**) Luminescence from transfections using a range of constructs incorporating different endogenous 5' and 3' UTRs flanking sNluc1. No-UTR mRNA contains flanking attB1 and attP1 sequences used for cloning. UTRs were cloned from ENO, enolase (dd_Smed_v6_510_0_1); RPL10, ribosomal protein L10 (dd_Smed_v6_130_0_1); YB1, YB1 cold shock protein (dd_Smed_v6_52_0_1); RPL15, ribosomal protein L15 (dd_Smed_v6_193_0_1), which were identified using the gene annotation provided by PlanMine (Rozanski et al., 2019). (**B**) Luminescence from alternative codon usage variants of Nluc flanked by YB1 UTR. (**C**) Luminescence from transfections using constructs that combine various UTRs with Nluc codon variants. Note the expression showed no significant differences. All transfections were performed across three biological replicates consisting of three technical replicates with 200,000 cells, from whole dissociated planarians, using 1 µg mRNA and 0.8 µL Viromer per well. Each biological replicate was performed with mRNA produced from independent *in vitro* synthesis reactions. Luminescence was assayed 24 hpt. (In **A** and **B**, p-values were calculated using two-sided Welch's t-test between experimental groups and negative controls with cells only; in **C** statistical insignificance was established using Tukey's HSD pairwise t-test.

**Figure 4**

**Figure 4: Heterologous reporter expression *in vivo*.** (**A**) Schematics showing the workflow of *in vivo* transfection experiments. Bulk luminescence was measured from dissociated worms injected with mRNA complexed with (**B**) Viromer mRNA or (**C**) Trans-IT mRNA. Dashed lines: 3 standard deviations above the background (based on the mRNA only condition) to discriminate negative and positive animals. Numbers of positive animals out of all animals injected are reported for each time point. Viromer was prepared with 0.8 µL of Viromer, 1 µg of RPL15-sNluc2 mRNA, and Viromer Buffer up to a final reaction volume of 25 µL. Trans-IT was prepared with 2 µL of Trans-IT, 1 µL of Boost, 1.5 µg of RPL15-sNluc2 mRNA, in L-15 up to a final volume of 25 µL. 30 worms, from two independent experiments, were injected along the tail midline with 0.9 µL of transfection mix per time point per transfection reagent. (**D**) Live planarians imaged on the Olympus LV200 luminescence microscope through a 20× air objective (NA 0.8, WD 0.6 mm, Olympus: UPLXAPO20X). 27 worms were injected per transfection reagent while negative control worms were injected with mRNA alone. Images were acquired at 12 hpt. Worms were anesthetized and incubated for 15 min in furimazine (NanoGlo-Live, Promega) supplemented with 1% DMSO to aid in permeabilization. Luminescence images were acquired using an exposure of 1 min. Brightfield images were acquired using an exposure of 10 ms. All experiments were performed using identical exposure time and digital gain. Dashed lines: worm tail boundary; arrows: the site of the injection. n: number of injections that produced visible, localized luminescence signal out of total injected animals.

 **Supplementary figures**

**Figure 1 – figure supplement 1: Nanostraw electro-delivery device.**

**Figure 1 – figure supplement 1: Nanostraw electro-delivery device.** (A) A photograph of the

nanostraw device. In frame is the power supply (left), the oscilloscope, which is used to monitor

the square wave pulses (top right), and the nanostraw electroporator device (center bottom)

showing the dials to set the pulse width, duration, and voltage. (B) An interior view of the

nanostraw device, showing the electrical contacts which interface with the nanostraw electro-

delivery cartridge and carry the electrical pulses to the titanium electrodes on the top and bottom

of the cartridge. (C) Individual components of the nanostraw cartridge, showing the cathode,

which dips into the cell culture medium from above, the anode, which sits just below the buffer

and mRNA to be delivered, and the nanostraw membrane, which is inserted atop the buffer and

anode. (D) A demonstration of a nanostraw membrane placed within the cartridge.

**Figure 2 – Figure supplement 1: CRNeoblast sorting and validation.**

**Figure 2 – figure supplement 1: CRNeoblast sorting and validation.** (A) FACS gates to sort

for CRNeoblasts and their susceptibility to irradiation. The cells are first gated by forward and

side-scatter (FSC-A, SSC-A) to remove larger somatic cells. Small cells are then gated for high

CellRox Green (it binds to DNA upon oxidation) and low LysoTracker Deep Red (it binds to

intracellular membranes) fluorescence intensity. This strategy is designed to select for cells with

high DNA content (i.e., cells in S or $G_2$/M phase) and low organelle complexity. Unlike other

DNA binding dyes, we found that CellRox Green has a very low toxicity level. CRNeoblasts

make up approximately 5% of the total events detected. Upon x-ray irradiation (20 Gy), this

population significantly diminishes in size and is nearly completely ablated by two days post

irradiation (dpi). (B) Sorted CRNeoblasts have uniform morphology. White arrowhead: an

example CRNeoblast in the process of dividing. Yellow arrowheads: processes that are

characteristic of planarian neoblasts. (C) Single cell RNA-sequencing analysis of CRNeoblasts

show that ~94% were positive for the neoblast marker *piwi-1*. Of these *piwi-1*⁺ cells, they were

split between neoblast subpopulations including epidermal, muscle, and gut progenitors

(N_epidermal, N_muscle, and N_gut), and clonogenic neoblasts (cNeoblasts, $tgs^+$).

**Figure 3 – Figure supplement 1: Cloning methodology.**

**Figure 3 – figure supplement 1: Cloning methodology.** (A) A diagram showing the steps of

generating pNHT7, the base vector used to clone endogenous planarian genes. The overall

design is a LacZ cassette, derived from pUC19, flanked by BsaI restriction sites, which allow for

subcloning of planarian genes into pNHT7. Upstream is a T7 promoter, which allows for *in vitro*

transcription (IVT), and all of these are flanked by M13 forward and reverse primers which are

used to generate linearized templates for IVT through PCR. The backbone for pNHT7 is derived

from the backbone from pDONOR221. (B) Isolating the 5' and 3' UTRs of a gene of interest

(GOI) from cDNA uses primers containing pNHT7 compatible BsaI restriction sites followed by

5' and 3' UTR sequences of the GOI. After amplification, the fragment containing the GOI can

be cloned into pNHT7 though a standard golden gate reaction. (C) The Nluc reporter is inserted

between the UTR sequences by using primers containing BsaI restriction sites to prime the 5'

and 3' ends of the reporter as well as outward facing primers which add complementary BsaI

restriction sites and prime the end of the 5' UTR and beginning of the 3' UTR for the gene

captured in pNHT7. These two amplicons can then be purified and assembled via a golden gate

reaction. All primer sequences are provided in **Supplementary Table 4**.

**Figure 3 – figure supplement 2: Expected *in vitro* transcription results.**

**Figure 3 – figure supplement 2: Expected *in vitro* transcription result**s. (A) Diagram of IVT.

(B) EtBr stained 1% agarose gel showing the linear amplification products. (C) Gel images

showing the mRNA product before and after polyadenylation. Upon successful polyadenylation,

the mRNA product should be shifted in size. Samples should be free of excessive degradation

products, which we found critical for successful transfections. The ladder used is the NEB 1 kb

DNA ladder.

**Figure 3 – figure supplement 3: Assessment of m1Ψ on mRNA expression in planarians**

**Figure 3 – figure supplement 3: Assessment of m1Ψ on mRNA expression in planarians.**

Recent reports suggest that the base analog 1-methyl pseudouridine (m1Ψ) may increase

translation, improve mRNA stability, and decrease immunogenicity (Andries et al., 2015; Svitkin

et al., 2017). This modification is also currently being used in recently developed mRNA

vaccines against SARS-CoV-2. We compared transfections with RPL15-sNluc2 mRNA

containing either rUTP or m1Ψ (chemical structures shown on the left). We found both to be

active, with no significant difference between the two samples, suggesting that the observed

benefits of m1Ψ modified mRNA in mammalian systems may not be directly transferrable to

more basal invertebrates like planarians. Arrow: the uracil group is flipped 180 degrees and

attached by a carbon-carbon bond rather than the usual carbon-nitrogen bond in pseudouridine.

Arrowhead: m1Ψ contains an additional methyl group. All transfections were conducted with 0.8

μL of Viromer mRNA and 1 μg of mRNA. 200,000 cells were incubated at 20°C, and

luminescence was measured at 24 hpt. Data were collected from a single batch of cells

transfected from three independent mRNA synthesis reactions across three technical replicates

per batch of mRNA. We anticipate that other chemical modifications may be tested similarly in

the future to identify ones which regulate gene expression in planarians.

**Supplementary table legends**

**Supplementary Table 1: Nluc variants used in this study.** Coding sequences with different codon usage bias are listed with their respective GC content, and codon adaptation index (CAI) calculated from http://genomes.urv.es/CAIcal/E-CAI/.

**Supplementary Table 2: Conditions for screening transfection reagents.** The conditions followed the manufacturer's protocol. Reagent 1 is the main transfection reagent provided with the kit (e.g., Lipofectamine 3000, Trans-IT mRNA, Viromer mRNA). Reagent 2 is a reagent provided by some but not all kits. For Trans-IT this is the Boost reagent, and for Lipofectamine 3000, this is the Transfection Enhancer reagent. Diluting Buffer is the solution used to dilute the mRNA and transfection reagents. Some kits provide a specific buffer to use (Viromer) while others require serum-free medium, which is L-15 in our experiments. Incubation time was done in room temperature and is recommended to allow the reagent and mRNA to complex before adding to cells.

**Supplementary Table 3: List of constructs used in this study.** Corresponding plasmid maps are provided in GenBank (.gb) format as **Supplementary File 1**. All constructs will be made available through Addgene.

**Supplementary Table 4: Primers used in this study.** Listed are primer ID, conventional name, a short description of the usage, the sequence from 5' to 3', and the predicted melting temperature of the primer.