Forecasting the effects of water regulation on the population viability of a threatened amphibian

Rupert Mathwin[1], Skye Wassens[2], Matthew S. Gibbs[3,4], Jeanne Young[5], Qifeng Ye[5,6], Frédérik Saltré[1,7], Corey J.A. Bradshaw[1,7]

1 Global Ecology *Partuyarta Ngadluku Wardli Kuu*, College of Science and Engineering, Flinders University, PO Box 2100, Tarndanya (Adelaide), South Australia 5001

2 School of Environmental Sciences, Charles Sturt University, PO Box 789, Albury, New South Wales 2640

3 School of Civil, Environmental and Mining Engineering, The University of Adelaide, Adelaide, South Australia 5005

4 Department for Environment and Water, GPO Box 2384, Adelaide, 5001 South Australia 5000

5 College of Science and Engineering, Flinders University, PO Box 2100, Adelaide, South Australia 5001, Australia

6 South Australian Research and Development Institute, 2 Hamra Ave, West Beach, South Australia 5024

7 ARC Centre of Excellence for Australian Biodiversity and Heritage, EpicAustralia.org.au

Corresponding Author: Rupert Mathwin

rupert.mathwin.ecology@gmail.com

Global Ecology, College of Science and Engineering, Flinders University, PO Box 2100, Adelaide, South Australia 5001

Author Contributions: RM and MSG completed the hydrological modelling, RM and CJB designed and constructed the population models. All authors contributed to the manuscript and provided editorial advice.

33 **Forecasting the effects of water regulation on the population**
34 **viability of a threatened amphibian**
35

Abstract

The regulation of river systems alters hydrodynamics and often reduces lateral connectivity

between river channels and floodplains. For taxa such as frogs that rely on floodplain

wetlands to complete their lifecycle, decreasing inundation frequency can reduce recruitment

and increase the probability of local extinction. We virtually reconstructed the inundation

patterns of wetlands under natural and regulated flow conditions and built stochastic

population models to quantify the probability of local extinction under different inundation

scenarios. Specifically, we explored the interplay of habitat size, inundation frequency, and

successive dry years on the local extinction probability of the threatened southern bell frog

Litoria raniformis in the Murray River floodplains of South Australia. We hypothesised that

the changes to wetland inundation resulting from river regulation are a principal driver of *L.*

raniformis declines in this semi-arid system.

Regulation has reduced the inundation frequency of essential habitats below critical

thresholds for the persistence of many fresh water-dependent species. Successive dry years

raise the probability of local extinction, and these effects are strongest in smaller wetlands.

Larger wetlands and those with more frequent average inundation are less susceptible to these

effects.

Elucidating these trends informs the prioritisation of treatment sites and the frequency of

conservation interventions. Environmental water provision (through pumping or the operation

of flow-regulating structures) is a promising tool to reduce the probability of breeding failure

and local extinction. Our modelling approach can be used to prioritise the delivery of

environmental water to *L. raniformis* and potentially many other frog species.

Keywords

environmental water provision, frog, *Litoria raniformis*, river regulation, stochastic

population model

Introduction

The global decline of amphibians (Blaustein and Wake 1990; Stuart et al. 2004) is stark, with the most recent figures showing 41% of assessed species are threatened with extinction (IUCN 2020). Human modifications of wetland networks on which many amphibian populations depend via river regulation can reduce lateral connectivity and the ecological function of wetland habitats (Castello and Macedo 2016). Being essential breeding and nursery areas for many amphibian species, such regulation can change amphibian community structure and drive local extinctions (Wassens and Maher 2011). Furthermore, amphibian extinctions and population declines are projected to increase through the 21st Century as the interactive perturbations increase in intensity (Hof et al. 2011); for example, the negative consequences of river regulation on amphibians are predicted to increase as climate change progressively overtakes land use (Narins and Meenderink 2014) as the main driver of species richness patterns and extinctions (Newbold 2018). Climate change-driven aridification reduces the availability of breeding habitats, resulting in lower species richness (McMenamin et al. 2008). The combination of aridification and increasing water consumption is therefore tlikely to reduce the availability of freshwater habitats further in some regions (Miller et al. 2018).

Manipulating water resources can help alleviate the effects of declining water availability and support amphibian recruitment (Shoo et al. 2011; Smith et al. 2019; Mathwin et al. 2021). Techniques for manipulating water to benefit amphibians vary in approach and success (Mathwin et al. 2021), but the best-supported intervention is the provision of water to breeding habitats to match the larval requirements of the target species ('environmental water provision'). This is because enhancing survival through breeding and early life stages can stabilise populations (Griffiths and Pavajeau 2008). The targeted delivery of environmental water is a common practice today (Kennen et al. 2018) and could become necessary to

conserve some species (Greenwood et al. 2016), especially those with limited phenotypic

plasticity or those near the edge of their ecological niche (Grant et al. 2020).

Australia's Murray-Darling Basin provides a model system to examine this process. The

catchment is heavily regulated and each year up to 61% of total flow is extracted for

consumptive use (CSIRO 2008). This has reduced the number and function of wetlands (Gell

and Reid 2014), resulting in the decline and fragmentation of water-dependent taxa

(including the southern bell frog, *Litoria raniformis*) (Clemann and Gillespie 2012). In

response to systemic environmental degradation, federal legislation mandates interventions

aimed at restoring ecological function (Docker and Robinson 2014), including environmental

water provision for species recovery.

In this paper we explore hydro-ecological thresholds and generate guidelines for

environmental water provision to benefit amphibians by constructing stochastic, hydro-

ecological population models. We hypothesise that the reduced frequency of wetland

inundation resulting from river regulation is driving local extinction events of *L. raniformis* in

this system. We hypothesise that larger wetlands and wetlands with greater average

inundation are less susceptible to local extinction during successive dry years. We posit that

the probability of local extinction can be reduced by environmental water provision and that

our modelling approach can be used to prioritise the location and frequency of environmental

water provision to at-risk amphibian populations.

Materials and Methods

Study area

The Murray-Darling Basin contains Australia's longest river system and is heavily regulated

to provide water for domestic and agricultural use. Regulating structures include a series of

14 main-channel weirs ('locks') that dissect the river. Our focus is 70 km of the Murray River

channel between Lock 3 (34° 11′ 16.95″ S, 140° 21′ 29.65″ E) and Lock 2 (34° 4′ 39.31″ S,

139° 55′ 52.81″ E) and the associated wetlands and floodplains along this reach (**Figure 1**).

This region receives an annual rainfall of 160–240 mm (semi-arid/arid), which is insufficient

to fill off-channel wetlands most years. As such, all naturally occurring *L. raniformis*

breeding in this reach results from elevated river level or the provision of environmental

water. Water is typically delivered using portable pumps that exclude most fish and aquatic

invertebrate predators.

Life history of southern bell frogs

The life history of *L. raniformis* is well described, with estimates for most vital rates

(survival, growth, reproduction, and recruitment). Eggs hatch two to four days after laying

(Anstis 2002). The likelihood of hatch (as a proxy for egg survival rate) is between 0.933 and

1.000/egg. This species ranges from north of latitude 35 °S to south of 45 °S, which spans a

range of thermal conditions (from 32.9 °C maximum daily summer temperatures in the north

of their range to 20.5 °C in the south). Being ectotherms, larval duration is strongly driven by

temperature and as such, larval duration varies from 10–12 weeks in the north of their range

to 12–15 months in the south (Anstis 2002). We used the larval duration of 70–80 days

calculated at a constant water temperature of 23 °C (Cree 1984), approximately the average

daily summer temperatures experienced in the study reach (which averages diurnal highs and

nocturnal lows). We used the estimates of survival to metamorphosis generated for *Crinia*

signifera (15–26% and 7–56%) (Williamson and Bull 1999), in the absence of species-

specific estimates, these are the most-relevant estimates available.

Both sexes reach maturity in their first year (Heard et al. 2012), and females breed

annually (Anstis 2017) and lay between 1885 and 4563 eggs each season (Humphries 1979,

Germano and White 2008). For adult life stages we used annual adult survival probability

from the closely related *Litoria aurea* (mean = 0.2172, standard deviation = 0.087) (Pickett et

al. 2016). These species are similar in size, appearance, and behaviour, although *L. aurea*

occurs along a more northerly latitude than *L. raniformis*. Using lines of arrested growth in

the shaft of the medial phalanx to determine age (skeletochronology), *L. raniformis* can

survive into its fifth year (Mann et al. 2010, G. Heard and A. Turner, Charles Sturt

University, pers comm.).

Demographic model

Within an R programming environment (R Version 4.0.2: Taking Off Again, R Project Team

2020), we constructed a stochastic population model based on a lifecycle graph (**Figure 2**)

and an age-classified (Leslie) population model (Leslie 1945) that considers only females

(**Figure S1**), where *f* represents fertility and *S* represents annual survival. Year 0 individuals

do not breed and are not assigned a fertility. We randomly resampled fertilities (*f*) for age

classes one to five years from a Normal distribution with a mean = 3245 and a standard

deviation = 897 (calculated from Humphries 1979 and Germano and White 2008), which we

halved to reflect the 50:50 sex ratio in this species.

We calculated survival *S* to the end of the first year as:

$$S_0 = S_e * S_t * ((365 - (D_e + D_t)) * S_{ad}) \qquad \text{[equation 2]}$$

156   where S_0 is the probability of survival from 0 to 1 year old. S_e is the probability of hatch

157   (randomly resampled from a uniform distribution from 0.933–1.000). S_t is the probability of

158   survival to metamorphosis, which we randomly resampled from a β distribution with a mean

159   = 0.26 and a standard deviation = 0.12 (Williamson and Bull 1999). We calculated the shape

160   parameters α and β of this distribution as:

$$\alpha = m\frac{\mu(1-\mu)}{\sigma^2} \qquad \text{[equation 3]}$$

$$\beta = (1-\mu)\left(\frac{m\mu(1-m\mu)}{\sigma^2} - 1\right) \qquad \text{[equation 4]}$$

where μ is the mean and σ is the standard deviation of the β distribution. D_e is the duration of

the egg stage (in days) resampled randomly from a uniform distribution between 2 and 4, and

D_t is the duration of the tadpole stage (in days) resampled randomly from a Normal

distribution between 70 and 80 days. S_{ad} is the daily adult survival probability calculated as:

$$S_{ad} = \sqrt[365]{S_a}$$ [equation 5]

where S_a is the annual probability of survival of an adult frog sampled from a β distribution

with a mean = 0.2172 and a standard deviation = 0.087 (Pickett et al. 2016). We assigned

frogs in their fifth year a survival = 0, reflecting senescence and death during their fifth year

(although the model permits breeding before death).

Population sizes

We modelled *L. raniformis* populations at each of four wetland sizes (*small*, *medium*, *large*,

very large). We derived these categories from typical wetlands in the region and the

corresponding population size is an estimate of their respective carrying capacity using the

maximum populations observed during monitoring. *Small* populations (~ 40 individuals)

reflect the carrying capacity and population dynamics present in a wetland pool several

metres in diameter. A *medium* population (~ 130 individuals) represents a wetland pool with

surface area similar to 1–2 domestic swimming pools. A *large* population (~ 300 individuals)

represents a wetland pool surface area similar to an Olympic swimming pool, and a *very*

large population (~ 1000 individuals) reflects a wetland several hundred metres in diameter.

The starting population of adult females was calculated as 50% of these values (as *L.*

raniformis has a 50:50 sex ratio), being 20, 65, 150 and 500 female frogs, respectively. We

assigned age classes for the starting female population by randomly resampling five adult

survival values and then dividing the total number of females among the five age classes in

these proportions. This created a more homogenous initial age structure than we might expect

in a wild population. We managed this by ignoring the first 10 generations of each run as a

'burn-in' period, which allowed the model to stabilise to a stochastic expression of the stable-

age distribution before analysis.

We allocated the initial number of eggs using an *a priori* number of spawning masses for

each wetland size ($n = 10, 30, 85, 150$, respectively). We then stochastically resampled the

number of eggs in each spawning mass (halved because the model only considers females).

The *a priori* assignment of spawning masses did not impact population dynamics after burn-

in.

Compensatory density feedback

To stabilise long-term population growth, we incorporated three compensatory density-

feedback relationships. These relationships are not available for *L. raniformis* so we

estimated these functions following the form demonstrated in other anurans. First, we

corrected rates of egg laying to reflect the maximum carrying capacity (K_{egg}) at each wetland.

As the number of eggs approaches the wetland's carrying capacity, eggs are reduced

following an exponential decay function of the form:

$$y = 1.01 - (x^{2+\frac{9}{x}})$$ [equation 6]

and the rate of egg reduction follows the equation:

$$R_{egg} = \int_0^{0.99} 1.01 - \left(K^{2+\frac{9}{K}}\right) d_K$$ [equation 7]

where R_{egg} is the total reduction in eggs laid at the wetland, K is the carrying capacity and d_K

is the differential K (carrying capacity) (**Figure S2**). Carrying capacity is calculated for each

wetland size category using the initial number of *a priori* spawning masses laid (10, 30, 85 or

150) at the maximum fecundity for the species (4563 eggs), which we halved to consider only

females. In this way, frogs lay without inhibition until the total egg count approaches the

carrying capacity. The highest inhibition rate corresponds to 0.99 of the wetland's carrying

capacity and all subsequent egg-laying events are reduced at this value (corresponding to a

reduction of 0.0078).

The second and third compensatory density-feedback functions reduce survival during

terrestrial life stages with increasing density (e.g., Harper and Semlitsch 2007, Berven, 2009).

The second feedback used the total number of tadpoles present to reduce survival

probabilities during the first year of life. This reflects predation and competition during larval

life stages, and in the first few months post-metamorphosis. We calculated density feedback

on survival probability using the function:

$$R_{0-1} = (-0.241 + 1.54K))^{-0.855}$$ [equation 8]

where R_{0-1} is the reduction factor in survival for age class 0 individuals (**Figure S3**). Here,

we assigned carrying capacity *a priori* based on the size of the wetland, *small* = 200 females,

medium = 600 females, *large* = 1800 females and *very large* = 3000 females. Density

feedback on survival from 0- to 1-year age classes is applied when the total adult population

> 0.3 of the wetland's carrying capacity. We applied the highest reduction in survival at 2.1

times carrying capacity that reduces survival probability to 0.0101 (Figure S3).

The third compensatory density-feedback function applies the abundance of all adult frogs

to reduce survival from the 1- to 2-year age class (R_{1-2}) to reflect competition for resources.

The form of the relationship followed the method applied for (R_{0-1}) presented above (equation

8, **Figure S3**).

Modelling hydrology

We considered two river-flow scenarios. The 'regulated flow' scenario is informed by the

historical flow record immediately downstream of Lock 3 (A4260517;

waterconnect.sa.gov.au). We used data starting in 1926, the year following the completion of

Lock 3. River levels above the operative range of this gauge 'drown out' the gauge, making

measurements inaccurate. During these periods, we used data from station A4260528 (6.5 km

downstream of Lock 3), which is unaffected by elevated river levels. This created a

continuous daily record of flow rate between Locks 3 and 2 for 83 years. The second river-

flow scenario is modelled 'natural flow', which covers the same time period, but in the

absence of extraction or regulation in the catchment (see Murray-Darling Basin Authority-

MDBA 2012).

A second-order polynomial derives river heights from mean daily flow rate (Ml day^{-1}) for

the two flow scenarios. Based on mean daily river height (metres with respect to Australian

River Height Datum, mAHD) at site A4260517 and the flow records above, the following

rating-curve equation:

$$H = -2f^2 10^{-10} + 8f 10^{-5} + 6.0144 \qquad \text{[equation 9]}$$

estimates river height, where H is river height (mAHD) and f is river flow (Ml day^{-1}).

Equation 9 represents an empirical 'inverse rating curve' for the site, where a rating curve is a

commonly used approach to derive streamflow based on a recorded river height. We used this

to create a continuous daily river height for this reach over 83 years under natural and

regulated flow.

Our model exposes wetland populations to one of two states each year. A 'wet' year is

when the wetland received sufficient water to support frog reproduction and recruitment;

conversely, during a 'dry' year, the wetland did not receive sufficient inputs to support frog

recruitment (often filling and then drying prematurely — see definition below). There is a

paucity of accurate sill-height (the river height at which a wetland begins to fill) data in the

reach, so rather than modelling the inundation of specific wetlands, we calculated inundation

of nine possible sill heights (7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5 and 11 mAHD). Following

discussion with local wetland managers, we used the rule that if mean daily river height is ≥

10 cm above sill height for ≥ 10 days during winter and spring, then the wetland is 'wet' and

can support frog recruitment that year (K. Mason – Department for Environment and Water,

Adelaide, pers comm.). It is not necessary that these ten days be consecutive to fulfil this

criterion. By using this criterion, we determined which of the nine sill heights were wet, and

which were dry for each of the 83 years under the natural and regulated scenarios.

Modelling flow scenarios

We modelled 18 hydrological scenarios — these being sill heights of 7, 7.5, 8, 8.5, 9, 9.5, 10,

10.5 and 11 mAHD — each modelled under both natural- and regulated-flow conditions. To

create stochastic expressions of these scenarios, we used the 83 years of daily river-height

data for the flow scenario (natural or regulated flow) and classified each of those 83 years as

either filling the nominated sill height (≥ 10 non-consecutive days exceeding the sill height

by ≥ 10 cm during the previous winter or spring) to create a wet year, or failing to fill the sill

height sufficiently and creating a dry year (when recruitment is unsuccessful). Using this

sequence of 83 wet and dry years (for each specific sill height and flow combination) we

created a discrete-time Markov chain (e.g., Supporting Information 1) to resample unique,

stochastic, 85-year wet/dry sequences for each model scenario run.

Finally, we modelled each of the four wetland sizes (*small*, *medium*, *large*, *very large*) at

each of nine sill heights (7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5 and 11 mAHD) and each of two flow

scenarios (natural and regulated flow). We ran each of these 72 wetland models stochastically

10000 times, each for 85 consecutive generations or until the population went extinct

(creating a combined total of 61.2 million stochastic generations). This was sufficient to

account for wetland variability in the study reach. We disregarded the first ten burn-in

generations of each model and used generations 11 to 85 (or to extinction) in our analyses.

For each model, we recorded the sequence of wet and dry years from generations 11 to 85

(or extinction). For each occurrence of two or more consecutive dry years, we recorded the

number of consecutive dry years and whether it resulted in local extinction. We organised

these data by the average frequency of dry years at the wetland and calculated the probability

of extinction for two to five consecutive dry years, noting that six consecutive dry years

exceeds the maximum reproductive lifespan of the species.

To quantify the relative influence of each variable on the likelihood of local extinction we

did a global sensitivity analysis on the model (Prowse *et al*, 2016). For this we modelled a

large wetland with a 7.5 m sill height under regulated flow conditions and used Latin

hypercube resampling to rerun 10000 discrete model iterations within the nine-dimensional

parameter space (using the `lhs` library in R). The ranges of uncertainty used were: clutch

size (1000 – 6000), duration of egg phase (0.5 – 14 days), hatch probability (0.5 – 0.98),

duration of tadpole phase (50 – 90 days), mean tadpole survival (0.03 – 0.6), mean annual

adult survival (0.03 – 0.35), maximum age (5 – 10 years), tadpole carrying capacity (500 –

5000 females) and juvenile carrying capacity (200 – 5000 females). We then used a boosted

regression tree emulator (implemented using the `dismo` library in R), setting the error

distribution family as Bernoulli, the bag fraction to 0.75, learning rate to 0.001, tolerance to

0.0001 and tree complexity to 2.

To examine the effect of maximum lifespan on the predictions, we also implemented a

univariate sensitivity analysis on a single, large wetland with a sill height of 7.5 m under

regulated flow conditions. We reran this model with senescence constrained to ages two,

three, four, and five years (Appendix S1).

Results

 Results

'Wetness' of wetlands under natural- and regulated-flow conditions

River regulation reduced the frequency of wet years by 19–34% compared to the natural-flow

scenario for the same 83-year period (**Figure 3**). This equates to a reduction of wet years by

19.5% at 7 mAHD, up to 60% at 11 mAHD. This changed both the proportions and pattern of

wet and dry years. Regulation increased the mean duration of successive dry years at all sill

heights (**Figure 4**). Under natural conditions the mean duration of dry years did not exceed

two years at any of the sill heights examined, whereas the regulated-flow scenario resulted in

mean duration of dry events approaching the maximum lifespan of the species at sill heights

≥ 10 mAHD. Under natural conditions the maximum duration of dry events did not exceed

the maximum reproductive lifespan of the species, except at a sill heights > 10 mAHD. The

maximum duration of dry events observed under river regulation exceeded the maximum

reproductive lifespan of this species at every sill height examined.

Extinction probability of each wetland scenario

Under the natural-flow scenario, the probability of extinction — Pr(Ext) — during the 85

modelled years was ~ 0 at sill heights ≤ 8.5 mAHD (**Figure 5**). Conversely, all wetland sizes

reached Pr(Ext) = 1 at sill heights ≥ 9 mAHD when flows were regulated. Thus, wetlands that

historically supported *L. raniformis* populations under natural-flow conditions (i.e., ≤ 8.5

mAHD) are unreliable under river regulation. Without intervention, wetlands with sill height

of ≥ 9 mAHD will not sustain *L. raniformis* populations under existing flow conditions.

The probabilities of extinction in *very large* wetlands were 0.25 to 0.70 lower than those in

small wetlands (**Figure 6**). This effect was more pronounced in the regulated-flow scenario

than under natural-flow conditions.

Two consecutive dry years resulted in Pr(Ext) < 0.25 in all treatments, including wetlands

with an 80% average frequency of dry years (**Figure 6**). Increasing the number of

consecutive dry years increases Pr(Ext) up to five consecutive dry years, which gives Pr(Ext)

> 0.5 in all treatments except for *very large* populations with an average frequency of dry

years < 20%. Smaller wetlands had increased extinction probability in all instances and the

disparity between wetland sizes became more pronounced with increasing successive dry

years. The average frequency of dry years at the wetland strongly influences the capacity to

survive extended dry periods. Wetter sites have lower extinction probability than drier sites

for each drought duration.

Boosted regression tree emulation of the nine-dimensional Latin hypercube space

accounted for 96.7% of the observed variation in the model. Only two parameters strongly

influenced the probability of local extinctions. These were annual mean adult survival

probability, which accounted for 58.3% of the model influence and mean tadpole survival,

which accounted for 21.6% of the observed variation in the model outputs (Figure 7).

Discussion

River regulation

Regulation and abstraction of flow in this catchment has resulted in wetlands that fill less

often (**Figure 3**), and the average duration of successive dry years has increased by up to

three years compared to natural flow conditions (**Figure 4**) (Maheshwari et al. 1995; Bice et

al. 2017). For species that rely on floodplain inundation to complete their lifecycle, dry years

result in reproductive failure. When breeding occurs and the site dries before completion of

larval life stages, this creates population sinks that are not uncommon in amphibians breeding

in ephemeral waterbodies. In isolation, reproductive failure causes a population fluctuation,

but does not always appreciably increase the probability of local extinction (Taylor et al.

2006), albeit with simplification of age structure and attrition of adult populations. However,

successive failures increase extinction risk, especially in short-lived species (Semlitsch et al.

1996). Without the capacity to extend lifespan through unfavourable periods (i.e.,

aestivation), droughts equalling or exceeding the reproductive lifespan of a species result in

local extinction, a process that is likely to have driven the local extinctions of 42% of

breeding sites of *Pseudophryne pengilleyi* in Kosciuszko and Brindabella National Parks in

New South Wales, Australia (Scheele et al. 2012).

Drought and the decline of Litoria raniformis

The flow record that informs our model includes a severe drought from 1996 to 2009. During

this time, south-eastern Australia experienced a region-wide reduction in rainfall and runoff,

below-average streamflow, and critical water shortages (Heberger 2011; van Dijk et al.

2013). This event, coupled with ongoing water extraction, resulted in seven or more

successive dry years at all wetlands in this reach (**Figure 4**). Modelled natural flow for the

same period indicates that the maximum number of successive dry years would not have

exceeded four years in wetlands up to 10 mAHD (**Figure 4**). Under natural flow conditions,

all wetlands < 10 mAHD could have supported *L. raniformis* through this perturbation. The

persistence of some few populations through the drought is, in part, due to targeted

environmental water provision during this time.

Anthropogenic shifts in the atmosphere have increased the frequency, intensity, and

duration of droughts across most of the globe and these trends are likely to continue (Chiang

et al. 2021). Droughts exacerbate the ecological impacts of river regulation and scientifically

defensible approaches to the manipulation of aquatic resources (like those presented here)

will become increasingly necessary to conserve some freshwater species, particularly

wetland- and pond-breeding amphibians.

Model assumptions

Our population model relies on two main assumptions: that (*i*) *L. raniformis* breed in their

fifth year before senescence, and (*ii*) there is no population exchange between wetlands.

However, our boosted regression tree analysis (**Figure 7**) and single-parameter perturbation

(sensitivity) analysis of age (**Appendix S1**) show little influence of maximum age on

extinction probability. The strong influence of survival probabilities (**Figure 7**) results in a

demography that is weighted strongly towards younger age classes and this first assumption

is unlikely to skew the results if incorrect. Similarly, the model outputs are unlikely to be

affected by inaccurate calculation of carrying capacities, clutch size, hatch probability, or

duration of egg or tadpole life stages (**Figure 7**).

The second major assumption is that wetlands are isolated and lack immigration or

emigration. As such, modelled local extinctions are an endpoint after which recolonisation

cannot occur. Amphibians are physiologically dependent on moist environments, have

relatively poor dispersal capacity (compared to other tetrapods), and can show strong site

fidelity, and these traits suggest limited capacity for recolonisation (Blaustein et al. 1994).

Despite limited dispersal, amphibians are frequently thought to exist in metapopulations

(Levins 1969; Smith and Green 2005). We propose that *L. raniformis* in this system meets the

four characteristics of a metapopulation (Hanski et al. 1995): (*i*) habitat patches support local

breeding populations, (*ii*) no single population is large enough to ensure long-term survival,

(*iii*) patches are not too isolated to prevent recolonisation, and (*iv*) local dynamics are

sufficiently asynchronous to make simultaneous extinction of all local populations unlikely.

Despite the semi-arid/arid climate, *L. raniformis* can move hundreds of metres in a single

night and can colonise over distances of 500 m (Herbert 2000; Department for Environment

and Heritage, 2004; Wassens et al. 2008). We anticipate that dispersal between favourable

patches (albeit infrequent and increasingly difficult in this system due to fragmentation, river

regulation, and aridification) can result in recolonisation and contributes to the persistence of

this species at the landscape level through the rescue effect (Brown and Kodric-Brown,

1977). We are developing metapopulation models for this reach to further explore these

dynamics.

Our methodology takes a cautionary approach to conservation. We deliberately model

wetlands without the stabilising effects of metapopulation structure. We acknowledge this

approach produces higher extinction probabilities than we might expect from wetlands within

a network, but this also produces cautionary conservation advice that supports wetlands

without reliance on stochastic phenomena. In systems where populations have become

fragmented, successful movement between populations is low, or movements vary broadly

between years, we recommend this approach to prioritise the active management of

threatened amphibians.

Environmental water provision

In highly regulated catchments, restoring large-scale historic inundation patterns is neither

possible nor desirable. However, providing specific reaches and wetlands with environmental

water can recreate critical components of historic flow regimes. The assumption is that

reconstructing specific components of 'natural' flow can maintain key ecosystem processes

through the timely, sequential ecological cues inherent in 'natural' flow (Poff et al. 1997). In

highly modified and non-stationary conditions, a more mechanistic understanding of

environmental water requirements is valuable (Poff 2018). An alternative approach uses

thresholds of inundation frequency, and in turn extinction probability to prioritise the delivery

of a 'designer' flow regime. When well-designed, this approach generates efficiencies in the

volumes of water delivered while still eliciting the desired ecological response (in this case, a

reduction in the probability of extinction for target amphibian populations).

Our results show that water deliveries maintaining no more than one sequential dry year

will reliably support *L. raniformis* (**Figure 6**). For wetlands with low extinction probability,

or when environmental water budgets are limiting, water provision to ensure no more than

two successive dry years will maintain extinction risk < 0.25 in all wetland sizes. Using this

approach, a rotating roster of wetlands could be watered to offset individual wetland risk,

noting that local extinctions will likely rely on dispersal from neighbouring wetlands to re-

establish. The highest priorities for intervention are wetlands that have experienced four or

five years without recruitment and where *L. raniformis* is still present (**Figure 6**). These

populations should be watered during the next breeding season to reduce the probability of

local extinction. These priorities are not intended to be cycled in perpetuity, especially in

situations where extinction risk is high. For example, a wetland with a 50% extinction

probability after five dry years could require several favourable years to recover before it

could be expected to persist through a second five-year drought.

Our model makes three main assumptions regarding wet years that influence the

techniques for environmental water delivery: that 1. wetlands support adult frogs between

breeding seasons, 2. breeding occurs during each wet year, and 3. wet years (and particularly

successive wet years) do not accumulate fish and invertebrate predators. Water delivery

should be tailored to meet these assumptions, including maintaining small pools to allow

adult frogs to rehydrate during dry periods, watering to support components of the vegetation

community that are important for breeding, and periodic drying to reduce predator densities.

This last point is particularly important as many systems do not dry naturally between wet

events and will accumulate predators more readily than this example. When modelling these

systems, an additional relationship is required which incorporates the effects of predator

densities. Caution is also required when designing flows to ensure the delivery method does

not damage existing habitat features, e.g., erosion or scouring of breeding habitats with high

delivery velocities (Kupferberg, 1996).

Conclusions

Our approach informs thresholds for environmental water provision that can be applied where

demographic data are available and where a clear relationship exists between intervention and

recruitment — for example, the delivery of water to extend hydroperiod in *Rana sevosa*

(Seigel et al. 2006) or *P. pengilleyi* (Scheele et al. 2012). This approach is not specific to

amphibians, nor to environmental water provision, and can also be applied where episodic

events are directly linked to reproductive outcomes, such as fire intermittency and

germination in pyrophytic plants, or supraseasonal flooding events in arid-zone seed

germination.

Environmental water provision to amphibian breeding habitats can reduce the probability

of extinction in populations with insufficient inundation frequency. We show that stochastic

population models, typically constructed for population viability analyses, can be used to

determine critical hydrological thresholds, and that these thresholds are sufficiently nuanced

to prioritise intervention locations and guide the frequency of environmental water provision.

Strategic *a priori* planning and efficient water use will become increasingly important

techniques in amphibian conservation, especially in regulated river systems.

Acknowledgements

We thank the Murraylands and Riverland Landscapes Board and the Nature Foundation SA

for their generous support and the Murray-Darling Basin Authority for providing time-series

data of natural flow.

Declarations

Funding per Acknowledgements, total funding value between the two bodies was small (< $2000 AUD). There are no corresponding grant numbers to cite.

Conflicts of interest/Competing interests None of the authors have any conflict of interest with regards to this work.

Ethics approval Not Applicable

Consent to participate Not Applicable

Consent for publication All authors consent to the publication of this manuscript, its supplementary information and the associated code.

Availability of data and material There are no novel datasets associated with this work. All data are cited and referenced herein

Code availability Data and code available from:
https://github.com/RupertLovesEcology/RiverRegulation_Frog_PopModel

Authors' contributions RM and MSG completed the hydrological modelling, RM and CJAB designed and constructed the population models. All authors contributed to the manuscript and provided editorial advice.

500 This article does not contain any studies with human participants or animals performed by any of the

501 authors.

503 *References*

Anstis M (2002) Tadpoles of south-eastern Australia. New Holland.

Anstis M (2017) Tadpoles and frogs of Australia. New Holland Publishers Pty Limited.

Bice CM, Gibbs MS, Kilsby NN, Mallen-Cooper M, Zampatti BP (2017) Putting the "river" back into

the Lower River Murray: quantifying the hydraulic impact of river regulation to guide

ecological restoration. Transactions of the Royal Society of South Australia, **141**, 108-131.

Berven KA (2009) Density dependence in the terrestrial stage of wood frogs: evidence from a 21-year

population study. Copeia, 328-338.

Bishop-Taylor R, Tulbure MG, Broich M (2015) Surface water network structure, landscape

resistance to movement and flooding vital for maintaining ecological connectivity across

Australia's largest river basin. Landscape Ecology, **30**, 2045-2065.

Blaustein AR, Wake DB (1990) Declining amphibian populations: a global phenomenon? Elsevier

Current Trends.

Blaustein AR, Wake DB, Sousa WP (1994) Amphibian declines: judging stability, persistence, and

susceptibility of populations to local and global extinctions. Conservation Biology, **8**, 60-71.

Brown JH, Kodric-Brown A. (1977) Turnover Rates in Insular Biogeography: Effect of Immigration

on Extinction. Ecology, **58**, 445-449.

Castello L, Macedo MN (2016) Large□scale degradation of Amazonian freshwater ecosystems.

Global Change Biology, **22**, 990-1007.

Chiang F, Mazdiyasni O, AghaKouchak A (2021) Evidence of anthropogenic impacts on global

drought frequency, duration, and intensity. Nature communications, **12**, 1-10.

Cree A (1984) Breeding biology, respiration, and larval development of two introduced frogs (*Litoria*

raniformis and *L. ewingi*). New Zealand Journal of Zoology, **11**, 179-187.

CSIRO (2008) Water availability in the Murray-Darling Basin. Summary of a report to the Australian

Government from the CSIRO Murray-Darling Basin Sustainable Yields Project. CSIRO,

Australia.

Docker B, Robinson I (2014) Environmental water management in Australia: experience from the

Murray-Darling Basin. International Journal of Water Resources Development, **30**, 164-177.

Gell P, Reid M (2014) Assessing change in floodplain wetland condition in the Murray-Darling

Basin, Australia. Anthropocene, **8**, 39-45.

Grant EHC, Miller DA, Muths E (2020) A Synthesis of Evidence of Drivers of Amphibian Declines.

Herpetologica.

Greenwood O, Mossman HL, Suggitt AJ, Curtis RJ, Maclean IMD (2016) Using in situ management

to conserve biodiversity under climate change. Journal of Applied Ecology, **53**, 885-894.

Griffiths RA, Pavajeau L (2008) Captive breeding, reintroduction, and the conservation of

amphibians. Conservation Biology, **22**, 852-861.

Hanski I, Pakkala T, Kuussaari M, Lei G (1995) Metapopulation persistence of an endangered

butterfly in a fragmented landscape. Oikos, 21-28.

Harper EB, Semlitsch RD (2007) Density dependence in the terrestrial life history stage of two

anurans. Oecologia **153**, 879-889.

Heard GW, Scroggie MP, Malone BS (2012) The life history and decline of the threatened Australian

frog, Litoria raniformis. Austral Ecology, **37**, 276-284.

Heberger M (2011) Australia's Millennium Drought: Impacts and Responses. The World's Water:

The Biennial Report on Freshwater Resources (ed. P.H. Gleick), pp. 97-125. Island

Press/Center for Resource Economics, Washington, DC.

Herbert T (2000) Frogs of the Loveday Wetland Complex (River Murray, South Australia): Species

Inventory and Observations on Abundance and Habitat Preferences. Final Report, prepared

for the Barmera-Moorook Field and Game Association. Barmera, South Australia.

Department for Environment and Heritage (2004) Lake Littra Draft Management Plan. Department

for Environment and Heritage, Berri, South Australia.

Germano J, and White H (2008) *Litoria raniformis* (Southern Bell Frog): clutch size. Herpetological

Review, **39**, 461-462.

Hof C, Araújo MB, Jetz W, Rahbek C (2011) Additive threats from pathogens, climate and land-use

change for global amphibian diversity. Nature, **480**, 516.

Humphries RB (1979) Dynamics of a breeding frog community. PhD dissertation, Australian National

University, Canberra

IUCN (2020) The IUCN Red List of Threatened Species. Version 2020-3.

https://www.iucnredlist.org. Downloaded on [13 March 2021].

Kennen JG, Stein ED, Webb JA (2018) Evaluating and managing environmental water regimes in a

water☐scarce and uncertain future. Freshwater Biology, **63,** 733-737.

Kupferberg SJ (1996) Hydrologic and geomorphic factors affecting conservation of a river breeding

frog (*Rana boylii*). Ecological applications **6**,1332-1344.

Leslie PH (1945) On the use of matrices in certain population mathematics. Biometrika, **33,** 183-212.

Levins R (1969) Some demographic and genetic consequences of environmental heterogeneity for

biological control. American Entomologist, **15**, 237-240.

Maheshwari BL, Walker KF, McMahon TA (1995) Effects of regulation on the flow regime of the

River Murray, Australia. Regulated Rivers: Research & Management, **10**, 15-38.

Mann RM, Hyne RV, Selvakumaraswamy P, Barbosa SS (2010) Longevity and larval development

among southern bell frogs (*Litoria raniformis*) in the Coleambally Irrigation Area-

implications for conservation of an endangered frog. Wildlife Research, **37**, 447-455.

Mason K (2013) Frog Monitoring in the Coorong, Lower Lakes and Murray Mouth Region 2012/13,

Department of Environment, Water and Natural Resources, Murray Bridge, South Australia.

Mathwin R, Wassens S, Young J, Ye Q, Bradshaw CJ (2021) Manipulating water for amphibian

conservation. Conservation Biology **35**, 24-34.

McMenamin SK, Hadly EA, Wright CK (2008) Climatic change and wetland desiccation cause

amphibian decline in Yellowstone National Park. Proceedings of the National Academy of

Sciences, **105**, 16988-16993.

Miller DAW, Grant EHC, Muths E, Amburgey SM, Adams MJ, Joseph MB, Waddle JH, Johnson PTJ, Ryan ME, Schmidt BR, Calhoun DL, Davis CL, Fisher RN, Green DM, Hossack BR, Rittenhouse TAG, Walls SC, Bailey LL, Cruickshank SS, Fellers GM, Gorman TA, Haas CA, Hughson W, Pilliod DS, Price SJ, Ray AM, Sadinski W, Saenz D, Barichivich WJ, Brand A, Brehme CS, Dagit R, Delaney KS, Glorioso BM, Kats LB, Kleeman PM, Pearl CA, Rochester CJ, Riley SPD, Roth M, Sigafus BH (2018) Quantifying climate sensitivity and climate-driven change in North American amphibian communities. Nature Communications, **9**, 3926.

Murray-Darling Basin Authority (2012) Hydrologic modelling to inform the proposed Basin Plan-methods and results., MDBA publication no: 17/12, Murray-Darling Basin Authority, Canberra.

Narins PM, Meenderink SW (2014) Climate change and frog calls: long-term correlations along a tropical altitudinal gradient. Proceedings of the Royal Society B: Biological Sciences, **281**, 20140401.

Newbold T (2018) Future effects of climate and land-use change on terrestrial vertebrate community diversity under different scenarios. Proceedings of the Royal Society B: Biological Sciences, **285**, 20180792.

Pickett EJ, Stockwell MP, Clulow J, Mahony MJ (2016) Modelling the population viability of a threatened amphibian with a fast life⬜history. Aquatic Conservation: Marine and Freshwater Ecosystems, **26**, 9-19.

Poff NL (2018) Beyond the natural flow regime? Broadening the hydro-ecological foundation to meet environmental flows challenges in a non-stationary world. Freshwater Biology, **63**, 1011-1021.

Poff NL, Allan JD, Bain MB, Karr JR, Prestegaard KL, Richter BD, Sparks RE, Stromberg JC (1997) The Natural Flow Regime. BioScience, **47**, 769-784.

Pyke G. (2002) A review of the biology of the Southern Bell Frog *Litoria raniformis* (Anura:Hylidae). Australian Zoologist, **32**, 32-48.

R Core Team (2020). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

Scheele B, Driscoll D, Fischer J, Hunter D (2012) Decline of an endangered amphibian during an

extreme climatic event. Ecosphere, **3**, 1-15.

Seigel RA, Dinsmore A, Richter SC (2006) Using well water to increase hydroperiod as a

management option for pond-breeding amphibians. Wildlife Society Bulletin (1973-2006),

34, 1022-1027.

Semlitsch RD, Scott D, Pechmann J, Gibbons J (1996) Structure and dynamics of an amphibian

community. Long-term studies of vertebrate communities**,** 217-248.

Shoo LP, Olson DH, McMenamin SK, Murray KA, Van Sluys M, Donnelly MA, Stratford D,

Terhivuo J, Merino-Viteri A, Herbert SM, Bishop PJ, Corn PS, Dovey L, Griffiths RA, Lowe

K, Mahony M, McCallum H, Shuker JD, Simpkins C, Skerratt LF, Williams SE, Hero J-M

(2011) Engineering a future for amphibians under climate change. Journal of Applied

Ecology, **48**, 487-492.

Smith MA, Green DM (2005) Dispersal and the metapopulation paradigm in amphibian ecology and

conservation: are all amphibian populations metapopulations? Ecography, **28**, 110-128.

Smith RK, Meredith H, Sutherland WJ (2019) Amphibian conservation. Pages 9–66 in Sutherland

WJ, Dicks LV, Ockenden N, Petrovan SO, Smith RK, editors. What works in conservation

2019. Open Book Publishers, Cambridge, United Kingdom

Stuart SN, Chanson JS, Cox NA, Young BE, Rodrigues AS, Fischman DL, Waller RW (2004) Status

and trends of amphibian declines and extinctions worldwide. Science, **306**, 1783-1786.

Taylor BE, Scott DE, Gibbons JW (2006) Catastrophic reproductive failure, terrestrial survival, and

persistence of the marbled salamander. Conservation Biology, **20**, 792-801.

van Dijk AI, Beck HE, Crosbie RS, de Jeu RA, Liu YY, Podger GM, Timbal B, Viney NR (2013)

The Millennium Drought in southeast Australia (2001–2009): Natural and human causes and

implications for water resources, ecosystems, economy, and society. Water Resources

Research, **49**, 1040-1057.

Wassens S, Maher M (2011) River regulation influences the composition and distribution of inland

frog communities. River Research and Applications, **27**, 238-246.

Wassens S, Watts RJ, Jansen A, Roshier D (2008) Movement patterns of southern bell frogs (*Litoria*

raniformis) in response to flooding. Wildlife Research, **35**, 50-58.

Williamson I, Bull CM (1999) Population ecology of the Australian frog *Crinia signifera*: larvae.

Wildlife Research, **26**, 81-99.

Figure 1 The reach between Locks 3 and 2 is at the downstream end of the Murray-Darling catchment (shaded grey). Flow is strongly influenced by regulation and extraction throughout the upstream reaches. Australian states are: SA= South Australia, Qld = Queensland, NSW = New South Wales, Vic = Victoria and Tas = Tasmania.

Figure 2 *Litoria raniformis* can live into the fifth year. Compensatory density-feedback reduction (*R*) in annual survival probability (*S*) and fertility (*f*) are calculated from both population and wetland size.

Figure 3. Proportion of the 83 years of observed river height (regulated) and modelled natural river height that potentially supported *Litoria raniformis* breeding (wet years) at sill heights from 7 to 11 mAHD.

Figure 4 Violin plot of wetland inundation pattern over 83 years of observed data under modelled natural flow conditions (a) and river regulation (b). Sill heights are presented in metres with respect to Australian River Height Datum (mAHD). The maximum duration of dry years exceeds the maximum reproductive lifespan of *Litoria raniformis* (six years) under all regulated sill height wetlands.

Figure 5 The probability of extinction at four wetland sizes under a) natural-flow and b) regulated-flow conditions. River regulation has increased the extinction probability of *Litoria raniformis* populations through most of their former habitats. These effects are strongest at smaller wetlands.

Figure 6 Successive dry years increase the probability of extinction and this effect increases with: increasing dry duration, decreasing wetland size, and increased average frequency of dry years at the wetland. Plots are a) 2 successive dry years, b) 3 successive dry years, c) 4 successive dry years and d) 5 successive dry years.

Figure 7 The boosted regression tree shows the relative influence of eight model parameters. Annual probability of survival in the adult and tadpole phases are the two parameters which appreciably drive model outputs.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Supporting Information

Additional Supporting Information can be found in the online version of this article.

Appendix S1. To examine the effects of maximum lifespan on the predicted probability of extinction, we applied a single-parameter perturbation (sensitivity) analysis on a single modelled scenario (a large wetland with a sill height of 7.5 metres with respect to the Australian Height Datum (mAHD) under a river regulation flow scenario).

Figure S1. The Leslie matrix (L1) assigns fertility to age classes 1 – 5 in the top row and age-specific annual survival probability on the sub-diagonal.

Figure S2. Inhibition of egg laying at a wetland follows an exponential decay function. If the total number of eggs laid is greater than 0.8 of the wetland's carrying capacity, subsequent eggs are reduced at an increasing rate.

Figure S3. Compensatory density feedback on survival rate of eggs to the 1-year age class and survival of the 1-year age class to the 2-year age class based on an exponential decay function. Reduction of survival probability starts when the population exceeds 0.3 of carrying capacity.