Title: The memory of airway epithelium damage in smokers and COPD patients

Authors: François M. Carlier[1,2,3], Bruno Detry[1], Marylène Lecocq[1], Amandine M. Collin[1],

Thomas Planté-Bordeneuve[1], Stijn E. Verleden[4], Monique Delos[5], Benoît Rondelet[3,6], Wim

Janssens[4], Jérôme Ambroise[7], Bart M. Vanaudenaerde[4], Sophie Gohy[1,8,9], and Charles

Pilette[1,8*].

Affiliations:

[1]Pole of Pneumology, ENT, and Dermatology, Institute of Experimental and Clinical

Research, Université Catholique de Louvain, Brussels, Belgium

[2]Department of Pneumology, CHU Mont-Godinne UCL Namur, Yvoir, Belgium

[3]Lung Transplant Centre, CHU Mont-Godinne UCL Namur, Yvoir, Belgium

[4]Department of chronic diseases, metabolism and ageing, Katholieke Universiteit Leuven,

Leuven, Belgium

[5]Department of Pathology, CHU Mont-Godinne UCL Namur, Yvoir, Belgium

[6]Deparment of Cardiovascular and Thoracic Surgery, CHU Mont-Godinne UCL Namur,

Yvoir, Belgium.

[7]Centre de Technologies Moléculaires Appliquées, Institute of Experimental and Clinical

Research, Université catholique de Louvain, Brussels, Belgium.

[8]Department of Pneumology, Cliniques Universitaires St-Luc, Brussels, Belgium

[9]Cystic Fibrosis Reference center, Cliniques Universitaires St-Luc, Brussels, Belgium

*__Corresponding author__

Charles Pilette, MD PhD

Pole of Pneumology, ENT, and Dermatology

Institut de Recherche expérimentale et clinique (IREC)

Université Catholique de Louvain

Tour Claude Bernard +5

Avenue Hippocrate, 54

Bte B1.54.04

1200 Brussels

Belgium

Phone : +3227649449

Fax : +3227649440

charles.pilette@saintluc.uclouvain.be

Author contributions

F.M.C. performed experiments, most data analysis, co-supervised the experimental design and

wrote the manuscript. B.D. helped with cell cultures and samples collection. M.L. processed

samples, designed PCR primers, and performed some PCR. T.P.B., W.J. and B.M.V. revised

the manuscript. S.E.V., C.M.S. and B.R. collected and provided samples. J.A. helped with

statistical analyses. A.M.C. helped with cell cultures. S.G. helped with manuscript redaction

and revision. C.P. supervised the design of the study and the writing of the manuscript.

Sources of support. This work was supported by the Fondation Mont-Godinne, Belgium,

grant to FC (N°FMG-2015-BC01, FMG-2016-BC01, and FMG-2017-BC01) and by the

Fonds National de Recherche Scientifique (FNRS), Belgium, grant to FC (N°1.L505.18) and

to CP (N°1.R016.16 and 1.R016.18). Funders were not involved in study design, data

collection, data analysis, interpretation, or writing of the manuscript.

Running title: Memory of airway epithelium pathology in COPD

Total word count: 3261 words

Abstract.

Rationale. Chronic obstructive pulmonary disease (COPD) is a devastating and mostly

irreversible lung disease. In COPD, the bronchial epithelium displays several structural and

functional abnormalities affecting barrier integrity, cell polarity, and differentiation, as well as

epithelial-to-mesenchymal transition and inflammation. Although COPD displays mostly

irreversible changes, the (ir)reversible nature of epithelium pathology *ex vivo* remains poorly

known and was the aim of this study.

Methods. The persistence of COPD epithelium abnormalities was addressed in long-term (10

weeks) primary cultures of air/liquid interface-reconstituted airway epithelium from non-

smoker controls, smoker controls, and COPD patients. Barrier function, epithelial polarity,

cell commitment, epithelial-to-mesenchymal transition and inflammation were assessed *in

vitro*, and certain features were compared *in situ* to the native epithelium. The role of

inflammation was explored by stimulating cultures with a cytokine mix consisting of TNF-α,

IL-6 and IL-1β.

Measurements and main results. Almost all epithelial defects (barrier dysfunction, impaired

polarity, lineage abnormalities) observed in cells from smokers and COPD patients persisted

in vitro up to week 10, except IL-8/CXCL-8 release and epithelial-to-mesenchymal transition

which declined over time. Cell lineage and polarity impairments matched abnormalities

observed *in situ* in the surgical samples from which the *in vitro* epithelium was derived.

Cytokine treatment induced COPD-like changes and, in COPD cells, reactivated epithelial-to-

mesenchymal transition.

Conclusions. The airway epithelium from smokers and COPD patients displays a memory of

its native state and previous injuries by cigarette smoking, which is multidimensional and

sustained for extended periods of time.

Abstract word count. 240

Key words. COPD, epithelial biology, barrier dysfunction, EMT, pIgR

Introduction

Chronic obstructive pulmonary disease (COPD) constitutes the third cause of death worldwide (1), and is mainly due to noxious airborne stimuli, most importantly cigarette smoke (2). COPD is characterized by the irreversible narrowing and disappearance of small conducting airways (3) and destruction of alveolar walls (4). Current therapies provide limited clinical and functional benefits and fail at targeting the underlying pathways leading to structural remodeling of the lungs (5).

The airway epithelium (AE), as first line barrier against inhaled particles, is constantly exposed to airborne pollutants, and fulfils multiple functions to maintain pulmonary homeostasis. It ensures adequate barrier function, cell differentiation and polarization, while maintaining a tight control on inflammatory mechanisms. In COPD, the AE fails at these duties, exhibiting altered physical barrier function (6, 7) underpinned by decreased expression of apical junction complexes (AJCs) proteins (8), abnormal differentiation and function of ciliated and club cells (9-12) as well as hyperplastic goblet cells (13), and aberrant epithelial-to-mesenchymal transition (EMT) that is triggered by cigarette smoke (14) and further enhanced in COPD (15). In addition, the polymeric immunoglobulin receptor (pIgR)/secretory component (SC) system, which ensures the transcytosis and release of polymeric immunoglobulins into mucosal secretions, witnessing epithelial polarization (16, 17) is also defective in COPD, with decreased epithelial pIgR expression (16) and local S-IgA deficiency in small airways (18, 19). Finally, epithelial inflammation control is dysregulated in COPD (20), with increased intraepithelial neutrophils (21) and sputum tumor necrosis factor (TNF)-α, interleukin-8 (IL-8/CXCL-8) and neutrophils (22, 23).

The air/liquid interface (ALI) culture model of primary human bronchial epithelial cells (HBEC) allows the reconstitution of the AE *in vitro* and recapitulates several of the alterations observed *in situ* in COPD. Cigarette smoke-exposed COPD HBEC show decreased E-

cadherin and ZO-1 expression as well as reduced transepithelial electric resistance (TEER)

when compared with smokers without COPD (14, 24) and after 4 weeks culture, the COPD

ALI-AE displays altered lineage differentiation, decreased pIgR expression, and EMT (8, 10,

16), along with increased cytokine release (25). Although these data suggest the persistence of

AE abnormalities, it remains unclear whether (and to what extent) these structural changes

persist on the long-term, matching the irreversible nature of COPD. COPD patients who quit

smoking for more than 3.5 years display less goblet cell hyperplasia (26) and smoking

cessation may improve lung function and survival (27, 28). Conversely, smoking cessation

does not influence COPD-related epidermal growth factor receptor activation (26) or protease

activity (29). In COPD, the inflammatory pattern shared with 'healthy' smokers is amplified

and persists even after smoking cessation, although with conflicting data (27, 30, 31).

Whereas small airway changes are importantly involved in COPD pathophysiology, the recent

Lancet Commission advocates for a broader definition including symptoms related to large

airways (e.g., bronchitis) and for active research in the field (32). The present study aims to

elucidate whether epithelial changes in large airways are persistently imprinted, questioning

the disease memory (31) retained in the COPD epithelium. To address this question, we

reconstituted HBEC-derived ALI-AE from non-smokers, non-COPD smokers and COPD

patients, and cultured it *in vitro* for up to 10 weeks. The spontaneous evolution of

abnormalities was assayed for epithelial readouts including barrier function, cell

differentiation, EMT, pIgR/SC-related polarity and production of inflammatory cytokines. In

addition, we assessed whether exogenous inflammation could trigger COPD-related changes.

Materials and methods

Study population and lung tissue samples

Lung surgical specimens were obtained from patients with or without COPD. Subjects were

sorted on basis of clinical diagnosis and pulmonary function tests, according to the GOLD

2001 classification (33). A primary proximal AE was reconstituted *in vitro* from all subjects

and subjected to long-term culture (10 weeks, n=26) or mid-term culture with cytokine

stimulation (5 weeks, n=25). Tables 1 and S1 recapitulate patients' characteristics for each

population. Detailed information of methods is available in the online data supplement.

In vitro **reconstitution of primary human AE on ALI culture**

ALI-cultures were conducted as previously published (17). TEER was assessed every week

for each sample, using the EMD Millipore™ Millicell-ERS Volt-Ohmmeter (Fisher

Scientific, Hampton, NH, USA) after transiently filling the apical pole with sterile PBS and

correcting for the resistance of the transwell membrane.

Reverse transcriptase quantitative polymerase chain reaction (RT-qPCR)

RNA extraction, reverse transcription, and RT-qPCR were performed as previously described

(34).

Western blot assays

HBEC lysates were analyzed by Western blot as previously described (35) except for

revelation, performed by chemiluminescence (GE Healthcare, Pittsburgh, PA) before

detection by Chemidoc XRS apparatus (Bio-Rad) and quantification by Quantity One

software (Bio-Rad).

Enzyme-linked immunosorbent assay (ELISA) for SC, IL-8/CXCL-8, IL-6, and

fibronectin

Levels of SC (in apical washes) and cytokines (in basolateral supernatants) were measured by

sandwich ELISA, as previously described (11, 35). Fibronectin release in the basal medium

was assessed by direct ELISA as previously reported (8, 36).

Immunofluorescence staining using tyramide signal amplification

Five micron-sections of surgical lung tissue or reconstituted ALI epithelium were prepared

and stained as previously described (17). Detailed information of methods and quantification

is provided in the online data supplement.

Statistical analysis

Data were expressed as means and standard deviation when passing normality tests, otherwise

as medians and interquartile ranges, and were analyzed with JMP® Pro, Version 14 (SAS

Institute Inc., Cary, NC, USA) and GraphPad Prism version 8.0.2 for Windows (GraphPad

Software, La Jolla, CA, USA). p-values < 0.05 were considered statistically significant.

Detailed information of statistical analysis is provided in the online data supplement.

Results

Readouts were assessed (for each sample, according to availability) at different timepoints of

redifferentiation, referred to as early (1 week ALI), short-term (2-3 weeks), mid-term (4-7

weeks) and long-term (8-10 weeks) cultures.

Barrier and junctional properties in the COPD AE

The AE physical barrier was assessed by measuring TEER of ALI-AE up to 10 weeks.

According to smoking status and lung function, the study population was divided into 4

groups: non-smoker controls (NS; n=5), smoker controls (Smo, n=7), mild/moderate COPD

(COPD1-2; n=7) and severe/very severe COPD (COPD3-4; n=7). The characteristics of the

study population are summarized in Table 1.

The COPD AE displayed decreased TEER as compared with NS, and to a lesser extent with

Smo, which also showed decreased TEER compared with NS (Figure 1a). This defect

appeared in early cultures, persisting in long-term cultures (Figure 1b). In addition, TEER

inversely correlated with the disease severity witnessed by the forced expiratory volume in

one second (FEV1). This correlation was significant from early up to long-term cultures

(Figure 1c).

The molecular substratum of this long-lasting barrier disruption was questioned by assessing

mRNA abundance and protein expression of major components of AJCs, namely claudin-1

(*CLDN1*), E-cadherin (*CDH1*), occludin (*OCLN*) and ZO-1/tight junction protein 1 (*TJP1*).

Although no difference was observed regarding mRNA (Figure S1), protein expression of E-

cadherin was reduced in COPD versus NS in early and short-term ALI (Figure 2a), while that

of occludin was decreased from early up to long-term cultures in Smo and COPD (Figure 2b),

and inversely correlated with FEV1 up to mid-term (Figure 2c). Figure 2d shows

representative blots for E-cadherin and occludin in NS and COPD3-4 AE.

These data globally depict epithelial barrier dysfunction that is engaged in Smo, further

worsens in COPD, and persists upon prolonged culture.

Lineage differentiation of the COPD AE

The persistence of differentiation alterations in COPD was questioned by assessing specific

markers and transcription factors of early differentiation towards intermediate cells, as well as

of goblet and ciliated cells.

Epithelial pre-differentiation

mRNA abundance of MYB Proto-Oncogene (*MYB*), a marker of early differentiation of basal

cells (37), was decreased up to long-term in Smo and COPD-derived AE (Figures S2a),

correlating with FEV1 at some time-periods (Figure S2b).

Differentiation towards ciliated cells

The differentiation towards ciliated cells was assessed by measuring the mRNA abundance of

Forkhead Box J1 (*FOXJ1*), a transcription factor involved in ciliated cells' commitment, and

of the dynein axonemal intermediate chain-1 (*DNAI1*), a marker of ciliated cells' terminal

differentiation. We also counted β-tubulin IV$^+$ cells in ALI-AE and in native tissues. Smo-

and COPD-derived cultures displayed persisting decreases of both gene transcripts, which

correlated with FEV1 at some time-periods (Figures 3a-b & S3a-b). β-tubulin IV$^+$ (ciliated)

cells numbers were decreased up to long-term ALI-cultures from Smo and COPD (Figures 3c-

d), in line with reduced β-tubulin IV$^+$ surface in the native COPD AE (Figure 3e).

Differentiation towards goblet cells

The mRNA abundance of SAM Pointed Domain Containing ETS Transcription Factor

(*SPDEF*) and Forkhead Box A3 (*FOXA3*) were assessed as transcription factors inducing

goblet cell differentiation (37-39), along with the protein expression of MUC5AC, a terminal

product of goblet cell differentiation. Compared to NS, *SPDEF* expression was persistently

increased in Smo as well as in COPD, although to a lesser extent (Figure 4a). Similarly,

MUC5AC$^+$ goblet cells numbers were increased in Smo ALI-cultures versus NS (Figure 4b),

matching *in situ* findings, where MUC5AC$^+$ surface in Smo-AE was considerably larger than

in NS (Figure 4c). As for *SPDEF*, *FOXA3* was upregulated in Smo compared with NS (Figure

S3c).

Both *SPDEF* and *FOXA3* expression were decreased at all timepoints in smokers who quit

smoking for more than 4 years, versus active smokers and ex-smokers with a smoking

cessation of less than 4 years (Figure 4d). These results corroborated *in situ* findings, with

reduced MUC5AC$^+$ surface in long-quitters' AE (Figure 4c, right graph).

These data show that goblet cell hyperplasia relates to active/recent smoking and demonstrate

its persistence over time *in vitro*.

EMT of the COPD AE

The expression of EMT protein markers (vimentin, fibronectin) was assessed in long-term

ALI cultures. Increased vimentin contents were observed in early, short-term and mid-term,

but not long-term COPD3-4 versus NS cultures, as well as in early Smo- and COPD1-2

cultures (Figure 5a-b). Accordingly, COPD and Smo-AE displayed increased fibronectin

release up to mid-term cultures versus NS (Figure 5c). No difference remained in long-term

cultures, while no difference was observed regarding vimentin mRNA abundance (*VIM*,

Figure S4). These data show that EMT features progressively vanish in COPD.

Polarity-related pIgR/SC expression

Smo- and COPD-derived AE released less apical SC than NS, from short- up to long-term

ALI cultures (Figure 6a), that was more prominent in COPD3-4 patients and correlated with

disease severity (Figure 6b). Concordantly, COPD3-4 AE displayed decreased *PIGR* mRNA

abundance throughout the long-term cultures. Although this was not significant on isolated

time points (Figure 6c), longitudinal data demonstrated decreased *PIGR* mRNA abundance in

COPD3-4 AE versus other groups (Figure 6c, right panel). In addition, considering that

epithelial differentiation in ALI is completed at 5 weeks, the acquisition of maximal pIgR

protein levels was delayed in COPD AE (Figure 6d). Concordantly, decreased pIgR levels

were also observed *in situ*, in bronchial sections from Smo and COPD AE (Figure 6e).

These data show that both pIgR expression and functionality (SC release) are impaired in

COPD. Along with impaired AJCs (see above), this indicates that the polarity of the AE in

COPD is persistently altered.

Inflammatory cytokine production by the COPD AE

IL-8/CXCL-8 and IL-6 are two main epithelial pro-inflammatory cytokines and were assayed

in ALI-AE. A trend for increased *CXCL8* mRNA abundance was seen in Smo and COPD, in

early to mid-term culture (Figure S5a). Accordingly, IL-8/CXCL-8 release was strongly

increased up to mid-term cultures from Smo- and COPD-derived versus NS, whilst this

difference disappeared afterwards (Figure 7a). Although no difference was observed

regarding *IL6* mRNA abundance (Figure S5b), IL-6 production was increased in Smo- and

COPD -AE. In contrast to IL-8/CXCL-8, IL-6 upregulation persisted in Smo and COPD long-

term cultures (Figure 7b).

Exogenous inflammation induces COPD-like AE features

Whether exogenous inflammation could promote COPD-like changes in the AE was assessed

by supplementing the culture medium with IL-6, TNF-α and IL-1β (all at 5ng/ml) for up to 5

weeks. First, TEER was decreased in inflammatory conditions in early cultures with this

defect persisting in subsequent weeks (Figure 8a). Cytokine-exposed cultures displayed

similar values irrespectively of the original phenotype, possibly indicating a maximal effect

on barrier (dys)function at the used concentrations. Second, cultures exposed to inflammatory

cytokines displayed EMT features, with increased fibronectin release and vimentin contents

(Figure 8b-c). Interestingly, this induction that was not significant in NS, was strikingly

increased in COPD compared to pooled controls. These results show that the COPD AE,

whilst losing its intrinsic EMT features, remains prone to develop EMT upon inflammatory

exposure. Third, cytokine-induced inflammation altered epithelial polarity, with SC apical

release being decreased in all cultures except in COPD3-4, probably due to pre-existent

pIgR/SC severe impairment in this group (Figure 8d).

In conclusion, exogenous inflammation induces COPD-like changes including barrier

dysfunction, EMT and altered polarity in NS- and Smo-AE. In addition, the COPD-derived

AE remained prone to develop EMT upon inflammation.

Discussion

Exploiting unprecedented long-term cultures of ALI-reconstituted human AE from COPD and

controls, we demonstrate that the COPD AE retains several abnormalities associated with the

disease for prolonged periods of time and reveal key temporal relationships between smoking

history and airway cell imprinting as assessed in ALI cultures. We also show that

inflammation may recapitulate a COPD-like phenotype and that COPD cells are prone to

EMT programming upon inflammatory stimulation.

COPD is a chronic and progressive disease due to repeated injury of the airway epithelial-

mesenchymal unit by toxics, leading to AE activation and remodeling. COPD patients who

quit smoking may benefit from decreases in mortality, respiratory symptoms and lung

function decline (27, 28, 40, 41), and display reduced AE remodeling and goblet cell

hyperplasia when smoking cessation exceeds 3.5 years (26). In contrast, no change was

observed following smoking cessation regarding axonemal abnormalities in ciliated cells (42),

airway mucosal inflammation (30), sputum IL-8/CXCL-8 or neutrophils (29, 43), suggesting

permanent alterations in these compartments.

This study explores whether the irreversible nature of the disease is imprinted in the AE in

such a way that aberrant features of the native epithelium persist in long-term cultures,

independently of signals provided by repeated insults and/or mesenchymal cells and

surrounding leukocytes. To study this, we chose to use the ALI model as it was previously

shown to recapitulate, at least to some extent, the native COPD phenotype (8, 10, 16, 17, 44),

and we prolonged the culture up to 10 weeks, an unusually long and so far unreported

duration in this model.

A major abnormality that persists in the COPD AE is the barrier and junctional defect (7), that

was initiated in Smo and was associated with decreased protein levels of E-cadherin and

occludin (Figures 1&2). This observation corroborates previous studies showing that CS

exposure disrupts AJCs (45). We here show that this alteration further persists in long-term

cultures for occludin. In addition, changes in TEER and E-cadherin/occludin expression are

further aggravated according to the presence and severity of COPD. No difference was

observed regarding mRNA abundance for the main AJCs' components, suggesting post-

transcriptional regulation.

In line with the requirement of AJCs' integrity to ensure baso-apical epithelial polarity (46),

we show that AE polarity is persistently impaired in COPD. Aside from abovementioned

AJCs disruption, the pIgR/SC system, that allows baso-apical transcytosis of polymeric

immunoglobulins (47), was also shown to be defective in COPD (16). Our study demonstrates

that this impairment persists over time, with decreased SC apical release and *PIGR* mRNA

abundance in COPD3-4 AE (Figure 6). Interestingly, those data are contrasting with previous

findings in asthma where pIgR downregulation, although being similarly present *in situ*, does

not persist in ALI (48), suggesting distinct mechanisms driving epithelium pathology in

asthma and COPD.

Ciliated cell hypoplasia – with decreased *FOXJ1* and *DNAI1* mRNA abundance and reduced

bêta-tubulin IV$^+$ cells numbers – also persists in long-term COPD-derived cultures, matching

in situ findings (Figures 3&S3). In addition, goblet cell hyperplasia persists in Smo, as

SPDEF and *FOXA3* expression, along with MUC5AC$^+$ cells numbers, remain higher in active

smokers and ex-smokers who quit for less than 4 years, compared with long-quitters (Figure

4). These results support *in situ* data, and corroborate previous findings indicating that the

muco-secretory trait relates more directly to smoking rather than to COPD (26).

In contrast, some abnormalities are observed only in short-term cultures. Indeed, we show

aberrant EMT features in Smo and COPD AE during the first weeks of culture, with increased

fibronectin release and vimentin contents, but these features vanish from mid-term cultures

onwards, with complete disappearance occurring earlier in Smo than in COPD (Figure 5).

Nevertheless, the COPD AE remains prone to reactivate EMT programming upon

inflammatory condition, suggesting the existence of imprinting of the COPD AE by previous

(*in vivo*) exposures conditioning its responses to further stimulation (Figure 8).

As observed for EMT, upregulated IL-8/CXCL-8 release by Smo and COPD AE also faded

away from mid-term cultures onwards, whereas IL-6 overproduction persisted in long-term

ALI-AE from both Smo and COPD patients (Figure 7). *In vitro* exposure to inflammatory

cytokines reproduced or aggravated alterations in barrier and polarity features as well as EMT

in controls and COPD AE, respectively.

The fundamental mechanisms of these observations question the nature of epithelial memory.

Inflammatory memory refers to memories of previous immune events enabling barrier tissues

to rapidly recall distinct environmental exposures, which may be stored not only in immune

cells but also in epithelial and mesenchymal cells (49, 50). While memory classically refers in

the immune system to somatic mutations underlying adaptive antibody responses to recall

antigens, "inflammatory memory" is less well defined and may include epigenetic

modifications and chromatin changes that may drive persistent alterations in damaged tissues.

In the airways, progenitor basal cells are prime candidates to retain epithelial memory, as do

epithelium progenitors in the skin towards inflammatory or mechanical stress (50) as well as

in the gut towards dietary components (51). Interestingly, WNT signalling, which is

upregulated in the COPD AE (17), has been shown to be involved in the latter form of

epithelial memory and may regulate stemness and tumorigenicity. In line with the recent

hypothesis that basal cells serve as repositories of allergic inflammatory memory in airway

epithelial cells (49), one could propose that airway stem cells also store the memory of

repeated previous injuries by inhaled toxics such as cigarette smoke.

Our study has several limitations. First, an effect of treatments (e.g., inhaled corticosteroids in

severe patients) on the findings cannot be excluded. However, *in vitro* studies on intestinal

cell lines and primary HBEC cultured at the ALI showed that dexamethasone induced

increased TEER, claudin-2 (52), and E-cadherin (53, 54) expression, while budesonide

exposure of ALI-HBEC counter-acted CS-induced barrier dysfunction (55). In addition,

dexamethasone and fluticasone propionate improved TGF-β_1-induced EMT in A549 cells

(56), suggesting that corticosteroids may rather improve barrier function and EMT changes.

Second, airway progenitor (basal) cells are suspected to display reduced self-renewal (57),

although conflicting data exist (58). One cannot exclude that COPD ALI-AE could originate

from basal cells with lower replication rate. However, no difference was observed regarding

time to reach confluence (both in flasks and in inserts) across study groups. In addition, one

could expect a "catch-up effect" in long-term cultures, which was not the case, arguing for a

progenitor imprinting in COPD, as already suggested by Ghosh and colleagues (57). Third,

the imprinting observed in this model of *ex vivo* reconstituted epithelium that implies

proliferation and differentiation of basal cells in prolonged ALI culture, could be confirmed in

other models such as 3D airway organoids.

In conclusion, this study demonstrates that the AE from Smo and COPD stores the memory of

its native state and previous insults from cigarette smoking. This memory is

multidimensional, including alterations in barrier function, epithelial polarity, lineage

differentiation, as well as IL-6 release and EMT reprogramming. Further studies are required

to explore the mechanisms of this memory, confirm the exact cellular niche retaining it, as

well as identify putative targets for future therapies.

Acknowledgements. The authors would like to thank J. Van Snick for his help with IL-6

ELISA, V. Lacroix, A. Belhaj, Ph. Eucher, and D. Hoton for providing surgical tissue, A.

Daumerie for technical advice, M. de Beukelaer and Ch. Fregimilicka for their help with

samples processing, and the IREC Pole of Microbiology (UCLouvain, Brussels, Belgium) for

sharing their molecular biology facility.

References

1.   Organization WH. Chronic respiratory diseases. 2017 [Available from: http://www.who.int/respiratory/copd/burden/en/.

2.   Salvi SS, Barnes PJ. Chronic obstructive pulmonary disease in non-smokers. Lancet. 2009;374(9691):733-43.

3.   McDonough JE, Yuan R, Suzuki M, Seyednejad N, Elliott WM, Sanchez PG, et al. Small-airway obstruction and emphysema in chronic obstructive pulmonary disease. N Engl J Med. 2011;365(17):1567-75.

4.   Vogelmeier CF, Criner GJ, Martinez FJ, Anzueto A, Barnes PJ, Bourbeau J, et al. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Lung Disease 2017 Report. GOLD Executive Summary. Am J Respir Crit Care Med. 2017;195(5):557-82.

5.   Celli BR, Anderson JA, Cowans NJ, Crim C, Hartley BF, Martinez FJ, et al. Pharmacotherapy and Lung Function Decline in Patients with Chronic Obstructive Pulmonary Disease: A Systematic Review. Am J Respir Crit Care Med. 2020.

6.   Aghapour M, Raee P, Moghaddam SJ, Hiemstra PS, Heijink IH. Airway Epithelial Barrier Dysfunction in Chronic Obstructive Pulmonary Disease: Role of Cigarette Smoke Exposure. Am J Respir Cell Mol Biol. 2018;58(2):157-69.

7.   Carlier FM, de Fays C, Pilette C. Epithelial Barrier Dysfunction in Chronic Respiratory Diseases. Frontiers in Physiology. 2021;12(861).

8.   Gohy ST, Hupin C, Fregimilicka C, Detry BR, Bouzin C, Gaide Chevronay H, et al. Imprinting of the COPD airway epithelium for dedifferentiation and mesenchymal transition. Eur Respir J. 2015;45(5):1258-72.

9.   Yaghi A, Dolovich MB. Airway Epithelial Cell Cilia and Obstructive Lung Disease. Cells. 2016;5(4).

10.   Gohy S, Carlier FM, Fregimilicka C, Detry B, Lecocq M, Ladjemi MZ, et al. Altered generation of ciliated cells in chronic obstructive pulmonary disease. Sci Rep. 2019;9(1):17963.

11.   Pilette C, Godding V, Kiss R, Delos M, Verbeken E, Decaestecker C, et al. Reduced epithelial expression of secretory component in small airways correlates with airflow obstruction in chronic obstructive pulmonary disease. Am J Respir Crit Care Med. 2001;163(1):185-94.

12. Gamez AS, Gras D, Petit A, Knabe L, Molinari N, Vachier I, et al. Supplementing

defect in club cell secretory protein attenuates airway inflammation in COPD. Chest.

2015;147(6):1467-76.

13. Saetta M, Turato G, Baraldo S, Zanin A, Braccioni F, Mapp CE, et al. Goblet cell

hyperplasia and epithelial inflammation in peripheral airways of smokers with both symptoms

of chronic bronchitis and chronic airflow limitation. Am J Respir Crit Care Med. 2000;161(3

Pt 1):1016-21.

14. Milara J, Peiro T, Serrano A, Cortijo J. Epithelial to mesenchymal transition is

increased in patients with COPD and induced by cigarette smoke. Thorax. 2013;68(5):410-20.

15. Mahmood MQ, Sohal SS, Shukla SD, Ward C, Hardikar A, Noor WD, et al. Epithelial

mesenchymal transition in smokers: large versus small airways and relation to airflow

obstruction. Int J Chron Obstruct Pulmon Dis. 2015;10:1515-24.

16. Gohy ST, Detry BR, Lecocq M, Bouzin C, Weynand BA, Amatngalim GD, et al.

Polymeric immunoglobulin receptor down-regulation in chronic obstructive pulmonary

disease. Persistence in the cultured epithelium and role of transforming growth factor-beta.

Am J Respir Crit Care Med. 2014;190(5):509-21.

17. Carlier FM, Dupasquier S, Ambroise J, Detry B, Lecocq M, Bietry-Claudet C, et al.

Canonical WNT pathway is activated in the airway epithelium in chronic obstructive

pulmonary disease. EBioMedicine. 2020;61:103034.

18. Polosukhin VV, Richmond BW, Du RH, Cates JM, Wu P, Nian H, et al. Secretory IgA

Deficiency in Individual Small Airways Is Associated with Persistent Inflammation and

Remodeling. Am J Respir Crit Care Med. 2017;195(8):1010-21.

19. Polosukhin VV, Cates JM, Lawson WE, Zaynagetdinov R, Milstone AP, Massion PP,

et al. Bronchial secretory immunoglobulin a deficiency correlates with airway inflammation

and progression of chronic obstructive pulmonary disease. Am J Respir Crit Care Med.

2011;184(3):317-27.

20. Barnes PJ. Inflammatory mechanisms in patients with chronic obstructive pulmonary

disease. J Allergy Clin Immunol. 2016;138(1):16-27.

21. Eapen MS, Myers S, Walters EH, Sohal SS. Airway inflammation in chronic

obstructive pulmonary disease (COPD): a true paradox. Expert Rev Respir Med.

2017;11(10):827-39.

22. Keatings VM, Collins PD, Scott DM, Barnes PJ. Differences in interleukin-8 and

tumor necrosis factor-alpha in induced sputum from patients with chronic obstructive

pulmonary disease or asthma. Am J Respir Crit Care Med. 1996;153(2):530-4.

23.   Rutgers SR, Timens W, Kaufmann HF, van der Mark TW, Koeter GH, Postma DS.
Comparison of induced sputum with bronchial wash, bronchoalveolar lavage and bronchial
biopsies in COPD. Eur Respir J. 2000;15(1):109-15.

24.   Heijink IH, Noordhoek JA, Timens W, van Oosterhout AJ, Postma DS. Abnormalities
in airway epithelial junction formation in chronic obstructive pulmonary disease. Am J Respir
Crit Care Med. 2014;189(11):1439-42.

25.   Schneider D, Ganesan S, Comstock AT, Meldrum CA, Mahidhara R, Goldsmith AM,
et al. Increased cytokine response of rhinovirus-infected airway epithelial cells in chronic
obstructive pulmonary disease. Am J Respir Crit Care Med. 2010;182(3):332-40.

26.   Lapperre TS, Sont JK, van Schadewijk A, Gosman MM, Postma DS, Bajema IM, et
al. Smoking cessation and bronchial epithelial remodelling in COPD: a cross-sectional study.
Respir Res. 2007;8:85.

27.   Willemse BW, Postma DS, Timens W, ten Hacken NH. The impact of smoking
cessation on respiratory symptoms, lung function, airway hyperresponsiveness and
inflammation. Eur Respir J. 2004;23(3):464-76.

28.   Bai JW, Chen XX, Liu S, Yu L, Xu JF. Smoking cessation affects the natural history
of COPD. Int J Chron Obstruct Pulmon Dis. 2017;12:3323-8.

29.   Louhelainen N, Rytila P, Haahtela T, Kinnula VL, Djukanovic R. Persistence of
oxidant and protease burden in the airways after smoking cessation. BMC Pulm Med.
2009;9:25.

30.   Gamble E, Grootendorst DC, Hattotuwa K, O'Shaughnessy T, Ram FS, Qiu Y, et al.
Airway mucosal inflammation in COPD is similar in smokers and ex-smokers: a pooled
analysis. Eur Respir J. 2007;30(3):467-71.

31.   Lapperre TS, Postma DS, Gosman MM, Snoeck-Stroband JB, ten Hacken NH,
Hiemstra PS, et al. Relation between duration of smoking cessation and bronchial
inflammation in COPD. Thorax. 2006;61(2):115-21.

32.   Stolz D, Mkorombindo T, Schumann DM, Agusti A, Ash SY, Bafadhel M, et al.
Towards the elimination of chronic obstructive pulmonary disease: a Lancet Commission.
Lancet. 2022.

33.   Pauwels RA, Buist AS, Ma P, Jenkins CR, Hurd SS, Committee GS. Global strategy
for the diagnosis, management, and prevention of chronic obstructive pulmonary disease:
National Heart, Lung, and Blood Institute and World Health Organization Global Initiative
for Chronic Obstructive Lung Disease (GOLD): executive summary. Respir Care.
2001;46(8):798-825.

34.   Bustin SA, Benes V, Garson J, Hellemans J, Huggett J, Kubista M, et al. The need for
transparency and good practices in the qPCR literature. Nat Methods. 2013;10(11):1063-7.

35.   Pilette C, Ouadrhiri Y, Dimanche F, Vaerman JP, Sibille Y. Secretory component is
cleaved by neutrophil serine proteinases but its epithelial production is increased by
neutrophils through NF-kappa B- and p38 mitogen-activated protein kinase-dependent
mechanisms. Am J Respir Cell Mol Biol. 2003;28(4):485-98.

36.   Collin AM, Lecocq M, Noel S, Detry B, Carlier FM, Aboubakar Nana F, et al. Lung
immunoglobulin A immunity dysregulation in cystic fibrosis. EBioMedicine.
2020;60:102974.

37.   Chen G, Korfhagen TR, Karp CL, Impey S, Xu Y, Randell SH, et al. Foxa3 induces
goblet cell metaplasia and inhibits innate antiviral immunity. Am J Respir Crit Care Med.
2014;189(3):301-13.

38.   Park KS, Korfhagen TR, Bruno MD, Kitzmiller JA, Wan H, Wert SE, et al. SPDEF
regulates goblet cell hyperplasia in the airway epithelium. J Clin Invest. 2007;117(4):978-88.

39.   Rajavelu P, Chen G, Xu Y, Kitzmiller JA, Korfhagen TR, Whitsett JA. Airway
epithelial SPDEF integrates goblet cell differentiation and pulmonary Th2 inflammation. J
Clin Invest. 2015;125(5):2021-31.

40.   Anthonisen NR, Connett JE, Kiley JP, Altose MD, Bailey WC, Buist AS, et al. Effects
of smoking intervention and the use of an inhaled anticholinergic bronchodilator on the rate of
decline of FEV1. The Lung Health Study. JAMA. 1994;272(19):1497-505.

41.   Fletcher C, Peto R. The natural history of chronic airflow obstruction. Br Med J.
1977;1(6077):1645-8.

42.   Verra F, Escudier E, Lebargy F, Bernaudin JF, De Cremoux H, Bignon J. Ciliary
abnormalities in bronchial epithelium of smokers, ex-smokers, and nonsmokers. Am J Respir
Crit Care Med. 1995;151(3 Pt 1):630-4.

43.   Willemse BW, ten Hacken NH, Rutgers B, Lesman-Leegte IG, Postma DS, Timens
W. Effect of 1-year smoking cessation on airway inflammation in COPD and asymptomatic
smokers. Eur Respir J. 2005;26(5):835-45.

44.   Carlier F, Detry B, Sibille Y, Pilette C. COPD epithelial phenotype shows partial
reversibility in long-term primary epithelial ALI-cultures. Eur Respir J. 2018;52(suppl 62).

45.   Shaykhiev R, Otaki F, Bonsu P, Dang DT, Teater M, Strulovici-Barel Y, et al.
Cigarette smoking reprograms apical junctional complex molecular architecture in the human
airway epithelium in vivo. Cell Mol Life Sci. 2011;68(5):877-92.

46.   Cereijido M, Valdes J, Shoshani L, Contreras RG. Role of tight junctions in establishing and maintaining cell polarity. Annu Rev Physiol. 1998;60:161-77.

47.   Carlier FM, Sibille Y, Pilette C. The epithelial barrier and immunoglobulin A system in allergy. Clin Exp Allergy. 2016;46(11):1372-88.

48.   Ladjemi MZ, Gras D, Dupasquier S, Detry B, Lecocq M, Garulli C, et al. Bronchial Epithelial IgA Secretion Is Impaired in Asthma. Role of IL-4/IL-13. Am J Respir Crit Care Med. 2018;197(11):1396-409.

49.   Ordovas-Montanes J, Beyaz S, Rakoff-Nahoum S, Shalek AK. Distribution and storage of inflammatory memory in barrier tissues. Nat Rev Immunol. 2020;20(5):308-20.

50.   Naik S, Larsen SB, Gomez NC, Alaverdyan K, Sendoel A, Yuan S, et al. Inflammatory memory sensitizes skin epithelial stem cells to tissue damage. Nature. 2017;550(7677):475-80.

51.   Beyaz S, Mana MD, Roper J, Kedrin D, Saadatpour A, Hong SJ, et al. High-fat diet enhances stemness and tumorigenicity of intestinal progenitors. Nature. 2016;531(7592):53-8.

52.   Fischer A, Gluth M, Weege F, Pape UF, Wiedenmann B, Baumgart DC, et al. Glucocorticoids regulate barrier function and claudin expression in intestinal epithelial cells via MKP-1. Am J Physiol Gastrointest Liver Physiol. 2014;306(3):G218-28.

53.   Carayol N, Vachier I, Campbell A, Crampette L, Bousquet J, Godard P, et al. Regulation of E-cadherin expression by dexamethasone and tumour necrosis factor-alpha in nasal epithelium. Eur Respir J. 2002;20(6):1430-6.

54.   Carayol N, Campbell A, Vachier I, Mainprice B, Bousquet J, Godard P, et al. Modulation of cadherin and catenins expression by tumor necrosis factor-alpha and dexamethasone in human bronchial epithelial cells. Am J Respir Cell Mol Biol. 2002;26(3):341-7.

55.   Heijink I, van den Berge M, Kliphuis N, ten Hacken N, Postma D, van Oosterhout A. Corticosteroids improve airway epithelial regeneration and restore oxidative stress-induced epithelial barrier dysfunction. Eur Respir J. 2012;40(Suppl 56).

56.   Yang HW, Lee SA, Shin JM, Park IH, Lee HM. Glucocorticoids ameliorate TGF-beta1-mediated epithelial-to-mesenchymal transition of airway epithelium through MAPK and Snail/Slug signaling pathways. Sci Rep. 2017;7(1):3486.

57.   Ghosh M, Miller YE, Nakachi I, Kwon JB, Baron AE, Brantley AE, et al. Exhaustion of Airway Basal Progenitor Cells in Early and Established Chronic Obstructive Pulmonary Disease. Am J Respir Crit Care Med. 2018;197(7):885-96.

533   58.   Staudt MR, Buro-Auriemma LJ, Walters MS, Salit J, Vincent T, Shaykhiev R, et al.

534   Airway Basal stem/progenitor cells have diminished capacity to regenerate airway epithelium

535   in chronic obstructive pulmonary disease. Am J Respir Crit Care Med. 2014;190(8):955-8.

537 **Table**

	Non-smoker controls (n=5)	Smoker controls (n=7)	COPD1-2 (n=7)	COPD3-4 (n=7)	
N (Male/Female)	5 (2/3)	7 (6/1)	7 (4/3)	7 (2/5)	ns
Age (years)	67.8 ± 13.8	66.1 ± 11.1	66.4 ± 12.3	58.8 ± 5.7	ns
Smoking history (never/former/current n)	5/0/0	0/5/2	0/1/6	0/7/0	p=0.001
Pack-years	NA	28.6 ± 14.4	43.5 ± 15.6	34.4 ± 10.8	ns
If applicable, duration since smoking cessation (months)	NA	73.1 ± 98.4	14.5 ± 23.9	75.4 ± 57.0	ns
FEV1 (% of PV)	100.0 ± 12.7	94.5 ± 18.9	77.3 ± 10.4*[#]	21.0 ± 2.3*[#¶]	p<0.0001
FEV1/VC ratio (%)	81.8 ± 4.7	83.4 ± 6.5	54.6 ± 6.9*[#]	31.6 ± 5.9*[#¶]	p<0.0001
DLCO (% of PV)	97.4 ± 13.0	90.2 ± 13.7	59.2 ± 13.1*	36.6 ± 8.0*[#¶]	p<0.0001
BMI (kg.m^{-2})	28.9 ± 3.5	26.6 ± 5.5	23.7 ± 5.1	22.5 ± 3.7	ns
Inhaled corticosteroids (n/total N)	0/5	1/7	0/7	7/7	p<0.0001
Surgical indication					
- **Neoplasia (n/total N)**	4/5	7/7	12/13	0/13	
○ **SCC**	0	5	2	0	
○ **AC**	2	2	4	0	
○ **Other thoracic cancers**	1	0	1	0	
○ **Pulmonary metastasis of other cancers**	1	0	0	0	
- **Lung transplant**	0	0	0	7	
- **Other**	1	0	0	0	

Table 1 | Patient cohort for ALI-cultures. Data are presented as mean ± SD, unless otherwise stated. Demographic data, lung function tests, smoking history and inhaled corticotherapy are stated for the patient groups, classified according to smoking history and the presence and severity of airflow limitation. AC, adenocarcinoma; ALI, air/liquid interface; BMI, body mass index; COPD, chronic obstructive pulmonary disease; DLCO, diffusing capacity of the lung for CO; FEV1, forced expiratory volume in 1 s; PV, predicted values; SCC, squamous cell cancer; SD, standard deviation; VC, vital capacity.
* = p<0.05 compared to non-smoker controls
[#] = p<0.05 compared to smoker controls
[¶] = p<0.05 compared to COPD stage 1-2 patients
ns, not significant

Figures

Figure 1 | COPD and Smo AE displays persistently decreased TEER compared with NS AE.
a) TEER in the ALI-AE from NS, Smo and COPD patients (GOLD classification from 1 to 4 according to the spirometric severity of the disease). ALI-AE derived from COPD patients display a sharp decrease in TEER as compared with (non)-smokers at all time periods, that is also observed to a lesser extent in mild and moderate COPD. $, £ and § highlight significant decreases in Smo, COPD1-2 and COPD3-4 patients respectively, as compared to NS.
b) Longitudinal analysis of the evolution of the TEER in the ALI-AE from NS, Smo, mild-to-moderate COPD and (very) severe COPD patients, showing a smoking-related persistent barrier dysfunction that is further enhanced in COPD.
c) The barrier dysfunction observed in COPD, witnessed by the TEER decrease, significantly correlates with the disease severity assessed by the FEV1, from early cultures up to long-term cultures. Graphs include only Smo and COPD samples to specifically assess the correlation with the disease severity.

, * indicate p-values of less than 0.001, and 0.0001, respectively. Bars indicate mean ± standard deviation.
AE, airway epithelium; ALI, air-liquid interface; COPD, chronic obstructive pulmonary disease; FEV1, forced expired volume in 1 second; NS, non-smokers; SEM, standard error of the mean; Smo, smokers; TEER, transepithelial electric resistance.

561
Figure 2 | COPD and Smo AE displays decreased E-cadherin and occludin protein expression compared
with NS.
a) Decreased E-cadherin protein levels in early and short-term ALI-AE, but no more at mid-term and long-term,
in Smo and COPD ALI-AE as compared with that of NS.
b) Decreased occludin protein levels, observable from early up to long-term cultures, in Smo and COPD ALI-AE
as compared with that of NS.
c) Occludin decrease observed in COPD, significantly correlates with the disease severity assessed by the FEV1,
from early cultures up to mid-term (but not long-term) cultures. Graphs include only Smo and COPD samples to
specifically assess the correlation with the disease severity.
d) Representative blots for E-cadherin and occludin in NS and very severe COPD AE, showing decrease in E-
cadherin expression in early and short-term ALI, and decrease in occludin expression in COPD ALI-AE from
early up to long-term cultures.

*, **, *** indicate p-values of less than 0.05, 0.01, and 0.001, respectively (analyzed using the Kruskal-Wallis
test followed by Dunn's post-hoc test). Bars indicate median ± interquartile range.
AE, airway epithelium; ALI, air-liquid interface; COPD, chronic obstructive pulmonary disease; FEV1, forced
expired volume in 1 second; NS, non-smokers; ns, not significant; SEM, standard error of the mean; Smo,
smokers.
579
580

Figure 3 | Altered differentiation programming towards ciliated cells in Smo and COPD patients.

a) Decreased *FOXJ1* expression in Smo and COPD AE, from early up to long-term ALI culture. Longitudinal analysis (middle graph) shows a strong, persistent downregulation in Smo and COPD, as compared with NS. At some time periods (as represented here in long-term ALI-AE), *FOXJ1* expression correlated with the severity of the disease witnessed by the FEV1 (left graph). Correlation graph includes only Smo and COPD samples to specifically assess the correlation with the disease severity.

$, £ and § highlight significant decreases in Smo, COPD1-2 and COPD3-4 patients respectively, as compared to NS.

b) Decreased *DNAI1* expression in Smo and COPD AE, from early up to long-term ALI culture. Longitudinal
analysis (middle graph) shows a strong, persistent downregulation in Smo and COPD, as compared with NS. At
some time periods (as represented here in long-term ALI-AE), *DNAI1* expression correlated with the severity of
the disease witnessed by the FEV1 (left graph). Correlation graph includes only Smo and COPD samples to
specifically assess the correlation with the disease severity.
c) Decreased numbers of bêta-tubulin IV positive ciliated cells in Smo and COPD-derived ALI at every time-
period.
d) Illustrative TSA-enhanced immunofluorescence staining for bêta-tubulin IV, demonstrating the differentiated
emergence of ciliated cells in NS and COPD-derived ALI cultures. Scale bar, 50μm.
e) Quantification of bêta-tubulin IV$^+$ surface in bronchial sections demonstrating reduced *in situ* expression of
bêta-tubulin IV in native Smo and COPD AE, as illustrated by TSA-enhanced immunofluorescence staining for
bêta-tubulin IV (right picture; scale bar, 50μm.).
*, **, ***, **** indicate p-values of less than 0.05, 0.01, 0.001, and 0.0001, respectively. Bars indicate median
± interquartile range (panels a, b, and e) or mean ± standard deviation (panel c).
AE, airway epithelium; ALI, air-liquid interface; COPD, chronic obstructive pulmonary disease; CT, control;
FEV1, forced expired volume in 1 second; HKG, housekeeping genes; NS, non-smokers; SEM, standard error of
the mean; Smo, smokers; y, years.
606

Figure 4 | Altered differentiation programming towards goblet cells in Smo and COPD patients.
a) Increased *SPDEF* expression in Smo AE as compared with NS in short-term, mid-term and long-term ALI
culture, as depicted for each timepoint (left graph) and in a longitudinal analysis (right graph, significance bar:
Smo *versus* NS). Of note, error bars were not added on the longitudinal data for reasons of readability.
#, £ and § highlight significant decreases in NS, COPD1-2, and COPD3-4 patients respectively, as compared to
NS.
b) Increased MUC5AC positive goblet cells in Smo AE as compared with NS and COPD patients in early, short-
term, mid-term and long-term ALI culture. Bottom left picture: illustrative TSA immunofluorescence staining
for MUC5AC, showing increased positive goblet cells in ALI cultures (here, in mid-term ALI-AE) from Smo as
compared with NS and COPD.
617

618   c) *In situ*, in bronchial surgical sections, MUC5AC positive surface is increased in Smo AE as compared with
619   NS and COPD. In addition, MUC5A positive surface is significantly higher in patients who quit smoking for less
620   than 4 years (plain dots) as compared with long-quitters (hollow dots).
621   d) At every time period, *SPDEF* (left graph) and FOXA3 (right graph) mRNA levels are decreased in ALI-AE
622   from Smo who quit smoking for more than 4 years (plain dots) as compared with long quitters (hollow dots).
623
624   *, **, *** indicate p-values of less than 0.05, 0.01, 0.001, and 0.0001. Bars indicate means ± standard deviation
625   (panels a, b, and c) or medians ± interquartile ranges (panel d).
626   AE, airway epithelium; ALI, air-liquid interface; COPD, chronic obstructive pulmonary disease; CT, control;
627   FEV1, forced expired volume in 1 second; HKG, housekeeping genes; NS, non-smokers; SEM, standard error of
628   the mean; Smo, smokers; y, years. Scale bar, 50µm.

Figure 5 | COPD-related EMT features vanish during long-term ALI cultures.

a) Vimentin contents are increased in early Smo cultures, and up to mid-term in COPD ALI-AE, with this increase vanishing in long-term cultures.

b) Illustrative gels from NS and very severe COPD-derived AE cultured at the ALI from early up to long-term, illustrating the vimentin increased contents up to mid-term in COPD.

c) Increased fibronectin release disappears in mid-term cultures from Smo and mild-to-moderate COPD patients, while it persists up to later (7 weeks) in (very) severe COPD AE.

*, ** indicate p-values of less than 0.05 and 0.01, respectively. Bars indicate median ± interquartile range. $, £, and § highlight significant decreases in Smo, COPD1-2 and COPD3-4 patients respectively, as compared to NS.

AE, airway epithelium; ALI, air-liquid interface; COPD, chronic obstructive pulmonary disease; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; NS, non-smokers; ns, not significant; Smo, smokers; w, weeks.

Figure 6 | Impaired polarity, witnessed by a disruption and a delayed acquisition of PIGR/SC expression,
is imprinted in the COPD AE.
a) SC apical release in ALI-AE from Smo and COPD patients is decreased as compared with NS, from short-
term ALI-AE up to long-term ALI-AE. \$, £ and § highlight significant decreases in Smo, COPD1-2 and
COPD3-4 patients respectively, as compared to NS.
b) SC decreased release in the COPD AE moderately correlates with the disease severity, witnessed by the
FEV1. Correlation graphs include only Smo and COPD samples to specifically assess the correlation with the
disease severity.
c) *PIGR* mRNA abundance is not significantly decreased in (very) severe COPD ALI-AE at separate time-points
(left panel), but longitudinal analysis shows a significant decrease in *PIGR* expression in (very) severe COPD
AE as compared with NS, Smo, and mild-to-moderate COPD (right panel).
d) pIgR acquisition is delayed during the differentiation of the AE in COPD as compared with NS, as shown in
the representative blots from NS and very severe COPD ALI-AE and summarized in the middle panel. The gap
was maximal at 2 weeks ALI culture (right panel) and catches it up only after 4 weeks.
e) TSA immunofluorescence staining for pIgR, showing reduced positive surface *in situ* (i.e., in bronchial
sections) in the COPD AE as compared with that of NS.
*, **, *** indicate p-values of less than 0.05, 0.01, and 0.001, respectively (analyzed using the Kruskal-Wallis
test followed by Dunn's post-hoc test, except for longitudinal analysis in B and C, mixed model). Bars indicate
mean ± SD (a, c [left], d [right], e), dots indicate mean (c [right], d [left]).
AE, airway epithelium; ALI, air-liquid interface; COPD, chronic obstructive pulmonary disease; FEV1, forced
expired volume in 1 second; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; HKG, housekeeping genes;
NS, non-smokers; ns, not significant; pIgR, polymeric immunoglobulin receptor; PV, predicted values; SC,
secretory component; SD, standard deviation; Smo, smokers; w, weeks.
666

667
Figure 7 | Epithelial release of inflammatory cytokines partly persists in COPD AE.
a) IL-8/CXCL-8 production by the reconstituted ALI-AE is increased in early, short-term and mid-term ALI-AE
from and COPD patients, but disappears afterwards, although a non-significant upward trend persists.
b) IL-6 production by the reconstituted ALI-AE is increased in Smo and COPD and persists in long-term
cultures.
**** indicates p-values of less 0.0001. Bars indicate mean ± SD (panel a) and median ± interquartile range
(panel b).
AE, airway epithelium; ALI, air-liquid interface; COPD, chronic obstructive pulmonary disease; IL, interleukin;
NS, non-smokers; ns, not significant; SD, standard deviation; Smo, smokers; w, weeks.
677
678

Figure 8 | Cytokine activation triggers COPD-like epithelial changes.
a) Epithelial inflammation, driven by exogenous TNF-α, IL-1β and IL-6, induces barrier dysfunction, witnessed
by a dramatic decrease in TEER in each group (NS, Smo, COPD 1-2, COPD3-4).
b-c) Cytokine-induced epithelial inflammation induces EMT in Smo and COPD-derived ALI AE, witnessed by
increased fibronectin release in early, short-term and mid-term cultures (b) and vimentin expression in mid-term
cultures (c). The absolute increase in fibronectin (b) and vimentin (c) was significantly higher in mid-term
COPD AE than in non-COPD AE (pooled Smo and NS, labeled "no COPD").
d) Cytokine-induced epithelial inflammation deteriorates the epithelial polarity, witnessed by the SC apical
release. No difference was seen in (very) severe COPD, due to low baseline levels.
*, **, **** indicate p-values of less than 0.05, 0.01, and 0.0001. Bars indicate median ± interquartile range (b, c,
d) and mean ± SEM (a, e).
AE, airway epithelium; ALI, air-liquid interface; COPD, chronic obstructive pulmonary disease; CT, controls;
EMT, epithelial-to-mesenchymal transition; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; IL,
interleukin; NS, non-smokers; ns, not significant; SC, secretory component; SEM, standard error of the mean;
Smo, smokers; TEER, transepithelial electric resistance; TNF-α, tumor necrosis factor α; w, weeks.
694