Evaluation of SARS-CoV-2 Entry, Inflammation and New Therapeutics in Human Lung Tissue Cells

Judith Grau-Expósito[1]*****, David Perea[1]*****, Marina Suppi[1], Núria Massana[1], Ander Vergara[2], Maria José Soler[2], Benjamin Trinite[3], Julià Blanco[3,4], Javier García-Pérez[5,6], José Alcamí[5,6,7], Anna Serrano-Mollar[8,9], Joel Rosado[10], Vicenç Falcó[1], Meritxell Genescà[1]*****, and Maria J. Buzon[1]*****

[1]Infectious Diseases Department, Vall d'Hebron Research Institute (VHIR), Hospital Universitari Vall d'Hebron, Universitat Autònoma de Barcelona, Barcelona, Spain. VHIR Task Force COVID-19.

[2]Nephrology Research Department, Vall d'Hebron Research Institute (VHIR), Hospital Universitari Vall d'Hebron, Universitat Autònoma de Barcelona, Barcelona, Spain. VHIR Task Force COVID-19.

[3]IrsiCaixa AIDS Research Institute, Germans Trias i Pujol Research Institute (IGTP), Can Ruti Campus, Autonomous University of Barcelona (UAB), 08916 Badalona, Spain.

[4]Chair of Infectious Diseases and Immunity, University of Vic–Central University of Catalonia (UVic-UCC), 08500 Vic, Spain.

[5]AIDS Immunopathology Unit, National Center of Microbiology, Instituto de Salud Carlos III, Madrid, Spain.

[6]Centro de Investigación Biomédica en Red de Enfermedades Infecciosas, Instituto de Salud Carlos III (ISCIII), Madrid, Spain

[7]Clinic HIV Unit, Hospital Clinic, IDIBAPS, Barcelona, Spain.

[8]Experimental Pathology Department, Institut d'Investigacions Biomèdiques de Barcelona, Consejo Superior de Investigaciones Científicas (IIBB-CSIC), Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain.

[9]Centro de Investigaciones Biomédicas en Red de Enfermedades Respiratorias (CIBERES), Madrid, Spain.

[10]Thoracic Surgery and Lung Transplantation Department, Vall d'Hebron Institut de Recerca (VHIR), Hospital Universitari Vall d'Hebron, Barcelona, Spain. VHIR Task Force COVID-19.

Running title: Human lung for the study of SARS-CoV-2

Corresponding authors*: Maria J. Buzon (mariajose.buzon@vhir.org) and Meritxell Genescà

(meritxell.genesca@vhir.org)

Abstract

The development of physiological models that reproduce SARS-CoV-2 infection in primary

human cells will be instrumental to identify host-pathogen interactions and potential

therapeutics. Here, using cell suspensions directly from primary human lung tissues (HLT), we

have developed a rapid platform for the identification of viral targets and the expression of viral

entry factors, as well as for the screening of viral entry inhibitors and anti-inflammatory

compounds. The direct use of HLT cells, without long-term cell culture and *in*

vitro differentiation approaches, preserves main immune and structural cell populations,

including the most susceptible cell targets for SARS-CoV-2; alveolar type II (AT-II) cells, while

maintaining the expression of proteins involved in viral infection, such as ACE2, TMPRSS2, CD147

and AXL. Further, antiviral testing of 39 drug candidates reveals a highly reproducible method,

suitable for different SARS-CoV-2 variants, and provides the identification of new compounds

missed by conventional systems, such as VeroE6. Using this method, we also show that

interferons do not modulate ACE2 expression, and that stimulation of local inflammatory

responses can be modulated by different compounds with antiviral activity. Overall, we present

a relevant and rapid method for the study of SARS-CoV-2.

Graphical Abstract

Highlights

Ex vivo physiological systems for the study of SARS-CoV-2-host interactions are scarce. Here, we

establish a method using primary human lung tissue (HLT) cells for the rapid analysis of cell

tropism and identification of therapeutics.

•   HLT cells preserve main cell subpopulations, including alveolar type-2 cells, and

expression of SARS-CoV-2 entry factors ACE2, CD147, TMPRSS2 and AXL.

•   HLT cells are readily susceptible to SARS-CoV-2 infection without the need of cell

isolation or further cell differentiation.

•   Antiviral testing in HLT cells allows the rapid identification of new drug candidates

against SARS-CoV-2 variants, missed by conventional systems.

• Local inflammation is supported in HLT cells and offers the identification of relevant
anti-inflammatory compounds for SARS-CoV-2 infection.

    Introduction

Only one antiviral against SARS-CoV-2, remdesivir, has been approved for the treatment of

COVID-19 in adults and pediatric patients (12 years of age and older) requiring hospitalization

[1, 2]. Moreover, molnupiravir, a ribonucleoside analog that inhibits SARS-CoV-2 replication, has

shown activity in several preclinical models of SARS-CoV-2, including prophylaxis, treatment, and

prevention of transmission [3, 4]. Promising human clinical trials in non-hospitalized patients are

ongoing (NCT04575597), and it could be the first oral antiviral medicine approved for COVID-19.

Further, the development of *Acute Respiratory Distress Syndrome* in severe COVID-19 patients

has been linked to dysregulated inflammatory responses. In this regard, treatment with the

glucocorticoid dexamethasone decreased 28-day mortality among patients receiving invasive

mechanical ventilation, but little benefit was observed in patients without respiratory support

[5]. Despite these major advances in treatment options for COVID-19, the rapid identification of

new antivirals that could be easily transferred into the clinic is still of paramount importance,

particularly with the potential emergence of drug-resistant variants.

Screening of novel drug candidates is often performed using cell lines. In this sense, the most

widely used cell lines for SARS-CoV-2 studies are epithelial cells derived either from lung (Calu-

3), kidney (VeroE6), or colon (CaCo-2) [6]. These immortalized systems are highly reproducible

and easy to handle but lack physiological relevance. The differential gene expression profiling of

cell lines compared with primary cells from tissues might significantly affect important enzymes

involved in the viral replication cycle. For instance, the level of ACE2 expression, the main

receptor used by SARS-CoV-2 for viral entry, is variable among several cell lines [7], while only a

small fraction of alveolar type II (AT-II) cells, the main target for SARS-CoV-2 in the distal lung,

express ACE2 [8, 9]. In addition, SARS-CoV-2 spike (S) glycoprotein, which is responsible for viral

entry into target cells, can be activated by several host proteases, such as furin, transmembrane

serine proteinase 2 (TMPRSS2) and cathepsin L, in a pH-dependent or independent manner [10,

11]. Whereas in some cell lines S protein is activated by endosomal pH-dependent protease

cathepsin L, in airway epithelial cells viral entry depends on the pH-independent TMPRSS2

protease [11]. Thus, it is currently not well defined if SARS-CoV-2 may utilize multiple cell-type-

specific host proteins for viral replication in primary target tissues and therefore, the potency of

therapeutics directed against these proteins may also differ.

Further, inflammatory immune responses might also impact viral dynamics in the lung by

affecting the expression of entry receptors. In this sense, early studies discovered that ACE2 was

a human interferon-stimulated gene (ISG); IFN-β and IFN-γ were shown to strongly upregulate

the expression of ACE2 at the mRNA and cell surface protein levels, indicating that inflammatory

molecules could shape cell susceptibility to viral infection [12]. However, how anti-inflammatory

drugs may affect ACE2 expression and facilitate SARS-CoV-2 infection remains to be elucidated.

One study reported that the use of nonsteroidal anti-inflammatory drugs (NSAIDs), such as

ibuprofen, was linked to enhanced ACE2 expression in a diabetic-induced rat model [13] and

other reports raised alarms regarding the possible role of NSAIDs at increasing susceptibility to

SARS-CoV-2 infection [14, 15]. On the contrary, experimental and clinical evidence showed that

medium-to-low-dose glucocorticoids may play a protective role in COVID-19 by activating ACE2

and suppressing the associated cytokine storm [16]. Overall, the use of more relevant and

physiological models for the study of SARS-CoV-2 infection, the identification of drug candidates,

or the impact of new therapeutics on the disease, could significantly advance the successful

translation of the results into the clinic.

Primary epithelial cell cultures of nasal and proximal airway epithelium have been used to study

SARS-CoV-2 infection in the upper airways [9, 17-19]. Similarly, organ on-chip and organoid

models of AT-II cells have been successfully developed [18, 20]. While very useful, these models

require long-term culture (sometimes several weeks) combined with the addition of cytokines

that might change cell functionality [21]. The direct use of human lung tissue (HLT) cells offers

important advantages over other *in vitro* and *in vivo* approaches for several reasons; it mimics

the main site of viral replication in the lung, contains all heterogeneous cell components present

in the tissue (with greater functional complexity compared to cell lines), and the cells were not

subjected to long-term culture nor exposed to in vitro differentiation approaches. In the past,

similar lung models have been successfully established to study the effect of allergens and

inflammatory stimuli [22, 23]. Importantly, using HLT cells allow mimicking an inflammatory

local response that could be attenuated by anti-inflammatory drugs, providing a low/medium

throughput screening of anti-inflammatory candidates for the treatment of airway diseases [24].

Significantly, lung tissues not only can be infected with SARS-CoV-2, but also generate local

immune responses to viral infection [25]. Considering all these factors, here we aimed to

characterize a physiological human lung tissue system, which could be used for the study of

virus-host interactions and the identification of potential antiviral compounds and their capacity

to modify local inflammation.

Results

Characterization of HLT cells

Non-neoplastic lung parenchyma was obtained from hospitalized non-COVID19 patients

undergoing thoracic surgery. First, we optimized cell culture and digestion conditions, since the

methodology used for tissue processing can significantly impact cell-type yield, viability and

function of target cell populations. We focused on the preservation of EpCAM$^+$ cells expressing

HLA-DR [26, 27], which in adult alveolar parenchyma is characteristic of AT-II cells (thereafter

referred as enriched AT-II cells). We also aimed to preserve several hematopoietic cell subsets,

as shown in the flow cytometry gating strategy (**Figure S1A**). We observed that collagenase

outperformed liberase and trypsin at preserving the enriched population of AT-II cells, the main

SARS-CoV-2 target (**Figure S2A-B**). Among the hematopoietic cells present in lung parenchyma,

we identified CD3 T lymphocytes (which represented 7.39% ± 6.97 out of the total living cells),

myeloid dendritic cells (0.10% ± 0.06), monocytes/macrophages subsets (1.36% ± 1.27),

neutrophils (5.00% ±4.28) and alveolar macrophages (0.21% ± 0.16). Moreover, out of the non-

hematopoietic cells (CD45-), enriched AT-II and endothelial cells represented 1.11% ± 1.16 and

0.78% ± 0.81 of the total living cells, respectively. Based on EpCAM expression and lack of CD31

and CD45, other epithelial cells including AT-I cells represented 0.52% ± 0.46 out of the living

fraction (**Figure 1A**). All these populations have been previously identified in human lungs [25,

28, 29]. Of note, other abundant structural cell subsets were not defined by specific phenotypic

markers within the same panel. Moreover, the nature of AT-II cells in lung cell suspensions was

also addressed by staining with phosphatase alkaline (**Figure 1B**), which is expressed in AT-II cells

both *in vitro* and *in vivo* [30], and by detection of Surfactant Protein C, which is expressed

exclusively by fully differentiated AT-II cells [31, 32] (**Supplementary Figure 1B**).

Next, we focus on the expression of previously identified proteins involved in viral entry.

Single-cell transcriptome studies have shown that ACE2, one of the main host cell surface

receptors for SARS-CoV-2 attachment and infection, is predominantly expressed by AT-II cells [9,

33]. Moreover, ACE2 expression wanes in distal bronchiolar and alveolar regions paralleled by

SARS-CoV-2 infectivity [9]. In human lung parenchyma-derived cells, we found ACE2 expression

mainly associated to the population enriched in AT-II cells (**Figure 1C**), a finding that was

confirmed by immunohistochemistry in concomitant tissue samples (**Figure 1D**). The percentage

of cells expressing ACE2 was rather small and varied between individuals (6.23% ± 3.47) (**Figure**

1C), as previously described [8]. We also studied the expression of CD147, which has been

reported as a route of SARS-CoV-2 infection *in vitro* and *in vivo* models [34]. CD147 was

ubiquitously expressed in several hematopoietic and non-hematopoietic cells (**Figure 1C**).

Importantly, 92.3% ± 2.4 of the population enriched in AT-II cells expressed CD147 (**Figure 1C**).

Similarly, and in agreement with other studies [35, 36], TMPRSS2 protease expression was also

identified in several subsets, including enriched AT-II cells (31.08% ± 7.06) (**Figure 1C**). When we

studied the double expression of ACE2 and TMPRSS2, or ACE2 and CD147, we found that only

1.31% ± 0.70 and 3.02% ± 1.84 of AT-II cells expressed both markers, respectively

(**Supplementary Figure 3B**). Last, we studied the expression of AXL, another candidate receptor

for SARS-CoV-2 entry in lung cells [37]. We detected high expression of AXL on myeloid dendritic

cells (45.27% ± 17.11), while only a fraction of the enriched AT-II cells (mean of 3.82% ± 4.71)

expressed AXL (**Supplementary Figure 3A**). Overall, we found that human lung cell suspensions

preserved critical populations and factors required for SARS-CoV-2 infection.

Susceptibility of HLT cells to SARS-CoV-2 viral entry

Next, we assessed if HLT cells were susceptible to viral infection. We generated pseudotyped

vesicular stomatitis virus (VSV) viral particles bearing the D614G form of the S protein of SARS-

CoV-2 and expressing either luciferase (VSV*ΔG (Luc)-S) or GFP (VSV*ΔG (GFP)-S) reporter genes

upon cell entry. As a control, we used the VSV-G virus, which has very broad cell tropism. As

expected, VeroE6 cells were highly susceptible to SARS-CoV-2 entry, as demonstrated for

VSV*ΔG (GFP)-S and VSV-ΔG (Luc)-S (**Figure 2A**). Of note, camostat, an inhibitor of the host

protease TMPRSS2, did not inhibit cell entry in this cell line, consistent with its lack of expression

of this protein (**Figure 2B**). Anti-ACE2 antibody blocked more than 90% of VSV*ΔG (Luc)-S, yet

was inactive for VSV-G (Luc) infection (**Figure 2B**). This observation has been widely reported

before [11, 38, 39], and identifies ACE2 as the main cell receptor required for viral entry in

VeroE6. Importantly for viral pathogenesis, it has been postulated that SARS-CoV-2 S protein

might downregulate ACE2 expression, as previously observed for SARS-CoV [40]. We

consistently observed a significant strong reduction in ACE2 expression after viral entry (**Figure**

2C).

We then evaluated the susceptibility of HLT cells to viral entry, using the same viral

constructs. HLT cells were readily infected with pseudotyped S particles (VSV*ΔG (Luc)-S and

VSV*ΔG (GFP)-S), with the natural donor variation representative of primary samples (**Figure**

2D). As expected [9], lung cells enriched with the AT-II phenotype were identified as the main

SARS-CoV-2 cell targets in steady conditions (**Figure 2E**). Blockade of ACE2 resulted in a donor-

dependent reduction of viral infectivity, ranging from 50 to 100% (**Figure 2F**). Camostat

significantly inhibited viral entry in all HLT assays, although the entry process was not always

completely abrogated (**Figure 2F**), suggesting that AT-II cells may become infected through the

use of alternative factors [41]. Similarly, the presence of an anti-CD147 antibody and the

recombinant protein AXL inhibited SARS-CoV-2 entry (**Figure 2F and Supplementary Figure 3D**).

Collectively, these data indicate that HLT cells are susceptible to SARS-CoV-2 viral entry, and that

ACE2, CD147, TMPRSS2 and AXL are important proteins required for viral entry in human lung

cells. Thus, these results support the value of the direct use of HLT cells to successfully study

SARS-CoV-2 viral entry, and related mechanisms, in a more physiological system compared to

immortalized cell lines.

Antiviral assays in HLT cells

To validate the HLT system as a platform for the rapid screening of antiviral candidates, we

assayed potential antiviral compounds, most of them previously identified by computational

methods with predicted ability to inhibit SARS-CoV-2 cell entry due to their interaction with S

protein or with the interface S-ACE2 [42]. A detailed description of the 39 selected drugs is

available in **Table S1**. HLT cells were exposed to VSV*ΔG (Luc)-S virus in the presence of 1/5

serial dilution of the different tested compounds. 20h post-exposure, antiviral activity and cell

viability were measured by luminescence. Cell viability for the individual HLT populations, before

and after SARS-CoV-2 infection, was measured by flow cytometry and is shown in

Supplementary Figure 3C. Antiviral results in HLT cells were systematically compared with

parallel testing in the cell line VeroE6. Among the 39 drugs that were evaluated in our study, 15

of them (38%) showed some antiviral activity against SARS-CoV-2 with EC_{50} values ranging from

0.37μM to 90μM (**Table S2**). From these, 25% had concordant results between both models

(**Figure 3**). Cepharanthine, a naturally occurring alkaloid reported to have potent anti-

inflammatory and antiviral properties, was one of the most potent antivirals identified in both

systems, with EC_{50} of 0.46μM and 6.08μM in VeroE6 and HLT cells, respectively (**Figure 3A and

B**). Of note, we observed cell toxicity at the highest concentrations ($CC_{50\ VeroE6}$= 22.3μM; $CC_{50\ HLT}$=

13.8μM), which translated in satisfactory *selectivity indexes* (SI= CC_{50}/EC_{50} of SI_{VeroE6} = 48.47 and

SI_{HLT} = 2.64 [43]. The anti-SARS-CoV-2 activity of hydroxychloroquine, a compound known to

interfere with endosomal acidification, which is necessary for cathepsin activity, has been

extensively reported [44, 45]. In our study, we observed that hydroxychloroquine was equally

effective at inhibiting viral entry in VeroE6 and HLT cells ($EC_{50\ VeroE6}$= 1.58μM; $EC_{50\ HLT}$= 3.22μM)

(**Figure 3A, B**), with no apparent cytotoxicity (**Figure 3C**). Ergoloid, an approved drug used for

dementia, and recently identified as a potential inhibitor of main protease of SARS-CoV-2 [46],

induced ~90% of viral entry inhibition in HLT cells at non-toxic concentrations (**Figure 3A-C**).

Indeed, *SI* for this compound was higher in the HLT model than in VeroE6 cells (SI_{VeroE6} = 3.9; SI_{HLT}

= 11.38). Similarly, ivermectin, a broad-spectrum anti-parasitic compound, showed very similar

antiviral potency in both models, however *SI* greatly differed between them (SI_{VeroE6} = 1.4; SI_{HLT} =

7.75). Additionally, we plot the individual EC_{50} values obtained from the different donors. We

show that the assay was reproducible (**Supplementary Figure 3E**), highlighting the suitability of

the HLT system for the rapid identification of antivirals.

We also detected discordant antiviral results between both models (**Figure 4**). Four drugs

inhibited SARS-CoV-2 entry in HLT cells without affecting cell viability, with no antiviral effect in

VeroE6 (**Figure 4A-C**). As expected and mentioned before, camostat, a TMPRSS2 inhibitor [47],

was not active in VeroE6 cells due to the lack of TMPRSS2 expression in this cell line. However,

camostat was highly active in HLT cells (EC_{50}=3.3μM). However, in a clinical trial with Covid-19

patients receiving camostat within the first 48h of admission, no significant benefit was

observed [48]. Yet, the potential benefit of camostat during the early phase of infection remains

to be addressed. Further, valaciclovir, an antiviral drug, presented some inhibitory potential at

100 μM only in HLT cells. Interestingly, phenformin, an antidiabetic drug and an mTOR inhibitor,

has been postulated as an inhaled drug candidate against influenza and coronavirus infections

[49]. Phenformin reduced the incidence of influenza infection in diabetic patients during the

1971 outbreak [49]. Here, we detected that phenformin significantly reduced viral entry at 20-

100μM only in HLT cells, supporting previous recommendations as inhaled treatment. Finally,

eriodictyol, a flavonoid used as a medicinal plant [50], demonstrated certain activity starting at

4μM. In contrast, quercetin induced some viral entry suppression only in VeroE6 cells and at

high concentrations (**Figure 4A-C**). To further demonstrate the suitability of our model to test

antivirals against other SARS-CoV-2 variants, we tested the inhibitory capacity of the drugs with

the lowest IC_{50} values, cepharanthine, camostat, ivermectin, hydroxychloroquine, and

ciclesonide using a pseudovirus containing the SARS-CoV-2 spike of the delta variant. Similar

inhibitory dynamics for all tested drugs were shown using both viral variants, indicating the

versatility of our model (**Figure 4D)**.

Last, we verified our main findings in HLT cells using a replication-competent SARS-CoV-2

virus. Although we initially also included ergoloid, we observed high toxicity in HLT cells (>60%

cell toxicity) at longer incubation times (48h), which precluded the execution of the antiviral

assay (**Supplementary Figure 3F**). HLT cells were infected with a SARS-CoV-2 clinical isolate and

viral genomes were measured in the supernatant of cell cultures by RT-PCR in the presence of

the different drugs. SARS-CoV-2 successfully replicated in HLT samples (**Figure 4E**), and viral

replication was partially inhibited by all five drugs (**Figure 4F**). These results are consistent with

previous reports describing the antiviral potency of camostat using precision-cut lung slices from

donors [47, 51]. Overall, these results indicate that HLT cells represent a reproducible and

relevant system for the screening of antivirals in a physiological model. This system not only

recapitulates the main antiviral activities observed in cell models, but also allows the

identification of new compounds missed by conventional systems.

Impact of inflammation and anti-inflammatory drugs on ACE2 expression and SARS-CoV-2

viral entry

Since SARS-CoV-2 viral infection rapidly induces an inflammatory response, we wondered if

certain components of this response could modulate ACE2 expression, potentially increasing

viral binding of SARS-CoV-2 and thus, enhancing infection. Further, ACE2 has been previously

identified as an ISG or a component of the IFN-signaling pathway [12, 52], and a recent

investigation showed that cultured human primary basal epithelial cells treated with IFN-α2 and

IFN-γ led to upregulation of ACE2 [12]. Moreover, IL-1β and IFN-β upregulated ACE2 in large

airway epithelial cell cultures [9]. Thus, considering that type I interferons represent a first line

of defence against viral infections and that several cytokines are rapidly induced and associated

with disease severity in COVID-19 patients [53], we tested the effects of different molecules on

ACE2 expression in HLT cells. In initial experiments, we tested three different doses of a wider

range of molecules (including tumor necrosis factor (TNF), IL-6 and IFN-γ that were subsequently

discarded), which were used to select doses and compounds of interest. Finally, the effect of

IFN-α2, IFN-β1, IL-1β, IL-10 and GM-CSF on ACE2 expression was evaluated in HLT cells. Cells

were then treated with selected immune stimuli and cultured for 20h, when the expression of

ACE2 in enriched AT-II cells was evaluated by flow cytometry. The only significant change we

observed was for IL-1β stimulation, which decreased the fraction of AT-II cells expressing ACE2

(**Figure 5A**). No other significant changes were observed, indicating that relevant inflammatory

stimuli, besides IL-1β, have a limited impact on ACE2 expression in AT-II cells.

Moreover, it is currently not well documented if anti-inflammatory drugs could modulate

ACE2 expression, and consequently, impact susceptibility to SARS-CoV-2 infection [54]. Several

glucocorticoids have shown to impart activating effects on ACE2 expression in cell lines; cortisol

showed the strongest effect on ACE2 activation, followed by prednisolone, dexamethasone, and

methylprednisolone [16]. Moreover, NSAIDs, compounds that inhibit cyclooxygenase-1 and 2

mediating the production of prostaglandins, which play a role in inflammatory responses, have

been linked to ACE2 upregulation [16]. Here, we use HLT cells to study the effect of several anti-

inflammatory drugs on both ACE2 expression and SARS-CoV-2 viral entry. 1/5 dilutions of

ibuprofen, cortisol, dexamethasone and prednisone were added to HLT cells for 20h. Overall, no

effect on ACE2 expression was observed after the addition of these anti-inflammatory drugs
(**Figure 5B**). Consequently, tested anti-inflammatory compounds showed no major impact of the
viral entry assay; however, high concentrations of prednisone and dexamethasone showed a
partial reduction of viral entry in HLT cells, without any apparent impact on VeroE6 cells (**Figure**
5C). Thus, selected anti-inflammatory drugs had limited impact on ACE2 expression within
enriched AT-II cells from the HLT model, as well as in SARS-CoV-2 viral entry.

Anti-inflammatory properties of selected compounds

Last, we were interested in modelling the anti-inflammatory properties of several drugs in
HLT cells. Based on their previous antiviral potency, we selected cepharanthine, ergoloid,
camostat, ivermectin, hydroxychloroquine and ciclesonide for further evaluation. Of note, some
of these drugs have been previously identified as immunomodulators with anti-inflammatory
effects (**Table S1**). However, their direct impact on inflammatory molecules directly secreted by
human lung cells have not been evaluated. HLT cells were stimulated with lipopolysaccharides
(LPS) and IFN-γ in the presence of these antivirals and, 20h after, the expression of IL-6 and
CXCL10, a potent pro-inflammatory cytokine and a chemokine respectively, were intracellularly
measured by flow cytometry. IL-6 and CXCL10 were selected as molecules significantly increased
in severe patients during acute infection, with a prediction value for hospitalization [55]. A
representative flow cytometry gating strategy is shown in **Figure S4**. As shown in **Figure 6A,** two
major subpopulations of myeloid cells contributed to the upregulation of CXCL10 and IL-6
expression upon stimulation. Myeloid CD11b$^+$CD14$^+$ were the cells with a greater response, with
50% of these cells expressing IL-6 and 30% expressing CXCL10 after stimulation (**Figure 6B**).
Using this model of local inflammation, we tested the capacity of the selected compounds to
attenuate this response. We observed that camostat had the most potent effect, which
significantly reduced the expression of CXCL10 in CD11b$^+$CD14$^-$ and of IL-6 in CD11b$^+$CD14$^+$
myeloid subsets (**Figure 6B**). Ergoloid, which has not been linked to modulation of inflammation
before, significantly reduced the expression of cytokines in CD11b$^+$CD14$^+$ myeloid cells, and
cepharanthine reduced IL-6 production within this same subset. In contrast, ciclesonide induced
CXCL10 secretion in CD11b$^+$CD14$^-$ myeloid cells (**Figure 6B**). Altogether, our results validate the
use of HLT cells as a relevant method for the identification of anti-inflammatory compounds
impacting specific pro-inflammatory cell populations located in the lung parenchyma.

Discussion

The emergency created by the fast spread of SARS-CoV-2 infection worldwide required a

quick response from physicians treating these patients, who adapted to the rapid knowledge

being generated by both clinical practice and basic research. However, up to date, only one

antiviral drug against SARS-CoV-2 has been approved for clinical use. New antivirals are urgently

needed, and the choice of the cell and animal models used to test the efficacy of drugs will

impact its rapid translation into the clinics. Here, we propose the use of human lung tissue (HLT)

cells as a method that can be safely performed in a BSL2 facility, which allows i) the identification

of cell targets and expression of viral entry factors, ii) the impact of inflammation on host-

pathogen interactions and iii) a rapid medium-high throughput drug screening of entry inhibitors

against SARS-CoV-2 variants and local anti-inflammatory candidates.

Using pseudotyped viral particles expressing the SARS-CoV-2 spike, we first corroborated that

a fraction of CD45$^-$ CD31$^-$ HLA-DR$^+$ and EpCam$^+$ is enriched in AT-II cells and are the primary cell

target in lung tissue in steady conditions. This agrees with several studies using different

approximations [12, 36, 56] and validates our primary model for viral tropism identification.

While cell lines have been traditionally used for the screening of potential antiviral compounds

due to their reproducibility, as well as being quick and user-friendly assays, they lack

physiological relevance. Similarly, entry receptors and viral factors have been identified using

immortalized cell lines [11, 57], and cell targets for SARS-CoV-2 in tissues have been mainly

determined by analyzing the expression of viral entry factors in RNA-seq datasets [58] or using

replication-competent SARS-CoV-2 isolates in BSL3 facilities [59]. Importantly, these studies

have identified AT-II cells as main viral targets for SARS-CoV-2 infection in the lungs, and the

molecules ACE2, CD147, TMPRSS2 and AXL as important factors for viral entry [11, 34, 37, 60].

However, the development of more refined and translational *ex vivo* models of SARS-CoV-2

entry will not only have implications for understanding viral pathogenesis, but also will be useful

for the characterization of cell targets under specific conditions or for the identification of

potential antivirals blocking viral entry in primary cells. The direct use of HLT cells allows the

maintenance of cell type diversity and it may represent a significant advantage over previous

models [18, 20, 61].

Moreover, we showed that the HLT cells can be successfully used for drug screening

purposes, not only against the D614G virus but also against the delta variant. We tested 39 drugs

and compared the results with antiviral testing in VeroE6 cells. Not surprisingly, we showed

discordant results between both methods. Indeed, we found that 33.3% of the tested

compounds had discordant results between HLT and VeroE6 cells; 26.66% of drugs showed some
antiviral effect in HLT but no activity was detected in VeroE6, and 6.67% showed only antiviral
effects in VeroE6 cells. Among other reasons, the differential expression of several key proteins
needed for viral entry, might explain current discrepancies between cell types. Importantly,
using HLT cells, we identified several compounds with antiviral activity; cepharanthine showed
an EC_{50} of 6.08μM and concordantly, it was recently identified in a high throughput screening as
one of the most potent drugs against SARS-CoV-2 [62], likewise several other studies have
pointed towards this drug as a potent entry and post-entry SARS-CoV-2 inhibitor [63]. Instead,
for hydroxychloroquine, an early report suggested no antiviral activity in human lung cells due
to different expression of the required proteases for viral entry [64]. Furthermore, clinical trials
failed to show effectiveness of this drug as a treatment for COVID-19 [65-67]. A strong
dependency of SARS-CoV-2 on TMPRSS2 for viral entry, rather than on cathepsin L, was
identified as a possible mechanistic explanation for its failure *in vivo* [68]. In our study, however,
we observed that this drug was equally effective at inhibiting viral entry in VeroE6 and HLT cells,
and also was effective when using replication competent viral isolates in HLT cells. Concordantly,
in differentiated air-liquid interface cultures of proximal airway epithelium and 3D organoid
cultures of alveolar epithelium, hydroxychloroquine significantly reduced viral replication [69].
The multiple mechanisms of action postulated for hydroxychloroquine action, including
interference in the endocytic pathway, blockade of sialic acid receptors and restriction of pH
mediated S protein cleavage at the ACE2 binding site [70], could help to explain its antiviral effect
in primary lung cells. Similarly, we identified ivermectin as an effective antiviral in HLT cells. Of
note, ivermectin received limited attention as a potential drug to be repurposed against COVID-
19 based on its limited ability to reach lung tissue *in vivo* [71]. Further, a clinical trial failed to
show a reduction in the proportion of PCR-positive patients seven days after ivermectin
treatment [72].

Importantly, HLT cells also provide a platform for testing anti-inflammatory drugs and the
modulation of viral entry factors with drug candidates and immunomodulatory stimuli. We
showed that IL-1β was able to reduce ACE2 expression in the fraction of enriched AT-II cells, in
contrast to other cytokines induced during SARS-CoV-2 infection like IFN-α2, IFN-β1, IL-10 and
GM-CSF, which did not impact ACE2 protein production. In primary epithelial cells derived from
healthy nasal mucosa, Ziegler et al. [12] showed a significant induction of ACE2 transcripts after
IFN-α2 and IFN-γ stimulation, as well as and in a human bronchial cell line treated with either
type I or type II IFN. Moreover, the authors showed that influenza A virus infection
increased ACE2 expression in lung resections [12], strongly suggesting that ACE2 was an

ISG. However, following studies showed that ACE2 transcription and protein production was not

responsive to IFN. Instead, they described a new RNA isoform, MIRb-ACE2, that was highly

responsive to IFN stimulation, but importantly, encoded a truncated and unstable protein

product [73, 74]. These results highlight the need to validate scRNA-seq data with orthogonal

approaches, such as the confirmation of protein expression levels in relevant systems. In HLT

cells, we quantified ACE2 protein expression and importantly, focused our analysis on putative

AT-II cells, the main SARS-CoV-2 targets in lung parenchyma. Also, in agreement with our results,

a primary human bronchial epithelial cell model, type I (β), II (γ), or III (λ1) IFNs did not induced

ACE2 expression [75]. Moreover, a study performed by Lang et al [76], showed that IFN-γ and

IL-4 downregulate the SARS-CoV receptor ACE2 in VeroE6 cells, and similarly, stimulation of

A549 cells with IFN-α, IFN-γ, and IFN-α+IFN-γ did not identify ACE2 as an ISG [77].

A feasible explanation for the decrease of ACE2 protein production upon IL-1 β treatment is

that IL-1β activates disintegrin and metalloproteinase domain-containing protein 17 (ADAM17)

[78], which mediates the shedding of ACE2 [79]. Although this effect would seem positive to

reduce SARS-CoV-2 infection, ACE2 is a lung-protective factor, as it converts Angiotensin (Ang)

II to Ang-(1–7); while Ang II promotes harmful effects in the lung, e. g. fibrosis, vasoconstriction,

inflammation, endothelial dysfunction, edema, and neutrophil accumulation[80], Ang-(1-7) has

counter-regulatory effects protective of lung injury. Moreover, Ang-(1–7) plays an essential role

in hemostasis, as it favors anti-thrombotic activity in platelets [81]. In any case, treatment of

COVID-19 patients with respiratory insufficiency and hyper inflammation with IL-1 inhibitors was

associated with a significant reduction of mortality [82], indicating that at least during severe

COVID-19 the overall effect of IL-1β is detrimental. While the reduction of ACE2 expression in

AT-II cells by IL-1β may be of interest, it needs to be determined if in combination with other

cytokines rapidly induced during viral respiratory infection [83], this effect would remain.

Further, glucocorticoids and NSAIDS have been linked to ACE2 upregulation previously [16]. In

contrast, we did not observe any significant impact of ibuprofen, cortisol, dexamethasone and

prednisone on ACE2 protein expression. These results are concordant with a recent report

showing that suppression of cyclooxygenase (COX)-2 by two commonly used NSAIDs, ibuprofen

and meloxicam, had no effect on ACE2 expression, viral entry, or viral replication in a mouse

model of SARS-CoV-2 infection [84]. Moreover, dexamethasone incompletely reduced viral

entry. This observation partially agrees with a study using lung cells previously treated with

dexamethasone, which showed significant suppression of SARS-CoV-2 viral growth [25].

We additionally show the interest of the HLT model to test local inflammation and evaluate

potential anti-inflammatory drugs. The culture of diverse cell subsets localized in the lung

parenchyma, without further cell separation, allows the detection of inflammatory responses
generated by different resident subpopulations, which is a significant advantage over
monoculture. Several resident-myeloid subsets, together with newly recruited ones, may
contribute to the rapid cytokine storm detected in COVID-19 patients [85-87]. Thus, the
identification of antiviral drugs that can also limit the extent of these initial pro-inflammatory
events may offer added value to the overall therapeutic effect of a given drug. In this sense, we
observed that camostat significantly reduced the expression of proinflammatory molecules IL-6
and CXCL10 in several myeloid CD11b$^+$ subsets. Concordantly, in a previous study using primary
cultures of human tracheal epithelial cells infected with H1N1 virus, camostat also reduced the
concentrations of the cytokines IL-6 and TNF in cell supernatants [88], suggesting a potent anti-
inflammatory potential. In contrast, ivermectin did not affect the expression of cytokines in our
model. However, ivermectin was previously shown to have protective anti-inflammatory effects
in mice, reducing the production of TNF, IL-1 and IL-6 *in vivo* and *in vitro* [89]. Of note, in our
system we optimized the detection of changes for the intracellular expression of IL-6 and CXCL10
by local myeloid cells, and thus, how these intracellular changes reflect total production in
supernatant needs further evaluation.

Finally, it is also important to note the potential limitations of the model. First, we did not
maintain the cells in an air-liquid interface, which may alter cell function. Other limitations
include the limited availability of human lung samples, inter-patient variation (age, smoking,
etc.), the effects on lung biology of the medical condition instigating surgery, and the exact
location of the sample resection, which may affect the proportion of cell subsets such as AT-II.
However, this variability is what shapes the HLT into a more physiological and relevant model in
comparison to current methods based on immortalized cell cultures. Besides the interest of the
different readouts from HLT cell-based system as proposed here, our results highlight drugs with
antiviral activity together with immunomodulatory properties, which could increase the benefit
of a given treatment during COVID-19 disease progression. For instance, camostat,
cepharanthine and ergoloid were three of the most potent drugs inhibiting SARS-CoV-2 entry,
and remarkably, also exerted a significant anti-inflammatory effect on myeloid cells. Clinical
trials with camostat, ergoloid and cepharanthine, ideally administrated during early infection,
should shed light on their use as both antivirals and anti-inflammatory compounds.

Materials and methods

Cells and virus

VeroE6, isolated from kidney epithelial cells of an African green monkey, were grown in DMEM medium supplemented with 10% fetal bovine serum (FBS; Gibco) 100 U/ml penicillin, and 100 µg/ml streptomycin (Capricorn Scientific) (D10) and maintained at 37°C in a 5% CO_2 incubator.

The spike of the SARS-CoV-2 virus (D614G variant) was generated (GeneArt Gene Synthesis, ThermoFisher Scientific) from the codon-optimized sequence obtained by Ou et al. [41] and inserted into pcDNA3.1D/V5-His-TOPO (pcDNA3.1-S-CoV2Δ19-G614). The spike of the SARS-CoV-2.SctΔ19 B.1.617.2 (delta) virus was generated (GeneArt Gene Synthesis, ThermoFisher Scientific) from the full protein sequence of the original SARS-CoV-2 isolate Wuhan-Hu-1 (WH1) modified to include the mutations specific to the delta variant (VOC-21APR-02: T19R, 157-158del, L452R, T478K, D614G, P681R, D950N). These plasmids present the mutation D614G and a deletion in the last 19 amino acids from the original spike. Pseudotyped viral stocks of VSV*ΔG(Luc)-S were generated following the protocol described by Whitt [90] with some modifications. Briefly, 293T cells were transfected with 3µg of the plasmid encoding the SARS-CoV-2 spike. Next day, cells were infected with a VSV-G-Luc virus (MOI=1) (generated from a lentiviral backbone plasmid that uses a VSV promoter to express luciferase) for 2h and gently washed with PBS. Cells were incubated overnight in D10 supplemented with 10% of I1 hybridoma (anti-VSV-G) supernatant (ATCC CRL-2700) to neutralize contaminating VSV*ΔG(Luc)-G particles. Next day, the resulting viral particles were collected and titrated in VeroE6 cells by enzyme luminescence assay (Britelite plus kit; PerkinElmer), as described previously [91].

Lung tissue

Lung tissues were obtained from patients without previous COVID-19 history and a recent negative PCR test for SARS-CoV-2 infection undergoing thoracic surgical resection at the Thoracic Surgery Service of the Vall d'Hebron University Hospital. Study protocol was approved by the Ethical Committee (Institutional Review Board number PR(AG)212/2020). Non-neoplastic tissue areas were collected in antibiotic-containing RPMI 1640 and immediately dissected into approximately 8-mm^3 blocks. These blocks were first enzymatically digested with 5 mg/ml collagenase IV (Gibco) and 100 µg/ml of DNase I (Roche) for 30 min at 37ºC and 400 rpm and, then, mechanically digested with a pestle. The resulting cellular suspension was filtered through

a 70µm-pore size cell strainer (Labclinics) and washed twice with PBS. Pellet recovered after

centrifugation was resuspended in fresh medium (RPMI 1640 supplemented with 5% FBS, 100

U/ml penicillin, and 100 µg/ml streptomycin) and DNase I to dissolve cell aggregates, and the

resulting cell suspension was then filtered through a 40µm-pore size cell strainer (Labclinics).

Cell number and viability were assessed with LUNA™ Automated Cell Counter (Logos

Biosystems). For cell phenotyping the following antibodies were used: anti-CD31 (PerCP-Cy5.5,

BioLegend), anti-CD11b (FITC, BioLegend), anti-CD11c (Pe-Cy7, BD Biosciences), anti-E-cadherin

(Pe-CF594, BD Biosciences), primary goat anti-ACE2 (R&D systems), anti-CD14 (APC-H7, BD

Biosciences), anti-CD45 (AF700, BioLegend), anti-EpCAM (APC, BioLegend), anti-CD3 (BV650, BD

Biosciences), anti-CD15 (BV605, BD Biosciences) and anti-HLA-DR (BV421, BioLegend). For ACE2

detection, after surface staining, cells were stained with secondary donkey anti-goat IgG (PE,

R&D Systems) for 30 min at 4 ºC. Cell viability was determined using an AQUA viability dye for

flow cytometry (LIVE/DEAD fixable AQUA, Invitrogen). In some experiments, instead of CD11b

or CD15, we used a primary rabbit anti-TMPRSS2 or anti-CD147 (BV605, BD Biosciences),

respectively. For TMPRSS2 detection, after ACE2 staining with the appropriate secondary

antibody, cells were washed twice with PBS 1% NMS (normal mouse serum) and then stained

with a secondary goat anti-rabbit IgG (AF488, Thermofisher) for 30 min at 4ºC. For SPC detection,

after surface staining with a primary rabbit anti-SPC antibody (Biorbyt) and instead of ACE2

staining, cells were stained with a secondary donkey anti-rabbit IgG (PE, Biolegend) for 30 min

at 4 ºC. After fixation with PBS 2% PFA, cells were acquired in an LSR Fortessa (BD Biosciences),

and data were analyzed using the FlowJo v10.6.1 software (TreeStar).

Cytospin and alkaline phosphatase staining

Cytospin preparations were obtained from freshly isolated human lung cells at an

approximate density of 150,000 cells/slide, and air-dried during 15 min. Cells were stained with

alkaline phosphatase, as an enzyme marking epithelial type II cells, following manufacturer's

instructions (Alkaline phosphatase Kit, Sigma). The intensity of pink stain reflects the amount of

alkaline phosphatase in positive cells.

ACE2 immunohistochemical staining in human lung tissue sections

Human lungs were maintained in 10% formalin for 24 hours and then embedded in paraffin.

Paraffin-embedded lungs were cut into 4 µm sections. After removing the paraffin, endogenous

peroxidases were inactivated in an aqueous solution containing 3% H_2O_2 and 10% methanol and

antigen retrieval was performed heating the samples in citrate buffer (10mM citric acid, pH 6.0).

The sections were then blocked in bovine serum albumin (5%), incubated with anti-ACE2
antibody (R&D Systems cat. nº AF933, dilution 1:100) and with biotinylated secondary antibody
against goat IgGs (Vector Laboratories cat. nº BA-9500, dilution 1:250). Proteins were visualized
using the ABC Peroxidase Standard Staining Kit (ThermoFisher) followed by 3,3'-
Diaminobenzidine (DAB) Enhanced Liquid Substrate System (Sigma Aldrich). Counterstaining
was done with hematoxylin.

Antiviral screening assay

The complete list of compounds tested in this study, including information about its clinical
use, product reference and vendors is shown in **Table S1**. Duplicates of five-fold serial dilutions
of 39 antiviral compounds were tested in both VeroE6 cell line and in human lung tissue (HLT)
cells using at least 2 different donors. For VeroE6, five-fold serial dilutions of the compounds,
ranging from 100µM to 0.25nM, were prepared in D10 in a 96-well flat-bottom plates. VeroE6
cells were added at a density of 30.000 cells/well and incubated with the drug for at least 1 h
before infection. Subsequently, cells were infected with 1,500 $TCID_{50}$ of VSV*ΔG(Luc)-S virus. In
parallel, drug cytotoxicity was monitored by luminescence. To evaluate the antiviral activity of
drugs in HLT cells, five-fold serial dilutions of the compounds, ranging from 100µM to 0.8µM or
6.4nM, were prepared in R10 in a 96-well conic-bottom plates. HLT cells were added at a density
of 300,000 cells/well and incubated with the compound for at least 1h before infection. Then,
MOI 0.1 of VSV*ΔG(Luc)-S virus were added to wells, and plates were spinoculated at 1,200g
and 37ºC for 2h. After infection, fresh medium was added to the wells and cell suspensions were
transferred into a 96-well flat-bottom plate. Cells were then cultured overnight at 37°C in a 5%
CO_2 incubator. Each plate contained the following controls: no cells (background control), cells
treated with medium (mock infection) and cells infected but untreated (infection control). After
20h, cells were incubated with Britelite plus reagent (Britelite plus kit; PerkinElmer) and then
transferred to an opaque black plate. Luminescence was immediately recorded by a
luminescence plate reader (LUMIstar Omega). To evaluate cytotoxicity, we used the CellTiter-
Glo® Luminescent kit (Promega), following the manufacturer's instructions. Data was
normalized to the mock-infected control, after which EC_{50} and CC_{50} values were calculated using
Graph-Pad Prism 7.

Drug validation with replication competent SARS-CoV-2

These experiments were performed in a BSL3 facility (Viral Vector Production Unit,
Universitat Autònoma de Barcelona, UAB). The SARS-CoV-2 virus was isolated from a

nasopharyngeal swab from an infected patient hospitalized at the Vall d'Hebron Hospital.
VeroE6 cells were cultured on a cell culture flask (25 cm^2) at 1.5×10^6 cells overnight prior to
inoculation with 1 mL of medium from a Deltaswab VICUM® tub containing the swab. Cells were
cultured for 1h at 37°C and 5% CO$_2$. Afterwards, DMEM containing 2% FCS were added to the
cells and incubated for 48h. Cells were assessed daily for cytopathic effect and the supernatant
was recollected and subjected to viral titration in VeroE6 by plaque assay.

For antiviral drug validation, HLT samples were incubated with different drugs at 20µM for
at least 1h before infection. Tested drugs were camostat, cepharanthine, ergoloid,
hydroxychloroquine, ivermectin and ciclesonide. Then, cells were infected with a MOI 0.5 of the
SARS-CoV-2 viral isolate, and the plate was incubated for 2h at 37°C and 5% CO2. After infection,
samples were extensively washed with PBS 1X to eliminate residual virus and suspended in fresh
media containing antiviral drugs and transferred into a new plate. 24 or 48h post infection, 140µl
of supernatant was collected in tubes containing 140µl of DNA/RNA Shield (Zymo Research) for
SARS-CoV-2 inactivation. For each experiment, a negative control, cells treated with only
medium, and a positive control, cells incubated in the presence of the virus alone, were included.
Percentage of viral infection was calculated by RT-PCR. Briefly, viral RNA from the supernatant
was extracted using the QIAamp Viral RNA Mini Kit (Qiagen), following the manufacturer's
instructions. RNA was reverse transcribed with SuperScript III (Invitrogen), in accordance with
the instructions provided by the manufacturer, and cDNA was quantified by qPCR using the
2019-nCoV CDC RUO Kit (IDT, catalog #10006713) for the detection of viral RNA of the
nucleocapsid region N1 from the SARS-CoV-2 (N1 forward 5'-GACCCCAAAATCAGCGAAAT-3' and
N1 reverse 5'-TCTGGTTACTGCCAGTTGAATCTG-3'; N1 probe 5'-FAMACCCCGCAT/ZEN/TACGTTT
GGTGGACC-3IABkFQ-3'). Copies of SARS-CoV-2 RNA were quantified using a standard (2019-
nCoV_N_Positive Control from IDT, catalog #10006625). Samples were run on a 7000 SDS
instrument (Applied Biosystems).

Modulation of ACE2 expression by anti-inflammatory drugs and immune stimuli

VeroE6 and lung cells were incubated with five-fold serial dilutions of selected anti-
inflammatory compounds (ranging from 100µM to 0.8µM) for 20h. Tested drugs included
cortisol, ibuprofen, prednisone and dexamethasone. Lung cells were also incubated with the
following cytokines: GM-CSF (100 ng/ml, Immunotools), IL-1β (10 ng/ml, Immunotools), IL-10
(100 ng/ml, Immunotools), IFN-β (100 U/ml, Immunotools), or IFN-α2 (100 U/ml, Sigma Aldrich).
For determination of ACE2 expression, the following surface staining antibodies were used:
primary goat anti-ACE2 (R&D Systems), anti-CD45 (AF700, BioLegend), anti-EpCAM (APC,

BioLegend), and anti-HLA-DR (BV421, BioLegend). For ACE2 detection, cells were then stained

with secondary donkey anti-goat IgG (PE, R&D Systems) for 30 min at 4 ºC. A Fluorescent Minus

One control (FMO) without primary anti-ACE2 antibody was used as a control. Cell viability was

determined using an AQUA viability dye for flow cytometry (LIVE/DEAD fixable AQUA,

Invitrogen). After fixation with PBS 2% PFA, cells were acquired in an LSR Fortessa (BD

Biosciences) and analyzed using the FlowJo v10.6.1 software (TreeStar).

Immunomodulatory capacity of selected drugs

HLT cells were cultured in a round-bottom 96-well plate containing 20 µM of cepharanthine,

ergoloid, ciclesonide, hydroxychloroquine, ivermectin, or camostat mesylate alone or in

combination with the stimuli LPS (50 ng/ml) and IFN-γ (100 ng/ml). For each patient, a negative

control, cells treated with only medium, and a positive control, cells incubated in the presence

of LPS and IFN-γ, were included. Immediately, brefeldin A (BD Biosciences) and monensin (BD

Biosciences) were added to cells and cultured overnight at 37 ºC in 5% CO_2. Next day, cellular

suspensions were stained with the following antibodies: anti-CD11b (FITC, BioLegend), anti-

CD69 (PE-CF594, BD Biosciences), anti-CD14 (APC-H7, BD Biosciences), anti-EpCAM (APC,

BioLegend), anti-CD3 (BV650, BD Biosciences), anti-CD45 (BV605, BioLegend), and anti-HLA-DR

(BV421, BioLegend). Cells were subsequently fixed and permeabilized using the

Cytofix/Cytoperm™ kit (BD Biosciences) and intracellularly stained with anti-IL-6 (PE-Cy7,

BioLegend), and anti-CXCL10 (PE, BioLegend). Cell viability was determined using an AQUA

viability dye for flow cytometry (LIVE/DEAD fixable AQUA, Invitrogen). After fixation with PBS

2% PFA, cells were acquired in an LSR Fortessa (BD Biosciences), and data were analyzed using

the FlowJo v10.6.1 software (TreeStar).

Statistical analyses

Statistical analyses were performed with Prism software, version 6.0 (GraphPad). A P value

<0.05 was considered significant.

Author contributions

Conceptualization, MJ.B. and M.G.; Sample Collection, J.R.; Methodology J.G-E, D.P, M.S, N.

M, A.V, MJ.S, J.G-P, J.A, J.B, B.T, A.S.M and V.F; Formal Analysis, J.G-E., D.P, M.S, M.G and MJ.B;

Writing-Original Draft J.G-E and MJ.B; Writing- Review & Editing, J.G-E, M.G. and MJ.B; Funding

Acquisition, M.G and MJ.B.; all authors revised the manuscript; Supervision, M.G and MJ.B.

Acknowledgments

This work was primarily supported by a grant from the Health Department of the
Government of Catalonia (DGRIS 1_5). This work was additionally supported in part by the
Spanish Health Institute Carlos III (ISCIII, PI17/01470; PI19CIII/00004; PI21CIII/00025 and COV20-
00679 (MPY 222-20)), the Spanish Secretariat of Science and Innovation and FEDER funds (grant
RTI2018-101082-B-I00 [MINECO/FEDER]), the Spanish AIDS network Red Temática Cooperativa
de Investigación en SIDA (RD16/0025/0007 and RD16CIII/0002/0001), the European Regional
Development Fund (ERDF), the Fundació La Marató TV3 (grants 201805-10FMTV3 and 201814-
10FMTV3), the Gilead fellowships GLD19/00084 and GLD18/00008 and the Becas Taller Argal
2020. M.J.B is supported by the Miguel Servet program funded by the Spanish Health Institute
Carlos III (CP17/00179). N.M. is supported by a Ph.D. fellowship from the Vall d'Hebron Institut
de Recerca (VHIR). The funders had no role in study design, data collection and analysis, the
decision to publish, or preparation of the manuscript.

References

1. Goldman, J.D., et al., *Remdesivir for 5 or 10 Days in Patients with Severe Covid-19.* N Engl J Med, 2020. **383**(19): p. 1827-1837.
2. Beigel, J.H., et al., *Remdesivir for the Treatment of Covid-19 - Final Report.* N Engl J Med, 2020. **383**(19): p. 1813-1826.
3. Cox, R.M., J.D. Wolf, and R.K. Plemper, *Therapeutically administered ribonucleoside analogue MK-4482/EIDD-2801 blocks SARS-CoV-2 transmission in ferrets.* Nat Microbiol, 2021. **6**(1): p. 11-18.
4. Painter, W.P., et al., *Human Safety, Tolerability, and Pharmacokinetics of Molnupiravir, a Novel Broad-Spectrum Oral Antiviral Agent with Activity Against SARS-CoV-2.* Antimicrob Agents Chemother, 2021.
5. Group, R.C., et al., *Dexamethasone in Hospitalized Patients with Covid-19.* N Engl J Med, 2021. **384**(8): p. 693-704.
6. Takayama, K., *In Vitro and Animal Models for SARS-CoV-2 research.* Trends Pharmacol Sci, 2020. **41**(8): p. 513-517.
7. Mossel, E.C., et al., *Exogenous ACE2 expression allows refractory cell lines to support severe acute respiratory syndrome coronavirus replication.* J Virol, 2005. **79**(6): p. 3846-50.
8. Ortiz, M.E., et al., *Heterogeneous expression of the SARS-Coronavirus-2 receptor ACE2 in the human respiratory tract.* EBioMedicine, 2020. **60**: p. 102976.
9. Hou, Y.J., et al., *SARS-CoV-2 Reverse Genetics Reveals a Variable Infection Gradient in the Respiratory Tract.* Cell, 2020. **182**(2): p. 429-446 e14.
10. Koch, J., et al., *Host Cell Proteases Drive Early or Late SARS-CoV-2 Penetration.* bioRxiv, 2020.
11. Hoffmann, M., et al., *SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor.* Cell, 2020. **181**(2): p. 271-280 e8.
12. Ziegler, C.G.K., et al., *SARS-CoV-2 Receptor ACE2 Is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is Detected in Specific Cell Subsets across Tissues.* Cell, 2020. **181**(5): p. 1016-1035 e19.
13. Qiao, W., et al., *Ibuprofen attenuates cardiac fibrosis in streptozotocin-induced diabetic rats.* Cardiology, 2015. **131**(2): p. 97-106.
14. Moore, N., et al., *Does Ibuprofen Worsen COVID-19?* Drug Saf, 2020. **43**(7): p. 611-614.
15. Kutti Sridharan, G., et al., *COVID-19 and Avoiding Ibuprofen. How Good Is the Evidence?* Am J Ther, 2020. **27**(4): p. e400-e402.
16. Xiang, Z., et al., *Glucocorticoids improve severe or critical COVID-19 by activating ACE2 and reducing IL-6 levels.* Int J Biol Sci, 2020. **16**(13): p. 2382-2391.
17. Ravindra, N.G., et al., *Single-cell longitudinal analysis of SARS-CoV-2 infection in human airway epithelium identifies target cells, alterations in gene expression, and cell state changes.* PLoS Biol, 2021. **19**(3): p. e3001143.
18. Si, L., et al., *A human-airway-on-a-chip for the rapid identification of candidate antiviral therapeutics and prophylactics.* Nat Biomed Eng, 2021.
19. Blanco-Melo, D., et al., *Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19.* Cell, 2020. **181**(5): p. 1036-1045 e9.
20. Lamers, M.M., et al., *An organoid-derived bronchioalveolar model for SARS-CoV-2 infection of human alveolar type II-like cells.* EMBO J, 2021. **40**(5): p. e105912.
21. Kim, J., B.K. Koo, and J.A. Knoblich, *Human organoids: model systems for human biology and medicine.* Nat Rev Mol Cell Biol, 2020. **21**(10): p. 571-584.
22. Chang, Y., et al., *Upregulation of IL-17A/F from human lung tissue explants with cigarette smoke exposure: implications for COPD.* Respir Res, 2014. **15**: p. 145.

23.   Hackett, T.L., et al., *Oxidative modification of albumin in the parenchymal lung tissue of*
current smokers with chronic obstructive pulmonary disease. Respir Res, 2010. **11**: p.
180.
24.   Rimington, T.L., et al., *Defining the inflammatory signature of human lung explant tissue*
in the presence and absence of glucocorticoid. F1000Res, 2017. **6**: p. 460.
25.   Schaller, M.A., et al., *In vitro infection of human lung tissue with SARS-CoV-2:*
Heterogeneity in host defense and therapeutic response. bioRxiv, 2021: p.
2021.01.20.427541.
26.   Hasegawa, K., et al., *Fraction of MHCII and EpCAM expression characterizes distal lung*
epithelial cells for alveolar type 2 cell isolation. Respir Res, 2017. **18**(1): p. 150.
27.   Wosen, J.E., et al., *Epithelial MHC Class II Expression and Its Role in Antigen Presentation*
in the Gastrointestinal and Respiratory Tracts. Front Immunol, 2018. **9**: p. 2144.
28.   Singer, B.D., et al., *Flow-cytometric method for simultaneous analysis of mouse lung*
epithelial, endothelial, and hematopoietic lineage cells. Am J Physiol Lung Cell Mol
Physiol, 2016. **310**(9): p. L796-801.
29.   Baharom, F., et al., *Human Lung Mononuclear Phagocytes in Health and Disease.* Front
Immunol, 2017. **8**: p. 499.
30.   Katsumiti, A., et al., *Immortalisation of primary human alveolar epithelial lung cells using*
a non-viral vector to study respiratory bioreactivity in vitro. Sci Rep, 2020. **10**(1): p.
20486.
31.   Stephan W. Glasser, J.E.B., Thomas R. Korfhagen, *Surfactant Protein-C in the*
Maintenance of Lung Integrity and Function. J Aller Ther, 2011. **S7**(001).
32.   Beers, M.F. and Y. Moodley, *When Is an Alveolar Type 2 Cell an Alveolar Type 2 Cell? A*
Conundrum for Lung Stem Cell Biology and Regenerative Medicine. Am J Respir Cell Mol
Biol, 2017. **57**(1): p. 18-27.
33.   Qi, F., et al., *Single cell RNA sequencing of 13 human tissues identify cell types and*
receptors of human coronaviruses. Biochem Biophys Res Commun, 2020. **526**(1): p. 135-
140.
34.   Wang, K., et al., *CD147-spike protein is a novel route for SARS-CoV-2 infection to host*
cells. Signal Transduct Target Ther, 2020. **5**(1): p. 283.
35.   Qi, J., et al., *The scRNA-seq Expression Profiling of the Receptor ACE2 and the Cellular*
Protease TMPRSS2 Reveals Human Organs Susceptible to SARS-CoV-2 Infection. Int J
Environ Res Public Health, 2021. **18**(1).
36.   Lukassen, S., et al., *SARS-CoV-2 receptor ACE2 and TMPRSS2 are primarily expressed in*
bronchial transient secretory cells. EMBO J, 2020. **39**(10): p. e105114.
37.   Wang, S., et al., *AXL is a candidate receptor for SARS-CoV-2 that promotes infection of*
pulmonary and bronchial epithelial cells. Cell Res, 2021. **31**(2): p. 126-140.
38.   Condor Capcha, J.M., et al., *Generation of SARS-CoV-2 Spike Pseudotyped Virus for Viral*
Entry and Neutralization Assays: A 1-Week Protocol. Front Cardiovasc Med, 2020. **7**: p.
618651.
39.   Fukushi, S., et al., *Vesicular stomatitis virus pseudotyped with severe acute respiratory*
syndrome coronavirus spike protein. J Gen Virol, 2005. **86**(Pt 8): p. 2269-2274.
40.   Kuba, K., et al., *A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS*
coronavirus-induced lung injury. Nat Med, 2005. **11**(8): p. 875-9.
41.   Ou, X., et al., *Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its*
immune cross-reactivity with SARS-CoV. Nat Commun, 2020. **11**(1): p. 1620.
42.   Smith, M. and J.C. Smith, *Repurposing Therapeutics for COVID-19: Supercomputer-Based*
Docking to the SARS-CoV-2 Viral Spike Protein and Viral Spike Protein-Human ACE2
Interface. ChemRxiv, 2020.
43.   Suffness, M. and J. Pezzuto, *Methods in plant biochemistry: assays for bioactivity.*
Academic Press, London, 1990: p. 71-133.

44.   Yao, X., et al., *In Vitro Antiviral Activity and Projection of Optimized Dosing Design of*
726   *Hydroxychloroquine for the Treatment of Severe Acute Respiratory Syndrome*
727   *Coronavirus 2 (SARS-CoV-2).* Clin Infect Dis, 2020. **71**(15): p. 732-739.

728   45.   Maisonnasse, P., et al., *Hydroxychloroquine use against SARS-CoV-2 infection in non-*
729   *human primates.* Nature, 2020. **585**(7826): p. 584-587.

730   46.   Alméciga-Díaz, C.J., et al., *Virtual Screening of Potential Inhibitors for SARS-CoV-2 Main*
731   *Protease*. 2020, Preprints.org.

732   47.   Hoffmann, M., et al., *Camostat mesylate inhibits SARS-CoV-2 activation by TMPRSS2-*
733   *related proteases and its metabolite GBPA exerts antiviral activity.* EBioMedicine, 2021.
734   **65**: p. 103255.

735   48.   Gunst, J.D., et al., *Efficacy of the TMPRSS2 inhibitor camostat mesilate in patients*
736   *hospitalized with Covid-19-a double-blind randomized controlled trial.*
737   EClinicalMedicine, 2021. **35**: p. 100849.

738   49.   Lehrer, S., *Inhaled biguanides and mTOR inhibition for influenza and coronavirus*
739   *(Review).* World Acad Sci J, 2020. **2**(3).

740   50.   Islam, A., et al., *The pharmacological and biological roles of eriodictyol.* Arch Pharm Res,
741   2020. **43**(6): p. 582-592.

742   51.   Hempel, T., et al., *Molecular mechanism of inhibiting the SARS-CoV-2 cell entry facilitator*
743   *TMPRSS2 with camostat and nafamostat.* Chemical Science, 2021. **12**(3): p. 983-992.

744   52.   Su, S. and S. Jiang, *A suspicious role of interferon in the pathogenesis of SARS-CoV-2 by*
745   *enhancing expression of ACE2.* Signal Transduct Target Ther, 2020. **5**(1): p. 71.

746   53.   Vabret, N., et al., *Immunology of COVID-19: Current State of the Science.* Immunity,
747   2020. **52**(6): p. 910-941.

748   54.   Smart, L., et al., *A narrative review of the potential pharmacological influence and safety*
749   *of ibuprofen on coronavirus disease 19 (COVID-19), ACE2, and the immune system: a*
750   *dichotomy of expectation and reality.* Inflammopharmacology, 2020. **28**(5): p. 1141-
751   1152.

752   55.   Grau-Exposito, J., et al., *Peripheral and lung resident memory T cell responses against*
753   *SARS-CoV-2.* Nat Commun, 2021. **12**(1): p. 3010.

754   56.   Hamming, I., et al., *Tissue distribution of ACE2 protein, the functional receptor for SARS*
755   *coronavirus. A first step in understanding SARS pathogenesis.* J Pathol, 2004. **203**(2): p.
756   631-7.

757   57.   Shang, J., et al., *Cell entry mechanisms of SARS-CoV-2.* Proc Natl Acad Sci U S A, 2020.
758   **117**(21): p. 11727-11734.

759   58.   Valyaeva, A.A., et al., *Expression of SARS-CoV-2 entry factors in lung epithelial stem cells*
760   *and its potential implications for COVID-19.* Sci Rep, 2020. **10**(1): p. 17772.

761   59.   Kaufer, A.M., et al., *Laboratory biosafety measures involving SARS-CoV-2 and the*
762   *classification as a Risk Group 3 biological agent.* Pathology, 2020. **52**(7): p. 790-795.

763   60.   Walls, A.C., et al., *Structure, Function, and Antigenicity of the SARS-CoV-2 Spike*
764   *Glycoprotein.* Cell, 2020. **183**(6): p. 1735.

765   61.   Ogando, N.S., et al., *SARS-coronavirus-2 replication in Vero E6 cells: replication kinetics,*
766   *rapid adaptation and cytopathology.* J Gen Virol, 2020. **101**(9): p. 925-940.

767   62.   Jeon, S., et al., *Identification of Antiviral Drug Candidates against SARS-CoV-2 from FDA-*
768   *Approved Drugs.* Antimicrob Agents Chemother, 2020. **64**(7).

769   63.   Rogosnitzky, M., P. Okediji, and I. Koman, *Cepharanthine: a review of the antiviral*
770   *potential of a Japanese-approved alopecia drug in COVID-19.* Pharmacol Rep, 2020.
771   **72**(6): p. 1509-1516.

772   64.   Hoffmann, M., et al., *Chloroquine does not inhibit infection of human lung cells with*
773   *SARS-CoV-2.* Nature, 2020. **585**(7826): p. 588-590.

774   65.   Omrani, A.S., et al., *Randomized double-blinded placebo-controlled trial of*
775   *hydroxychloroquine with or without azithromycin for virologic cure of non-severe Covid-*
776   *19.* EClinicalMedicine, 2020. **29**: p. 100645.

777   66.   Boulware, D.R., et al., *A Randomized Trial of Hydroxychloroquine as Postexposure*
778   *Prophylaxis for Covid-19.* N Engl J Med, 2020. **383**(6): p. 517-525.

779   67.   Mitja, O., et al., *Hydroxychloroquine for Early Treatment of Adults with Mild Covid-19: A*
780   *Randomized-Controlled Trial.* Clin Infect Dis, 2020.

781   68.   Ou, T., et al., *Hydroxychloroquine-mediated inhibition of SARS-CoV-2 entry is attenuated*
782   *by TMPRSS2.* PLoS Pathog, 2021. **17**(1): p. e1009212.

783   69.   Mulay, A., et al., *SARS-CoV-2 infection of primary human lung epithelium for COVID-19*
784   *modeling and drug discovery.* Cell Rep, 2021. **35**(5): p. 109055.

785   70.   Satarker, S., et al., *Hydroxychloroquine in COVID-19: Potential Mechanism of Action*
786   *Against SARS-CoV-2.* Curr Pharmacol Rep, 2020: p. 1-9.

787   71.   Bray, M., et al., *Ivermectin and COVID-19: A report in Antiviral Research, widespread*
788   *interest, an FDA warning, two letters to the editor and the authors' responses.* Antiviral
789   Res, 2020. **178**: p. 104805.

790   72.   Chaccour, C., et al., *The effect of early treatment with ivermectin on viral load, symptoms*
791   *and humoral response in patients with non-severe COVID-19: A pilot, double-blind,*
792   *placebo-controlled, randomized clinical trial.* EClinicalMedicine, 2021. **32**: p. 100720.

793   73.   Onabajo, O.O., et al., *Interferons and viruses induce a novel truncated ACE2 isoform and*
794   *not the full-length SARS-CoV-2 receptor.* Nat Genet, 2020. **52**(12): p. 1283-1293.

795   74.   Ng, K.W., et al., *Tissue-specific and interferon-inducible expression of nonfunctional*
796   *ACE2 through endogenous retroelement co-option.* Nat Genet, 2020. **52**(12): p. 1294-
797   1302.

798   75.   Busnadiego, I., et al., *Antiviral Activity of Type I, II, and III Interferons Counterbalances*
799   *ACE2 Inducibility and Restricts SARS-CoV-2.* mBio, 2020. **11**(5).

800   76.   de Lang, A., A.D. Osterhaus, and B.L. Haagmans, *Interferon-gamma and interleukin-4*
801   *downregulate expression of the SARS coronavirus receptor ACE2 in Vero E6 cells.*
802   Virology, 2006. **353**(2): p. 474-81.

803   77.   Russell, A.B., C. Trapnell, and J.D. Bloom, *Extreme heterogeneity of influenza virus*
804   *infection in single cells.* Elife, 2018. **7**.

805   78.   Hall, K.C. and C.P. Blobel, *Interleukin-1 stimulates ADAM17 through a mechanism*
806   *independent of its cytoplasmic domain or phosphorylation at threonine 735.* PLoS One,
807   2012. **7**(2): p. e31600.

808   79.   Heurich, A., et al., *TMPRSS2 and ADAM17 cleave ACE2 differentially and only proteolysis*
809   *by TMPRSS2 augments entry driven by the severe acute respiratory syndrome*
810   *coronavirus spike protein.* J Virol, 2014. **88**(2): p. 1293-307.

811   80.   Domingo, P., et al., *The four horsemen of a viral Apocalypse: The pathogenesis of SARS-*
812   *CoV-2 infection (COVID-19).* EBioMedicine, 2020. **58**: p. 102887.

813   81.   Fraga-Silva, R.A., et al., *The angiotensin-converting enzyme 2/angiotensin-(1-7)/Mas*
814   *receptor axis: a potential target for treating thrombotic diseases.* Thromb Haemost,
815   2012. **108**(6): p. 1089-96.

816   82.   Cavalli, G., et al., *Interleukin-1 and interleukin-6 inhibition compared with standard*
817   *management in patients with COVID-19 and hyperinflammation: a cohort study.* Lancet
818   Rheumatol, 2021. **3**(4): p. e253-e261.

819   83.   Liao, M., et al., *Single-cell landscape of bronchoalveolar immune cells in patients with*
820   *COVID-19.* Nat Med, 2020. **26**(6): p. 842-844.

821   84.   Chen, J.S., et al., *Nonsteroidal Anti-inflammatory Drugs Dampen the Cytokine and*
822   *Antibody Response to SARS-CoV-2 Infection.* Journal of Virology, 2021. **95**(7): p. e00014-
823   21.

824   85.   Zhang, D., et al., *Frontline Science: COVID-19 infection induces readily detectable*
825   *morphologic and inflammation-related phenotypic changes in peripheral blood*
826   *monocytes.* J Leukoc Biol, 2021. **109**(1): p. 13-22.

827   86.   Fahlberg, M.D., et al., *Cellular events of acute, resolving or progressive COVID-19 in*
828   *SARS-CoV-2 infected non-human primates.* Nat Commun, 2020. **11**(1): p. 6078.

829   87.   Aggarwal, N.R., L.S. King, and F.R. D'Alessio, *Diverse macrophage populations mediate*
830   *acute lung inflammation and resolution.* Am J Physiol Lung Cell Mol Physiol, 2014.
831   **306**(8): p. L709-25.

832   88.   Yamaya, M., et al., *The serine protease inhibitor camostat inhibits influenza virus*
833   *replication and cytokine production in primary cultures of human tracheal epithelial*
834   *cells.* Pulm Pharmacol Ther, 2015. **33**: p. 66-74.

835   89.   Zhang, X., et al., *Ivermectin inhibits LPS-induced production of inflammatory cytokines*
836   *and improves LPS-induced survival in mice.* Inflamm Res, 2008. **57**(11): p. 524-9.

837   90.   Whitt, M.A., *Generation of VSV pseudotypes using recombinant DeltaG-VSV for studies*
838   *on virus entry, identification of entry inhibitors, and immune responses to vaccines.* J
839   Virol Methods, 2010. **169**(2): p. 365-74.

840   91.   Li, M., et al., *Human immunodeficiency virus type 1 env clones from acute and early*
841   *subtype B infections for standardized assessments of vaccine-elicited neutralizing*
842   *antibodies.* J Virol, 2005. **79**(16): p. 10108-25.

Figure 1. Phenotyping of human lung cells. (**A**). *t*-distributed Stochastic Neighbor

Embedding (t-SNE) representation displaying the major cell clusters present in the CD45$^+$ and

CD45$^-$ EpCAM$^+$ fractions of a representative human lung tissue. The vertical bars in the right

panel show the frequency of each subset relative to live cells. All cell subsets were identified as

shown in Figure S1A. mDCs, myeloid dendritic cells; enriched AT-II, enriched fraction in alveolar

type 2. (**B**). Phosphatase alkaline positive AT-II cells (pink staining) were detected in a cytospin

obtained from human lung tissue cells and observed at 10x. Lower panel shows a high

magnification (40x) of the black square. Scale bars are 100 µm and 10 µm in top and bottom

panels, respectively. (**C**). *t*-distributed Stochastic Neighbor Embedding (tSNE) representation for

ACE2, CD147 and TMPRSS2 expression in CD45$^+$ and CD45$^-$EpCAM$^+$ fractions from a

representative lung tissue. Right graphs show the percentage of expression of each entry factor

in the different cell subpopulations, which were identified as in Figure 1A with some

modifications for the identification of myeloid cells and neutrophils (From big cells:

monocytes/macrophages, CD11c$^+$HLA-DR$^+$CD14$^+$; Alveolar macrophages and mDCs, CD11c$^+$HLA-

DR$^+$CD14$^-$; Neutrophils, CD11c$^-$HLA-DR$^-$CD14$^-$CD3$^-$). (**D**). Images of ACE2 immunohistochemical

staining in human lung tissue sections at 40x magnification, counterstained with haematoxylin

(top) or without (bottom). Black arrows indicate staining of ACE2 in AT-II cells (upper panel).

Mean±SEM is shown for all graphs.

Figure 2. Susceptibility of VeroE6 and the HLT model to SARS-CoV-2 viral entry. VeroE6 and

HLT cells were infected with two different viral constructs (GFP and Luciferase) expressing the

spike protein upon viral entry; VSV*ΔG(GFP)-Spike and VSV*ΔG(Luc)-Spike. (**A**) Representative

flow cytometry plots of VeroE6 cells infected with VSV*ΔG(GFP)-Spike or the background form

VSV*ΔG(GFP)-empty (left panel); and luciferase activity (RLUs; relative light units) at 20h post-

infection with the pseudotyped VSV*G(Luc)-G, the VSV*ΔG(Luc)-Spike or the background form

VSV*ΔG(Luc)-empty (right panel). (**B**) Percentage of viral entry after treatment with anti-ACE2

antibody (10µg/ml) and camostat (100µM) in VeroE6 cells infected with the pseudotyped virus

expressing the control G protein or the spike from SARS-CoV-2. (**C**) A flow cytometry plot

showing ACE2 expression in GFP$^+$ VeroE6 cells. Right graph shows mean fluorescence intensity

(MFI) of ACE2 in both infected and uninfected fractions, based on GFP expression. (**D**)

Representative flow cytometry plots of HLT cells infected with the viral construct expressing the

spike protein (VSV*ΔG(GFP)-Spike) or the background form (VSV*ΔG(GFP)-empty) (left panel);

and luciferase activity (RLUs; relative light units) at 20h post-infection with the VSV*ΔG(Luc)-

Spike or the background form VSV*ΔG(Luc)-empty (right panel). Infection was measured as the

percentage of GFP or RLUs, respectively. (**E**) Susceptible HLT cells to viral entry (identified as

GFP+ cells) compatible with an AT-II phenotype, determined by the co-expression of HLA-DR and

EpCAM in the CD45-CD31- fraction of live cells. (**F**) Bar plots showing the percentage of viral entry

inhibition on HLT cells in the presence of anti-ACE2 antibody (15µg/ml), camostat (100µM) or

anti-CD147 antibody (25µg/ml) after cell challenge with VSV*ΔG(Luc)-Spike (left graph) or

VSV*ΔG(GFP)-Spike (right graph). Mean±SEM is shown for all graphs. Data in panel 2C were

analyzed by a Wilcoxon matched-pairs test; *p<0,05. Data in panel 2F were analyzed by one

sample t-test; *p<0.05, **p<0.01, **p<0.0001

Figure 3. Antiviral assays with concordant results between models. (**A**). Percentage of viral

entry in VeroE6 and HLT cells exposed to VSV*ΔG(Luc)-Spike in the presence of cepharanthine,

ergoloid, hydroxychloroquine, hypericin, licofelone, ivermectin, ciclesonide, quercetin,

vidarabine and celecoxib. Drugs were used at concentrations ranging from 100µM to 0.256nM,

in VeroE6, and to 0.8µM in lung cells. Non-linear fit model with variable response curve from at

least three independent experiments in replicates is shown (red lines). Cytotoxic effect on

VeroE6 cells and HLT exposed to drug concentrations in the absence of virus is also shown (green

lines). (**B**). EC$_{50}$ values of each drug in VeroE6 and HLT cells. (**C**). CC$_{50}$ values of each drug are

shown for VeroE6 and for HLT cells.

Figure 4. Antiviral assays with discordant results between models. (**A**). Percentage of viral

entry in VeroE6 and HLT cells exposed to VSV*ΔG(Luc)-Spike in the presence of luteolin,

eriodictyol, phenformin, camostat, sulindac, and valaciclovir. Drugs were used at concentrations

ranging from 100µM to 0.256 nM, in VeroE6, and to 0.8µM in lung cells. Non-linear fit model

with variable response curve from at least three independent experiments in replicates is shown

(red lines). Cytotoxic effect on VeroE6 cells and HLT exposed to drug concentrations in the

absence of virus is also shown (green lines). (**B**). EC$_{50}$ values of each drug in VeroE6 and HLT cells.

(**C**)**.** CC$_{50}$ values of each drug are shown for VeroE6 and HLT cells. (**D**). Percentage of viral entry

in HLT cells exposed to VSV*ΔG(Luc)-Spike-delta and VSV*ΔG(Luc)-Spike-D614G variants in the

presence of ivermectin, camostat, hydroxychloroquine, cepharanthine and ciclesonide. Drugs

were used at concentrations ranging from 100µM to 0.8µM. (**E**) Number of viral genomes/µl at

24h and 48h after infection with replication-competent SARS-CoV-2. (**F**). Percentage of viral

replication in the presence of 20µM of camostat, cepharanthine, hydroxychloroquine,

ivermectin and ciclesonide after infection with replication-competent SARS-CoV-2. Mean±SEM

is shown. Data in panel 4F were analyzed by one sample t-test; *p<0.05, **p<0.01.

Figure 5. Impact of inflammation and anti-inflammatory drugs on SARS-CoV-2 viral entry

and ACE2 expression. Both models, HLT cells and VeroE6 cells, were incubated in the presence

of different anti-inflammatory drugs to evaluate the modulation of ACE2 expression by flow

cytometry and the antiviral effect by luminescence. (**A**). HLT cells were treated with different

stimuli for 20h and the percentage of protein expression (left) or the mean fluorescence

intensity (MFI, right) of ACE2 receptor was evaluated in the enriched AT-II fraction by flow

cytometry. (**B**) Modulation of ACE2 protein expression was assessed by flow cytometry in both

models, Vero E6 and HLT cells, in the presence of different concentrations of each anti-

inflammatory drug, ranging from 100µM to 0.8µM. Percentage of ACE2 expression was

quantified in AT-II cells from at least six independent lung samples, and in VeroE6 cells from 2

independent experiments. (**C**). Cytotoxic effect on Vero E6 and HLT cells exposed to

VSV*ΔG(Luc)-Spike in the presence of different concentrations of the anti-inflammatory drugs

prednisone, cortisol, ibuprofen and dexamethasone. Drugs were used at a concentration

ranging from 100µM to 0.256 nM, in VeroE6, and to 0.8µM in lung cells. Non-linear fit with

variable response curve from at least two experiments in replicates is shown (red lines).

Cytotoxic effect on Vero E6 cells and HLT cells exposed to different concentrations of drugs in

the absence of virus is also shown (green lines). Mean±SEM are shown and statistical

comparisons with the control medium were performed using the Wilcoxon test. *p<0.05.

Figure 6. Anti-inflammatory effect of compounds with antiviral activity against SARS-CoV-

2. HLT cells were cultured in the presence of 20µM of cepharanthine, ergoloid mesylate,

ciclesonide, hydroxychloroquine sulfate, ivermectin or camostat mesylate, alone or in

combination with the stimuli LPS (50 ng/ml) and IFN-γ (100 ng/ml). (**A**) *t*-distributed Stochastic

Neighbor Embedding (t-SNE) representations displaying the major cell clusters present in live

CD45[+] myeloid gate, based on FSC and SSC, of a representative human lung tissue in baseline

conditions and after stimulation with LPS and IFN-γ. Two major subsets of myeloid cells are

shown (CD11b[+] CD14[+], in blue-green, and CD11b[+] CD14[-], in orange). The expression of CXCL10

and IL-6 among the different populations is shown in maroon and green, respectively. (**B**)

Expression of CXCL10 and IL-6 was measured in HLT cells in response to stimuli in the presence

of selected drugs in both myeloid subpopulations, CD11b[+] CD14[+] (left panel) and CD11b[+] CD14[-]

(right panel). HQ, hydroxychloroquine. Mean±SEM are represented and statistical comparisons

with the control medium were performed using the One sample t test. *p<0.05, **p<0.01.

Figure S1. Gating strategy for the identification of cell subpopulations in the human lung

tissue model. (**A**) General gating strategy used to identify different cell subsets in lung samples.

A gate based on FSC vs. SSC was followed by doublet and dead cells exclusion. From live CD45[-]

cells, endothelial cells (CD31$^+$, purple) and epithelial cells (EpCAM$^+$, grey) were gated, and within

epithelial cells, AT-II cells (EpCAM$^+$ and HLA-DR$^+$, pink) were identified. Out of live CD45$^+$ cells

and based on FSC vs. SSC, we identified a lymphocyte population in which we distinguished

between non-T lymphocytes (turquoise) and T cells (dark green) based on CD3 expression; and

big cells, where we identified three major subsets based on their expression of CD11b and CD11c

and, subsequently, CD14 and HLA-DR markers. We identified alveolar macrophages (blue),

monocytes (violet), myeloid dendritic cells (mDCs, fuchsia) and neutrophils (orange). (**B**)

Representative flow cytometry plots showing Surfactant Protein C (SPC) staining and its

respective fluorescence minus one (FMO) control. (**C**) Representative flow cytometry plots

showing ACE2 staining and its respective fluorescence minus one (FMO) control.

Figure S2. Optimization of lung tissue enzymatic digestion visualized by *t*-distributed

Stochastic Neighbor Embedding (tSNE). (**A**) Representative tSNE maps showing concatenated

flow cytometry standard files for three different protocols based on different digestion enzymes

(collagenase, liberase or trypsin) from total live cells (upper), CD45$^+$ cells (middle) and CD45$^-$ cells

(lower). (**B**) Bar plots showing cell-type composition (count) analyzed by flow cytometry for each

tissue protocol.

Figure S3. (**A**) *t*-distributed Stochastic Neighbor Embedding (tSNE) representation for AXL

expression in CD45$^+$ and CD45$^-$EpCAM$^+$ fractions from a representative lung tissue. Right graphs

show the percentage of expression of the AXL entry factor in the different cell populations,

which were identified as in Figure 1A. (**B**) Percentage of enriched AT-II cells co-expressing the

entry factors ACE and CD147 (in blue), and ACE and TMPRSS2 (in purple). (**C**) Frequency of each

subset relative to live cells at 0h and 24h with and without the presence of virus . All cell subsets

were identified as shown in Figure S1A. (**D**) Bar plots showing the percentage of viral entry

inhibition on HLT cells in the presence of anti-ACE2 antibody (15μg/ml) or recombinant human

AXL (50μg/ml) after cell challenge with VSV*ΔG(Luc)-Spike. Data were analyzed by one sample

t-test; *p<0.05, **p<0.01. (**E**) EC$_{50}$ values for in the HLT model obtained from 3 different lung

donors and performed in replicates. (**F**) Cells from 1 donor were cultured with 20 μM of selected

drugs for 48h, and cell toxicity was measured using the CellTiter-Glo® Luminescent kit

(Promega), following the manufacturer's instructions. Data was normalized to the untreated

control. Mean±SEM is shown for all graphs.

Figure S4. Gating strategy for the identification of anti-inflammatory effects of selected

compounds. General gating strategy used to evaluate the expression of inflammatory molecules

in lung samples. A gate based on FSC vs. SSC was followed by doublet and dead cells exclusion.

From live CD45$^+$ cells and based on FSC vs. SSC, we identified lymphocyte population and big

cells, in which we identified two subsets based on their expression of CD11b and CD14: myeloid

CD11b$^+$CD14$^+$ cells (blue-green) and myeloid CD11b$^+$CD14$^-$ cells (orange) are shown.

Figure 1

Figure 2

VeroE6

A

VSV*ΔG(GFP)-empty

VSV*ΔG(GFP)-Spike

B

VSV*G Luc-G   VSV*ΔG Luc-S

C

HLT cells

D

VSV*ΔG(GFP)-empty

VSV*ΔG(GFP)-Spike

VSV*ΔG(Luc)

E

☐ Uninfected (GFP⁻)
■ Infected (GFP⁺)

F

VSV*ΔG(Luc)-S

VSV*ΔG(GFP)-S

Figure 3

A

VeroE6 | HLT | VeroE6 | HLT

Cepharanthine

Hydroxychloroquine

Licofelone

Ciclesonide

Vidarabine

Ergoloid

Hypericin

Ivermectin

Luteolin

Celecoxib

% viral entry

% cell death

Log [drug], µM

B

	VeroE6	HLT
Drugs	EC$_{50}$ (µM)	
Cepharanthine	0.46	6.08
Ergoloid	4.78	9.17
Hydroxychloroquine	1.58	3.22
Hypericin	1.24	0.31
Licofelone	87.50	32.16
Ivermectin	13.94	12.98
Ciclesonide	20.52	16.41
Luteolin	~70.7	~82.39
Vidarabine	>100	51.60
Celecoxib	12.99	55.72

C

	VeroE6	HLT
Drugs	CC$_{50}$ (µM)	
Cepharanthine	22.37	16.15
Ergoloid	18.67	~100
Hydroxychloroquine	>100	~100
Hypericin	4.80	0.14
Licofelone	>100	~53.61
Ivermectin	20.23	~100
Ciclesonide	>100	>100
Luteolin	>100	>100
Vidarabine	>100	>100
Celecoxib	12.98	~100

Figure 4

A

VeroE6   HLT

Camostat

Valaciclovir

Phenformin

Eriodictyol

Quercetin

% viral entry

% cell death

Log [drug], µM

B

Drugs	VeroE6	HLT
	EC$_{50}$ (µM)	
Camostat	no effect	3.30
Valaciclovir	no effect	>100
Phenformin	no effect	38.16
Eriodictyol	no effect	90
Quercetin	>100	no effect

C

Drugs	VeroE6	HLT
	CC$_{50}$ (µM)	
Camostat	>100	>100
Valaciclovir	>100	>100
Phenformin	>100	>100
Eriodictyol	>100	>100
Quercetin	>100	>100

D

% viral entry

log[Ivermectin], uM

log[camostat], uM

log[hydroxychloroquine], uM

log[cepharanthine], uM

log[ciclesonide], uM

D614G
Delta

E

copies/µl of SARS-CoV-2

0h   24h   48h

F

% viral inhibition

C+   Camostat   Cepharanthine   Hydroxychloroquine   Ivermectin   Ciclesonide

Figure 5

Figure 6

A

B