1

Single-molecule imaging of chromatin remodelers reveals role of ATPase in promoting fast kinetics of target search and dissociation from chromatin

Jee Min Kim[1], Pat Visanpattanasin[1,5], Vivian Jou[1,5], Sheng Liu[1], Xiaona Tang[1], Qinsi Zheng[2], Kai Yu Li[1], Jonathan Snedeker[1], Luke D. Lavis[2], Timothée Lionnet[3], Carl Wu[1,4,*]

[1]Department of Biology, Johns Hopkins University, Baltimore, MD 21218, USA

[2]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA 20147, USA

[3]Institute of Systems Genetics, Langone Medical Center, New York University, New York, NY 10016, USA

[4]Department of Molecular Biology and Genetics, Johns Hopkins School of Medicine, Baltimore, MD 21287, USA

[5]These authors contributed equally

[*]Correspondence: wuc@jhu.edu

ABSTRACT

Conserved ATP-dependent chromatin remodelers establish and maintain genome-wide chromatin architectures of regulatory DNA during cellular lifespan, but the temporal interactions between remodelers and chromatin targets have been obscure. We performed live-cell single-molecule tracking for RSC, SWI/SNF, CHD1, ISW1, ISW2, and INO80 remodeling complexes in budding yeast and detected hyperkinetic behaviors for chromatin-bound molecules that frequently transition to the free state for all complexes. Chromatin-bound remodelers display notably higher diffusion than nucleosomal histones, and strikingly fast dissociation kinetics with 4-7 s mean residence times. These enhanced dynamics require ATP binding or hydrolysis by the catalytic ATPase, uncovering an additional function to its established role in nucleosome remodeling. Kinetic simulations show that multiple remodelers can repeatedly occupy the same promoter region on a timescale of minutes, implicating an unending 'tug-of-war' that controls a temporally shifting window of accessibility for the transcription initiation machinery.

KEYWORDS

ATP-dependent chromatin remodelers, single-molecule tracking, live-cell imaging, search and residence times, promoter region occupancy

INTRODUCTION

Eukaryotic chromatin is assembled in nucleosomes and higher order structures that compact the DNA for genome folding in the cell nucleus. Nucleosomes are actively organized at promoter and enhancer elements that are hypersensitive to nuclease digestion (Almer & Hörz, 1986; Heintzman et al., 2007; Wu, 1980). In the budding yeast *Saccharomyces cerevisiae*, gene promoters contain nucleosome-depleted regions (NDRs), approximately 150 base-pair stretches of DNA that are depleted of nucleosomes (Yuan et al., 2005). Non-canonical nucleosome conformations, often called 'fragile nucleosomes', and non-histone protein-DNA complexes, are also observed by limited MNase treatment and occupy a subset of NDRs (Floer et al., 2010; Kubik et al., 2015; Prajapati, Ocampo, & Clark, 2020). NDRs are flanked by well-positioned +1 and -1 nucleosomes, with the +1 nucleosome overlapping the transcription start site (TSS) in

yeast (Albert et al., 2007; Yuan et al., 2005). The +1 nucleosome also phases

downstream nucleosome positions in regularly spaced locations which become

progressively less well-positioned into the gene body (Lai & Pugh, 2017a; Mavrich et al.,

2008). This arrangement of nucleosomes is important for the accurate engagement of

transcription regulators and the transcription pre-initiation complex [PIC], as well as the

progression of the transcription machinery after initiation.

ATP-dependent chromatin remodelers are key *trans*-acting factors in establishing and

maintaining nucleosome organization around genes (Becker & Workman, 2013; Rando

& Winston, 2012; Zhang et al., 2011). As specialized members of the superfamily 2 (SF2)

translocases, chromatin remodeling enzymes share a highly conserved ATPase motor

that utilizes DNA translocation as the fundamental mechanism to restructure DNA-histone

contacts within nucleosomes. In addition to the core ATPase domain, chromatin

remodelers harbor additional functional domains and accessory subunits, forming

multiprotein complexes up to ~1 MDa in size that show substantial functional diversity.

They are further classified into four sub-families based on sequence homology of the

catalytic ATPase and possession of shared components, namely the SWI/SNF [Switch

defective/sucrose non-fermenting], CHD [Chromodomain helicase DNA-binding], ISWI

[Imitation switch], and INO80 [Inositol requiring 80] sub-families.

In vivo studies of remodelers in yeast revealed their distinct genome-wide specificities

and functions in the multi-stage transcription process (Yen, Vinayachandran, Batta,

Koerber, & Pugh, 2012). In this context, remodelers can be distinguished based on their

in vivo specificities for nucleosome targets genome-wide. The first group of remodelers,

RSC, SWI/SNF, INO80, and ISW2, mainly act at gene promoter regions to define the +1

and -1 nucleosome positions. RSC and SWI/SNF mobilize the +1 and -1 nucleosomes

away from the NDR relative to the TSS to promote proper engagement of transcription

initiation machinery (Ganguli, Chereji, Iben, Cole, & Clark, 2014; Klein-Brill, Joseph-

Strauss, Appleboim, & Friedman, 2019; Kubik et al., 2018). Specifically, RSC assists NDR

formation for the majority of yeast genes, and the consequence of conditional RSC

inactivation is a global loss of transcription (Brahma & Henikoff, 2019; Ganguli et al., 2014;

Kubik et al., 2018; Yen et al., 2012). This has led to the concept of RSC (and SWI/SNF)
as nucleosome 'pushers,' widening the NDR (Kubik et al., 2019). Antagonizing the
pushing actions of RSC and SWI/SNF are INO80 and ISW2 (Klein-Brill et al., 2019; Kubik
et al., 2019; Shimada et al., 2008; Yen et al., 2012). Both ISW2 and INO80 remodelers
reposition the +1 and -1 nucleosomes towards the NDR *in vivo*, which is important for
suppressing yeast cryptic transcription via noncanonical TSS usage (Klein-Brill et al.,
2019; Kubik et al., 2019; Whitehouse, Rando, Delrow, & Tsukiyama, 2007).
97
The second group of remodelers, CHD1 and ISW1, act primarily in the gene body where
they maintain proper nucleosome spacing and density relative to the +1 nucleosome.
Their actions are coupled to transcription elongation by interacting with the elongating
polymerase to maintain nucleosome density and thus suppress cryptic initiation within the
gene body (Cheung et al., 2008; Radman-Livaja et al., 2012; Smolle et al., 2012a).
Remodelers with similar *in vivo* activities are functionally redundant as shown by stronger
effects due to multiple deletions or depletions, compared to single deletion or depletion
(Kubik et al., 2019; Ocampo, Chereji, Eriksson, & Clark, 2016). Furthermore, remodelers
act competitively to fine-tune nucleosome positions around genes, leading to proper
transcriptional regulation (Kubik et al., 2019; Ocampo et al., 2016; Ocampo, Chereji,
Eriksson, & Clark, 2019; Parnell, Schlichter, Wilson, & Cairns, 2015). These results
further highlight the current perspective that nucleosomes located around genes are
highly dynamic rather than static, and that the concerted actions of multiple remodelers
result in the striking steady-state nucleosome organization observed by genome-wide
mapping experiments. However, despite this knowledge, a gap still lies in our
understanding of their real-time dynamics and timescales of remodeler interactions on
their chromatin targets.
115
Here we utilize single-molecule tracking (SMT) to directly observe and characterize the
chromatin-binding kinetics of ATP-dependent chromatin remodelers in living cells
(Lionnet & Wu, 2021). We investigated a comprehensive set of remodelers (RSC,
SWI/SNF, CHD1, ISW1, ISW2, INO80) acting at gene promoter regions and gene bodies,
allowing us to quantify and compare their *in vivo* dynamics. We show that remodelers

have varying but substantial frequencies of chromatin binding, while exhibiting a common

target search strategy of frequently engaging in highly transient (sub-second) chromatin

interactions and stable residence times of only several seconds. We also discovered that

the catalytic ATPase is responsible for enhancing their chromatin-associated diffusion

and fast dissociation rates. By integrating the kinetic parameters measured for individual

chromatin remodelers with values from genomic studies, we could simulate substantial

temporal occupancies at yeast chromatin targets, leading to a tug-of-war model for the

organization and dynamic positioning of the nucleosome landscape.

RESULTS

Chromatin remodelers exist in chromatin bound and free populations

We tagged the catalytic subunits of 6 major chromatin remodeling complexes, RSC,

SWI/SNF, CHD1, ISW1 (ISW1a, and ISW1b), ISW2, and INO80 at the C-terminus with

the self-labeling HaloTag by engineering the endogenous loci and expressed the fusion

proteins as the sole source under natural promoter control. The fusion proteins were

localized in the nucleus and did not display detectable cleavage of the tag by SDS-PAGE

(Figure 1-figure supplement 1A,B). Furthermore, no phenotypes were observed for all

strains containing tagged constructs (Figure 1-figure supplement 1C). We then

investigated their endogenous, real-time dynamics as representative subunits of

chromatin remodeling complexes by single-molecule tracking (Figure 1A).

In order to quantify a broad range of kinetic behaviors displayed by remodelers, two

imaging regimes were applied. 'Fast-tracking' acquires 10 ms frame-rate movies to

directly measure a range of single-molecule diffusivities from 'slow' (chromatin-bound) to

'fast' (chromatin-free) and determine fractional representation (Figure 1B). However, high

laser power and extensive photobleaching precludes measurement of chromatin

residence times. 'Slow-tracking' with a longer 250 ms frame-rate and lower laser power

motion-blurs fast diffusing molecules to selectively visualize the chromatin-bound state

and report dwell times (Figure 1C). Combining the two imaging regimes provides a holistic

and quantitative view of a range of diffusive behaviors and kinetic subpopulations.

We applied two independent methods for visualization and quantification of fast-tracking

datasets. First, we determined the diffusion coefficient D for trajectories ≥ 6 frames (i.e.

≥ 60 ms) based on their mean squared displacements (MSD), and present frequency

histograms based on the log(D) values of each trajectory. The histograms were fit to two

Gaussian distributions, representing slow and fast subpopulations (Figure 1-source data

1). For more robust quantification, we applied Spot-On analytics, which uses kinetic

modeling based on distribution of displacements for trajectories lasting ≥ 3 frames

(Hansen et al., 2018) (Figure 1-figure supplement 2A,B). Hereafter, we refer to diffusive

values derived from Spot-On in the text. As previously reported for biological controls,

H2B histone (Halo-H2B) and free HaloTag (Halo-NLS, nuclear localization signal), exhibit

two distinct, well-separated diffusion states representing chromatin-bound and chromatin-

free molecules (Ranjan et al., 2020). We found that the majority of H2B molecules (79.4

$\pm$ 1.9%) are slow-moving with average D of 0.026 $\mu m^2 s^{-1}$ (Figure 1-figure supplement 2A)

consistent with incorporation into chromatin, whereas most of the chromatin-free Halo-

NLS molecules show greatly increased diffusivity (D ~ 5 $\mu m^2 s^{-1}$) (Ranjan et al., 2020).

Compared to H2B, chromatin remodelers exhibit a slow D_{bound} fraction (average 0.036 $\pm$

0.007 to 0.067 $\pm$ 0.004 $\mu m^2 s^{-1}$) as would be expected for molecules associated with

largely immobile chromatin (Figure 1-figure supplement 2A). However, as discussed later,

the D_{bound} values are ~2-fold higher than H2B. Furthermore, we also observed a separable

chromatin-free fraction whose D_{free} values (0.464 $\pm$ 0.043 to 1.014 $\pm$ 0.024 $\mu m^2 s^{-1}$) are

~10-fold higher, but distinctly lower than the D_{free} for Halo-NLS, indicating that our imaging

regime captures essentially the full range of potential diffusive behaviors for this family.

In addition, the D_{free} values show an inverse correlation with the estimated total molecular

weights of chromatin remodeling complexes, consistent with expectations that the tagged

catalytic subunits are associated within larger complexes (Figure 1-figure supplement 1D).

Figure 1. Chromatin-binding and chromatin-free fractions of RSC, SWI/SNF, INO80, and ISW2.
Diffusion coefficient histograms show the chromatin-binding fraction is highest for RSC among gene promoter-acting remodelers. **(A)** Experimental scheme. **(B)** Fast-tracking imaging regime uses short exposures (10 ms) at high laser power to distinguish slow (chromatin-bound) and fast (chromatin-free) diffusing populations. **(C)** Slow-tracking regime directly observes the dwell times of chromatin-bound molecules using 250 ms exposures at low laser power. **(D-G)** Fast-tracking diffusion histograms for Sth1-Halo **(D)**, Snf2-Halo **(E)**, Ino80-Halo **(F)**, and Isw2-Halo **(G)**. Left: normalized histograms of $\log_{10}$ diffusion coefficients of single-molecule trajectories fitted to two Gaussian distribution functions (solid gray line: sum of two Gaussians; dashed lines: individual Gaussian curves representing chromatin-bound and chromatin-free populations). Histograms combined from 2 or 3 biological replicates are resampled 100 times by the bootstrap method for resampling errors. Right: Spot-On kinetic modeling results based on displacement distribution histograms. Solid colored bar with indicated value represents % chromatin-bound molecules; open bar represents % chromatin-free. Error bars are standard deviations from 2 or 3 biological replicates.

Figure supplement 1. Cell growth, integrity, and localization of HaloTagged remodeler subunits.
Figure supplement 2. Spot-On kinetic modeling analyses.
Source data 1. MSD-based kinetic analysis results.
Source data 2. Original gel images for Figure supplement 1A.

We next assessed how the fractions of chromatin-bound and chromatin-free molecules

vary among subgroups of chromatin remodeling enzymes. RSC and SWI/SNF mobilize

+1 and -1 nucleosomes to increase promoter accessibility, while INO80 and ISW2

mobilize them to reduce accessibility (Hartley & Madhani, 2009; Kubik et al., 2019). We

found that the majority of both RSC and SWI/SNF molecules are associated with

chromatin (RSC: 66.0 ± 1.1%; SWI/SNF: 55.9 ± 1.3%) (Figure 1D,E). INO80 and ISW2

exhibit F_{bound} values of 48.3 ± 0.2% and 34.8 ± 0.4%, respectively (Figure 1F,G). Overall,

these NDR-acting remodelers display a broad range of chromatin-binding fractions

(inclusive of stable and transient binding), with RSC showing the highest overall

chromatin binding.

Figure 2. Chromatin-binding and chromatin-free populations of CHD1 and ISW1.
(A-B) Diffusion coefficient histograms and Spot-On analysis as described in Figure 1 for the catalytic subunits Chd1-Halo (A) and Isw1-Halo **(B)**. **(C-D)** Diffusion coefficient histograms and Spot-On analysis of the accessory subunits of ISW1a and ISW1b complexes: Ioc3-Halo **(C)** and Ioc4-Halo **(D)**.

CHD1 and ISW1 act primarily on nucleosomes located in the gene body (Kubik et al.,

2019; Ocampo et al., 2016). The two remodelers show comparable F_{bound} values (CHD1:

47.8 $\pm$ 4.9%; ISW1a/b: 52.0 $\pm$ 1.1%) (Figure 2A,B). However, the catalytic subunit Isw1

is shared by two distinct chromatin remodeling complexes called ISW1a and ISW1b (Vary

et al., 2003), in addition to potentially un-complexed Isw1 catalytic subunit (Tsukiyama,

Palmer, Landel, Shiloach, & Wu, 1999). The ISW1a complex localizes near the

transcription start and end of genes, whereas the ISW1b complex occupies more mid-

coding regions (Morillon et al., 2003; Smolle et al., 2012b; Yen et al., 2012). Since Isw1

catalytic subunit dynamics represent a composite of the two remodeling complexes, we

also tagged Ioc3 and Ioc4 accessory subunits unique to ISW1a and ISW1b complexes,

respectively. The gene-body acting ISW1b (Ioc4-Halo) complex exhibits higher F_{bound}

compared to ISW1a (Ioc3-Halo) complex (ISW1b: 55.5 $\pm$ 2.6%; ISW1a: 39.6 $\pm$ 0.0%)

(Figure 2C,D).

Chromatin remodelers frequently transition between bound and free states

Unlike the two well-separated Gaussian distributions for H2B histones and HaloTag

protein (Figure 3A), the Log(D) histograms of all imaged chromatin remodelers display

less distinct bound and free populations, with a noticeable fraction showing an

intermediate range of diffusion coefficients (Figure 1D-G, Figure 2). This population could

either represent remodeler complexes transitioning between chromatin-bound and

chromatin-free states, or chromatin-free molecules of intermediate diffusivity due to

association with additional factors or confined inside a subnuclear compartment (Hansen,

Amitai, Cattoglio, Tjian, & Darzacq, 2020; Izeddin et al., 2014; McSwiggen et al., 2018;

Strom et al., 2017).

To distinguish between these possibilities, we analyzed single-particle trajectories using

vbSPT, a variational Bayesian Hidden Markov Model (HMM) algorithm, which models

state kinetics and assigns diffusive states to each displacement (Persson, Lindén,

Unoson, & Elf, 2013). We classified every displacement as either State 1 ('bound') or

State 2 ('free') (Figure 3-source data 1), and sub-classified all trajectories as bound, free,

or transitioning (Figure 3B). The median bound and free displacement lengths between

transitioning and non-transitioning trajectories are highly similar or identical for each

remodeler, validating the vbSPT state assignments and essentially excluding a dominant

It is striking that the population of transitioning trajectories is more prominent for

remodelers (from $20.9 \pm 2.5\%$ to $30.2 \pm 1.5\%$) compared to free HaloTag ($10.4 \pm 0.7\%$)

and H2B histone ($7.1 \pm 2.1\%$), (Figure 3C-I, Figure 3-figure supplement 1D-E). We

observed comparable frequencies for remodeler dissociation (bound to free transition:

45.3 ± 1.3 to 50.24 ± 0.01 %) and association (free to bound: 49.76 ± 0.01 to $54.7 \pm 1.3\%$),

indicating that there is little bias in the direction of state transitions (Figure 3-figure

supplement 1F). Furthermore, the frequent detection of state transitions over short

trajectory lifetimes suggests that the duration of each state is short-lived. We concluded

that transient but frequent chromatin interactions are characteristic of the six remodeling

complexes.

All remodelers have remarkably short *in vivo* residence times of 4-7 s

The chromatin-bound remodeler population measured by fast tracking consists of both

transiently and stably bound molecules. We acquired long-exposure movies [250

ms/frame] under slow tracking (Chen et al., 2014) to generate survival curves revealing

the apparent dissociation of chromatin-bound molecules as a function of time (Figure 1C).

Particle dissociation can be due to molecules truly disengaging from chromatin, or to

fluorophore photobleaching and chromatin movements out of focus, which can corrected

using the survival curve of H2B histone as a standard (Hansen, Pustova, Cattoglio, Tjian,

& Darzacq, 2017). The remodeler survival plots fit well to a double exponential decay

model (Figure 4-figure supplement 1A-F,H), from which the average lifetimes (τ_{sb}, τ_{tb}) and

fractions (f_{sb}, f_{tb}) of stable-binding and transient-binding species were extracted (Figure

4). All τ values presented in the text and figures are corrected based on H2B decay

kinetics.

The stable-binding subpopulations (f_{sb}) of RSC ($27 \pm 2\%$) and SWI/SNF ($24 \pm 6\%$) display

strikingly short lifetimes (RSC: 5.0 ± 0.7 s; SWI/SNF 4.4 ± 1.2 s) (Figure 4A,B), consistent

with a previous measurement for the Rsc2 subunit of RSC (Mehta et al., 2018). Similarly,

INO80 and ISW2 exhibit stable-binding fractions (f_{sb} $20 \pm 3\%$ and $13 \pm 3\%$, respectively)

and similarly short residence times (τ_{sb} 3.7 ± 0.8 s and 4.9 ± 2.2 s, respectively) (Figure

4C,D). Hence, all NDR-acting remodelers bind stably for less than 5 s in live yeast,

whereas transient-binding populations are more short-lived by almost an order of
magnitude (Figure 4-source data 1).
261

Figure 4. All remodelers have short-lived stable-binding residence times of 4-7 s.
(A-F) Fitted double exponential decay curves from 1-CDF plots of observed dwell times from individual binding events (n) imaged by slow-tracking, for Sth1-Halo **(A)** Snf2-Halo **(B)**, Ino80-Halo **(C)**, and Isw2-Halo **(D)**, Chd1-Halo **(E)**, and Isw1-Halo **(F)**. Solid colored and dashed black fitted curves for indicated remodelers and H2B, respectively. Pie charts show the percentage (f_{sb}) and average residence time (τ_{sb}) of the stable binding population after photobleaching correction. Errors represent bootstrap resampling errors after resampling 100 times (sb: stable-binding; tb: transient-binding).

Figure supplement 1. Survival plots [1-CDF] of dwell times showing 1- vs 2-component exponential decay fits.
Source data 1. Kinetic parameters determined by Slow-tracking.

For gene body-acting remodelers, CHD1 and ISW1 complexes exhibit stable-binding
fractions (f_{sb} 15 ± 3% and 19 ± 2%, respectively) and short dwell times (τ_{sb} 7.2 ± 3.3 s and
5.6 ± 1.3 s, respectively) (Figure 4E,F). Interestingly, ISW1b shows 2.5-fold higher
residence times compared to ISW1a (τ_{sb} 5.9 ± 2.5 s and 2.2 ± 1.0 s, respectively) with
comparable stable-binding fractions (Figure 4-figure supplement 1G). These remodelers
also exhibit very short transient-binding residence times (τ_{tb}< 0.65 s). Hence, the majority
of chromatin binding events by remodelers is transient, and stable binding, on the order
of several seconds, is notably short-lived.

Figure 5. ATP hydrolysis is responsible for rapid chromatin dissociation.
(A) Bar diagram and cartoons for remodelers mutated in the 'Walker A' and 'Walker B' motifs, respectively.
(B) Representative 3D plots of trajectories imaged by slow-tracking for wildtype (Chd1-Halo, black) and ATPase-dead mutant (Chd1K407R-Halo, red). Each plot shows all trajectories (≥ 3 frames) from single nucleus where lines represent apparent durations of chromatin-binding events. **(C-F)** 1-CDF plot, pie chart, and residence times of wildtype (top) and ATPase-dead mutants (bottom) for Isw2 **(C)**, Isw1 **(D)**, and Chd1 **(E,F)**.

Figure supplement 1. Expression levels and 1-CDF plots for wildtype and mutant ATPase-dead Isw2D312N.
Source data 1. Slow-tracking results for ATPase-dead mutants.
Source data 2. Original gel images for Figure supplement 1A.

 ATPase activity is coupled to fast dissociation rates

 To examine whether the measured dissociation kinetics are intrinsic to chromatin

 remodeling complexes or functionally related to their ATP-dependent remodeling

 activities, we made strains harboring a point mutation in the ATPase domains of Isw2,

Isw1, and Chd1; these mutations have previously been shown to abolish their ATPase
activities (Isw2K215R, Isw1K227R, Chd1K407R, and Chd1D513N) (Figure 5A)
(Fitzgerald et al., 2004; Gelbart, Rechsteiner, Timothy, Tsukiyama, & Richmond, 2001;
Hauk, McKnight, Nodelman, & Bowman, 2010; Tsukiyama et al., 1999). We then acquired
slow-tracking movies to compare the dwell times of mutant to those of wildtype
remodeling enzymes (Figure 5B)
283
We found that the stable-binding average residence time increased by ~2-fold (from 4.9
$\pm$ 2.2 to 9.7 $\pm$ 3.1 s) for the Isw2K215R mutant (Figure 5C). Similarly, we observed
increased residence time (from 5.6 $\pm$ 1.3 to 9.1 $\pm$ 3.6 s) for the Isw1K227R (Figure 5D).
The two ATPase-dead Chd1 mutants both showed increased stable-binding residence
times (Chd1K407R from 7.2 $\pm$ 3.3 to ~52 s; Chd1D513N from 7.2 $\pm$ 3.3 to 16.1 $\pm$ 7.3 s)
(Figure 5E,F). Interestingly, the tail of the Chd1K407R survival curve approaches that of
H2B, which indicates its longevity, but precludes precise determination of dwell time
(Figure 5-figure supplement 1B). All four mutants (Figure 5-source data 1) exhibit little to
no changes in the transient-binding residence times compared to wildtype. In all, our
results indicate that after chromatin association, the mutant ATPases exhibit slower
dissociation rate (the reciprocal of residence time), consistent with previous genome-wide
ChIP and biochemical studies (Fitzgerald et al., 2004; Gelbart et al., 2001).
296

ATP binding enhances chromatin-bound mobility of remodelers
Chromatin imaged by several distinct methods in living cells displays heterogeneous
mobility, which is dependent on its compaction state, subnuclear localization, and ATP-
dependent processes (Gasser, 2002; Gu et al., 2018; Marshall et al., 1997; Soutoglou &
Misteli, 2007). Remodelers may undergo 1D translocation on DNA (Sirinakis et al., 2011),
and alter either local chromatin movement (Basu et al., 2020; Neumann et al., 2012) or
higher-order chromatin structure (Lusser, Urwin, & Kadonaga, 2005; Maier, Chioda,
Rhodes, & Becker, 2008) in an ATP-dependent fashion. We assessed the diffusive
behavior of the chromatin-bound fraction of remodelers relative to the average dynamics
of incorporated Halo-H2B histone. From each trajectory classified as bound by vbSPT,
the apparent D value and the R_c [radius of confinement] were calculated to characterize

its diffusivity and the confined domain encompassing the observed trajectory, respectively
(Lerner et al., 2020). Importantly, chromatin-bound remodelers exhibit ~2-fold higher
mobility than H2B histone, as revealed by the average MSD plot and the distribution of
individual D values of each trajectory under fast-tracking (Figure 6A,B). The mean R_c
values are also substantially higher for remodelers compared to the global mean
measured for H2B (Figure 6-figure supplement 1A). This is further supported by the
higher apparent D values to varying degrees [2- to 4-fold] of stably-bound remodelers
measured by slow-tracking (Figure 6C). Such greater mobility of chromatin-bound
remodelers may be due to the combined effects of remodeler diffusion on chromatin and
movement of the chromatin fiber caused by remodeling activity, or alternatively, may
reflect the intrinsic dynamics of genomic loci being targeted.
319
To distinguish between these two alternatives, we measured the chromatin-associated
mobility of the four aforementioned ATPase-dead mutants. Three mutants Isw2K215R,
Isw1K227R, Chd1K407R harboring substitutions in the catalytic ATPase Walker A motif
responsible for ATP binding display strikingly lower diffusivity as revealed by the average
MSD plot of stably bound molecules, which approaches or substantially overlaps the
global H2B curve (Figure 6D-F). This is supported by the violin plots of individual D values
for stably bound trajectories (Figure 6G). Surprisingly, Chd1D513N bearing a substitution
in the Walker B motif of Chd1 shows no substantial changes in the average MSD curve
and apparent D values for stably bound trajectories compared to wildtype as measured
by slow-tracking (Figure 6F,H). To rule out Chd1-specific effects, we also made a strain
harboring the corresponding D-to-N substitution in ISW2 (Isw2D312N), and found no
substantial changes in the average MSD curve and apparent D values compared to
wildtype (Figure 6D,H). As expected for a mutation in the catalytic ATPase, the
Isw2D312N mutant exhibits an approximately two-fold increase in its residence time
compared to wildtype (Figure 5-figure supplement 1C). Because the Walker B motif is
important for ATP hydrolysis (via coordinating Mg^{2+} ion and a water molecule), but not for
ATP binding (Singleton, Dillingham, & Wigley, 2007; Walker, Saraste, Runswick, & Gay,
1982) (Figure 5A), this result suggests that the ATP-bound state may be adequate to

induce enhanced diffusion on chromatin as part of the mechanism of target search by

remodeling enzymes.

Figure 6. ATP utilization is responsible for enhanced mobility of chromatin-bound remodeler.
(A-B) Average MSD plot **(A)** and violin plot **(B)**, of individual D values for 'bound only' trajectories imaged by fast-tracking, shown for six remodelers and H2B histone. **(C)** Violin plot showing distribution of individual D values imaged by slow-tracking for six remodelers and H2B histone. For **A-C**, each wildtype remodeler is compared to H2B by the ordinary one-way ANOVA test (****$p<0.0001$, ***$p<0.001$, **$p<0.01$, *$p<0.05$). **(D-H)** MSD plot **(D-F)** and violin plot **(G,H)** of individual D values for trajectories imaged by slow-tracking for wildtype, ATPase-dead mutant, and H2B. For violin plots, thick red and dotted gray lines represent the median and two quartiles, respectively. For **D-H**, mutants are compared to wildtype by the unpaired t test (****$p<0.0001$, ***$p<0.001$, ns: not significant). **(I)** Representative trajectories imaged by slow-tracking for H2B and remodelers. H2B displays low mobility, whereas remodelers display higher chromatin-associated diffusivity that is enhanced by ATP utilization.

Figure supplement 1. Chromatin-bound remodelers display higher radius of confinement (Rc) values than H2B.
Source data 1. Number of molecules (N), statistical tests, and source data for Figure 6.

Promoter-enriched remodelers have robust chromatin occupancies

Chromatin remodelers are key regulators of the +1 nucleosome position genome-wide, whose accurate location is crucial for the PIC (pre-initiation complex) formation and TSS fidelity (Lai & Pugh, 2017b; Zhang et al., 2011). RSC and SWI/SNF mobilize the +1 nucleosome away from the NDR, opposed by INO80 and ISW2 activities, which slide the +1 nucleosome towards the NDR. As a quantitative indicator of nucleosome engagement, we determined the occupancies of the four remodelers, i.e. the percent average occupancy at a chromatin target by each remodeler over a given time period. To calculate temporal occupancy, we utilized the measured overall chromatin-binding fraction [F_{sb}] and the temporal parameters for stable [τ_{sb}, f_{sb}] and transient [τ_{tb}, f_{tb}] chromatin-binding (Figure 7A). Here, we assume that stable binding, which is almost an order of magnitude longer than transient binding, represents binding at 'specific' target sites within promoter regions including -1, +1 nucleosomes and the intervening NDR, while transient binding represents non-specific chromatin interactions (Ball et al., 2016).

The fraction of stable-binding depends on both the number of molecules per nucleus ($N_{molecules}$) and number of the target sites in the genome ($N_{targets}$) (Chen et al., 2014). For $N_{molecules}$, we used published values compiled from orthogonal approaches to determine protein abundance in *S. cerevisiae* (Ho, Baryshnikova, & Brown, 2018). As regards the number of chromatin targets ($N_{targets}$), many studies have investigated the genome-wide specificities of chromatin remodelers using multiple approaches including ChIP-Seq (Cutler, Lee, & Tsukiyama, 2018; Floer et al., 2010; Shimada et al., 2008; Spain et al., 2014), MNase-ChIP (Yen et al., 2012), ChIP-exo (Rossi et al., 2021), Native-ChIP-Seq (Ramachandran, Zentner, & Henikoff, 2015; Zentner, Tsukiyama, & Henikoff, 2013), CUT&RUN (Brahma & Henikoff, 2019), and ChEC-seq (Kubik et al., 2019). For this paper, we utilized the $N_{targets}$ values reported by Kubik *et al.*, who investigated the binding sites for four yeast remodelers of interest in this study, and explicitly quantified the number of mRNA gene promoters enriched for each remodeler. Assuming that the +1 or -1 nucleosomes and the NDR together represent the main interaction substrates at promoter regions, this value of $N_{targets}$ can be considered as a lower bound estimate. Accordingly,

371 RSC binding is the most widespread ($N_{targets}$= 3702), SWI/SNF binds only a small subset

372 ($N_{targets}$= 466), and INO80 ($N_{targets}$= 1646) and ISW2 ($N_{targets}$= 1802) each bind to

373 approximately a third of all yeast promoters (Kubik et al., 2019).

Figure 7. Remodelers show substantial temporal occupancies at chromatin targets.
(A) Key parameters measured in this study and acquired from the literature (Ho et al., 2018; Kubik et al., 2019) are used to calculate occupancy levels for gene promoter-acting remodelers. **(B)** Time trace simulations of temporal occupancy for individual remodelers at a target promoter region based on average τ_{sb} and sampling interval. Top and bottom bars represent occupied (on) and vacant (off) states, respectively, and vertical lines depict transitions between the two states. **(C)** Time trace simulations of occupancy at a RSC- and INO80-bound promoter region based on average τ_{sb} and sampling interval. Individual time trace simulations are shown above, and the cumulative simulated occupancy plot (black) shows either one or both remodelers bound in the time course of 500 s.

Figure supplement 1. Time trace simulations of temporal occupancies at promoters bound by multiple remodelers, and analysis of CHD1 DNA-binding mutant.

To calculate occupancy values, we used the τ_{search} (search time), the time it takes for a

molecule to go from one stable target site to the next [i.e. time bound non-specifically plus

time in free diffusion], the SI (sampling interval) [i.e. the time between initial binding of

one molecule and binding of the second molecule], and the estimated values for $N_{molecules}$

and $N_{targets}$ per cell (Figure 7A; see Methods). RSC shows substantial occupancy (32 $\pm$

7.5%) at stably bound chromatin targets, despite its short residence time (τ_{sb}) of 5.0 $\pm$ 0.7

s. Thus, rather than individual RSC molecules residing for long periods of time, the high

occupancy rate can be attributed to the short τ_{search} (15 $\pm$ 2.6 s) and comparable SI (16 $\pm$

3.1 s) values coupled to high $N_{molecules}$ (>4000) (Figure 7A). SWI/SNF exhibits the highest

occupancy (93 $\pm$ 41%) among the four remodelers, consistent with the highest raw ChEC

signals reported for Swi3 (Kubik et al., 2019). Strikingly, our estimate indicates that

SWI/SNF also maintains its occupancy at target sites by coupling short residence time

(4.4 $\pm$ 1.2 s) with short SI (4.7 $\pm$ 1.0 s) (Figure 7A).

We next assessed the occupancy values for INO80 and ISW2, which oppose the actions

of RSC and SWI/SNF. Comparable to RSC, INO80 displays substantial occupancy (32 $\pm$

9.2%) at its targets while ISW2 displays a lower occupancy (11 $\pm$ 5.0%). INO80 exhibits

short τ_{search} (23 $\pm$ 6.0 s) and SI (16 $\pm$ 4.3 s) values, while ISW2 has a relatively longer

τ_{search} (61 $\pm$ 30 s) and SI (47 $\pm$ 22 s) values (Figure 7A). Average time trace simulations

of stably bound occupancies for each of the four remodelers over several hundred

seconds (Figure 7B) show that at promoter regions targeted by multiple remodelers such

as genes in 'cluster IV' enriched for RSC and INO80 (Kubik et al., 2019), the occupancy

by any one remodeler is strikingly high, and more than one remodeler can simultaneously

engage a promoter repeatedly over several minutes (Figure 7C; see also Figure 7-figure

supplement 1A,B).

Figure 8. Nucleosome remodeling cycle at NDRs.
Model for nucleosome remodeling cycle at a gene promoter region targeted by RSC and INO80. The promoter region transitions between remodeler-occupied [solid arrow] and -vacant [dashed arrow] states, and their durations are indicated. After association of RSC or INO80 to the NDR, remodelers undergo 1-D diffusion on chromatin in an ATP-dependent manner, resulting in higher chromatin-associated mobility. Upon engaging its nucleosome substrate [e.g. the +1 nucleosome], RSC and INO80 uses the energy of ATP hydrolysis to push or pull the nucleosome away from NDR, respectively. ATP hydrolysis facilitates remodeler dissociation, and the promoter region becomes vacant for other factor interactions. The order of remodeler visitation is arbitrary, and simultaneous co-occupancy within the NDR can occur infrequently (see text for details).

DISCUSSION

Imaging chromatin remodeler diffusion by the fast-tracking mode in yeast shows that they

bind to chromatin at substantial frequencies [F_{bound}: 35 - 66%], and with a notable

population [21 - 30%] displaying intermediate D values resulting from transitions between

bound and free states (Figure 1-3). This suggests remodelers frequently undergo highly

short-lived chromatin interactions and is consistent with our slow-tracking measurements

of transient-binding frequency (f_{tb}: 73 - 87%) (Figure 4), and with FRAP and FCS

measurements of over-expressed mammalian ISWI (Erdel, Schubert, Marth, Längst, &

Rippe, 2010). The high frequency of transient interactions, and direct evidence for
transitioning trajectories is also consistent with the model of 1D-3D facilitated diffusion, a
proposed mechanism to increase the target search efficiency of nuclear proteins (Von
Hippel & Berg, 1989),

By slow-tracking, two chromatin-associated populations, 'stable-binding' and 'transient-
binding', were observed for all six remodelers. Previous SMT studies on the mammalian
Sox2 and yeast Ace1 transcription factors showed that stable-binding subpopulation
represent interactions with cognate target sequences (Chen et al., 2014; Mehta et al.,
2018). Comparable to reported values for the Rsc2 subunit of RSC and the yeast
transcription factors, Ace1 and Gal4 (Donovan et al., 2019; Mehta et al., 2018), all imaged
remodelers show stable and transient residence times of 4-7 s and 0.4-0.7 s, respectively.
Furthermore, the effect of mutating the DNA-binding domain of CHD1 monomer (Ryan,
Sundaramoorthy, Martin, Singh, & Owen-Hughes, 2011; Tran, Steger, Iyer, & Johnson,
2000) results in a 3-fold reduction in the τ_{sb} value (from 7.2 ± 3.3 to 2.4 ± 0.7 s) (Figure 7-
figure supplement 1C,D). Unlike sequence-specific transcription factors, a complete loss
of stable-binding would not be expected for remodeling complexes, whose recruitment
relies on multiple interactions with gene-specific transcription factors, histone modification
recognition domains, and interaction with components of the transcription machinery
(Becker & Workman, 2013). Indeed, we speculate that the multiplicity of interaction motifs
has a central role in the unusual diffusive behaviors shown by chromatin remodelers.

Importantly, the fast dissociation rates of remodelers are facilitated by ATP hydrolysis.
Five tested ATPase-dead mutants (for ISW2, ISW1, CHD1) show two-fold or greater
increase in their stable-binding residence times (Figure 5C-F), highlighting a new role of
for ATP-utilization in coupling nucleosome remodeling to rapid enzyme dissociation from
chromatin. This also suggests that their mean residence times can reflect timescales for
the diverse reactions performed by remodeling enzymes on chromatin *in vivo*. Assuming
that the +1 or -1 nucleosomes and the NDR are the main targets for promoter-acting RSC,
SWI/SNF, INO80 and ISW2, their 4-7 s stable residence time would include time for
diffusion on the NDR as well as time expended for nucleosome remodeling. Biochemical

studies have shown that remodelers undertake small translocation steps with remodeling
rates of a few bp/sec (Blosser, Yang, Stone, Narlikar, & Zhuang, 2009; Deindl et al., 2013;
Harada et al., 2016; Qiu et al., 2018; Sabantsev, Levendosky, Zhuang, Bowman, & Deindl,
2019). For example, with an enzymatic rate of 2 bp/s for ISWI (Blosser et al., 2009), an
ISWI stable-binding event would allow octamer sliding by roughly 12 bp, which is within
range of *in vivo* nucleosome position changes after conditional inactivation of RSC,
SWI/SNF, INO80 and ISW2 (Ganguli et al., 2014; Kubik et al., 2019).

Under our imaging conditions, the chromatin-bound populations of the six remodelers
exhibit higher mobility than H2B measured for bulk incorporated histones (Figure 6A-C).
We further showed that this enhanced mobility is dependent on the ATPase domain.
Mutations in the ISW1, ISW2, and CHD1 Walker A motif implicated in nucleotide binding
(Singleton et al., 2007) substantially decreases in vivo mobility. Further analysis of
mutations in the CHD1 and ISW2 Walker B motif, implicated in ATP hydrolysis [but not
ATP binding] (Singleton et al., 2007), displays a milder decrease of its chromatin mobility,
suggesting that nucleotide binding is largely sufficient for promoting diffusion for the two
remodelers. Previous observations for other DNA-binding ATPase enzymes have noted
ATP binding-dependent, hydrolysis-independent 1D diffusion or sliding on DNA (Cho et
al., 2012; Mazur, Mendillo, & Kolodner, 2006; Tóth, Bollins, & Szczelkun, 2015),
suggesting that this mode of diffusion to enhance target search may be shared among
remodelers. In addition, RSC and Drosophila ISWI remodelers undergo ATP hydrolysis-
dependent translocation on ssDNA and dsDNA in vitro (Saha, Wittmeyer, & Cairns, 2005;
Whitehouse, Stockdale, Flaus, Szczelkun, & Owen-Hughes, 2003), with processivities of
20-70 bp/translocation event (Fischer, Saha, & Cairns, 2007; Saha et al., 2005; Sirinakis
et al., 2011; Whitehouse et al., 2003). Finally, the absence of any change in ISW1
chromatin-bound mobility upon treatment with a general transcription inhibitor thiolutin
rules out transcription *per* se as a source of enhanced remodeler diffusion (Figure 6-figure
supplement 1B). In all, our results suggest that chromatin remodelers use the catalytic
ATPase not only for nucleosome remodeling but also to enhance target search kinetics
by promoting 1D diffusion on chromatin and rapid detachment after reaction.

Yeast promoter regions can be classified into different groups enriched either for no
remodeler or a combination of RSC, SWI/SNF, INO80 and ISW2, with about half of
promoters genome-wide harboring at least two distinct remodelers that harbor
nucleosome pushing and pulling activities relative to the NDR (Kubik et al., 2019). At
promoter regions where opposing remodelers bind, we expect a consecutive "tug-of-war"
between the pushing and pulling activities, in which successive engagements would
ultimately result in fine-tuning the steady-state nucleosome position, with the final
outcome dependent on remodeler occupancy and nucleosome remodeling activity.
Based on occupancy estimates, two remodelers may be found to simultaneously engage
promoter chromatin (Figure 7C, Figure 7-figure supplement 1A,B), but steric
considerations likely preclude two remodelers binding to the same nucleosome or the
same face of a nucleosome. Alternatively, at promoter regions where none or only one
remodeler binds, other mechanisms are likely to have more substantial roles in
nucleosome positioning. These include the sequence-dependent bendability of promoter
DNA as well as the binding of general regulatory factors (GRFs), such as Reb1, Abf1,
and Rap1, acting as barriers to nucleosome mobility (Struhl & Segal, 2013).

A temporal model for nucleosome remodeling at NDRs

By integrating our live-cell SMT measurements with available genome-wide localization
and protein expression data, we estimate temporal occupancies ranging from 11 ± 5.0 to
$93 \pm 41\%$ for RSC, SWI/SNF, INO80, and ISW2 at target promoter regions including the
NDR and flanking nucleosomes. Our findings of highly dynamic and frequent remodeler-
nucleosome interactions are consistent with recent genomics studies showing substantial
changes in nucleosome positions upon rapid, conditional inactivation of remodelers in
yeast and mammalian systems (Iurlaro et al., 2021; Klein-Brill et al., 2019; Kubik et al.,
2019; Schick et al., 2021). Accordingly, we envision a 'nucleosome remodeling cycle' in
which remodeler combinations undergo frequent association, ATP-dependent
mobilization and dissociation from chromatin to dynamically fine-tune -1 and +1
nucleosome positions.

At the subset of genes targeted by RSC and INO80, we anticipate stochastic recruitment
e.g. of RSC (Figure 8). RSC recognizes general promoter characteristics, such as the
long DNA stretch of the NDR (Wagner et al., 2020), histone acetylation marks potentially
read by 8 bromodomains in four RSC subunits (Josling, Selvarajah, Petter, & Duffy, 2012),
and the Rsc3 DNA-binding sequence motif found in several hundred promoters (Badis et
al., 2008). Upon binding within the accessible NDR, RSC undergoes 1-D diffusion in an
ATP-dependent manner, manifesting higher chromatin-associated mobility. On
engagement with either flanking nucleosome substrate [+1 nucleosome shown], RSC
uses the energy of ATP hydrolysis to reposition the nucleosome away from NDR,
enlarging NDR length. Importantly, this remodeling activity facilitates RSC dissociation.
Subsequent stochastic recruitment of INO80, ATP-dependent 1-D diffusion, and
nucleosome engagement remodels the nucleosome to move in the opposing direction
and narrow the NDR, coupled with INO80 dissociation. Cycles of sequential or
simultaneous binding and activity by RSC and INO80 provides a dynamic temporal
window of accessibility for promoter chromatin. A similar tug-of-war is anticipated for other
combinations of opposing remodelers.

In a related study (Nguyen et al., 2020), the average promoter occupancy of the yeast
PIC that forms upstream and overlapping the +1 nucleosome was found to be in the range
of 10%, i.e. on the same order of magnitude but lower than three of four remodelers
examined. Similar to chromatin remodelers, a full PIC lasts only several seconds before
dissociation from chromatin, but the average promoter is vacant for ~100 seconds before
PIC reformation. Thus, we suggest that there may be robust and dynamic competition
between PIC components and mobilized NDR-flanking nucleosomes with chromatin
exposure of key promoter elements such as the TATA box occurring for only a limited
time window allowing proper assembly of downstream PIC components. This temporally
positioned +1 nucleosome would enable Pol II to scan and start transcription at the proper,
canonical TSS. In this way, the dynamic interactions of remodeling enzymes with their
promoter targets provides a temporal, chromatin accessibility-based regulatory
mechanism for eukaryotic transcription.

Taken together, our SMT study elucidates the dynamic behaviors of this family of nuclear
proteins and offers insights into additional kinetic functions for the remodeling ATPase
and the timescales that govern nucleosome repositioning in relation to transcription
events. Outstanding questions for future studies include determining the kinetic
parameters for other chromatin regulators such as histone acetyltransferases,
methyltransferases, histone de-modification enzymes, and histone chaperones, to gain a
comprehensive view of the overall competition for engagement of promoter-proximal
nucleosomes, their effects on nucleosome positioning and the dynamics of transcription
complexes. Our findings in live cells provide a temporal framework for further testing of
proposed models and should facilitate development of in vitro single-molecule assays that
allow direct observation of physical and functional interactions between transcription
regulators, chromatin, and the transcription machinery.

METHODS

Yeast strains

All *Saccharomyces cerevisiae* strains used in this study are isogenic derivatives of W303
strain carrying *pdr5∆* for efficient JF ligand labeling, and are listed in Supplementary file
1. HaloTag was fused to the C-terminus of the protein of interest using standard methods
for yeast transformation, using pBS-SK-Halo-NatMX plasmid (Ranjan et al., 2020). Point
mutations were introduced by either the traditional "pop-in pop-out" (Rothstein, 1991) or
the "50:50" method (Horecka & Davis, 2014), using pUG72 plasmid (P30117, Euroscarf,
Germany), and list of primers are provided in Supplementary file 2.

Yeast Growth Assays

The cell growth of strains carrying HaloTag fusion constructs were compared to their
derived parental genotype. The strains grown to saturation in YPAD (Yeast Extract-
Peptone-Dextrose + 40 mg/L Ade-SO$_4$) were serially diluted (5-fold dilutions) on YPAD
plates. Plates were imaged after 48 or 72 h growing at 3 different temperatures (25^{o}C,
30^{o}C, and 38^{o}C).

Cell Lysate Preparation to check integrity of HaloTag fusion proteins

Yeast cultures growing at early log phase (OD600 0.2) were treated with JF646 dye at a

saturating dye concentration of 20 nM JF646 (Grimm et al., 2015) was used instead of

JF552 for better dye labeling (Ranjan et al., 2020). The cell extract was prepared using

the NaOH method (Cold Spring Harbor Protocols). Total protein concentration was

measured using the Bradford Assay, and 45 ng of total protein was loaded per well in

SDS-PAGE. Gels were imaged on Tecan 5 scanner, with Cy5 excitation. After imaging,

gels were stained with Coomassie dye for loading control.

Yeast culture preparation for single molecule imaging

Yeast cultures growing in CSM+Ade (complete synthetic media) were treated with dyes

at early log phase (D600 0.2 - 0.3) for 2 h. For fast-tracking, saturating dye concentrations

ranging from 10 to 20 nM JF552 (Zheng et al., 2019) were used depending on factor

abundance. For slow-tracking, we used 5 - 7.5 nM JF552. In some instances, we also

added JF646 (~5 nM) to visualize nuclear fluorescence without JF552 excitation and to

partially reduce JF552 labelling. Cells growing in CSM+Ade medium were harvested

around mid-log phase by brief centrifugation (3500 rpm for 2 min), washed at least 3 times,

and finally resuspended in CSM medium. Resuspended cells were loaded on

Concanavalin A-treated coverslip (#1.5 Micro Coverglass -25 mm Diameter, Electron

Microscopy Sciences, Cat. No. 72225-01) assembled on imaging cell chamber (Invitrogen,

Cat. No. A7816), where coverslips were flamed prior to the treatment in order to reduce

single-to-noise background. After 5 min incubation time, the immobilized cells were

further washed 3-5 times with a final resuspension in 1 mL CSM.

Live cell, single molecule imaging using wide-field microscopy

Microscope setup: All yeast imaging was performed using a custom-built Zeiss widefield

microscope (Axio Observer Z1) with a 150X glycerin immersion objective (NA 1.35) as

previously described (Ranjan et al., 2020). Data was acquired with EM-CCD (Hamamatsu

C9100-13) camera with FF01-750/SP and NF03-405/488/561/635E quad-notch filters for

a final x-y pixel size of 107 nm. All imaging was performed with a single excitation channel.

For JF552 dye excitation, 555 nm laser (Crystalaser) at (TTL pulsed) with 561 beam-

splitter and 612/69 nm filter was used. For JF646 dye excitation, a 639 nm laser with 648

beamsplitter and 676/29 nm filter was used. Microscope manipulations (i.e. Z-focus, X/Y

translation, filter cube switch) was performed by Zen software (Zeiss, Germany) and

camera and data acquisition was controlled by HCImage software (Hamamatsu

Photonics, Japan).

Data acquisition

After yeast immobilization on the coverslip, cells were imaged for around 2 h at room

temperature.

Fast tracking: 10 ms frame-rate movies were recorded with continuous 555 nm laser

irradiation at ~1 kW/cm^2. A field of view of 128 x 128 pixels was used to capture 4-6 yeast

nuclei. About 40 movies each lasting about 1.5 min were acquired per imaging session,

and at least 2 biological replicates were obtained for each sample.

Slow tracking: 250 ms frame-rate movies were acquired using continuous 555 nm laser

irradiation at 0.05 kW/cm^2 for sufficient signal-to-noise while minimizing photobleaching.

A focal plane of 256 x 256 pixels was used to capture 15-20 yeast nuclei. In the beginning

of each movie, the 639 nm excitation channel was briefly used to fine-tune the focus, and

then immediately switched to 555 nm excitation to start data acquisition. Around 15-20

movies were taken per imaging session, with each movie lasting typically 7-9 min. At least

2 biological replicates were obtained for each sample.

Single molecule image analysis

For each raw movie, we first manually selected a 'substack' where ~1 single molecule per

nucleus per frame was observed in order to minimize tracking errors resulting from

connecting different molecules as one trajectory. Substack lengths of 5000 frames (50 s)

and 750 frames (3.125 min) were selected for fast and slow tracking movies, respectively,

using ImageJ (1.52p) custom-written script. The substacks were then applied to the

Diatrack software (ver. 3.05) to localize and track single particles(Vallotton & Olivier,

2013). For localization, the following parameters were applied: Remove dim: 75 - 85,

Remove blurred: 0.1, Activate High Precision mode: ON (HWHM=1 pixel). For tracking,

we used max jump of '6' (642 nm) and '3' pixels (321 nm) for fast and slow tracking

datasets, respectively. Furthermore, we masked the nuclear regions based on the

maximum intensity Z-projection of the selected substacks to filter out trajectories found
outside of the nucleus in the subsequent analysis steps. The Diatrack output file
containing information about the x, y coordinate and frame number were then applied for
further downstream analysis.

Fast tracking:

MSD-based diffusion coefficient histograms: All "masked" trajectories with at least 5
displacements were analyzed, using the lab custom-written R package, Sojourner
(https://rdrr.io/github/sheng-liu/sojourner/). Briefly, for each trajectory, MSD plot for time
lags from 2 to 5 $\triangle t$ ($\triangle t$ = 10 ms) were generated, then fit to linear regression (filtering out
$R^2 < 0.8$ plots). From the slope, the diffusion coefficient was calculated as (where d is the
number of dimensions, or 2):

638
$$D = 2d\frac{MSD}{dt}$$

Spot-On (Hansen et al., 2018): All "masked" trajectories with at least 2 displacements
were analyzed. The following parameters were applied for Jump Length Distribution: Bin
width (µm): 0.01, Number of time-points: 6, Jumps to consider: 4, Use entire trajectories
No, Max jump (µm): 2. Additionally, the following parameters were applied for 2-state
Model Fitting: D_{bound} (µm^2/s): 0.0005-0.1, D_{free} (µm^2/s): 0.15-25, F_{bound}: 0-1, Localization
error (µm): Fit from data (0.01-0.1), dZ (µm): 0.6, Use Z correction, Model Fit: CDF,
Iterations: 3.

vbSPT (variational Bayesian) HMM (Persson et al., 2013): All "masked" trajectories with
at least 2 displacements were analyzed. The following parameters were used to run
vbSPT-1.1.3 to classify each displacement into two states, "Bound" or "Free" (Hansen *et*
al., 2020, https://gitlab.com/anders.sejr.hansen/anisotropy): timestep= 0.01; dim= 2;
trjLmin= 2; runs= 3; maxHidden= 2; stateEstimate= 1; bootstrapNum=10; fullBootstrap=
0; init_D = [0.001, 16]; init_tD = [2, 20]*timestep; and default prior choices according to
(Persson et al., 2013).

Then each trajectory was sub-classified as "Bound only" if all displacements are classified

as bound state; "Free only" if all displacements are classified as free state; and

"Transitioning" if the trajectory contains both bound and free displacements with at least

two consecutive displacements in each state. To validate that the transitioning trajectories

consist of bound and free states, we calculated and compared the displacement length

between "bound only" and bound segments of transitioning trajectories, and between

"free only" and free segments of transitioning trajectories. Finally, the sub-classified

trajectories were used to regenerate the diffusion coefficient histograms.

Radius of confinement: All "masked" trajectories with at least 4 displacements were

analyzed, as described previously (Lerner et al., 2020). To determine the radius of

confinement exhibited by chromatin-bound molecules, we analyzed trajectories classified

as "bound only" by vbSPT (as described above). Since many confined trajectories with

low D do not pass the $R^2 \geq 0.8$ filter, we used all trajectories whose MSD plots passed

the more lenient $R^2 \geq 0.1$ filtering. The MSD plot was then fit to the circular confined

diffusion model:

672
$$MSD_{circle} = R^2 \cdot \left(1 - e^{\frac{-4 \cdot D \cdot t_{lag}}{R^2}}\right)$$

where R is the radius of confinement, D is the short-term diffusion coefficient. Specifically,

the first 10 time points of the MSD plot were used to fit to the model, and trajectories with

squared norm of residual (RSS) higher than 10^{-5} and Rc higher than 300 nm were

discarded.

Slow tracking:

Residence times: Using Sojourner package, the apparent lifetimes (temporal length of

trajectories) were determined for all "masked" trajectories lasting at least 3 frames. To

account for blinking or mislocalizations, we allowed for gaps up to 2 frames between two

localizations and linked them as one trajectory if they were less than 3 pixels apart. 1-

CDF curves were generated and fit to a double exponential decay model:

684
$$P(t) = f_{sb}\, e^{-k_{sb}t} + f_{tb}\, e^{-k_{tb}t}$$

where k_{sb} and k_{tb} correspond to dissociation rates for stable- and transient-binding events,

respectively, and $1 = f_{sb} + f_{tb}$ for the 2 components.

The apparent k_{sb} and k_{tb} values are affected by technical and imaging limitations such as

photobleaching and chromatin movements. To correct for this bias, we used apparent

dissociation rates of H2B imaged under same conditions as described previously(Hansen

et al., 2017). The corrected residence times for stable- (τ_{sb}) and transient binding (τ_{tb})

were calculated as follows:

692
$$\tau_{sb} = \frac{1}{k_{sb} - k_{sb,H2B}}$$

694
$$\tau_{tb} = \frac{1}{k_{tb} - k_{sb,H2B}}$$

Apparent diffusion coefficient values for stably bound trajectories: All "masked"

trajectories lasting at least 5 frames (not allowing for gaps) were analyzed, using

Sojourner package. For each trajectory, MSD plot for time lags from 2 to 5 Δt (Δt = 10

ms) were generated, then fit to linear regression (filtering out R^2 < 0.8 plots). From the

slope, the diffusion coefficient was calculated as (where d is the number of dimensions,

or 2):

702
$$D = 2d\frac{MSD}{dt}$$

Occupancy calculation

To calculate temporal occupancy, we integrated approaches from previous studies (Chen

et al., 2014; Loffreda et al., 2017; Tatavosian et al., 2018).

Search time (τ_{search}) is the average time it takes for a molecule to go from one specific site

to its next specific site. The two specific binding events (lasting for τ_{sb}) are interspersed

by a number of trials (N_{trials}) binding to non-specific sites (lasting for τ_{tb}). τ_{free} is the average

free time between 2 binding events. Assuming equal probability of binding to all specific

and non-specific sites, the search time is calculated as follows:

712
$$\tau_{search} = N_{trials} \times \tau_{tb} + (N_{trials} + 1) \times \tau_{free}$$

713   N_{trials} depends on the ratio of number of non-specific (N_{ns}) to specific sites (N_s), or r_s:

$$714 \qquad N_{trials} = \frac{N_s + N_{ns}}{N_s} = 1 + r_s$$

715 Here, r_s can be determined based on two assumed scenarios for bound molecules
716 observed during slow tracking (as described in Nguyen et al. 2020). First, f_{sb} determined
717 by slow tracking depends on the time a molecule spends bound to specific sites compared
718 to nonspecific sites:

$$719 \qquad f_{sb} = \frac{N_s \times \tau_{sb}}{N_s \times \tau_{sb} + N_{ns} \times \tau_{tb}} = \frac{\tau_{sb}}{\tau_{sb} + r_{s,1} \times \tau_{tb}}$$

720 Thus r_s is equal to:

$$721 \qquad r_{s,1} = \frac{\tau_{sb}}{\tau_{tb}} \times (\frac{1}{f_{sb}} - 1)$$

722 In the second scenario, f_{sb} depends on the probability that a free molecule binds to a
723 specific site over all sites:

$$724 \qquad f_{sb} = \frac{N_s}{N_s + N_{ns}} = \frac{1}{1 + r_{s,2}}$$

726 In this case r_s is:

$$727 \qquad r_{s,2} = \frac{1}{f_{sb}} - 1$$

728 We take the average value calculated from the 2 proposed scenarios to finally determine
729 r_s:

$$730 \qquad r_s = \frac{1}{2}(\frac{1}{f_{sb}} - 1)(\frac{\tau_{sb}}{\tau_{tb}} + 1)$$

731 In fast tracking, F_{bound} is percentage or fraction of the time a molecule spends bound to
732 chromatin either specifically or non-specifically:

$$733 \qquad F_{bound} = \frac{N_{trials} \times \tau_{tb} + \tau_{sb}}{N_{trials} \times \tau_{tb} + \tau_{sb} + (N_{trials} + 1) \times \tau_{free}}$$

734 Thus τ_{free} is (in terms of r_s):

$$735 \qquad \tau_{free} = \frac{\frac{(1 + r_s) \times \tau_{tb} + \tau_{sb}}{F_{bound}} - (1 + r_s) \times \tau_{tb} - \tau_{sb}}{2 + r_s}$$

Using the values derived for r_s and τ_{free}, we then calculated the search time as shown
above.

Sampling interval (SI) is the time interval between two specific binding events at a given
site as described previously (Chen et al., 2014):

$$Sampling\ Interval\ (SI) = \frac{(\tau_{search} + \tau_{sb}) \times N_{targets}}{N_{molecules}}$$

We used $N_{targets}$ values presented by (Kubik et al., 2019). $N_{molecules}$ was determined as the
median and standard error values (Ho et al., 2018), and their standard error was used for
error propagation.

Finally, occupancy is the temporal probability that a given specific site is occupied by the
protein of interest:

$$Occupancy = \frac{\tau_{sb}}{SI}$$

Target occupancy simulation

Remodeler occupancy at a target promoter region was simulated as described previously
(Nguyen et al. 2020). Briefly, experimentally determined τ_{sb} and estimated sampling
interval (SI) values were used to simulate sequential promoter-occupied and vacant
states over the time trace (500 s). The duration for each occupied and vacant state was
randomly chosen from exponential distributions of the average τ_{sb} and (SI- τ_{sb}) values,
respectively. For promoter regions targeted by multiple remodelers, each remodeler was
independently subject to the occupancy simulation, and the number of any single or
multiple remodeler(s) co-occupying each timepoint was calculated throughout the time
trace.

ACKNOWLEDGEMENTS

We thank Vu Q. Nguyen, Anand Ranjan, and Gaku Mizuguchi for experimental guidance
at the initial stages of this project, Sun Jay Yoo, Yick Hin Ling and Taibo Li for
computational assistance, Pascal Vallotton for support with Diatrack software, Jonathan
Lerner, Ken Zaret, and Melike Lakadamyali for assistance with the Two-parameter single-
molecule analysis, Slawomir Kubik and David Shore for advice on analysis of genomic
data on remodeling enzymes, Toshio Tsukiyama and Brad Cairns for yeast strains,
Anders Hansen and Greg Bowman for discussions and Wu lab members for helpful
comments. This study was supported by funds from a Korean Foundation for Advanced

Studies Fellowship (J.M.K.), a Johns Hopkins Bloomberg Distinguished Professorship
(C.W.) and National Institute of Health grant GM132290-01 (C.W.).
770

771 **AUTHOR CONTRIBUTIONS**
772 Conceptualization, J.M.K. and C.W.; Software, S.L., X.T., J.M.K., and T.L.; Investigation,
773 J.M.K., P.V., V.J., K.Y.L., and J.S.; Formal Analysis, J.M.K., P.V., and V.J.; Writing –
774 Original Draft, J.M.K. and C.W.; Writing – Review & Editing, J.M.K and C.W.; Funding
775 Acquisition, C.W.; Resources, Q.Z. and L.D.L.; Supervision, J.M.K. and C.W.
776

777 **COMPETING INTERESTS**
778 T.L. holds intellectual property rights related to Janelia Fluor dyes used in this publication.
779 L.D.L. and Q.Z. are listed as inventors on patents and patent applications whose value
780 might be affected by publication. The remaining authors declare no competing interests.
781

REFERENCES
Albert, I., Mavrich, T. N., Tomsho, L. P., Qi, J., Zanton, S. J., Schuster, S. C., & Pugh,
B. F. (2007). Translational and rotational settings of H2A.Z nucleosomes across the
Saccharomyces cerevisiae genome. *Nature*. https://doi.org/10.1038/nature05632
Almer, A., & Hörz, W. (1986). Nuclease hypersensitive regions with adjacent positioned
nucleosomes mark the gene boundaries of the PHO5/PHO3 locus in yeast. *The*
EMBO Journal, *5*(10), 2681–2687. https://doi.org/10.1002/j.1460-
2075.1986.tb04551.x
Badis, G., Chan, E. T., van Bakel, H., Pena-Castillo, L., Tillo, D., Tsui, K., … Hughes, T.
R. (2008). A Library of Yeast Transcription Factor Motifs Reveals a Widespread
Function for Rsc3 in Targeting Nucleosome Exclusion at Promoters. *Molecular Cell*,
32(6), 878–887. https://doi.org/10.1016/j.molcel.2008.11.020
Ball, D. A., Mehta, G. D., Salomon-Kent, R., Mazza, D., Morisaki, T., Mueller, F., …
Karpova, T. S. (2016). Single molecule tracking of Ace1p in Saccharomyces
cerevisiae defines a characteristic residence time for non-specific interactions of
transcription factors with chromatin. *Nucleic Acids Research*, *44*(21), 1–12.
https://doi.org/10.1093/nar/gkw744

Basu, S., Shukron, O., Ponjavic, A., Parruto, P., Boucher, W., Zhang, W., … Laue, E. D.
(2020). Live-cell 3D single-molecule tracking reveals how NuRD modulates
enhancer dynamics. *BioRxiv*, 2020.04.03.003178.
https://doi.org/10.1101/2020.04.03.003178

Becker, P. B., & Workman, J. L. (2013). Nucleosome remodeling and epigenetics. *Cold*
Spring Harbor Perspectives in Biology.
https://doi.org/10.1101/cshperspect.a017905

Blosser, T. R., Yang, J. G., Stone, M. D., Narlikar, G. J., & Zhuang, X. (2009). Dynamics
of nucleosome remodelling by individual ACF complexes. *Nature*, *462*(7276),
1022–1027. https://doi.org/10.1038/nature08627

Brahma, S., & Henikoff, S. (2019). RSC-Associated Subnucleosomes Define MNase-
Sensitive Promoters in Yeast. *Molecular Cell*.
https://doi.org/10.1016/j.molcel.2018.10.046

Chen, J., Zhang, Z., Li, L., Chen, B. C., Revyakin, A., Hajj, B., … Liu, Z. (2014). Single-
molecule dynamics of enhanceosome assembly in embryonic stem cells. *Cell*.
https://doi.org/10.1016/j.cell.2014.01.062

Cheung, V., Chua, G., Batada, N. N., Landry, C. R., Michnick, S. W., Hughes, T. R., &
Winston, F. (2008). Chromatin- and transcription-related factors repress
transcription from within coding regions throughout the Saccharomyces cerevisiae
genome. *PLoS Biology*. https://doi.org/10.1371/journal.pbio.0060277

Cho, W. K., Jeong, C., Kim, D., Chang, M., Song, K. M., Hanne, J., … Lee, J. B. (2012).
ATP alters the diffusion mechanics of MutS on mismatched DNA. *Structure*.
https://doi.org/10.1016/j.str.2012.04.017

Cutler, S., Lee, L. J., & Tsukiyama, T. (2018). Chromatin remodeling factors isw2 and
ino80 regulate chromatin, replication, and copy number of the saccharomyces
cerevisiae ribosomal DNA locus. *Genetics*.
https://doi.org/10.1534/genetics.118.301579

Deindl, S., Hwang, W. L., Hota, S. K., Blosser, T. R., Prasad, P., Bartholomew, B., &
Zhuang, X. (2013). ISWI remodelers slide nucleosomes with coordinated multi-
base-pair entry steps and single-base-pair exit steps. *Cell*, *152*(3), 442–452.
https://doi.org/10.1016/j.cell.2012.12.040

Donovan, B. T., Huynh, A., Ball, D. A., Patel, H. P., Poirier, M. G., Larson, D. R., …

Lenstra, T. L. (2019). Live-cell imaging reveals the interplay between transcription

factors, nucleosomes, and bursting. *The EMBO Journal*.

https://doi.org/10.15252/embj.2018100809

Erdel, F., Schubert, T., Marth, C., Längst, G., & Rippe, K. (2010). Human ISWI

chromatin-remodeling complexes sample nucleosomes via transient binding

reactions and become immobilized at active sites. *Proceedings of the National*

Academy of Sciences of the United States of America.

https://doi.org/10.1073/pnas.1003438107

Fischer, C. J., Saha, A., & Cairns, B. R. (2007). Kinetic model for the ATP-dependent

translocation of Saccharomyces cerevisiae RSC along double-stranded DNA.

Biochemistry. https://doi.org/10.1021/bi700930n

Fitzgerald, D. J., DeLuca, C., Berger, I., Gaillard, H., Sigrist, R., Schimmele, K., &

Richmond, T. J. (2004). Reaction cycle of the yeast Isw2 chromatin remodeling

complex. *EMBO Journal*. https://doi.org/10.1038/sj.emboj.7600364

Floer, M., Wang, X., Prabhu, V., Berrozpe, G., Narayan, S., Spagna, D., … Ptashne, M.

(2010). A RSC/nucleosome complex determines chromatin architecture and

facilitates activator binding. *Cell*. https://doi.org/10.1016/j.cell.2010.03.048

Ganguli, D., Chereji, R. V., Iben, J. R., Cole, H. A., & Clark, D. J. (2014). RSC-

dependent constructive and destructive interference between opposing arrays of

phased nucleosomes in yeast. *Genome Research*.

https://doi.org/10.1101/gr.177014.114

Gasser, S. M. (2002). Nuclear architecture: Visualizing chromatin dynamics in

interphase nuclei. *Science*. https://doi.org/10.1126/science.1067703

Gelbart, M. E., Rechsteiner, T., Timothy, J., Tsukiyama, T., & Richmond, T. J. (2001).

Interactions of Isw2 Chromatin Remodeling Complex with Nucleosomal Arrays :

Analyses Using Recombinant Yeast Histones and Immobilized Templates

Interactions of Isw2 Chromatin Remodeling Complex with Nucleosomal Arrays :

Analyses Using Recombinant Yeast Hi. *Molecular and Cellular Biology*, (June

2014). https://doi.org/10.1128/MCB.21.6.2098

Grimm, J. B., English, B. P., Chen, J., Slaughter, J. P., Zhang, Z., Revyakin, A., …

Lavis, L. D. (2015). A general method to improve fluorophores for live-cell and

862   single-molecule microscopy. *Nature Methods*. https://doi.org/10.1038/nmeth.3256

863   Gu, B., Swigut, T., Spencley, A., Bauer, M. R., Chung, M., Meyer, T., & Wysocka, J.

864   (2018). Transcription-coupled changes in nuclear mobility of mammalian cis-

865   regulatory elements. *Science*. https://doi.org/10.1126/science.aao3136

866   Hansen, A. S., Amitai, A., Cattoglio, C., Tjian, R., & Darzacq, X. (2020). Guided nuclear

867   exploration increases CTCF target search efficiency. *Nature Chemical Biology*,

868   *16*(3), 257–266. https://doi.org/10.1038/s41589-019-0422-3

869   Hansen, A. S., Pustova, I., Cattoglio, C., Tjian, R., & Darzacq, X. (2017). CTCF and

870   cohesin regulate chromatin loop stability with distinct dynamics. *ELife*, *6*.

871   https://doi.org/10.7554/eLife.25776.001

872   Hansen, A. S., Woringer, M., Grimm, J. B., Lavis, L. D., Tjian, R., & Darzacq, X. (2018).

873   Robust model-based analysis of single-particle tracking experiments with spot-on.

874   *ELife*. https://doi.org/10.7554/eLife.33125

875   Harada, B. T., Hwang, W. L., Deindl, S., Chatterjee, N., Bartholomew, B., & Zhuang, X.

876   (2016). Stepwise nucleosome translocation by RSC remodeling complexes. *ELife*,

877   *5*(FEBRUARY2016), 1–20. https://doi.org/10.7554/eLife.10051

878   Hartley, P. D., & Madhani, H. D. (2009). Mechanisms that Specify Promoter

879   Nucleosome Location and Identity. *Cell*. https://doi.org/10.1016/j.cell.2009.02.043

880   Hauk, G., McKnight, J. N., Nodelman, I. M., & Bowman, G. D. (2010). The

881   Chromodomains of the Chd1 Chromatin Remodeler Regulate DNA Access to the

882   ATPase Motor. *Molecular Cell*. https://doi.org/10.1016/j.molcel.2010.08.012

883   Heintzman, N. D., Stuart, R. K., Hon, G., Fu, Y., Ching, C. W., Hawkins, R. D., … Ren,

884   B. (2007). Distinct and predictive chromatin signatures of transcriptional promoters

885   and enhancers in the human genome. *Nature Genetics*.

886   https://doi.org/10.1038/ng1966

887   Ho, B., Baryshnikova, A., & Brown, G. W. (2018). Unification of Protein Abundance

888   Datasets Yields a Quantitative Saccharomyces cerevisiae Proteome. *Cell Systems*.

889   https://doi.org/10.1016/j.cels.2017.12.004

890   Horecka, J., & Davis, R. W. (2014). The 50:50 method for PCR-based seamless

891   genome editing in yeast. *Yeast (Chichester, England)*.

https://doi.org/10.1002/yea.2992

Iurlaro, M., Stadler, M. B., Masoni, F., Jagani, Z., Galli, G. G., & Schübeler, D. (2021).

Mammalian SWI/SNF continuously restores local accessibility to chromatin. *Nature*

Genetics. https://doi.org/10.1038/s41588-020-00768-w

Izeddin, I., Récamier, V., Bosanac, L., Cissé, I. I., Boudarene, L., Dugast-Darzacq, C.,

… Darzacq, X. (2014). Single-molecule tracking in live cells reveals distinct target-

search strategies of transcription factors in the nucleus. *ELife*, *2014*(3), 1–27.

https://doi.org/10.7554/eLife.02230

Josling, G. A., Selvarajah, S. A., Petter, M., & Duffy, M. F. (2012). The role of

bromodomain proteins in regulating gene expression. *Genes*, *3*(2), 320–343.

https://doi.org/10.3390/genes3020320

Klein-Brill, A., Joseph-Strauss, D., Appleboim, A., & Friedman, N. (2019). Dynamics of

Chromatin and Transcription during Transient Depletion of the RSC Chromatin

Remodeling Complex. *Cell Reports*. https://doi.org/10.1016/j.celrep.2018.12.020

Kubik, S., Bruzzone, M. J., Challal, D., Dreos, R., Mattarocci, S., Bucher, P., … Shore,

D. (2019). Opposing chromatin remodelers control transcription initiation frequency

and start site selection. *Nature Structural and Molecular Biology*.

https://doi.org/10.1038/s41594-019-0273-3

Kubik, S., Bruzzone, M. J., Jacquet, P., Falcone, J. L., Rougemont, J., & Shore, D.

(2015). Nucleosome Stability Distinguishes Two Different Promoter Types at All

Protein-Coding Genes in Yeast. *Molecular Cell*.

https://doi.org/10.1016/j.molcel.2015.10.002

Kubik, S., O'Duibhir, E., de Jonge, W. J., Mattarocci, S., Albert, B., Falcone, J. L., …

Shore, D. (2018). Sequence-Directed Action of RSC Remodeler and General

Regulatory Factors Modulates +1 Nucleosome Position to Facilitate Transcription.

Molecular Cell. https://doi.org/10.1016/j.molcel.2018.05.030

Lai, W. K. M., & Pugh, B. F. (2017a). Understanding nucleosome dynamics and their

links to gene expression and DNA replication. *Nature Reviews Molecular Cell*

Biology. https://doi.org/10.1038/nrm.2017.47

Lai, W. K. M., & Pugh, B. F. (2017b). Understanding nucleosome dynamics and their

links to gene expression and DNA replication. *Nature Reviews Molecular Cell*

Biology, *18*(9), 548–562. https://doi.org/10.1038/nrm.2017.47

Lerner, J., Gomez-Garcia, P. A., McCarthy, R. L., Liu, Z., Lakadamyali, M., & Zaret, K. S. (2020). Two-Parameter Mobility Assessments Discriminate Diverse Regulatory Factor Behaviors in Chromatin. *Journal of Cleaner Production*. https://doi.org/10.1016/j.molcel.2020.05.036

Lionnet, T., & Wu, C. (2021). Single-molecule tracking of transcription protein dynamics in living cells: seeing is believing, but what are we seeing? *Current Opinion in Genetics & Development*, *In press*. Retrieved from https://www.journals.elsevier.com/current-opinion-in-genetics-and-development

Loffreda, A., Jacchetti, E., Antunes, S., Rainone, P., Daniele, T., Morisaki, T., … Mazza, D. (2017). Live-cell p53 single-molecule binding is modulated by C-terminal acetylation and correlates with transcriptional activity. *Nature Communications*. https://doi.org/10.1038/s41467-017-00398-7

Lusser, A., Urwin, D. L., & Kadonaga, J. T. (2005). Distinct activities of CHD1 and ACF in ATP-dependent chromatin assembly. *Nature Structural and Molecular Biology*, *12*(2), 160–166. https://doi.org/10.1038/nsmb884

Maier, V. K., Chioda, M., Rhodes, D., & Becker, P. B. (2008). ACF catalyses chromatosome movements in chromatin fibres. *EMBO Journal*. https://doi.org/10.1038/sj.emboj.7601902

Marshall, W. F., Straight, A., Marko, J. F., Swedlow, J., Dernburg, A., Belmont, A., … Sedat, J. W. (1997). Interphase chromosomes undergo constrained diffusional motion in living cells. *Current Biology*. https://doi.org/10.1016/S0960-9822(06)00412-X

Mavrich, T. N., Ioshikhes, I. P., Venters, B. J., Jiang, C., Tomsho, L. P., Qi, J., … Pugh, B. F. (2008). A barrier nucleosome model for statistical positioning of nucleosomes throughout the yeast genome. *Genome Research*. https://doi.org/10.1101/gr.078261.108

Mazur, D. J., Mendillo, M. L., & Kolodner, R. D. (2006). Inhibition of Msh6 ATPase Activity by Mispaired DNA Induces a Msh2(ATP)-Msh6(ATP) State Capable of Hydrolysis-Independent Movement along DNA. *Molecular Cell*. https://doi.org/10.1016/j.molcel.2006.02.010

McSwiggen, D., Hansen, A., Marie-Nelly, H., Teves, S., Heckert, A., Dugast-Darzacq,

C., … Darzacq, X. (2018). Transient DNA Binding Induces RNA Polymerase II

Compartmentalization During Herpesviral Infection Distinct From Phase

Separation. *BioRxiv*. https://doi.org/10.1101/375071

Mehta, G. D., Ball, D. A., Eriksson, P. R., Chereji, R. V., Clark, D. J., McNally, J. G., &

Karpova, T. S. (2018). Single-Molecule Analysis Reveals Linked Cycles of RSC

Chromatin Remodeling and Ace1p Transcription Factor Binding in Yeast. *Molecular*

Cell. https://doi.org/10.1016/j.molcel.2018.09.009

Morillon, A., Karabetsou, N., O'Sullivan, J., Kent, N., Proudfoot, N., & Mellor, J. (2003).

Isw1 Chromatin Remodeling ATPase Coordinates Transcription Elongation and

Termination by RNA Polymerase II. *Cell*. https://doi.org/10.1016/S0092-

8674(03)00880-8

Neumann, F. R., Dion, V., Gehlen, L. R., Tsai-Pflugfelder, M., Schmid, R., Taddei, A., &

Gasser, S. M. (2012). Targeted INO80 enhances subnuclear chromatin movement

and ectopic homologous recombination. *Genes and Development*.

https://doi.org/10.1101/gad.176156.111

Ocampo, J., Chereji, R. V., Eriksson, P. R., & Clark, D. J. (2016). The ISW1 and CHD1

ATP-dependent chromatin remodelers compete to set nucleosome spacing in vivo.

Nucleic Acids Research. https://doi.org/10.1093/nar/gkw068

Ocampo, J., Chereji, R. V., Eriksson, P. R., & Clark, D. J. (2019). Contrasting roles of

the RSC and ISW1/CHD1 chromatin remodelers in RNA polymerase II elongation

and termination. *Genome Research*. https://doi.org/10.1101/gr.242032.118

Parnell, T. J., Schlichter, A., Wilson, B. G., & Cairns, B. R. (2015). The chromatin

remodelers RSC and ISW1 display functional and chromatin-based promoter

antagonism. *ELife*. https://doi.org/10.7554/eLife.06073

Persson, F., Lindén, M., Unoson, C., & Elf, J. (2013). Extracting intracellular diffusive

states and transition rates from single-molecule tracking data. *Nature Methods*.

https://doi.org/10.1038/nmeth.2367

Prajapati, H. K., Ocampo, J., & Clark, D. J. (2020). Interplay among atp-dependent

chromatin remodelers determines chromatin organisation in yeast. *Biology*, *9*(8), 1–

23. https://doi.org/10.3390/biology9080190

Qiu, Y., Levendosky, R., Chakravarthy, S., Patel, A., Myong, S., States, U., … States,

U. (2018). HHS Public Access, *68*(1), 76–88.

https://doi.org/10.1016/j.molcel.2017.08.018.The

Radman-Livaja, M., Quan, T. K., Valenzuela, L., Armstrong, J. A., van Welsem, T., Kim,

T. S., … Hartzog, G. A. (2012). A key role for Chd1 in histone H3 dynamics at the 3′

ends of long genes in yeast. *PLoS Genetics*.

https://doi.org/10.1371/journal.pgen.1002811

Ramachandran, S., Zentner, G. E., & Henikoff, S. (2015). Asymmetric nucleosomes

flank promoters in the budding yeast genome. *Genome Research*, *25*(3), 381–390.

https://doi.org/10.1101/gr.182618.114

Rando, O. J., & Winston, F. (2012). Chromatin and transcription in yeast. *Genetics*.

https://doi.org/10.1534/genetics.111.132266

Ranjan, A., Nguyen, V. Q., Liu, S., Wisniewski, J., Kim, J. M., Tang, X., … Wu, C.

(2020). Live-cell single particle imaging reveals the role of RNA polymerase II in

histone H2A.Z eviction. *ELife*. https://doi.org/10.7554/elife.55667

Rossi, M. J., Kuntala, P. K., Lai, W. K. M., Yamada, N., Badjatia, N., Mittal, C., … Pugh,

B. F. (2021). A high-resolution protein architecture of the budding yeast genome.

Nature. https://doi.org/10.1038/s41586-021-03314-8

Rothstein, R. (1991). Targeting, Disruption, Replacement, and Allele Rescue:

Integrative DNA Transformation in Yeast. *Methods in Enzymology*.

https://doi.org/10.1016/0076-6879(91)94022-5

Ryan, D. P., Sundaramoorthy, R., Martin, D., Singh, V., & Owen-Hughes, T. (2011). The

DNA-binding domain of the Chd1 chromatin-remodelling enzyme contains SANT

and SLIDE domains. *EMBO Journal*, *30*(13), 2596–2609.

https://doi.org/10.1038/emboj.2011.166

Sabantsev, A., Levendosky, R. F., Zhuang, X., Bowman, G. D., & Deindl, S. (2019).

Direct observation of coordinated DNA movements on the nucleosome during

chromatin remodelling. *Nature Communications*, *10*(1), 1–12.

https://doi.org/10.1038/s41467-019-09657-1

Saha, A., Wittmeyer, J., & Cairns, B. R. (2005). Chromatin remodeling through

directional DNA translocation from an internal nucleosomal site. *Nature Structural*

and Molecular Biology, *12*(9), 747–755. https://doi.org/10.1038/nsmb973

Schick, S., Grosche, S., Kohl, K. E., Drpic, D., Jaeger, M. G., Marella, N. C., … Kubicek,

S. (2021). Acute BAF perturbation causes immediate changes in chromatin

accessibility. *Nature Genetics*. https://doi.org/10.1038/s41588-021-00777-3

Shimada, K., Oma, Y., Schleker, T., Kugou, K., Ohta, K., Harata, M., & Gasser, S. M.

(2008). Ino80 Chromatin Remodeling Complex Promotes Recovery of Stalled

Replication Forks. *Current Biology*. https://doi.org/10.1016/j.cub.2008.03.049

Singleton, M. R., Dillingham, M. S., & Wigley, D. B. (2007). Structure and Mechanism of

Helicases and Nucleic Acid Translocases. *Annual Review of Biochemistry*, *76*(1),

23–50. https://doi.org/10.1146/annurev.biochem.76.052305.115300

Sirinakis, G., Clapier, C. R., Gao, Y., Viswanathan, R., Cairns, B. R., & Zhang, Y.

(2011). The RSC chromatin remodelling ATPase translocates DNA with high force

and small step size. *EMBO Journal*, *30*(12), 2364–2372.

https://doi.org/10.1038/emboj.2011.141

Smolle, M., Venkatesh, S., Gogol, M. M., Li, H., Zhang, Y., Florens, L., … Workman, J.

L. (2012a). Chromatin remodelers Isw1 and Chd1 maintain chromatin structure

during transcription by preventing histone exchange. *Nature Structural and*

Molecular Biology. https://doi.org/10.1038/nsmb.2312

Smolle, M., Venkatesh, S., Gogol, M. M., Li, H., Zhang, Y., Florens, L., … Workman, J.

L. (2012b). Chromatin remodelers Isw1 and Chd1 maintain chromatin structure

during transcription by preventing histone exchange. *Nature Structural and*

Molecular Biology. https://doi.org/10.1038/nsmb.2312

Soutoglou, E., & Misteli, T. (2007). Mobility and immobility of chromatin in transcription

and genome stability. *Current Opinion in Genetics and Development*.

https://doi.org/10.1016/j.gde.2007.08.004

Spain, M. M., Ansari, S. A., Pathak, R., Palumbo, M. J., Morse, R. H., & Govind, C. K.

(2014). The RSC Complex Localizes to Coding Sequences to Regulate Pol II and

Histone Occupancy. *Molecular Cell*. https://doi.org/10.1016/j.molcel.2014.10.002

Strom, A. R., Emelyanov, A. V., Mir, M., Fyodorov, D. V., Darzacq, X., & Karpen, G. H.

(2017). Phase separation drives heterochromatin domain formation. *Nature*.

https://doi.org/10.1038/nature22989

Struhl, K., & Segal, E. (2013). Determinants of nucleosome positioning. *Nature Structural and Molecular Biology*, *20*(3), 267–273. https://doi.org/10.1038/nsmb.2506

Tatavosian, R., Duc, H. N., Huynh, T. N., Fang, D., Schmitt, B., Shi, X., … Ren, X. (2018). Live-cell single-molecule dynamics of PcG proteins imposed by the DIPG H3.3K27M mutation. *Nature Communications*. https://doi.org/10.1038/s41467-018-04455-7

Tóth, J., Bollins, J., & Szczelkun, M. D. (2015). Re-evaluating the kinetics of ATP hydrolysis during initiation of DNA sliding by Type III restriction enzymes. *Nucleic Acids Research*. https://doi.org/10.1093/nar/gkv1154

Tran, H. G., Steger, D. J., Iyer, V. R., & Johnson, A. D. (2000). The chrome domain protein Chd1p from budding yeast is an ATP-dependent chromatin-modifying factor. *EMBO Journal*. https://doi.org/10.1093/emboj/19.10.2323

Tsukiyama, T., Palmer, J., Landel, C. C., Shiloach, J., & Wu, C. (1999). Characterization of the imitation switch subfamily of ATP-dependent chromatin-remodeling factors in Saccharomyces cerevisiae. *Genes and Development*. https://doi.org/10.1101/gad.13.6.686

Vallotton, P., & Olivier, S. (2013). Tri-track: Free software for large-scale particle tracking. *Microscopy and Microanalysis*. https://doi.org/10.1017/S1431927612014328

Vary, J. C., Gangaraju, V. K., Qin, J., Landel, C. C., Kooperberg, C., Bartholomew, B., & Tsukiyama, T. (2003). Yeast Isw1p forms two separable complexes in vivo. *Molecular and Cellular Biology*. https://doi.org/10.1128/MCB.23.1.80-91.2003

Von Hippel, P. H., & Berg, O. G. (1989). Facilitated target location in biological systems. *Journal of Biological Chemistry*.

Wagner, F. R., Dienemann, C., Wang, H., Stützer, A., Tegunov, D., Urlaub, H., & Cramer, P. (2020). Structure of SWI/SNF chromatin remodeller RSC bound to a nucleosome. *Nature*. https://doi.org/10.1038/s41586-020-2088-0

Walker, J. E., Saraste, M., Runswick, M. J., & Gay, N. J. (1982). Distantly related sequences in the alpha- and beta-subunits of ATP synthase, myosin, kinases and other ATP-requiring enzymes and a common nucleotide binding fold. *The EMBO*

Journal. https://doi.org/10.1002/j.1460-2075.1982.tb01276.x

Whitehouse, I., Rando, O. J., Delrow, J., & Tsukiyama, T. (2007). Chromatin

remodelling at promoters suppresses antisense transcription. *Nature*, *450*(7172),

1031–1035. https://doi.org/10.1038/nature06391

Whitehouse, I., Stockdale, C., Flaus, A., Szczelkun, M. D., & Owen-Hughes, T. (2003).

Evidence for DNA Translocation by the ISWI Chromatin-Remodeling Enzyme.

Molecular and Cellular Biology. https://doi.org/10.1128/mcb.23.6.1935-1945.2003

Wu, C. (1980). The 5' ends of drosophila heat shock genes in chromatin are

hypersensitive to DNase I. *Nature*. https://doi.org/10.1038/286854a0

Yen, K., Vinayachandran, V., Batta, K., Koerber, R. T., & Pugh, B. F. (2012). Genome-

wide nucleosome specificity and directionality of chromatin remodelers. *Cell*.

https://doi.org/10.1016/j.cell.2012.04.036

Yuan, G.-C., Liu, Y.-J., Dion, M. F., Slack, M. D., Wu, L. F., Altschuler, S. J., & Rando,

O. J. (2005). Genome-scale identification of nucleosome positions in

Saccharomyces cerevisiae. *Nature Methods*. https://doi.org/10.1038/nmeth0805-

567

Zentner, G. E., Tsukiyama, T., & Henikoff, S. (2013). ISWI and CHD Chromatin

Remodelers Bind Promoters but Act in Gene Bodies. *PLoS Genetics*, *9*(2),

e1003317. https://doi.org/10.1371/journal.pgen.1003317

Zhang, Z., Wippo, C. J., Wal, M., Ward, E., Korber, P., & Pugh, B. F. (2011). A packing

mechanism for nucleosome organization reconstituted across a eukaryotic

genome. *Science*. https://doi.org/10.1126/science.1200508

Zheng, Q., Ayala, A. X., Chung, I., Weigel, A. V., Ranjan, A., Falco, N., … Lavis, L. D.

(2019). Rational Design of Fluorogenic and Spontaneously Blinking Labels for

Super-Resolution Imaging. *ACS Central Science*.

https://doi.org/10.1021/acscentsci.9b00676

1104

FIGURE LEGENDS

Figure 1. Chromatin-binding and chromatin-free fractions of RSC, SWI/SNF, INO80, and

ISW2. Diffusion coefficient histograms show the chromatin-binding fraction is highest for

RSC among gene promoter-acting remodelers. **(A)** Experimental scheme. **(B)** Fast-

tracking imaging regime uses short exposures (10 ms) at high laser power to distinguish

slow (chromatin-bound) and fast (chromatin-free) diffusing populations. **(C)** Slow-tracking

regime directly observes the dwell times of chromatin-bound molecules using 250 ms

exposures at low laser power. (**D-G**) Fast-tracking diffusion histograms for Sth1-Halo (**D**),

Snf2-Halo (**E**), Ino80-Halo (**F**), and Isw2-Halo (**G**). Left: normalized histograms of $\log_{10}$

diffusion coefficients of single-molecule trajectories fitted to two Gaussian distribution

functions (solid gray line: sum of two Gaussians; dashed lines: individual Gaussian curves

representing chromatin-bound and chromatin-free populations). Histograms combined

from 2 or 3 biological replicates are resampled 100 times by the bootstrap method for

resampling errors. Right: Spot-On kinetic modeling results based on displacement

distribution histograms. Solid colored bar with indicated value represents % chromatin-

bound molecules; open bar represents % chromatin-free. Error bars are standard

deviations from 2 or 3 biological replicates.

Figure supplement 1. Cell growth, integrity, and localization of HaloTagged remodeler

subunits.

Figure supplement 2. Spot-On kinetic modeling analyses.

Source data 1. MSD-based kinetic analysis results.

1126

1127   **Figure 2.** Chromatin-binding and chromatin-free populations of CHD1 and ISW1. **(A-B)**

1128   Diffusion coefficient histograms and Spot-On analysis as described in Figure 1 for the

1129   catalytic subunits Chd1-Halo (**A**) and Isw1-Halo (**B**). **(C-D)** Diffusion coefficient

1130   histograms and Spot-On analysis of the accessory subunits of ISW1a and ISW1b

1131   complexes: Ioc3-Halo (**C**) and Ioc4-Halo (**D**).

1132

Figure 3. Remodelers undergo frequent transitions between bound and free states. **(A)**

Halo-H2B (brown) and Halo-NLS (pink) molecules display well-separated peaks in their

diffusion coefficient histograms. **(B)** An overview of displacement-based HMM

classification (vbSPT) to identify transitioning trajectories. After classifying each

displacement as either in bound or free state, each trajectory is sub-classified as 'bound

only', 'free only', or 'transitioning'. (**C-I)** Left: Overlay of raw histograms of $\log_{10}$ diffusion

coefficients for 'Bound only' (turquoise), 'Free only' (yellow), 'Transitioning' (purple), and

total trajectories (thin black). Right: Quantification (%) of transitioning trajectories in the
diffusion coefficient histogram, where errors represent standard deviation between 2 or 3
biological replicates. **(C)** Transitioning trajectories for Halo-H2B (top) and Halo-NLS
(bottom). (**D-I**) Transitioning trajectories for remodelers: Sth1-Halo (**D**), Snf2-Halo (**E**),
Ino80-Halo (**F**), and Isw2-Halo (**G**), Chd1-Halo (**H**), and Isw1-Halo (**I**).
Figure supplement 1. Validation of two diffusive states classified by vbSPT, and
quantification of transitioning frequencies.
Source data 1. vbSPT analysis results.
1148
Figure 4. All remodelers have short-lived stable-binding residence times of 4-7 s. **(A-F)**
Fitted double exponential decay curves from 1-CDF plots of observed dwell times from
individual binding events (n) imaged by slow-tracking, for Sth1-Halo (**A**) Snf2-Halo (**B**),
Ino80-Halo (**C**), and Isw2-Halo (**D**), Chd1-Halo (**E**), and Isw1-Halo (**F**). Solid colored and
dashed black fitted curves for indicated remodelers and H2B, respectively. Pie charts
show the percentage (f_{sb}) and average residence time (τ_{sb}) of the stable binding
population after photobleaching correction. Errors represent bootstrap resampling errors
after resampling 100 times (sb: stable-binding; tb: transient-binding).
Figure supplement 1. Survival plots [1-CDF] of dwell times showing 1- vs 2-component
exponential decay fits.
Source data 1. Kinetic parameters determined by Slow-tracking.
1160
Figure 5. ATP hydrolysis is responsible for rapid chromatin dissociation. **(A)** Bar diagram
and cartoons for remodelers mutated in the 'Walker A' and 'Walker B' motifs, respectively.
(B) Representative 3D plots of trajectories imaged by slow-tracking for wildtype (Chd1-
Halo, black) and ATPase-dead mutant (Chd1K407R-Halo, red). Each plot shows all
trajectories (≥ 3 frames) from single nucleus where lines represent apparent durations of
chromatin-binding events. **(C-F)** 1-CDF plot, pie chart, and residence times of wildtype
(top) and ATPase-dead mutants (bottom) for Isw2 (**C**), Isw1 (**D**), and Chd1 (**E,F**).
Figure supplement 1. Expression levels and 1-CDF plots for wildtype and mutant
ATPase-dead Isw2D312N.
Source data 1. Slow-tracking results for ATPase-dead mutants.

1171
Figure 6. ATP utilization is responsible for enhanced mobility of chromatin-bound
remodeler. (A-B) Average MSD plot (**A**) and violin plot (**B**), of individual D values for
'bound only' trajectories imaged by fast-tracking, shown for six remodelers and H2B
histone. **(C)** Violin plot showing distribution of individual D values imaged by slow-tracking
for six remodelers and H2B histone. For (**A-C**) each wildtype remodeler is compared to
H2B by the ordinary one-way ANOVA test (****$p<0.0001$, ***$p<0.001$, **$p<0.01$, *$p<0.05$).
(D-H) MSD plot (**D-F**) and violin plot (**G,H**) of individual D values for 'trajectories imaged
by slow-tracking for wildtype, ATPase-dead mutant, and H2B. For violin plots, thick red
and dotted gray lines represent the median and two quartiles, respectively. For **D-H**,
mutants are compared to wildtype by the unpaired t test (****$p<0.0001$, ***$p<0.001$, ns:
not significant). **(I)** Representative trajectories imaged by slow-tracking for H2B and
remodelers. H2B displays low mobility, whereas remodelers display higher chromatin-
associated diffusivity that is enhanced by ATP utilization.
Figure supplement 1. Chromatin-bound remodelers display higher radius of
confinement (Rc) values than H2B.
Source data 1. Number of molecules (N), statistical tests, and source data for Figure 6.
1188
Figure 7. Remodelers show substantial temporal occupancies at chromatin targets.
(A) Key parameters measured in this study and acquired from the literature (Ho et al.,
2018; Kubik et al., 2019) are used to calculate occupancy levels for gene promoter-acting
remodelers. **(B)** Time trace simulations of temporal occupancy for individual remodelers
at a target promoter region based on average τ_{sb} and sampling interval. Top and bottom
bars represent occupied (on) and vacant (off) states, respectively, and vertical lines depict
transitions between the two states. **(C)** Time trace simulations of occupancy at a RSC-
and INO80-bound promoter region based on average τ_{sb} and sampling interval. Individual
time trace simulations are shown above, and the cumulative simulated occupancy plot
(black) shows either one or both remodelers bound in the time course of 500 s.
Figure supplement 1. Time trace simulations of temporal occupancies at promoters
bound by multiple remodelers, and analysis of CHD1 DNA-binding mutant.
1201

Figure 8. Nucleosome remodeling cycle at NDRs. Model for nucleosome remodeling cycle at a gene promoter region targeted by RSC and INO80. The promoter region transitions between remodeler-occupied [solid arrow] and -vacant [dashed arrow] states, and their durations are indicated. After association of RSC or INO80 to the NDR, remodelers undergo 1-D diffusion on chromatin in an ATP-dependent manner, resulting in higher chromatin-associated mobility. Upon engaging its nucleosome substrate [e.g. the +1 nucleosome], RSC and INO80 uses the energy of ATP hydrolysis to push or pull the nucleosome away from NDR, respectively. ATP hydrolysis facilitates remodeler dissociation, and the promoter region becomes vacant for other factor interactions. The order of remodeler visitation is arbitrary, and simultaneous co-occupancy within the NDR can occur infrequently (see text for details).

SUPPLEMENTAL INFORMATION

Figure 1-figure supplement 1. Cell growth, integrity, and localization of HaloTagged remodeler subunits.

(A) SDS-PAGE gel scanned for JF646 dye fluorescence (top) and imaged after Coomassie staining (bottom). Cell lysates are prepared after treatment with JF646 at a saturating dye concentration (20 nM) for 2 h at 30°C. **(B)** Overlay of Phase Contrast image and 'nuclear glow' captured by initial JF552 dye excitation of yeast stained with JF552. **(C)** Five-fold dilutions of HaloTag fusion and wildtype strains are plated on YPAD plates at the indicated temperatures for 2-3 days. **(D)** Relationship between D_{free} values determined by Spot-On analysis and the calculated molecular weights of chromatin remodeling complexes.

Figure 1-figure supplement 2. Spot-On kinetic modeling analyses.

(A) Kinetic parameters (F_{bound}, D_{bound}, D_{free}) determined by Spot-On analyses. Errors represent standard deviation between 2 or 3 biological replicates. **(B)** Raw displacement histograms for individual biological replicates over the first 5 time frames (Δt: 10, 20, 30, 40, 50 ms). A two-state kinetic model was used for fitting the CDF [black lines] in Spot-On. F_{bound} values (shown above the histograms) are highly reproducible between replicates.

1233

Figure 3-figure supplement 1. Validation of two diffusive states classified by vbSPT, and quantification of transitioning frequencies.

(A) Pipeline for classification and sub-classification of trajectories using vbSPT. Individual displacement lengths were determined for 'bound only', 'free only', and 'transitioning'- subclassified trajectories to validate the two states and state transitions. **(B-C)** Violin plot (**B**) and median value (**C**) of individual displacement lengths for the Bound and Free states in non-transitioning and transitioning trajectories. For violin plots, thick red and dotted gray lines represent the median and two quartiles, respectively. **(D-E)** Transitioning trajectories for Ioc3-Halo (**D**) and Ioc4-Halo (**E**). Left: Overlay of raw histograms of $\log_{10}$ diffusion coefficients for 'Bound only' (turquoise), 'Free only' (yellow), 'Transitioning' (purple), and total trajectories (thin black). Right: Quantification (%) of transitioning trajectories in the diffusion coefficient histogram. **(F)** For all classified transitioning trajectories, 'FREE' to 'BOUND' and 'BOUND' to 'FREE' transition frequencies are indicated. For **D-F,** errors represent standard deviation between 2 or 3 biological replicates.

Figure 4-figure supplement 1. Survival plots [1-CDF] of dwell times showing 1- vs 2-component exponential decay fits.

(A-F)1- and 2-component exponential decay fits to survival plots of dwell times for Sth1-Halo (**A**), Snf2-Halo (**B**), Ino80-Halo (**C**), and Isw2-Halo (**D**), Chd1-Halo (**E**), and Isw1-Halo (**F**). **(G)** 1-CDF plot, pie chart as in Figure 4, and residence times of Ioc3-Halo (Left) and Ioc4-Halo (Right). **(H)** 1- and 2-component exponential decay fits to survival plots of dwell times for Ioc3-Halo (Left) and Ioc4-Halo (Right).

Figure 5-figure supplement 1. Expression levels and 1-CDF plots for wildtype and mutant ATPase-dead Isw2D312N.

(A) SDS-PAGE analysis; gel scanned for JF646 dye fluorescence (top) and imaged after Coomassie staining (bottom). Cell lysates of ATPase mutants (Isw1K227R-Halo, Isw2K215R-Halo and Chd1D513N-Halo) and their wildtype strains were prepared after treatment with JF646 at a saturating dye concentration (20 nM) for 2 h at 30oC. **(B-E)**1-

1264   CDF plot in log-log scale for Isw2K215R (**B**), Isw1K227R (**C**), Chd1K407R (**D**), and

1265   Chd1D513N (**E**) compared to wildtype. Colored dashed lines represent 95% confidence

1266   interval.

1267

Figure 6-figure supplement 1. Chromatin-bound remodelers display higher radius

of confinement (Rc) values than H2B.

(A) Radius of confinement values of bound trajectories in 10-ms exposure movies for

histone H2B and chromatin remodelers. Violin plot showing distribution of R_c values, and

comparison between histone H2B and each of wildtype remodelers by ordinary one-way

ANOVA test. **(B)** Violin plot of individual D values by slow-tracking for Isw1-Halo after 30

min pre-treatment with 3 or 5 ug/mL thiolutin, and comparison between wildtype and

thiolutin-treated samples by unpaired t test (ns: not significant). Thick red and dotted gray

lines represent the median and two quartiles, respectively.

1277

Figure 7-figure supplement 1. Time trace simulations of temporal occupancies at

promoters bound by multiple remodelers, and analysis of CHD1 DNA-binding

mutant.

(A-B) Time trace simulations of occupancy as in Figure 6, at a promoter region bound by

RSC, INO80, and ISW2 remodelers (**A**) or by RSC, SWI/SNF, INO80, and ISW2

remodelers (**B**). Individual time trace simulations are shown above, and the cumulative

simulated occupancy time trace (black) shows any one or multiple remodelers bound in

the time course of 500 s. **(C-D)** Fast-tracking and slow-tracking results for CHD1 DNA-

binding domain mutant (Chd1R1016A/K1020A/R1255A-Halo). (wildtype: dashed gray

lines; DBD mutant: solid green lines). (**C**) Normalized histogram $\log_{10}$ diffusion coefficients

(Left) and Spot-On kinetic modeling results (Right). (**D**) 1-CDF plot, pie chart, and

residence times.

1290

1291   **Supplementary file 1. List of Yeast strains used in the study**

1292   **Supplementary file 2. List of oligonucleotides used in the study**

Figure 1-figure supplement 1. Cell growth, integrity, and localization of HaloTagged remodeler subunits.
(A) SDS-PAGE gel scanned for JF646 dye fluorescence (top) and imaged after Coomassie staining (bottom). Cell lysates are prepared after treatment with JF646 at a saturating dye concentration (20 nM) for 2 h at 30°C. **(B)** Overlay of Phase Contrast image and 'nuclear glow' captured by initial JF552 dye excitation of yeast stained with JF552. **(C)** Five-fold dilutions of HaloTag fusion and wildtype strains are plated on YPAD plates at the indicated temperatures for 2-3 days. **(D)** Relationship between D_{free} values determined by Spot-On analysis and the calculated molecular weights of chromatin remodeling complexes.

A

Remodeler	F_{bound} (%)	D_{bound} ($\mu m^2 s^{-1}$)	D_{free} ($\mu m^2 s^{-1}$)
RSC (Sth1)	66.0 ± 1.1	0.038 ± 0.002	0.577 ± 0.069
SWI/SNF (Snf2)	55.9 ± 1.3	0.036 ± 0.007	0.464 ± 0.043
INO80 (Ino80)	48.3 ± 0.2	0.050 ± 0.004	0.639 ± 0.058
ISW2 (Isw2)	35.4 ± 0.4	0.062 ± 0.001	1.014 ± 0.024
CHD1 (Chd1)	47.8 ± 4.9	0.047 ± 0.007	0.915 ± 0.038
ISW1a/b (Isw1)	52.0 ± 1.1	0.050 ± 0.001	0.806 ± 0.010
ISW1a (Ioc3)	39.6 ± 0.0	0.067 ± 0.004	0.791 ± 0.059
ISW1b (Ioc4)	55.5 ± 2.6	0.044 ± 0.005	0.532 ± 0.042
Histone H2B	79.4 ± 1.9	0.026 ± 0.000	0.970 ± 0.158

B

Figure 1-figure supplement 2. Spot-On kinetic modeling analyses.
(A) Kinetic parameters (F_{bound}, D_{bound}, D_{free}) determined by Spot-On analyses. Errors represent standard deviation between 2 or 3 biological replicates. **(B)** Raw displacement histograms for individual biological replicates over the first 5 time frames (Δt: 10, 20, 30, 40, 50 ms). A two-state kinetic model was used for fitting the CDF [black lines] in Spot-On. F_{bound} values (shown above the histograms) are highly reproducible between replicates.

Figure 3-figure supplement 1. Validation of two diffusive states classified by vbSPT, and quantification of transitioning frequencies.
(A) Pipeline for classification and sub-classification of trajectories using vbSPT. Individual displacement lengths were determined for 'bound only', 'free only', and 'transitioning'-subclassified trajectories to validate the two states and state transitions. **(B-C)** Violin plot (**B**) and median value (**C**) of individual displacement lengths for the Bound and Free states in non-transitioning and transitioning trajectories. For violin plots, thick red and dotted gray lines represent the median and two quartiles, respectively. **(D-E)** Transitioning trajectories for Ioc3-Halo (**D**) and Ioc4-Halo (**E**). Left: Overlay of raw histograms of $\log_{10}$ diffusion coefficients for 'Bound only' (turquoise), 'Free only' (yellow), 'Transitioning' (purple), and total trajectories (thin black). Right: Quantification (%) of transitioning trajectories in the diffusion coefficient histogram. **(F)** For all classified transitioning trajectories, 'FREE' to 'BOUND' and 'BOUND' to 'FREE' transition frequencies are indicated. For **D-F**, errors represent standard deviation between 2 or 3 biological replicates.

Figure 4-figure supplement 1. Survival plots [1-CDF] of dwell times showing 1- vs 2-component exponential decay fits.

(A-F) 1- and 2-component exponential decay fits to survival plots of dwell times for Sth1-Halo (**A**), Snf2-Halo (**B**), Ino80-Halo (**C**), and Isw2-Halo (**D**), Chd1-Halo (**E**), and Isw1-Halo (**F**). **(G)** 1-CDF plot, pie chart as in Fig. 4, and residence times of Ioc3-Halo (Left) and Ioc4-Halo (Right). **(H)** 1- and 2-component exponential decay fits to survival plots of dwell times for Ioc3-Halo (Left) and Ioc4-Halo (Right).

Figure 5-figure supplement 1. Expression levels and 1-CDF plots for wildtype and mutant ATPase-dead Isw2D312N.
(A) SDS-PAGE analysis; gel scanned for JF646 dye fluorescence (top) and imaged after Coomassie staining (bottom). Cell lysates of ATPase mutants (Isw1K227R-Halo, Isw2K215R-Halo and Chd1D513N-Halo) and their wildtype strains were prepared after treatment with JF646 at a saturating dye concentration (20 nM) for 2 h at 30°C. **(B)** 1-CDF plot in log-log scale for Chd1K407R compared to wildtype. Colored dashed lines represent 95% confidence interval. **(C)** 1-CDF plot, pie chart, and residence times of wildtype Isw2-Halo (top) and Isw2D312N-Halo (bottom).

Figure 6-figure supplement 1. Chromatin-bound remodelers display higher radius of confinement (R$_c$) values than H2B.
(A) Radius of confinement values of bound trajectories in 10-ms exposure movies for histone H2B and chromatin remodelers. Violin plot showing distribution of R$_c$ values, and comparison between histone H2B and each of wildtype remodelers by ordinary one-way ANOVA test. **(B)** Violin plot of individual D values by slow-tracking for Isw1-Halo after 30 min pre-treatment with 3 or 5 ug/mL thiolutin, and comparison between wildtype and thiolutin-treated samples by unpaired t test (ns: not significant). Thick red and dotted gray lines represent the median and two quartiles, respectively.

Figure 7-figure supplement 1. Time trace simulations of temporal occupancies at promoters bound by multiple remodelers, and analysis of CHD1 DNA-binding mutant.

(A-B) Time trace simulations of occupancy as in Fig. 6, at a promoter region bound by RSC, INO80, and ISW2 remodelers (**A**) or by RSC, SWI/SNF, INO80, and ISW2 remodelers B). Individual time trace simulations are shown above, and the cumulative simulated occupancy time trace (black) shows any one or multiple remodelers bound in the time course of 500 s. **(C-D)** Fast-tracking and slow-tracking results for CHD1 DNA-binding domain mutant (Chd1R1016A/K1020A/R1255A-Halo). (wildtype: dashed gray lines; DBD mutant: solid green lines). (**C**) Normalized histogram $\log_{10}$ diffusion coefficients (Left) and Spot-On kinetic modeling results (Right). (**D**) 1-CDF plot, pie chart, and residence times.