The genetic basis of structural colour variation in mimetic *Heliconius* butterflies

Melanie N. Brien[1*], Juan Enciso Romero[1,2,*], Emma V. Curran[1], Andrew J. Parnell[3], Carlos Morochz[4],

Patricio A. Salazar[1], Pasi Rastas[5], Thomas Zinn[6], Nicola J. Nadeau[1]

[1]Department of Animal and Plant Sciences, The University of Sheffield, Alfred Denny Building,

Western Bank, Sheffield, S10 2TN, United Kingdom

[2]Biology Program, Faculty of Natural Sciences, Universidad del Rosario, Bogotá, Colombia

[3]Department of Physics and Astronomy, The University of Sheffield, Hicks Building, Hounsfield

Road, Sheffield, S3 7RH, United Kingdom

[4]Biology & Research Department, Mashpi Lodge, Ecuador

[5]Institute of Biotechnology, University of Helsinki, Finland

[6]ESRF - The European Synchrotron, 38043, Grenoble Cedex 9, France

[*]These authors contributed equally. Correspondence: M. Brien (Current address: University of

Jyväskylä, Finland, melanie.n.brien@jyu.fi) and J. Enciso Romero (juan.enciso@urosario.edu.co).

Short title: Structural colour genetics in *Heliconius* butterflies.

Keywords: Structural colour, iridescence, mimicry, *Heliconius,* QTL, convergence.

Abstract

Structural colours, produced by the reflection of light from ultrastructures, have evolved multiple times in butterflies. Unlike pigmentary colours and patterns, little is known about the genetic basis of these colours. Reflective structures on wing-scale ridges are responsible for iridescent structural colour in many butterflies, including the Müllerian mimics *Heliconius erato* and *Heliconius melpomene.* Here we quantify aspects of scale ultrastructure variation and colour in crosses between iridescent and non-iridescent subspecies of both of these species and perform quantitative trait locus (QTL) mapping. We show that iridescent structural colour has a complex genetic basis in both species, with offspring from crosses having wide variation in blue colour (both hue and brightness) and scale structure measurements. We detect two different genomic regions in each species that explain modest amounts of this variation, with a sex-linked QTL in *H. erato* but not *H. melpomene.* We also find differences between species in the relationships between structure and colour. Our results suggest that these species have followed different evolutionary trajectories in their convergent evolution of similar structural colour. This study provides a starting point for determining the genetic basis of structural colouration more broadly.

Introduction

Structural colours are some of the most vivid and striking colours found in nature. They are formed from the reflection and refraction of light from physical ultrastructures and examples of these can be found in nearly all groups of organisms. The structural colours of butterflies and moths are among the best described and play diverse roles, including initiation of courtship and mating behaviour [1,2], sex and species discrimination [1,3], long distance mate recognition [4] and signalling of quality and adult condition [5].

Butterflies and moths have evolved several mechanisms of structural colour production by modifying different components of scale morphology [6,7]. Scales typically consist of a flat lower lamina connected to an upper lamina by pillar-like trabeculae, with a small space separating the upper and lower laminae. The lower lamina can act as a thin film reflector that produces hues ranging from violet to green depending on its thickness [8–10]. The upper lamina has a more complex structure; it consists of a parallel array of ridges connected by cross-ribs, and modifications to these can yield diverse optical effects. For example, a lamellar structure in the ridges forms multilayer reflectors that produce the iridescent blue in *Morpho* butterflies [11] and the UV reflectance found in *Colias eurytheme* [12,13]. The variations in hue and brightness of colour produced in the intricate structures of the upper lamina depend on an interplay between the number of lamellae, the thickness of each layer and the spacing between the ridges [14]. It has also been shown that butterflies can tune the structures of the upper lamina to achieve the opposite effect, resulting in a substantial reduction of light reflection. Species with ultra-black wings use a combination of thicker trabeculae, reduced cross-rib spacing, and ridges with a steep shape to funnel the light into melanin granules, thus strongly increasing its absorption [15,16].

Recent studies have begun to uncover the genetic and developmental basis of structural colours in some species [7], revealing a common pattern in *Bicyclus anynana* and *Junonia coenia*; artificial selection for colourful phenotypes quickly resulted in changes in lower lamina thickness, and consequently hue, in a relatively small number of generations [8,10]. On the other hand, knockouts of known colour pattern genes [17], and genes involved in pigment synthesis pathways [18,19], have

shown that modification of these can result in altered scale ultrastructure, and moreover has brought

out unexpected instances of structural colour [18]. Interestingly, there are groups of butterflies for

which the gene *optix,* a known major colour pattern gene [20]*,* can jointly control pigment based

colouration and thickness of the lower lamina, producing blue structural colour [10]. It remains

unclear whether this joint control extends to other butterfly taxa that produce structural colour using

lower lamina reflection. In species with ridge reflectors, such as *Heliconius,* this does not seem to be

the case [18]. More efforts are required to understand the genetic underpinnings and evolution of

upper lamina architecture and the organisation of reflective nanostructures in the scales of iridescent

butterflies.

Wing colour patterns have been widely studied in the *Heliconius* butterflies, a group of butterflies

with a diverse set of aposematic colour patterns. These patterns show examples of both convergent

evolution between distantly related species, and divergent evolution within species. Some species

form mimicry rings, in which wing patterning is under strong positive frequency-dependent selection

due to predation [21]. Pigment colour patterns are largely determined by a small number of genes

which are homologous across species. Extensive research has uncovered a toolkit of five loci which

control much of the colour pattern variation in *Heliconius* species, and some other Lepidoptera [22].

Heliconius also display structural colour, and in comparison to the well-studied pigment colours, very

little is known about the development and genetic basis of these. From previous work, we know that

certain aspects of scale morphology affect the production of iridescent colouration, a type of angle-

dependent structural colour [14,23]. While overall scale morphology is similar between iridescent and

non-iridescent scales, those with blue structural colour have overlapping ridge lamellae which act as

multilayer reflectors (as in *Morpho*), along with a greater density of ridges on the scale (narrower

ridge spacing).

Structural colour has evolved multiple times within the *Heliconius* genus [14]. In some species, all

subspecies have iridescent colour, while others exhibit interspecific variation in iridescence.

Heliconius erato and *Heliconius melpomene* are two co-mimicking species which diverged around

10-13 Mya [24] with each evolving around 25 different colour pattern morphs [25]. Most of the

different colour patterns are produced by pigment colours, but subspecies found west of the Andes in

Ecuador and Colombia also have an iridescent blue structural colour. *H. erato cyrbia* and *H.*

melpomene cythera found in Western Ecuador have the brightest iridescence, while subspecies *H.*

erato demophoon and *H. melpomene rosina,* found to the north in Panama, are matt black in the

homologous wing regions (Figure 1). A hybrid zone forms between the iridescent and non-iridescent

groups where they meet near the border between Panama and Colombia, and here, populations with

intermediate levels of iridescence can be found [26]. Continuous variation in iridescent colour is

observed in the centre of the hybrid zone and in experimental crosses [23], suggesting that this trait is

controlled by multiple genes. The evolution of pigmentation and simple colour pattern traits have

frequently been shown to involve the re-use of a small number of genes across animal species

[22,27,28]. However, we may expect the genetic basis of a quantitative trait controlled by multiple

genes, such as iridescence in these species, to be less predictable [29]. In addition, iridescence in *H. e.*

cybria is much brighter than in *H. m. cythera* [14], suggesting some differences in scale structure and

presumably genetic control of this structure formation process.

Here we use crosses between subspecies of iridescent and non-iridescent *Heliconius* to determine the

genetics of both colour and scale ultrastructure traits for the first time. We measure the intensity of

blue colour and overall luminance (brightness) to assess variation in colour. We complement our

estimates of colour variation with high throughput measurements of ridge spacing and cross-rib

spacing using ultra small-angle X-ray scattering (USAXS). Using a quantitative trait locus (QTL)

mapping approach, we can identify the location and effect sizes of loci in the genome that are

controlling variation in iridescent colour. Comparison of the genetic basis of these traits between *H.*

melpomene and *H. erato*, two distantly related mimetic species, allows us to ask whether, like pigment

colour patterns, variation in iridescent colour and scale structure is also an example of gene reuse.

Results

Experimental crosses

Experimental crosses were performed using geographical morphs of both *Heliconius erato* and *Heliconius melpomene*. In both species, morphs from Panama (*H. e. demophoon* and *H. m. rosina*) were crossed with morphs from Western Ecuador (*H. e. cyrbia* and *H. m. cythera*), then the F1 generation crossed with each other to produce an F2. For *H. erato*, we also analysed a backcross between the F1 and *H. e. cyrbia* (Figure 1). Due to a mix-up in the insectary, one of our largest *H. melpomene* broods, named 'EC70', was obtained from a cross between an F1 father and a mother of unknown parentage, likely an F2 individual. Further details of the crosses are in Table S1.

In the offspring of these crosses, we measured four phenotypes - blue colour (BR), luminance, ridge spacing and cross-rib spacing. 'BR' is the average blue pixel value relative to red, extracted from RGB values from photographs. This was used as a measure of blue hue, while luminance measured overall brightness. Scale structure measurements were extracted from small angle X-ray scattering data from a single family of each species.

Figure 1. (A) Crosses between iridescent and non-iridescent morphs of *Heliconius melpomene* and *Heliconius erato*. For *H. melpomene* we used F2 crosses, plus one cross thought to be F1xF2 (not shown). For *H. erato*, we used F2 crosses and a backcross to the iridescent race. (B) Electron microscope image of a *Heliconius erato cyrbia* wing scale showing the analysed scale structure measurements.

Phenotypic variation

Based on current understanding of the reflective structures in *Heliconius* and other butterflies [14,30], we predict that reduced ridge spacing should increase the brightness of the reflectance from the wing due to the increased density of reflective structures (ridges) on the wing. However, there are other aspects of scale nanostructure that we would also expect to affect colour hue and brightness, particularly lamellae layering. These aspects could correlate with ridge spacing if they are controlled by the same genetic and developmental mechanism, or they could be controlled independently. The extent to which the different aspects of colour and scale structure are correlated in our broods gives an initial approximation of the extent of genetic correlation between these traits, and therefore the extent to which we are likely to find that they are controlled by the same or different genetic loci. For the *H. erato* family for which we have USAXS measurements, we found significant correlations between scale structure and colour measurements: ridge spacing has a negative correlation of moderate

strength with both BR value (Spearman's rho = -0.411, n = 56, p = 0.0013) and luminance

(Spearman's rho = -0.46, n = 56, p = 0.0002, Figure 2), as expected. We also found significant

correlations between ridge and cross-rib spacing (Spearman's rho = 0.342, n = 56, p = 0.008) and

between BR value and luminance (Spearman's rho = 0.341, n = 56, p = 0.008). Cross-rib spacing is

negatively correlated with BR value (Spearman's rho = -0.325, n = 56, p = 0.012) but not with

luminance (Spearman's rho = -0.18, n = 56, p = 0.17).

On the other hand, the *H. melpomene* family only showed significant correlations for BR value and

luminance (Spearman's rho = 0.295, n =73, p = 0.008) but no correlations were found for BR value

and ridge spacing (Spearman's rho = 0.0012, n = 73, p = 0.991), luminance and ridge spacing

(Spearman's rho = -0.117, n = 73, p =0.306, Figure 2), ridge spacing and cross-rib spacing

(Spearman's rho = -0.088, n = 73, p = 0.441), BR value and cross-rib spacing (Spearman's rho = 0.05,

n = 73, p = 0.622) or luminance and cross-rib spacing (Spearman's rho = 0.13, n = 73, p = 0.226).

This separation of ridge spacing and colour variation may have arisen in *H. melpomene* as a result of a

higher number of generations of mixing in the brood we used (EC70) in comparison to the *H. erato*

brood. However, it would also be consistent with a greater genetic correlation between multiple

aspects of scale structure in *H. erato* than in *H. melpomene*.

Figure 2. Relationship between ridge spacing and luminance in F2 offspring of *H. erato* (A) and *H.*
melpomene (B). A significant correlation was found for *H. erato* only.

We tested for sexual dimorphism in the four traits and did not find any significant differences between

male and female *H. melpomene* offspring from the largest brood (EC70) in all four traits (Table S2).

Sexual dimorphism in *H. erato* was analysed in our previous study, finding differences in all four

phenotypes in offspring with a maternal *cyrbia* grandfather [23]. Our sample size from the parental

subspecies is small but did not reveal any large difference between the sexes for any of the traits

measured, suggesting that the difference we see between the sexes in the crosses are largely due to

sex-linkage, as we previously proposed [23]. Nevertheless, we included sex as a covariate in all our

subsequent analyses to control for the possibility of slight sexual dimorphism.

QTL mapping in H. erato

Significant QTL were found for three phenotypes in *H. erato* - BR colour, luminance and ridge

spacing (Figure 3, Table 1, Table S3). When analysing the colour measurements, F2 and backcross

genome scans were combined, and for BR values showed 2 significant QTL on chromosomes 20 and

the Z sex chromosome. These were also found when analysing the F2 broods separately from the

backcross brood (Figure S1). The intervals around these QTL each contain around 150 genes. At both

markers, individuals with Panama-type genotypes (Pan/Pan and Pan(W)) had lower BR values than

Ecuador-type and heterozygous genotypes, following the expected trend (Figure 4). Only two

individuals have homozygous Panama-type *demophoon* genotypes at the Z chromosome marker due

to the small number of individuals with a *demophoon* maternal grandfather (Table S1). The QTL on

the Z chromosome explained the largest proportion of the phenotypic variation in BR colour in both

the F2 crosses (19.5%) and the backcross (24.6%), and the chromosome 20 QTL explained a further

12.3%.

Luminance (overall brightness of the wing colour) was highly associated with the Z chromosome

(Figure 3B). The significant marker did not map exactly to the same position as for the BR values but

was apart by only 3.6cM, and confidence intervals for each overlap. Individuals with Ecuador-type

alleles had higher luminance values than those with Panama-type alleles, showing the same trend as

the BR values (Figure 4). This QTL explained 40.2% of the variance in luminance values in the F2

crosses and 24.2% in the backcross. This was the only significant QTL for luminance, with nothing

appearing on chromosome 20.

A single QTL on the Z chromosome was also significant for ridge spacing (Figure 3C). This marker

was at a different position to the markers for BR and luminance, which may be due to using only one

family for this analysis, although the confidence intervals overlap. When we repeated the analysis for

luminance using only this family, the same marker showed the highest LOD score for both luminance

and ridge spacing (Figure S1). All genotypes with one or two Ecuador-type alleles had similar ridge

spacing, but those with a hemizygous Panama-type genotype ('Pan(W)' in Figure 4) had significantly

wider ridge spacing. This QTL explained 34.8% of variance in ridge spacing in this family. No

significant QTL were found for cross-rib spacing, although the highest LOD score was seen on the Z

chromosome (Figure 3D).

208
209 Figure 3. *H. erato* QTL plots including all families for BR colour (A) and luminance (B). QTL plots
210 for a single family for ridge spacing (C) and cross-rib spacing (D). Dotted line shows p=0.05
211 significance level.

212
213 Figure 4. The phenotypes of F2 individuals with different genotypes at the most significant markers
214 within each *H. erato* QTL. The QTL for BR on the Z chromosome (A) and chromosome 20 (B).
215 Significant Z markers for luminance (C)and ridge spacing (D). 'Pan' denotes alleles from the Panama
216 race *demophoon*, and 'Ec' the Ecuador race *cyrbia*. Marker positions are shown in Table 1.
217
218

Phenotype	Marker	Chromosome	Position (cM)	LOD	p
Heliconius erato					
BR colour (all families)	Herato2101_12449252	Z	38.0	7.07	0.001
	Herato2001_12633065	20	32.9	4.75	0.022
Luminance (all families)	Herato2101_12449398	Z	41.6	14.50	<0.001
Ridge Spacing (single family)	Herato2101_7491127	Z	23.0	5.21	0.013
Heliconius melpomene					
BR (all families)	Hmel203003o_2119654	3	15.22	7.26	0.001
Luminance (all families)	Hmel203003o_2635435	3	17.97	13.61	<0.001
Ridge spacing (EC70)	Hmel207001o_11550301	7	53.61	5.71	<0.001

Table 1: Significant QTL found for three phenotypes in *H. erato* and *H. melpomene*.

Genome scans for three F2 families were combined with scans of the EC70 brood. Since the

parentage of the mother of the EC70 brood is unknown, the maternal alleles in the offspring could not

be assigned as being from either a *cythera* or a *rosina* grandparent. Therefore, in this family only

paternal alleles were taken into account (and all maternal alleles set to 'A'), and the cross was treated

as if a backcross. A single significant QTL for BR colour was found on chromosome 3 (Figure 5A,

Table 1, Table S4). The interval around this marker relates to a genomic region of approximately

3Mbp and contains 130 genes. The marker explains 15.3% of phenotypic variation in EC70 (which

will be an underestimate due to all maternal alleles being assigned to the same genotype) and 9.2% in

the three F2 families. This suggests that there will be other loci in the genome with effects on colour

that we lack the power to detect. A marker on chromosome 10 shows a high LOD score that does not

reach significance, but could be suggestive of a locus on this chromosome. This is clearer when

looking only at the three F2 families separated from the EC70 brood (Figure S2). This marker

explains 17% of variation in the F2 families. Chromosome 10 contains the *WntA* gene, which controls

pigment colour pattern elements, however the QTL interval does not overlap with this. Luminance

was also strongly associated with markers on chromosome 3 (Figure 5B). The significant marker was

2.75cM from the marker for BR colour, and the confidence intervals overlap. In contrast, for ridge

spacing we found a significant QTL on chromosome 7 (using just the EC70 brood), explaining 30.3%

of variation (Figure 5C). Again, no significant QTL were found for cross-rib spacing (Figure 5D).

Individuals with homozygous Panama-type genotypes at the mapped chromosome 3 markers had

lower BR and luminance values (Figure 6). Individuals carrying Ecuador-type alleles at the mapped

chromosome 7 marker showed reduced ridge spacing, consistent with the observation that the Panama

race has greater ridge spacing and that this is associated with a lack of iridescence.

243
244 Figure 5. *H. melpomene* QTL plots including all families for BR colour (A) and luminance (B). QTL
245 plots for a single family for ridge spacing (C) and cross-rib spacing (D).
246

247
248 Figure 6. Genotype x phenotype plots for significant *H. melpomene* QTL. For the F2 families: (A) BR
249 colour and (B) luminance. 'Pan' denotes alleles from the Panama race *rosina*, and 'Ec' the Ecuador
250 race *cythera*. (C) Ridge spacing in the EC70 brood. Here we only know the origin of the paternal
251 allele. Marker positions shown in Table 1.

 Association analysis for H. melpomene

Treating the EC70 cross as a backcross is not ideal as it ignores potentially interesting genetic

variation that is segregating from the mother (which phenotypically and genetically appeared to be an

F2 individual), and could lead to artefacts. Therefore, we also conducted a SNP association analysis

including the full set of *H. melpomene* broods (F2 + EC70) and for the EC70 family alone, using a

family-based score test for association [31] and controlling for family structure [32].

The GWAS of the EC70 family did not reveal significant genetic associations for either of the scale

structure traits (Figure S3). The SNPs with the lowest p-values were found on chromosomes 1, 7, 17

and Z for ridge spacing, and on chromosomes 13 and Z for cross-rib spacing. Similarly, we did not

find significant associations between SNPs and BR values in EC70, although the lowest p-values were

found on chromosome 3 and chromosome 10, which aligns with the QTL results. We did find SNPs

significantly associated with variation in luminance on chromosome 3, with a single SNP scoring

above both conservative genome-wide Bonferroni corrected thresholds. When analysing all of *H.*

melpomene broods together accounting for pedigree structure, we found significant associations for

both colour traits (BR value and luminance) on chromosome 3. Stronger support for association was

found for luminance in comparison to BR value. Overall, the association analysis results supported

the QTL results but did not reveal additional loci.

Location of pigmentation pathway genes

Previous studies have shown that genes belonging to pigmentation pathways have an effect on both

pigmentation and scale structure in butterflies. Matsuoka and Monteiro [19] tested the effect of genes

from melanin and ommochrome synthesis pathways and found that 2 out of 8 genes tested had an

effect on the modification of scale structure. We queried the position of these 8 genes (*DDC, yellow,*

aaNAT, pale, ebony, vermillion, white and *scarlet*) in the genomes of *H. erato* and *H. melpomene* to

assess whether our QTL overlapped any of these genes (Table S5). In addition, we queried the *tan*

gene which is involved in cuticular melanisation [33] and is up-regulated in melanic regions of

Heliconius butterflies [34]. We found that the pigmentation genes belonging to the ommochrome and

melanin pathways have similar physical locations in the genomes of *H. erato* and *H. melpomene*, as

expected due to their high level of synteny. None of the pigmentation pathway genes are located

within the Bayesian confidence intervals estimated for the QTL found in either species for structural

colour or scale structure variation. However, we found that the *tan* and *pale* genes, both belonging to

the melanin pathway, are located on the Z sex chromosome. The *pale* gene is located at a physical

position that is 1 Mbp and 724 kbp from the upper limits of the Bayesian confidence intervals of the

QTL found for ridge spacing and luminance respectively in *H. erato,* and *tan* is located approximately

1.76 Mbp and 1.2 Mbp from the lower limits of the confidence intervals of the same QTL.

Discussion

In one of the first studies to look at the genetics of structural colour variation in terms of both colour

and structure, we show that the trait is controlled by multiple genes in the co-mimics *Heliconius erato*

and *Heliconius melpomene*. While we found only a small set of significant loci, these explain

relatively little of the overall phenotypic variation, suggesting there are more loci which remain

undetected. We found evidence for a genetic correlation between two aspects of structural colour

variation – BR value for hue and luminance for brightness. Within each species these were controlled

by loci on the same chromosomes. In *H. erato*, this was on the Z chromosome, confirming our

previous phenotypic analysis [23], and in *H. melpomene*, on chromosome 3. An additional locus on

chromosome 20 was also found to affect blue colour but not brightness in *H. erato*. Brightness of

iridescent colour is predicted to be controlled primarily by the number of layers within the multilayer

reflector structure, while hue is predicted to be controlled by the thickness and spacing of the layers

[35]. A genetic correlation between these two aspects of colour would suggest that both aspects of

scale nanostructure are controlled by these genetic loci. This would be advantageous in the evolution

of structural colour by allowing concerted changes in both aspects required for colour production. In

contrast, the presence of a possible independent hue modifier locus on chromosome 20 in *H. erato*

would allow more evolutionary flexibility in the colours that could be produced.

We also found correlations between colour, brightness and ridge spacing in *H. erato*. The Z

chromosome locus appears to control ridge spacing, which could have a direct effect on the brightness

of the reflectance by increasing the density of reflective structures. As further proof of this, in the

single-family analyses, luminance and ridge spacing mapped to exactly the same marker.

Alternatively, the observed correlation between brightness and ridge spacing in *H. erato* may be a

product of an unobserved association between tighter ridge spacing and other aspects of scale

nanostructure, specifically the number of lamellae layers within the ridges. Theoretical analyses and

simulations of the optical properties of multilayers have revealed that increasing the number of layers

will result in a rapid increase of brightness; adding even a small number of layers produces a

significant increase in the amount of reflected light [36]. Therefore, the Z chromosome locus may be

affecting multiple aspects of scale structure, producing the observed correlations between the different

colour and structure measurements. Also consistent with this is the evidence for a genetic correlation

between ridge and cross-rib spacing in *H. erato.* The effect of the Z locus on cross-rib spacing did not

reach significance, but had the highest genome-wide LOD score. Cross-rib spacing is not predicted to

have an effect on colour but could have a developmental correlation with ridge spacing.

In contrast, these correlations are not seen in *H. melpomene*, with different loci controlling colour and

ridge spacing, suggesting a more dispersed genetic architecture and different loci controlling different

aspects of scale structure. We found strong evidence for a locus on chromosome 3 controlling BR and

luminance, but this locus appeared to have no effect on our measurements of scale structure and so is

likely controlling other aspects of scale structure not quantified here. Instead, we find a locus on

chromosome 7 that partially controls ridge spacing. We see a small, but not significant, effect of this

chromosome on BR colour, suggesting that ridge spacing may have a small and relatively weak effect

on colour in *H. melpomene*, despite the parental populations showing a similar difference in ridge

spacing to that seen in *H. erato.* Differences in the experimental crosses between species may have

influenced these results, with the analysis in *H. melpomene* using only paternal alleles, and so having

reduced power. Nevertheless, the Z chromosome QTL has an effect of similar size on ridge spacing

and brightness in *H. erato* and no other loci affecting ridge spacing were detected in *H. erato*,

suggesting that the detection of different loci controlling colour and ridge spacing in *H. melpomene*

cannot be explained by reduced power. If *H. erato* has a locus on the Z chromosome that can control

multiple aspects of scale structure, while *H. melpomene* requires mutations at loci dispersed around

the genome, this could provide one explanation for how *H. erato* has been able to evolve brighter

structural colour than that observed in *H. melpomene*. Measuring and quantifying the lamellar

structure in large numbers of individuals is a major technical challenge but could be an important next

step in understanding the genetic control of butterfly structural colour.

In contrast to many of the loci for pigment colour patterns which are homologous across multiple

Heliconius species, the loci controlling iridescence in *H. erato* and *H. melpomene* appear to be

different. Differences in the physical scale architecture and brightness of colour between the species

perhaps makes these genetic differences unsurprising [14,26]. A lack of genetic parallelism may also

be more likely for a quantitative trait such as iridescence [29]. Shared standing genetic variation can

lead to repeated use of certain genes, as is seen in the repeated adaptation of sticklebacks to freshwater

environments [37,38]. The differences in genetic architecture between species suggest this is unlikely

to be the case for iridescence in *Heliconius* and that the trait likely evolved due to multiple,

independent genetic mutations. Nevertheless, on the Z chromosome in *H. melpomene*, we do observe

elevated LOD scores in the QTL analysis and low p-value SNPs in the GWAS for both scale structure

traits, but neither of the colour traits. This suggests that *H. melpomene* may have a locus homologous

to that in *H. erato,* which is controlling some aspects of scale structure variation, but with apparently

little or no effect on colour variation.

One limitation of our approach is that the accuracy of QTL analyses is limited, particularly when

sample sizes are relatively small, as they are here, resulting in inflated estimates of explained variance

[39]; the percentages of explained variance we report are probably an overestimate due to the Beavis

effect [40]. Therefore, the QTL we detected likely explain the most variation in these particular

families, and other QTL that remain undetected possibly could be shared between species. Thus, we

cannot discard a scenario of a partially shared genetic architecture for scale structure and luminance.

Differences in the ecology and selective forces shaping colour traits between species could also

explain differences in the genetic basis of structural colour. Comparisons of the hybrid zones in both

species have revealed much broader clines in iridescent colour in *H. melpomene* compared to *H.*

erato, suggesting that it is under weaker selection in *H. melpomene* [26]. In addition, loci controlling

sexually selected traits are expected to accumulate on the sex chromosomes [41,42]. Sex-linkage of

this trait in *H. erato* could therefore be consistent with iridescent colour being sexually selected in this

species. Unpublished reflectance data also support greater sexual dimorphism in iridescent colour in

H. erato than in *H. melpomene* [43]. Disentangling cause and effect in this case is clearly complex

and experimental studies are needed to determine what function iridescence plays in these species.

In recent years, reverse genetics research has revealed a surprising connection between the molecular

machinery underlying the development of pigmented wing patterns and the ultrastructure of butterfly

scales in various species [17–19,44,45]. In particular, knock-outs of two of the major colour pattern

controlling loci, *optix* and *WntA,* have been shown to affect aspects of scale structure [10,17,18,44].

However, our QTL are not associated with any known colour pattern gene of large or small effect.

The known toolkit of colour-pattern controlling loci are located on chromosomes 1, 10, 13, 15 and 18

(*aristaless, WntA, vvl, cortex* and *optix* respectively) in *Heliconius* [22] while the QTL we detect are

on chromosomes 3, 7, 20 and 21 (the Z chromosome). Hence, our findings show that *H. erato* and *H.*

melpomene do not use the known molecular machinery of wing pattern production for sculpting

specialised nanostructures and iridescent wings.

Genes involved in pigment production pathways in insects have also been found to have an effect on

the architecture of the butterfly scale. Knockouts for two genes of the melanin-dopamine pathway

produced modifications of the scale architecture [19,46]. Our examination of the locations of melanin

and ommochrome production genes did not reveal an overlap with the QTL or their estimated

confidence intervals in any of our crosses. We found QTL on the same linkage group as two melanin

pathway genes, but despite this, the evidence suggests that these genes are not involved in scale

structure modification in *Heliconius,* as they fall outside the confidence intervals of the QTL. The

evidence we present here, together with previous reverse genetics research on *Heliconius* [17,18]

suggest that the genetic architecture of iridescent structural colour produced by ridge reflection in *H.*

erato and *H. melpomene* is completely decoupled from that of mimicry related wing pattern regulation

and that of pigment production. Furthermore, the QTL that we found here do not overlap with QTL

for pigment pattern variation that were found in a super-set of the families that we have used here

[47], further supporting the idea that structural colour variation and wing pattern variation is unlinked

in both species.

Overall, we show that evolution has targeted different sets of genes to develop scale architecture

suitable for structural colour production in *H. erato* and *H. melpomene*. The lack of gene reuse in the

evolution of this complex quantitative trait may be due to a joint effect of the ecological role of

structural colour shaping the mutational landscape differently in both species, and also of a presumed

less constrained evolutionary pathway due to the increased availability of loci of small effect that can

potentially evolve iridescent blue phenotypes. Although our forward genetics approach does not

single out specific genes, it has revealed novel loci of morphological evolution in butterflies. Gene

expression and functional studies could follow up genes in these QTL intervals to understand the role

these play in sculpting colour-producing nanostructures.

Methods

Sampling

Non-iridescent *H. erato demophoon* and *H. melpomene rosina* were collected from Gamboa, Panama (9.12ºN, 79.67ºW). Iridescent *H. erato cyrbia* and *H. melpomene cythera* were collected from the area around Mashpi, Ecuador (0.17ºN, 78.87ºW). Within a few hours of emergence, wings of offspring were removed and stored in glassine envelopes and bodies were stored in NaCl saturated 20% dimethyl sulfoxide (DMSO) 0.25M EDTA solution to preserve the DNA. A total of 155 *H. erato* individuals from 5 broods were used to generate linkage maps and perform QTL mapping (3 *demophoon* and 3 *cyrbia* grandparents, 11 F1 parents, and 40 backcross and 99 F2 offspring). For *H. melpomene*, data from 4 broods made up of 228 individuals were used (1 *rosina* and 2 *cythera* grandparents, 5 F1 and 219 offspring, and 1 unknown parent). Many samples analysed in this study were previously used in Brien et al. [23] and Bainbridge et al. [47]. Further details of crosses and samples used in each analysis can be found in Table S1.

DNA extraction and sequencing

Genomic DNA was extracted from approximately half of the thorax of each individual using the Qiagen DNeasy Blood & Tissue kit. Single-digest Restriction site-associated DNA (RAD) library preparation and sequencing were carried out by the Edinburgh Genomics facility. DNA was digested with the enzyme *PstI*, and libraries were sequenced on an Illumina HiSeq 2500 producing 125bp paired-end reads. Parents of the crosses (16 *H. erato* and 9 *H. melpomene*) were included at 2x higher concentration within the pooled library to produce a higher depth of coverage.

Data processing and linkage map construction

Sequence data processing and linkage map construction are described in detail in Bainbridge et al. [47]. Briefly, RAD sequences were demultiplexed using the RADpools function in RADtools (v1.2.4, [48]). FASTQ files were mapped to the *H. erato* v1 reference genome [49] or the *H. melpomene* v2.5 reference genome [50] using bowtie2 v2.3.2 [51]. BAM files were then sorted and indexed with

SAMtools (v1.3.1, [52]) and PCR duplicates were removed using Picard tools MarkDuplicates

(v1.102). SAMtools mpileup and Lep-MAP data processing scripts [53] were used to call genotype

posteriors.

Linkage maps were constructed with Lep-MAP3 [53], using the modules ParentCall2 (with options

ZLimit=2 and removeNonInformative=1), Filtering2, SeparateChromsomes2 (using sizeLimit = 50

and a LOD score of 12 for *H. erato* and 23 for *H. melpomene*) and JoinSingles2All. Finally, the

module OrderMarkers2 was used to order the markers along the linkage groups, taking into account

the genomic positions of the markers. Female recombination was set to 0 and male recombination set

to 0.05 (following [54]). We used LMPlot to visualise the maps and check for errors in marker order.

Any erroneous markers that caused long gaps at the beginning or ends of the linkage groups were

manually removed. Genotypes were phased using Lep-MAP's map2genotypes.awk script, and

markers were named using the map.awk script and a list of the SNPs used to provide the scaffold

name and position. The final linkage maps contained 5648 markers for *H. erato*, and 2163 markers for

H. melpomene.

Measurements of colour variation

Wings were photographed under standard lighting conditions (full details in [23]). A colour checker in

each photograph was used to standardise the photographs using the levels tool in Adobe Photoshop

(CS3). RGB values were extracted from two areas of each wing (proximal areas of both the forewing

and hindwing) and averaged. Blue-red (BR) values were used as a measure of blue iridescent colour.

These were calculated as (B-R)/(B+R), where +1 is completely blue and -1 is completely red.

Luminance measured overall brightness and was calculated as R+G+B, with each colour having a

maximum value of 255.

Measurements of scale structure variation using ultra small-angle X-ray scattering (USAXS)

The USAXS experiment was carried out at the ID02 beamline at the European Synchrotron (ESRF,

Grenoble, France). We examined scale structure variation for one family of each species (n=56 *H.*

erato F2 and n=73 *H. melpomene* (unknown mother)). We measured between 33 and 113 points per

individual along a linear proximodistal path across the proximal part of the forewing, which has the

most vivid iridescence in the blue subspecies (Figure S4). The X-ray beam wavelength was $\lambda = 1\text{Å}$

and it had 12.45 keV energy. The beam was collimated to a section of 50 μm × 50 μm to probe a

small area within a single scale. The scattered radiation was recorded with a high-sensitivity FReLon

16M Kodak CCD detector with an effective area of 2048 × 2048 pixels and a 24 μm pixel size, a

sample to detector distance of 30.7m, accessible q range between $8.5\text{x}10\text{-4 nm}^{-1}$ and $3.8\text{x}10\text{-2 nm}^{-1}$,

and limited interpretation of structure spacing to above 160 nm. The raw images were corrected for

dark current and spatial distortion.

Scanning electron microscopy (SEM)

SEM on a small subset of individuals was used to confirm our scale structure measures from the

USAXS data. We sampled four iridescent *H. e. cyrbia* and three *H. m. cythera*, and eight *H. e.*

demophoon and three *H. m. rosina* with black pigment colour. Representative SEM images of dorsal

cover scales for each of the four races used are in Figure S5. First, we compared ridge spacing of blue

individuals vs. black individuals on the dorsal side. This allowed us to gauge the expected range of

scale structure variation in blue and black coloured scales. Secondly, we compared ridge spacing of

dorsal vs. ventral scales within each race. This comparison allowed us to separate signals coming

from overlapping scales in the USAXS experiment.

Wing samples were cut from the region of interest shown in Figure S4. We mounted samples on

aluminium SEM stubs using 9mm adhesive carbon tabs and coated them with a thin layer of gold

using vacuum evaporation. Samples were imaged on a JEOL JSM-6010LA microscope and we used

the InTouchScope software to take images of the regions of interest at a magnification level of 900X.

Images included both cover and ground scales.

At the midpoint of the scale length, we drew a linear transect perpendicular to the orientation of the

ridges (Figure S6). Ridge spacing was calculated using the PeakFinder tool [55]; a plugin for the Fiji

[56] image analysis package, using a tolerance level of 60 and a minimum distance threshold of 10

pixels between peaks. PeakFinder detects spikes of intensity along the linear transect, each peak

corresponding to a single ridge. It then yields the distances, in pixels, between subsequent peaks

(Figure S6). Using the scale bar of SEM slides we determined the variable rate of conversion from

pixels to nm to be $25 \times 10^{-3} / M$ where M is the level of magnification and we used this variable rate

to convert distances between ridges to nm and average all distances between ridges to obtain average

ridge spacing per scale.

Analysis of USAXS data to extract ridge spacing values

We first did a qualitative joint assessment of our USAXS and SEM data to get an idea of the scale

structures we captured in our experiment and how to best get their information. Our interpretation of

scale structures in a USAXS scattering pattern is shown in Figure S7. Two important observations

come from this joint assessment. Firstly, it can be observed that blobs of scattered radiation distribute

along linear paths from the centre of the picture with equal spacing between them (Figure S7 A). This

periodic scattering can be understood as the signature of the interaction of the beam with the array of

parallel ridges on the surface of the scale (Figure S7 B) [57]. Secondly, the beam goes through the

dorsal and ventral sides of the wing, projecting onto the detector, with the scattering due to structures

of as many as four scales: cover and ground scales on both dorsal and ventral sides of the wing, in the

case that all of them overlap. This can be observed in Figure S7 A, where at least three overlapping

scales were picked up by the beam. Our SEM analysis revealed that scales with smaller ridge spacing

are found on the dorsal side compared to the ventral side of the wing regardless of the structural

colour of the butterfly analysed (Figure S8). These differences in ridge spacing between dorsal and

ventral scales should be discernible in the USAXS data.

We used these two observations to implement a script that extracts relevant scale structure

information from the USAXS data. The script scans the 2D scattering pattern in non-overlapping

regions of 5°. If the region contains periodic scattering it is assumed to correspond to the ridges of the

scale and is kept for comparison with other sections that also show periodic scattering. The script

keeps the section with the highest magnitude of the scattering vector q based on the observation that q

should be larger for dorsal scales because they have tighter ridge spacing. The chosen region is

recorded and the scanning section is rotated 85° anti-clockwise to capture the scattering due to cross-

ribs, which are perpendicular to ridges in the scale. Azimuthal integration is then performed on the

two regions of interest, reducing them to a one-dimensional representation of scattered intensity I as a

function of the magnitude of the vector q in a similar way to previous studies [14,23], applying

corrections for transmitted flux and masking the pixels corresponding to the beam stop. The script

uses the pyFAI [58] and fabio [59] libraries to do the azimuthal integration and USAXS data

handling.

Despite dividing the scattering pattern in small regions to pick apart the signals due to different

structures and scales, some frames still contained information of more than one scale. This happens in

the case that overlapping scales have highly similar orientations (i.e. their ridges are parallel or nearly

parallel). Thus, a visual inspection was done and we filtered out 1D scattering patterns that contained

signals from more than one scale. We also aimed to reduce noise in the scattering signals to improve

estimation of dimensions of scale structures. We ran a PCA on the 1D scattering data and retained

PCs that cumulatively explained 90% or more of the variation in the data. The original data were

projected back onto the retained PCs. The choice of threshold for the percentage variance explained is

arbitrary, but we found 90% to be a sensible balance to retain signal but reduce noise.

The retained 1D scattering patterns were fit to a Lorentzian curve (Figure S9) using the FitManager

module of the silx package [60]. The "center" parameter of the fitting procedure gives the magnitude

of the scattering vector q. This magnitude contains information on the distance between peaks of

intensity in reciprocal space. To get estimates of structure sizes in real space, we use the expression d

$= 2\pi/q$ where d is the distance between ridges or cross-ribs in real space. We averaged estimates of

each structure on a per-individual basis to get estimates of average ridge spacing and average cross-rib

spacing per butterfly.

Quantitative Trait Locus mapping

The R package R/qtl was used for the QTL analysis [61]. For *H. erato*, initially the F2 crosses were

analysed together and the backcross analysed separately. Genotype probabilities were calculated for

these two groups using *calc.genoprob*. We ran standard interval mapping to estimate LOD (logarithm

of the odds) scores using the *scanone* function with the Haley-Knott regression method. In the F2

analysis, sex and family were included as additive covariates, and family was included as an

interactive covariate, to allow multiple families to be analysed together. Sex was included as a

covariate in the backcross analysis to account for any sexual dimorphism. To determine the

significance level for the QTL, we ran 1000 permutations, with *perm.Xsp=T* to get a separate

threshold for the Z chromosome. A single F2 family (n=56) was used to analyse scale structure

variation (ridge spacing and cross-rib spacing) using the same method, albeit a higher number of

permutations was used for determining the significance level of the QTL (4000). For analyses of BR

colour and luminance, LOD scores for the F2 crosses and the backcross were added together, to allow

analysis of all individuals together to increase power, and the significance level recalculated in R/qtl.

Confidence intervals for the positions of QTL were determined with the *bayesint* function and we

used a *fitqtl* model to calculate the phenotypic variance that each QTL explained. Genome scan plots

and genotype plots were made with R/qtl2 [62]. Genetic distances in the QTL results are based on the

observed recombination rate and expressed in centimorgans (cM), which is the distance between two

markers that recombine once per generation. These were related to physical distances based on the

marker positions in the assembled reference genome of each species.

The same method was used to run genome scans for BR colour and luminance in *H. melpomene*. LOD

scores of the three F2 families were added to the LOD score from the EC70 family, as in *H. erato*, and

the significance level recalculated. Again, a single family was used for analysis of scale structures

(n=73).

Association analysis for H. melpomene

The test was done in two steps: in the first step a mixed model is used to get estimates of trait values

after adjusting for covariates and random polygenic effects using the linear model:

$Y_i = \mu + \sum_j \quad \beta_j C_{ji} + G_i + e_i$ (1)

In this model we used a genomic kinship matrix to account for pedigree structure. In the model Y_i is a

vector of phenotypes for each individual, μ is the mean phenotype for the population, C_{ji} is the value

of covariate j for individual i, β_j is the estimated effect for each covariate, G_i are random additive

polygenic effects and e_i are the residual effects for each individual. The result of this first part are the

trait residuals after accounting for covariates and random polygenic effects and is denoted by $\hat{e}_i$. The

estimates of trait variation that were not explained by kinship or covariates in the first step are then fit

to a second model:

$$\hat{e}_i = \mu + \beta g + e_i \quad (2)$$

In the second step, β is the parameter for the effect of each SNP tested, g is a single SNP and e_i is the

residual variation in phenotype for each individual i. We did the association tests on two sets: the first

set included all *H. melpomene* families and two phenotypes; BR value and luminance. The second set

included the EC70 family only and four phenotypes: BR value, luminance, ridge spacing and cross-rib

spacing.

Genotypes were called using the 'mpileup' command in samtools v1.8 [52] with minimum mapping

quality set to 20, and the 'call' command in bcftools v1.8 [52] run in consensus mode, with SNP p-

value threshold set to 0.05 and calling only variant sites. For our association analyses we kept biallelic

SNPs only. We applied the following quality control criteria to our SNP data: we excluded individuals

and sites that were not at least 95% genotyped. Sites were also excluded from our analysis if they

were not in Hardy-Weinberg equilibrium ($p<0.01$). The MAF threshold was set to the default value

(5/2n, where n is the number of individuals). After quality control, we retained 146,122 markers

genotyped across 211 individuals for the first set, and 63,132 sites genotyped across 69 individuals for

the second set. We set sites called as heterozygous on the sex chromosome in females as missing

(females are the heterogametic sex). Multiple testing was accounted for using two thresholds: A

Bonferroni corrected threshold calculated as 0.05/k where k is the number of sites analysed, and a less

conservative 5% false discovery rate threshold estimated using the GenABEL qvaluebh95 function.

Acknowledgements

We thank the governments of Ecuador and Panama for permission to collect butterflies. Thanks to

Darwin Chalá, Juan López and Gabriela Irazábal for their assistance with the crosses. We are grateful

to the European Synchrotron Radiation Facility for provision of X-ray beamtime under proposal

LS2720, and to Andrew Dennison for assistance with USAXS data collection. Library preparation and

sequencing was performed by staff at the Edinburgh Genomics Facility, at the University of

Edinburgh. We thank Vicky Lloyd for assistance with the collection and analysis of SEM data and for

helpful comments on the manuscript, and Alan Dunbar for use of the scanning electron microscope.

Data availability

Sequence data have been deposited in the European Nucleotide Archive under project number

PRJEB38330. Photographs of all samples can be found at https://doi.org/10.5281/zenodo.3799188.

USAXS data handling scripts can be found at https://github.com/juanenciso14/butterfly_usaxs.

Linkage maps, QTL scripts and all phenotypic measurements used will be available on publication.

Author contributions

MNB and JER performed the QTL analyses and wrote the manuscript with NJN. PR and MNB constructed the linkage maps. AJP co-ordinated collection of the USAXS data, which were collected by TZ, AJP, EVC, NJN and MNB. JER analysed the USAXS and SEM data and ran the association analysis. PAS, CM, MNB, EVC and NJN performed the crosses. NJN devised and co-ordinated the study. All authors read and commented on the manuscript.

Funding disclosure

This work was funded by the UK Natural Environment Research Council (NERC) through an Independent Research Fellowship (NE/K008498/1) to NJN. MNB and EVC were funded by the NERC doctoral training partnership, ACCE (NE/L002450/1). JER is funded through the Leverhulme Centre for Advanced Biological Modelling as well as scholarships from Universidad del Rosario and the University of Sheffield. The scanning electron microscope is funded via a EPSRC 4CU grant (EP/K001329/1).

References

1. Silberglied RE, Taylor OR. Ultraviolet reflection and its behavioral role in the courtship of the sulfur butterflies *Colias eurytheme* and *C. philodice* (Lepidoptera, Pieridae). Behav Ecol Sociobiol. 1978;3: 203–243.

2. Obara Y, Ozawa G, Fukano Y, Watanabe K, Satoh T. Mate preference in males of the cabbage butterfly, *Pieris rapae crucivora*, changes seasonally with the change in female UV color. Zoolog Sci. 2008;25: 1–5.

3. Rutowski RL. The use of visual cues in sexual and species discrimination by males of the small sulphur butterfly *Eurema lisa* (Lepidoptera, Pieridae). J Comp Physiol. 1977;115: 61–74.

4. Sweeney A, Jiggins C, Johnsen S. Insect communication: Polarized light as a butterfly mating signal. Nature. 2003;423: 31–32.

5. Kemp DJ, Rutowski RL. Condition dependence, quantitative genetics, and the potential signal content of iridescent ultraviolet butterfly coloration. Evolution. 2007;61: 168–183.

6. Ghiradella H. Light and color on the wing: structural colors in butterflies and moths. Appl Opt. 1991;30: 3492–3500.

7. Lloyd VJ, Nadeau NJ. The evolution of structural colour in butterflies. Curr Opin Genet Dev. 2021;69: 28–34.

8. Wasik BR, Liew SF, Lilien DA, Dinwiddie AJ, Noh H, Cao H, et al. Artificial selection for structural color on butterfly wings and comparison with natural evolution. Proc Natl Acad Sci U S A. 2014;111: 12109–12114.

9. Stavenga DG, Leertouwer HL, Wilts BD. Coloration principles of nymphaline butterflies - thin films, melanin, ommochromes and wing scale stacking. J Exp Biol. 2014;217: 2171–2180.

10. Thayer RC, Allen FI, Patel NH. Structural color in *Junonia* butterflies evolves by tuning scale lamina thickness. Elife. 2020;9. doi:10.7554/eLife.52187

11. Vukusic P, Sambles JR, Lawrence CR, Wootton RJ. Quantified interference and diffraction in single *Morpho* butterfly scales. Proc R Soc Lond B Biol Sci. 1999;266: 1403–1411.

12. Ghiradella H. Development of ultraviolet-reflecting butterfly scales: How to make an interference filter. J Morphol. 1974;142: 395–409.

13. Rutowski RL, Macedonia JM, Merry JW, Morehouse NI, Yturralde K, Taylor-Taft L et al. Iridescent ultraviolet signal in the orange sulphur butterfly (*Colias eurytheme)*: spatial, temporal and spectral properties. Biological J. Linnean Soc. 2007;90: 349-364.

14. Parnell AJ, Bradford JE, Curran EV, Washington AL, Adams G, Brien MN, et al. Wing scale ultrastructure underlying convergent and divergent iridescent colours in mimetic butterflies. J R Soc Interface. 2018;15. doi:10.1098/rsif.2017.0948

15. Davis AL, Nijhout HF, Johnsen S. Diverse nanostructures underlie thin ultra-black scales in butterflies. Nat Commun. 2020;11: 1294.

16. Vukusic P, Sambles JR, Lawrence CR. Structurally assisted blackness in butterfly scales. Proc Biol Sci. 2004;271 Suppl 4: S237–9.

17. Concha C, Wallbank RWR, Hanly JJ, Fenner J, Livraghi L, Rivera ES, et al. Interplay between
Developmental Flexibility and Determinism in the Evolution of Mimetic *Heliconius* Wing
Patterns. Curr Biol. 2019;29: 3996–4009.e4.

18. Zhang L, Mazo-Vargas A, Reed RD. Single master regulatory gene coordinates the evolution and
development of butterfly color and iridescence. Proc Natl Acad Sci U S A. 2017;114: 10707–
10712.

19. Matsuoka Y, Monteiro A. Melanin Pathway Genes Regulate Color and Morphology of Butterfly
Wing Scales. Cell Rep. 2018;24: 56–65.

20. Reed RD, Papa R, Martin A, Hines HM, Counterman BA, Pardo-Diaz C, et al. *optix* drives the
repeated convergent evolution of butterfly wing pattern mimicry. Science. 2011;333: 1137–1141.

21. Mallet J, Barton NH. Strong natural selection in a warning-color hybrid zone. Evolution.
1989;43: 421–431.

22. Nadeau NJ. Genes controlling mimetic colour pattern variation in butterflies. Curr Opin Insect
Sci. 2016;17: 24–31.

23. Brien MN, Enciso-Romero J, Parnell AJ, Salazar PA, Morochz C, Chalá D, et al. Phenotypic
variation in crosses shows that iridescent structural colour is sex-linked and controlled by
multiple genes. Interface Focus. 2019;9: 20180047.

24. Kozak KM, Wahlberg N, Neild AFE, Dasmahapatra KK, Mallet J, Jiggins CD. Multilocus
species trees show the recent adaptive radiation of the mimetic H*eliconius* butterflies. Syst Biol.
2015;64: 505–524.

25. Sheppard PM. Genetics and the evolution of Müllerian mimicry in *Heliconius* butterflies.
Philosophical Transactions of the Royal Society of London. B, Biological Sciences. 1985: 433–
610. doi:10.1098/rstb.1985.0066

26. Curran EV, Stankowski S, Pardo-Diaz C, Salazar C, Linares M, Nadeau NJ. Müllerian mimicry
of a quantitative trait despite contrasting levels of genomic divergence and selection. Mol Ecol.
2020;29: 2016–2030.

27. Manceau M, Domingues VS, Linnen CR, Rosenblum EB, Hoekstra HE. Convergence in
pigmentation at multiple levels: mutations, genes and function. Philosophical Transactions of the
Royal Society B: Biological Sciences. 2010. pp. 2439–2450. doi:10.1098/rstb.2010.0104

28. Hubbard JK, Uy JAC, Hauber ME, Hoekstra HE, Safran RJ. Vertebrate pigmentation: from
underlying genes to adaptive function. Trends Genet. 2010;26: 231–239.

29. Conte GL, Arnegard ME, Peichel CL, Schluter D. The probability of genetic parallelism and
convergence in natural populations. Proc Biol Sci. 2012;279: 5039–5047.

30. Wilts BD, Vey AJM, Briscoe AD, Stavenga DG. Longwing (*Heliconius*) butterflies combine a
restricted set of pigmentary and structural coloration mechanisms. BMC Evol Biol. 2017;17: 226.

31. Chen W-M, Abecasis GR. Family-based association tests for genome wide association scans. Am
J Hum Genet. 2007;81: 913–926.

32. Aulchenko YS, Ripke S, Isaacs A, van Duijn CM. GenABEL: an R library for genome-wide
association analysis. Bioinformatics. 2007;23: 1294–1296.

33. Thurmond J, Goodman JL, Strelets VB, Attrill H, Gramates LS, Marygold SJ, et al. FlyBase 2.0:
the next generation. Nucleic Acids Res. 2018;47: D759–D765.

34. Ferguson LC, Maroja L, Jiggins CD. Convergent, modular expression of ebony and tan in the
mimetic wing patterns of *Heliconius* butterflies. Dev Genes Evol. 2011;221: 297–308.

35. Gruson H, Andraud C, Daney de Marcillac W, Berthier S, Elias M, Gomez D. Quantitative
characterization of iridescent colours in biological studies: a novel method using optical theory.
Interface Focus. 2019;9: 20180049.

36. Kinoshita S, Yoshioka S. Structural colors in nature: the role of regularity and irregularity in the
structure. ChemPhysChem. 2005;6: 1442–1459.

37. Colosimo PF, Hosemann KE, Balabhadra S, Villarreal G Jr, Dickson M, Grimwood J, et al.
Widespread parallel evolution in sticklebacks by repeated fixation of Ectodysplasin alleles.
Science. 2005;307: 1928–1933.

38. Burke MK, Liti G, Long AD. Standing genetic variation drives repeatable experimental evolution
in outcrossing populations of *Saccharomyces cerevisiae*. Mol Biol Evol. 2014;31: 3228–3239.

39. Slate J. From beavis to beak color: a simulation study to examine how much QTL mapping can
reveal about the genetic architecture of quantitative traits. Evolution. 2013;67: 1251–1262.

40. Xu S. Theoretical Basis of the Beavis Effect. Genetics. 2003;165: 2259–2268.

41. Rice WR. Sex Chromosomes and the Evolution of Sexual Dimorphism. Evolution. 1984. p. 735.
doi:10.2307/2408385

42. Reinhold K. Sex linkage among genes controlling sexually selected traits. Behav Ecol Sociobiol.
1998;44: 1–7.

43. Brien MN. The genetics and evolution of iridescent structural colour in *Heliconius* butterflies.
PhD thesis, University of Sheffield. 2019.

44. Fenner J, Benson C, Rodriguez-Caro L, Ren A, Papa R, Martin A, et al. *Wnt* Genes in Wing
Pattern Development of Coliadinae Butterflies. Frontiers in Ecology and Evolution. 2020.
doi:10.3389/fevo.2020.00197

45. Livraghi L, Hanly JJ, Van Belleghem SM, Montejo-Kovacevich G, van der Heijden ESM, Loh
LS, et al. *Cortex cis*-regulatory switches establish scale colour identity and pattern diversity in
Heliconius. bioRxiv. 2021. doi:10.1101/2020.05.26.116533

46. Zhang L, Martin A, Perry MW, van der Burg KRL, Matsuoka Y, Monteiro A, et al. Genetic
Basis of Melanin Pigmentation in Butterfly Wings. Genetics. 2017. pp. 1537–1550.
doi:10.1534/genetics.116.196451

47. Bainbridge HE, Brien MN, Morochz C, Salazar PA, Rastas P, Nadeau NJ. Limited genetic
parallels underlie convergent evolution of quantitative pattern variation in mimetic butterflies. J
Evol Biol. 2020. doi:10.1111/jeb.13704

48. Baxter SW, Davey JW, Johnston JS, Shelton AM, Heckel DG, Jiggins CD, et al. Linkage
mapping and comparative genomics using next-generation RAD sequencing of a non-model
organism. PLoS One. 2011;6: e19315.

49. Van Belleghem SM, Rastas P, Papanicolaou A, Martin SH, Arias CF, Supple MA, et al.
Complex modular architecture around a simple toolkit of wing pattern genes. Nat Ecol Evol.
2017;1: 52.

50. Davey JW, Chouteau M, Barker SL, Maroja L, Baxter SW, Simpson F, et al. Major
Improvements to the *Heliconius melpomene* Genome Assembly Used to Confirm 10
Chromosome Fusion Events in 6 Million Years of Butterfly Evolution. G3. 2016;6: 695–708.

51. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:
357–359.

52. Li H. A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. Bioinformatics. 2011;27: 2987–
2993.

53. Rastas P. Lep-MAP3: robust linkage mapping even for low-coverage whole genome sequencing
data. Bioinformatics. 2017;33: 3726–3732.

54. Morris J, Navarro N, Rastas P, Rawlins LD, Sammy J, Mallet J, et al. The genetic architecture of
adaptation: convergence and pleiotropy in *Heliconius* wing pattern evolution. Heredity.
2019;123: 138–152.

55. Vischer N. PeakFinder. 2013. Available:
https://sils.fnwi.uva.nl/bcb/objectj/examples/PeakFinder/PeakFinderTool.txt

56. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.

57. Sivia DS. Elementary Scattering Theory: For X-ray and Neutron Users. OUP Oxford; 2011.

58. Kieffer J, Karkoulis D. PyFAI, a versatile library for azimuthal regrouping. J Phys Conf Ser.
2013;425: 202012.

59. Knudsen EB, Sørensen HO, Wright JP, Goret G, Kieffer J. FabIO: easy access to two-
dimensional X-ray detector images in Python. J Appl Crystallogr. 2013;46: 537–539.

60. Vincent T, Valls V, payno, Kieffer J, Solé VA, Paleo P, et al. silx-kit/silx: v0.13.0 beta 0:
21/02/2020. 2020. doi:10.5281/zenodo.3678538

61. Broman KW, Wu H, Sen S, Churchill GA. R/qtl: QTL mapping in experimental crosses.
Bioinformatics. 2003. pp. 889–890. doi:10.1093/bioinformatics/btg112

62. Broman KW, Gatti DM, Simecek P, Furlotte NA, Prins P, Sen Ś, et al. R/qtl2: Software for
Mapping Quantitative Trait Loci with High-Dimensional Data and Multiparent Populations.
Genetics. 2019. pp. 495–502. doi:10.1534/genetics.118.301595

63. Challis RJ, Kumar S, Dasmahapatra KK, Jiggins CD, Blaxter M. Lepbase: the Lepidopteran
genome database. 2016. doi:10.1101/056994

64. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol
Biol. 1990;215: 403–410.