Evolutionary rates of testes-expressed genes differ between monogamous and promiscuous *Peromyscus* species

Authors: Landen Gozashti[1,2,3*], Russell Corbett-Detig[4] and Scott W. Roy[5*]

[1]Department of Organismic & Evolutionary Biology, Cambridge, MA, USA
[2]Museum of Comparative Zoology, Harvard University, Cambridge, MA, USA
[3]Howard Hughes Medical Institute, Cambridge, MA, USA
[4]Department of Biomolecular Engineering and Genomics Institute, University of California Santa Cruz, Santa Cruz, CA, USA
[5]Department of Biology, San Francisco State University, San Francisco, CA, USA

[*]Correspondence

Abstract

Reproductive proteins, including those expressed in the testes, are among the fastest evolving proteins across the tree of life. Sexual selection on traits involved in sperm competition is thought to be a primary driver of testes gene evolution and is expected to differ between promiscuous and monogamous species due to intense competition between males to fertilize females in promiscuous lineages and lack thereof in monogamous ones. Here, we employ the rodent genus *Peromyscus* as a model to explore differences in evolutionary rates between testes-expressed genes in monogamous and promiscuous species. We find candidate genes that may be associated with increased sperm production in promiscuous species and gene ontology categories that show patterns of molecular convergence associated with phenotypic convergence in independently evolved monogamous species. Overall, our results highlight the possible molecular consequences mating system, likely due to differences in selective pressures.

Introduction

Reproductive genes evolve faster than other genes (Swanson and Vacquier 2002; Clark et al. 2006). This pattern is ubiquitous across diverse lineages, spanning from microbes to mammals (Makalowski and Boguski 1998; Armbrust and Galindo 2001; Sato et al. 2002; Torgerson et al. 2002; Wik et al. 2008; Vacquier and Swanson 2011). Testes-expressed genes in particular are fast evolving, likely due to sexual selection driven by sperm competition (Parker 1970; Civetta and Singh 1995; Rice and Holland 1997; Swanson and Vacquier 2002; Torgerson et al. 2002; Ramm and Stockley 2009; Firman and Simmons 2010a; Harrison et al. 2015; Moyle

et al. 2020). Rapid evolution of testes-expressed genes is thought to generally reflect positive selection (Ramm et al. 2008; Ramm and Stockley 2009; Teng et al. 2017), but could also be explained by relaxed selection (Dapper and Wade 2016).

Differences in mating system, that is the general pattern by which males and females mate within a species, are expected to drive differences in rates of evolution of testes-expressed genes (Lüpold et al. 2016; Clutton-Brock 2017). The vast majority of mammalian species are promiscuous, meaning that females mate with multiple males and vice versa (Garcia-Gonzalez 2017). However, genetic monogamy (where males and females only mate with one individual for life) has evolved multiple times independently across mammalian lineages (Lukas and Clutton-Brock 2013). Promiscuous and monogamous species often exhibit morphological, physiological and behavioral differences associated with mate fidelity (Stanyon and Bigoni 2014; Tidière et al. 2015; Dapper and Wade 2016; Wey et al. 2017; Ambaryan et al. 2019; Civetta and Ranz 2019). Morphological and physiological differences are often associated with the male testes and with sperm performance, consistent with promiscuous but not monogamous males competing to fertilize females (Firman and Simmons 2010a; Dapper and Wade 2016). Faster, more viable sperm and higher sperm counts have been shown to yield an increased chance of fertilization and thus a fitness advantage in promiscuous species (Pizzari 2006). Indeed, promiscuous males often possess larger testes and produce more abundant and more competitive sperm than monogamous males, even in the case of recently diverged species (Heske and Ostfeld 1990; Shuster 2009; Firman and Simmons 2010a; Claw et al. 2018; Fisher et al. 2018). However, the genetic basis of morphological and physiological differences related to differences in mating system remain largely unexplored.

Social/sexual monogamy has evolved at least twice independently within the rodent genus *Peromyscus* (Figure 1) (Turner et al. 2010; Bedford and Hoekstra 2015), with behavioral and physical traits differing consistently between monogamous and promiscuous *Peromyscus* species (Fisher and Hoekstra 2010; Fisher et al. 2014, 2016, 2018; Bendesky et al. 2017). *Peromyscus* is a powerful model for interrogating the drivers of convergent evolution and adaptation due to its well-resolved phylogeny (Greenbaum et al. 2017; Sullivan et al. 2017), well documented convergence of phenotypes (Steiner et al. 2009; Manceau et al. 2010; Bedford and Hoekstra 2015), and extensive ecological, morphological, and behavioral variation (Gering et al. 2009; Fisher and Hoekstra 2010; Shorter et al. 2012; Bedford and Hoekstra 2015; Guralnick et al. 2020). Here, we combine evolutionary rate modeling and gene expression analysis to identify genes and gene ontology categories with differential evolution in promiscuous and monogamous species.

Methods

Obtaining and reconstructing transcriptomic data
We employed a conservative but robust pipeline for detecting signatures of parallel evolutionary rates in transcripts of *P. polionotus* and *P. eremicus* relative to *P. leucopus* and *P.*

maniculatus. First, we obtained *P. maniculatus, P. polionotus, P. leucopus,* and *P. eremicus* testes transcriptome data from (Lindsey et al. 2020), (https://www.ncbi.nlm.nih.gov/bioproject/584485), (https://datadryad.org/stash/dataset/doi:10.5061/dryad.r8ns3) and (Kordonowy and MacManes 2016) respectively. We used as a reference the testes specific transcriptome was previously assembled for *P. eremicus (Kordonowy and MacManes 2016)*. For the other three species, we aligned each species' testes transcriptome data to reference CDS data (GCF_000500345.1 for *P. maniculatus* and *P. polionotus* since no reference exists for *P. polionotus* and *P. maniculatus* is its closest relative with a reference CDS file, and GCF_004664715.2 for *P. leucopus*) using *hisat2 (Kim et al. 2015)* with default parameters and called variants using *bcftools* with default parameters. We then reconstructed each transcript in a variant aware manner using a custom python script that replaces the reference allele at a given site with any alternate allele at higher frequency in the considered testes dataset (see https://github.com/lgozasht/Peromyscus-reproductive-genetic-differences).

Identification of orthologous gene sets

Once we possessed reconstructed transcriptomes for each species, we performed an all vs. all BLAST (Altschul et al. 1990) with a minimum percent identity of 90 to identify putative homologous transcripts between species. We filtered our BLAST output for a minimum 80 percent overlap relative to each transcript length and concatenated all BLAST output into one file. Then, we employed *silixx* (Miele et al. 2011), a graph theory based clustering software, to cluster transcripts into families of highly similar sequences from homologous genes. We filtered for families that contained transcripts from all four species and included the transcript with the lowest e value when a species possessed multiple candidate transcripts for a given family. Through this process we identified 942 transcripts expressed in reproductive tissue that shared homology between all 4 of our considered species.

Generating and curating PAML input files

We employed the classic program *PAML* (Phylogenetic Analysis by Maximum Likelihood) (Yang 2007) to search for shared signatures of evolutionary rates in *P. eremicus* and *P. polionotus* relative to *P. leucopus* and *P. maniculatus*. For each transcript cluster, we used *MAFFT (Katoh and Standley 2013)* to generate a multiple sequence alignment and *IQ-TREE (Minh et al. 2020)* to infer a maximum likelihood based tree, as required for *PAML* input and confirmed that tree topology remained consistent. Since indels and poor alignment quality can impair *PAML* accuracy and functionality, we performed additional curation and filtering on our input alignments. We used the tool *Alignment_Refiner_v2,* (see https://github.com/dportik/Alignment_Refiner) (Young et al. 2019) to trim alignments above an overall missing data threshold of 1% and abandoned clusters with greater than 50 gaps in a particular sequence or 100 total gaps thereafter. We also ignored alignments containing

sequences with lengths not divisible by 3. We were left with 199 transcripts suitable for PAML input after our stringent filtering and curation.

Comparing evolutionary rates of transcripts in P. eremicus and P. maniculatus

Short branch lengths between *P. maniculatus* and *P. polionotus* make it challenging to conduct accurate pairwise *dn/ds* comparisons (Yang 2007; Kryazhimskiy and Plotkin 2008; Mugal et al. 2014), so instead we performed comparisons between *P. maniculatus* and *P. eremicus*. For each transcript cluster, we applied *PAML* first with a null model (*model = 2,* see http://abacus.gene.ucl.ac.uk/software/pamlDOC.pdf) in which *dN/dS* differs in *P. maniculatus* and *P. eremicus* relative to *P. polionotus* and *P. leucopus* then with an alternative model (*model = 2,* see http://abacus.gene.ucl.ac.uk/software/pamlDOC.pdf) in which each *dN/dS* differs between *P. maniculatus* and *P. eremicus* relative to *P. polionotus* and *P. leucopus*. We then calculated log likelihood ratios using our *null* and *alternative* output and performed likelihood ratio tests for each transcript cluster. After further manual alignment quality filtering and applying a false-discovery rate correction for multiple testing (Benjamini and Hochberg 1995), we were left with 7 transcripts under significant differences in rates of evolution for downstream analysis (α = .05).

Differential expression analysis of P. eremicus and P. maniculatus

We obtained testes RNA data from three runs of separate replicates for *P. eremicus* (SRA run excessions: ERR1353571, ERR1353572, and ERR1353573) and one run of pooled replicates for *P. maniculatus* (SRA run excession: SRR8587279) and aligned to respective testes transcriptomes using *hisat2* (Kim et al. 2015). Next, we obtained read counts for each transcript using *htseq-count* (Anders et al. 2014) and calculated *RPKM* for each transcript. We calculated delta RPKM for homologous transcripts between *P. eremicus* and *P. maniculatus*. Then, we performed independent Mann–Whitney U tests for delta *RPKMs* of transcripts corresponding to each gene in which we observed significant differences in evolutionary rates between *P. eremicus* and *P. maniculatus*. We aggregated p-values for transcripts corresponding to each gene using Fisher's combined probability to obtain a final p-value for each gene, in similarity to a method described in (Yi et al. 2018).

Comparing parallel rates of evolution between monogamous and promiscuous species

We used *PAML* first to compute the likelihood of a null model (*model = 0,* see http://abacus.gene.ucl.ac.uk/software/pamlDOC.pdf) in which *dn/ds* is equal across all branches of the tree with our alternative model. Then we fit a branch model (*model = 2,* see http://abacus.gene.ucl.ac.uk/software/pamlDOC.pdf) in which the *dn/ds* ratio is allowed to differ between monogamous and promiscuous lineages, but where the two ratios are constrained to be identical across all monogamous or promiscuous branches. We included the monogamous *P. eremicus* and *P. polionotus* and the promiscuous *P. maniculatus* and *P. leucopus* for this analysis.

To assess significance, we used a likelihood ratio test, which is X2 distributed with a single degree of freedom.

Finding evolutionary trends in different gene ontologies

We interrogated trends of evolutionary rates by aligning each gene that passed our upstream filtering to the *Mus musculus* GRCm38 genome (Genbank accession GCF_000001635.26) and cross referencing the MGI ontology database (see http://www.informatics.jax.org/mgihome/projects/aboutmgi.shtml). We then grouped genes by mouse gene ontology (GO) terms. For each group of genes, we performed a two-sided binomial test with regard to the number of genes evolving at slower (or faster) rates in monogamous species relative to promiscuous species. For our null, we used the proportion of genes evolving at slower (or faster) rates between monogamous and promiscuous species among all considered loci: P = .519. We filtered for groups in which we observed a significant trend of different evolutionary rates ($\alpha < 0.05$). To determine that these signatures of correlated evolutionary rates within GO categories are not driven by species-specific effects, we performed a test for idiosyncratic evolution in each species. To do this, for each species, we ran *PAML* with a branch model in which that species possessed its own α that differed from to the other three species.

Results and Discussion

We employ a maximum likelihood approach to compare evolutionary rates of transcripts expressed in the testes of the monogamous *P. eremicus* and the promiscuous *P. maniculatus*. Our analysis reveals 6 genes exhibiting significant differences in evolutionary rates between species. Intriguingly, 5 of these are implicated in spermatogenesis and are evolving faster in *P. maniculatus*: Znf644, KDM3A, Ddx25, Spata16 and ESSBP (Epididymis Secretory Sperm Binding Protein) (Figure 2, Table 1) (Dam et al. 2007; Liu et al. 2010; Tsai-Morris et al. 2010; Souza et al. 2017; Wilson et al. 2017). Sequence level changes leading to increased sperm abundance or competitive ability might be favored by selection in promiscuous species due to the selective pressures of mate competition. Thus, we hypothesize that these genes may have experienced positive selection in *P. maniculatus*. Consistent with this hypothesis, previous studies have suggested that sperm production is sensitive to local adaptation in light of the pressures associated with competition (Ramm and Stockley 2009; Winkler et al. 2019; Lindsey et al. 2020).

Arhgef18, the other gene with a significantly higher evolutionary rate in the promiscuous *P. maniculatus*, lacks known reproductive implications. Although its role in spermatogenesis remains unexplored, Arhgef18 has been shown to play a role in maintaining apico-basal polarity and localization of tight junctions and cortical actin, and thus plays a role in cellular morphology and cell-cell interactions, which are essential to sperm cell organization and sertoli-spermatid interactivity during spermatogenesis (Wong and Cheng 2009; Gao and Cheng 2016). Its rapid evolution in *P. maniculatus* suggests it too may evolve under positive selection.

Differential gene expression analysis lends further support to the evidence for positive selection of Spata16 and ESSBP. We calculated delta RPKMs for homologous transcripts between species, focusing specifically on the 6 genes exhibiting significantly different rates of evolution. Interestingly, we find that ESSBP and Spata16 exhibit significantly increased expression in *P. maniculatus* (*Mann-Whitney U* P = 0.0294 and 0.0376 respectively). The relationship between sequence divergence and expression divergence between species has been shown to be positively correlated in mammals (Liao and Zhang 2006; Warnefors and Kaessmann 2013). Furthermore, this pattern is consistent with changes in gene expression levels resulting from positive selection (Jordan et al. 2005), further raising the possibility that ESSBP and Spata16 might be under positive selection in *P. maniculatus* as a result of their roles in spermatogenesis.

We also used branch models to compare shared rates of evolution between two species in which monogamy evolved independently, *P. eremicus* and *P. polionotus* and two relatively divergent promiscuous *Peromyscus* species, *P. maniculatus* and *P. leucopus*. Due to the short branch lengths separating *P. maniculatus* and *P. polionotus*, branch tests using individual genes may be unreliable. Therefore, rather than focusing on significantly different rates between independent homologous loci, we used MGI's gene ontology database (Smith and Eppig 2009) to cluster all considered genes based on ontology, and explored trends of shared evolutionary rates between genes in monogamous species relative to promiscuous species with involvement in particular ontologies. We find significant differences in evolutionary trends between monogamous and promiscuous species in eight gene ontology categories, seven of which are important for sperm function (Table 2, Figure 3). We find that genes within ontologies with implications in sperm function generally have slightly increased evolutionary rates in monogamous species. In fact, all considered genes involved in "flagellated sperm motility," "motile cilium," "centrosome" and "cell projection" are evolving at faster rates in monogamous species. One possible explanation for this is that monogamous species exhibit reduced selective constraint on genes integral to sperm function. Mutations are likely filtered more frequently by purifying selection in promiscuous species than in monogamous species where negative selection is relaxed (Firman and Simmons 2010a). An alternative explanation is positive selection on these genes in monogamous species. However, in contrast with the expected observation for loci under positive selection, we do not find significant differences in evolutionary rates between monogamous and promiscuous species when we compare genes within these ontologies independently, and instead find that they exhibit dN/dS values closer to 1. These observations may constitute a molecular signature of convergent evolution in protein evolutionary rates for genes involved in sperm competitiveness in monogamous species.

Our results illuminate the molecular correlates of changes in reproductive functions associated with changes in mating systems. Sperm abundance, motility and morphology are critical for fertility (Sharpe 2012; Baptissart et al. 2013). We identify six genes that exhibit significantly increased evolutionary rates in the promiscuous *P. maniculatus* relative to the monogamous *P. eremicus*. Five of these are known to be involved in spermatogenesis and the

sixth has plausible connections. Additionally, two of these genes also exhibit significantly increased expression in *P. maniculatus*, providing additional evidence that these loci experience positive selection in *P. maniculatus* due to competition between males to fertilize females. Consistent with this interpretation, previous studies have found that sperm in the promiscuous *P. maniculatus* exhibit adaptive morphological traits and increased abundance when compared to that of monogamous *Peromyscus* species *(Fisher et al. 2016)*.

Furthermore, we observe that genes implicated in sperm function exhibit marginally faster evolutionary rates in two independently evolved monogamous species relative to promiscuous ones when clustered by ontology, highlighting the possible role of relaxed selection in propelling the molecular convergence of genes integral to sperm competitiveness. However, future work is required to investigate roles of selection, and we propose that these genes serve as candidates for future studies on adaptation and convergent evolution with regard to reproductive behavior. Future studies should include nucleotide and transcript-abundance polymorphism data to enable direct estimation of the mode of selection and increased sampling of monogamous species across many independent origins in diverse lineages to test for molecular convergence associated with differences in mating system.

Acknowledgments

The authors thank Jenny Chen for her input on optimizing statistical analyses and Hopi Hoekstra for advice. The authors declare no conflicts of interest. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Figure 1: Phylogenetic relationship between our four considered *Peromyscus* species with the addition of *Mus musculus* and *Rattus norvegicus* as outgroups. We note that we only include a small subset of known *Peromyscus* species in our phylogeny. Our tree was produced with *IQ-TREE* (Minh et al. 2020) using 15 concatenated orthologs in each respective species. Genetic distances are measured in units of the number of base substitutions in proportion to alignment length (Sievers et al. 2011). Blue colored nodes represent promiscuity and red colored nodes represent monogamy (Xia and Millar 1991; Shurtliff et al. 2005; Costa et al. 2016). Monogamy has evolved independently at least twice in the *Peromyscus* genus (Turner et al. 2010; Bedford and Hoekstra 2015).

Figure 2: Respective evolutionary rates (ω) for genes exhibiting different evolutionary rates between in the monogamous *P. eremicus* and the promiscuous *P. maniculatus*. Colored points correspond to genes exhibiting statistically significant differences in evolutionary rates. Blue points correspond to genes that play crucial roles in spermatogenesis.

Gene	Relative Evolutionary Rate	Involvement in Reproduction	Q value	Source for sexual implication
KDM3A	P>M	Spermatogenesis	<0.0001	(Wilson et al. 2017)
Spata16	P>M	Spermatogenesis	0.0006	(Chang et al. 2004)
Ddx25	P>M	Spermatogenesis	0.0020	(Tsai-Morris et al. 2010)

Znf644	P>M	Spermatogenesis	0.0002	(Liu et al. 2010)
ESSBP	P>M	Spermatogenesis	<0.0001	(Souza et al. 2017)
Arhgef18	P>M	Cell morphology and cell-cell interactions	<0.0001	(Niu et al. 2003)

Table 1: Genes exhibiting significantly different rates of evolution in considered monogamous and promiscuous species. Column 1 displays gene name. Column 2 provides the trend of evolution (monogamous (M) species vs. promiscuous (P) species). Column 3 summarizes respective known involvement in reproduction, column 4 provides FDR corrected X^2 P-values (Q-values), and column 5 contains sources for sexual implications.

Figure 3: Proportion of genes evolving faster in monogamous species relative to the number of genes within each respective ontology cluster. The seven ontology groups are colored orange or blue (Binomial test alpha = 0.05). The five Ontology groups with significant trends of evolution

in monogamous species relative to promiscuous ones and known specific implications in sperm performance are colored blue. Other significant ontologies are colored in orange. The dotted line represents the null expected proportion of genes evolving faster in monogamous species. The "*" denotes an ontology group that is not specifically involved in sperm performance but whose data point overlaps with an ontology group that is.

MGI Ontology	Relative Evolutionary Rate	Sexual Implication	P value	Source for sexual implication
cell projection	M>P	Sperm function (motility)	0.0078	(Avidor-Reiss et al. 2017)
centrosome	M>P	Sperm function (motility)	0.0078	(Terada et al. 2010; Schatten et al. 2011; Schatten and Stearns 2015; Gunes et al. 2020)
cilium	M>P	Sperm function (motility)	0.0390	(Gadella 2008; Inaba and Mizuno 2016; Yuan et al. 2019; Sironen et al. 2020)
flagellated sperm motility	M>P	Sperm function (motility)	0.0156	(Vladić et al. 2002; Lüpold et al. 2009; Firman and Simmons 2010b; Fisher et al. 2016)
hydrolase activity	M>P	NA	0.0351	NA
motile cilium	M>P	Sperm function (motility)	0.0312	(Gadella 2008; Inaba and Mizuno 2016; Yuan et al. 2019; Sironen et al.

				2020)
nucleic acid binding	M>P	NA	0.0352	NA
ribonucleoprotein complex	M>P	NA	0.0313	NA

Table 2: MGI Ontology categories in which we observe differences in trends of evolutionary rates. Column 1 presents ontology categories. Column 2 provides trends in evolutionary rate (monogamous (M) species vs. promiscuous (P) species). Column 3 provides respective sexual implications, column 4 contains P-values (*binomial test*) and column 5 contains sources for sexual implications.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403–410.

Ambaryan AV, Voznessenskaya VV, Kotenkova EV. 2019. Mating behavior differences in monogamous and polygamous sympatric closely related species Mus musculus and Mus spicilegus and their role in behavioral precopulatory isolation. *Russ. J. Theriol.* 18:67–79.

Anders S, Pyl PT, Huber W. 2014. HTSeq – A Python framework to work with high-throughput sequencing data. *Cold Spring Harbor Laboratory* [Internet]:002824. Available from: https://www.biorxiv.org/content/10.1101/002824v2

Armbrust EV, Galindo HM. 2001. Rapid evolution of a sexual reproduction gene in centric diatoms of the genus Thalassiosira. *Appl. Environ. Microbiol.* 67:3501–3513.

Avidor-Reiss T, Ha A, Basiri ML. 2017. Transition Zone Migration: A Mechanism for Cytoplasmic Ciliogenesis and Postaxonemal Centriole Elongation. *Cold Spring Harb. Perspect. Biol.* [Internet] 9. Available from: http://dx.doi.org/10.1101/cshperspect.a028142

Baptissart M, Vega A, Martinot E, Volle DH. 2013. Male fertility: Is spermiogenesis the critical step for answering biomedical issues? *Spermatogenesis* [Internet] 3. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3710220/

Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Series B Stat. Methodol.* 57:289–300.

Chang C, Chen Y-T, Yeh S-D, Xu Q, Wang R-S, Guillou F, Lardy H, Yeh S. 2004. Infertility with defective spermatogenesis and hypotestosteronemia in male mice lacking the androgen receptor in Sertoli cells. *Proc. Natl. Acad. Sci. U. S. A.* 101:6876–6881.

Civetta A, Ranz JM. 2019. Genetic Factors Influencing Sperm Competition. *Front. Genet.* 10:820.

Civetta A, Singh RS. 1995. High divergence of reproductive tract proteins and their association with postzygotic reproductive isolation in Drosophila melanogaster and Drosophila virilis group species. *J. Mol. Evol.* 41:1085–1095.

Clark NL, Aagaard JE, Swanson WJ. 2006. Evolution of reproductive proteins from animals and plants. *Reproduction* 131:11–22.

Claw KG, George RD, MacCoss MJ, Swanson WJ. 2018. Quantitative evolutionary proteomics of seminal fluid from primates with different mating systems. *BMC Genomics* 19:488.

Clutton-Brock T. 2017. Reproductive competition and sexual selection. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* [Internet] 372. Available from: http://dx.doi.org/10.1098/rstb.2016.0310

Costa F, Richardson JL, Dion K, Mariani C, Pertile AC, Burak MK, Childs JE, Ko AI, Caccone A. 2016. Multiple Paternity in the Norway Rat, Rattus norvegicus, from Urban Slums in Salvador, Brazil. *J. Hered.* 107:181–186.

Dapper AL, Wade MJ. 2016. The evolution of sperm competition genes: The effect of mating system on levels of genetic variation within and between species. *Evolution* 70:502–511.

Firman RC, Simmons LW. 2010a. Experimental evolution of sperm quality via postcopulatory sexual selection in house mice. *Evolution* 64:1245–1256.

Firman RC, Simmons LW. 2010b. Sperm midpiece length predicts sperm swimming velocity in house mice. *Biol. Lett.* 6:513–516.

Fisher HS, Hook KA, Weber WD, Hoekstra HE. 2018. Sibling rivalry: Males with more brothers develop larger testes. *Ecol. Evol.* 8:8197–8203.

Fisher HS, Jacobs-Palmer E, Lassance J-M, Hoekstra HE. 2016. The genetic basis and fitness consequences of sperm midpiece size in deer mice. *Nat. Commun.* 7:13652.

Gadella BM. 2008. Sperm membrane physiology and relevance for fertilization. *Anim. Reprod. Sci.* 107:229–236.

Gao Y, Cheng CY. 2016. Does cell polarity matter during spermatogenesis? *Spermatogenesis* 6:e1218408.

Garcia-Gonzalez F. 2017. Promiscuity. In: Vonk J, Shackelford T, editors. Encyclopedia of Animal Cognition and Behavior. Cham: Springer International Publishing. p. 1–7.

Gunes S, Sengupta P, Henkel R, Alguraigari A, Sinigaglia MM, Kayal M, Joumah A, Agarwal A. 2020. Microtubular Dysfunction and Male Infertility. *World J. Mens Health* 38:9–23.

Harrison PW, Wright AE, Zimmer F, Dean R, Montgomery SH, Pointer MA, Mank JE. 2015. Sexual selection drives evolution and rapid turnover of male gene expression. *Proc. Natl. Acad. Sci. U. S. A.* 112:4393–4398.

Heske EJ, Ostfeld RS. 1990. Sexual Dimorphism in Size, Relative Size of Testes, and Mating Systems in North American Voles. *J. Mammal.* 71:510–519.

Inaba K, Mizuno K. 2016. Sperm dysfunction and ciliopathy. *Reprod. Med. Biol.* 15:77–94.

Jordan IK, Mariño-Ramírez L, Koonin EV. 2005. Evolutionary significance of gene expression divergence. *Gene* 345:119–126.

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Mol. Biol. Evol.* 30:772–780.

Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low memory requirements. *Nat. Methods* 12:357–360.

Kordonowy L, MacManes M. 2016. Characterization of a Male Reproductive Transcriptome for Peromyscus eremicus (Cactus mouse). *Cold Spring Harbor Laboratory* [Internet]:048348. Available from: https://www.biorxiv.org/content/10.1101/048348v1.full

Kryazhimskiy S, Plotkin JB. 2008. The population genetics of dN/dS. *PLoS Genet.* 4:e1000304.

Liao B-Y, Zhang J. 2006. Evolutionary conservation of expression profiles between human and mouse orthologous genes. *Mol. Biol. Evol.* 23:530–540.

Lindsey LL, Platt RN, Phillips CD, Ray DA, Bradley RD. 2020. Differential Expression in Testis and Liver Transcriptomes from Four Species of Peromyscus (Rodentia: Cricetidae). *Genome Biol. Evol.* 12:3698–3709.

Liu Y-Q, Bai G, Zhang H, Su D, Tao D-C, Yang Y, Ma Y-X, Zhang S-Z. 2010. Human RING finger protein ZNF645 is a novel testis-specific E3 ubiquitin ligase. *Asian J. Androl.* 12:658–666.

Lukas D, Clutton-Brock TH. 2013. The evolution of social monogamy in mammals. *Science* 341:526–530.

Lüpold S, Calhim S, Immler S, Birkhead TR. 2009. Sperm morphology and sperm velocity in passerine birds. *Proceedings of the Royal Society B: Biological Sciences* 276:1175–1181.

Lüpold S, Manier MK, Puniamoorthy N, Schoff C, Starmer WT, Luepold SHB, Belote JM, Pitnick S. 2016. How sexual selection

can drive the evolution of costly sperm ornamentation. *Nature* 533:535–538.

Makalowski W, Boguski MS. 1998. Evolutionary parameters of the transcribed mammalian genome: an analysis of 2,820 orthologous rodent and human sequences. *Proc. Natl. Acad. Sci. U. S. A.* 95:9407–9412.

Miele V, Penel S, Duret L. 2011. Ultra-fast sequence clustering from similarity networks with SiLiX. *BMC Bioinformatics* 12:116.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* 37:1530–1534.

Moyle LC, Wu M, Gibson MJS. 2020. Reproductive proteins evolve faster than non-reproductive proteins among Solanum species. *Cold Spring Harbor Laboratory* [Internet]:2020.11.30.405183. Available from: https://www.biorxiv.org/content/10.1101/2020.11.30.405183v1

Mugal CF, Wolf JBW, Kaj I. 2014. Why time matters: codon evolution and the temporal dynamics of dN/dS. *Mol. Biol. Evol.* 31:212–231.

Niu J, Profirovic J, Pan H, Vaiskunaite R, Voyno-Yasenetskaya T. 2003. G Protein betagamma subunits stimulate p114RhoGEF, a guanine nucleotide exchange factor for RhoA and Rac1: regulation of cell shape and reactive oxygen species production. *Circ. Res.* 93:848–856.

Parker GA. 1970. Sperm competition and its evolutionary consequences in the insects. *Biol. Rev. Camb. Philos. Soc.* 45:525–567.

Pizzari T. 2006. Of mice and sperm. *Proc. Natl. Acad. Sci. U. S. A.* 103:14983–14984.

Ramm SA, Oliver PL, Ponting CP, Stockley P, Emes RD. 2008. Sexual selection and the adaptive evolution of mammalian ejaculate proteins. *Mol. Biol. Evol.* 25:207–219.

Ramm SA, Stockley P. 2009. Adaptive plasticity of mammalian sperm production in response to social experience. *Proc. Biol. Sci.* 276:745–751.

Rice WR, Holland B. 1997. The enemies within: intergenomic conflict, interlocus contest evolution (ICE), and the intraspecific Red Queen. *Behav. Ecol. Sociobiol.* 41:1–10.

Sato K, Nishio T, Kimura R, Kusaba M, Suzuki T, Hatakeyama K, Ockendon DJ, Satta Y. 2002. Coevolution of the S-locus genes SRK, SLG and SP11/SCR in Brassica oleracea and B. rapa. *Genetics* 162:931–940.

Schatten G, Stearns T. 2015. Sperm Centrosomes: Kiss Your Asterless Goodbye, for Fertility's Sake. *Curr. Biol.* 25:R1178–R1181.

Schatten H, Rawe VY, Sun Q-Y. 2011. The Sperm Centrosome: Its Role and Significance in Nature and Human Assisted Reproduction. *Journal of Reproductive and Stem Cell Biotechnology* 2:121–127.

Sharpe RM. 2012. Sperm counts and fertility in men: a rocky road ahead: Science & Society Series on Sex and Science. *EMBO Rep.* 13:398.

Shurtliff QR, Pearse DE, Rogers DS. 2005. Parentage Analysis of the Canyon Mouse (Peromyscus crinitus): Evidence for Multiple Paternity. *J. Mammal.* 86:531–540.

Shuster SM. 2009. Sexual selection and mating systems. *Proc. Natl. Acad. Sci. U. S. A.* 106 Suppl 1:10009–10016.

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M, Söding J, et al. 2011. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.* 7:539.

Sironen A, Shoemark A, Patel M, Loebinger MR, Mitchison HM. 2020. Sperm defects in primary ciliary dyskinesia and related causes of male infertility. *Cell. Mol. Life Sci.* 77:2029–2048.

Smith CL, Eppig JT. 2009. The mammalian phenotype ontology: enabling robust annotation and comparative analysis. *Wiley Interdiscip. Rev. Syst. Biol. Med.* 1:390–399.

Souza APB de, Schorr-Lenz ÂM, Lucca F, Bustamante IC Filho. 2017. The epididymis and its role on sperm quality and male fertility. *Anim. Reprod.* 14:1234–1244.

Stanyon R, Bigoni F. 2014. Sexual selection and the evolution of behavior, morphology, neuroanatomy and genes in humans and

other primates. *Neurosci. Biobehav. Rev.* 46P4:579–590.

Swanson WJ, Vacquier VD. 2002. The rapid evolution of reproductive proteins. *Nat. Rev. Genet.* 3:137–144.

Teng L, Fan X, Xu D, Zhang X, Mock T, Ye N. 2017. Identification of Genes under Positive Selection Reveals Differences in Evolutionary Adaptation between Brown-Algal Species. *Front. Plant Sci.* 8:1429.

Terada Y, Schatten G, Hasegawa H, Yaegashi N. 2010. Essential roles of the sperm centrosome in human fertilization: developing the therapy for fertilization failure due to sperm centrosomal dysfunction. *Tohoku J. Exp. Med.* 220:247–258.

Tidière M, Gaillard J-M, Müller DWH, Lackey LB, Gimenez O, Clauss M, Lemaître J-F. 2015. Does sexual selection shape sex differences in longevity and senescence patterns across vertebrates? A review and new insights from captive ruminants. *Evolution* 69:3123–3140.

Torgerson DG, Kulathinal RJ, Singh RS. 2002. Mammalian sperm proteins are rapidly evolving: evidence of positive selection in functionally diverse genes. *Mol. Biol. Evol.* 19:1973–1980.

Tsai-Morris C-H, Sheng Y, Gutti RK, Tang P-Z, Dufau ML. 2010. Gonadotropin-regulated testicular RNA helicase (GRTH/DDX25): a multifunctional protein essential for spermatogenesis. *J. Androl.* 31:45–52.

Vacquier VD, Swanson WJ. 2011. Selection in the rapid evolution of gamete recognition proteins in marine invertebrates. *Cold Spring Harb. Perspect. Biol.* 3:a002931.

Vladić TV, Afzelius BA, Bronnikov GE. 2002. Sperm Quality as Reflected Through Morphology in Salmon Alternative Life Histories. *Biol. Reprod.* 66:98–105.

Warnefors M, Kaessmann H. 2013. Evolution of the correlation between expression divergence and protein divergence in mammals. *Genome Biol. Evol.* 5:1324–1335.

Wey TW, Vrana PB, Mabry KE. 2017. Mating system as a possible driver of behavioral diversity in Peromyscus. *Behav. Ecol. Sociobiol.* 71:163.

Wik L, Karlsson M, Johannesson H. 2008. The evolutionary trajectory of the mating-type (mat) genes in Neurospora relates to reproductive behavior of taxa. *BMC Evol. Biol.* 8:109.

Wilson S, Fan L, Sahgal N, Qi J, Filipp FV. 2017. The histone demethylase KDM3A regulates the transcriptional program of the androgen receptor in prostate cancer cells. *Oncotarget* 8:30328–30343.

Winkler L, Kirch LM, Reinhold K, Ramm SA. 2019. Impact of low sperm competition on male reproductive trait allometries in a bush-cricket. *BMC Evol. Biol.* 19:185.

Wong EWP, Cheng CY. 2009. Polarity proteins and cell-cell interactions in the testis. *Int. Rev. Cell Mol. Biol.* 278:309–353.

Xia X, Millar JS. 1991. Genetic evidence of promiscuity in Peromyscus leucopus. *Behav. Ecol. Sociobiol.* 28:171–178.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* 24:1586–1591.

Yi L, Pimentel H, Bray NL, Pachter L. 2018. Gene-level differential analysis at transcript-level resolution. *Genome Biol.* 19:53.

Young RL, Ferkin MH, Ockendon-Powell NF, Orr VN, Phelps SM, Pogány Á, Richards-Zawacki CL, Summers K, Székely T, Trainor BC, et al. 2019. Conserved transcriptomic profiles underpin monogamy across vertebrates. *Proc. Natl. Acad. Sci. U. S. A.* 116:1331–1336.

Yuan S, Liu Y, Peng H, Tang C, Hennig GW, Wang Z, Wang L, Yu T, Klukovich R, Zhang Y, et al. 2019. Motile cilia of the male reproductive system require miR-34/miR-449 for development and function to generate luminal turbulence. *Proc. Natl. Acad. Sci. U. S. A.* 116:3584–3593.