Physical Characterization of Triolein and Implications for Its Role in Lipid Droplet Biogenesis

Siyoung Kim[1] and Gregory A. Voth[2]*

1) Pritzker School of Molecular Engineering, The University of Chicago, Chicago, IL, 60637 USA

2) Department of Chemistry, Chicago Center for Theoretical Chemistry, James Franck Institute, and Institute for Biophysical Dynamics, The University of Chicago, Chicago, IL 60637, USA

*Corresponding Author: gavoth@uchicago.edu

ABSTRACT

Lipid droplets (LDs) are neutral lipid storing organelles surrounded by a phospholipid (PL) monolayer. At present, how LDs are formed in the endoplasmic reticulum (ER) bilayer is poorly understood. In this study, we present a revised triolein (TG) model, the main constituent of the LD core, and characterize its properties in a bilayer membrane to demonstrate the implications of its behavior in LD biogenesis. In all-atom (AA) bilayer simulations, TG resides at the surface, adopting PL-like conformations (denoted in this work as SURF-TG). Free energy sampling simulation results estimate the barrier for TG relocating from the bilayer surface to the bilayer center to be ~2 kcal/mol in the absence of an oil lens. Conical SURF-TG is able to modulate membrane properties by increasing PL ordering, decreasing bending modulus, and creating local negative curvature. The other conical lipid, dioleoyl-glycerol (DAG), also reduces the membrane bending modulus and populates the negative curvature regions. A phenomenological coarse-grained (CG) model is also developed to observe larger scale SURF-TG-mediated membrane deformation. The CG simulations confirm that TG nucleates between the bilayer leaflets at a critical concentration when SURF-TG is evenly distributed. However, when one monolayer contains more SURF-TG, the membrane bends toward the other leaflet. The central conclusion of this study is that SURF-TG is a negative curvature inducer, as well as a membrane modulator. To this end, a model is proposed in which the accumulation of SURF-TG in the luminal leaflet bends the ER bilayer toward the cytosolic side, followed by TG nucleation.

INTRODUCTION

Lipid droplets (LDs) are energy- and lipid-storing organelles surrounded by a phospholipid (PL) monolayer.[1-4] Enriched with neutral lipids such as triacylglycerol (TG) or sterol esters in their core, LDs are known to be formed from the endoplasmic reticulum (ER). However, little is known about LD biogenesis in spite of its implications for a number of metabolic diseases. The current model posits three distinct steps[5-7]: 1) synthesis of TG by diacylglycerol o-acyltransferase (DGAT) at the ER, 2) phase separation of TG at a critical concentration to form nascent LDs, and 3) LD budding toward the cytosol. From nucleation to budding, the lipid droplet assembly complex is known to mediate LD biogenesis. Afterwards, the complex is separated into the ER resident protein seipin and the lipid droplet assembly factor (LDAF1).[8] The absence of this complex results in ectopic LDs; however, how this machinery catalyzes LD formation remains poorly understood.[9]

Computer simulations have clearly proven to be highly valuable for the study of biomolecules such as lipid bilayers and proteins at the scale of up to ~100 M atoms and ~1 milliseconds, largely enabled by rapid advances in hardware, software, enhanced sampling, and supporting theories.[10] As such, molecular dynamics (MD) simulations have provided valuable insights on both dynamical and structural properties. For instance, the physical properties of lipid bilayer membranes have been extensively studied involving varying sizes and lipid compositions, as well as lipid-protein interactions at various resolutions (see, e.g., refs 11-15). In contrast, computational studies of LDs are only just beginning to receive attention; this is partly because the available force fields do not support certain neutral lipids. For this reason, prior computational studies have produced a new TG topology with or without a new parameter set. For instance, the Vanni group has been incorporating the Shinoda-DeVane-Klein (SDK) force field in their LD studies, which involves the parametrization of the glycerol group to reproduce interfacial tension at the water interface.[16-19] The more common (and easier) approach is to create a new TG topology by replacing the head group of a PL with one of its tails without creating a new parameter set. This approach has been used in all-atom (AA) MD[20-23] and coarse-grained (CG) MD simulations, the latter run with the MARTINI CG force field.[24-27]

Recent 10 µs-long AA simulations of LDs report surface-oriented TG (called SURF-TG in this work),[22] which was also implicated in NMR experiments.[28] The relative amount of SURF-TG to PLs, which ranges from 5-8% in the unstressed LD[22] to ~20% in the stressed LD,[23] has been shown to modulate the area per lipid (APL) and interdigitation with PLs. The display of SURF-

TG at the LD surface creates a unique composition and defect type that does not exist in other bilayer-bounded organelles, thereby providing a plausible explanation for LD-specific targeting or conditional targeting by certain proteins. For instance, the amount of SURF-TG increases in expanding LDs to reduce the surface tension.[23] The highly amphipathic, autoinhibitory motif of CTP:phosphocholine cytidylyltransferase (CCT) is predicted computationally to sense defects created by SURF-TG and preferentially binds to those defects when LDs are expanding.[23] Once fully associated, the insertion depth of the Trp278 of CCTα is aligned with the glycerol moiety of SURF-TG to form hydrogen bonds, which may increase the binding affinity.[23] Although additional research will be needed to confirm that surface-oriented neutral lipids at the LD surface promotes specificity in LD targeting, this latter CCT binding simulation study reinforces the possible importance of SURF-TG in peptide targeting.

In this paper, the focus is not on investigating SURF-TG as a stress reducer and peptide-targeting mediator but instead on SURF-TG as a membrane property modulator. In particular, the way in which its molecular geometry is implicated in LD biogenesis is discussed. Lipids that have a conical shape such as phosphatidylethanolamine (PE), diacylglycerol (DAG), or TG do not usually form a lamellar structure by themselves.[29-31] One notable example is 2,3-dioleoyl-D-glycero-1-phosphatidylcholine (DOPC) vs 2,3-dioleoyl-D-glycero-1-phosphatidylethanolamine (DOPE). The only difference between those two molecular structures is the head group, with DOPC having $-CH_2-N(CH_3)_3^+$ in comparison to the $-CH_2-NH_3^+$ head group of DOPE. Therefore, a single DOPC molecule features a cylindrical shape, while DOPE has a conical shape. This difference in molecular topology leads to two distinct phases when DOPC and DOPE exist in single-component bulk systems: DOPC forms a planar bilayer while DOPE mostly forms an inverse hexagonal phase.[32-33] The physical properties of these two phases have been characterized experimentally[32, 34-36] and computationally.[37] Compared to DOPC or DOPE, DAG is more conical because of the absence of the head group, and SURF-TG is even more so due to the addition of the acyl chain. Using AA-MD simulations, we demonstrate here that these conical lipids are able to modulate membrane properties, such as bending modulus, and populate negative curvature regions of the membrane.

The present study also describes the development of a phenomenological CG model for PL and TG to access larger effective simulation time scales, wherein we show that TG nucleates at a critical TG concentration between the leaflets as opposed to the asymmetric distribution of SURF-

TG, which induces membrane bending first. Collectively, we demonstrate that SURF-TG is a negative curvature inducer. To this end, we propose a model in which SURF-TG-driven bending helps budding and determines the directionality of LD budding.

METHODS

AA-MD simulations and system setup

In this study, dioleoyl-glycerol (DOGL) and triolein were used to represent DAG and TG, respectively. A system was prepared using either the CHARMM-GUI membrane builder[38-41] or PACKMOL.[42] The details for each simulation and system are summarized in Table 1. The TG molecular structure used for the PACKMOL input was based on a prior LD simulation featuring a trident structure.[22] Three different TG models were used (Fig. 1). The first one was made based on the CHARMM36 (C36) lipid force field[43] by replacing the head group of DOPE with its *sn-1* acyl chain. However, the structural and thermodynamic properties of nonpolar molecules are sensitive to the cutoff distance of Lennard-Jones (LJ) interactions, which was discussed in ref 44. Indeed, the bulk TG simulations, using the 1.2 nm cutoff distance of LJ interactions, which is consistent with C36, resulted in the low density and low surface tension at the vacuum interface (See Fig. 2 in the Results). Recently, a LJ cutoff-free version of C36 (C36/LJ-PME) was released.[45-46] The second TG model was made based on the C36/LJ-PME parameters in the same fashion used in the first model. However, both first and second models suffer from the low surface tension at the water interface (Fig. 2 in the Results). Therefore, we introduced the revised TG model (C36/LJ-PME-r), which was made based on C36/LJ-PME, however, with the reduced partial charge distribution at the ester group (Fig. 1). The partial charges of PL were unchanged. Therefore, for the simulations that do not include TG, C36/LJ-PME and C36/LJ-PME-r are the same. All of those TG topologies are available in https://github.com/ksy141/TG. Of note, DAG is also not included in the original C36 force field release; nonetheless, the CHARMM-GUI developers (The Im group) created a new topology for DAG based on the existing parameter set.

Figure 1. The glycerol moiety of three TG models with partial charges. The revised TG model (C36/LJ-PME-r) has a significantly reduced charge distribution in the ester group (shown in red).

Table 1. Description of AA simulations.

AA Simulations	Lipid composition [a]	Time	Ensemble	Replicas	Setup
Bulk TG	216 TG	100 ns	NPT	6 [b]	PACKMOL
Bulk TG-vacuum	216 TG	200 ns	NVT	6 [b]	From above sim.
POPC bilayer	128 POPC	200 ns	NPT	1	PACKMOL
POPC + TG bilayer	128 POPC + 1 TG	1 μs	NPT	1	PACKMOL
POPC + TG bilayer	128 POPC + 1 TG	300 ns	NPT + biased	12 [c]	From above sim.
POPC + TG bilayer	128 POPC + 1 TG	2 μs	NPT + biased	5	From above sim.
POPC + TG bilayer	124 POPC + 8 TG	1 μs	NPT	1	PACKMOL
POPC + TG bilayer	1984 POPC + 128 TG	600 ns	NPT	1	From above sim.
POPC bilayer	4050 POPC	300 ns	NPT	1	CHARMM-GUI
POPC + DAG bilayer (referred to as DAG 10%)	3200 POPC + 320 DAG	1 μs	NPT	1	CHARMM-GUI
POPC + DAG bilayer (referred to as DAG 20%)	3200 POPC + 640 DAG	1 μs	NPT	1	CHARMM-GUI
POPC + DAG bilayer (referred to as DAG 30%)	3200 POPC + 960 DAG	1 μs	NPT	1	CHARMM-GUI
DAG 30% membrane deformation and then NPAT	3200 POPC + 960 DAG	20 ns + 1 μs	NPγT and then NPAT	1	From above sim.
DOPC bilayer	128 DOPC	200 ns	NPT	1	CHARMM-GUI

a) The total number of lipid molecules of each type in a system.

b) With varying LJ cutoff distances

c) 12 windows

All simulations were run by GROMACS 2018 and 2020[47] with the C36,[43] C36/LJ-PME,[45-46] or C36/LJ-PME-r force field. Simulations using C36 used the 1.2 nm LJ cutoff distance with a force-switching function between 1.0 nm to 1.2 nm, unless otherwise noted. All simulations including DAG were performed with C36. Simulations using C36/LJ-PME and C36/LJ-PME-r used the real-space cutoff distance of 1.0 nm. Biased simulations (described below) were performed with the external plugin, PLUMED2 (v.2.6).[48] Simulations were evolved with a 2-fs timestep. The long-range electrostatic interactions were evaluated with the Particle Mesh Ewald algorithm,[49] with the distance cutoff set to be the same as the LJ interaction. Every bond involving a hydrogen atom was constrained using the LINCS algorithm.[50] For our constant pressure simulations, the Parrinello-Rahman barostat was used to maintain a pressure of 1 bar with a compressibility of 4.5×10^{-5} bar^{-1} and a coupling time constant of 5 ps.[51] For membrane simulations, the pressure was coupled semi-isotropically, whereas during bulk simulations the pressure was coupled isotropically. For constant area simulations (NPAT), only the pressure of the Z dimension was controlled. The Nose-Hoover thermostat was used to maintain a temperature of 310 K with a coupling time constant of 1 ps.[52-53] In order to deform the DAG 30% membrane, a constant surface tension simulation (NPγT) was conducted for 20 ns with the Berendsen thermostat and barostat

using the same parameters.[54] The pressure was coupled semi-isotropically with the pressure in the X and Y dimension set to -100 bar during membrane deformation.

Density, isothermal compressibility, and surface tension

Density was obtained with the GROMACS analysis tool, which was outputted every 2 ps. Isothermal compressibility was evaluated from

$$\beta = -\frac{1}{V}\left(\frac{\partial V}{\partial p}\right)_T = \frac{1}{k_B T}\frac{\langle V^2 \rangle - \langle V \rangle^2}{\langle V \rangle} \quad (1)$$

where the bracket represents the ensemble average and V denotes the volume of the system. Similarly, GROMACS was used to analyze the surface tension at the vacuum and water interfaces. Bilayer area compressibility was determined using the following equation

$$K_A = k_B T \frac{\langle A \rangle}{\langle A^2 \rangle - \langle A \rangle^2} \quad (2)$$

where *A* is the instantaneous total area of a membrane.[55]

Order parameters

Order parameters were evaluated from $S_{CD} = 0.5 \times |\langle 3\cos^2\theta - 1 \rangle|$, where the angle ($\theta$) is between the Z axis and the position vector of an acyl carbon atom to a bonded hydrogen atom; the bracketed portion indicates the ensemble average.[56-58]

Potential of mean force of TG flip-flop

In order to estimate the preferential location of TG in a bilayer and its tendency to become SURF-TG, the potential of mean force (PMF) of the TG as a function of its Z position in the bilayer was calculated. A 3-palmitoyl-2-oleoyl-D-glycero-1-Phosphatidylcholine (POPC) bilayer membrane consisting of 128 POPC molecules and one TG molecule was prepared. Biased simulations were performed with the collective variable representing the Z distance between the center of the mass of phosphorus atoms and the center of the mass of the glycerol moiety of the TG molecule. Replica-exchange[59] umbrella sampling[60] (REUS) simulations were run for 300 ns with a harmonic restraint of a force constant of 47.8 kcal/mol/nm² for each window. The total of 12 windows was prepared with a 0.2 nm spacing over a range of 0.0 to 2.2 nm (0 is the average Z position of the center of the mass of the phosphorus atoms, which is comparable to the center of

the membrane). An initial structure for each window was prepared with a steered MD simulation. The exchange between windows was attempted every 2,000 steps. The weighted histogram analysis method (WHAM) was used for the calculation of the PMF with a bin spacing of 0.0275 nm.[61] The Grossfield WHAM implementation was used.[62] The simulations were divided into five equal-length blocks to estimate the errors. The PMF was calculated for each block and then the standard deviation of block PMFs was reported as the errors. To support the PMF result, we also carried out efficient, faster converging transition-tempered metadynamics (TT-MetaD) simulations with the same collective variable.[63-64] The Gaussian hill was deposited every 5K steps at a height of 0.29 kcal/mol and width of 0.08 nm or 0.1 nm. The bias factor of 6 and threshold of 2.4 kcal/mol for TT-MetaD were used. The position of the two basins were set to -1.1 nm and +1.1 nm, when the center of the mass of phosphorus was set to zero. Five simulation replicas, each 2 μs long, were performed to estimate the averaged potential of mean force (PMF). The errors represent the standard deviation of the PMFs obtained from the five replicas.

CG simulations

In order to study SURF-TG-driven membrane deformation and nucleation, a new phenomenological implicit solvent CG model was developed for the PL and TG. Each PL and TG molecule consisted of 11 and 13 CG beads, respectively. Each acyl chain was composed of 4 CG beads, with 2 different CG types. The approximate molecular groupings of the PL and TG structures are shown in Fig. S1. In this model, the attraction was modeled with the Gaussian function, $-A\,exp(-Br^2)$ where $B = 2\,nm^{-2}$, and the repulsion with the repulsive LJ interaction, $4\varepsilon\,(\sigma/r)^{12}$ where $\varepsilon = 0.0028\,kcal/mol$. No CG bead carried any charge. The attraction only occurs between hydrophobic CG beads (beads in the acyl chain or CT1 and CT2 in Fig. S1 of the Supporting Information), between the glycerol CG beads of TG, and between the glycerol CG beads of TG and PL (No attraction exists between the glycerol CG beads of PL); in contrast, repulsion occurs between all CG beads unless the pair is bonded. The attraction and repulsion parameters, A and σ, were primarily parameterized to reproduce the radial distribution function (RDF) from the mapped AA (C36) RDF (Figs. S2 and S3). Unless otherwise indicated, the repulsion parameter (σ) of a pair composed of two different types of CG beads was determined using an arithmetic average, while the corresponding attraction parameter (A) for such a pair was set explicitly. All CG bonds were modeled with the harmonic potential, with the equilibrium

distance of 0.5 nm and the force constant of 5 kcal/mol. The angle potentials were modeled with the harmonic potential with a small force constant of 0.12 kcal/mol. However, we note that the properties of PL and TG in general were not sensitive to the bond and angle force constants. The details of this CG model can be found in Fig. S1. The potentials and inputs are available at https://github.com/ksy141/SK_CGFF.git.

All CG simulations were conducted with GROMACS 2018 using a 20-fs timestep.[47] The cutoff distance of 2.4 nm was used for non-bonded interactions. Temperature was controlled via velocity rescaling with a stochastic term with the target temperature of 310 K and a time constant of 1 ps.[65] A pressure of 1 bar was maintained with the Parrinello-Rahman barostat with a compressibility of 4.5×10^{-5} bar^{-1} and a coupling time constant of 10 ps. For the PL or PL+TG bilayer simulation, the pressure was semi-isotopically coupled with the X and Y dimensions but was not coupled with the Z dimension (by setting compressibility in the Z dimension to 0). For the bulk TG simulation, the pressure was isotopically coupled. Finally, for bilayer simulations that have asymmetric distribution of TG between two leaflets, weak pressure and temperature couplings were used to minimize perturbations of the evolution of the membrane remodeling. For those simulations that convey membrane deformation, time constants of 1 ns and 10 ns for temperature and pressure were used, respectively. Note that a previously reported implicit solvent CG model that detected membrane deformation used weak couplings as well.[66] The initial structures were prepared using PACKMOL[42] and MDAnalysis[67]; simulation details are provided in Table 2.

Table 2. Description of CG simulations.

CG Simulations	Lipid composition [a]	Nsteps	Ensemble	Replicas	To compute
Bulk TG	1000 TG	5M	NPT	1	ρ, β, RDF
Bulk TG-water [b]	1000 TG	1.5M	NVT	1	γ_{wat}
PL bilayer	200 PL	1.5M	NPT	1	K_A, APL, RDF
PL bilayer	7200 PL	5M	NPT	1	K_C
PL + TG bilayer (2%) [c]	1764 PL + 36 TG	50M	NPT	3	Nucleation
PL + TG bilayer (5%) [c]	1710 PL + 90 TG	50M	NPT	3	Nucleation
PL + TG bilayer (1%) [d]	7128 PL + 72 TG	5M	NPT [e]	2	Bending
PL + TG bilayer (5%) [d]	6840 PL + 360 TG	5M	NPT [e]	2	Bending

a) The total number of lipid molecules of each type in a system.
b) For implicit CG models, it is ambiguous to distinguish water and vacuum interfaces. Specific to TG, the surface tensions at those two interfaces happen to be comparable (See Fig. 2).
c) Symmetric SURF-TG distribution
d) Asymmetric SURF-TG distribution
e) Weak temperature and pressure couplings

Analysis and visualization

Simulations were analyzed using MDAnalysis[67] and GROMACS.[47] Unless otherwise noted, the standard errors (*se*) were reported in this work. Equilibrated trajectories were first divided into *M* blocks of equal length and the block average for each block was calculated. The standard errors (*se*) were estimated by $se = 1/\sqrt{M} \, sd$, where *sd* is the standard deviation of the block averages. Five blocks ($M = 5$) were used in this work. Molecular images were captured using PyMOL.

RESULTS

AA-MD

Physical properties of bulk TG

In this section, we explore the quality of three TG models (C36, C36/LJ-PME, and C36/LJ-PME-r; see Fig. 1 and Methods) by characterizing four physical properties of bulk TG (Fig. 2): density (ρ), isothermal compressibility (β), surface tension at the vacuum (γ_{vac}), and surface tension at the water interface (γ_{wat}). With a prior study demonstrating that the first three properties are sensitive to the LJ cutoff distances,[44] we performed MD simulations (C36) of a bulk TG consisting of 216 TG molecules with varying LJ cutoff distances. The computed density and isothermal compressibility of the bulk TG were shown to converge when the LJ cutoff distance was increased from 1.0 nm to 2.0 nm (Fig. 2). The density of the bulk TG approached the experimental value

Figure 2. Physical properties of the bulk TG evaluated with three different TG models. For C36, the properties were calculated as a function of LJ cutoff distances and the force switching function was applied between $r_c -$ 0.2 nm and r_c. For the surface tension at the water interface, the values were averaged every 100 ns. C36 (1.2 nm) represent the simulation carried out with C36 using the cutoff distance of 1.2 nm.

(0.8991 g/cm^3 at 313 K)[68] when simulated with the LJ cutoff of 2.0 nm (0.898 g/cm^3). The same calculations were carried out with C36/LJ-PME and C36/LJ-PME-r. The extended LJ interactions led to a higher density in the C36/LJ-PME results; however, the extra increase became compensated by the reduced charge distribution in C36/LJ-PME-r. The same behavior was observed in the RDF of the TG glycerol moiety (Fig. S4). While C36/LJ-PME increases the first peak due to the increased LJ range, the reduced charge distribution in C36/LJ-PME-r lowers the increased peak. Next, we performed MD simulations of the TG-vacuum interface by adding an empty space in the Z dimension to the bulk TG system. Consistent with density and isothermal compressibility, the surface tension of the TG-vacuum interface converges with the increased LJ cutoff distance and also agrees reasonably well with the results of C36/LJ-PME and C36/LJ-PME-r. In all cases, no TG molecules were released from the bulk TG. Although there are no experimental TG data for β and γ$_{vac}$, the results of C36 with the 2.0 nm cutoff distance, C36/LJ-PME, and C36/LJ-PME-r show the reasonable agreement for ρ, β, and γ$_{vac}$.

On the other hand, the experimental surface tension at the water interface (32 mN/m)[69] was not reproduced except for C36/LJ-PME-r (Fig. 2). The surface tension became reduced to < 20 mN/m when using C36 or C36/LJ-PME-r, verifying the necessity of reducing the distribution of partial charges in the ester group. Although the additive force field cannot take into account polarization therefore cannot accurately produce the water density in nonpolar environment unless further modified,[70-71] it is worth nothing the large difference found in the water density at the TG layer. While C36 and C36/LJ-PME demonstrate the high water density (~10^{-2} g/cm^3), C36/LJ-PME-r shows the low water density (~10^{-3} g/cm^3). Although no experimental measurement of the water amount in the pure TG is available, a wide range of water density from 3×10^{-4} g/cm^3 to up to 7×10^{-3} g/cm^3 was reported in olive oil.[72-74]

TG flip-flop

We characterized the physical properties of TG in a bilayer membrane in order to investigate its implication in LD biogenesis. A bilayer membrane containing 128 POPC molecules and 1 TG molecule was prepared. We then calculated the PMF of the TG molecule (Fig. 3) as a function of its Z position within a bilayer membrane by performing the REUS simulations. Both C36 and C36/LJ-PME-r demonstrate the stability of SURF-TG in the absence of an oil lens although the C36 result overestimates the stability at the surface by ~1.5 kcal/mol compared to the

C36/LJPME-r result. The preferential location of the TG molecule is slightly different by 0.2 nm as well. We further verified our results with another biased methodology. Five replicas of TT-MetaD simulations were carried out with the same collective variable, each run for 2 µs. The PMFs obtained from the REUS and TT-MetaD simulations show good agreement (Fig. S5). Finally, the 1-µs unbiased MD simulation behaves as consistent with our findings. For the C36 simulation, the

TG molecule was initially located at the center of the bilayer (CORE-TG), however, became SURF-TG rapidly (< 10 ns). During 1 µs, the TG molecule did not flip-flop but resided at the same leaflet. In contrast, the TG molecule visited the membrane center more frequently and eventually flip-flopped at ~820 ns with C36/LJ-PME-r. Taken together, both unbiased and biased results suggest that a single TG molecule is principally SURF-TG in a bilayer membrane, adopting the PL-like conformation, in the absence of an oil lens.

Figure 3. PMF of a TG molecule in a POPC bilayer as a function of the Z distance from the bilayer center (TG flip-flop), calculated from the REUS simulations. The Z position of the center of the phosphorus atoms was zeroed. The errors represent the standard deviation of the PMFs obtained from the five equal-length blocks.

Local membrane deformation by SURF-TG

Conical molecules such as PE or DAG are known to be responsible for amphipathic peptide targeting by creating packing defects.[75-79] However, the precise mechanisms for how those molecules modulate membrane properties are poorly understood. The molecular shape of SURF-TG is inevitably conical, with the glycerol moiety forming the vertex of the cone and the end group of three acyl chains forming a flat base while POPC features a quasi-cylindrical shape. In order to study how SURF-TG modulates the local properties of bilayers, we computed the height field of the phosphorus atoms of the upper leaflet in the bilayer containing 128 POPC molecules and 1 TG molecule. Histogram analysis was used for the 1-µs C36/LJ-PME-r simulation to investigate the relative positions of the phosphorus atoms of the upper leaflet with respect to the glycerol moiety of the SURF-TG molecule (Fig. 4). Approximately one-fifth of the frames were not used for this analysis because in those frames SURF-TG transitioned to CORE-TG or resided at the other leaflet

(flip-flop occurred at ~820 ns). The height of the phosphorus atoms near the SURF-TG molecule can be as low as ~0.15 nm compared to those of the other phosphorus atoms (Fig. 4). The range of the local deformation created by one SURF-TG molecule can be more than 2 nm in both the X and Y dimensions (Fig. 4). We anticipate that if SURF-TG molecules cluster at the surface there will be stronger and longer PL packing discontinuity. We confirmed that the same conclusion can be made with the C36 trajectory (data not shown). Our data indicate that SURF-TG locally pulls PLs toward the bilayer center, thereby creating local negative curvature. How this process serves to remodel bilayer membranes will be discussed later via our CG simulation results.

Figure 4. *(left)* Height field, z(x, y), of the phosphorus atoms of the upper leaflet respective to the glycerol moiety of the SURF-TG molecule (circle at the origin). Frames in which SURF-TG resided at the upper leaflet were used for the analysis. *(right)* Snapshot demonstrating the local deformation by SURF-TG. The red spheres are the phosphorus atoms and PLs are shown in blue lines. The glycerol moiety and acyl chains of TG are shown with green and yellow spheres, respectively. The simulation was carried out with C36/LJ-PME-r.

Molecular and hydration properties of SURF-TG

In order to characterize the molecular and hydration properties of SURF-TG, a new POPC bilayer membrane containing 6% TG (62 POPC and 4 TG molecules per leaflet) was prepared and compared with a pure POPC bilayer membrane (64 POPC molecules per leaflet). The system contained a higher concentration of TG than the critical nucleation concentration, which was reported to be 2.8% g TG or 2.4% mol TG.[28] However, within the AA simulation timescale, TG nucleation did not occur, and all the TG molecules remained SURF-TG most of the times. The simulations were carried out with C36/LJ-PME-r. For a pure POPC bilayer membrane, C36/LJ-PME and C36/LJ-PME-r are equivalent.

Figure 5. Order parameters of *sn-1* (top) and *sn-2* (bottom) chains. The POPC order parameters of the pure POPC bilayer (served as a reference) are shown in the dashed lines. The POPC order parameters of the POPC + TG bilayer are indicated in the continuous lines. The TG order parameters of the same system are shown in squares. The *sn-3* chain of TG is not included for visual clarity. The simulations were carried out with C36/LJ-PME-r.

The order parameters of SURF-TG and POPC were computed (Fig. 5). Interestingly, the order parameters of POPC in the POPC+TG bilayer (continuous lines) were found to be larger than those of POPC in the pure POPC bilayer (dashed lines), suggesting that SURF-TG increases the order of POPC molecules. As reported in a previous study, the increase in PL ordering in LDs compared to the bilayers can be attributed to the fact that CORE-TG interdigitates with the PL monolayer and increases the density of the low-density tail region.[22] Similarly, short LD simulations lacking any SURF-TG also demonstrated increased PL ordering.[80] Our results suggest that SURF-TG increased PL ordering as well. The SURF-TG order parameters have the same trend with PL; however, they are more reduced than PL. The reduced order parameters of TG compared to PL can be attributable to the higher degree of freedom in the Z dimension of TG molecules. While PL moves little in the Z dimension and therefore its order parameters are usually determined by the lipid-packing density in the XY dimensions, TG is relatively free in moving in the Z dimension.

The hydration properties of SURF-TG were evaluated in the same POPC+TG bilayer. It was suggested that the SURF-TG has the primary carbonyls in *sn-1* and *sn-3* chains that are more exposed to water than the secondary carbonyl in *sn-2* chain, which would explain specificity for hydrolysis at the primary carbonyl position.[28] The computed RDF between each TG oxygen atom (O11, O21, and O31) and water shows reduced accessibility of water to the secondary carbonyl

(Fig. 6). However, the RDFs between the other oxygen atoms (O12, O22, O32) and water were identical at the first peak (data not shown).

Figure 6. The RDFs of TG oxygen atoms (O11, O21, O31) with the oxygen atom of water. The structure of the TG glycerol moiety is illustrated in the inlet. The simulation was carried out with C36/LJ-PME-r.

Impact of conical molecules on membrane properties

Figure 7. Bending modulus of five POPC bilayer membranes, containing different amounts of DAG or TG. The number of POPC and DAG or TG molecules in each system is as follows: POPC (POPC 4050), DAG 10% (POPC 3200 + DAG 320), DAG 20% (POPC 3200 + DAG 640), DAG 30% (POPC3200 + DAG 960), TG 6% (POPC 1984 + TG 128). The errors represent the standard errors.

In order to systematically study the impact of conical lipids on bending modulus (K_C), we assembled large POPC bilayer membranes of varying DAG concentrations ranging from 0% to 30% and computed the bending modulus using the undulation spectrum.[81-83] On our simulation timescale (1 μs), there was no evidence for phase separation in the DAG, despite its high concentration. With an increased concentration of DAG, the bending modulus decreased (Fig. 7), suggesting that DAG lowers bending rigidity. We also confirmed a similar correlation for the bilayer containing 6% SURF-TG with both

C36 and C36/LJ-PME-r (Fig. 7). The bending modulus of this particular bilayer was determined to be $23.1 \pm 1.0\ k_BT$ (C36) and $22.4 \pm 2.5\ k_BT$ (C36/LJ-PME-r), which is smaller than $32\ k_BT$ (ref 12) or $36 \pm 5\ k_BT$ (Fig. 7), the bending modulus of the pure POPC bilayer.

Next, constant surface tension was applied semi-isotropically for 20 ns to induce deformation of the 30% DAG membrane. The resulting structure displayed a buckling that resembled a cubic function in the X axis (Fig. 8a). We then ran constant area simulation (NPAT) for 1 μs, confirming that the overall buckling was maintained (Fig. 8a and 8b). As shown in Fig. 8b, the height field region in the upper leaflet where X is between 5 nm and 15 nm displayed a negative curvature; conversely, the region where X is between 25 nm and 30 nm displayed a positive curvature. In contrast, the curvature becomes inverted in the lower leaflet (Fig. S6). Consistent with our expectations, a conical DAG is less populated at the positive curvature (for instance, the region where X is between 25 nm and 30 nm in the upper leaflet), but more populated at the negative curvature (Fig. 8c and Fig. S6). The central conclusion derived from these simulations is that a conical lipid reduces the bilayer bending modulus and behaves as a negative curvature inducer.

Figure 8. Buckled POPC+DAG membrane (DAG 30%). a) The snapshot of the MD simulation. Water is in red and DAG in blue. b) The height field, $z(x,y)$, of the upper leaflet, averaged over the trajectory. The averaged Z position of the upper phosphorous atoms was zeroed. c) The DAG distribution, $p(x,y)$, in the upper leaflet, averaged over the trajectory.

CG-MD

TG bulk and PL bilayer

In order to further investigate the properties of SURF-TG in a bilayer membrane and establish their implications in LD biogenesis, we developed a new implicit solvent CG model for PL and TG as noted earlier. In this model, each PL and TG molecule consisted of 11 and 13 CG atoms, respectively, where each acyl chain was composed of 4 CG atoms and of 2 CG types. No charges were associated with CG beads. The interactions were modeled with a Gaussian function for attraction, and the LJ repulsive potential for repulsion. The details of the CG model can be found in Fig. S1.

The attraction and repulsion parameters were then primarily parameterized to reproduce the RDFs from the AA trajectories with the atoms mapped on the CG sites of the CG model (the so-called "mapped AA model"). A PL bilayer simulation consisting of 200 residues was performed for 1.5M steps, after which the computed 2-dimensional RDFs were compared with those from the mapped AA trajectory provided in Fig. S2. Similarly, a bulk TG simulation consisting of 1,000 residues was conducted for 5M steps; the computed RDFs from this trajectory were compared with those from the mapped AA trajectory shown in Fig. S3. The results showed reasonable agreement between the CG RDFs and the mapped AA RDFs, which were achieved both in the PL bilayer and bulk TG systems. We then determined the physical properties of the bulk TG and PL bilayer from the same simulations as shown in Table 3. Although the isothermal compressibility of the bulk TG obtained from the CG simulation was found to be a factor of three larger, and the bending modulus of the PL bilayer was recorded to be 1.4 times higher than the AA results, the remaining properties agreed well with the experimental or AA data.

Table 3. Physical properties of DOPC bilayer and TG bulk from experiments, AA simulations, and CG simulations. Standard errors are given for the simulations that we performed.

		Exp.	AA	CG
DOPC bilayer				
	APL $[nm^2]$	0.674[a], 0.724[b]	0.70 ± 0.00	0.72 ± 0.00
	K_A $[mN/m]$	300[c]	251 ± 20, 290[g]	307 ± 9
	K_C $[k_B T]$	20.75[d], 20.26[e], 26.6[f]	29[g]	41 ± 1
TG bulk				
	ρ $[g/cm^3]$	0.8991[h]	0.90 ± 0.00[i]	0.90 ± 0.00

$\beta \times 10^{-9}\ [m^2/N]$	N/A	$0.88 \pm 0.01^{i)}$	2.33 ± 0.07
$\gamma_{wat}\ [mN/m]^{j)}$	$32^{k)}$	$32 \pm 0^{i)}$	38 ± 0

a) ref [84] b) ref [85] c) ref [86] d) ref [87] e) ref [88] f) ref [89] g) ref [12] h) ref [68] i) Taken from the C36/LJ-PME-r simulations. j) As it is an implicit CG model (no explicit solvent molecules), it is ambiguous to distinguish surface tension at the air interface and water interface. Specific to TG, those two values are comparable (see Fig.2) k) ref [69]

TG Nucleation

TG nucleation between ER leaflets represents the first step in LD biogenesis. A previous NMR study determined the maximum solubility of TG in a PL phase to be 2.8% g TG or 2.4% mol TG.[28] Therefore, we expected that a bilayer containing TG at a higher than critical concentration would undergo TG nucleation. Accordingly, we prepared CG bilayer systems with varying TG concentrations to study TG nucleation. Consistent with our expectations, our CG simulations confirmed that TG nucleates and forms an oil lens between the leaflets within 20M time steps when the TG concentration is 5% but does not if it is 2% (Fig. 9). In order to estimate the degree of nucleation, the nucleation % was calculated by dividing the number of TG molecules in the biggest cluster by the number of total TG molecules. If two TG molecules are within 2 nm,

Figure 9. CG simulations of a symmetric bilayer and TG nucleation. The last snapshots (50M steps) of two membranes and nucleation % are shown. The dashed lines indicate the standard deviation of the quantity from three replicas for each concentration. In the snapshots, PL head group and acyl chains are shown in light blue and dark blue, respectively. The TG glycerol moiety and TG acyl chains are shown in green and yellow, respectively.

they were assumed to be in the same cluster. Results for the 5% simulation indicate that the oil cluster is stable and recruits more TG with increasing simulation time until equilibrium is reached (Fig. 9). Importantly, TG molecules that are not in the oil cluster but dissolved in the membrane

Figure 10. CG simulation of an asymmetric bilayer and membrane bending toward the lower leaflet. a) The initial distribution of SURF-TG in the upper leaflet (red) and in the lower leaflet (blue). The upper leaflet contains more SURF-TG than the lower leaflet; the scaled X, Y coordinates were used. The regions where the local density of SURF-TG is high in the upper leaflet are circled. b) The height of the phosphorus atoms of the upper leaflet at 20K steps. The scaled X, Y, and Z coordinates were used. c) The CG-MD snapshot at 20K time steps; the upper phosphorus atoms are shown in red and the lower phosphorus atoms in blue. TG glycerol moiety and acyl chains are shown in green and yellow, respectively.

are primarily SURF-TG. This can be also seen in the 2% simulation where most of the TG molecules resided at the surface (Fig. 9).

SURF-TG-mediated membrane bending

As confirmed by our AA simulation results, SURF-TG locally induces a negative curvature. Using CG simulations, we investigated how local membrane deformation leads to global membrane deformation. First, we prepared an asymmetric bilayer wherein each leaflet contained a different number of SURF-TG molecules. For instance, in one of our CG simulations, the upper leaflet of the bilayer contained 3552 PL and 48 SURF-TG molecules, while the lower leaflet contained 3576 PL and 24 SURF-TG molecules. The biological rationale for the asymmetric distribution of SURF-TG between two leaflets will be discussed in the Discussion section later. Initially from a flat bilayer, the bilayer then rapidly (at 20K time

steps) bent toward the lower leaflet containing less SURF-TG than the upper leaflet (Fig. 10). Interestingly, one of the regions in the upper leaflet where the local concentration of SURF-TG was initially high (x = 0.6 and y = 0.1 in Fig. 10a) became the lowest point, as indicated in Fig. 10b and Fig. 10c. The other region that initially had the high local concentration of SURF-TG in the upper leaflet (x = 0.5 and y = 0.9 in Fig. 10a) became curved to the lower leaflet. In this particular simulation, TG nucleation did not occur because of the low TG concentration. Consistent with the AA results, TG remained SURF-TG, populated at the negative curvature during 5M steps (data not shown). The same simulation but containing the higher TG concentration demonstrates the same bending behavior, followed by nucleation. We then compared the timescale between membrane bending and TG nucleation, which for our CG simulations was much faster for membrane bending compared to TG nucleation. The timescale comparison can be also confirmed via AA simulation results. Specifically, in our AA simulations membrane undulation was found to occur within several nanoseconds, thereby enabling us to evaluate the bending modulus from the undulation spectrum. In comparison, the bilayer containing 5% TG did not undergo TG nucleation within the AA timescale. Both our AA and CG simulations suggest the timescale of TG nucleation is slower than that of membrane undulation. The slow timescale of TG nucleation was also shown by other CG simulations.[18-19, 24, 27] Taken together, SURF-TG will be important in the LD biogenesis because it can modulate the membrane properties before transitioning into an oil lens. In particular, membrane deformation driven by SURF-TG, which would precede TG nucleation, may determine the LD budding directionality.

DISCUSSION

Using a revised AA-TG model that reproduces the experimental surface tension at the water interface, we have examined herein how a conical lipid modulates membrane properties and induces membrane remodeling. The AA simulations demonstrate that SURF-TG decreases bending rigidity and increases PL ordering. Due to its conical shape, SURF-TG induces a local negative curvature. We also confirmed using biased simulations that the energy barrier for relocating SURF-TG to the bilayer center is ~2 kcal/mol in the absence of an oil lens. This finding is consistent with our experience that TG initially locates at the bilayer center, while becoming SURF-TG within 10 ns of unbiased MD simulations and it resides principally at the surface. Finally, we systematically demonstrated the impact of conical lipids on the membrane properties with varying DAG concentrations by determining that conical lipids populate and induce a negative curvature and reduce the bending modulus.

In order to access SURF-TG-mediated membrane remodeling, we developed a phenomenological implicit solvent CG model for PL and TG using a Gaussian function for attraction and the LJ potential for repulsion. Primarily parameterized to reproduce the RDFs from the atomistic CG mapped trajectories, the CG model was able to reproduce most of the physical properties, with the exception of the isothermal compressibility of TG, which was found to be a factor of three higher; additionally, the bilayer bending modulus was noted to be 1.4 times higher compared with our AA results. Using our phenomenological CG model, we showed TG forms an oil lens between the leaflets when the TG concentration is 5% but does not if it is 2%. We also demonstrated that asymmetric SURF-TG composition induces membrane bending because SURF-TG works as a negative curvature inducer. For instance, when the upper leaflet of a bilayer contained more SURF-TG, the membrane would bend toward the lower direction. Consistent with the AA results, TG molecules that do not belong to the oil cluster are mostly SURF-TG.

Although a related hypothesis suggesting that asymmetric surface tension determines the directionality of LD budding has recently been proposed,[90-91] the cause of any asymmetric surface tension was not linked to lipid type or lipid geometry. A possible physiological explanation for the asymmetric distribution of SURF-TG between the luminal leaflet and cytosolic leaflet of the ER bilayer can be found in the recently resolved structure of DGAT1.[92-93] Catalyzing the last step of TG synthesis, DGAT1 intakes DAG and outputs TG. The lateral gate, which is located closer to the luminal leaflet, is anticipated to be a pathway for the flow of the reactant and product. Therefore,

we can propose a model in which a newly synthesized TG molecule bends the membrane toward the cytosol and determines the budding directionality based on the following steps: 1) The synthesized TG molecule is released through the lateral gate of DGAT1. 2) The released TG molecule becomes SURF-TG since it has an energy penalty of ~2 kcal/mol when residing at the ER bilayer center. 3) Given that the lateral gate is closer to the ER luminal leaflet, it is more likely that the luminal leaflet will contain more SURF-TG than the cytosolic leaflet. 4) The accumulation of SURF-TG in the luminal leaflet bends the ER bilayer toward the cytosolic side. 5) The curved ER bilayer recruits the lipid droplet assembly complex.[94]

This report makes reference to a number of papers that have described comparable results, although using different lipids and in different contexts.[87, 95-99] In those studies, the amount of polyunsaturated fatty acids or polyunsaturated PLs were correlated with bending modulus and the distribution of those lipids. The central conclusion from those studies is that polyunsaturated acyl chains add fluidity in the Z dimension, which implies that they can be more flexibly adapted for curvature than saturated acyl chains, thereby reducing bending modulus, facilitating lipid trafficking, and modulating membrane dynamics. In this paper, TG and DAG were modeled with triolein and dioleoyl-glycerol, wherein each chain was mono-unsaturated. Therefore, we suggest that our findings are more related to the molecular shape rather than to the degree of unsaturation. Nonetheless, a future study should be designed to investigate the interplay between those two factors.

Finally, we discuss here our future work on CG modelling. In this study, a phenomenological CG model was developed via hand-tuning. A follow-on study may incorporate a bottom-up approach that systematically constructs CG models based on underlying AA interactions. Such a methodology will include the force-matching method,[100-104] relative entropy minimization,[105] or a hybrid approach that utilizes both methods.[106] Using a highly coarse-grained membrane, large-scale membrane deformations mediated by the Bin/amphiphysin/Rvs (BAR) domain-containing proteins have been successfully described with such methods.[107-111] Also, one interesting potential direction would be to introduce internal "states" to lipid CG beads using the ultra-coarse-graining (UCG) theory in order to modulate the CG interactions in different chemical or physical environments.[112-114] In particular, by assuming that the internal state dynamics remain in quasi-equilibrium,[115-117] we expect that the UCG modeling can further capture different TG solubilities at PL bilayers and LD surfaces.

CONCLUSIONS

In this study we show how neutral lipids are able to modulate the physical properties of bilayers. Results from our AA-MD bilayer simulations indicate that TG principally resides at the bilayer surface, adopting PL-like conformations, in the absence of an oil lens. SURF-TG is an innately conical lipid because of the addition of the acyl chain and the elimination of the head group compared to POPC; as such, a conical SURF-TG produces a local negative curvature and lowers bending modulus. We also find a conical molecule, DAG, populates the negative curvature and lowers bending modulus. In order to increase the accessible simulation time scale, a phenomenological CG model for PL and TG was developed by parameterizing the non-bonded interactions to reproduce the RDFs from the mapped atomistic trajectories and physical properties. In the CG simulations, TG molecules form an oil cluster when TG concentration is above the critical concentration. Our CG simulations of the bilayers, wherein each monolayer surface contained a different number of SURF-TG, confirmed SURF-TG-driven membrane deformation, which may determine the directionality of LD budding and catalyze TG nucleation. Consistent with the AA results, TG molecules that are not in the oil cluster but dissolved in the PL phase are mostly SURF-TG and populate the negative curvature regions. To conclude, this paper demonstrates how the conical shape of neutral lipids are implicated in the early step of LD biogenesis.

SUPPORTING INFORMATION

Figures S1-S6 are included in Supporting Information.

ACKNOWLEDGEMENTS

This research was supported by a grant from the National Institute of General Medical Science (NIGMS) of the National Institutes of Health (NIH), grant R01-GM063796. The computer simulations were performed on the high-performance GPU cluster (GM4) at the University of Chicago Research Computing Center, supported by NSF grant DMR-1828629 and the Stampede2 supercomputer at the Texas Advanced Computing Center (TACC) through allocation TG-MCA94P017 with resources provided by the Extreme Science and Engineering Discovery Environment (XSEDE) supported by NSF grant ACI-1548562. We thank Chenghan Li, Jeeyun

Chung, Zack Jarin, Myong In Oh, Jessica Swanson, Robert Farese, Jr., Tobias Walther, and Richard Pastor for their useful perspectives and discussions.

REFERENCES

1.   Ducharme, N. A.; Bickel, P. E., Minireview: Lipid Droplets in Lipogenesis and Lipolysis. *Endocrinology* **2008,** *149* (3), 942-949.
2.   Farese, R. V.; Walther, T. C., Lipid Droplets Finally Get a Little R-E-S-P-E-C-T. *Cell* **2009,** *139* (5), 855-860.
3.   Walther, T. C.; Farese, R. V., Lipid Droplets and Cellular Lipid Metabolism. *Annual Review of Biochemistry* **2012,** *81* (1), 687-714.
4.   Olzmann, J. A.; Carvalho, P., Dynamics and functions of lipid droplets. *Nature Reviews Molecular Cell Biology* **2019,** *20* (3), 137-155.
5.   Walther, T. C.; Chung, J.; Farese, R. V., Lipid Droplet Biogenesis. *Annual Review of Cell and Developmental Biology* **2017,** *33* (1), 491-510.
6.   Thiam, A. R.; Farese Jr, R. V.; Walther, T. C., The biophysics and cell biology of lipid droplets. *Nature Reviews Molecular Cell Biology* **2013,** *14* (12), 775-786.
7.   Wilfling, F.; Haas, J. T.; Walther, T. C.; Jr, R. V. F., Lipid droplet biogenesis. *Current Opinion in Cell Biology* **2014,** *29*, 39-45.
8.   Chung, J.; Wu, X.; Lambert, T. J.; Lai, Z. W.; Walther, T. C.; Farese, R. V., Jr., LDAF1 and Seipin Form a Lipid Droplet Assembly Complex. *Dev Cell* **2019,** *51* (5), 551-563.
9.   Wang, H.; Becuwe, M.; Housden, B. E.; Chitraju, C.; Porras, A. J.; Graham, M. M.; Liu, X. N.; Thiam, A. R.; Savage, D. B.; Agarwal, A. K.; Garg, A.; Olarte, M.-J.; Lin, Q.; Fröhlich, F.; Hannibal-Bach, H. K.; Upadhyayula, S.; Perrimon, N.; Kirchhausen, T.; Ejsing, C. S.; Walther, T. C.; Farese, R. V., Seipin is required for converting nascent to mature lipid droplets. *eLife* **2016,** *5*, e16582.
10.   Brini, E.; Simmerling, C.; Dill, K., Protein storytelling through physics. *Science* **2020,** *370* (6520).
11.   J. Sodt, A.; Logan Sandar, M.; Klaus Gawrisch; W. Pastor, R.; Edward Lyman, The Molecular Structure of the Liquid-Ordered Phase of Lipid Bilayers. *Journal of the American Chemical Society* **2014,** *136* (2), 725-732.
12.   Venable, R. M.; Brown, F. L. H.; Pastor, R. W., Mechanical properties of lipid bilayers from molecular dynamics simulation. *Chem Phys Lipids* **2015,** *192*, 60-74.
13.   Enkavi, G.; Javanainen, M.; Kulig, W.; Rog, T.; Vattulainen, I., Multiscale Simulations of Biological Membranes: The Challenge To Understand Biological Phenomena in a Living Substance. *Chem Rev* **2019,** *119* (9), 5607-5774.
14.   Corradi, V.; Sejdiu, B. I.; Mesa-Galloso, H.; Abdizadeh, H.; Noskov, S. Y.; Marrink, S. J.; Tieleman, D. P., Emerging Diversity in Lipid-Protein Interactions. *Chem Rev* **2019,** *119* (9), 5775-5848.
15.   Marrink, S. J.; Corradi, V.; Souza, P. C. T.; Ingólfsson, H. I.; Tieleman, D. P.; Sansom, M. S. P., Computational Modeling of Realistic Cell Membranes. *Chemical Reviews* **2019,** *119* (9), 6184-6226.
16.   Bacle, A.; Gautier, R.; Jackson, C. L.; Fuchs, P. F. J.; Vanni, S., Interdigitation between Triglycerides and Lipids Modulates Surface Properties of Lipid Droplets. *Biophys J* **2017,** *112* (7), 1417-1430.
17.   Campomanes, P.; Zoni, V.; Vanni, S., Local accumulation of diacylglycerol alters membrane properties nonlinearly due to its transbilayer activity. *Communications Chemistry* **2019,** *2* (1).
18.   Zoni, V.; Khaddaj, R.; Campomanes, P.; Thiam, A. R.; Schneiter, R.; Vanni, S., Pre-existing bilayer stresses modulate triglyceride accumulation in the ER versus lipid droplets. *Elife* **2021,** *10*.
19.   Zoni, V.; Khaddaj, R.; Lukmantara, I.; Shinoda, W.; Yang, H.; Schneiter, R.; Vanni, S., Seipin accumulates and traps diacylglycerols and triglycerides in its ring-like structure. *Proc Natl Acad Sci U S A* **2021,** *118* (10), e2017205118.
20.   Coline Prévost; E. Sharp, M.; Nora Kory; Qingqing Lin; A. Voth, G.; V. Farese, R.; C. Walther, T., Mechanism and Determinants of Amphipathic Helix-Containing Protein Targeting to Lipid Droplets. *Developmental Cell* **2018,** *44* (1), 73-86.
21.   Olarte, M. J.; Kim, S.; Sharp, M. E.; Swanson, J. M. J.; Farese, R. V., Jr.; Walther, T. C., Determinants of Endoplasmic Reticulum-to-Lipid Droplet Protein Targeting. *Dev Cell* **2020,** *54* (4), 471-487.
22.   Kim, S.; Swanson, J. M. J., The Surface and Hydration Properties of Lipid Droplets. *Biophys J* **2020,** *119* (10), 1958-1969.
23.   Kim, S.; Oh, M. I.; Swanson, J. M. J., Stressed lipid droplets: How neutral lipids relieve surface tension and membrane expansion drives protein association. *J. Phys. Chem. B* **2021,** *submitted*.

24.   Khandelia, H.; Duelund, L.; Pakkanen, K. I.; Ipsen, J. H., Triglyceride Blisters in Lipid Bilayers: Implications for Lipid Droplet Biogenesis and the Mobile Lipid Signal in Cancer Cell Membranes. *PLoS ONE* **2010,** *5* (9), e12811.
25.   Chaban, V. V.; Khandelia, H., Distribution of Neutral Lipids in the Lipid Droplet Core. *The Journal of Physical Chemistry B* **2014,** *118* (38), 11145-11151.
26.   Chaban, V. V.; Khandelia, H., Lipid Structure in Triolein Lipid Droplets. *The Journal of Physical Chemistry B* **2014,** *118* (35), 10335-10340.
27.   Prasanna, X.; Salo, V. T.; Li, S.; Ven, K.; Vihinen, H.; Jokitalo, E.; Vattulainen, I.; Ikonen, E., Seipin traps triacylglycerols to facilitate their nanoscale clustering in the endoplasmic reticulum membrane. *PLoS Biol* **2021,** *19* (1), e3000998.
28.   Hamilton, J. A.; Small, D. M., Solubilization and localization of triolein in phosphatidylcholine bilayers: a 13C NMR study. *Proc Natl Acad Sci U S A* **1981,** *78* (11), 6878-82.
29.   Cornell, R. B.; Ridgway, N. D., CTP:phosphocholine cytidylyltransferase: Function, regulation, and structure of an amphitropic enzyme required for membrane biogenesis. *Prog Lipid Res* **2015,** *59*, 147-71.
30.   Cornell, R. B., Membrane lipid compositional sensing by the inducible amphipathic helix of CCT. *Biochim Biophys Acta* **2016,** *1861* (8 Pt B), 847-861.
31.   Cornell, R. B., Membrane Lipids Assist Catalysis by CTP: Phosphocholine Cytidylyltransferase. *Journal of Molecular Biology* **2020,** *432* (18), 5023-5042.
32.   Gawrisch, K.; Parsegian, V. A.; Hajduk, D. A.; Tate, M. W.; Graner, S. M.; Fuller, N. L.; Rand, R. P., Energetics of a hexagonal-lamellar-hexagonal-phase transition sequence in dioleoylphosphatidylethanolamine membranes. *Biochemistry* **1992,** *31* (11), 2856-64.
33.   Ding, W.; Palaiokostas, M.; Wang, W.; Orsi, M., Effects of Lipid Composition on Bilayer Membranes Quantified by All-Atom Molecular Dynamics. *The Journal of Physical Chemistry B* **2015,** *119* (49), 15263-15274.
34.   Chen, Z.; Rand, R. P., The influence of cholesterol on phospholipid membrane curvature and bending elasticity. *Biophys J* **1997,** *73* (1), 267-76.
35.   Kozlov, M. M.; Leikin, S.; Rand, R. P., Bending, hydration and interstitial energies quantitatively account for the hexagonal-lamellar-hexagonal reentrant phase transition in dioleoylphosphatidylethanolamine. *Biophys J* **1994,** *67* (4), 1603-11.
36.   Rand, R. P.; Fuller, N. L., Structural dimensions and their changes in a reentrant hexagonal-lamellar transition of phospholipids. *Biophysical Journal* **1994,** *66* (6), 2127-2138.
37.   Alexander J. Sodt; Richard W. Pastor, Bending Free Energy from Simulation: Correspondence of Planar and Inverse Hexagonal Lipid Phases. *Biophysical Journal* **2013,** *104* (10), 2202-2211.
38.   Jo, S.; Kim, T.; Iyer, V. G.; Im, W., CHARMM-GUI: A web-based graphical user interface for CHARMM. *Journal of Computational Chemistry* **2008,** *29* (11), 1859-1865.
39.   Jo, S.; Lim, J. B.; Klauda, J. B.; Im, W., CHARMM-GUI Membrane Builder for Mixed Bilayers and Its Application to Yeast Membranes. *Biophysical Journal* **2009,** *97* (1), 50-58.
40.   Wu, E. L.; Cheng, X.; Jo, S.; Rui, H.; Song, K. C.; Dávila-Contreras, E. M.; Qi, Y.; Lee, J.; Monje-Galvan, V.; Venable, R. M.; Klauda, J. B.; Im, W., CHARMM-GUI Membrane Builder toward realistic biological membrane simulations. *Journal of Computational Chemistry* **2014,** *35* (27), 1997-2004.
41.   Lee, J.; Cheng, X.; Swails, J. M.; Yeom, M. S.; Eastman, P. K.; Lemkul, J. A.; Wei, S.; Buckner, J.; Jeong, J. C.; Qi, Y.; Jo, S.; Pande, V. S.; Case, D. A.; Brooks, C. L., 3rd; MacKerell, A. D., Jr.; Klauda, J. B.; Im, W., CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J Chem Theory Comput* **2016,** *12* (1), 405-13.
42.   Martínez, L.; Andrade, R.; Birgin, E. G.; Martínez, J. M., PACKMOL: A package for building initial configurations for molecular dynamics simulations. *Journal of Computational Chemistry* **2009,** *30* (13), 2157-2164.
43.   Klauda, J. B.; Venable, R. M.; Freites, J. A.; O'Connor, J. W.; Tobias, D. J.; Mondragon-Ramirez, C.; Vorobyov, I.; Mackerell, A. D.; Pastor, R. W., Update of the CHARMM All-Atom Additive Force Field for Lipids: Validation on Six Lipid Types. *The Journal of Physical Chemistry B* **2010,** *114* (23), 7830-7843.
44.   Leonard, A. N.; Simmonett, A. C.; Pickard, F. C. t.; Huang, J.; Venable, R. M.; Klauda, J. B.; Brooks, B. R.; Pastor, R. W., Comparison of Additive and Polarizable Models with Explicit Treatment of Long-Range Lennard-Jones Interactions Using Alkane Simulations. *J Chem Theory Comput* **2018,** *14* (2), 948-958.
45.   Yu, Y.; Kramer, A.; Venable, R. M.; Brooks, B. R.; Klauda, J. B.; Pastor, R. W., CHARMM36 Lipid Force Field with Explicit Treatment of Long-Range Dispersion: Parametrization and Validation for Phosphatidylethanolamine, Phosphatidylglycerol, and Ether Lipids. *J Chem Theory Comput* **2021,** *17* (3), 1581-1595.
46.   Yu, Y.; Kramer, A.; Venable, R. M.; Simmonett, A. C.; MacKerell, A. D., Jr.; Klauda, J. B.; Pastor, R. W.; Brooks, B. R., Semi-automated Optimization of the CHARMM36 Lipid Force Field to Include Explicit Treatment of Long-Range Dispersion. *J Chem Theory Comput* **2021,** *17* (3), 1562-1580.

47.   Van Der Spoel, D.; Lindahl, E.; Hess, B.; Groenhof, G.; Mark, A. E.; Berendsen, H. J. C., GROMACS: Fast, flexible, and free. *Journal of Computational Chemistry* **2005,** *26* (16), 1701-1718.

48.   Tribello, G. A.; Bonomi, M.; Branduardi, D.; Camilloni, C.; Bussi, G., PLUMED 2: New feathers for an old bird. *Computer Physics Communications* **2014,** *185* (2), 604-613.

49.   Essmann, U.; Perera, L.; Berkowitz, M. L.; Darden, T.; Lee, H.; Pedersen, L. G., A smooth particle mesh Ewald method. *The Journal of Chemical Physics* **1995,** *103* (19), 8577-8593.

50.   Hess, B., P-LINCS: A Parallel Linear Constraint Solver for Molecular Simulation. *Journal of Chemical Theory and Computation* **2008,** *4* (1), 116-122.

51.   Parrinello, M.; Rahman, A., Polymorphic transitions in single crystals: A new molecular dynamics method. *Journal of Applied Physics* **1981,** *52* (12), 7182-7190.

52.   Nosé, S., A unified formulation of the constant temperature molecular dynamics methods. *The Journal of Chemical Physics* **1984,** *81* (1), 511-519.

53.   Hoover, W. G., Canonical dynamics: Equilibrium phase-space distributions. *Physical Review A* **1985,** *31* (3), 1695-1697.

54.   Berendsen, H. J. C.; Postma, J. P. M.; Van Gunsteren, W. F.; Dinola, A.; Haak, J. R., Molecular dynamics with coupling to an external bath. *The Journal of Chemical Physics* **1984,** *81* (8), 3684-3690.

55.   Feller, S. E.; Pastor, R. W., Constant surface tension simulations of lipid bilayers: The sensitivity of surface areas and compressibilities. *The Journal of Chemical Physics* **1999,** *111* (3), 1281-1287.

56.   Seelig, J.; Niederberger, W., Deuterium-labeled lipids as structural probes in liquid crystalline bilayers. Deuterium magnetic resonance study. *Journal of the American Chemical Society* **1974,** *96* (7), 2069-2072.

57.   Marčelja, S., Chain ordering in liquid crystals. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1974,** *367* (2), 165-176.

58.   Schindler, H.; Seelig, J., Deuterium order parameters in relation to thermodynamic properties of a phospholipid bilayer. Statistical mechanical interpretation. *Biochemistry* **1975,** *14* (11), 2283-2287.

59.   Sugita, Y.; Okamoto, Y., Replica-exchange molecular dynamics method for protein folding. *Chemical Physics Letters* **1999,** *314* (1-2), 141-151.

60.   Torrie, G. M.; Valleau, J. P., Nonphysical sampling distributions in Monte Carlo free-energy estimation: Umbrella sampling. *Journal of Computational Physics* **1977,** *23* (2), 187-199.

61.   Kumar, S.; Rosenberg, J. M.; Bouzida, D.; Swendsen, R. H.; Kollman, P. A., THE weighted histogram analysis method for free-energy calculations on biomolecules. I. The method. *Journal of Computational Chemistry* **1992,** *13* (8), 1011-1021.

62.   Grossfied, A., version 2.0.9. WHAM: an implementation of the weighted histogram analysis method.

63.   Dama, J. F.; Rotskoff, G.; Parrinello, M.; Voth, G. A., Transition-Tempered Metadynamics: Robust, Convergent Metadynamics via On-the-Fly Transition Barrier Estimation. *Journal of Chemical Theory and Computation* **2014,** *10* (9), 3626-3633.

64.   Sun, R.; Dama, J. F.; Tan, J. S.; Rose, J. P.; Voth, G. A., Transition-Tempered Metadynamics Is a Promising Tool for Studying the Permeation of Drug-like Molecules through Membranes. *Journal of Chemical Theory and Computation* **2016,** *12* (10), 5157-5169.

65.   Bussi, G.; Donadio, D.; Parrinello, M., Canonical sampling through velocity rescaling. *The Journal of Chemical Physics* **2007,** *126* (1), 014101.

66.   Madsen, J. J.; Grime, J. M. A.; Rossman, J. S.; Voth, G. A., Entropic forces drive clustering and spatial localization of influenza A M2 during viral budding. *Proc Natl Acad Sci U S A* **2018,** *115* (37), E8595-E8603.

67.   Michaud-Agrawal, N.; Denning, E. J.; Woolf, T. B.; Beckstein, O., MDAnalysis: A toolkit for the analysis of molecular dynamics simulations. *Journal of Computational Chemistry* **2011,** *32* (10), 2319-2327.

68.   Gouw, T. H.; Vlugter, J. C., Physical Properties of Triglycerides. I. Density and Refractive Index. *Fette, Seifen, Anstrichmittel* **1966,** *68* (7), 544-549.

69.   Mitsche, M. A.; Wang, L.; Small, D. M., Adsorption of egg phosphatidylcholine to an air/water and triolein/water bubble interface: use of the 2-dimensional phase rule to estimate the surface composition of a phospholipid/triolein/water surface as a function of surface pressure. *J Phys Chem B* **2010,** *114* (9), 3276-84.

70.   Kramer, A.; Pickard, F. C. t.; Huang, J.; Venable, R. M.; Simmonett, A. C.; Reith, D.; Kirschner, K. N.; Pastor, R. W.; Brooks, B. R., Interactions of Water and Alkanes: Modifying Additive Force Fields to Account for Polarization Effects. *J Chem Theory Comput* **2019,** *15* (6), 3854-3867.

71.   Venable, R. M.; Kramer, A.; Pastor, R. W., Molecular Dynamics Simulations of Membrane Permeability. *Chem Rev* **2019,** *119* (9), 5954-5997.

72.   Cerretani, L.; Giuliani, A.; Maggio, R. M.; Bendini, A.; Gallina Toschi, T.; Cichelli, A., Rapid FTIR determination of water, phenolics and antioxidant activity of olive oil. *European Journal of Lipid Science and Technology* **2010,** *112* (10), 1150-1157.

73.   Hatzakis, E.; Dais, P., Determination of water content in olive oil by 31P NMR spectroscopy. *J Agric Food Chem* **2008,** *56* (6), 1866-72.

74.   Ragni, L.; Berardinelli, A.; Cevoli, C.; Valli, E., Assessment of the water content in extra virgin olive oils by Time Domain Reflectometry (TDR) and Partial Least Squares (PLS) regression methods. *Journal of Food Engineering* **2012,** *111* (1), 66-72.

75.   Arnold, R. S.; Cornell, R. B., Lipid regulation of CTP: phosphocholine cytidylyltransferase: electrostatic, hydrophobic, and synergistic interactions of anionic phospholipids and diacylglycerol. *Biochemistry* **1996,** *35* (30), 9917-24.

76.   Cornell, R. B.; Arnold, R. S., Modulation of the activities of enzymes of membrane lipid metabolism by non-bilayer-forming lipids. *Chemistry and Physics of Lipids* **1996,** *81* (2), 215-227.

77.   Davies, S. M.; Epand, R. M.; Kraayenhof, R.; Cornell, R. B., Regulation of CTP: phosphocholine cytidylyltransferase activity by the physical properties of lipid membranes: an important role for stored curvature strain energy. *Biochemistry* **2001,** *40* (35), 10522-31.

78.   Vanni, S.; Vamparys, L.; Gautier, R.; Drin, G.; Etchebest, C.; Fuchs, F. J., Patrick; Antonny, B., Amphipathic Lipid Packing Sensor Motifs: Probing Bilayer Defects with Hydrophobic Residues. *Biophysical Journal* **2013,** *104* (3), 575-584.

79.   Harayama, T.; Riezman, H., Understanding the diversity of membrane lipid composition. *Nature Reviews Molecular Cell Biology* **2018,** *19* (5), 281-296.

80.   Gordon, S. M.; Pourmousa, M.; Sampson, M.; Sviridov, D.; Islam, R.; Perrin, B. S., Jr.; Kemeh, G.; Pastor, R. W.; Remaley, A. T., Identification of a novel lipid binding motif in apolipoprotein B by the analysis of hydrophobic cluster domains. *Biochim Biophys Acta Biomembr* **2017,** *1859* (2), 135-145.

81.   Brannigan, G.; Brown, F. L., Solvent-free simulations of fluid membrane bilayers. *J Chem Phys* **2004,** *120* (2), 1059-71.

82.   Brannigan, G.; Philips, P. F.; Brown, F. L., Flexible lipid bilayers in implicit solvent. *Phys Rev E Stat Nonlin Soft Matter Phys* **2005,** *72* (1 Pt 1), 011915.

83.   Brandt, E. G.; Braun, A. R.; Sachs, J. N.; Nagle, J. F.; Edholm, O., Interpretation of fluctuation spectra in lipid bilayer simulations. *Biophys J* **2011,** *100* (9), 2104-11.

84.   Kučerka, N.; Nagle, J. F.; Sachs, J. N.; Feller, S. E.; Pencer, J.; Jackson, A.; Katsaras, J., Lipid Bilayer Structure Determined by the Simultaneous Analysis of Neutron and X-Ray Scattering Data. *Biophysical Journal* **2008,** *95* (5), 2356-2367.

85.   Pan, J.; Tristram-Nagle, S.; Kučerka, N.; Nagle, J. F., Temperature Dependence of Structure, Bending Rigidity, and Bilayer Interactions of Dioleoylphosphatidylcholine Bilayers. *Biophysical Journal* **2008,** *94* (1), 117-124.

86.   Evans, E.; Rawicz, W.; Smith, B. A., Concluding remarks Back to the future: mechanics and thermodynamics of lipid biomembranes. *Faraday Discuss.* **2013,** *161*, 591-611.

87.   Rawicz, W.; Olbrich, K. C.; Mcintosh, T.; Needham, D.; Evans, E., Effect of Chain Length and Unsaturation on Elasticity of Lipid Bilayers. *Biophysical Journal* **2000,** *79* (1), 328-339.

88.   Jablin, M. S.; Akabori, K.; Nagle, J. F., Experimental Support for Tilt-Dependent Theory of Biomembrane Mechanics. *Physical Review Letters* **2014,** *113* (24).

89.   Gracià, R. S.; Bezlyepkina, N.; Knorr, R. L.; Lipowsky, R.; Dimova, R., Effect of cholesterol on the rigidity of saturated and unsaturated membranes: fluctuation and electrodeformation analysis of giant vesicles. *Soft Matter* **2010,** *6* (7), 1472.

90.   Chorlay, A.; Thiam, A. R., An Asymmetry in Monolayer Tension Regulates Lipid Droplet Budding Direction. *Biophysical Journal* **2018,** *114* (3), 631-640.

91.   Chorlay, A.; Monticelli, L.; Veríssimo Ferreira, J.; Ben M'Barek, K.; Ajjaji, D.; Wang, S.; Johnson, E.; Beck, R.; Omrane, M.; Beller, M.; Carvalho, P.; Rachid Thiam, A., Membrane Asymmetry Imposes Directionality on Lipid Droplet Emergence from the ER. *Developmental Cell* **2019,** *50* (1), 25-42.

92.   Sui, X.; Wang, K.; Gluchowski, N. L.; Elliott, S. D.; Liao, M.; Walther, T. C.; Farese, R. V., Jr., Structure and catalytic mechanism of a human triacylglycerol-synthesis enzyme. *Nature* **2020,** *581* (7808), 323-328.

93.   Lie Wang; Hongwu Qian; Yin Nian; Yimo Han; Zhenning Ren; Hanzhi Zhang; Liya Hu; Venkataram Prasad, B. V.; Arthur Laganowsky; Nieng Yan; Ming Zhou, Structure and mechanism of human diacylglycerol O-acyltransferase 1. *Nature* **2020,** *581* (7808), 329-332.

94.   Santinho, A.; Salo, V. T.; Chorlay, A.; Li, S.; Zhou, X.; Omrane, M.; Ikonen, E.; Thiam, A. R., Membrane Curvature Catalyzes Lipid Droplet Assembly. *Curr Biol* **2020,** *30* (13), 2481-2494 e6.

95.   Pinot, M.; Vanni, S.; Pagnotta, S.; Lacas-Gervais, S.; Payet, L.-A.; Ferreira, T.; Gautier, R.; Goud, B.; Antonny, B.; Barelli, H., Polyunsaturated phospholipids facilitate membrane deformation and fission by endocytic proteins. *Science* **2014,** *345* (6197), 693-697.

96.   Rong, X.; Wang, B.; Dunham, M. M.; Hedde, P. N.; Wong, J. S.; Gratton, E.; Young, S. G.; Ford, D. A.; Tontonoz, P., Lpcat3-dependent production of arachidonoyl phospholipids is a key determinant of triglyceride secretion. *Elife* **2015,** *4*.

97.   Hashidate-Yoshida, T.; Harayama, T.; Hishikawa, D.; Morimoto, R.; Hamano, F.; Tokuoka, S. M.; Eto, M.; Tamura-Nakano, M.; Yanobu-Takanashi, R.; Mukumoto, Y.; Kiyonari, H.; Okamura, T.; Kita, Y.; Shindou, H.; Shimizu, T., Fatty acid remodeling by LPCAT3 enriches arachidonate in phospholipid membranes and regulates triglyceride transport. *eLife* **2015,** *4*.

98.   Barelli, H.; Antonny, B., Lipid unsaturation and organelle dynamics. *Current Opinion in Cell Biology* **2016,** *41*, 25-32.

99.   Augenstreich, J.; Haanappel, E.; Ferre, G.; Czaplicki, G.; Jolibois, F.; Destainville, N.; Guilhot, C.; Milon, A.; Astarie-Dequeker, C.; Chavent, M., The conical shape of DIM lipids promotes Mycobacterium tuberculosis infection of macrophages. *Proc Natl Acad Sci U S A* **2019,** *116* (51), 25649-25658.

100.   Izvekov, S.; Voth, G. A., A Multiscale Coarse-Graining Method for Biomolecular Systems. *The Journal of Physical Chemistry B* **2005,** *109* (7), 2469-2473.

101.   Izvekov, S.; Voth, G. A., Multiscale Coarse-Graining of Mixed Phospholipid/Cholesterol Bilayers. *Journal of Chemical Theory and Computation* **2006,** *2* (3), 637-648.

102.   Noid, W. G.; Chu, J.-W.; Ayton, G. S.; Krishna, V.; Izvekov, S.; Voth, G. A.; Das, A.; Andersen, H. C., The multiscale coarse-graining method. I. A rigorous bridge between atomistic and coarse-grained models. *The Journal of Chemical Physics* **2008,** *128* (24), 244114.

103.   Noid, W. G.; Liu, P.; Wang, Y.; Chu, J.-W.; Ayton, G. S.; Izvekov, S.; Andersen, H. C.; Voth, G. A., The multiscale coarse-graining method. II. Numerical implementation for coarse-grained molecular models. *The Journal of Chemical Physics* **2008,** *128* (24), 244115.

104.   Izvekov, S.; Voth, G. A., Solvent-Free Lipid Bilayer Model Using Multiscale Coarse-Graining. *The Journal of Physical Chemistry B* **2009,** *113* (13), 4443-4455.

105.   Shell, M. S., The relative entropy is fundamental to multiscale and inverse thermodynamic problems. *J Chem Phys* **2008,** *129* (14), 144108.

106.   Pak, A. J.; Dannenhoffer-Lafage, T.; Madsen, J. J.; Voth, G. A., Systematic Coarse-Grained Lipid Force Fields with Semiexplicit Solvation via Virtual Sites. *J Chem Theory Comput* **2019,** *15* (3), 2087-2100.

107.   Simunovic, M.; Mim, C.; Marlovits, T. C.; Resch, G.; Unger, V. M.; Voth, G. A., Protein-mediated transformation of lipid vesicles into tubular networks. *Biophys J* **2013,** *105* (3), 711-9.

108.   Simunovic, M.; Srivastava, A.; Voth, G. A., Linear aggregation of proteins on the membrane as a prelude to membrane remodeling. *Proc Natl Acad Sci U S A* **2013,** *110* (51), 20396-401.

109.   Srivastava, A.; Voth, G. A., A Hybrid Approach for Highly Coarse-grained Lipid Bilayer Models. *J Chem Theory Comput* **2013,** *9* (1), 750-765.

110.   Simunovic, M.; Evergren, E.; Golushko, I.; Prevost, C.; Renard, H. F.; Johannes, L.; McMahon, H. T.; Lorman, V.; Voth, G. A.; Bassereau, P., How curvature-generating proteins build scaffolds on membrane nanotubes. *Proc Natl Acad Sci U S A* **2016,** *113* (40), 11226-11231.

111.   Jarin, Z.; Tsai, F. C.; Davtyan, A.; Pak, A. J.; Bassereau, P.; Voth, G. A., Unusual Organization of I-BAR Proteins on Tubular and Vesicular Membranes. *Biophys J* **2019,** *117* (3), 553-562.

112.   Dama, J. F.; Sinitskiy, A. V.; Mccullagh, M.; Weare, J.; Roux, B.; Dinner, A. R.; Voth, G. A., The Theory of Ultra-Coarse-Graining. 1. General Principles. *Journal of Chemical Theory and Computation* **2013,** *9* (5), 2466-2480.

113.   Davtyan, A.; Dama, J. F.; Sinitskiy, A. V.; Voth, G. A., The Theory of Ultra-Coarse-Graining. 2. Numerical Implementation. *Journal of Chemical Theory and Computation* **2014,** *10* (12), 5265-5275.

114.   Dama, J. F.; Jin, J.; Voth, G. A., The Theory of Ultra-Coarse-Graining. 3. Coarse-Grained Sites with Rapid Local Equilibrium of Internal States. *J Chem Theory Comput* **2017,** *13* (3), 1010-1022.

115.   Jin, J.; Voth, G. A., Ultra-Coarse-Grained Models Allow for an Accurate and Transferable Treatment of Interfacial Systems. *J Chem Theory Comput* **2018,** *14* (4), 2180-2197.

116.   Jin, J.; Han, Y.; Voth, G. A., Ultra-Coarse-Grained Liquid State Models with Implicit Hydrogen Bonding. *J Chem Theory Comput* **2018,** *14* (12), 6159-6174.

117.   Jin, J.; Yu, A.; Voth, G. A., Temperature and Phase Transferable Bottom-up Coarse-Grained Models. *J Chem Theory Comput* **2020,** *16* (11), 6823-6842.

TOC IMAGE