Running title: Functional and structural characterization of *Serratia marcescens* ExbB:

determinants of the interaction with HasB/TonB

Valérie Biou[1], Ricardo Jorge Diogo Adaixo[2], Mohamed Chami[2], Pierre-Damien Coureux[3],

Benoist Laurent[4], Yvette Ntsogo[1], Gisele Cardoso de Amorim[5], Nadia Izadi-Pruneyre[5], Christian

Malosse[6], Julia Chamot-Rooke[6], Henning Stahlberg[2], Philippe Delepelaire[1]

1: Laboratoire de Biologie Physico-Chimique des Protéines Membranaires, Université de
Paris, UMR 7099 CNRS, F-75005, Paris, France.
Institut de Biologie Physico-Chimique, F-75005, Paris, France.
2: Center for Cellular Imaging and NanoAnalytics, Biozentrum, University of Basel,
Mattenstrasse 26, CH-4058, Basel, Switzerland
3: Laboratoire de Biologie Structurale de la Cellule, BIOC, UMR7654 CNRS/Ecole
polytechnique, Palaiseau France
4: Plateforme de Bioinformatique, Université de Paris, FRC 550 CNRS, F-75005, Paris, France.
Institut de Biologie Physico-Chimique, F-75005, Paris, France.
5: Structural Bioinformatics Unit, Department of Structural Biology and Chemistry, C3BI,
Institut Pasteur, CNRS UMR3528, CNRS USR3756, Paris, France.
6: Mass Spectrometry for Biology Unit, CNRS USR 2000, Institut Pasteur, 75015 Paris, France
18
19 For correspondence: valerie.biou@ibpc.fr, philippe.delepelaire@ibpc.fr
20

Summary.

ExbBD is part of a cytoplasmic membrane molecular motor driven by the proton-motive force.

It belongs to the larger family of motors involved in nutriment import across the outer

membrane of Gram-negative bacteria (ExbBD), flagellar rotation (MotAB) or late steps of cell

division in Gram-negative bacteria (TolQR). ExbB and ExbD are integral membrane proteins

with three (ExbB) or one (ExbD) transmembrane segment. Here we have solved by single-

particle cryo-EM the structures of ExbB alone and of the ExbB-ExbD complex of the

opportunistic pathogen *Serratia marcescens*. ExbB$_{Sm}$ alone behaves as a stable pentamer, and

the complex displays the ExbB$_5$-ExbD$_2$ stoichiometry. This is similar to what has been observed

for ExbB-ExbD complexes from *Escherichia coli* and *Pseudomonas savastanoi* as well as MotAB

complexes from various species. We identified residues located in the first TM of ExbB$_{Sm}$ and

ExbB$_{Ec}$ that are likely involved in the interaction with TonB/HasB and that are essential for

function. ExbB$_{Sm}$ has a ca. 40 residues long periplasmic extension absent in *E. coli*. Such long

ExbB's are found in some Gammaproteobacteria, and several genera of Alphaproteobacteria.

We show that this extension interacts with HasB, a dedicated TonB paralog from the heme

acquisition system (Has) from *S. marcescens*. We also show that it is involved in heme

acquisition *via* the Has system from *S. marcescens*. ExbB$_{Sm}$ represents thus a new class of ExbB

protein and our results shed light on the specificity determinants between the ExbB-ExbD

complex and their associated TonB partners.

Introduction

Transport of nutrients across the Gram-negative outer membrane is either a diffusion-

facilitated or an active process, powered in that last case by the energy of the proton-motive

force (pmf) transmitted to specialized outer membrane transporters (the so-called TBDT's,

TonB dependent transporters). A complex of three cytoplasmic membrane proteins, TonB,

ExbB and ExbD, that together form the TonB complex (see [1] for a review) conveys the energy

of the pmf to the TBDT. In *Escherichia coli* K12, there is only one set of *tonB, exbB* and *exbD*

genes, whereas there are nine TBDT's [2], all energized by the same complex. ExbB and ExbD

respectively belong to the MotA-TolQ-ExbB and MotB-TolR-ExbD protein families, that are

involved in power generation and transmission in various processes (flagellar rotation for

MotAB, active transport of molecules across OM receptors for ExbBD and cell division for

TolQR), at the expense of pmf dissipation across the cytoplasmic membrane [3-6]. Those

proteins form complexes that associate with TonB (ExbBD), TolA (TolQR) or the flagellar rotor

(MotAB, stator of the flagellar motor). It has also been shown that ExbD does not accumulate

in the absence of ExbB, and that TonB does not accumulate either in the absence of the ExbBD

complex [7]. It is also known that the C-terminal domains of TonB and ExbD interact with each

other [8, 9]. ExbD TM has one essential strictly conserved aspartate residue [10], that is

thought to undergo cycles of protonation/deprotonation (coupled to pmf dissipation).

TBDT's comprise a 22-stranded ß-barrel anchored in the outer membrane. An N-terminal

domain folded inside the barrel and called the plug contains the main region of interaction

with TonB, the so-called TonB box located in the N-terminal periplasm-exposed part of TBDT.

TonB has its N-terminus localized in the cytoplasm, a single transmembrane segment,

followed by a Pro-Lys rich region long enough to span the periplasm and a structured C-

terminal domain interacting with the TBDT. Conformational changes elicited upon interaction

between TonB and the TonB box of the TBDT, and likely mediated via a rearrangement of the

plug domain, ultimately lead to entry of the substrate (initially bound to the extracellular side

of the receptor) into the periplasm by a yet unknown mechanism.

In *Serratia marcescens*, a close relative of *E. coli*, in which were identified about twenty potential TBDT's, there are at least two TonB homologs: an ortholog of $TonB_{Ec}$ (having 48% identity with $TonB_{Ec}$[11]) and HasB, a TonB paralog that is strictly dedicated to its cognate outer membrane receptor HasR[12, 13]. The Has (heme acquisition system) system includes the TBDT HasR, which recognizes both free heme and the high-affinity extracellular heme-binding HasA hemophore. HasB is encoded within the *has* locus and displays low sequence identity with either *E. coli* or *S. marcescens* TonB. Unlike $TonB_{Sm}$, HasB cannot complement $TonB_{Ec}$ functions, nor it is able to drive heme entry *via* the HasR receptor in the presence of $ExbBD_{Ec}$. A gain of function mutation in the TM domain of HasB was however isolated in *E. coli*, restoring heme entry *via* HasR in the presence of $ExbBD_{Ec}$ [12]. HasB has the same topology as TonB, and its C-terminal domain interacts specifically with HasR with a ca. 40-fold higher affinity than the corresponding TonB domain [14]. Those observations prompted us to identify the ExbBD complex that would function with HasB in *S. marcescens*. Hence, we first identified the resident orthologous $ExbBD_{Sm}$ and found that this complex is active with both HasB and TonB.

Recently several structures of *E. coli* ExbB and ExbBD have been solved, either by X-ray crystallography or cryo-EM[15-17]. All samples studied included ExbB and ExbD, but not all showed an ordered structure of ExbD. In these structures, $ExbB_{Ec}$[15] appeared as a pentamer, each monomer of ExbB having three TM helices that extend into the cytoplasm, and with a highly polarized charge distribution on the cytoplasmic side. The central pore is apolar, lined by TM helices 2 and 3 of each monomer, creating a large hydrophobic cavity inside the structure. Another study using X-ray crystallography, single-particle cryo-EM and electron diffraction on two-dimensional crystals concluded that ExbB could undergo a pH-dependent pentamer to hexamer transition, and that the hexameric ExbB could accommodate three ExbD TM segments in its pore[16]. A more recent cryo-EM study of ExbBD reconstituted in nanodiscs however confirmed the $ExbB_5ExbD_2$ stoichiometry consistent with previous DEER results and identified the two TM helices of ExbD in the ExbB pore [16]. Several structures of the flagellar MotAB motor complexes (related to ExbBD) from diverse bacteria (*Campylobacter jejuni, Bacillus subtilis, Clostridium sporogenes, Vibrio mimicus, Shewanella oneidensis,* PomAB from *Vibrio alginolyticus*) were recently published that also display a $MotA_5MotB_2$ stoichiometry with some common topology elements [18, 19]. Similarly, *Pseudomonas savastanoi* ExbBD exhibits the same stoichiometry.

Here we identify a new family of ExbB proteins with a long N-terminal extension whose

presence is strongly correlated to the presence of a *hasB* gene ortholog in the genome. We

characterized ExbB$_{Sm}$ alone and ExbBD$_{Sm}$ by single-particle cryo-EM and determined their

structures at 3.2 and 3.9Å, respectively. We show that in both cases, ExbB$_{Sm}$ behaves as a

stable pentamer; in the ExbBD$_{Sm}$ complex, we identify two TM helices of ExbD$_{Sm}$ in the pore

of ExbB pentamer, as it was shown for the *E. coli* case. Using NMR measurements, we also

show that the N-terminal periplasmic extension specific to ExbB$_{Sm}$ interacts with the globular

domain of HasB. Finally, *via* mutagenesis studies and bacterial growth assays we show that

the first transmembrane helix TM1 of ExbB contains specificity determinants for interaction

with HasB/TonB and that the periplasmic extension does play a role in heme acquisition *via*

the <u>has</u> system.

Results

1. Orthologs of *E. coli exbB* and *exbD* in *Serratia marcescens* define a new ExbB family with

a N-terminal extension.

Sequence analysis of strain Db11, a fully sequenced *S. marcescens* isolate (GenBank: *Serratia*

marcescens subsp. marcescens Db11, complete genome. ACCESSION HG326223,

https://www.ncbi.nlm.nih.gov/nuccore/HG326223.1), identified one putative operon

encoding orthologs of *E. coli* ExbB and ExbD (SMDB11_3479(ExbD) and SMDB11_3480(ExbB)).

In this case the *exbB*-like gene is close to and in the opposite orientation to a *metC* homolog,

as in *E. coli.* There is no conservation on the 3' end. Comparison of the coding sequences

indicated that the *exbD* gene encoded a 140 residue-long protein (71% identity with the 141

residue-long *E. coli* sequence). In contrast, the putative *exbB* gene encodes a much longer

protein than its *E. coli* counterpart (325 residues instead of 244 residues in *E. coli*, 73% identity

in the common part). The additional stretch of residues present in ExbB$_{Sm}$ is located at the N-

terminus and corresponds to a putative signal peptide followed by a ~50 residues N-terminal

extension of the mature protein (Supplementary Figure S1).

A BLAST search on bacterial genomes indicated that such "extralong" ExbB's are found in

different Gammaproteobacteria, including Serratia, Yersinia, Pseudomonas, Erwinia/Dickeya,

and many genera in the Alphaproteobacteria class. Table SI lists representative species of

Alphaproteobacteria in which such long ExbB's are found. About 90% of the long ExbB-

containing species listed in this table also have a *hasB* gene ortholog.

An aminoacid sequence alignment of these ExbBs (Clustal omega

https://www.ebi.ac.uk/Tools/msa/clustalo/, Figure 1) also shows that the mature N-terminal

extensions are of variable length between a few and 150 residues with an average of 50

residues. These extensions are quite rich in Ala (24,6%) and Pro (14,5%) residues and therefore

likely to be unstructured but could be involved in protein-protein interactions as proline-rich

regions often are in signalling processes [20-22].

The Conserved Domain Database [23] contains currently two ExbB subfamilies in the cl00568

MotA_ExbB superfamily, TIGR02797 (containing *E. coli* ExbB, as well as longer ExbB's) and

TIGR02805 (containing *Haemophilus influenzae* ExbB, with a very short cytoplasmic domain

between TM1 and TM2); based on sequence data, it is then possible to subdivide TIGR02797

into 2 groups with "extralong" ExbB's in one subfamily, and *E. coli*-like sequences in the other.

2. ExbBD$_{Sm}$ complements ExbBD$_{Ec}$ for iron acquisition and functionally associates with both

TonB and HasB for heme acquisition through HasR

As a first step in the characterization of the identified ExbBD$_{sm}$, we tested whether this

complex would complement ExbBD$_{Ec}$. To this end, the plasmids pBADExbBD$_{Sm}$ and

pBADExbBD$_{Ec}$ were constructed, introducing the two genes, from *S. marcescens* and *E. coli*,

into pBAD24 vector under the control of p$_{araBAD}$ promoter. By using growth under iron

starvation conditions as a test, we could show that *exbBD$_{Sm}$* complements *exbBD$_{Ec}$*. The *E. coli*

C600*ΔexbBD* strain is indeed more sensitive than its wild-type counterpart to iron starvation

(induced for example by the Fe^{2+} chelator dipyridyl) as ExbBD are required to transduce pmf

to TonB for siderophore uptake. Its growth in the presence of dipyridyl is restored by the

expression of either pBADExbBD$_{Ec}$ or pBADExbBD$_{Sm}$ (Supplementary Figure S2).

HasB is a HasR-specific TonB ortholog, and as ExbBD$_{Ec}$ does not function with HasB (for heme

acquisition through HasA-HasR [12]), we then verified that ExbBD$_{Sm}$ would do so. To this end,

pBADExbBD$_{Sm}$ was introduced into strain C600*ΔhemAΔtonBΔexbBD* harbouring the

phasISRADEB/238 recombinant plasmid containing the whole <u>has</u> locus with its own

regulatory elements. This strain could grow in the presence of heme (down to a concentration

of 0.2μM) and arabinose, whereas the C600*ΔhemAΔtonBΔexbBD*(phasISRADEB/238 +

pBADExbBD$_{Ec}$) could not (see Supplementary Figure S3 bottom part). The strain

C600*ΔhemAΔexbBD*(phasISRADE/238 + pBADExbBD$_{Ec}$) could grow in the presence of heme

and arabinose, indicating that pBADExbBD$_{Ec}$ is functional (see Supplementary Figure S3 top

part). Likewise C600Δ*hemA*Δ*exbBD*(phasISRADE/238 + pBADExbBD$_{Sm}$) could grow in the

presence of heme and arabinose. This indicates that ExbB$_{Sm}$ and ExbD$_{Sm}$ functionally associate

with either HasB or TonB$_{Ec}$ to allow HasA-mediated heme entry *via* HasR, whereas ExbBD$_{Ec}$

only functions with TonB leading to the heme entry *via* the HasA-HasR system (see

Supplementary Figure S3 top part). As already observed, a stronger iron starvation is

necessary when TonB is the link between the inner and outer membrane components [24].

Therefore, ExbBD$_{Sm}$ is the genuine *E. coli* ortholog, with a looser specificity.

3. Characterization of ExbB$_{Sm}$ and ExbBD$_{Sm}$ and their structural analysis by cryo-EM show

ExbB$_5$ and ExbB$_5$D$_2$ stoichiometries

To gain further insight into the possible differences between ExbBD$_{Sm}$ and ExbBD$_{Ec}$, we purified

both ExbB$_{Sm}$ and ExbBD$_{Sm}$ (as His-tagged proteins, see Materials and Methods section) in

LMNG micelles (Supplementary Figure S4). Both ExbB$_{Sm}$ and ExbBD$_{Sm}$ appear as homogeneous

oligomers at the last size exclusion chromatography (SEC) step. Mass spectrometry analysis of

intact ExbD$_{Sm}$ (16 131 Da) and ExbB$_{Sm}$ (29 557 Da) proteins, together with the determination

of the first amino-acid residues for ExbB (APAAN) showed that the predicted signal sequence

of ExbB$_{Sm}$ is absent from the mature sequence and has therefore been processed. It also

showed that ExbD initial Met residue was cleaved off (see Supplementary Figure S5). More

interestingly, mass spectrometry analysis of chloroform/methanol extracts of the purified

proteins also showed that both complexes contained native lipids (Supplementary Figure S6),

mostly phosphatidylglycerol (PG) and phosphatidylethanolamine (PE), and that ExbBD also

contained some cardiolipin (CL) (Supplementary Figure S7). Further analysis of aliphatic chain

composition of lipids shows evidence of specific composition (Supplementary Figure S8).

3a ExbB structure

As ExbB$_{Sm}$ presented specific features compared to ExbB$_{Ec}$, we set out to determine its

structure by single-particle cryo-EM. The 3D class average model clearly showed a pentameric

structure. It was refined and polished to obtain a resolution of 3.2Å. Maps were refined with

and without C5 symmetry and showed a 98% correlation. Therefore, the map used for model

building was the one with C5 symmetry.

The structure solved here and shown in Figure 2 A and B, is very similar to that of *E. coli* ExbB

5SV0 structure (that was co-purified with ExbD), with a 1.3Å rmsd over all Cα atoms (Table I).

This shows that ExbB is stable as a pentamer on its own. The periplasmic N-terminal extension

did not yield any visible electron density, precluding its structure determination. At the C-terminus of each monomer, however, electron density was clearly defined, allowing structure determination for an additional 10 residues in a helical conformation up to the last one (Gly 283, see Figure 2D, helix α8), before the tag. In the 5SV0 X-ray structure, a calcium ion is bound to the five Glu 106 (of helix α2 in the TM region) on the cytoplasmic side. In ExbB$_{Sm}$, this residue is replaced by an Asn, and we do not observe any density that could be assigned to a metal ion.

The structure of the whole pentamer with one monomer coloured as a function of sequence conservation in this sub-class of ExbB's (Supplementary figure 9) shows that the highest residue conservation is inside the transmembrane channel, indicative of high functional constraints, while the TM residues located at the membrane surface are more variable. The same observation was made for those ExbB's without N-terminal signal sequence and periplasmic extension [15].

Similarly to the ExbB$_{EC}$ structure, at the cytoplasmic junction with the membrane there are "fenestrations", potentially leading to solvent or ion passage [15]. These vents are located at the junction of adjacent monomers as shown in figure 2A. Additional density was also clearly observed on the external surface, which we attributed to the phospholipids present in the preparation. Each ExbB monomer appears to be associated with the equivalent of two PG molecules, located at the inner leaflet of the cytoplasmic membrane (Figure 2A and C). However, we were able to model only one PG molecule per monomer with confidence. Three PE (phosphatidylethanolamine, the major *E. coli* phospholipid) molecules and one PG molecule were identified as associated to the ExbBD pentamer in the *E. coli* complex after reconstitution in nanodiscs [17]. In summary, our cryo-EM structure shows that ExbB from *S. marcescens* is stable as a pentamer and associates strongly with specific phospholipids coming from the inner leaflet of the membrane.

3b ExbBD$_{Sm}$ structure

We also solved the structure of the ExbBD$_{Sm}$ complex by cryo-EM (Figure 3 and Table I). As for ExbB, the purified complex exhibited a symmetric peak on SEC (see Supplementary Figure S4). Cryo-EM data were collected and processed as described in the Material and Methods section. 2D classes showed preferential orientations with 92% top views and only 8% side views. Due to this very strong bias in particle orientation, the resolution achieved was 3.9 Å, precluding

reliable positioning of the side chains. As observed for ExbB alone, in the ExbBD complex, ExbB behaves as a pentamer and no sign of other assemblies was found. Similarly, no density could be attributed to the periplasmic extension. There are two clear densities inside the ExbB pore that were assigned to the pair of ExbD TM segments (Figure 3 B and C). The charge distribution is highly polarised on the cytoplasmic side: the region close to the membrane is positively charged as expected and the distal part is negatively charged (Supplementary Figure S10, top left). The central pore is apolar, lined by TM helices 2 and 3 of each monomer, creating a large hydrophobic cavity inside the structure (Supplementary Figure S10, top right). In the refined model, Asp 25 from ExbD monomer chain G faces Thr 218 from ExbB chain C, while Asp 25 from the ExbD chain F faces the interface between two ExbB monomers A and E (Supplementary Figure S10, bottom). The estimation of pKa by using the program Propka [25] server (http://server.poissonboltzmann.org/) shows that Asp 25 has a pKa of 7.3 for chain F and 7.42 for chain G, both very peculiar for solvent-accessible acidic residues but that can be found in buried active sites or membrane proteins [26]. This pKa should allow protonation and deprotonation of ExbD Asp 25 at physiological pH.

As compared to the structure of $ExbB_{Sm}$, there is a slight «opening» towards the periplasmic side (the distance of Ala 197 from one subunit to Leu 204 of the facing subunit varying from 25.5Å in ExbB to 29.8Å in ExbBD). This opening of the structure is limited to the periplasmic part (Supplementary Figure S11). The RMSD between $ExbB_{Sm}$ alone and in complex with $ExbD_{Sm}$ is 2Å over all Cα of the pentamer, and the RMSD between $ExbB_{Sm}$ in complex with $ExbD_{Sm}$ and $ExbB_{Ec}$ in complex with $ExbD_{Ec}$ (structure 6TYI chains A-E) is 1.7Å.

As compared to the 6TYI structure of *E. coli* ExbBD complex, there are some significant differences that could be related to the specificity of function of $ExbB_{Sm}$. Both $ExbB_{Sm}$ and $ExbBD_{Sm}$ inner pores are slightly wider than their *E. coli* counterparts at the periplasmic entrance. Consequently, two channels cross the membrane region, (extending from the periplasmic entrance to the Asp 25 residue from the ExbD TM deeply embedded in the ExbB pore) that are clearly seen in the $ExbBD_{Sm}$ (as detected by the MoleOnline Server) while the *E. coli* structure exhibits a single, much thinner channel as detected with the same parameters of MoleOnline (Supplementary Figure S12). It is therefore possible that this structure represents another state of the ExbBD motor complex, whereby the periplasmic side allows solvent access to the Asp 25 residue of ExbD TM. In summary, we show that ExbBD from *S.*

marcescens has a 5:2 stoichiometry with a larger channel allowing solvent or proton transport

from the periplasmic side as compared with the *E. coli* structure.

4. Specific functions of the N-terminal extension and of the first TM segment of ExbB_{Sm}

4a NMR shows an interaction between HasB and ExbB_{Sm} periplasmic regions.

The N-terminal extension of ExbB$_{Sm}$ contains almost 60% of Ala and Pro residues and is

predicted to be unstructured. In agreement with this, there was no observable electron

density for this region neither in the free form nor in the complex. Indeed, NMR analysis of a

44 residue-long synthetic peptide corresponding to this extension (ExbB$_{Sm}$ 1-44) did not show

any indication of secondary structure (unpublished results). Since this region is predicted to

be in the periplasm where the C-terminal domain of HasB, HasB$_{CTD}$ is also located, we

investigated the interaction of the ExbB$_{Sm}$ 1-44 synthetic peptide with HasB$_{CTD}$ by NMR. The

analysis of chemical shift perturbations (CSP) of amide resonances of HasB$_{CTD}$ upon addition

of the peptide, showed that the chemical environment of their corresponding residues was

modified as a consequence of their interaction (Supplementary Figure S13). The residues

showing the highest chemical shift perturbations are mapped on HasB$_{CTD}$ structure 2M2K in

Figure 4. They are mainly located on the helical face of HasB$_{CTD}$ forming a continuous surface

of interaction. An additional small pocket at the C-terminus of the protein is also involved in

this interaction.

4b An ExbB_{Sm} mutant devoid of the N-terminal extension is affected in its function

To assess the potential function of the N-terminal periplasmic extension of ExbB$_{Sm}$, we

engineered a mutant lacking this extension (see Materials and Methods for details) and tested

its effect on the growth of a C600Δ*hemA*Δ*exbBD*Δ*tonB* strain harbouring plasmids

pAMhasISRADEB and pBADexbBD$_{Sm}$ or its derivatives. To get high sensitivity and numerical

data in our growth tests, we switched to growth in liquid medium in 48-well plates (see

Materials and Methods for details). We could then show that the lag between the start of the

experiment and the onset of growth is delayed by about 2 hours, for the mutant strain (in

which ExbB is devoid of its periplasmic extension), as compared to the WT strain. In addition,

the maximum OD reached after growth is slightly but significantly decreased (Supplementary

Figure S14, top). In this kind of experiment, the growth is related first to the expression of the

various <u>has</u> components under the control of their own regulatory elements, and to the

efficiency of the heme uptake process itself. Although we did not measure kinetics and extent

of the transcriptional activation of the <u>has</u> locus under these conditions, which is well beyond

the scope of our study, one possible interpretation of our results points to a likely role of the

N-terminal periplasmic extension in the activation of the transcription of the <u>has</u> locus. To

ensure that this difference was not actually due to a difference of expression of the two

constructs, we determined by Western blot their amounts in membrane preparations

(Supplementary Figure 14, bottom). As our anti-ExbB$_{Sm}$ antibody was not sensitive enough for

this measurement, we used His-tagged versions of ExbB. The experiment shows that the

amount of the ExbB variant deleted of its N-terminal extension is at least equal, if not slightly

higher than the amount of the wild-type protein, ruling out a possible decrease in protein

concentration.

4c Swapping residues in TM1 between ExbB$_{Sm}$ and ExbB$_{Ec}$ strongly suggests that ExbB TM1

interacts with HasB/TonB.

As mentioned above, HasB is a dedicated Has system TonB ortholog, that is functional with

ExbBD$_{Sm}$, but not with ExbBD$_{Ec}$. This clearly points to the looser specificity of ExbBD$_{Sm}$, as

compared to ExbBD$_{Ec}$, as ExbBD$_{Sm}$ is functional with both HasB and TonB$_{Ec}$. One reason for this

could be the higher stability of HasB itself in the cytoplasmic membrane, as it is known that

TonB can only accumulate in the cytoplasmic membrane when its ExbB and ExbD partners are

present. It is hypothesized that TonB is proteolytically degraded in the absence of ExbBD. We

therefore tested to what level HasB accumulates in the presence of either ExbBD$_{Ec}$ or ExbBD$_{Sm}$

and found that in the absence of either complexes, HasB was hardly detectable, whereas it

readily accumulated in their presence (Supplementary Figure S15). This observation therefore

contradicts the interpretation that HasB could be inactive in the presence of ExbBD$_{Ec}$ due to a

lack of accumulation, and also indicated that most likely HasB also interacts with ExbBD$_{Ec}$ so

as to be stabilized (i. e. withstanding proteolytic degradation), but in a non-functional manner.

Given the general framework put forward for the Mot complexes where the MotA (equivalent

of ExbB) rotation around MotB (the equivalent of ExbD) drives the rotation of the flagellum

basal ring [18, 19] we tested the possibility that ExbB residues exposed at the surface of the

protein might be involved in this functional interaction with HasB/TonB. The structure of ExbB

highlighted the conservation of the residues across the different bacterial genomes and

showed that the membrane-facing residues of the TM region of helix α2 and the cytoplasmic

residues were the least conserved as shown in the alignment and the structure

(Supplementary Figures S1 and S9). There was one conspicuous stretch of residues located in

the cytoplasmic leaflet of TM helix 1 (see Supplementary Figure S9): residues 76-88 of ExbB$_{Sm}$

-TILFAKGSELLRA-, corresponding to residues 39-51 of ExbB$_{Ec}$ -AIFFSKSVEFFNQ-, that is

the least conserved one between the two proteins. In this stretch, the conserved I, K and E

residues point toward the innermost part of the protein and the non-conserved residues point

to the outside and are available for protein-protein interaction (Supplementary Figure S16). It

was tempting to speculate that this stretch might correspond to an interaction region between

ExbB$_{Sm}$ and HasB. We therefore made a chimeric ExbB$_{Ec}$ protein with its 39-51 region replaced

by residues 76-88 from ExbB$_{Sm}$ named ExbB$_{Ec-Sm76-88}$ and asked whether this chimeric ExbB$_{Ec}$

protein would allow growth in the presence of HasB (see Supplementary Figure S3bis for the

constructions). As shown in Supplementary Figure S3, this chimeric protein ExbB$_{Ec-Sm76-88}$ was

active with both TonB$_{Ec}$ and HasB as it promotes growth with both hasISRADE and hasISRADEB

plasmids, strongly suggesting that this region is involved in specific interactions with HasB. We

also made the converse change where the 76-88 region of ExbB$_{Sm}$ was replaced by the 39-51

region from ExbB$_{Ec}$ (ExbB$_{Sm-Ec39-51}$). In our *in vivo* tests there was no major difference between

ExbB$_{Sm}$ and ExbB$_{Sm-Ec39-51}$ and ExbB$_{Sm-Ec39-51}$ was still active with HasB, although to a slightly

lower extent. To better locate the region responsible for specificity, we produced one

additional mutant where only residues 76-84 (TILFAKGSE) from ExbB$_{Sm}$ were exchanged

for residues 39-47 (AIFFSKSVE) from ExbB$_{Ec}$ yielding ExbB$_{Ec-Sm76-84}$. ExbB$_{Ec-Sm76-84}$ was still

active with HasB, showing that this 9-residue stretch is sufficient to transform ExbBEc into a

Sm-like phenotype (see Supplementary Figure S3 and S3bis for the visualisation of the

constructions). This set of experiments shows that intramembrane functional zones of ExbB

are crucial for the interaction between ExbB and HasB.

In summary, we showed using NMR and culture growth assays that ExbB$_{Sm}$ N-terminal

extension and a few residues in the α2 helix TM part are likely involved in specific interactions

with HasB.

Discussion

1) ExbBD$_{Sm}$ is a 5:2 complex at pH 8. Our structures were solved at pH 8, and examination of

the 2D class averages only showed pentameric structures and not hexameric ones, contrarily

to what has been observed in the Maki-Yonekura study, where ExbB was mostly seen as a

hexamer in the high pH regime[16]. In their study, the solubilizing detergent was DDM, that

was subsequently exchanged for either C8E4 or C10E5. These shorter chain detergents might

have led to some destabilization of the pentameric structure and provoked the conversion to

the hexameric form. On the other hand, our data agree with those published by Lloubes,

Buchanan et al. [15, 17]. *In vivo*, a stoichiometry of 7ExbB, 2ExbD and 1TonB has been

determined in *E. coli* [27]. Although one cannot completely rule out some biases in this indirect

measurement, a physiologically relevant reason for an excess of ExbB would be to provide a

permanent scaffold to which TonB and/or ExbD could associate with, upon TonB complex

dissociation, that might occur during the catalytic cycle, as has already been proposed [28].

2) ExbB and ExbBD are co-purified with endogenous lipids

In their last study, Celia *et al.* reconstituted ExbBD from *E. coli* into nanodiscs with added

phospholipids from *E. coli*, and found mostly PE bound to ExbBD [17]. It is unclear whether

this phospholipid was a genuine tightly bound lipid co-purified or whether it came from the

lipids added during reconstitution in the nanodiscs. In our case, we did not add any lipid and

we found mostly PG and some PE in the ExbB case (and also CL for ExbBD) using mass

spectrometry. The cryo-EM structure shows that they are located in the inner leaflet, as in the

E. coli complex. PG and CL are negatively charged, whereas PE is zwitterionic. This might be

pertaining to the functioning of the complex, as the lipids are bound to a highly positively

charged interface, very close to the fenestrations seen in the structure and PE might therefore

be less tightly bound than PG or CL. Further analysis by mass spectrometry of the aliphatic

chains of the lipids present in our sample also revealed a specific composition in their length

and unsaturated nature.

One point also worth mentioning is the existence of "cavities" inside the ExbB tunnel,

potentially allowing the passage of proton/hydronium ion up to the Asp 25 residue of ExbD,

deeply embedded in the apolar medium of ExbB, and therefore with a pKa close to

physiological pH, allowing easy protonation/deprotonation steps of Asp 25 side chain. The

identified tunnels connect the cytoplasm to the periplasm *via* Asp 25, allowing us to propose

a possible trajectory for the proton transfer.

3) Motor model proposal

Regarding the coupling of the pmf dissipation with the mechanical work carried out by

MotAB/TolQR/ExbBD complexes, several models have been put forward that might be more

or less easy to reconcile with the structural data. A detailed mechanism was proposed for *C.*

jejuni MotAB complex, where cycles of protonation/deprotonation of the conserved Asp of

the MotB TM (equivalent to the conserved Asp 25 of the ExbD TM) are coupled to the rotation

in discrete steps of the MotA pentamer around MotB axis, provided that the periplasmic

domain of MotB is anchored to the peptidoglycan layer *via* its PG binding site [18]. Our ExbBD

structure indeed shows that ExbD monomers have two orientations relative to ExbB: one is

such that Asp 25 faces Thr 208 of ExbB and the other Asp 25 faces a hydrophobic region at the

interface of two ExbB monomers. These two states are compatible with

protonation/deprotonation cycles accompanied by a change in the environment upon

rotation.

Moreover, our NMR results show that ExbB interacts *via* its alanine and proline-rich

periplasmic N-terminal region with the HasB periplasmic domain, and growth studies of

mutant bacteria show that ExbB$_{Sm}$ α2 helix TM region likely carries specificity determinants of

the interaction with HasB; this last observation is in line with a previous TonB TM point mutant

partially suppressed by an ExbB α2 point mutant [29]. It also appears that HasB stabilisation

can occur without full functionality, as ExbBD$_{Ec}$ protects HasB from degradation, without

productively interacting with it. One possible interpretation is that stabilisation of HasB is

provided by its interaction with ExbD$_{Ec}$ periplasmic domain.

The MotA-MotB model also posits that MotA outer region interacts with the rotor of the

flagellum and thus that MotA rotation drives the rotation of the flagellum. A similar interaction

between the outermost TM of ExbB (α2) and the TM of HasB/TonB is supported by our

mutagenesis data, as we could swap the specificity of ExbB$_{Ec}$ towards HasB by exchanging a

short stretch of residues between ExbB$_{Sm}$ and ExbB$_{Ec}$. Moreover, the sequences of the TM

regions of HasB and TonB are different, apart from a conserved Histidine. Structure

comparison of the swapped regions shows that ExbB$_{Sm}$ has smaller residues that may be better

accommodated and less specific than the *E. coli* sequence (Supplementary Figure S16).

On the part of the periplasmic space close to the outer membrane, it is well known that the

C-terminal domain of HasB/TonB interacts with the "TonB/HasB box" of the outer membrane

receptor, and that this interaction is essential for the entry of the substrate bound to the

extracellular side of the TBDT into the periplasm [30]. AFM experiments also showed that by

exerting a pulling force on the C-terminal domain of TonB bound to the BtuB TBDT, a partial

unfolding of the TBDT plug occurs, potentially leading to the diffusion of the substrate in the

periplasm [31]. Kinetic measurements showed that the *in vivo* entry of a siderophore is a

rather slow process: during a cell division time a bacterial cell has to take up ca. 300000 iron

atoms from its surroundings. Under full induction, there are roughly 10-15000 FepA
siderophore receptors per cell and 1500 TonB complexes [27], meaning that each receptor
has to undergo 20 cycles during a generation, leading to a turnover time of the TonB complex
of ca 5-10 seconds. Several models have been proposed to account for the functioning of the
TonB complex, in conjunction with the entry of a substrate bound on the extracellular side of
a given receptor. Firstly, piston-like movements of TonB have been proposed, driving the
unfolding of the plug inside the receptor barrel. AFM experiments and steered molecular
dynamics also showed that monomeric TonB devoid of its TM segment was able to drive the
partial unfolding of the BtuB plug allowing thus vitamin B12 entry into the periplasm in a few
discrete steps [31]. It is also known that ExbD and TonB interact *via* their periplasmic parts
[32] and in particular residue 150 of TonB (positioned just upstream the C-terminal globular
domain) can make cross-links to the C-terminal domain of ExbD [33]. Secondly, the MotAB
model can be extrapolated to the ExbBD complex. In this model, ExbB would rotate around
ExbD, and this would drive the rotation of TonB, thanks to the specific interaction between
TonB TM and ExbB TM (in our case HasB TM and ExbB TMα2). It has also been hypothesized
that the C-terminal domain of TonB [34] and/or the C-terminal domain of ExbD have specific
binding sites for the peptidoglycan network [35]. This C-terminal domain can change
conformations along the catalytic cycle of protonation and deprotonation events at the
conserved Asp residue in ExbD TM. In addition, the periplasmic linker between the TM helix
and the C-terminal domain of TonB is rich in Pro and Lys residues and might adopt an extended
conformation of a sufficient length to span the periplasm [36]. We therefore propose a model
in which the force generated through rotation of TonB, driven by ExbB rotation, is not directly
transmitted to the TonB box of the receptor, but through the intermediary of the C-terminal
domain of ExbD, that might act as a wedge or a linchpin allowing force transmission and
converting the rotation into a pulling force exerted on the TonB/HasB box of the receptor.
Further studies are needed to test this model.
446
447 **Material and Methods**
448
Strains and plasmid construction
Strains, plasmids, and oligonucleotides used are shown Table S2.
-Plasmid pBADExbBD$_{Sm}$ was constructed after amplification on the genomic DNA from strain

S. marcescens Db11 of a ca. 1.42kb fragment using primers ExbBDSm5' and ExbBDSm3'. The

PCR product was purified digested with *Eco*RI and *Sph*I, and ligated with pBAD24 digested with

*Eco*RI and *Sph*I. Correct clones were selected after sequencing of the insert.

-Plasmid pBADExbBD$_{Sm}$His6 was constructed by first amplifying on pBADExbBDSm a ca. 0.4kb

fragment with the following oligonucleotides SphIHisCtexbDSm and BglIIexbDSm; after

amplification, the fragment was purified, digested with *Bgl*II and *Sph*I, and ligated with

pBADExbBDSm digested with the same enzymes. The correct clones were selected after

sequencing of the insert. In biological tests, this plasmid was undistinguishable from its parent

plasmid pBADExbBDSm.

-Plasmid pBADExbBD$_{Ec}$ was constructed by amplification on genomic MG1655 DNA of a ca.

1.7kb fragment with the following oligonucleotides ExbBD5c, and ExbBD3c; the PCR product

was purified, digested with *Eco*RI and *Sph*I, and ligated with pBAD24 digested with the same

enzymes. Correct clones were selected after sequencing.

-Plasmid pBADExbB$_{Sm}$His6 was constructed as follows: a PCR fragment was amplified on

pBADExbBDSm with the oligos PBADFOR and ExbBHis6, digested with *Eco*RI and *Sph*I, and

ligated with pBAD24 digested with the same enzymes. Correct clones were selected by

sequencing.

Plasmid pBADExbBD$_{Ec-Sm76-88}$ was constructed as follows: Plasmid pBADExbBD$_{Ec}$ was amplified

with the two following 5' phosphorylated oligonucleotides: ExbBEcSm76-88.1 and

ExbBEcSm76-88.2. After digestion with *Dpn*I and self-ligation, correct clones were selected by

sequencing and the mutated fragments recloned in pBAD24.

Plasmid pBADExbBD$_{Ec-Sm76-84}$ was constructed in the same manner, with the following

oligonucleotides pairs: ExbBEcSm 76-84.1 and ExbBEcSm 76-84.2.

In the same manner, plasmid pBADExbBD$_{Sm-Ec39-51}$ was constructed by PCR on the

pBADExbBD$_{Sm}$ plasmid with the following couple of 5' phosphorylated oligonucleotides:

ExbBSmEc39-51.1 and ExbBSmEc39-51.2.

Plasmid pBADExbB$_{Smdelextss}$D was constructed similarly, by using as a template pBADExbBDSm,

with the two phosphorylated oligonucleotides ExbBdelextss1 and ExbBdelextss2.

To construct pBADExbB$_{Smdelextss}$His6, a ca. 0.6kb *Eco*RI-*Kpn*I fragment from pBADExbB$_{Smdelextss}$D

was exchanged for the corresponding fragment of pBADExbBHis6.

Plasmid pAMhasISRADEB, was constructed by digesting pAMhasISRADE and pSYC7 [37] by

*Kpn*I and *Hin*dIII, and purifying fragments of respectively ca. 9 and 7kbases, ligating them

together to obtain a plasmid with the whole *has* locus on a low copy number plasmid
(pAM238) under the control of its endogenous regulation signals.

All constructions were carried out in *E. coli* strain XL1-Blue.

Strains and other plasmids used in this study are shown in Table S2 and are from the laboratory
collection.

Protein expression and purification

The BL21DE3(pExbBD$_{Sm}$His6/pBAD24) or BL21DE3(pExbB$_{Sm}$His6/pBAD24) were grown at 37°C
either in TB or MDM medium, and induced with 0.2% arabinose at 1.5-2OD600nm (TB) or 5-
6OD600nm (MDM) and the incubation continued for 3hrs. The cells were harvested by
centrifugation (10000g 20minutes 4°C), washed once in 20mMTris-HCl pH8.0, flash-frozen in
liquid N$_2$ and kept at -80°C. Cells were broken in a Cell disruptor (Constant, UK) at 1kbar (10g
of cells in 40ml final of 20mM Tris pH 8.0 containing protease inhibitor (Roche, EDTA free), at
4°C. Benzonase was added and after ca. 15 minutes, the solution was centrifuged for 1hr at
100000g at 4°C. The pellet (crude membrane preparation) was resuspended in 20mM Tris pH
8.0 plus protease inhibitor cocktail (Roche EDTA free), flash-frozen in liquid N$_2$ and kept at -
80°C.

The crude membrane preparation was solubilized in 20mM Tris pH 8.0, 20mM Imidazole,
100mM NaCl, 10% Glycerol, 0.8% LMNG (10g of equivalent whole-cell pellet solubilized in
40ml), plus protease inhibitor cocktail (Roche EDTA-free) for 30 minutes at 15°C. After
centrifugation (1hr 100000xg), the supernatant was incubated with 2.5ml Ni-Agarose beads
(Thermo-Fischer His-Pure Ni-NTA #88222) preequilibrated with the same buffer except for the
detergent concentration (0.0015% LMNG). After three hours of incubation on a rotating wheel
at 4°C, the beads were washed three times with 25ml of pre-equilibration buffer and then
eluted with two times 25ml of the pre-equilibration buffer containing 200mM Imidazole. The
eluate was concentrated and washed on 100kDa cut-off centrifugal device, in pre-
equilibration buffer without NaCl and Imidazole. The resulting sample was loaded on a monoQ
HR10-100 column equilibrated with 20mM Tris pH 8.0, 10% glycerol, 0.0015%LMNG and
eluted with a gradient from 0 to 1M NaCl in the same buffer. Peak fractions were collected
and concentrated as before. The concentrated sample was then loaded on a Superose 6
increase column equilibrated with 20mM Tris pH8.0, 100mM NaCl, 0.0015%LMNG. The peaks
fractions were collected, concentrated and their concentration determined by using the

theoretical absorption coefficient of either ExbB5D2 ($113790 M^{-1}cm^{-1}$), or ExbB5 ($104850 M^{-1}cm^{-1}$). They were then kept frozen at -80°C in aliquots until use. The SEC profiles as well as a gel of a representative sample after purification are shown in supplementary data. Previous attempts with DDM instead of LMNG yielded similar results, and LMNG was chosen, owing to its very low CMC and its lower background in cryo-EM.

Activity tests

Three types of tests were used, either in liquid medium or on agar plates.

1. Growth tests in liquid medium: *E. coli* C600 and its *ΔexbBD* derivatives were transformed with specified plasmids and their growth tested in liquid LB medium at 37°C, at various dipyridyl concentrations to induce iron starvation. OD_{600nm} was measured after overnight growth.

2. Growth tests on plates: the relevant plasmids were also transformed into *E. coli* C600 *ΔhemA*, a heme auxotroph strain (and derivatives thereof), and growth of the strains was assayed as follows. Briefly, cells were first grown in LB medium (supplemented with delta-aminolevulinic acid (50μg/ml) to bypass the effect of the *ΔhemA* mutation) at 37°C up to an OD_{600nm} of 1, and then mixed with melted top agar (0.6%agar in LB), and poured onto LB agar plates containing the appropriate antibiotics with arabinose at a concentration of 40μg/ml to induce expression of the genes under the P_{araBAD} promoter[38]. Wells, punched with Pasteur pipettes were filled with heme-albumin (at 5, 1 or 0.2μM), at two dipyridyl concentrations (100 or 200μM) to induce iron starvation. Plates were incubated overnight at 37°C and scored for growth around the wells. All experiments were performed in triplicate.

3. Growth curves in liquid medium: a few colonies of *E. coli* C600 *ΔhemAΔtonBΔexbBD*(pAMhasISRADEB + pBAD24 or derivatives thereof) were first inoculated in 4ml of LB medium at 37°C with the corresponding antibiotics, and 100μM dipyridyl, 4μg/ml arabinose but without delta-aminolevulinic acid. Once the culture reached an OD_{600nm} of ca. 1.2-1.5, it was diluted and inoculated in 48 well Greiner plates, in the same medium to which was added 0.4μM He-BSA, as a heme source. The initial OD_{600nm} of the cultures was 0.001. Each well contained 300μl of growth medium. Duplicates of each strain were made, and the plate was incubated at 37°C with vigorous shaking (500rpm) in a Clariostar Plus Microplate reader. OD_{600nm} was recorded every 30 minutes for 70 hours.

N-terminal sequencing

The N-terminus of ExbB$_{Sm}$ was determined at the Plateforme Protéomique de l'institut de microbiologie de la Méditerranée (Marseille), after blotting on a PVDF membrane a purified sample of ExbBD$_{His6}$ run on SDS-PAGE and determined to be APAAN.

Lipid extraction

Chloroform (0.20 mL) and methanol (0.40 mL) were sequentially added to a sample of the ExbB or ExbBD complex (0.1 mL). The sample was vortexed for 10 min at room temperature and chloroform (0.2 mL) and water (0.2 mL) were further added. The organic phase was collected and the extraction procedure was repeated on the remaining aqueous phase. Combined organic layers were evaporated to dryness under argon and stored at -20 °C.

Mass Spectrometry

Dry lipid extracts obtained from the ExbBD complex and from an *E. coli* lysate were solubilized in chloroform/methanol (50/50). Samples were then nano-electrosprayed using a TriVersa NanoMate (Advion Biosciences, Ithaca, USA) coupled to a Synapt G2-Si mass spectrometer (Waters Corporation, Manchester, UK). The instrument was calibrated in negative ion mode (for lipids) and positive ion mode (for proteins) from 50 m/z to 2000 m/z with NaI (50 mg/ml) with an accuracy of 0.8 ppm in resolution mode. The following settings were chosen: sampling cone 40 V, source offset 40 V, source temperature 80°C, trap gas flow 5 mL/min, helium cell gas flow 180 mL/min. MS/MS spectra were recorded using CID (Collision Induced Dissociation) with a normalized collision energy set up to 30.

To measure the mass of the intact proteins of the ExbBD and purified ExbB complexes, a desalting step was performed using micro Bio-Spin™ 6 (BIO-RAD) with 500mM ammonium acetate. Both samples were analyzed in denaturing conditions after a two-fold dilution with acetonitrile 4% formic acid. Mass spectra were acquired in positive ion mode. All molecular weights were measured after MaxEnt1 deconvolution into neutral species.

Cryo-EM grid preparation and data acquisition

3 µL of either purified ExbB$_{Sm}$ or ExbBD$_{Sm}$ complex at ca. 1 mg/mL was applied to C-Flat 1.2/1.3 holey carbon grids (Protochips Inc., USA) previously glow-discharged in air for 30s. Grids were blotted for 2s at blot force 1 and vitrified in liquid ethane using a Vitrobot mark IV (FEI company) operated at 10 °C and 100% relative humidity.

All data collection was performed with a Titan Krios (ThermoFisher Scientific) operated at 300 kV equipped with a K2 Summit direct electron detector (ThermoFisher Scientific) at the European synchrotron research facility, ESRF (Grenoble, France). Movies were recorded in electron counting mode with EPU software (ThermoFisher Scientific), aligned with Motioncor2 [39] and aligned images were processed with Gctf [40] using Scipion interface[41].

For ExbB data collection, 4567 movies were collected at a magnification of 165000x with a nominal pixel size of 0.827Å using a defocus range from -1.5 to -2.5µm. Movies of 56 frames were acquired using a dose rate of 8 electrons/Å^2/second over 7 seconds, yielding a cumulative exposure of 55.95 electrons/Å^2.

For ExbBD data collection, 4043 movies were collected at a nominal magnification of 139000 with a pixel size of 1.067Å. Movies of 48 frames were acquired using a dose rate of 4.6 electrons/Å^2/second over 12 seconds yielding a cumulative exposure of 55.2 electrons/Å^2.

Cryo-EM image processing and analysis

a) ExbB

For ExbB, aligned movies were processed with Gctf [40] and only images with a resolution higher than 4Å were kept; after visual inspection of the remaining images, processing was carried out with Relion-3 [42]. Particles were extracted using a 2-fold binning, issued from a manual picking and a 2-D classification of particles picked out from on 50 images. Automatic extraction was performed using the selected 2D class averages. After several rounds of 2D and 3D classification, 161k particles were selected for 3D refinement. They were corrected for local motion using Bayesian polishing option in Relion-3 and a post-refined map produced a 3.1Å overall resolution with a 5-fold symmetry (Supplementary Figure S17).

A homology model was built with the *Serratia marcescens* ExbB sequence using Phyre2 server from 5SV0 monomer and docked into the refined map. Refinement was done with Phenix Real Space Refine option with secondary structure, "non-crystallographic symmetry" and Ramachandran restraints [43] and graphically adjusted with Coot [44]. Lipid phosphatidyl

glycerol starting structure and geometry were built using Phenix eLBOW [45] and was fit in

the map using Coot then refined along with the protein model in Phenix.real_space_refine.

b) ExbBD

Movies were drift-corrected and dose-weighted using MotionCorr2 [39]. Aligned dose

weighted averages were imported into Cryosparc2 [46] and contrast transfer function was

estimated using CTFFIND4.1 [47]. Micrographs with poor CTF estimation statistics or high drift

profiles were discarded. The remaining 3028 micrographs were used for automated particle

picking. Particles were extracted, Fourier cropped to 2 Å/px and 2D classified. The best 2D

classes were used as templates for automated particle picking resulting in 1.3 million particles.

After several rounds of classification, the best 600k particles were submitted to 3D

classification by means of multi-class ab-initio Reconstruction and Heterogeneous

Refinement. 158 k particles belonging to the best resolved classes were corrected for local

motion, re-extracted and used in Non-Uniform Refinement. The resulting refined map has a

nominal resolution of 4.56 Å.

Based on the previous map a soft mask lining the micelle was designed in UCSF Chimera [48]

and used to signal subtract the corresponding micelle density of particles in the refined map.

Localized refinement of the signal subtracted particles produced a map of the complex with

an estimated resolution of 3.96 Å judging by FSC at 0.143 criterion (Supplementary Figure

S18). Data have been deposited both at the EMDB and PDB databases (EMDB 10789 and

11806, PDB 6YE4 and 7AJQ, for $ExbB_{Sm}$ and $ExbBD_{Sm}$ respectively, see Table I for refinement

statistics).

Unless otherwise specified, the structure figures were made using UCSF Chimera software

[48].

NMR experiments

The C-terminal periplasmic domain of HasB, $HasB_{CTD}$ comprising residues 133-263, was

produced and purified as previously reported [14, 30].The peptide corresponding to the

periplasmic   extension   of $ExbB_{sm}$

(A_1PAANPAVTESVAPTTAPAPAAAAPESITPVNPAPTIQPPETRG$_{44}$- numbering with reference to

the mature protein) was synthetized by *Proteogenix*.

NMR experiments were acquired at 293 K on a 600 MHz Bruker Avance III spectrometer equipped with a TCI cryoprobe. The spectra were processed with NMRpipe [49] and analyzed with CcpNmr Analysis 2.4 software [50]. Proton chemical shifts were referenced to 2,2-dimethyl-2-silapentane-5 sulfonate as 0 ppm. ^{15}N were referenced indirectly to DSS [51] (Wishart et al., 1995). ^{1}H–^{15}N HSQC experiments were acquired on 0.15 mM $HasB_{CTD}$ in 50mM sodium phosphate pH 7, 50 mM NaCl with or without the peptide. Aliquots from a solution of peptide at 25mg/ml in the same buffer were added to the protein sample at 2:1 and 10:1 ratios. Chemical shift perturbations (CSPs) of backbone amide cross-peaks were quantified by using the equation $CSP = [\Delta \delta H^2 + (\Delta \delta N \times 0.15)^2]^{1/2}$, where $\Delta \delta H$ and $\Delta \delta N$ are the observed ^{1}H and ^{15}N chemical shift differences between the two experimental conditions. The ^{1}H and ^{15}N resonance assignments were from Lefevre et al 2007 [52].

Other biochemical methods

SDS-PAGE and immunodetection with anti-HasB or anti-His6 antibodies (Abcam [HIS.H8] (ab18184)) on whole cells or membrane preparations were carried out following standard protocols. Secondary antibodies were coupled to alkaline phosphatase.

BLAST search

$ExbB_{Sm}$ was used as a search for BLAST for orthologs in complete bacterial genomes, focusing on "long" ExbB's. Top hits were in *Serratia*, *Yersinia*, *Dickeya*, *Erwinia* and *Pseudomonas*. Those genera were later excluded from successive BLAST searches to obtain orthologs in other genera with higher p-value.

Acknowledgments This work was supported by grants from the ANR (HEMESTOCKEXCHANGE ANR-12-BSV3-0022-01 and LABEX DYNAMO ANR-11-LABEX-0011-01). We thank Nathalie Dautin for helpful suggestions and critical reading of the manuscript. We acknowledge Emmanuel Frachon and Christophe Thomas from the Institut Pasteur platform "Production of recombinant proteins" for large scale cultures used in this work, European Synchrotron Radiation Facility for the provision of time on the Titan Krios at beamline CM01, the Institut Pasteur Biological NMR Technological Platform for the use of the NMR spectrometers, the staff at the ESRF CM01 facility in Grenoble for help in the data collection, the staff at the Marseille proteomics facility and Djalila Djouadi and Diana Basto who initially contributed to this project. We thank M. Nilges for interest in this work.

Figure legends

Figure 1:

Weblogo (http://weblogo.threeplusone.com/) representation of 131 « long » ExbB aminoacid sequences; those with much shorter periplasmic extensions have been excluded from the alignment. The orange boxes show the position of the three TM segments from the *E. coli* 5SV0 structure.

Figure 2:

Representation of the pentameric structure of ExbB$_{Sm}$ solved by cryo-EM. A: the density map is shown, each monomer with distinct shades of blue, view from the side. The grey densities represent PG molecules. B: view from the cytoplasmic side. C: representation of one PG density together with the fitting of the PG molecule inside the density, and with the stick representation of the nearby ExbB residues using the Coot software [44]. D: superposition of a monomer of ExbB$_{Ec}$ (dark grey) and ExbB$_{Sm}$ (red), represented as ribbon. The membrane thickness is represented by the dotted lines and the transmembrane segments TM1 (part of α2 helix), TM2 (part of α6 helix) and TM3 (part of α7 helix) are indicated.

Figure 3:

Structure of ExbBD$_{Sm}$ solved by cryo-EM. A: side view, the same color code as in Figure 2 for the ExbB monomers is used. A: side view; B: view from the periplasmic space; the two ExbD helices are seen respectively in red and magenta. C: clipped side view, with the helix of each ExbD monomer represented (red and magenta). D and E, side and top views showing the model built in the density maps.

Figure 4:

Interaction of HasB$_{CTD}$ with the periplasmic fragment of ExbB$_{Sm}$, as detected by NMR. HasB$_{CTD}$ residues shown on its structure (PDB code 2M2K) exhibiting the highest CSP in the presence of the periplasmic peptide of ExbB$_{sm}$ are coloured in red (Top: surface representation; bottom: cartoon representation). Right: Superposed ^{1}H–^{15}N HSQC spectra of 0.15 mM ^{15}N-labelled HasB$_{CTD}$ in 50mM sodium phosphate, pH 7, 50 mM NaCl in the presence (red) or absence (black) of the periplasmic peptide of ExbB$_{sm}$.

Complex	ExbB	ExbB-ExbD
Deposition ID	EMD-10789, PDB 6YE4	EMD-11806, PDB ID 7AJQ
Data collection and processing		
Microscope type, camera	ESRF Titan Krios, Gatan K2	ESRF Titan Krios, Gatan K2
Magnification	165000	139000
Voltage (kV)	300	300
Electron exposure (e–/Å^2)	56	55
Defocus range (μm)	-1.5 to -2.5	-1 to -3
Pixel size (Å)	0.87	1.067
Initial particle images (no.)	850000	1373000
Final particle images (no.)	157000	158000
Map resolution (Å)	2.97	4
FSC threshold	0.143	0.143
Map resolution range (Å)	3.2-5.5	3.97-8
Symmetry imposed	C5	C1
Initial model used	Pdb id. 5SV0	Our ExbB structure
Model composition		
Non-hydrogen atoms	9145	9256
Protein residues	1180	1229
Ligands	PGT : 5	0
B factors (Å^2)		
Protein min/max/mean	21.58/ 145.06/ 45.97	93.07/ 411.58/ 169.35
Ligand min/max/mean	45.43/ 46.91/ 46.12	
R.m.s. deviations		
Bond length (Å) (# >4σ)	0.026 (3)	0.003 (0)
Bond angles (°) (# > 4σ)	1.56 (8)	0.573 (4)
Validation		
Refined model CC	0.85	0.77
MolProbity score	1.1	2.03
Clashscore	3	15.3
Poor rotamers (%)	0	0
Ramachandran plot		
Favored (%)	99	95
Allowed (%)	1	5
Disallowed (%)	0	0

723 Table 1: Data collection, processing, and refinement statistics for ExbB$_{Sm}$ and ExbBD$_{Sm}$.

733   Figure 1

761   Figure 2
762
763

Figure 3

806 Figure 4

C-ter

N-ter

848   **References**
849

1.   Noinaj, N., et al., *TonB-dependent transporters: regulation, structure, and function.*
Annu Rev Microbiol, 2010. **64**: p. 43-60.
2.   Grinter, R. and T. Lithgow, *The structure of the bacterial iron-catecholate transporter*
Fiu suggests that it imports substrates via a two-step mechanism. J Biol Chem, 2019.
294(51): p. 19523-19534.
3.   Cascales, E., R. Lloubes, and J.N. Sturgis, *The TolQ-TolR proteins energize TolA and*
share homologies with the flagellar motor proteins MotA-MotB. Mol Microbiol, 2001.
42(3): p. 795-807.
4.   Braun, V., *The structurally related exbB and tolQ genes are interchangeable in*
conferring tonB-dependent colicin, bacteriophage, and albomycin sensitivity. J
Bacteriol, 1989. **171**(11): p. 6387-90.
5.   Eick-Helmerich, K. and V. Braun, *Import of biopolymers into Escherichia coli:*
nucleotide sequences of the exbB and exbD genes are homologous to those of the
tolQ and tolR genes, respectively. J Bacteriol, 1989. **171**(9): p. 5117-26.
6.   Braun, V. and C. Herrmann, *Evolutionary relationship of uptake systems for*
biopolymers in Escherichia coli: cross-complementation between the TonB-ExbB-ExbD
and the TolA-TolQ-TolR proteins. Mol Microbiol, 1993. **8**(2): p. 261-8.
7.   Fischer, E., K. Gunter, and V. Braun, *Involvement of ExbB and TonB in transport across*
the outer membrane of Escherichia coli: phenotypic complementation of exb mutants
by overexpressed tonB and physical stabilization of TonB by ExbB. J Bacteriol, 1989.
171(9): p. 5127-34.
8.   Ahmer, B.M., et al., *Characterization of the exbBD operon of Escherichia coli and the*
role of ExbB and ExbD in TonB function and stability. J Bacteriol, 1995. **177**(16): p.
4742-7.
9.   Higgs, P.I., P.S. Myers, and K. Postle, *Interactions in the TonB-dependent energy*
transduction complex: ExbB and ExbD form homomultimers. J Bacteriol, 1998.
180(22): p. 6031-8.
10.   Braun, V., et al., *Energy-coupled transport across the outer membrane of Escherichia*
coli: ExbB binds ExbD and TonB in vitro, and leucine 132 in the periplasmic region and
aspartate 25 in the transmembrane region are important for ExbD activity. J
Bacteriol, 1996. **178**(10): p. 2836-45.
11.   Gaisser, S. and V. Braun, *The tonB gene of Serratia marcescens: sequence, activity*
and partial complementation of Escherichia coli tonB mutants. Mol Microbiol, 1991.
5(11): p. 2777-87.
12.   Paquelin, A., et al., *Characterization of HasB, a Serratia marcescens TonB-like protein*
specifically involved in the haemophore-dependent haem acquisition system. Mol
Microbiol, 2001. **42**(4): p. 995-1005.
13.   Benevides-Matos, N., C. Wandersman, and F. Biville, *HasB, the Serratia marcescens*
TonB paralog, is specific to HasR. J Bacteriol, 2008. **190**(1): p. 21-7.
14.   Lefevre, J., et al., *Modulation by substrates of the interaction between the HasR outer*
membrane receptor and its specific TonB-like protein, HasB. J Mol Biol, 2008. **378**(4):
p. 840-51.
15.   Celia, H., et al., *Structural insight into the role of the Ton complex in energy*
transduction. Nature, 2016. **538**(7623): p. 60-65.

16. Maki-Yonekura, S., et al., *Hexameric and pentameric complexes of the ExbBD energizer in the Ton system.* Elife, 2018. **7**.

17. Celia, H., et al., *Cryo-EM structure of the bacterial Ton motor subcomplex ExbB–ExbD provides information on structure and stoichiometry.* Communications Biology, 2019. **2**(1): p. 358.

18. Santiveri, M., et al., *Structure and Function of Stator Units of the Bacterial Flagellar Motor.* Cell, 2020. **183**(1): p. 244-257 e16.

19. Deme, J.C., et al., *Structures of the stator complex that drives rotation of the bacterial flagellum.* Nat Microbiol, 2020.

20. Ihsan, M.Z., et al., *Gene Mining for Proline Based Signaling Proteins in Cell Wall of Arabidopsis thaliana.* Front Plant Sci, 2017. **8**: p. 233.

21. Mishra, A.K., et al., *Tryptophan-Rich and Proline-Rich Antimicrobial Peptides.* Molecules, 2018. **23**(4).

22. Ball, L.J., et al., *Recognition of proline-rich motifs by protein-protein-interaction domains.* Angew Chem Int Ed Engl, 2005. **44**(19): p. 2852-69.

23. Marchler-Bauer, A., et al., *CDD: a curated Entrez database of conserved domain alignments.* Nucleic Acids Res, 2003. **31**(1): p. 383-7.

24. Letoffe, S., P. Delepelaire, and C. Wandersman, *Free and hemophore-bound heme acquisitions through the outer membrane receptor HasR have different requirements for the TonB-ExbB-ExbD complex.* J Bacteriol, 2004. **186**(13): p. 4067-74.

25. Li, H., A.D. Robertson, and J.H. Jensen, *Very fast empirical prediction and rationalization of protein pKa values.* Proteins, 2005. **61**(4): p. 704-21.

26. Langsetmo, K., J.A. Fuchs, and C. Woodward, *The conserved, buried aspartic acid in oxidized Escherichia coli thioredoxin has a pKa of 7.5. Its titration produces a related shift in global stability.* Biochemistry, 1991. **30**(30): p. 7603-9.

27. Higgs, P.I., R.A. Larsen, and K. Postle, *Quantification of known components of the Escherichia coli TonB energy transduction system: TonB, ExbB, ExbD and FepA.* Mol Microbiol, 2002. **44**(1): p. 271-81.

28. Ollis, A.A., et al., *Cytoplasmic membrane protonmotive force energizes periplasmic interactions between ExbD and TonB.* Mol Microbiol, 2009. **73**(3): p. 466-81.

29. Larsen, R.A., et al., *Partial suppression of an Escherichia coli TonB transmembrane domain mutation (delta V17) by a missense mutation in ExbB.* Mol Microbiol, 1994. **13**(4): p. 627-40.

30. de Amorim, G.C., et al., *The structure of HasB reveals a new class of TonB protein fold.* PLoS One, 2013. **8**(3): p. e58964.

31. Hickman, S.J., et al., *Gating of TonB-dependent transporters by substrate-specific forced remodelling.* Nat Commun, 2017. **8**: p. 14804.

32. Garcia-Herrero, A., et al., *The solution structure of the periplasmic domain of the TonB system ExbD protein reveals an unexpected structural homology with siderophore-binding proteins.* Mol Microbiol, 2007. **66**(4): p. 872-89.

33. Ollis, A.A. and K. Postle, *The same periplasmic ExbD residues mediate in vivo interactions between ExbD homodimers and ExbD-TonB heterodimers.* J Bacteriol, 2011. **193**(24): p. 6852-63.

34. Klebba, P.E., *ROSET Model of TonB Action in Gram-Negative Bacterial Iron Acquisition.* J Bacteriol, 2016. **198**(7): p. 1013-21.

35. Boags, A.T., F. Samsudin, and S. Khalid, *Binding from Both Sides: TolR and Full-Length*
OmpA Bind and Maintain the Local Structure of the E. coli Cell Wall. Structure, 2019.
27(4): p. 713-724 e2.

36. Kohler, S.D., et al., *The proline-rich domain of TonB possesses an extended polyproline*
II-like conformation of sufficient length to span the periplasm of Gram-negative
bacteria. Protein Sci, 2010. **19**(4): p. 625-30.

37. Létoffé, S., J.M. Ghigo, and C. Wandersman, *Identification of two components of the*
Serratia marcescens metalloprotease transporter: protease SM secretion in
Escherichia coli is TolC dependent. J Bacteriol, 1993. **175**(22): p. 7321-8.

38. Guzman, L.M., et al., *Tight regulation, modulation, and high-level expression by*
vectors containing the arabinose PBAD promoter. J Bacteriol, 1995. **177**(14): p. 4121-
30.

39. Zheng, S.Q., et al., *MotionCor2: anisotropic correction of beam-induced motion for*
improved cryo-electron microscopy. Nat Methods, 2017. **14**(4): p. 331-332.

40. Zhang, K., *Gctf: Real-time CTF determination and correction.* J Struct Biol, 2016.
193(1): p. 1-12.

41. Gomez-Blanco, J., et al., *Using Scipion for stream image processing at Cryo-EM*
facilities. J Struct Biol, 2018. **204**(3): p. 457-463.

42. Zivanov, J., et al., *New tools for automated high-resolution cryo-EM structure*
determination in RELION-3. Elife, 2018. **7**.

43. Afonine, P.V., et al., *Real-space refinement in PHENIX for cryo-EM and*
crystallography. Acta Crystallogr D Struct Biol, 2018. **74**(Pt 6): p. 531-544.

44. Emsley, P., et al., *Features and development of Coot.* Acta Crystallogr D Biol
Crystallogr, 2010. **66**(Pt 4): p. 486-501.

45. Moriarty, N.W., R.W. Grosse-Kunstleve, and P.D. Adams, *electronic Ligand Builder*
and Optimization Workbench (eLBOW): a tool for ligand coordinate and restraint
generation. Acta Crystallogr D Biol Crystallogr, 2009. **65**(Pt 10): p. 1074-80.

46. Punjani, A., et al., *cryoSPARC: algorithms for rapid unsupervised cryo-EM structure*
determination. Nat Methods, 2017. **14**(3): p. 290-296.

47. Rohou, A. and N. Grigorieff, *CTFFIND4: Fast and accurate defocus estimation from*
electron micrographs. J Struct Biol, 2015. **192**(2): p. 216-21.

48. Pettersen, E.F., et al., *UCSF Chimera--a visualization system for exploratory research*
and analysis. J Comput Chem, 2004. **25**(13): p. 1605-12.

49. Delaglio, F., et al., *NMRPipe: a multidimensional spectral processing system based on*
UNIX pipes. J Biomol NMR, 1995. **6**(3): p. 277-93.

50. Vranken, W.F., et al., *The CCPN data model for NMR spectroscopy: development of a*
software pipeline. Proteins, 2005. **59**(4): p. 687-96.

51. Wishart, D.S., et al., *1H, 13C and 15N chemical shift referencing in biomolecular NMR.*
J Biomol NMR, 1995. **6**(2): p. 135-40.

52. Lefèvre, J., et al., *(1)H, (13)C and (15)N resonance assignments of the C-terminal*
domain of HasB, a specific TonB like protein, from Serratia marcescens. Biomol NMR
Assign, 2007. **1**(2): p. 197-9.