Reflectance spectroscopy allows rapid, accurate, and non-destructive estimates of functional traits from pressed leaves

Shan Kothari[1,2,4], Rosalie Beauchamp-Rioux[1], Etienne Laliberté[1], Jeannine Cavender-Bares[2,3]

[1]Institut de recherche en biologie végétale, Département de sciences biologiques, Université de Montréal, 4101 rue Sherbrooke Est, Montréal, QC, Canada H1X 2B2
[2]Department of Plant and Microbial Biology, University of Minnesota, 1479 Gortner Ave, St. Paul, MN, United States 55108
[3]Department of Ecology, Evolution, and Behavior, University of Minnesota, 1479 Gortner Ave, St. Paul, MN, United States 55108
[4]Corresponding author: shan.kothari@umontreal.ca

Running headline: Estimating traits from pressed-leaf spectra

Abstract

1. More than ever, ecologists seek to employ herbarium collections to estimate plant functional traits from the past and across biomes. However, many trait measurements are destructive, which may preclude their use on valuable specimens. Researchers increasingly use reflectance spectroscopy to estimate traits from fresh or ground leaves, and to delimit or identify taxa. Here, we extend this body of work to non-destructive measurements on pressed, intact leaves, like those in herbarium collections.

2. Using a dataset with 618 samples from 68 species, we used partial least-squares regression to build models linking pressed-leaf reflectance spectra to a broad suite of traits, including leaf mass per area (LMA), leaf dry matter content (LDMC), equivalent water thickness, carbon fractions, pigments, and twelve elements. We compared these models to those trained on fresh- or ground-leaf spectra of the same samples.

3. Our pressed-leaf models were best at estimating LMA ($R^2 = 0.917$; %RMSE = 7.30), C ($R^2 = 0.774$; %RMSE = 11.8), and N ($R^2 = 0.772$; %RMSE = 11.5), followed by water-related traits, carbon fractions, and pigments (all $R^2 = 0.554$-0.763; %RMSE = 12.3-18.6). Other than Ca, Na, and P, remaining elements were predicted poorly ($R^2 < 0.5$, %RMSE > 20). For most chemical traits, pressed-leaf models performed better than fresh-leaf models, but worse than ground-leaf models. Pressed-leaf models were worse than fresh-leaf models for estimating LMA and LDMC,

but better than ground-leaf models for LMA. Finally, in a subset of samples, we used partial

least-squares discriminant analysis to classify specimens among 10 species with near-perfect

accuracy from pressed- (>97%), and ground-leaf (>96%) spectra, and slightly lower accuracy

from fresh-leaf spectra (>89%).

4. These results show that applying spectroscopy to pressed leaves is a promising way to estimate

leaf functional traits and identify species without destructive analysis. Pressed-leaf spectra might

combine advantages of fresh and ground leaves: like fresh leaves, they retain some of the spectral

expression of leaf structure; but like ground leaves, they avoid the masking effect of water

absorption. Our study has far-reaching implications for capturing the wide range of functional and

taxonomic information in the world's preserved plant collections.

Keywords: functional traits; herbarium collections; leaf chemistry; partial least-squares regression

(PLSR); reflectance spectroscopy; species identification

Introduction

The world's herbaria together contain more than 390 million specimens (Thiers 2020), which are a rich

source of information about global plant diversity. Herbarium specimens are collected for many

reasons—for example, to document where a species is present, or to serve as vouchers for taxonomic

studies. But these specimens are often repurposed for new ends, unforeseen by the collectors (Heberling

& Isaac 2017; Meineke et al. 2018). More than ever, ecologists and evolutionary biologists seek to use

herbarium specimens to measure functional traits (Heberling 2021): for example, to evaluate long-term

changes, often in response to human activity (Willis et al. 2017; Lang et al. 2018); to fill in gaps in sparse

trait databases (Perez et al. 2020); or to conduct comparative studies of clades (e.g. Jardine et al. 2020).

Measuring functional traits on herbarium specimens carries the promise of letting us reach the

inaccessible, including the past or distant parts of the world. Using herbarium specimens also allows

researchers to benefit from the taxonomic skill of collectors and curators and refer back to the same

specimens for further use—for example, as sources of genetic data, or as references for species

identification from new collections (Heberling 2021). By repurposing specimens towards new ends,

researchers can address ecological and evolutionary questions that require merging functional, genetic,

and distributional data at global scales.

Many functional trait measurements require destructive sampling—for example, by grinding up tissue for

chemical analyses. Such measurements include most protocols to determine the elemental or molecular

composition of a sample (Pérez-Harguindeguy et al. 2013). Because herbarium specimens are

irreplaceable—especially those from historical collections—curators may hesitate to let them be

destroyed for ecological research, even in part. Non-destructive techniques to estimate leaf traits from

herbarium specimens would make it more feasible to use specimens in functional ecology.

Reflectance spectroscopy is a technique often used to estimate foliar functional traits non-destructively

(Curran 1989; Jacquemoud & Ustin 2020). A reflectance spectrum is a measurement of the reflectance of

light from a surface across a range of wavelengths. For leaf tissue, researchers tend to measure reflectance

between about 350 and 2500 nm, a range that includes over 98% of solar radiation reaching Earth's

surface (American Society for Testing and Materials, 2006). Because the leaf's chemical and structural

makeup determines how it absorbs, reflects, and transmits solar radiation, reflectance within this range

carries information about many plant traits. These traits are shaped by evolutionary history, such that leaf

reflectance spectra can also be used to understand trait evolution or distinguish lineages (Meireles et al.

2020a).

Two main approaches exist to estimate traits from entire reflectance spectra. First, physics-based radiative

transfer models like PROSPECT can be inverted on fresh-leaf spectra to estimate a handful of traits

whose optical properties are well understood (Jacquemoud & Baret 1990; Féret et al. 2017). Second,

statistical models, often created using machine learning techniques like partial least-squares regression

(PLSR), can estimate an even wider range of traits, albeit in a less mechanistic (and perhaps less general)

way (Serbin & Townsend 2020). Many ecologically important traits lack a unique set of strong absorption

features, complicating the development of radiative transfer models. For example, nitrogen is found alike

in various proteins (e.g. RuBisCO) and pigments (Wang et al. 2018) in ratios that vary from plant to

plant. Moreover, absorption features of compounds often overlap, especially after being broadened by

multiple scattering (Curran 1989). A multivariate empirical approach based on techniques like PLSR or

stepwise regression gives researchers the flexibility to predict complex traits (Curran 1989). Likewise,

multivariate classification techniques like partial least-squares discriminant analysis (PLS-DA) are often

used to take advantage of the full spectrum in discriminating species, lineages, or other kinds of biological

classes (Clark et al. 2005; Meireles et al. 2020b).

Empirical approaches like PLSR have been widely used to estimate plant traits from spectroscopic data

measured on fresh or ground leaves. These traits include leaf N and leaf mass per area (LMA; Asner et al.

2011; Serbin et al. 2014; Serbin et al. 2019; Streher et al. 2020), pigments (Asner et al. 2011; Yang et al.

2016), defense compounds (Couture et al. 2016; Nakaji et al. 2019), non-structural carbohydrates (Ely et

al. 2019), and even physiological rates like the maximum rate of RuBP carboxylation in photosynthesis

(V_{cmax}; Dechant et al. 2017; Meacham-Hensold et al. 2019; Yan et al. 2021). Leaf-level PLSR models

have been used to address such varied ecological topics as defense responses to herbivory (Kula et al.

2020), the role of biodiversity in ecosystem function (Schweiger et al. 2018), and functional trait

responses to El Niño (Nunes et al. 2019). Although this multivariate statistical approach is flexible, it is

sensitive to the kind of leaf tissue used to train the model. Existing PLSR models have mostly been

trained on spectra of fresh leaves (e.g. Serbin et al. 2019) or dried, ground leaves (e.g. Serbin et al. 2014).

Such models are not expected to transfer to dried, intact leaves like herbarium specimens because both

drying and grinding cause major changes in the reflectance spectrum.

We built PLSR models to estimate traits from the spectra of pressed leaves, like herbarium specimens,

and compared their accuracy to models built from fresh or ground leaves. Costa et al. (2018) showed that

models based on a related technique, Fourier Transform-Near Infrared Spectroscopy (FT-NIR), could

predict five leaf structural traits and a proxy for chlorophyll content from pressed leaves of tropical trees.

Here, we explicitly compare the accuracy of trait estimation from the spectra of fresh, pressed, and

ground leaves for a wide variety of leaf chemical and structural traits from multiple biomes and functional

groups.

For most chemical traits, such as elemental composition or carbon fractions, we conjectured that ground-

leaf spectral models would be the most accurate, as prior studies have found (Serbin et al. 2014; Couture

et al. 2016; Wang et al. 2020). Both drying and grinding might be important in achieving this high

accuracy. Drying may reveal minor absorption features of compounds in dry matter within the short-wave

infrared (SWIR) range that, in fresh leaves, are obscured by the dominant effect of water absorption

(Peterson et al. 1988; Couture et al. 2016; Wang et al. 2018). Grinding homogenizes the variation in

structure and composition throughout the leaf lamina, which may allow us to capture a more even and

representative sample of tissue (Richardson et al. 2021). Since pressed leaves are dried but not ground, we

predicted they would yield intermediate accuracy for chemical traits.

For structural traits like LMA, we instead expected fresh and pressed leaves to outperform ground leaves

because grinding disrupts the leaf structure. For water-related traits like leaf dry matter content (LDMC;

dried mass divided by fresh mass), we expected fresh leaves to outperform both pressed and ground

leaves because only they retain the water absorption features that allow direct, quantitative prediction of

water content (Carter 1991; Rapaport et al. 2015). Likewise, because pigments tend to degrade after

sample collection, we expected to estimate their concentration best from fresh leaves. We also assessed

sample discoloration in preparation for mounting or in storage and considered whether it reduces the

accuracy of trait estimates, which may indicate whether spectroscopy is useful on old or degraded

specimens.

Finally, we asked whether pressed-leaf spectra can be used to identify samples to species. Reflectance

spectra often show phylogenetic signal, at least in particular wavelength ranges, because of phylogenetic

conservatism in the evolution of their underlying traits (McManus et al. 2016; Diniz et al. 2020; Meireles

et al. 2020a). This phylogenetic signal is what often makes it possible to classify species or higher-level

taxa from fresh-leaf spectra (Cavender-Bares et al. 2016; Meireles et al. 2020b), but it may or may not be

retained in pressed-leaf spectra. Studies with tropical forest species have indeed demonstrated that FT-

NIR absorbance spectra of pressed herbarium specimens can be used to classify species or higher-level

taxa (Durgante et al. 2013; Lang et al. 2015; Lang et al. 2017; Prata et al. 2018). Based on these studies,

Draper et al. (2020) proposed using spectra of herbarium specimens as part of an integrative process of

species delimitation and identification. Here, we compare the accuracy of supervised classification from

fresh-, pressed-, and ground-leaf spectra among the common species in our dataset. Because they preserve

some of the spectral expression of leaf structure (unlike ground leaves) and reveal the distinctive SWIR

absorption features of macromolecules and other compounds (unlike fresh leaves), pressed leaves may

represent the best of both worlds for distinguishing species using spectroscopy.

Methods

Sampling

We trained PLSR models on leaf reflectance spectra and traits measured as part of four projects

conducted by the Canadian Airborne Biodiversity Observatory (CABO). The projects are:

1. Beauchamp-Rioux: Deciduous forest trees sampled throughout the growing season in 2018 at
   various sites in southern Québec and Ontario ($n = 405$; 9 species).
2. Dessain: Forbs, grasses, shrubs, and broadleaf trees from forests and open areas sampled
   throughout the growing season in 2017 at four sites in southern Québec ($n = 72$; 59 species).
3. Boucherville: Forbs, shrubs, and grasses sampled throughout the growing season in 2018 from
   the Parc national des Îles-de-Boucherville in southern Québec ($n = 73$; 12 species).
4. Warren: A single broadleaf evergreen tree species (*Agonis flexuosa* (Willd.) Sweet) collected
   across a soil chronosequence (Turner et al. 2018) in southwestern Australia ($n = 68$).

The majority of samples were from trees ($n = 504$; 25 species), but herbs ($n = 60$; 19 species), shrubs ($n =

48$; 23 species), and vines ($n = 6$; 1 species) were also represented. The full sampling procedure is

described in the Supplementary Materials. We validated the models both internally and on an independent

dataset of pressed tree and herb samples collected at Cedar Creek Ecosystem Science Reserve (East

Bethel, MN, USA).

Spectral and leaf trait measurements

For each CABO sample, we measured full-range reflectance spectra (350-2500 nm) of the leaves at three

stages: (1) freshly sampled, (2) pressed and dried, and (3) ground into a fine powder. We measured fresh-

leaf directional-hemispherical reflectance spectra on the adaxial surface of multiple leaves or (for small-

or narrow-leaved plants) leaf arrays from each sample using spectroradiometers equipped with integrating

spheres. We pressed a portion of the sample, and measured reflectance spectra on the adaxial surface of

pressed leaves between six months and three years later using a spectroradiometer with a leaf clip. Lastly,

we measured ground-leaf spectra using a spectroradiometer with a benchtop reflectance probe that

pressed loose leaf powder into a smooth, even pellet that we assume has very low transmittance. We

trimmed all spectra to 400-2400 nm. Detailed notes on measurement and post-measurement processing of

reflectance spectra are found in the Supplementary Materials.

While measuring pressed-leaf spectra, we inspected each pressed specimen by eye to note signs of

discoloration in preparation or storage. Typical forms of discoloration include blackening, browning, or

the development of a silvery or whitish finish on the leaf surface. We scored each leaf on a discrete scale

from 0 to 4. A score of 0 indicates no noticeable discoloration. Scores 1 through 4 indicate increasing

discoloration, from 1 (either <5% blackening/browning or development of a silvery finish to the leaf) to 4

(>80% blackening/browning).

We measured the following leaf structural and chemical traits on all CABO samples: Leaf mass per area

(LMA; kg m^{-2}), leaf dry matter content (LDMC; mg g^{-1}), equivalent water thickness (EWT; cm), carbon

fractions (soluble cell contents, hemicellulose, cellulose, and lignin; %), pigments (chlorophyll *a*,

chlorophyll *b*, and total carotenoids; mg g^{-1}), and concentrations of a variety of elements (Al, C, Ca, Cu,

Fe, K, Mg, Mn, N, Na, P, Zn; % or mg g^{-1}). Protocol summaries are in the Supplementary Materials. We

kept chemical traits on a mass basis for consistency with the usual basis on which they are measured and

reported (pigments excepted). However, some traits are most often distributed proportionally to area

(Osnas et al. 2013), and retrievals on an area basis may be better for some purposes (Kattenborn et al.

2019).

PLSR modeling for trait estimation

We used a PLSR modeling framework to link the fresh-, pressed-, and ground-leaf spectra to the traits we

measured. PLSR is well-suited to handle leaf spectral datasets, which have large numbers of collinear

predictors, because it projects the spectral matrix onto a smaller number of orthogonal latent components

in a way that maximizes the ability to predict the response variable. We implemented PLSR modeling

using *R* package *pls v. 2.7.1* (Mevik et al. 2019). We conducted analyses separately for fresh-, pressed-,

and ground-leaf spectra, using the full spectrum (400-2400 nm) to predict each trait. We simply used

reflectance values as predictors, since calculating various transformations of reflectance (first- or second-

order differences, or pseudoabsorbance defined as –log[reflectance]) made little difference for predictive

accuracy in preliminary tests. We also did not perform any transformations to reduce the skew of trait

distributions, again finding in preliminary tests that they did not improve predictive accuracy enough to

warrant the added complexity. Some studies restrict the wavelengths used in prediction—for example, to

ranges known or assumed to contain features relevant to a given trait (e.g. Serbin et al. 2014). Given the

potential for photosynthetic pigments to degrade and produce brown pigments in pressed leaves (Fourty et

al. 1996), we also built pressed-leaf models restricted to the range 1300-2400 nm, which might be

expected to generalize better to datasets that include more degraded leaves. We present these results

mainly in the Supplementary Materials.

Our methods for model calibration and validation largely follow Serbin et al. (2014) and Burnett et al.

(2021). First, we randomly divided the data into calibration (75%) and validation (25%) datasets,

stratified by project. We began by fitting a model for each trait on the calibration dataset. We selected the

smallest number of components for which the root mean squared error of prediction (RMSEP) from 10-

fold cross-validation was within one standard deviation of the global minimum. We used this number of

components—a different number for each trait—in further analyses to predict traits on the internal

validation dataset. We calculated the variable importance in projection (VIP) metric for calibration

models to see which parts of the spectrum were most important for predicting each trait (Wold et al.

1994).

To test how well we could predict traits on the internal validation subset, we did a jackknife analysis by

repeatedly (100×) dividing the 75% calibration data further into random 70% training and 30% testing

subsets. For each trait, we trained models on the 70% using the previously determined optimal number of

components and predicted the remaining 30%, which gave us a distribution of model performance

statistics across iterates (R^2, %RMSE). This distribution reveals how sensitive model performance is to

randomly varying sets of training and testing data.

Next, we applied the 100 jackknife models for each trait to the 25% internal validation dataset: for each

sample in the validation dataset, we generated a distribution of predictions from the 100 models. We

quantified model performance using R^2 and root mean squared error (RMSE) between measurements and

mean predictions. We also report the RMSE as a percentage of the 2.5% trimmed range (%RMSE), which

we used rather than the full range (as in e.g. Burnett et al. 2021) for robustness to outliers. For each trait,

we also tested whether the magnitude of residuals (observed minus predicted) in the validation dataset

varied among leaves with different discoloration scores. We performed all statistical analyses in *R v. 3.6.3*

(R Core Team 2020).

Finally, to test how well our model would transfer to a fully independent dataset, we applied the ensemble

of pressed-leaf models for five traits (LMA, LDMC, EWT, N, and C) to pressed-leaf spectra from Cedar

Creek and compared the model-derived trait estimates to measured values. Like most of the CABO

dataset, the Cedar Creek dataset comprises trees and herbs from northeastern temperate North America,

but it includes an entire functional group (needleleaf conifers) absent among the CABO projects in this

study. We collected the spectra with the same instrument and foreoptic as the pressed-leaf spectra in the

CABO dataset, but used different sampling, preparation, and trait measurement protocols. Full details on

this dataset are provided in Supplementary Materials.

PLS-DA modeling for species classification

We tested the potential to classify species with fresh-, pressed-, and ground-leaf spectra using partial

least-squares discriminant analysis (PLS-DA; Barker & Rayens 2003). We took spectra from the ten most

common species in our dataset—all of which are deciduous trees except *A. flexuosa*, which is evergreen.

Each species was represented by at least 20 specimens (~480 total). For each tissue type, we divided the

full dataset into 60% calibration and 40% validation subsets. In the *R* library *caret v. 6.0.84* (Kuhn 2020),

we trained models on the calibration subset using 10-fold cross-validation. We chose the number of PLS

components during cross-validation by maximizing Cohen's kappa (κ), which describes the agreement

between the true and predicted species identities while accounting for the probability of agreement by

chance. Imbalanced training data can bias classification algorithms (Sun et al. 2009; Lindström et al.

2011), so we used a two-step procedure to balance classes while maintaining enough training data and

avoiding overfitting. First, we downsampled within better-represented species classes at random so that

all classes had equal size, then chose the number of components (*n*) that maximized κ. Second, we

upsampled from less-represented classes at random with replacement so that classes had equal size—

again maximizing κ, but restricting the range of components evaluated to no more than the *n* chosen

during the downsampling step. We applied the cross-validated PLS-DA model from the upsampling step

to the validation subset and summarized its performance using raw classification accuracy and κ.

Results

Patterns in traits and reflectance spectra

We saw large variation among samples in each of our target traits, ranging from 1.4-fold variation in C to

more than 20-fold variation in traits like lignin, P, K, and Zn (Table 1). The ranges of most traits in our

dataset covered a large portion of the global distributions in the TRY dataset, but tended to be narrower at

both extremes (Kattge et al. 2020). Trees tended to have higher LDMC, C, and lignin than other growth

forms. Among the herbs, grasses had particularly high hemicellulose and cellulose and low lignin content,

while forbs generally had high N. Some of the trait variation was driven by projects that occupied

particular portions of trait space; for example, *A. flexuosa* in the Warren project tended to have

particularly high LMA, Na, and C and low N.

Both pressed and ground leaves had higher median reflectance across nearly the entire spectrum (Fig. 1),

as expected based on changes in water content and structure (Carter 1991). Indeed, water absorption

features (the largest of which are centered around 1450 and 1930 nm) largely disappeared in pressed and

ground leaves. The red edge between the visible and near-infrared (NIR) regions was also blunted by both

pressing and grinding, causing the global maximum of median reflectance to shift from 872 nm (fresh) to

954 nm (pressed) to 1313 nm (ground). The coefficient of variation (CV) of reflectance was generally

highest where average reflectance was lowest.

PLSR modeling for trait estimation

Tested on the internal validation dataset, pressed-leaf models performed best at predicting LMA (R^2 =

0.917; %RMSE = 7.30), C (R^2 = 0.774; %RMSE = 11.8), and N (R^2 = 0.772; %RMSE = 11.5). These

traits were followed by a mixture of water-related traits, carbon fractions, and pigments, as well as Ca (all

R^2 = 0.554-0.763; %RMSE = 12.3-18.6). Although some other elements (Na, P) could be estimated with

R^2 > 0.5 and %RMSE < 20, most models for other elements showed lower accuracy (Table 1; Figs. 2-3

and S1-7). In most cases, the models restricted to 1300-2500 nm had slightly lower predictive accuracy

than the full range models (Table S1).

We compared pressed-leaf models to both fresh-leaf and ground-leaf models. The optimal number of

components selected to predict each trait was between 3 and 23. For any given trait, ground-leaf models

always had the most components, usually followed by pressed-leaf models (Table 1). Fresh-leaf models

were best for predicting the structural and water-related traits—LMA, LDMC, and EWT (Figs. 2-3 and

S1-7). On the other hand, ground-leaf models were best for predicting chemical traits, like carbon

fractions and most elements. Fresh- and ground-leaf models had similar performance for pigments.

Statistics from jackknife analyses showed that model performance was more variable for traits that were

predicted less accurately (Figs. S11-S13). There was no correlation between the magnitude of residuals

and our discoloration index for any trait ($p > 0.05$).

For all traits except LMA and Fe, the VIP metric for fresh-leaf spectra showed a global maximum

between 710 and 720 nm (Fig. 4 and S8-10)—wavelengths slightly longer than the typical inflection point

of the red edge (Gitelson et al. 1996). Many traits also show high VIP across the wavelengths ~530-570

nm, corresponding to the green hump. Bands in the NIR range were less important for predicting most

traits than much of the visible range. The SWIR range was generally important for predicting LMA,

EWT, Na, and pigments, and many other traits showed several local peaks of importance, most

prominently at about 1880 nm, but also near 1480 and 1720 nm.

For predicting traits from pressed-leaf spectra, the general trend held that visible reflectance and certain

ranges in the SWIR were important for predicting most traits, while the NIR and much of the shorter

SWIR (800-1750 nm) were less important (Fig. 4 and S8-10). The red edge peak of importance for most

traits was near 705 nm. Other prominent local maxima for many traits lay close to 1440, 1720, 1920,

2130, and 2300 nm. We saw broadly similar patterns in ground-leaf spectra, except that VIP for most

traits was lower at longer SWIR wavelengths (2000-2400 nm).

External validation

For most traits, pressed-leaf model performance on the external validation dataset from Cedar Creek was

comparable to, but not quite as strong as, the internal validation (Table 2; Fig. 5). The exception was C,

for which the models performed very poorly ($R^2 < 0.05$). Among the remaining traits, R^2 ranged from

0.557 (N) to 0.874 (LMA). For N in particular, %RMSE was high (37.8%) due to bias—N concentrations

were underestimated for conifers but slightly overestimated for remaining samples.

Since conifers were absent from the CABO training dataset, we considered whether the models we built

would extend to this new functional group. For N, models performed better when excluding conifers ($R^2 =$

0.594, %RMSE = 37.8) than when retaining them ($R^2 = 0.557$; %RMSE = 29.5). In contrast, for LMA,

models performed worse when excluding conifers ($R^2 = 0.762$; %RMSE = 15.7) than when retaining them

($R^2 = 0.874$; %RMSE = 10.3), since their LMA values could be predicted well and extended the trait

range. Models restricted to 1300-2500 nm did not consistently outperform those trained on the full

range—they yielded somewhat better external validation R^2 for N, but worse for LDMC, both including

and excluding conifers (Table S2; Fig. S14).

PLS-DA modeling for species classification

PLS-DA models using pressed- and ground-leaf spectra showed near-perfect performance at classifying

species (Fig. 6). Models using fresh-leaf spectra were slightly worse but still showed strong performance.

The optimal fresh-leaf model, which had 29 PLS components, correctly predicted the taxonomic identity

of 170 of the 189 samples in the training dataset ($\kappa = 0.884$; $p < 0.0001$). The best pressed-leaf model,

which had 29 PLS components, correctly predicted 184 of the 188 samples ($\kappa = 0.975$; $p < 0.0001$). The

best ground-leaf model, which had 30 PLS components, correctly predicted 183 out of 189 samples ($\kappa =$

0.963; $p < 0.0001$). In all cases, the majority (75-100%) of misclassifications were between congenerics.

Discussion

We show that we can estimate a wide range of leaf functional traits for 68 woody and herbaceous species

from reflectance spectra of pressed leaves (Table 1; Figs. 2-3 and S1-7). Model performance was highest

for LMA, C, and N, followed by a mixture of water-related traits, carbon fractions, pigments, and a few

important elements (Ca, Na, and P). Finally, other elements could only be estimated with fairly low

accuracy. These results show that pressed-leaf spectra provide an integrative measure of leaf phenotypes,

much like fresh-leaf spectra (Cavender-Bares et al. 2017), but with even stronger potential to characterize

variation in chemical traits. Perhaps as a result, we could use pressed-leaf spectra to classify species as

accurately as ground-leaf spectra, and better than fresh-leaf spectra. Our results point to the potential that

using reflectance spectroscopy on herbarium specimens could yield rapid and non-destructive estimates of

many functional traits, enabling more expansive studies of trait variation across space and time.

We compared pressed-leaf models to fresh-leaf and ground-leaf models from the same samples. Our

findings about which kind of tissue was best for predicting each trait mostly supported our hypotheses.

Ground-leaf spectra showed the strongest performance for chemical traits, likely because grinding

homogenizes the lamina and removes the potentially confounding influence of leaf structure (Table 1).

Pressed-leaf spectra showed intermediate performance for most chemical traits, perhaps because, like

ground leaves, they lack the major water absorption features that mask the smaller features of other

compounds in the SWIR range (Peterson et al. 1988). Contrary to our predictions, ground-leaf spectra

performed about as well as fresh-leaf spectra, and better than pressed-leaf spectra, for estimating pigment

concentrations. The same factors that provided an advantage to ground-leaf spectra in estimating chemical

traits—removal of water absorption and leaf structure—also perhaps explain why pressed- and (especially

for LMA) ground-leaf spectra performed worse for estimating water-related and structural traits (LMA,

LDMC, and EWT). Pressed-leaf models may represent a good compromise in allowing many traits to be

estimated with mostly intermediate but nonetheless quite high accuracy.

It may seem perplexing that we could succeed at all in predicting LMA from ground-leaf spectra, or

LDMC and EWT from pressed- and ground-leaf spectra. Many nutrients also have weak or no absorption

features. The ability to estimate these traits must not result from the optical expression of the traits

themselves. We suggest that we instead sense these traits via their correlations with other traits that have a

stronger optical expression. This kind of effect—a "constellation effect" (*sensu* Chadwick & Asner 2016;

Nunes et al. 2017)—has been invoked to explain the ability to estimate traits like leaf nutrients (Nunes et

al. 2017) and $\delta^{15}N$ (Serbin et al. 2014) that are not known to have strong absorption features in the

measured range of wavelengths. For example, some nutrients are sensed mainly through symptoms of

deficiency (Jacquemoud & Ustin 2020). However, models that rely on such constellation effects could

break down when patterns of trait covariance vary, which could make models for certain traits fail beyond

the domain of the training data.

The performance of pressed-leaf models in this study was comparable to the successes obtained here and

elsewhere using fresh- and ground-leaf spectra. For example, the ground-leaf models in Serbin et al.

(2014) had a validation RMSE of 1.4 and 2.4 for cellulose and lignin percentages. In comparison, our

pressed-leaf models achieved an RMSE of 1.52 and 2.30 (Table 1). Our models for LMA also had an

RMSE (0.0115 kg m^{-2}) lower than fresh-leaf models from the literature, including Serbin et al. (2019;

0.015 kg m^{-2}), Nakaji et al. (2019; 0.015 kg m^{-2}), and Streher et al. (2020; 0.051 kg m^{-2}). On the other

hand, our pressed-leaf model for N performed worse than Serbin et al. (2014; validation RMSE = 0.13 vs

0.273). Although these studies sampled different kinds of vegetation with different trait ranges, such

comparisons establish that pressed-leaf models are often comparable to ground- or fresh-leaf models. Our

models' error could be within acceptable bounds for addressing a wide range of questions about large-

scale ecological or evolutionary patterns that encompass a wide range of trait variation. Moreover, our

external validation analyses indicate that our models for some important traits—like LMA, LDMC, EWT,

and N, but not C—can transfer reliably to other datasets.

The VIP metric helps us understand what features allow these models to work, but interpreting it can be

challenging because reflectance at a given band usually cannot be understood in terms of a single trait.

Fresh-leaf VIP for most traits had peaks in the visible and SWIR ranges, with a global maximum at 705

nm (Fig. 4 and S8-10). The pattern of high VIP along the green hump and at the red edge is common in

PLSR models from fresh leaves (Ely et al. 2019; Streher et al. 2020; Yan et al. 2021). The red edge

(particularly 700-725 nm) may be so important because of its sensitivity to both chlorophyll content and

leaf structure (Gitelson et al. 1996; Richardson et al. 2002). Much of the visible range was proportionally

even more important for pressed- and ground-leaf models, and most of the NIR was less important

(except for LMA; Fig. 4 and S8-10). There were multiple small peaks in the SWIR, most notably one at

1920 nm that appears similar to a VIP feature in Serbin et al. (2014). Reflectance at 1920 nm is often

associated with water absorption in fresh tissue, but any causal link to the leaf's fresh water content is

unlikely for pressed and ground leaves. Bands near 1920 nm also include broad absorption features for

many components of dry matter, including protein, cellulose, lignin, and starch, any or all of which may

contribute to the importance of this region (Curran et al. 1989; Fourty et al. 1996).

With some exceptions, the VIP metric showed that the same bands are often important for predicting

different traits. This pattern might be taken as an artifact of trait covariance: For example, the three

pigment pools covaried strongly ($R^2 = 0.827$-0.969) and had nearly identical VIP across the spectrum

(Figs. S8-S10). In some cases, similarities in VIP might be taken further to imply that there are a small

number of traits whose tight coordination with others underlies the performance of all models through

constellation effects. Nevertheless, across the whole dataset, many traits covaried only weakly but still

shared important VIP features. For example, EWT, cellulose, N, and K were not tightly coordinated ($R^2 =$

0.003-0.152) but shared similar patterns of pressed-leaf VIP across the spectrum (Fig. 4), including major

peaks at 705 and 1920 nm.

PLS-DA models showed that fresh-, pressed-, and ground-leaf spectra alike could be used to classify

species with near-perfect accuracy (>96%) for pressed or ground leaves, and excellent (>89%) accuracy

for fresh leaves (Fig. 6), much as in earlier work on tropical flora (e.g. Durgante et al. 2013). In contrast

to prior work that deliberately selected many congenerics, our most common species were often distantly

related. Among the misidentified samples, most were mistaken for congenerics, which implies that related

species are more spectrally similar (Schweiger et al. 2018; Meireles et al 2020a). Hence, if we had

included more closely related species, we might have seen a higher error rate. Nevertheless, our analysis

reinforces that spectral models can often distinguish species or higher-level phylogenetic groups

(Meireles et al. 2020b).

As we predicted, pressed-leaf models showed the highest classification accuracy (>97%). This result is

noteworthy, since pressed specimens are the standard way that field botanists preserve plant samples for

later identification, but the outcome may be surprising because pressed leaves were never best for

predicting any single trait. We conjecture that pressed- and ground-leaf models have an advantage

because drying reveals the absorption features of potentially distinctive compounds in the SWIR range.

While water content can only vary in one dimension, there are many macromolecules and metabolites

whose absorption features in this range might vary independently and allow finer discrimination of

species.

The future of spectroscopic trait estimation

Although trait predictions from spectral models are not perfect, they have a few advantages over

conventional trait measurements: they (1) can be non-destructive; (2) are fast and require relatively little

training; and (3) have very low marginal cost, despite the high capital cost of buying a spectrometer

(Costa et al. 2018). These advantages could make it easier to address questions that require large datasets

of functional traits. But researchers may be deterred if they must each build their own models tailored to

particular uses—and for herbarium specimens, it might not be possible to do the conventional trait

measurements needed to train the models. Ideally, spectral models would be general enough that

researchers could confidently use them without further validation, but this aim is not easy to achieve: for

several reasons, a model trained on any particular spectral dataset may make poor trait predictions on new

data. As with any other technique, the goal for spectroscopic trait estimation is to improve model

accuracy and generality as much as they can be jointly improved without making the technique harder to

use. Below, we discuss some challenges one by one, particularly as they concern herbarium specimens.

One concern is that the new data could be outside the range of traits or optical properties in the training

dataset (Schweiger 2020). A fully general model, if such a thing is possible, would need to represent the

full range of leaf functional traits and optical properties. Another kind of concern about model generality

concerns sample preparation before spectral measurements. For example, particle size influences ground-

leaf spectra (Foley et al. 1998). For pressed leaves, it may be particularly important to prepare samples in

consistent ways that preserve the leaves' morphological integrity. In our external validation analyses, we

found that pressed-leaf models yielded reasonably accurate predictions of most traits, even though the

validation dataset differed in sample preparation protocols and included conifers, which were absent from

the training dataset. Nevertheless, even setting aside conifers, external validation for one trait (C) was

very poor, and for another (N) showed noticeable bias, such that researchers might need to develop their

own correction factors to use the model in practice.

Another class of challenges concerns spectrometers and their foreoptics. Spectra of fresh leaves can be

measured with different foreoptics, including integrating spheres, contact probes, or leaf clips. In this

study, we used a leaf clip with pressed specimens because mounting delicate pressed leaves in an

integrating sphere could damage them. However, leaf clips and contact probes are less likely to produce

consistent measurements among instruments or replicate samples due to variation in viewing geometry

and anisotropic surface reflectance (Petibon et al. 2021). The logistical constraint of having to use a leaf

clip or contact probe could thus make it harder to compare across datasets. Moreover, measuring

reflectance with a leaf clip or probe usually requires placing a black absorbing background under the

sample to keep transmitted light from being reflected back into the sensor, but many herbarium specimens

are glued to a paper backing. When loose leaf material is not available, using spectra from these

specimens may require new methods to correct for reflectance from the mounting paper.

Spectrometers and their software may also differ subtly in their sensors and techniques for processing

spectra (Castro-Esau et al. 2006), and in some cases, researchers must take steps to reconcile spectra

measured from different instruments (Meireles et al. 2020b). Assuming that these kinds of technical

challenges can be overcome, our results and others encourage confidence about building general models

to estimate traits from a wide variety of plants (Serbin et al. 2019). The creation of open libraries for

spectral data (like EcoSIS, https://ecosis.org/; or the CABO data portal, https://data.caboscience.org/leaf)

and spectral models (like EcoSML; https://ecosml.org/) will contribute to this goal.

Even though species classification models are more likely to remain domain-specific, some of the same

concerns apply. For example, a training dataset for a species classification model must include strong

representation for all the species that one might encounter. Researchers may also want to use existing

data, as from open libraries, to train models, which would require resolving concerns about sampling—

including phenotypic plasticity—as well as the potential influence of instrumentation and measurement

protocols (Meireles et al. 2020b; Draper et al. 2020).

What should herbarium managers do?

Perhaps the most concerning challenge for interpreting spectra from herbarium specimens is that

specimens' optical and chemical properties may degrade during preparation or storage, especially light-

sensitive pigments. Such degradation could make it hard to distinguish changes in the traits of living

plants over time from changes in storage. Even in this study, where no specimens were collected before

2017, many underwent visible changes in color, including browning or blackening; ~12% were scored at

2 or higher. Certain preparation and storage conditions could accelerate changes in optical properties. For

example, when collecting herbarium specimens in the wet tropics, researchers often treat leaves with an

alcohol solution as a preservative (Blanco et al. 2006). Such treatments could discolor leaves—but so

could the spoilage that would occur in their absence. Exposure to high light, temperature, or humidity

could also hasten discoloration, which underscores the importance of using best practices for preparation

and storage. Some species may also be more susceptible than others; in our dataset, ~42% of *Populus*

grandidentata specimens were scored at 2 or higher, but no *Betula papyrifera* specimens were.

Whether such degradation could affect trait estimation is an open empirical question. We find no evidence

that it does, even for badly discolored specimens, but our specimens were only collected as early as 2017,

while ecologists may want to use specimens collected decades ago. Long-term studies of the same

specimens—especially with sets of specimens subjected to varying preparation and storage techniques—

would help establish how their optical properties tend to change over time. In any case, it seems likely

that estimating traits in old or degraded specimens will require including equally old or degraded

specimens in the dataset used to build the model.

Some of the challenges we describe pertain to projects that would seek to measure spectra on samples

already collected, but spectroscopy, among countless other novel uses for herbarium specimens

(Heberling & Isaac 2017), could also prompt changes in collection practice. The best way to incorporate

spectroscopy into the operation of a working herbarium may be to measure incoming specimens as they

are pressed, but before they are mounted, which could also mitigate the effects of degradation. Collecting

and storing extra leaf material would also circumvent the potential challenge of measuring mounted

leaves and aid other novel uses of herbarium specimens (Heberling 2021).

Collections-based research has been accelerated by advances in biodiversity informatics that make it

easier to gather and synthesize data (Heberling et al. 2021). If it becomes common practice to measure

spectra on herbarium specimens—either at the time of collection (fresh) or after (pressed)—new tools

may be needed to link spectral data to specimen metadata and other phenotypic or genetic information

(Draper et al. 2020). The hyperdimensionality of the reflectance spectrum could make it hard to

accommodate within existing standards for biodiversity informatics such as the Darwin Core (Wieczorek

et al. 2012), at least without extensions. One could link records to external databases like EcoSIS or

SPECCHIO (Hueni et al. 2020), which are also equipped to store spectral metadata about instrumentation

and processing. Coordination between herbarium managers and researchers who use reflectance

spectroscopy could advance protocol development and build agreement about best practices for

measurement and curation.

We show that non-destructively measured pressed-leaf spectra retain much of the information about many

leaf functional traits found in fresh-leaf spectra. While validating this technique on older specimens will

require extensive further research, our findings suggest that leveraging reflectance spectroscopy could

allow herbaria to take on a much greater role in plant functional ecology and evolution. Our study has far-

reaching implications for capturing the wide range of functional and phenotypic information in the

world's preserved plant collections.

Acknowledgements

We performed this research on the ancestral and contemporary land—mostly unceded—of Native people: primarily the Kanien'kehá꞉ka (Mohawk), Omàmiwinini (Algonquin), and Abenaki First Nations for the Beauchamp-Rioux, Dessain, and Boucherville projects, the Noongar peoples for the Warren project, and the Dakota and Ojibwe peoples for the Cedar Creek dataset. For permissions and assistance, we thank the Jardin botanique de Montréal; la Station de biologie des Laurentides (SBL) de l'Université de Montréal; l'Institut de recherche d'Hydro-Québec (IREQ); the National Capital Commission for sampling at Mer Bleue Bog; la Société des établissements de plein air du Québec (SÉPAQ) for sampling at Parc national d'Oka, Parc national du Mont-Saint-Bruno, and Parc national des Îles-des-Boucherville; and the Parks and Wildlife Service of Western Australia for sampling in D'Entrecasteaux National Park. We thank Antoine Mathieu, Alexandra Massey, Florence Blanchard, Elisabeth Hardy-Lachance, Zachary Bélisle, Sandra Jooty, Myriam Cloutier, Fabien Cichonski, Madeleine Trickey-Massé, Vincent Fournier, Aurélie Dessain, Xavier Guilbeault-Mayers, Jocelyne Ayotte, Megan Erding, and especially Sabrina Demers-Thibeault for contributing to trait measurements. Carole Sinou, Phil Townsend, Dudu Meireles, Mason Heberling, and members of the Cavender-Bares lab and the Laboratory of Plant Functional Ecology (LEFO) contributed to insightful discussions. Data from Canadian and Australian sites were collected as part of the Canadian Airborne Biodiversity Observatory (CABO) funded by NSERC Discovery Frontiers grant 509190-2017, as well as NSERC Discovery grants (RGPIN-2014-06106, RGPIN-2019-04537) and a Canada Research Chair to EL. Cedar Creek collections were supported by the NSF/NASA (DEB-1342778) and the NSF ASCEND Biology Integration Institute (DBI-2021898). Cedar Creek is supported by the NSF Long-Term Ecological Research program, currently under DEB-1831944. Travel funding was provided by an International Thesis Research Travel Grant from the University of Minnesota's Graduate School and a grant distributed through an NSF-funded Research Coordination Network (DEB-1745562). SK was supported by an NSF Graduate Research Fellowship (Grant No. 00039202) and a UMN Doctoral Dissertation Fellowship. RBR was supported by NSERC through the Canada Graduate Scholarships—Master's program and the FRQNT through a Master's Research Scholarship.

Conflict of interest statement

The authors have no conflicts of interest to declare.

Author contributions

SK, JCB, and EL conceived the ideas and designed the methodology. RBR and SK collected the data. SK

analyzed the data and wrote the first draft of the manuscript. All authors made substantial contributions to

further drafts and gave final approval for publication.

Tables

Table 1: Summary statistics of PLSR model internal validations from fresh-, pressed-, and ground-leaf spectra. %RMSE is calculated as RMSE divided by the

2.5% trimmed range of data within either the training data set (calibration) or the testing data set (validation). Global ranges are derived from the TRY database

(Kattge et al. 2020; accessed 13 April 2021), omitting data with error risk greater than 3. Carbon fractions (solubles, hemicellulose, cellulose, and lignin) and

carotenoids are each represented by <1250 datapoints in TRY, which may limit the estimates of the global range. EWT is not available on TRY.

Trait	Dataset range	Global range	Fresh No. comps	R^2	RMSE	%RMSE	Pressed No. comps	R^2	RMSE	%RMSE	Ground No. comps	R^2	RMSE	%RMSE
LMA (kg m^{-2})	0.0233-0.217	0.00866-0.354	15	0.960	0.00795	5.05	9	0.917	0.0115	7.30	20	0.824	0.0165	10.5
LDMC (mg g^{-1})	155-573	68.2-857	16	0.853	26.0	8.82	17	0.713	36.4	12.3	20	0.764	33.1	11.2
EWT (cm)	0.00488-0.0282	NA	7	0.871	0.00167	9.99	14	0.763	0.00213	12.8	22	0.691	0.00242	14.5
Solubles (%)	38.5-88.9	20.0-74.6	15	0.438	6.12	19.5	12	0.701	4.31	13.6	23	0.814	3.36	10.6
Hemicellulose (%)	2.77-34.1	8.71-45.0	11	0.483	3.47	22.0	12	0.731	2.48	15.6	20	0.755	2.35	14.6
Cellulose (%)	5.19-25.8	7.65-42.6	20	0.482	2.29	27.4	15	0.735	1.52	18.1	21	0.872	1.04	12.4
Lignin (%)	0.918-21.5	2.27-32.4	10	0.497	2.42	19.6	12	0.554	2.30	18.6	20	0.777	1.65	13.3
C (%)	39.5-53.6	25.5-60.3	17	0.716	1.22	13.4	14	0.774	1.08	11.8	20	0.908	0.689	7.53
N (%)	0.883-5.62	0.14-7.27	10	0.729	0.301	12.6	15	0.772	0.273	11.5	22	0.952	0.127	5.31
Chl a (mg g^{-1})	1.24-15.5	1.38-22.1 (total Chl)	5	0.719	1.12	14.1	7	0.626	1.30	16.4	9	0.719	1.14	14.4
Chl b (mg g^{-1})	0.486-5.29		5	0.696	0.397	14.7	6	0.555	0.484	17.9	8	0.688	0.410	15.2
Carotenoids (mg g^{-1})	0.193-3.05	0.00102-5.1	5	0.638	0.264	16.9	7	0.615	0.274	17.5	16	0.712	0.240	15.3
Al (mg g^{-1})	0-0.379	0.004-19.7	5	0.220	0.0415	24.5	6	0.243	0.0409	24.1	10	0.593	0.0304	17.9
Ca (mg g^{-1})	1.64-36.9	0.68-69	11	0.425	5.34	19.0	17	0.744	3.57	12.7	23	0.868	2.58	9.16
Cu (mg g^{-1})	0-0.052	0.00154-0.064	8	0.472	0.00577	20.6	10	0.301	0.00653	23.2	7	0.340	0.00641	23.1
Fe (mg g^{-1})	0.0150-0.266	0.0128-1.1	4	0.267	0.0380	20.4	7	0.215	0.0392	21.0	9	0.358	0.0356	19.1
K (mg g^{-1})	1.06-38.2	0.3-78.1	7	0.598	2.92	17.6	14	0.488	3.18	20.1	17	0.694	2.55	15.3
Mg (mg g^{-1})	0.482-7.86	0.277-15.6	7	0.253	0.802	22.1	13	0.373	0.759	20.9	23	0.529	0.656	18.0
Mn (mg g^{-1})	0-1.02	0.00560-5.37	3	0.198	0.158	25.3	12	0.349	0.144	22.9	17	0.655	0.107	18.1
Na (mg g^{-1})	0-4.87	0.00499-48.7	4	0.360	0.612	21.9	9	0.514	0.537	19.2	14	0.426	0.579	20.7

P (mg g⁻¹)	0.222-7.18	0.1-9.9	7	0.389	0.866	21.3	19	0.547	0.738	18.1	13	0.598	0.699	17.2
Zn (mg g⁻¹)	0.00142-0.647	0.00221-0.242	9	0.250	0.0744	26.2	8	0.310	0.0704	24.8	15	0.437	0.0634	22.3

585 Table 2: Summary statistics for external validation of pressed-leaf PLSR models. The models were trained on CABO data (see Table 1) and applied to a dataset

586 collected at Cedar Creek.

	Including conifers				Excluding conifers			
Trait	Range	R^2	RMSE	%RMSE	Range	R^2	RMSE	%RMSE
LMA (kg m⁻²)	0.0179-0.229	0.874	0.0186	10.3	0.0179-0.136	0.762	0.0126	15.7
LDMC (mg g⁻¹)	133-525	0.559	47.8	16.9	133-504	0.571	49.7	17.2
EWT (cm)	0.00602-0.0372	0.609	0.00449	18.1	0.00602-0.0363	0.567	0.00354	19.6
C (%)	35.7-59.8	0.033	4.42	25.2	35.7-54.1	0.036	4.21	27.1
N (%)	0.75-4.68	0.557	0.725	37.8	0.75-4.68	0.594	0.754	29.5

Figure captions

Fig. 1: Distributions of spectral reflectance and its coefficient of variation (CV) among fresh (A), pressed (B), and ground (C) leaf samples. A solid black line connects the median reflectance at each wavelength, around which dark blue and light blue ribbons denote the middle 50% and 95%. The solid red line shows the coefficient of variation among samples at each wavelength.

Fig. 2: Internal validation results for predictions of LMA, EWT, cellulose, and chlorophyll *a* from fresh-(left), pressed- (middle), and ground-leaf (right) spectra. In each panel, a separate ordinary least-squares regression line is shown for each functional group, overlaid on top of the thick dashed 1:1 line. The error bars for each data point are 95% confidence intervals calculated from the distribution of predictions based on the model coefficients from the 100 jackknife iterations. Figs. 2 and 3 display a subset of traits that represent various functions and span the range of model performance. Plots for remaining traits are shown in Supplementary Materials.

Fig. 3: Internal validation results for predictions of three elements (N, K, and Mn) from fresh- (left), pressed- (middle), and ground-leaf (right) spectra, displayed as in Fig. 2.

Fig. 4: The variable importance of projection (VIP) metric calculated based on fresh- (left), pressed-(middle), and ground-leaf (right) models. As in Figs. 2 and 3, we selected seven traits that represent a range of different patterns in VIP across the tissue types (cell = cellulose). The dashed horizontal line at 0.8 represents the heuristic threshold for importance recommended by Burnett et al. (2021). Plots for traits not shown here are found in Supplementary Materials.

Fig. 5: External validation results for predictions of various traits in Cedar Creek data based on models trained on CABO data. Panels are displayed as in Figs. 2 and 3.

Fig. 6: Confusion matrices for partial least-squares discriminant analysis from fresh- (A), pressed- (B),

and ground-leaf (C) spectra. The row specifies the true species identity, while the column specifies the

model's prediction. Numbers on the diagonal represent correctly classified species, while off-diagonal

numbers represent misclassified species. Full binomials: *Agonis flexuosa*, *Acer rubrum* L., *A.*

saccharinum L., *A. saccharum* Marshall, *Betula papyrifera* Marshall, *B. populifolia* Marshall, *Fagus*

grandifolia Ehrh., *Populus grandidentata* Michx., *Populus tremuloides* Michx., and *Quercus rubra* L.

Figures

Fig. 1

Fig. 2

Fresh-leaf spectra · Pressed-leaf spectra · Ground-leaf spectra

Growth form — herb — shrub — tree —

627 **Fig. 3**

636 **Fig. 4**

655 **Fig. 5**

Fig. 6

A

Fresh−leaf spectra

B

Pressed−leaf spectra

C

Ground−leaf spectra

References

American Society for Testing and Materials. (2006). *Standard Tables for Reference Solar Spectral Irradiances: Direct Normal and Hemispherical on 37° Tilted Surface* (Standard G173-06).

Asner, G. P., Martin, R. E., Knapp, D. E., Tupayachi, R., Anderson, C., Carranza, L., Martinez, P., Houcheime, M., Sinca, F., & Weiss, P. (2011). Spectroscopy of canopy chemicals in humid tropical forests. *Remote Sensing of Environment, 115*(12), 3587–3598.

Barker, M., & Rayens, W. (2003). Partial least squares for discrimination. *Journal of Chemometrics*, 17(3), 166–173.

Blanco, M. A., Whitten, W. M., Penneys, D. S., Williams, N. H., Neubig, K. M., & Endara, L. (2006). A Simple and Safe Method for Rapid Drying of Plant Specimens Using Forced-Air Space Heaters. *Selbyana*, *27*(1), 83–87.

Burnett, A. C., Anderson, J., Davidson, K. J., Ely, K. S., Lamour, J., Li, Q., Morrison, B. D., Yang, D., Rogers, A., & Serbin, S. P. (2021). A best-practice guide to predicting plant traits from leaf-level hyperspectral data using partial least squares regression. *Journal of Experimental Botany*, erab295.

Carter, G. A. (1991). Primary and Secondary Effects of Water Content on the Spectral Reflectance of Leaves. *American Journal of Botany, 78*(7), 916–924.

Castro-Esau, K. L., Sánchez-Azofeifa, G. A., & Rivard, B. (2006). Comparison of spectral indices obtained using multiple spectroradiometers. *Remote Sensing of Environment, 103*(3), 276–288.

Cavender-Bares, J., Gamon, J. A., Hobbie, S. E., Madritch, M. D., Meireles, J. E., Schweiger, A. K., &

Townsend, P. A. (2017). Harnessing plant spectra to integrate the biodiversity sciences across biological

and spatial scales. *American Journal of Botany, 104*(7), 966–969.

Cavender-Bares, J., Meireles, J. E., Couture, J. J., Kaproth, M. A., Kingdon, C. C., Singh, A., Serbin, S.

P., Center, A., Zuniga, E., Pilz, G., & Townsend, P. A. (2016). Associations of Leaf Spectra with Genetic

and Phylogenetic Variation in Oaks: Prospects for Remote Detection of Biodiversity. *Remote Sensing*,

8(3), 221.

Chadwick, K. D., & Asner, G. P. (2016). Organismic-Scale Remote Sensing of Canopy Foliar Traits in

Lowland Tropical Forests. *Remote Sensing*, *8*(2), 87.

Clark, M. L., Roberts, D. A., & Clark, D. B. (2005). Hyperspectral discrimination of tropical rain forest

tree species at leaf to crown scales. *Remote Sensing of Environment, 96*(3), 375–398.

Costa, F. R. C., Lang, C., Almeida, D. R. A., Castilho, C. V., & Poorter, L. (2018). Near-infrared

spectrometry allows fast and extensive predictions of functional traits from dry leaves and branches.

Ecological Applications, 28(5), 1157–1167.

Couture, J. J., Singh, A., Rubert-Nason, K. F., Serbin, S. P., Lindroth, R. L., & Townsend, P. A. (2016).

Spectroscopic determination of ecologically relevant plant secondary metabolites. *Methods in Ecology*

and Evolution, *7*(11), 1402–1412.

Curran, P. J. (1989). Remote sensing of foliar chemistry. *Remote Sensing of Environment*, *30*(3), 271–

278.

Dechant, B., Cuntz, M., Vohland, M., Schulz, E., & Doktor, D. (2017). Estimation of photosynthesis

traits from leaf reflectance spectra: Correlation to nitrogen content as the dominant mechanism. *Remote*

Sensing of Environment, 196, 279–292.

Diniz, É. S., Amaral, C. H., Sardinha, S. T., Thiele, J., & Meira-Neto, J. A. A. (2020). Phylogenetic

signatures in reflected foliar spectra of regenerating plants in Neotropical forest gaps. *Remote Sensing of*

Environment, 112172.

Draper, F. C., Baker, T. R., Baraloto, C., Chave, J., Costa, F., Martin, R. E., Pennington, R. T., Vicentini,

A., & Asner, G. P. (2020). Quantifying Tropical Plant Diversity Requires an Integrated Technological

Approach. *Trends in Ecology & Evolution, 35*(12), 1100–1109.

Durgante, F. M., Higuchi, N., Almeida, A., & Vicentini, A. (2013). Species Spectral Signature:

Discriminating closely related plant species in the Amazon with Near-Infrared Leaf-Spectroscopy. *Forest*

Ecology and Management, *291*, 240–248.

Ely, K. S., Burnett, A. C., Lieberman-Cribbin, W., Serbin, S. P., & Rogers, A. (2019). Spectroscopy can

predict key leaf traits associated with source–sink balance and carbon–nitrogen status. *Journal of*

Experimental Botany, 70(6), 1789–1799.

Féret, J.-B., Gitelson, A. A., Noble, S. D., & Jacquemoud, S. (2017). PROSPECT-D: Towards modeling

leaf optical properties through a complete lifecycle. *Remote Sensing of Environment*, *193*, 204–215.

Foley, W. J., McIlwee, A., Lawler, I., Aragones, L., Woolnough, A. P., & Berding, N. (1998). Ecological

applications of near infrared reflectance spectroscopy – a tool for rapid, cost-effective prediction of the

composition of plant and animal tissues and aspects of animal performance. *Oecologia, 116*(3), 293–305.

Fourty, Th., Baret, F., Jacquemoud, S., Schmuck, G., & Verdebout, J. (1996). Leaf optical properties with

explicit description of its biochemical composition: Direct and inverse problems. *Remote Sensing of*

Environment, 56(2), 104–117.

Gitelson, A. A., Merzlyak, M. N., & Lichtenthaler, H. K. (1996). Detection of Red Edge Position and

Chlorophyll Content by Reflectance Measurements Near 700 nm. *Journal of Plant Physiology*, *148*(3),

501–508.

Heberling, J. M., & Isaac, B. L. (2017). Herbarium specimens as exaptations: New uses for old

collections. *American Journal of Botany*, 104, 963–965.

Heberling, J. M. (2021). Herbaria as Big Data Sources of Plant Traits. *International Journal of Plant*

Sciences. (in press)

Heberling, J. M., Miller, J. T., Noesgaard, D., Weingart, S. B., & Schigel, D. (2021). Data integration

enables global biodiversity synthesis. *Proceedings of the National Academy of Sciences, 118*(6),

e2018093118.

Hueni, A., Chisholm, L. A., Ong, C., Malthus, T. J., Wyatt, M., Trim, S. A., Schaepman, M. E., &

Thankappan, M. (2020). The SPECCHIO Spectral Information System. *IEEE Journal of Selected Topics*

in Applied Earth Observations and Remote Sensing, 13, 5789–5799.

Jacquemoud, S., & Baret, F. (1990). PROSPECT: A model of leaf optical properties spectra. *Remote*

Sensing of Environment, 34(2), 75–91.

Jacquemoud, S., & Ustin, S. (2019). Leaf Optical Properties. Cambridge University Press, New York.

Jardine, E. C., Thomas, G. H., Forrestel, E. J., Lehmann, C. E. R., & Osborne, C. P. (2020). The global

distribution of grass functional traits within grassy biomes. *Journal of Biogeography*, *47*(3), 553–565.

Kattenborn, T., Schiefer, F., Zarco-Tejada, P., & Schmidtlein, S. (2019). Advantages of retrieving

pigment content [µg/cm2] versus concentration [%] from canopy reflectance. *Remote Sensing of*

Environment, 230, 111195.

Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Tautenhahn, S., Werner, G. D.

A., Aakala, T., Abedi, M., et al. (2020). TRY plant trait database – enhanced coverage and open access.

Global Change Biology, 26(1), 119–188.

Kuhn, M. (2020). caret: Classification and Regression Training. R package version 6.0-86.

https://CRAN.R-project.org/package=caret

Kula, A. A. R., Hey, M. H., Couture, J. J., Townsend, P. A., & Dalgleish, H. J. (2020). Intraspecific

competition reduces plant size and quality and damage severity increases defense responses in the

herbaceous perennial, *Asclepias syriaca. Plant Ecology*, *221*(6), 421–430.

Lang, C., Almeida, D. R. A., & Costa, F. R. C. (2017). Discrimination of taxonomic identity at species,

genus and family levels using Fourier Transformed Near-Infrared Spectroscopy (FT-NIR). *Forest*

Ecology and Management, *406*, 219–227.

Lang, C., Costa, F. R. C., Camargo, J. L. C., Durgante, F. M., & Vicentini, A. (2015). Near Infrared

Spectroscopy Facilitates Rapid Identification of Both Young and Mature Amazonian Tree Species. *PLOS*

ONE, *10*(8), e0134521.

Lang, P. L. M., Willems, F. M., Scheepens, J. F., Burbano, H. A., & Bossdorf, O. (2019). Using herbaria

to study global environmental change. *New Phytologist*, *221*(1), 110–122.

Lindström, S. W., Geladi, P., Jonsson, O., & Pettersson, F. (2011). The Importance of Balanced Data Sets

for Partial Least Squares Discriminant Analysis: Classification Problems Using Hyperspectral Imaging

Data. *Journal of Near Infrared Spectroscopy, 19*(4), 233–241.

McManus, K. M., Asner, G. P., Martin, R. E., Dexter, K. G., Kress, W. J., & Field, C. B. (2016).

Phylogenetic Structure of Foliar Spectral Traits in Tropical Forest Canopies. *Remote Sensing*, *8*(3), 196.

Meacham-Hensold, K., Montes, C. M., Wu, J., Guan, K., Fu, P., Ainsworth, E. A., Pederson, T., Moore,

C. E., Brown, K. L., Raines, C., & Bernacchi, C. J. (2019). High-throughput field phenotyping using

hyperspectral reflectance and partial least squares regression (PLSR) reveals genetic modifications to

photosynthetic capacity. *Remote Sensing of Environment*, *231*, 111176.

Meineke, E. K., Davis, C. C., & Davies, T. J. (2018). The unrealized potential of herbaria for global

change biology. *Ecological Monographs*, *88*(4), 505–525.

Meireles, J. E., B. O'Meara, and J. Cavender-Bares. (2020a). Linking leaf spectra to the plant tree of life.

Pages 155-172 *in* J. Cavender-Bares, J. A. Gamon, and P. A. Townsend, editors. *Remote Sensing of Plant*

Biodiversity. Springer.

Meireles, J. E., J. Cavender-Bares, P. A. Townsend, S. Ustin, J. A. Gamon, A. K. Schweiger, M. E.

Schaepman, G. P. Asner, R. E. Martin, A. Singh, F. Schrodt, A. Chlus, and B. O'Meara. (2020b). Leaf

reflectance spectra capture the evolutionary history of seed plants. *New Phytologist* 228:485-493.

Mevik, B.-H., Wehrens, R., & Liland, K. H. (2019). pls: Partial Least Squares and Principal Component

Regression. R package version 2.7-1. https://CRAN.R-project.org/package=pls

Nakaji, T., Oguma, H., Nakamura, M., Kachina, P., Asanok, L., Marod, D., Aiba, M., Kurokawa, H.,

Kosugi, Y., Kassim, A. R., & Hiura, T. (2019). Estimation of six leaf traits of East Asian forest tree

species by leaf spectroscopy and partial least square regression. *Remote Sensing of Environment, 233*,

111381.

Nunes, M. H., Davey, M. P., & Coomes, D. A. (2017). On the challenges of using field spectroscopy to

measure the impact of soil type on leaf traits. *Biogeosciences*, *14*(13), 3371–3385.

Nunes, M. H., Both, S., Bongalov, B., Brelsford, C., Khoury, S., Burslem, D. F. R. P., Philipson, C.,

Majalap, N., Riutta, T., Coomes, D. A., & Cutler, M. E. J. (2019). Changes in leaf functional traits of

rainforest canopy trees associated with an El Niño event in Borneo. *Environmental Research Letters*,

14(8), 085005.

Osnas, J. L. D., Lichstein, J. W., Reich, P. B., & Pacala, S. W. (2013). Global Leaf Trait Relationships:

Mass, Area, and the Leaf Economics Spectrum. *Science, 340*(6133), 741–744.

Perez, T. M., Rodriguez, J., & Heberling, J. M. (2020). Herbarium-based measurements reliably estimate

three functional traits. *American Journal of Botany*, *107*(10), 1457–1464.

Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M.

S., Cornwell, W. K., Craine, J. M., Gurvich, D. E., et al. (2013). New handbook for standardised

measurement of plant functional traits worldwide. *Australian Journal of Botany, 61*(3), 167–234.

Peterson, D. L., Aber, J. D., Matson, P. A., Card, D. H., Swanberg, N., Wessman, C., & Spanner, M.

(1988). Remote sensing of forest canopy and leaf biochemical contents. *Remote Sensing of Environment,*

24(1), 85–108.

Petibon, F., Czyż, E. A., Ghielmetti, G., Hueni, A., Kneubühler, M., Schaepman, M. E., & Schuman, M.

C. (2021). Variation in reflectance spectroscopy of European beech leaves captures phenology and

biological hierarchies despite measurement uncertainties. *Remote Sensing of Environment*, *264*, 112601.

Prata, E. M. B., Sass, C., Rodrigues, D. P., Domingos, F. M. C. B., Specht, C. D., Damasco, G., Ribas, C.

C., Fine, P. V. A., & Vicentini, A. (2018). Towards integrative taxonomy in Neotropical botany:

Disentangling the Pagamea guianensis species complex (Rubiaceae). *Botanical Journal of the Linnean*

Society, 188(2), 213–231.

R Core Team. (2020). R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rapaport, T., Hochberg, U., Shoshany, M., Karnieli, A., & Rachmilevitch, S. (2015). Combining leaf

physiology, hyperspectral imaging and partial least squares-regression (PLS-R) for grapevine water status

assessment. *ISPRS Journal of Photogrammetry and Remote Sensing, 109*, 88–97.

Richardson, A. D., Duigan, S. P., & Berlyn, G. P. (2002). An evaluation of noninvasive methods to

estimate foliar chlorophyll content. *New Phytologist, 153*(1), 185–194.

Richardson, A. D., Aubrecht, D. M., Basler, D., Hufkens, K., Muir, C. D., & Hanssen, L. (2021).

Developmental changes in the reflectance spectra of temperate deciduous tree leaves and implications for

thermal emissivity and leaf temperature. *New Phytologist, 229*(2), 791–804.

Schweiger, A. K., Cavender-Bares, J., Townsend, P. A., Hobbie, S. E., Madritch, M. D., Wang, R.,

Tilman, D., & Gamon, J. A. (2018). Plant spectral diversity integrates functional and phylogenetic

components of biodiversity and predicts ecosystem function. *Nature Ecology & Evolution*, *2*(6), 976–982.

Schweiger, A. K. (2020). Spectral Field Campaigns: Planning and Data Collection. Pages 385-423 *in* J.

Cavender-Bares, J. A. Gamon, and P. A. Townsend, editors. *Remote Sensing of Plant Biodiversity*.

Springer.

Serbin, S. P., Singh, A., McNeil, B. E., Kingdon, C. C., & Townsend, P. A. (2014). Spectroscopic

determination of leaf morphological and biochemical traits for northern temperate and boreal tree species.

Ecological Applications, *24*(7), 1651–1669.

Serbin, S. P. & Townsend, P. A. (2020). Scaling Functional Traits from Leaves to Canopies. Pages 43-82

in J. Cavender-Bares, J. A. Gamon, and P. A. Townsend, editors. *Remote Sensing of Plant Biodiversity*.

Springer.

Serbin, S. P., Wu, J., Ely, K. S., Kruger, E. L., Townsend, P. A., Meng, R., Wolfe, B. T., Chlus, A.,

Wang, Z., & Rogers, A. (2019). From the Arctic to the tropics: Multibiome prediction of leaf mass per

area using leaf reflectance. *New Phytologist*, *224*(4), 1557–1568.

Streher, A. S., Torres, R. da S., Morellato, L. P. C., & Silva, T. S. F. (2020). Accuracy and limitations for

spectroscopic prediction of leaf traits in seasonally dry tropical environments. *Remote Sensing of*

Environment, *244*, 111828.

Sun, Y., Wong, A. K. C., & Kamel, M. S. (2009). Classification of imbalanced data: A review.

International Journal of Pattern Recognition and Artificial Intelligence, 23(04), 687–719.

Thiers, B. 2020. The world's herbaria 2019: A summary report based on data from Index Herbariorum.

Available at http://sweetgum.nybg.org/science/ih/

Turner, B. L., Hayes, P. E., & Laliberté, E. (2018). A climosequence of chronosequences in southwestern

Australia. *European Journal of Soil Science, 69*(1), 69–85.

Wang, Z., Chlus, A., Geygan, R., Ye, Z., Zheng, T., Singh, A., Couture, J. J., Cavender-Bares, J., Kruger,

E. L., & Townsend, P. A. (2020). Foliar functional traits from imaging spectroscopy across biomes in

eastern North America. *New Phytologist, 228*(2), 494–511.

Wang, Z., Skidmore, A. K., Darvishzadeh, R., & Wang, T. (2018). Mapping forest canopy nitrogen

content by inversion of coupled leaf-canopy radiative transfer models from airborne hyperspectral

imagery. *Agricultural and Forest Meteorology, 253–254*, 247–260.

Wieczorek, J., Bloom, D., Guralnick, R., Blum, S., Döring, M., Giovanni, R., Robertson, T., & Vieglais,

D. (2012). Darwin Core: An Evolving Community-Developed Biodiversity Data Standard. *PLOS ONE,*

7(1), e29715.

Willis, C. G., Ellwood, E. R., Primack, R. B., Davis, C. C., Pearson, K. D., Gallinat, A. S., Yost, J. M.,

Nelson, G., Mazer, S. J., Rossington, N. L., Sparks, T. H., & Soltis, P. S. (2017). Old Plants, New Tricks:

Phenological Research Using Herbarium Specimens. *Trends in Ecology & Evolution, 32*(7), 531–546.

Wold, S. (1994). PLS for multivariate linear modeling. In H. van de Waterbeemd, editor. Chemometric

methods in molecular design, methods and principles in medicinal chemistry, Verlag-Chemie (Weinheim,

Germany), 195–218.

Yan, Z., Guo, Z., Serbin, S. P., Song, G., Zhao, Y., Chen, Y., Wu, S., Wang, J., Wang, X., Li, J., Wang,

B., Wu, Y., Su, Y., Wang, H., Rogers, A., Liu, L., & Wu, J. (2021). Spectroscopy outperforms leaf trait

relationships for predicting photosynthetic capacity across different forest types. *New Phytologist, 232*(1),

134–147.

Yang, X., Tang, J., Mustard, J. F., Wu, J., Zhao, K., Serbin, S., & Lee, J.-E. (2016). Seasonal variability

of multiple leaf traits captured by leaf spectroscopy at two temperate deciduous forests. *Remote Sensing*

of Environment, 179, 1–12.

Supplementary Materials

Sampling

For each sample, we collected a large group of sunlit leaves (>3 h estimated sun exposure per day) at

about the same vertical position—for trees, the same branch or neighboring branches from the same

individual. We avoided leaves with noticeable damage from herbivores or pathogens. We measured fresh-

leaf spectra and fresh-leaf mass in the field, froze leaf disks for pigment analyses, and transported samples

in a cooler to the lab. We then measured other whole-leaf traits and divided the remainder of the sample

into two groups: (1) a portion to be pressed, and (2) a portion whose leaves were dried at 65 °C for three

days, ground using a cyclone mill with 2 mm mesh, redried at 65 °C, and stored for chemical analyses.

We measured pressed-leaf spectra on the former portion prior to mounting, and ground-leaf spectra and

chemical traits on the latter. Although these are distinct subsets of leaves, we treat them as identical in

their trait values since they are divided at random from the larger, relatively homogeneous bulk sample. A

minimum of one representative pressed specimen of each species at each site and date is prepared as a

specimen at the Marie-Victorin Herbarium in Montréal, Québec, Canada.

We classified species' growth forms following the Database of Vascular Plants of Canada (VASCAN;

Desmet & Brouillet 2013). We classified *Agonis flexuosa* as a tree and manually disambiguated some

species listed as adopting either shrub or three growth forms.

Spectral measurements

For the Beauchamp-Rioux, Boucherville, and Warren projects, we measured fresh-leaf directional-

hemispherical reflectance spectra on each sample in 2018 using an HR-1024i spectroradiometer outfitted

with a DC-R/T integrating sphere from Spectra Vista Corporation (SVC; Poughkeepsie, NY, USA). We

measured spectra on the adaxial surface of six leaves or, for small or narrow leaves, single-layer leaf

arrays following the protocol in Noda et al. (2013) as adapted for our instrument (Laliberté & Soffer

2018a; Laliberté and Soffer 2018b). For the Dessain project, we measured fresh-leaf reflectance spectra in

2017 using a FieldSpec 4 spectroradiometer equipped with an RTS-3ZC integrating sphere from

Analytical Spectral Devices, Inc. (ASD; Boulder, CO, USA) following the leaf spectroscopy protocol

from the Global Airborne Observatory (https://gao.asu.edu/spectranomics-protocols). We collected

spectra from a variable number of leaves or arrays per sample—six in most cases, but as few as three or

as many as eighteen from certain samples. For all projects, each reflectance measurement was calibrated

against a white Spectralon 99% reflectance panel and corrected for stray light.

For all projects, we measured reflectance spectra on the adaxial surface of pressed, intact leaf samples

using a PSR+ 3500 spectroradiometer with a leaf clip foreoptic (Spectral Evolution, Haverhill, MA)

between six months and three years after collection. We calibrated reflectance against a white Spectralon

reflectance panel and measured 3-7 spectra per sample. Although all samples were pressed in a plant

press, some nevertheless dried in a way that created an uneven leaf surface. We noted any visible

discoloration that samples underwent during drying or storage, including browning or blackening. In

general, we aimed to avoid uneven or discolored areas when measuring spectra, although it was

sometimes not entirely possible when discoloration was pervasive.

For all projects, we also measured ground-leaf spectra using the PSR+ 3500 spectroradiometer with a

benchtop reflectance probe foreoptic that pressed loose powder into a smooth, even pellet. For each

sample, we added at least 0.6 g of leaf powder into a sample tray; in preliminary tests, adding additional

material did not change the spectra, suggesting that transmittance was close to zero. We calibrated

reflectance against a white Spectralon reflectance panel placed in an identical sample tray. We measured

three spectra per sample on the same pellet, turning the sample tray 120° between the first two spectra,

and loosening and mixing the powder before reshaping the pellet between the second and third spectra.

A small number of specimens did not have usable spectra at all three stages, usually either due to lack of

enough pressed or ground material or to poor spectral quality at any stage. To ensure fair comparisons

among the stages, we removed any sample from the dataset that was not available for at least two stages.

Among the remaining samples, any sample that was not available for all three stages (<4%) was placed in

the training subset for PLSR analyses.

Our spectral processing pipeline varied by the type of data. Although the actual spectral resolution was

more variable and coarser, the ASD Field Spec 4 and Spectral Evolution PSR+ 3500 spectrometers'

software automatically resampled the spectra to 1 nm resolution and interpolated over the overlap region

between sensors (Schweiger & Laliberté 2020). For data from the SVC HR-1024i spectrometer, we

performed these steps using linear interpolation. For pressed-leaf spectra only, we detected jumps in

reflectance at the sensor overlap region and removed them using the function *match_sensors()* in *R*

package *spectrolab v. 0.0.10* (Meireles et al. 2017). Next, we averaged all spectra for a given sample and

tissue type (fresh, pressed, or ground). Spectra measured with integrating spheres tend to have greater

noise, especially at the ends. To reduce noise in fresh-leaf spectra, we applied a Savitzky-Golay filter

using *R* package *signal 0.7.6* (Signal Developers, 2013), with varying order and length: order 3 and length

21 from 350-715 nm, order 3 and length 35 from 715-1390 nm, order 3 and length 75 from 1390-1880

nm, and order 5 and length 175 from 1880-2500 nm. Finally, we trimmed all spectra to 400-2400 nm,

removing the extremes where sensors show greater noise. We did all spectral processing in *R v. 3.6.3* (R

Core Team 2020) using *spectrolab v. 0.0.10* (Meireles et al. 2017).

1010

Leaf trait measurements

We performed all leaf trait measurements excluding petioles, but including the rachis for compound

leaves, since the rachis is functionally analogous to the midrib of a simple leaf. We measured the

following leaf structural and chemical traits: Leaf mass per area (LMA; $kg\ m^{-2}$), leaf dry matter content

(LDMC; $mg\ g^{-1}$), equivalent water thickness (EWT; cm), carbon fractions (soluble cell contents,

hemicellulose, cellulose, and lignin) on a mass basis, pigments (chlorophyll *a*, chlorophyll *b*, and total

carotenoids) on a mass basis, and concentrations of a variety of elements (Al, C, Ca, Cu, Fe, K, Mg, Mn,

N, Na, P, Zn).

1019

1020 From a subset of the leaves for each sample whose fresh-leaf spectra we measured, we used a 7 mm cork

1021 borer to punch leaf disks (or cut fragments of narrow leaves) in the field and stored them on ice in the

1022 darkness before transferring them to a -80 °C freezer in the lab. We extracted pigments under darkness in

1023 an $MgCO_3$-MeOH solution and estimated pigment concentrations using a spectrophotometric protocol on

1024 a SPECTROstar Nano microplate reader (BMG LABTECH, Ortenburg, Germany; Girard et al. 2020).

1025

From a separate subset of leaves from each sample, we weighed fresh mass in the field shortly after

collection. We then rehydrated the leaves in sealed plastic bags with damp paper towels for at least 12 h

and weighed them for rehydrated mass. We scanned leaves and estimated their area using WinFOLIA

(Regent Instruments, Québec, QC, CA). Lastly, we dried them for at least 72 h in a drying oven at 65°C

before reweighing for dry mass. We calculated LDMC as total dry mass divided by total fresh mass and

LMA as total dry mass divided by total rehydrated area (Laliberté 2018). We further calculated EWT

from LDMC and LMA.

1033

We analyzed ground bulk tissue samples for N and C using a Vario MICRO Cube combustion analyzer

(Elementar, Langenselbold, Germany; Ayotte et al. 2019). We analyzed the same samples for other

elements (Al, Ca, Cu, Fe, K, Mg, Mn, Na, P, Zn) using inductively coupled plasma-mass spectrometry

(ICP-MS). We removed five data points from the ICP-MS output as extreme outliers that were not

representative of the range of concentrations of their respective elements across our samples. We also set

values below the detection limit, which sometimes returned as negative concentrations, to zero.

1040

Finally, we analyzed the samples for carbon fractions using an ANKOM 2000 Fiber Analyzer (Ankom

Technology, Macedon, New York, USA) to perform a sequence of digestions (Ayotte & Laliberté 2019).

The first digestion in a heated neutral detergent washes off soluble cell contents. The second digestion in

a heated acidic detergent removes hemicellulose and bound proteins—which, for simplicity, we just refer

to as hemicellulose. The third digestion in 70% sulfuric acid removes cellulose, leaving behind lignin and

recalcitrant compounds. We used the change in each bag's mass between stages to determine the mass of

each fraction. We ashed all samples and used ash content as a correction factor to calculate the ash-free

lignin fraction.

1049

External validation dataset

From August to November 2018, we collected 333 samples for an external validation dataset at Cedar

Creek Ecosystem Science Reserve (East Bethel, MN, USA). These samples included broadleaf trees (n =

177; 10 species), coniferous needleleaf trees (n = 101; 4 species), and herbs (n = 55; 14 species). Most

samples were from the Forests and Biodiversity (FAB) experiment (Grossman et al. 2017), the

Biodiversity in Willows and Poplars (BiWaP) experiment (Grossman & Cavender-Bares 2019), and the

Big Biodiversity (BioDIV or e120) experiment (Tilman et al. 1997), although some were from other parts

of the site.

1058

Each sample included multiple leaves from the same individual, taken when possible from a nearby

position on the plant. Unlike in the CABO dataset, we collected samples from any part of an individual,

not just sunlit leaves. From each sample, we designated one leaf for fresh mass to be measured

immediately following collection; this leaf ('fresh-mass subsample') was stored separately and its spectra

were labeled to keep track of its identity. (For conifers or other plants with small or narrow leaves, we

often included multiple leaves in this fresh mass sample.) We took and flash-froze 3-5 hole punches from

another leaf (or group of leaves) for pigment analyses ('pigment subsample'; pigment data not shown);

this leaf and its spectra were also labeled separately. We collected more leaves to have enough tissue for a

bulk chemical sample, but did not keep track of them individually.

1068

In the field, we measured multiple fresh-leaf spectra per sample (not shown here) using a PSR+ 3500

spectroradiometer with a leaf clip foreoptic (Spectral Evolution, Haverhill, MA) and placed the leaves in

sealed plastic bags with a damp paper towel. These bags were kept in coolers until the end of the

sampling day. We weighed the fresh-mass subsample immediately upon return to the lab and kept all

leaves in a refrigerator until we scanned the fresh-mass and pigment subsamples within the next few days.

After scanning, we placed all leaves in flat paper bags and pressed them for 2-3 hours in a plant press

before removing them and placing them in a drying shed at 40 °C for at least one week. This deviates

from most standard herbarium protocols, including the one used in the CABO project, where leaves are

kept in the press while drying and subjected to a higher temperature for a shorter time. After we dried the

leaves, we weighed the fresh-mass and pigment subsamples. Certain species, especially *Populus*

tremuloides, seemed especially prone to discoloration during sample preparation.

1080

In May 2020, we measured pressed-leaf spectra using a PSR+ 3500 spectroradiometer with a leaf clip

foreoptic. We did not use a specific protocol for small or narrow leaves, but aimed to arrange leaves in a

way that minimized overlaps or gaps. We removed spectra with especially low (<40%) or high (>80%)

peak reflectance, and any with major visible gaps in the sensor overlap regions. We measured three

spectra per subsample except in cases where getting high-quality spectra proved especially difficult. We

then removed the petioles (when applicable) and put leaves in scintillation vials with ball bearings to

grind the tissue in a paint shaker, combining all subsamples from a given sample. We then dried samples

again and measured C and N with an elemental analyzer (Costech ECS 4010 Analyzer, Valencia,

California, USA).

1090

1091 For estimating LMA, LDMC, and EWT, we only used the spectra from the fresh-mass subsample. For C

1092 and N, we used all spectra from a given sample, ignoring the subsample labels. As among the CABO

1093 pressed-leaf spectra, we first matched sensors, then averaged spectra within the relevant sample or

1094 subsample, and lastly trimmed them to 400-2400 nm.

1095

1096 We applied the ensemble of models resulting from our 100× jackknife analysis on the pressed-leaf CABO

1097 data and applied them to the Cedar Creek data. We also applied the model ensemble from the pressed-leaf

1098 CABO data restricted to 1300-2400 nm to the Cedar Creek data restricted to the same range. We

1099 calculated summary statistics and plotted results as in the internal validation.

1100 **Supplementary references**

1101

1102 Ayotte, J., Guilbeault-Mayers, X., & Laliberté, E. (2019). Measuring CN content in leaf samples using

1103 Elementar Vario MICRO Cube. https://doi.org/10.17504/protocols.io.udces2w

1104

1105 Ayotte, J., & Laliberté, E. (2019). Measuring leaf carbon fractions with the ANKOM2000 Fiber

1106 Analyzer. https://doi.org/10.17504/protocols.io.yinfude

1107

1108 Desmet, P., & Brouillet, L. (2013). Database of Vascular Plants of Canada (VASCAN): A community

1109 contributed taxonomic checklist of all vascular plants of Canada, Saint Pierre and Miquelon, and

1110 Greenland. *PhytoKeys, 25*, 55–67.

1111

1112 Girard, A., Ayotte, J., & Laliberté, E. (2020). Measuring chlorophylls and carotenoids in plant tissue.

1113

1114 Grossman, J. J., & Cavender-Bares, J. (2019). Consequences of biodiversity shift across phylogenetic

1115 scales for aspen and willow growth, survival, and herbivory. *Journal of Vegetation Science, 30*(2), 301–

1116 311.

1117

1118 Grossman, J. J., Cavender-Bares, J., Hobbie, S. E., Reich, P. B., & Montgomery, R. A. (2017). Species

1119 richness and traits predict overyielding in stem growth in an early-successional tree diversity experiment.

1120 *Ecology, 98*(10), 2601–2614.

1121

1122 Laliberté, E. (2018). Measuring specific leaf area and water content.

1123 https://doi.org/10.17504/protocols.io.p3tdqnn

1124

1125   Laliberté, E., & Soffer, R. (2018a). Measuring spectral reflectance and transmittance (350-2500 nm) of

1126   large leaves using the Spectra Vista Corporation (SVC) DC-R/T Integrating Sphere.

1127   https://doi.org/10.17504/protocols.io.p8pdrvn

1128

1129   Laliberté, E., & Soffer, R. (2018b). Measuring spectral reflectance and transmittance (350-2500 nm) of

1130   small and/or narrow leaves using the Spectra Vista Corporation (SVC) DC-R/T Integrating Sphere.

1131   https://doi.org/10.17504/protocols.io.q56dy9e

1132

1133   Meireles, J., Schweiger, A., & Cavender-Bares, J. (2017). spectrolab: Class and Methods for

1134   Hyperspectral Data in R. R package version 0.0.10. https://CRAN.R-project.org/package=spectrolab

1135

1136   Noda, H. M., Motohka, T., Murakami, K., Muraoka, H., & Nasahara, K. N. (2013). Accurate

1137   measurement of optical properties of narrow leaves and conifer needles with a typical integrating sphere

1138   and spectroradiometer. *Plant, Cell & Environment*, *36*(10), 1903–1909.

1139

1140   R Core Team. (2020). R: A language and environment for statistical computing. R Foundation for

1141   Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

1142

1143   Schweiger, A. K., & Laliberté, E. (2020). Processing of leaf spectra.

1144   https://doi.org/10.17504/protocols.io.bhsdj6a6

1145

1146   Signal Developers (2013). signal: Signal processing. http://r-forge.r-project.org/projects/signal/.

1147

1148   Tilman, D., Knops, J., Wedin, D., Reich, P., Ritchie, M., & Siemann, E. (1997). The Influence of

1149   Functional Diversity and Composition on Ecosystem Processes. *Science, 277*(5330), 1300–1302.

Supplementary tables

Table S1: Summary statistics of PLSR model internal validation from pressed-leaf spectra restricted to 1300-2500 nm. %RMSE is calculated as RMSE divided by the 2.5% trimmed range of data within either the training data set (calibration) or the testing data set (validation).

Trait	No. comps	R^2	RMSE	%RMSE
LMA (kg m^{-2})	7	0.902	0.0125	7.94
LDMC (mg g^{-1})	10	0.694	37.6	12.8
EWT (cm)	10	0.767	0.00222	13.3
Solubles (%)	10	0.712	4.25	13.5
Hemicellulose (%)	10	0.733	2.48	15.6
Cellulose (%)	11	0.745	1.50	17.9
Lignin (%)	11	0.638	2.07	16.7
C (%)	11	0.811	0.987	10.8
N (%)	9	0.772	0.273	11.5
Chl a (mg g^{-1})	7	0.582	1.37	17.3
Chl b (mg g^{-1})	7	0.514	0.506	18.7
Carotenoids (mg g^{-1})	7	0.599	0.280	17.9
Al (mg g^{-1})	6	0.210	0.0420	24.7
Ca (mg g^{-1})	11	0.723	3.72	13.2
Cu (mg g^{-1})	8	0.243	0.00676	24.0
Fe (mg g^{-1})	4	0.223	0.0391	20.9
K (mg g^{-1})	11	0.566	2.93	18.6
Mg (mg g^{-1})	11	0.413	0.733	20.1
Mn (mg g^{-1})	10	0.368	0.140	22.4
Na (mg g^{-1})	12	0.434	0.577	20.6
P (mg g^{-1})	11	0.507	0.772	19.0
Zn (mg g^{-1})	9	0.295	0.0712	25.1

Table S2: Summary statistics for external validation of pressed-leaf PLSR models, restricting the range of spectra to 1300-2500 nm. The models were trained on CABO data (see Table S1) and applied to a dataset collected at Cedar Creek.

Trait	Including conifers			Excluding conifers		
	R^2	RMSE	%RMSE	R^2	RMSE	%RMSE
LMA (kg m^{-2})	0.867	0.0195	10.8	0.746	0.0111	13.7
LDMC (mg g^{-1})	0.468	51.2	18.1	0.461	53.4	18.5
EWT (cm)	0.608	0.00512	20.7	0.633	0.00318	17.6
C (%)	0.007	4.54	26.0	0.023	4.38	28.2
N (%)	0.622	0.666	34.7	0.614	0.772	30.3

1159 **Supplementary figure captions**

1160

1161 Fig. S1: Internal validation results for predictions of LMA, LDMC, and EWT from fresh- (left), pressed-

1162 (middle), and ground-leaf (right) spectra. In each panel, a separate ordinary least-squares regression line

1163 is shown for each functional group, overlaid on top of the thick dashed 1:1 line. The error bars for each

1164 data point are 95% confidence intervals calculated from the distribution of predictions based on the model

1165 coefficients from the 100 jackknife iterations.

1166

1167 Fig. S2: Internal validation results for predictions of C, N, and P from fresh- (left), pressed- (middle), and

1168 ground-leaf (right) spectra, displayed as in Fig. S1.

1169

1170 Fig. S3: Internal validation results for predictions of soluble cell contents, hemicellulose, cellulose, and

1171 lignin from fresh- (left), pressed- (middle), and ground-leaf (right) spectra, displayed as in Fig. S1.

1172

1173 Fig. S4: Internal validation results for predictions of chlorophyll a, chlorophyll b, and total carotenoids

1174 from fresh- (left), pressed- (middle), and ground-leaf (right) spectra, displayed as in Fig. S1.

1175

1176 Fig. S5: Internal validation results for predictions of Al, Ca, and Cu from fresh- (left), pressed- (middle),

1177 and ground-leaf (right) spectra, displayed as in Fig. S1.

1178

1179 Fig. S6: Internal validation results for predictions of Fe, K, and Mg from fresh- (left), pressed- (middle),

1180 and ground-leaf (right) spectra, displayed as in Fig. S1

1181

1182 Fig. S7: Internal validation results for predictions of Mn, Na, and Zn from fresh- (left), pressed- (middle),

1183 and ground-leaf (right) spectra, displayed as in Fig. S1

1184

1185 Fig. S8: The variable importance of projection (VIP) metric calculated for all traits from fresh-leaf

1186 spectra. Abbreviations: sol = solubles, hemi = hemicellulose, cell = cellulose, lign = lignin, car = total

1187 carotenoids. The dashed horizontal line at 0.8 represents the heuristic threshold for importance

1188 recommended by Burnett et al. (2021).

1189

1190 Fig. S9: The variable importance of projection (VIP) metric calculated for all traits from pressed-leaf

1191 spectra. Abbreviations: sol = solubles, hemi = hemicellulose, cell = cellulose, lign = lignin, car = total

1192 carotenoids. The dashed horizontal line at 0.8 represents the heuristic threshold for importance

1193 recommended by Burnett et al. (2021).

1194

1195 Fig. S10: The variable importance of projection (VIP) metric calculated for all traits from ground-leaf

1196 spectra. Abbreviations: sol = solubles, hemi = hemicellulose, cell = cellulose, lign = lignin, car = total

1197 carotenoids. The dashed horizontal line at 0.8 represents the heuristic threshold for importance

1198 recommended by Burnett et al. (2021).

1199

1200 Fig. S11: The distributions of fresh-leaf spectral model performance statistics for each trait based on 100

1201 jackknife resamples from the calibration data set. The red dots show the R^2 and %RMSE from applying

1202 the ensemble of models to the internal validation subset. Abbreviations: sol = solubles, hemi =

1203 hemicellulose, cell = cellulose, lign = lignin, car = total carotenoids.

1204

1205 Fig. S12: The distributions of pressed-leaf spectral model performance statistics for each trait based on

1206 100 jackknife resamples from the calibration data set. The red dots show the R^2 and %RMSE from

1207 applying the ensemble of models to the internal validation subset. Abbreviations: sol = solubles, hemi =

1208 hemicellulose, cell = cellulose, lign = lignin, car = total carotenoids.

1209

1210   Fig. S13: The distributions of ground-leaf spectral model performance statistics for each trait based on

1211   100 jackknife resamples from the calibration data set. The red dots show the R^2 and %RMSE from

1212   applying the ensemble of models to the internal validation subset. Abbreviations: sol = solubles, hemi =

1213   hemicellulose, cell = cellulose, lign = lignin, car = total carotenoids.

1214

1215   Fig. S14: External validation results for predictions of various traits in Cedar Creek data based on models

1216   trained on CABO data, restricted to 1300-2500 nm. Each panel is displayed as in Figs. S1-7.

Supplementary figures

Fig. S1

Fig. S2

Fig. S3

235 **Fig. S4**

Fig. S5

Fig. S6

Fig. S7

Ground