Reflectance spectroscopy allows rapid, accurate, and non-destructive estimates of functional traits from pressed leaves

Shan Kothari[1,2,4], Rosalie Beauchamp-Rioux[1], Etienne Laliberté[1], Jeannine Cavender-Bares[2,3]

[1]Institut de recherche en biologie végétale, Département de sciences biologiques, Université de Montréal
[2]Department of Plant and Microbial Biology, University of Minnesota
[3]Department of Ecology, Evolution, and Behavior, University of Minnesota
[4]Corresponding author: shan.kothari@umontreal.ca

Abstract

More than ever, ecologists seek to employ herbarium collections as tools to estimate plant functional traits

from the past or from places that are hard to sample. However, many functional trait measurements are

destructive, which may preclude their use on valuable herbarium specimens. Reflectance spectroscopy is

increasingly used to estimate traits rapidly from fresh or dried, ground leaves, as well as to differentiate

and identify taxa. Here, we extend this body of work to pressed, intact leaves such as those in herbarium

collections. Using a dataset with 619 plant samples belonging to 70 woody and herbaceous species, we

used partial least-squares regression to build validated models linking pressed-leaf reflectance spectra to a

broad suite of traits, including leaf mass per area (LMA), leaf dry matter content (LDMC), carbon (C_{mass})

and nitrogen (N_{mass}) concentrations, and carbon fractions such as cellulose and lignin. We compared the

accuracy of these trait estimates to those from fresh- or ground-leaf spectra of the same samples. Our

pressed-leaf models predicted these traits with moderate-to-high accuracy (R^2 = 0.586-0.924; %RMSE =

5.7-11.7%). For estimating chemical traits, pressed-leaf models performed better than fresh-leaf models

but slightly worse than ground-leaf models. Pressed-leaf models did not perform as well as fresh-leaf

models for estimating LMA and LDMC, but outperformed ground-leaf models for LMA. Accuracy was

no worse for pressed leaves that underwent discoloration in storage. Finally, on a subset of common

species in the dataset, we used partial least-squares discriminant analysis to classify specimens to species

with near-perfect accuracy from pressed- (>97%), and ground-leaf (>96%) spectra and slightly lower

accuracy from fresh-leaf spectra (>89%). The success of trait estimation and species classification from

pressed-leaf spectra may owe to the fact that they combine advantages of fresh and ground leaves: like

32 fresh leaves, they retain some of the spectral expression of internal leaf structure; like ground leaves, they

33 reveal minor absorption features of chemical constituents that would otherwise be masked by water.

34 These results show that applying spectroscopy to pressed leaves is a promising way to estimate leaf

35 functional traits non-destructively. Our study has far-reaching implications for capturing the wide range

36 of functional, phenotypic, and taxonomic information in the world's preserved plant collections.

Introduction

The world's herbaria together contain more than 390 million specimens (Thiers 2020), which are collectively a rich source of information about global plant diversity. Herbarium specimens are collected for many reasons—for example, to document that a species is present in a certain location, or to serve as vouchers or sources for DNA sequences used to estimate phylogenies. But these specimens are often repurposed for new ends, unforeseen by the collectors (Heberling and Isaac 2017; Meineke et al. 2018; Heberling et al. 2019). More than ever, ecologists and evolutionary biologists are interested in using herbarium specimens to measure functional traits: for example, to evaluate long-term changes, often in response to human activity (e.g. Hietz et al. 2011; Willis et al. 2017; Koski et al. 2020; reviewed in Lang et al. 2018); to fill in gaps in sparse trait databases (Queenborough and Porras 2014; Perez et al. 2020); or to conduct comparative studies of clades whose species are hard to sample all at once (e.g. Jardine et al. 2020). Measuring functional traits on herbarium specimens carries the promise of letting us reach the inaccessible, including the past or distant parts of the world. Moreover, using herbarium specimens allows researchers to benefit from the taxonomic skill of collectors and curators and refer back to the same specimens for further use—for example, as prototypes for identification of taxa from new collections. By repurposing herbarium specimens towards new ends, researchers can address ecological and evolutionary questions that require merging functional, genetic, and distributional data at global scales.

Many functional trait measurements require destructive sampling—for example, by grinding up tissue for chemical analyses. Such methods include most techniques to determine the content or concentration of various elements, isotopes, macromolecules, or metabolites in a sample. Because of the high value of herbarium specimens—especially ones from historical collections—herbarium curators may hesitate to let samples be destroyed for ecological research, even in part. The development of non-destructive techniques to estimate leaf traits from herbarium specimens would make it more acceptable to use them in functional ecology.

Reflectance spectroscopy is a well-established technique used to estimate a variety of foliar functional

traits non-destructively (Curran 1989). A reflectance spectrum is a measurement of the reflectance of light

from a surface at a range of wavelengths. For leaf tissue, researchers tend to measure reflectance between

about 350 and 2500 nm, a range that includes more than 98% of solar radiation reaching Earth's surface

(American Society for Testing and Materials, 2006). Because the leaf's chemical and structural makeup

determines how it absorbs, reflects, and transmits solar radiation, reflectance within this range carries

information about many critical plant traits. These underlying chemical and structural traits are shaped by

past evolutionary forces, raising the possibility that leaf reflectance spectra can also be used to distinguish

lineages and study trait evolution (Meireles et al. 2020a).

Two main approaches exist to estimate traits from entire reflectance spectra. First, physics-based radiative

transfer models like PROSPECT can be applied to fresh-leaf spectra to estimate traits like chlorophyll and

water content and the number of mesophyll cell layers (Jacquemoud & Baret 1990; Féret et al. 2017).

Second, statistical models, often created using machine learning techniques like partial least-squares

regression (PLSR), can be used to estimate an even wider range of traits, albeit in a less mechanistic way

(Serbin & Townsend 2020). Many ecologically important traits, like leaf nitrogen concentration (N_{mass}),

lack a unique set of strong absorption features, complicating the development of physical models. For

example, nitrogen is found alike in nucleic acids, chlorophyll, and proteins (e.g. RuBisCO), in ratios that

vary from plant to plant. Moreover, absorption features of chemical bonds and compounds often overlap,

especially after being broadened by multiple scattering (Curran 1989). A multivariate empirical approach

based on statistical techniques like PLSR or stepwise regression gives researchers the flexibility needed to

predict complex traits (Curran 1989). Likewise, multivariate classification techniques like partial least-

squares discriminant analysis (PLS-DA) are often used to take advantage of the full spectrum in

discriminating species, lineages, or other kinds of biological classes (Castro-Esau et al. 2006; Gold et al.

2020; Meireles et al. 2020b).

Empirical approaches like PLSR have been widely used by ecologists to estimate plant traits from

spectroscopic data measured on fresh or ground leaves. These traits include N_{mass} and LMA (Doughty et

al. 2011; Asner et al. 2011a; Serbin et al. 2014; Serbin et al. 2019; Streher et al. 2020), pigments (Asner et

al. 2011a; Yang et al. 2016), condensed tannins and other defense compounds (Couture et al. 2016), non-

structural carbohydrates (Ramirez et al. 2015; Ely et al. 2019), and even physiological rates like the

maximum rate of RuBP carboxylation in photosynthesis (V_{cmax}; Serbin et al. 2012; Wu et al. 2019;

Meacham-Hensold et al. 2019). Such leaf-level PLSR models have been used to address such varied

ecological subjects as defense responses to herbivory (Kula et al. 2020), the role of biodiversity in

ecosystem function (Schweiger et al. 2018), and functional trait responses to El Niño (Nunes et al. 2019).

Although this multivariate statistical approach is flexible, it is sensitive to the kind of leaf tissue used for

model development. For example, existing PLSR models have mostly been trained on spectra of fresh

leaves (e.g. Serbin et al. 2019) or dried, ground leaves (e.g. Serbin et al. 2014). Such models cannot be

expected to transfer to dried, intact leaves like herbarium specimens because both drying and grinding

produce major changes in the features of the reflectance spectrum.

We tested whether we could use a PLSR modeling approach to estimate traits from the spectra of pressed

leaves like herbarium specimens. We ask: Could models calibrated and validated on pressed leaves be as

accurate as those from fresh or ground leaves? For most chemical traits, such as N_{mass} or leaf carbon

fractions, we conjectured that ground-leaf spectral models would be the most accurate, as prior studies

have found (Serbin et al. 2014; Couture et al. 2016; Wang et al. 2020). Both drying and grinding might be

important in achieving this high accuracy. Drying may reveal minor absorption features of compounds in

dry matter within the short-wave infrared range (SWIR) that, in fresh leaves, are obscured by the

dominant effect of water absorption (Peterson et al. 1988; Ecarnot et al. 2013; Couture et al. 2016).

Grinding homogenizes the leaf lamina, which otherwise varies in its structure and composition across its

surface and along its cross-section, altering reflectance in complex ways due to internal scattering (Grant

1987). By evening out this variation, grinding may allow us to capture a more even and representative

sample of tissue. Since pressed leaves are dried but not ground, we predict their spectra would produce

estimates of intermediate accuracy.

For structural traits like LMA, we instead expect fresh and pressed leaves to outperform ground leaves

because grinding disrupts the leaf structure. Moreover, for water-related traits like leaf dry matter content

(LDMC; dried mass divided by fresh mass), we expect fresh leaves to outperform both pressed and

ground leaves because only they retain the water absorption features that allow direct, quantitative

prediction of water content (Carter 1991; Rapoport et al. 2015). An additional consideration for herbarium

specimens and other pressed-leaf tissue collections is that their optical properties may change in

preparation or storage. We considered whether discoloration reduces the accuracy of trait estimates,

which may indicate whether spectroscopy is useful on old or degraded specimens.

Finally, we asked how accurately pressed-leaf spectra can be used to identify samples to species.

Reflectance spectra often show phylogenetic signal, at least in particular wavelength ranges, because of

phylogenetic conservatism in the evolution of their underlying traits (McManus et al. 2016; Diniz et al.

2020; Meireles et al. 2020a; Schweiger et al. 2021). This phylogenetic signal is what often makes it

possible to classify species or higher-level taxa from fresh-leaf spectra (Cavender-Bares et al. 2016;

Meireles et al. 2020b), but it may or may not be retained in pressed-leaf spectra. A few studies with

tropical forest species have indeed demonstrated that absorbance spectra of pressed herbarium specimens

can be used to classify species or higher-level taxa (Durgante et al. 2013; Lang et al. 2015; Lang et al.

2017; Prata et al. 2018). Here, we build on these studies by comparing the accuracy of classification from

fresh-, pressed-, and ground-leaf spectra among the common species in our dataset. Because they preserve

some of the spectral expression of leaf structure (unlike ground leaves) and reveal the distinctive SWIR

absorption features of macromolecules and other compounds (unlike fresh leaves), pressed leaves may

represent the best of both worlds for distinguishing species using spectroscopy.

Methods

Sampling

We compiled data from four projects, all part of the Canadian Airborne Biodiversity Observatory

(CABO). Each project includes reflectance spectra and several structural and chemical leaf traits. The

projects are:

1. Beauchamp-Rioux: Deciduous forest trees sampled throughout the growing season in 2018 at

various sites in southern Québec and Ontario ($n = 417$).

2. Dessain: Forbs, grasses, shrubs, and broadleaf trees from forests and open areas sampled

throughout the growing season in 2017 at four sites in southern Québec ($n = 75$).

3. Boucherville: Forbs, shrubs, and grasses sampled throughout the growing season in 2018 from

the Parc national des Îles-de-Boucherville in southern Québec ($n = 72$).

4. Warren: A single broadleaf evergreen tree species (*Agonis flexuosa* (Willd.) Sweet) collected

across a soil chronosequence in southwestern Australia (Turner et al. 2018; $n = 67$).

The full dataset includes 70 species, whose growth forms we classified following the Database of

Vascular Plants of Canada (VASCAN; Desmet & Brouillet 2013). We classified *Agonis flexuosa* as a tree

and manually disambiguated some species listed as adopting either shrub or three growth forms. The

majority of samples were from trees ($n = 518$), with shrubs ($n = 47$) and forbs ($n = 36$) also moderately

represented.

In general, we aimed to avoid leaves with noticeable damage from herbivores or pathogens. For each

dataset, we measured reflectance spectra (350-2500 nm) of the leaves at three stages: (1) freshly sampled,

(2) pressed and dried, and (3) ground into a fine powder. We also measured a suite of leaf functional traits

on each of these samples. Further details on spectral and trait measurements are provided in later

subsections, but we first lay out our sampling procedure.

For each sample, we collected a group of sunlit leaves (>3 h estimated sun exposure per day) at about the

same vertical position of the same individual—the same branch or neighboring branches, for trees. After

measuring fresh-leaf spectra and fresh mass in the field, we took leaves to the lab to measure leaf

structural traits, such as leaf dry matter content (LDMC) and LMA. We then divided the leaves into two

groups: (1) a portion to be pressed, and (2) a portion to be dried at 65 °C for three days, ground using a

cyclone mill with 2 mm mesh, and stored for chemical analyses. We measured pressed-leaf spectra on the

former portion prior to mounting, and ground-leaf spectra and chemical traits on the latter. Although these

are distinct subsets of leaves, we treat them as identical in their trait values and spectra under the

assumption that they are very close in canopy position and unlikely to differ systematically. A minimum

of one representative pressed specimen of each species at each site and date is prepared as a specimen at

the Marie-Victorin Herbarium in Montréal, Québec, Canada.

Spectral measurements

For the Beauchamp-Rioux, Boucherville, and Warren projects, we measured fresh-leaf directional-

hemispherical reflectance spectra on each sample in 2018 using an HR-1024i spectroradiometer outfitted

with a DC-R/T integrating sphere from Spectra Vista Corporation (SVC; Poughkeepsie, NY, USA). We

measured spectra on the adaxial surface of six leaves or, for small or narrow leaves, single-layer leaf

arrays following the protocol in Noda et al. (2013) as adapted for our instrument (Laliberté & Soffer

2018a; Laliberté and Soffer 2018b). For the Dessain project, we measured fresh-leaf reflectance spectra in

2017 using a FieldSpec 4 spectroradiometer equipped with an RTS-3ZC integrating sphere from

Analytical Spectral Devices, Inc. (ASD; Boulder, CO, USA) following the leaf spectroscopy protocol

from the Global Airborne Observatory (https://gao.asu.edu/spectranomics-protocols). We collected

spectra from a variable number of leaves or arrays per sample—six in most cases, but as few as three or

as many as eighteen from certain samples. For all projects, each reflectance measurement was calibrated

against a white Spectralon 99% reflectance panel and corrected for stray light.

For all projects, we measured reflectance spectra on the adaxial surface of pressed, intact leaf samples

using a PSR+ 3500 spectroradiometer with a leaf clip foreoptic (Spectral Evolution, Haverhill, MA)

between six months and three years after collection. We calibrated reflectance against a white Spectralon

reflectance panel and measured 3-7 spectra per sample. Although all samples were pressed in a plant

press, some nevertheless dried in a way that created an uneven leaf surface. We noted any visible

discoloration that samples underwent during drying or storage, including browning or blackening. In

general, we aimed to avoid uneven or discolored areas when measuring spectra, although it was

sometimes not entirely possible when discoloration was pervasive.

For all projects, we also measured ground-leaf spectra using the PSR+ 3500 spectroradiometer with a

benchtop reflectance probe foreoptic that pressed loose powder into a smooth, even pellet. For each

sample, we added at least 0.6 g of leaf powder into a sample tray; in preliminary tests, adding additional

material did not change the spectra, suggesting that transmittance was close to zero. We calibrated

reflectance against a white Spectralon reflectance panel placed in an identical sample tray. We measured

three spectra per sample on the same pellet, turning the sample tray 120° between the first two spectra,

and loosening and mixing the powder before reshaping the pellet between the second and third spectra. A

small number (<15) of specimens could not be measured because there was not enough ground leaf

material.

Our spectral processing pipeline varied by the type of data. Although the actual spectral resolution was

more variable and coarser, the ASD Field Spec 4 and Spectral Evolution PSR+ 3500 spectrometers'

software automatically resampled the spectra to 1 nm resolution and interpolated over the overlap region

between sensors (Schweiger & Laliberté 2020). For data from the SVC HR-1024i spectrometer, we

performed these steps using linear interpolation. For pressed-leaf spectra only, we detected jumps in

reflectance at the sensor overlap region and removed them using the function *match_sensors()* in *R*

package *spectrolab v. 0.0.10* (Meireles et al. 2017). Next, we averaged all spectra for a given sample and

tissue type (fresh, pressed, or ground). Spectra measured with integrating spheres tend to have greater

noise, especially at the ends. To reduce noise in fresh-leaf spectra, we applied a Savitzky-Golay filter

using *R* package *signal 0.7.6* (Signal Developers, 2013), with varying order and length: order 3 and length

21 from 350-715 nm, order 3 and length 35 from 715-1390 nm, order 3 and length 75 from 1390-1880

nm, and order 5 and length 175 from 1880-2500 nm. Finally, we trimmed all spectra to 400-2400 nm,

removing the extremes where sensors show greater noise. We did all spectral processing in *R v. 3.6.1* (R

Core Team 2020) using package *spectrolab v. 0.0.10* (Meireles et al. 2017).

Trait measurements

We performed all trait measurements excluding petioles, but including the rachis for compound leaves,

since the rachis is functionally analogous to the midrib of a simple leaf. We measured the following leaf

structural and chemical traits: Leaf mass per area (LMA; kg m^{-2}), leaf dry matter content (LDMC; mg g^{-1}),

leaf nitrogen per unit mass (N_{mass}), leaf carbon per unit mass (C_{mass}), and carbon fractions (soluble cell

contents, hemicellulose, cellulose, and lignin) on a mass basis.

We weighed six leaves per sample for fresh mass shortly after collection. We then rehydrated them in a

plastic bag with a damp paper towel for at least 12 h and scanned and weighed them for rehydrated mass

and area. Lastly, we dried them for 72 h in a drying oven at 65°C before reweighing for dry mass. We

calculated LDMC as total dry mass divided by total fresh mass and LMA as total dry mass divided by

total rehydrated area (Laliberté 2018).

We analyzed ground bulk tissue samples for N_{mass} and C_{mass} using a Vario MICRO Cube combustion

analyzer (Elementar, Langenselbold, Germany; Ayotte et al. 2019). We analyzed the same samples for

carbon fractions using an ANKOM 2000 Fiber Analyzer (Ankom Technology, Macedon, New York,

USA) to perform a sequence of digestions (Ayotte & Laliberté 2019). The first digestion in a heated

neutral detergent washes off soluble cell contents. The second digestion in a heated acidic detergent

removes hemicellulose and bound proteins—which, for simplicity, we just refer to as hemicellulose. The

third digestion in 70% sulfuric acid removes cellulose, leaving behind lignin and recalcitrant compounds.

We ashed all samples and used ash content as a correction factor to calculate the ash-free lignin fraction.

Scoring discoloration

We inspected each pressed specimen visually to note signs of discoloration that could have occurred in

preparation or storage. Typical forms of discoloration include blackening or browning or the development

of a silvery or whitish finish on the leaf surface. Based on our descriptive notes, we scored each leaf on a

discrete scale from 0 to 4. A score of 0 indicates no noticeable discoloration. Scores 1 through 4 indicate

increasing discoloration, from 1 (either <5% blackening/browning or development of a silvery finish to

the leaf) to 4 (>80% blackening/browning).

PLSR modeling for trait estimation

We used a PLSR modeling framework to link the fresh-, pressed-, and ground-leaf spectra to the traits we

measured. PLSR is well-suited to handle leaf spectral datasets, which have large numbers of collinear

predictors, because it projects the spectral matrix to a smaller number of orthogonal latent components in

a way that maximizes the ability to predict the response variable. Moreover, it does not require assuming

that the variables predicted are measured without error. We implemented PLSR modeling using *R*

package *pls v. 2.7.1* (Mevik et al. 2019). We conducted analyses separately for fresh-, pressed-, and

ground-leaf spectra, using the full spectrum (400-2400 nm) to predict each trait. Some previous studies

have restricted the ranges of wavelengths used in prediction to those known or assumed to contain

features relevant to a given trait (e.g. Serbin et al. 2014). We avoided this approach because the features

important for prediction may vary depending on whether the tissue is fresh, pressed, or ground. In

preliminary analyses, restricting the range of wavelengths, using first or second derivatives of reflectance,

or calculating pseudoabsorbance of ground tissue (Norris et al. 1976) made little difference for predictive

accuracy.

Our methods for model calibration and validation largely follow Serbin et al. (2014). First, we divided the

data into calibration (80%) and validation (20%) datasets, stratified by project. We began by fitting a

model for each trait on the full calibration dataset, using 10-fold cross-validation to select the optimum

number of components. For each trait, we selected the smallest number of components for which the

cross-validation root mean squared error of prediction (RMSEP) was within one standard deviation of the

global minimum RMSEP at any number of components. This number of components—a different number

for each trait (10-27)—was used in further analyses that aimed to predict traits in the independent

validation dataset. We used the variable importance in projection (VIP) metric calculated for calibration

models to see which parts of the spectrum were most important for predicting each trait (Wold et al.

1994).

To test how well we could predict traits from the independent validation dataset, we developed a new set

of models for each trait. We did a jackknife analysis by repeatedly ($100\times$) dividing the 80% calibration

data further into random 70% training and 30% testing subsets. For each trait, we trained models on the

70% using the previously determined optimal number of components and predicted the remaining 30% to

get a distribution of model performance statistics (R^2, RMSE). The distribution of model predictions

reveals how sensitive model performance is to varying sets of training and testing data.

Next, we applied the 100 jackknife models for each trait to the 20% independent validation dataset. For

each sample in the validation dataset, we generated a distribution of predictions from the 100 models.

After averaging predictions of each validation sample's traits from the 100 models, we quantified model

performance using R^2 and root mean squared error (RMSE) using OLS regression. In all cases, we

calculated RMSE relative to the 1:1 line rather than the observed-predicted regression line, as

recommended by Piñeiro et al. (2008) to account for bias. We also report the RMSE as a percentage of

the sample data range (%RMSE), following Feilhauer et al. (2010). For each trait, we also tested whether

the magnitude of residuals (observed minus predicted) in the validation dataset varied among leaves

scored with different amounts of discoloration. We performed all statistical analyses in *R* v. 3.6.1.

PLS-DA modeling for species classification

We tested the ability to use fresh-, pressed-, and ground-leaf spectra in species classification using partial

least-squares discriminant analysis (PLS-DA; Barker and Rayens 2003). We took spectra from the ten

most common species in our dataset: *Acer rubrum* L., *A. saccharinum* L., *A. saccharum* Marshall, *Agonis*

flexuosa, *Betula papyrifera* Marshall, *B. populifolia* Marshall, *Fagus grandifolia* Ehrh., *Populus*

grandidentata Michx., *Populus tremuloides* Michx., and *Quercus rubra* L. All are deciduous trees except

A. flexuosa, which is evergreen. Each of these species was represented by at least 20 specimens (~480

total).

We divided the full dataset into 60% calibration and 40% validation datasets. In the *R* library *caret v.*

6.0.84 (Kuhn 2020), we trained models on the calibration dataset using 10-fold cross-validation.

Imbalanced groups within the calibration dataset can bias classification algorithms (Sun et al. 2009;

Lindström et al. 2011), so prior to the cross-validation procedure, we upsampled within less-represented

species classes at random with replacement to the point that each species had representation equal to the

best-represented species within the calibration dataset. We chose the number of PLS components during

cross-validation by maximizing Cohen's kappa (κ). This statistic describes the agreement between the

true and predicted species identities while accounting for the probability of agreement by chance. To

reduce the chance of overfitting due to our upsampling procedure, we capped the number of components

at the optimal number we would have selected by running the same procedure, but instead downsampling

classes to the size of the smallest class. (For example, if cross-validation with downsampling showed

maximum κ at *n* components, we would allow the cross-validation with upsampling to select a model

with no more than *n* components.) We then applied the cross-validated model (with upsampling) to the

validation dataset and summarized PLS-DA model performance using raw classification accuracy and κ.

Results

Patterns in traits and reflectance spectra

We saw large variation among samples in each of our target traits, ranging from 1.4-fold variation in C_{mass} to 23-fold variation in lignin (Table 1). The range of most traits in our dataset is generally narrower than in the global TRY dataset (Kattge et al. 2020), excluding the lower part of the range for soluble carbon, the upper part for remaining carbon fractions, and both extremes for remaining traits. Trees tended to have higher LDMC, C_{mass}, and lignin than other growth forms, consistent with prior research (Ma et al. 2018); grasses had particularly high hemicellulose and cellulose and low lignin content. Forbs generally had high N_{mass}. Some of the trait variation was driven by projects that occupied particular portions of trait space; for example, *A. flexuosa* in the WCS project tended to have particularly high LMA and C_{mass} and low N_{mass}.

The coefficient of variation (CV) of fresh-leaf reflectance was usually largest where reflectance itself was lowest—namely, in the visible range (400-700 nm) and the water absorption features (the largest of which are centered around 1450 and 1930 nm; Fig. 1A). Within the visible range, this pattern did not always hold. The CV remained high along the green reflectance hump (~530-570 nm) even as reflectance increased; it also has a local minimum at 682 nm and a local maximum along the red edge at 697 nm. These patterns may be a consequence of chlorophyll's absorbance spectrum. Where chlorophyll absorbs most, as at 660-685 nm, leaf absorptance may be nearly saturated regardless of chlorophyll content. Where chlorophyll absorbs less strongly, as along the green hump and red edge, leaf absorptance is much more sensitive to chlorophyll content (Curran 1989; Gitelson et al. 1996; Gitelson & Merzlyak 1997).

Median reflectance across the entire spectrum was higher for pressed leaves than fresh leaves, as expected based on changes in water content and leaf structure (Carter 1991; Fig. 1B). Indeed, water absorption features largely disappeared in pressed leaves. The red edge became less sharp, such that the global

maximum of median reflectance was at a longer wavelength (954 nm vs 872 nm for fresh-leaf spectra).

Mean reflectance again drove most variation in the CV, and a small number of specimens, primarily the

grasses *Phragmites australis* and *Phalaris arundinacea*, had much higher visible reflectance than all other

specimens. The CV of pressed-leaf reflectance once again increased into the short-wave infrared (SWIR)

range as mean reflectance declined, although this increase was less pronounced than for fresh leaves due

to the near-absence of water absorption.

Ground-leaf reflectance was higher than pressed-leaf reflectance across nearly the entire spectrum (except

400-425 nm) because transmittance was nearly zero (Fig. 1C). The red edge was even more strongly

blunted than for pressed leaves, with the global maximum of median reflectance occurring at an even

longer wavelength (1313 nm). The CV again was inversely related to mean reflectance, although the rise

in the SWIR was less pronounced than among pressed-leaf spectra in part because the standard deviation

of reflectance was low.

PLSR modeling for trait estimation

In general, our models predicted traits in the independent validation dataset with moderate-to-high

accuracy (Tables 1, 2, 3; Figs. 2 and 3). The optimal number of components selected to predict each trait

was between 10 and 27, but was always higher for a given trait in ground-leaf models than the others.

Fresh-leaf models were best for predicting structural or water-related traits, including LMA and LDMC.

On the other hand, ground-leaf models were best for predicting chemical traits, including soluble cell

contents, hemicellulose, cellulose, lignin, N_{mass}, and C_{mass} (validation $R^2 = 0.736$-0.954, compared to

0.513-0.708 for fresh leaves and 0.586-0.821 for pressed leaves). Statistics from jackknife analyses show

that model performance was most variable and dependent on the training/testing datasets for traits that

were predicted less accurately (Figs. S1, S2, and S3).

We tested for relationships between discoloration and the magnitude (absolute value) of the residuals for

each trait. Our independent validation dataset contained only one specimen scored at a 4 for discoloration.

After removing this specimen as a potential outlier with undue influence on the slope estimates, there was

no correlation between the magnitude of residuals and discoloration for any trait ($p > 0.05$).

For all traits except LMA, the VIP metric for fresh-leaf spectra showed a global maximum between 710

and 720 nm (Fig. 4A)—wavelengths slightly longer than the typical inflection point of the red edge

between the visible and near-infrared (NIR) regions (Gitelson et al. 1996). Many traits, especially LDMC,

N_{mass}, and C_{mass}, show high VIP also across the wavelengths ~530-570 nm, corresponding to the green

hump. Bands in the NIR range were less important for predicting most traits than much of the visible

range. The SWIR range was generally important for predicting LMA, and many other traits showed

several local peaks of importance, most prominently at about 1880 nm, but also near 1460 and 1720 nm.

For predicting traits from pressed-leaf spectra, the general trend held that visible reflectance and several

band ranges in the SWIR were important for predicting most traits, while NIR reflectance was less

important (Fig. 4B). In fresh-leaf models, LMA had particularly low VIP in the visible range and high

VIP and in the SWIR, but in pressed-leaf models, it fell much more in line with other traits. The red edge

peak of importance for most traits was near 705 nm, and another prominent local maximum of importance

lay close to 1920 nm. We saw broadly similar patterns in VIP from ground-leaf spectra (Fig. 4C).

PLS-DA modeling for species classification

Our PLS-DA analyses showed that models using pressed- and ground-leaf spectra showed near-perfect

performance at classifying species, with misclassifications only among species within the same genus

(Fig. 5). Models using fresh-leaf spectra were slightly worse but still showed excellent performance. The

optimal fresh-leaf model, which had 29 PLS components, correctly predicted the taxonomic identity of

170 of the 189 samples in the training dataset ($\kappa = 0.884$; $p < 0.0001$; Fig. 5). The best pressed-leaf

model, which had 29 PLS components, correctly predicted 184 of the 188 samples ($\kappa = 0.975$; $p <$

0.0001).The best ground-leaf model, which had 30 PLS components, correctly predicted 183 out of 189

samples ($\kappa = 0.963$; $p < 0.0001$). In all cases, a majority of misclassifications were between congenerics.

Discussion

We show that we could accurately estimate a wide range of leaf functional traits for 70 woody and

herbaceous species from reflectance spectra of fresh leaves, pressed leaves, and ground leaf material

(Tables 1 and 2; Figs. 2 and 3). Importantly, using pressed-leaf spectra may provide an accurate, non-

destructive method for estimating the traits of herbarium specimens. Ground-leaf estimates were most

accurate for chemical traits—consistent with the success of such models in prior research (Serbin et al.

2014)—and fresh-leaf estimates were most accurate for structural or water-related traits. Pressed-leaf

estimates may represent a good compromise in allowing both classes of traits to be estimated with

intermediate but quite high accuracy. These results show that pressed-leaf spectra can provide an

integrative measure of leaf phenotypes, much like fresh-leaf spectra (Cavender-Bares et al. 2017). We

also showed that pressed-leaf spectra could be used to classify species as well as ground-leaf spectra and

better than fresh-leaf spectra.

Our findings about which kind of tissue was best for predicting each trait supported our predictions. For

estimating chemical traits, ground-leaf spectra showed strong performance, likely resulting from the way

that grinding homogenizes the lamina and removes the potentially confounding influence of leaf structure.

Pressed-leaf spectra showed intermediate performance in estimating such traits because, like ground

leaves, they lack the major water absorption features that mask the smaller absorption features of other

compounds in the SWIR range (Peterson et al. 1988; Ecarnot et al. 2013). On the other hand, these same

reasons may explain why pressed- and (especially for LMA) ground-leaf spectra performed worse at

predicting LMA and LDMC. Indeed, it may seem perplexing that we could succeed at all in predicting

LMA from ground-leaf spectra, or LDMC from pressed- or ground-leaf spectra. The ability to estimate

these traits must not be a result of optical features driven by the traits themselves. Instead, we suggest that

we sense these traits via their correlations with traits whose optical signatures are not destroyed by

pressing or grinding. This kind of effect—a "constellation effect" (*sensu* Chadwick and Asner 2016;

Nunes et al. 2017)—has been invoked to explain the ability to estimate other traits like leaf micronutrients

(Nunes et al. 2017) and $\delta^{15}N$ (Serbin et al. 2014) that do not have strong absorption features in the

measured range of wavelengths. However, models that rely on such constellation effects would be less

robust in situations where such trait covariance breaks down.

The performance of pressed-leaf models in this study was comparable to the successes obtained here and

elsewhere using fresh- and ground-leaf spectra. For example, the ground-leaf models in Serbin et al.

(2014) had a validation RMSE of 1.4 and 2.4 for cellulose and lignin. In comparison, our pressed-leaf

models achieved an RMSE of 1.42 and 2.16 for these traits (Table 2). On the other hand, our pressed-leaf

model for N_{mass} performed worse than that in Serbin et al. (2014; validation RMSE = 0.13 vs 0.247), but

better than the fresh-leaf models in Doughty et al. (2011; RMSE = 0.45). Our models for LDMC from

pressed leaves had a lower RMSE (33.8 mg g^{-1}) than Streher et al. (2020) obtained from fresh leaves (53

mg g^{-1}). Our models for LMA also had an RMSE (0.010 kg m^{-2}) lower than full-range fresh-leaf models

from the literature, including Asner et al. (2011b; 0.015 kg m^{-2}), Doughty et al. (2011; 0.019 kg m^{-2}),

Serbin et al. (2019; 0.015 kg m^{-2}), and Streher et al. (2020; 0.051 kg m^{-2}). Although these studies sampled

from different kinds of vegetation with different ranges of trait values, such comparisons establish that

pressed-leaf models are not categorically worse than ground- or fresh-leaf models, and in fact are often

better. It may depend on the research question whether the error obtained from these models is acceptable,

but we suggest that it would be for a wide range of questions concerning large-scale ecological or

evolutionary patterns that encompass a wide range of trait variation.

The VIP metric helps us understand what features allow these models to work, but interpreting its patterns

can be a challenge because reflectance at a given band usually cannot be understood in terms of a single

chemical or structural trait. Fresh-leaf VIP for most traits had peaks in the visible and SWIR ranges, with

a global maximum at 705 nm. Streher et al. (2020) found a similar peak for LMA and LDMC, among

others scattered primarily in the SWIR (Fig. 4A). The importance of red edge reflectance (particularly

700-725 nm) may owe to this region's sensitivity to both chlorophyll content and leaf structure (Gitelson

et al. 1996; Richardson et al. 2002). Much of the visible range was proportionally even more important

for pressed- and ground-leaf models, and the NIR was less important (except for LMA; Fig. 4B-C). There

are multiple small peaks in the SWIR, most notably one at 1920 nm that appears similar to a VIP feature

in Serbin et al. (2014). Reflectance at 1920 nm is often associated with absorption by water in fresh

tissue, but any causal link to water content is unlikely because the pressed and ground leaves were dried

in preparation; whatever water might remain comes only from the air. Bands near 1920 nm include broad

absorption features for many components of dry matter, including protein, cellulose, lignin, and starch,

any or all of which may contribute to the importance of this region (Curran et al. 1989; Fourty et al.

1996).

With some exceptions, the VIP metric showed that the same bands are often important for predicting

different traits (Fig. 2). This might be taken as an artifact of trait covariance, but surprisingly, across the

whole dataset, many traits covaried only weakly. While soluble cell contents, hemicellulose, and cellulose

all showed correlations, positive or negative (R^2 = 0.463-0.791), the others often did not. For example,

while N_{mass}, C_{mass}, LMA, and LDMC all showed broadly similar patterns of pressed-leaf VIP across the

spectrum, including peaks at 705 and 1920 nm, they covaried weakly with each other (R^2 = 0.029-0.380).

The fact that some of the same regions of the spectrum were important for predicting traits may not be

purely as an artifact of trait covariance. The VIP statistic is just a heuristic, and does not show the

direction in which a band's reflectance alters trait estimates; the same bands may matter for different traits

in different ways, like the red edge, which is affected by both chlorophyll and leaf structure.

PLS-DA models showed that fresh-, pressed-, and ground-leaf spectra alike could be used to classify

species with near-perfect accuracy (>96%) for pressed or ground leaves, and excellent (>89%) accuracy

for fresh leaves, much as in work on tropical flora (Durgante et al. 2013; Lang et al. 2015; Lang et al.

2017; Prata et al. 2018). In contrast to prior work that deliberately selected many congenerics, our most

common species were not all closely related. Among the samples misidentified, most were mistaken for

congenerics (Fig. 5), particularly for pressed and ground specimens, which implies that related species are

more spectrally similar (Schweiger et al. 2018; Meireles et al 2020a; Schweiger et al. 2021). Hence,

discriminating more closely related species might pose a greater challenge and lead to a higher error rate.

Nevertheless, our analysis reinforces that spectral models can often distinguish among species or higher-

level taxonomic groups (Meireles et al. 2020b).

The fact that pressed-leaf models showed the highest classification accuracy (>97%) is noteworthy, since

pressed specimens are the standard way that field botanists preserve plant samples for later identification.

This outcome is surprising since pressed leaves were never best for predicting any single trait. Although

all three models performed well, we conjecture that pressed-leaf models have an advantage because they

retain aspects of leaf structure that vary among species while, more importantly, also revealing the

absorption features of potentially distinctive compounds in the SWIR range. While water content can only

vary in one dimension, there are many macromolecules and metabolites whose absorption features in this

range might allow finer discrimination of species. The partial preservation of leaf structure, at least

relative to ground leaves, might allow for another dimension along which species can be distinguished.

The future of spectroscopic trait estimation

Although trait predictions from spectral models are not perfect, they have a few advantages over

conventional trait measurements: they (1) are potentially non-destructive; (2) are fast and require little

training, especially compared to measuring multiple traits; and (3) have very low marginal cost, despite

the high capital cost of buying a spectrometer. Because of these advantages, the widespread adoption of

leaf spectroscopy could make it easier to address questions that require large datasets of functional traits.

But researchers who are attracted to reflectance spectroscopy by the promise of quick, cheap trait

estimates may be deterred if researchers must each make their own trait prediction model, which would

imply that they must still measure traits conventionally on a subset of their samples. For that reason, we

should aim for spectral models to be general.

This aim is not easy to achieve: for multiple reasons, a model trained on one spectral dataset may not

always make good trait predictions from a second dataset. The reasons may include, but are not limited to,

differences between the training and target data sets in: (1) their representation of plant traits and optical

properties; (2) plant sample preparation prior to spectral measurements; and (3) the instruments and

foreoptics used. Here we discuss these issues one by one; in the following section, we describe some

concerns particular to herbarium specimens.

First, the ranges and the covariance structure of the traits may differ among the datasets. For example, if a

researcher wants to apply a model trained on boreal conifers alone to tropical broadleaf species, the

conifer model may have to extrapolate well beyond its range. A general trait model must include a broad

range of functional variation, which poses a logistical challenge because any one site only encompasses a

fraction of global trait variation. Moreover, even portions of trait space that are represented within the

training dataset may be poorly predicted—for example, Streher et al. (2020) found that models for LMA

among plants in the seasonally dry Neotropics performed poorly above 300 g m^{-2}. A fully general model,

if such a thing is possible, would need to represent the full range of leaf structure and chemistry.

Another class of issues with model generality concerns sample preparation before spectral measurements.

For example, ground-leaf spectra vary based on the size of the particles in the ground tissue (Foley et al.

1998). Other decisions, such as the choice to include or exclude petioles in the ground tissue sample,

could alter the spectrum and compromise trait estimation from models trained on samples prepared

another way.

Yet another class of issues concerns spectrometers and their foreoptics. Spectra of fresh leaves can be

measured with different foreoptics, including integrating spheres, contact probes, or leaf clips. Integrating

spheres are designed for making measurements that are nearly independent of the angle of the light source

or the sensor relative to the sample. They also allow the use of light traps to absorb nearly all transmitted

light; in contrast, most leaf clips or contact probes use black backgrounds that invariably reflect a small

fraction of light. These features make integrating spheres likely to produce more consistent measurements

between instruments or replicate samples (Petibon et al. 2021). Finally, integrating spheres also allow

researchers to measure both reflectance and transmittance spectra. On the other hand, leaf clips and

contact probes are much faster to use, often produce higher signal-to-noise ratios, and allow measurement

of angle-dependent (anisotropic) reflectance when it is desired. The data from these foreoptics are not

often directly comparable: spectra measured with leaf clips or contact probes tend to show higher

reflectance, probably due to the effects of the viewing and illumination angles (Potůčková et al. 2016;

Hovi et al. 2018). Spectrometers and their software may also differ in their sensors and techniques for

processing spectra.

In this study, we use integrating spheres with fresh leaves, but leaf clips with pressed specimens; using an

integrating sphere requires placing specimens either inside the sphere or in one of the ports, each of which

could damage the delicate pressed specimens. Such logistical constraints could limit the choices that

researchers can make in data collection. In theory, our use of different foreoptics for measuring fresh-,

pressed-, and ground-leaf spectra could confound our comparisons of model accuracy. But given that we

calibrated the models separately, we argue that the physical changes caused by pressing or grinding are

more important than the foreoptic type for explaining which kind of spectra work best for each trait in this

study. Some studies (e.g. Serbin et al. 2019; Meireles et al. 2020) have successfully built models that

combine data collected with multiple kinds of instruments—and indeed, here we use two different models

of spectrometers for fresh-leaf estimates. Assuming that these kinds of technical challenges can be

overcome—for example, by developing tools to reconcile measurements with different setups, as in

Meireles et al. (2020)—we show that there are good reasons to be confident about building general

models to estimate traits from a wide variety of plants.

The challenges of reducing the error and improving the generality of trait estimates are not unique to

reflectance spectroscopy (Clark & Kellner 2012). Conventional wet lab trait measurements, too, are prone

to varying amounts of error, including those used to train the spectral models. For example, different

methods of measuring lignin yield different results, and some have high variance between replicate

samples (Brinkmann et al. 2002). As with any other technique, the goal for spectroscopic trait estimation

is to improve model accuracy and generality as much as they can be jointly improved without making the

technique harder to use. The creation of open libraries for spectral data (like EcoSIS; https://ecosis.org/)

and spectral models (like EcoSML; https://ecosml.org/) will contribute to this goal.

Some of the same concerns apply to species classification models, even though they are likely to remain

unavoidably specific to particular applications. For example, a species classification model must include

strong representation in the training dataset for all the species that one might encounter. Researchers may

also want to use existing data, as from open libraries, to train models, which would require resolving

concerns about sampling—including phenotypic plasticity—as well was the potential influence of

instrumentation and measurement protocols (Meireles et al. 2020).

What should herbarium managers do?

If spectroscopic models can reliably estimate traits from pressed-leaf spectra, researchers may be able to

use herbarium specimens non-destructively to address core questions in functional trait ecology and

evolution, including studies of historical changes in trait composition and diversity (Lang et al. 2018).

This technique may allow researchers to reduce their reliance on measuring traits or reflectance spectra

through intensive, logistically challenging field campaigns in order to fill in trait datasets, speeding up

progress in plant functional ecology. It could also keep researchers from having to do destructive and

time-consuming conventional trait measurements on herbarium specimens. Likewise, spectroscopy may

be able to assign species identifications to specimens that are otherwise hard to identify.

However, measuring and interpreting spectra from herbarium specimens presents additional challenges.

Perhaps the most concerning is the potential for specimens' optical and chemical properties to degrade

during preparation or storage, especially light-sensitive photosynthetic pigments. Such degradation could

make it hard to distinguish secular trends in the traits of living plants over time from changes in storage.

Even in this study, where no specimens were collected before 2017, many underwent noticeable changes

in color, including browning or blackening; ~12% were scored at 2 or higher. Certain preparation and

storage conditions could accelerate changes in optical properties. For example, when collecting herbarium

specimens in the wet tropics, researchers often treat leaves with an alcohol solution as a preservative

(Blanco et al. 2006). Such treatments could discolor leaves—as could the spoilage that would occur in

their absence. Exposure to high light, temperature, or humidity could also contribute to discoloration,

which underscores the importance of developing and using best practices for preparation and storage.

Some species may also be more susceptible than others; for example, in our dataset, ~42% of *Populus*

grandidentata specimens were scored at 2 or higher, but no *Betula papyrifera* specimens were.

Whether such degradation could affect trait estimation is an open empirical question. We find no evidence

that it does, even for badly discolored specimens, but our specimens were only collected as early as 2017,

while ecologists may want to use specimens collected decades ago. Long-term studies of the same

specimens over time—especially with sets of specimens subjected to varying preparation and storage

techniques—would help establish how optical properties tend to change, and may establish predictable

patterns of degradation over time that can be corrected to estimate the original spectrum. In any case, it

seems likely that estimating traits in old or degraded specimens will require including equally old or

degraded specimens in the dataset used to build the trait model.

There may be other challenges besides sample degradation. Measuring reflectance usually requires a

black absorbing background so that any transmitted light is not reflected back into the sensor. However,

many herbarium specimens are glued to a paper backing. The availability of loose whole leaf material

would make measuring reflectance spectra much easier. However, researchers may be able to develop and

validate methods to measure and correct for reflectance from the paper backing, a potential subject for

future research.

If taking spectral measurements during field collections is too challenging, the best way to incorporate

spectroscopy into the operation of a working herbarium may be to measure the spectra of recently pressed

and collected specimens that have not yet been mounted—ideally, immediately after drying. The

spectrum could then be linked to the herbarium record for future analyses and made available to

download, together with metadata about the instrument, foreoptic, specimen preparation, and date of

spectral measurements. While the ideal situation would be for specimen collectors to gather abundant

loose tissue for trait (or spectral) measurements (Heberling and Isaac 2017), doing so is not always

feasible. Linking spectra to herbarium records may also require new developments in biodiversity

informatics, since the hyperdimensionality of the reflectance spectrum could make it hard to

accommodate within existing standards such as the Darwin Core (Wieczorek et al. 2012) or the ABCD

schema (Holetschek et al. 2012). Coordination between herbarium managers and researchers who use

reflectance spectroscopy could advance protocol development and build agreement about best practices

for measurement and curation (Heberling and Isaac 2017).

Validating the spectral estimation of traits from older herbarium specimens will require extensive further

research, but we take a first step towards this goal by showing that non-destructively measured pressed-

leaf spectra retain information about many leaf functional traits. This finding raises the possibility of

using herbaria as tools to understand plant functional trait ecology and evolution in a much more

comprehensive way. Our study has far-reaching implications for capturing the wide range of functional

and phenotypic information in the world's preserved plant collections and broadly advancing digital taxon

identification in collected specimens.

Acknowledgements

We performed this research on the ancestral and contemporary land of Native people: primarily the

Kanien'kehá꞉ka (Mohawk) and Omàmiwinini (Algonquin) First Nations for the Beauchamp-Rioux,

Dessain, and Boucherville projects, and the Noongar people for the Warren project. For permissions and

assistance, we thank le Jardin botanique de Montréal; la Station de biologie des Laurentides (SBL) de

l'Université de Montréal; l'Institut de recherche d'Hydro-Québec (IREQ); the National Capital

Commission for sampling at Mer Bleue Bog; la Société des établissements de plein air du Québec

(SÉPAQ) for sampling at Parc national d'Oka, Parc national du Mont-Saint-Bruno, and Parc national des

Îles-des-Boucherville; and the Parks and Wildlife Service of Western Australia for sampling in

D'Entrecasteaux National Park. We would like to thank Antoine Mathieu, Alexandra Massey, Florence

Blanchard, Elisabeth Hardy-Lachance, Zachary Bélisle, Sandra Jooty, Myriam Cloutier, Fabien

Cichonski, Madeleine Trickey-Massé, Aurélie Dessain, Xavier Guilbeault-Mayers, Jocelyne Ayotte, and

especially Sabrina Demers-Thibeault for contributing to trait measurements. Trait and fresh-leaf spectral

data from Canadian and Australian sites were collected as part of the Canadian Airborne Biodiversity

Observatory (CABO), which is funded by NSERC Discovery Frontiers grant 509190-2017, as well as

NSERC Discovery grants (RGPIN-2014-06106, RGPIN-2019-04537) and a Canada Research Chair to

EL. Other spectral data were collected with support from the NSF/NASA under DEB #1342778, an

International Thesis Research Travel Grant from the University of Minnesota's Graduate School, and a

grant distributed through an NSF-funded Research Coordination Network (DEB-1745562). S.K. was

supported by an NSF Graduate Research Fellowship (Grant No. 00039202) and a UMN Doctoral

Dissertation Fellowship. R. B.-R. was supported by NSERC through the Canada Graduate Scholarships—

Master's program.

Tables

Table 1. Summary statistics of PLSR model calibration and validation from fresh-leaf spectra. %RMSE is

calculated as RMSE divided by the range of data within either the training data set (calibration) or the

testing data set (validation), each of which may be smaller than the range reported in the complete data

set. Global ranges are derived from the TRY database (Kattge et al. 2020; accessed 13 April 2021),

omitting data with error risk greater than 3. Carbon fractions (solubles, hemicellulose, cellulose, and

lignin) are each represented by <1250 datapoints in TRY, which may curb the estimates of the global

range.

	Range	Global range	No. comps	R^2		RMSE		%RMSE	
				Cal.	Val.	Cal.	Val.	Cal.	Val.
Solubles (%)	38.5-88.9	20.0-74.6	13	0.526	0.457	5.63	5.92	11.2	15.2
Hemicellulose (%)	2.77-34.1	8.71-45.0	10	0.517	0.479	3.36	3.43	10.7	14.6
Cellulose (%)	5.19-25.8	7.65-42.6	24	0.644	0.597	1.90	2.14	9.46	11.0
Lignin (%)	0.918-21.5	2.27-32.4	10	0.513	0.498	2.34	2.56	11.4	13.0
C_{mass} (%)	39.5-53.6	25.5-60.3	12	0.708	0.715	1.16	1.21	8.67	10.0
N_{mass} (%)	0.883-5.62	0.14-7.27	11	0.689	0.695	0.328	0.338	7.92	7.32
LMA (kg m^{-2})	0.0233-0.217	0.00866-0.354	17	0.955	0.950	0.00783	0.00818	4.03	4.58
LDMC (mg g^{-1})	159-573	68.2-857	19	0.877	0.800	22.5	28.2	5.50	7.42

668 Table 2. Summary statistics of PLSR model calibration and validation from pressed-leaf spectra. Data

669 ranges in the complete data set are reported in Table 1.

	No. comps	R^2		RMSE		%RMSE	
		Cal.	Val.	Cal.	Val.	Cal.	Val.
Solubles (%)	12	0.690	0.657	4.57	4.54	9.07	11.7
Hemicellulose (%)	12	0.711	0.666	2.64	2.65	8.42	11.3
Cellulose (%)	18	0.803	0.701	1.42	1.63	7.05	8.78
Lignin (%)	13	0.586	0.621	2.16	2.24	10.5	11.4
C_{mass} (%)	17	0.815	0.844	0.935	0.881	7.00	7.31
N_{mass} (%)	16	0.821	0.744	0.247	0.312	5.97	6.75
LMA (kg m^{-2})	10	0.924	0.916	0.0102	0.0102	5.24	5.72
LDMC (mg g^{-1})	19	0.738	0.701	33.8	34.9	8.16	9.20

671   Table 3. Summary statistics of PLSR model calibration and validation from ground-leaf spectra. Data

672   ranges in the complete data set are reported in Table 1.

	No. comps	R^2		RMSE		%RMSE	
		Cal.	Val.	Cal.	Val.	Cal.	Val.
Solubles (%)	24	0.806	0.809	3.66	3.54	7.26	9.12
Hemicellulose (%)	20	0.736	0.716	2.54	2.54	8.12	10.8
Cellulose (%)	27	0.912	0.914	0.916	0.941	4.56	4.83
Lignin (%)	23	0.764	0.812	1.64	1.57	7.98	8.01
C_{mass}	22	0.914	0.932	0.641	0.589	4.80	4.89
N_{mass}	23	0.954	0.966	0.123	0.114	2.98	2.47
LMA (kg m^{-2})	19	0.866	0.804	0.0136	0.0154	6.97	8.67
LDMC (mg g^{-1})	20	0.797	0.778	29.9	30.3	7.21	8.44

Figure captions

Fig. 1: Distributions of spectral reflectance and its coefficient of variation (CV) among fresh (left), pressed (middle), and ground (right) leaf samples. A solid black line connects the median reflectance at each wavelength, around which dark blue and light blue ribbons denote the middle 50% and 95%. The solid red line shows the coefficient of variation among spectra at each wavelength.

Fig. 2: Independent validation results for predictions of soluble cell contents, hemicellulose, cellulose, and lignin from fresh- (left), pressed- (middle), and ground-leaf (right) spectra. In each panel, a separate regression line is shown for each functional group, overlaid on top of the thick dashed 1:1 line. The error bars for each data point are 95% confidence intervals calculated from the distribution of predictions based on the model coefficients from the 100 jackknife iterations.

Fig. 3: Independent validation results for predictions of C_{mass}, N_{mass}, LMA, and LDMC from fresh- (left), pressed- (middle), and ground-leaf (right) spectra, displayed as in Fig. 3.

Fig. 4: The variable importance of projection (VIP) metric calculated based on fresh- (A), pressed- (B), and ground-leaf (C) models. Each panel shows carbon fractions (top) and traits related to structure or elemental composition (bottom). Abbreviations: sol = solubles, hemi = hemicellulose, cell = cellulose, lign = lignin.

Fig. 5: Confusion matrices for partial least-squares discriminant analysis from fresh- (left), pressed- (middle), and ground-leaf (right) spectra. Numbers on the diagonal represent correctly classified species, while off-diagonal numbers represent misclassified species; the row denotes the true species identity, while the column denotes the model's prediction.

Figures

702 **Fig. 1**

Fig. 2

707 **Fig. 3**

Fresh-leaf spectra | Pressed-leaf spectra | Ground-leaf spectra

Growth Form
- Forb
- Grass
- Shrub
- Tree
- Vine

710   **Fig. 4A**

722   **Fig. 4B**

734 **Fig. 4C**

746   **Fig. 5A**

748   **Fig. 5B**

752   **Fig. 5C**

C

References

American Society for Testing and Materials. (2006). *Standard Tables for Reference Solar*

Spectral Irradiances: Direct Normal and Hemispherical on 37° Tilted Surface (Standard G173-

06).

Anderegg, L. D. L., Berner, L. T., Badgley, G., Sethi, M. L., Law, B. E., & HilleRisLambers, J.

(2018). Within-species patterns challenge our understanding of the leaf economics spectrum.

Ecology Letters, *21*(5), 734–744.

Asner, G. P., Martin, R. E., Knapp, D. E., Tupayachi, R., Anderson, C., Carranza, L., Martinez,

P., Houcheime, M., Sinca, F., & Weiss, P. (2011a). Spectroscopy of canopy chemicals in humid

tropical forests. *Remote Sensing of Environment, 115*(12), 3587–3598.

Asner, G. P., Martin, R. E., Tupayachi, R., Emerson, R., Martinez, P., Sinca, F., Powell, G. V.

N., Wright, S. J., & Lugo, A. E. (2011b). Taxonomy and remote sensing of leaf mass per area

(LMA) in humid tropical forests. *Ecological Applications, 21*(1), 85–98.

Ayotte, J., Guilbeault-Mayers, X., & Laliberté, E. (2019). Measuring CN content in leaf samples

using Elementar Vario MICRO Cube. https://doi.org/10.17504/protocols.io.udces2w

Ayotte, J., & Laliberté, E. (2019). Measuring leaf carbon fractions with the ANKOM2000 Fiber

Analyzer. https://doi.org/10.17504/protocols.io.yinfude

Barker, M., & Rayens, W. (2003). Partial least squares for discrimination. *Journal of*

Chemometrics, 17(3), 166–173.

Blanco, M. A., Whitten, W. M., Penneys, D. S., Williams, N. H., Neubig, K. M., & Endara, L.

(2006). A Simple and Safe Method for Rapid Drying of Plant Specimens Using Forced-Air

Space Heaters. *Selbyana*, *27*(1), 83–87.

Brinkmann, K., Blaschke, L., & Polle, A. (2002). Comparison of Different Methods for Lignin

Determination as a Basis for Calibration of Near-Infrared Reflectance Spectroscopy and

Implications of Lignoproteins. *Journal of Chemical Ecology*, *28*(12), 2483–2501.

Carter, G. A. (1991). Primary and Secondary Effects of Water Content on the Spectral

Reflectance of Leaves. *American Journal of Botany, 78*(7), 916–924.

Castro-Esau, K. L., Sánchez-Azofeifa, G. A., Rivard, B., Wright, S. J., & Quesada, M. (2006).

Variability in leaf optical properties of Mesoamerican trees and the potential for species

classification. *American Journal of Botany, 93*(4), 517–530.

Cavender-Bares, J., Gamon, J. A., Hobbie, S. E., Madritch, M. D., Meireles, J. E., Schweiger, A.

K., & Townsend, P. A. (2017). Harnessing plant spectra to integrate the biodiversity sciences

across biological and spatial scales. *American Journal of Botany, 104*(7), 966–969.

Cavender-Bares, J., Meireles, J. E., Couture, J. J., Kaproth, M. A., Kingdon, C. C., Singh, A.,

Serbin, S. P., Center, A., Zuniga, E., Pilz, G., & Townsend, P. A. (2016). Associations of Leaf

Spectra with Genetic and Phylogenetic Variation in Oaks: Prospects for Remote Detection of

Biodiversity. *Remote Sensing*, *8*(3), 221.

Chadwick, K. D., & Asner, G. P. (2016). Organismic-Scale Remote Sensing of Canopy Foliar

Traits in Lowland Tropical Forests. *Remote Sensing*, *8*(2), 87.

Clark, D. B., & Kellner, J. R. (2012). Tropical forest biomass estimation and the fallacy of

misplaced concreteness. *Journal of Vegetation Science, 23*(6), 1191–1196.

Couture, J. J., Singh, A., Rubert-Nason, K. F., Serbin, S. P., Lindroth, R. L., & Townsend, P. A.

(2016). Spectroscopic determination of ecologically relevant plant secondary metabolites.

Methods in Ecology and Evolution, *7*(11), 1402–1412.

Curran, P. J. (1989). Remote sensing of foliar chemistry. *Remote Sensing of Environment*, *30*(3),

271–278.

Desmet, P., & Brouillet, L. (2013). Database of Vascular Plants of Canada (VASCAN): A

community contributed taxonomic checklist of all vascular plants of Canada, Saint Pierre

and Miquelon, and Greenland. *PhytoKeys, 25*, 55–67.

Diniz, É. S., Amaral, C. H., Sardinha, S. T., Thiele, J., & Meira-Neto, J. A. A. (2020).

Phylogenetic signatures in reflected foliar spectra of regenerating plants in Neotropical forest

gaps. *Remote Sensing of Environment*, 112172.

Doughty, C. E., Asner, G. P., & Martin, R. E. (2011). Predicting tropical plant physiology from

leaf and canopy spectroscopy. *Oecologia, 165*(2), 289–299.

Durgante, F. M., Higuchi, N., Almeida, A., & Vicentini, A. (2013). Species Spectral Signature:

Discriminating closely related plant species in the Amazon with Near-Infrared Leaf-

Spectroscopy. *Forest Ecology and Management*, *291*, 240–248.

Ecarnot, M., Compan, F., & Roumet, P. (2013). Assessing leaf nitrogen content and leaf mass

per unit area of wheat in the field throughout plant cycle with a portable spectrometer. *Field*

Crops Research, *140*, 44–50.

Ely, K. S., Burnett, A. C., Lieberman-Cribbin, W., Serbin, S. P., & Rogers, A. (2019).

Spectroscopy can predict key leaf traits associated with source–sink balance and carbon–nitrogen

status. *Journal of Experimental Botany, 70*(6), 1789–1799.

Feilhauer, H., Asner, G. P., Martin, R. E., & Schmidtlein, S. (2010). Brightness-normalized

Partial Least Squares Regression for hyperspectral data. *Journal of Quantitative Spectroscopy*

and Radiative Transfer, 111(12), 1947–1957.

Féret, J.-B., Gitelson, A. A., Noble, S. D., & Jacquemoud, S. (2017). PROSPECT-D: Towards

modeling leaf optical properties through a complete lifecycle. *Remote Sensing of Environment*,

193, 204–215.

Foley, W. J., McIlwee, A., Lawler, I., Aragones, L., Woolnough, A. P., & Berding, N. (1998).

Ecological applications of near infrared reflectance spectroscopy – a tool for rapid, cost-effective

prediction of the composition of plant and animal tissues and aspects of animal performance.

Oecologia, 116(3), 293–305.

Fourty, Th., Baret, F., Jacquemoud, S., Schmuck, G., & Verdebout, J. (1996). Leaf optical

properties with explicit description of its biochemical composition: Direct and inverse problems.

Remote Sensing of Environment, 56(2), 104–117.

Gitelson, A. A., Merzlyak, M. N., & Lichtenthaler, H. K. (1996). Detection of Red Edge Position

and Chlorophyll Content by Reflectance Measurements Near 700 nm. *Journal of Plant*

Physiology, *148*(3), 501–508.

Gitelson, A. A., & Merzlyak, M. N. (1997). Remote estimation of chlorophyll content in higher

plant leaves. *International Journal of Remote Sensing, 18*(12), 2691–2697.

Gold, K. M., Townsend, P. A., Chlus, A., Herrmann, I., Couture, J. J., Larson, E. R., & Gevens,

A. J. (2020). Hyperspectral Measurements Enable Pre-Symptomatic Detection and

Differentiation of Contrasting Physiological Effects of Late Blight and Early Blight in Potato.

Remote Sensing, 12(2), 286.

Grant, L. (1987). Diffuse and specular characteristics of leaf reflectance. *Remote Sensing of*

Environment, 22(2), 309–322.

Heberling, J. M., & Isaac, B. L. (2017). Herbarium specimens as exaptations: New uses for old

collections. *American Journal of Botany*, 104, 963–965.

Heberling, J. M., Prather, L. A., & Tonsor, S. J. (2019). The Changing Uses of Herbarium Data

in an Era of Global Change: An Overview Using Automated Content Analysis. *BioScience*,

69(10), 812–822.

Hietz, P., Turner, B. L., Wanek, W., Richter, A., Nock, C. A., & Wright, S. J. (2011). Long-

Term Change in the Nitrogen Cycle of Tropical Forests. *Science, 334*(6056), 664–666.

Holetschek, J., Dröge, G., Güntsch, A., & Berendsohn, W. G. (2012). The ABCD of primary

biodiversity data access. *Plant Biosystems, 146*(4), 771–779.

Hovi, A., Forsström, P., Mõttus, M., & Rautiainen, M. (2018). Evaluation of Accuracy and

Practical Applicability of Methods for Measuring Leaf Reflectance and Transmittance Spectra.

Remote Sensing, *10*(1), 25.

Jacquemoud, S., & Baret, F. (1990). PROSPECT: A model of leaf optical properties spectra.

Remote Sensing of Environment, 34(2), 75–91.

Jardine, E. C., Thomas, G. H., Forrestel, E. J., Lehmann, C. E. R., & Osborne, C. P. (2020). The

global distribution of grass functional traits within grassy biomes. *Journal of Biogeography*,

47(3), 553–565.

Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I. C., Leadley, P., Tautenhahn, S., Werner,

G. D. A., Aakala, T., Abedi, M., et al. (2020). TRY plant trait database – enhanced coverage and

open access. *Global Change Biology, 26*(1), 119–188.

Koski, M. H., MacQueen, D., & Ashman, T.-L. (2020). Floral Pigmentation Has Responded

Rapidly to Global Change in Ozone and Temperature. *Current Biology*, *30*(22), 4425-4431.e3.

Kuhn, M. (2020). caret: Classification and Regression Training. R package version 6.0-86.

https://CRAN.R-project.org/package=caret

Kula, A. A. R., Hey, M. H., Couture, J. J., Townsend, P. A., & Dalgleish, H. J. (2020).

Intraspecific competition reduces plant size and quality and damage severity increases defense

responses in the herbaceous perennial, *Asclepias syriaca*. *Plant Ecology*, *221*(6), 421–430.

Laliberté, E. (2018). Measuring specific leaf area and water content.

https://doi.org/10.17504/protocols.io.p3tdqnn

Laliberté, E., & Soffer, R. (2018a). Measuring spectral reflectance and transmittance (350-2500

nm) of large leaves using the Spectra Vista Corporation (SVC) DC-R/T Integrating Sphere.

https://doi.org/10.17504/protocols.io.p8pdrvn

Laliberté, E., & Soffer, R. (2018b). Measuring spectral reflectance and transmittance (350-2500

nm) of small and/or narrow leaves using the Spectra Vista Corporation (SVC) DC-R/T

Integrating Sphere. https://doi.org/10.17504/protocols.io.q56dy9e

Lang, C., Almeida, D. R. A., & Costa, F. R. C. (2017). Discrimination of taxonomic identity at

species, genus and family levels using Fourier Transformed Near-Infrared Spectroscopy (FT-

NIR). *Forest Ecology and Management*, *406*, 219–227.

Lang, C., Costa, F. R. C., Camargo, J. L. C., Durgante, F. M., & Vicentini, A. (2015). Near

Infrared Spectroscopy Facilitates Rapid Identification of Both Young and Mature Amazonian

Tree Species. *PLOS ONE*, *10*(8), e0134521.

Lang, P. L. M., Willems, F. M., Scheepens, J. F., Burbano, H. A., & Bossdorf, O. (2019). Using

herbaria to study global environmental change. *New Phytologist*, *221*(1), 110–122.

Lindström, S. W., Geladi, P., Jonsson, O., & Pettersson, F. (2011). The Importance of Balanced

Data Sets for Partial Least Squares Discriminant Analysis: Classification Problems Using

Hyperspectral Imaging Data. *Journal of Near Infrared Spectroscopy, 19*(4), 233–241.

Ma, S., He, F., Tian, D., Zou, D., Yan, Z., Yang, Y., Zhou, T., Huang, K., Shen, H., & Fang, J.

(2018). Variations and determinants of carbon content in plants: A global synthesis.

Biogeosciences, *15*(3), 693–702.

McManus, K. M., Asner, G. P., Martin, R. E., Dexter, K. G., Kress, W. J., & Field, C. B. (2016).

Phylogenetic Structure of Foliar Spectral Traits in Tropical Forest Canopies. *Remote Sensing*,

8(3), 196.

Meacham-Hensold, K., Montes, C. M., Wu, J., Guan, K., Fu, P., Ainsworth, E. A., Pederson, T.,

Moore, C. E., Brown, K. L., Raines, C., & Bernacchi, C. J. (2019). High-throughput field

phenotyping using hyperspectral reflectance and partial least squares regression (PLSR) reveals

genetic modifications to photosynthetic capacity. *Remote Sensing of Environment*, *231*, 111176.

Meineke, E. K., Davis, C. C., & Davies, T. J. (2018). The unrealized potential of herbaria for

global change biology. *Ecological Monographs*, *88*(4), 505–525.

Meireles, J., Schweiger, A., & Cavender-Bares, J. (2017). spectrolab: Class and Methods for

Hyperspectral Data in R. R package version 0.0.10. https://CRAN.R-

project.org/package=spectrolab

Meireles, J. E., B. O'Meara, and J. Cavender-Bares. (2020a). Linking leaf spectra to the plant tree

of life. Pages 155-172 *in* J. Cavender-Bares, J. A. Gamon, and P. A. Townsend, editors. *Remote*

Sensing of Plant Biodiversity. Springer.

Meireles, J. E., J. Cavender-Bares, P. A. Townsend, S. Ustin, J. A. Gamon, A. K. Schweiger, M.

E. Schaepman, G. P. Asner, R. E. Martin, A. Singh, F. Schrodt, A. Chlus, and B. O'Meara.

(2020b). Leaf reflectance spectra capture the evolutionary history of seed plants. *New*

Phytologist 228:485-493.

Mevik, B.-H., Wehrens, R., & Liland, K. H. (2019). pls: Partial Least Squares and Principal

Component Regression. R package version 2.7-1. https://CRAN.R-project.org/package=pls

Noda, H. M., Motohka, T., Murakami, K., Muraoka, H., & Nasahara, K. N. (2013). Accurate

measurement of optical properties of narrow leaves and conifer needles with a typical integrating

sphere and spectroradiometer. *Plant, Cell & Environment*, *36*(10), 1903–1909.

Norris, K. H., Barnes, R. F., Moore, J. E., & Shenk, J. S. (1976). Predicting Forage Quality by

Infrared Replectance Spectroscopy. *Journal of Animal Science, 43*(4), 889–897.

Nunes, M. H., Davey, M. P., & Coomes, D. A. (2017). On the challenges of using field

spectroscopy to measure the impact of soil type on leaf traits. *Biogeosciences*, *14*(13), 3371–

3385.

Nunes, M. H., Both, S., Bongalov, B., Brelsford, C., Khoury, S., Burslem, D. F. R. P., Philipson,

C., Majalap, N., Riutta, T., Coomes, D. A., & Cutler, M. E. J. (2019). Changes in leaf functional

traits of rainforest canopy trees associated with an El Niño event in Borneo. *Environmental*

Research Letters, *14*(8), 085005.

Queenborough, S. A., & Porras, C. (2014). Expanding the coverage of plant trait databases – A

comparison of specific leaf area derived from fresh and dried leaves. *Plant Ecology & Diversity*,

7(1–2), 383–388.

Perez, T. M., Rodriguez, J., & Heberling, J. M. (2020). Herbarium-based measurements reliably

estimate three functional traits. *American Journal of Botany*, *107*(10), 1457–1464.

Peterson, D. L., Aber, J. D., Matson, P. A., Card, D. H., Swanberg, N., Wessman, C., & Spanner,

M. (1988). Remote sensing of forest canopy and leaf biochemical contents. *Remote Sensing of*

Environment, 24(1), 85–108.

Petibon, F., Czyż, E. A., Ghielmetti, G., Hueni, A., Kneubühler, M., Schaepman, M. E., &

Schuman, M. C. (2021). Variation in reflectance spectroscopy of European beech leaves captures

phenology and biological hierarchies despite measurement uncertainties. BioRxiv,

https://doi.org/10.1101/2021.03.09.434578

1000

1001   Piñeiro, G., Perelman, S., Guerschman, J. P., & Paruelo, J. M. (2008). How to evaluate models:

1002   Observed vs. predicted or predicted vs. observed? *Ecological Modelling, 216*(3), 316–322.

1003

1004 Potůčková, M., Červená, L., Kupková, L., Lhotáková, Z., Lukeš, P., Hanuš, J., Novotný, J., &

1005 Albrechtová, J. (2016). Comparison of Reflectance Measurements Acquired with a Contact

1006 Probe and an Integration Sphere: Implications for the Spectral Properties of Vegetation at a Leaf

1007 Level. *Sensors, 16*(11), 1801.

1008

1009 Prata, E. M. B., Sass, C., Rodrigues, D. P., Domingos, F. M. C. B., Specht, C. D., Damasco, G.,

1010 Ribas, C. C., Fine, P. V. A., & Vicentini, A. (2018). Towards integrative taxonomy in

1011 Neotropical botany: Disentangling the Pagamea guianensis species complex (Rubiaceae).

1012 *Botanical Journal of the Linnean Society, 188*(2), 213–231.

1013

1014 R Core Team. (2020). R: A language and environment for statistical computing. R Foundation

1015 for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

1016

1017 Ramirez, J. A., Posada, J. M., Handa, I. T., Hoch, G., Vohland, M., Messier, C., & Reu, B.

1018 (2015). Near-infrared spectroscopy (NIRS) predicts non-structural carbohydrate concentrations

1019 in different tissue types of a broad range of tree species. *Methods in Ecology and Evolution*, *6*(9),

1020 1018–1025.

1021

1022 Rapaport, T., Hochberg, U., Shoshany, M., Karnieli, A., & Rachmilevitch, S. (2015). Combining

1023 leaf physiology, hyperspectral imaging and partial least squares-regression (PLS-R) for

1024 grapevine water status assessment. *ISPRS Journal of Photogrammetry and Remote Sensing, 109*,

1025 88–97.

1026

1027 Richardson, A. D., Duigan, S. P., & Berlyn, G. P. (2002). An evaluation of noninvasive methods

1028 to estimate foliar chlorophyll content. *New Phytologist, 153*(1), 185–194.

1029

1030 Schweiger, A. K., Cavender-Bares, J., Townsend, P. A., Hobbie, S. E., Madritch, M. D., Wang,

1031 R., Tilman, D., & Gamon, J. A. (2018). Plant spectral diversity integrates functional and

1032 phylogenetic components of biodiversity and predicts ecosystem function. *Nature Ecology &*

1033 *Evolution*, *2*(6), 976–982.

1034

1035 Schweiger, A. K., Cavender-Bares, J., Kothari, S., Townsend, P. A., Madritch, M. D., Grossman,

1036 J. J., Gholizadeh, H., Wang, R., & Gamon, J. A. (2021). Coupling spectral and resource-use

1037 complementarity in experimental grassland and forest communities. BioRxiv,

1038 https://doi.org/10.1101/2020.04.24.060483

1039

1040 Schweiger, A. K., & Laliberté, E. (2020). Processing of leaf spectra.

1041 https://doi.org/10.17504/protocols.io.bhsdj6a6

1042

1043 Serbin, S. P., Dillaway, D. N., Kruger, E. L., & Townsend, P. A. (2012). Leaf optical properties

1044 reflect variation in photosynthetic metabolism and its sensitivity to temperature. *Journal of*

1045 *Experimental Botany*, *63*(1), 489–502.

1046

1047   Serbin, S. P., Singh, A., McNeil, B. E., Kingdon, C. C., & Townsend, P. A. (2014).

1048   Spectroscopic determination of leaf morphological and biochemical traits for northern temperate

1049   and boreal tree species. *Ecological Applications*, *24*(7), 1651–1669.

1050

1051   Serbin, S. P., Wu, J., Ely, K. S., Kruger, E. L., Townsend, P. A., Meng, R., Wolfe, B. T., Chlus,

1052   A., Wang, Z., & Rogers, A. (2019). From the Arctic to the tropics: Multibiome prediction of leaf

1053   mass per area using leaf reflectance. *New Phytologist*, *224*(4), 1557–1568.

1054

1055   Signal Developers (2013). signal: Signal processing. http://r-forge.r-project.org/projects/signal/.

1056

1057   Streher, A. S., Torres, R. da S., Morellato, L. P. C., & Silva, T. S. F. (2020). Accuracy and

1058   limitations for spectroscopic prediction of leaf traits in seasonally dry tropical environments.

1059   *Remote Sensing of Environment*, *244*, 111828.

1060

1061   Sun, Y., Wong, A. K. C., & Kamel, M. S. (2009). Classification of imbalanced data: A review.

1062   *International Journal of Pattern Recognition and Artificial Intelligence, 23*(04), 687–719.

1063

1064   Thiers, B. 2020. The world's herbaria 2019: A summary report based on data from Index

1065   Herbariorum. Available at http://sweetgum.nybg.org/science/ih/

1066

1067   Turner, B. L., Hayes, P. E., & Laliberté, E. (2018). A climosequence of chronosequences in

1068   southwestern Australia. *European Journal of Soil Science, 69*(1), 69–85.

1069

1070 Wang, Z., Chlus, A., Geygan, R., Ye, Z., Zheng, T., Singh, A., Couture, J. J., Cavender-Bares, J.,

1071 Kruger, E. L., & Townsend, P. A. (2020). Foliar functional traits from imaging spectroscopy

1072 across biomes in eastern North America. *New Phytologist, 228*(2), 494–511.

1073

1074 Wieczorek, J., Bloom, D., Guralnick, R., Blum, S., Döring, M., Giovanni, R., Robertson, T., &

1075 Vieglais, D. (2012). Darwin Core: An Evolving Community-Developed Biodiversity Data

1076 Standard. *PLOS ONE, 7*(1), e29715.

1077

1078 Willis, C. G., Ellwood, E. R., Primack, R. B., Davis, C. C., Pearson, K. D., Gallinat, A. S., Yost, J. M.,

1079 Nelson, G., Mazer, S. J., Rossington, N. L., Sparks, T. H., & Soltis, P. S. (2017). Old Plants, New Tricks:

1080 Phenological Research Using Herbarium Specimens. *Trends in Ecology & Evolution, 32*(7), 531–546.

1081

1082 Wold, S. (1994). PLS for multivariate linear modeling. In H. van de Waterbeemd, editor. Chemometric

1083 methods in molecular design, methods and principles in medicinal chemistry, Verlag-Chemie (Weinheim,

1084 Germany), 195–218.

1085

1086 Wu, J., Rogers, A., Albert, L. P., Ely, K., Prohaska, N., Wolfe, B. T., Oliveira, R. C., Saleska, S.

1087 R., & Serbin, S. P. (2019). Leaf reflectance spectroscopy captures variation in carboxylation

1088 capacity across species, canopy environment and leaf age in lowland moist tropical forests. *New*

1089 *Phytologist*, *224*(2), 663–674.

1090

1091 Yang, X., Tang, J., Mustard, J. F., Wu, J., Zhao, K., Serbin, S., & Lee, J.-E. (2016). Seasonal

1092 variability of multiple leaf traits captured by leaf spectroscopy at two temperate deciduous

1093 forests. *Remote Sensing of Environment*, 179, 1–12.

1094

 Supplementary figure captions

 Fig. S1: Fresh-leaf spectral model performance statistics for each trait based on 100 jackknife resamples

 from the calibration data set. Abbreviations: sol = solubles, hemi = hemicellulose, cell = cellulose, lign =

 lignin.

 Fig. S2: Pressed-leaf spectral model performance statistics for each trait based on 100 jackknife resamples

 from the calibration data set. Abbreviations: sol = solubles, hemi = hemicellulose, cell = cellulose, lign =

 lignin.

 Fig. S3: Ground-leaf spectral model performance statistics for each trait based on 100 jackknife resamples

 from the calibration data set. Abbreviations: sol = solubles, hemi = hemicellulose, cell = cellulose, lign =

 lignin.

 Supplementary figures

 Fig. S1

1111

1112

1113

1114

1115

1116

1117   **Fig. S2**

Pressed−leaf spectra

 Fig. S3

1126