Lipid droplets modulate proteostasis, SQST-1/SQSTM1 dynamics, and lifespan in *C.*

elegans

Anita V. Kumar[#,1], Joslyn Mills[#, 1], Wesley M. Parker[#, 1], Joshua A. Leitão[#, 1], Diego I. Rodriguez[, 1],

Celeste Ng[, 1], Rishi Patel[, 1], Joseph L. Aguilera[, 1], Joseph R. Johnson[, 1], Shi Quan Wong[, 1] and

Louis R. Lapierre[*, 1,2]

[#]: Equal contributions

[*]: Correspondence addressed to L.R.L. (louis.rene.lapierre@umoncton.ca)

1: Department of Molecular Biology, Cell Biology and Biochemistry, Brown University, 185

Meeting St., Providence, Rhode Island 02912, USA

2 : Centre de Médecine de Précision du Nouveau Brunswick, Département de chimie et biochimie,

Université de Moncton, 27 rue Providence, Moncton, New Brunswick, E1C 8X3, Canada

ABSTRACT

The ability of organisms to live long depends largely on the maintenance of proteome stability via

proteostatic mechanisms including translational regulation, protein chaperoning and degradation

machineries. In several long-lived *Caenorhabditis elegans* strains, such as insulin/IGF-1 receptor

daf-2 mutants, enhanced proteostatic mechanisms are accompanied by elevated intestinal lipid

stores, but the role of lipid droplets in longevity has remained obscure. Here, while determining

the regulatory network of the selective autophagy receptor SQST-1/SQSTM1, we unexpectedly

uncovered a novel role for lipid droplets in proteostasis and longevity. Using an unbiased genome-

wide RNAi screening approach, we identified several SQST-1 modulators, including proteins

found on lipid droplets and those prone to aggregate with age. SQST-1 accumulated on lipid

droplets when autophagy was inhibited, suggesting that lipid droplets may serve a role in

facilitating selective autophagy. Expansion of intestinal lipid droplets by silencing the conserved

cytosolic triacylglycerol lipase gene *atgl-1/ATGL* enhanced autophagy, and extended lifespan in

an HSF-1/HSF1-dependent and CDC-48/VCP-dependent manner. Silencing *atgl-1* mitigated the

age-related accumulation of SQST-1 and reduced overall ubiquitination of proteins. Reducing

atgl-1 also improved proteostasis in proteotoxicity models of neurodegenerative diseases.

Subcellular analyses revealed that lipid droplets unexpectedly harbor more soluble ubiquitinated

proteins than the cytosol. Accordingly, low lipid droplet levels exacerbated the proteostatic

collapse when autophagy or proteasome function was compromised. Altogether, our study

uncovers a key role for lipid droplets in *C. elegans* as a proteostatic mediator that reduces protein

ubiquitination, facilitates autophagy, and promotes longevity.

INTRODUCTION

One of the major hallmarks of aging is the accumulation of damaged proteins that progressively

compartmentalize in inclusions and aggregates[1]. In the nematode *C. elegans*, numerous proteins

display impaired solubility with age[2,3]. This phenomenon suggests that mechanisms that mediate

protein stabilization, such as chaperone-mediated folding and post translational modifications,

actively contribute to somatic maintenance by preventing the collapse of the proteome[4,5]. The

enhanced proteasomal[6] and autophagic[7,8] capacities, the increased chaperone function[9] and

alterations in ribosomal biogenesis[10] and function[11,12] in long-lived nematodes support the notion

that the balance between synthesis, folding, and efficient clearance of proteins is important for

conferring organismal longevity.

Generally, proteins destined for degradation are tagged via poly-ubiquitination and recognized by

ubiquitin binding domain-containing proteins that direct cargo toward proteasomal or autophagic

degradation. The autophagy receptor, Sequestosome 1 (SQST-1/SQSTM1) is a well-established

and conserved mediator of cargo recognition, which includes ubiquitinated targets[13]. The SQST-

1-cargo complex can interact with the autophagosome proteins LGG-1/GABARAP and LGG-

2/LC3 to enable cargo sequestration in the nascent autophagosome[14]. Notably, SQST-

1/SQSTM1 has emerged as a potential lifespan modulator in nematodes and flies[15-17].

Additionally, selective autophagy of mitochondria has been linked to longevity[18], suggesting that

autophagic degradation of specific cargos and organelles is beneficial against organismal aging.

Although SQST-1/SQSTM1 has been implicated in the selective clearance of aggregated

proteins[19], much remains to be understood about the modulation of this autophagy receptor, its

cargoes, and the nature and extent of its contribution to lifespan.

In nematodes, intestinal cells have a unique role in autophagy-mediated longevity as they manage

nutrient influx and signals from neurons[20] and the germline[21] to coordinate appropriate responses

necessary for organismal survival. In particular, intestinal autophagy genes are required for

lifespan extension[22] by maintaining intestinal tissue integrity and proteostasis. Intestinal cells are

also sites of dynamic lysosomal function[23-26] that mediate specific longevity-associated lipid

signaling[27-29]. Intriguingly, several established long-lived animals including *daf-2*, *glp-1* and *rsks-*

1 mutants also maintain unusually large intestinal lipid droplet stores throughout their life[30-33].

Accordingly, redistributing lipids intended for secreted lipoproteins to intestinal lipid droplets is

sufficient to extend lifespan[29]. However, in long-lived animals, it is unclear whether adult intestinal

lipid stores are simply a consequence of reduced demand from the germline[34], a long-term energy

store rationing strategy[35] or a more direct mediator of somatic maintenance. Here, while

identifying the regulatory network of SQST-1, we found that lipid droplet accumulation in long-

lived nematodes represents a novel proteostatic mechanism that reduces overall SQST-1 and

ubiquitination levels, and mitigates age-related proteostatic stress.

RESULTS

SQST-1 over-expression is detrimental to lifespan in C. elegans

We initially considered the possibility that increasing SQST-1 may improve proteostasis by

enhancing the ability of cells to mediate selective autophagy. We hypothesized that increasing

the expression of a receptor that recognizes ubiquitinated cargoes may suffice in driving their

degradation. While this study was being conducted, the Hansen laboratory showed that over-

expressing SQST-1 can extend lifespan at 20°C[16]. Since longevity interventions and temperature

mechanistically interact[36], we examined lifespan at both 20°C and 25°C using several over-

expressing strains (*psqst-1::sqst-1::rfp*, *psqst-1::sqst-1::gfp::rfp*), including two created in the

Hansen laboratory (*psqst-1::sqst-1::gfp* and *psqst-1::sqst-1*)[16]. Over-expression of SQST-1 was

unexpectedly detrimental at 25°C (**Figure 1a-c, Supplemental Figure 1a-c**) and was not

sufficient to extend lifespan at 20°C (**Figure 1d-f**). While our findings at 20°C differ from recent

literature[16], highlighting potential lab-to-lab experimental variations, our findings are in line with

most studies about the autophagy pathway and aging, i.e. that over-expressing an autophagy

receptor might not necessarily be sufficient to improve lifespan[37]. At 25°C, SQST-1 clearly

accumulates in the head, as well as the gonadal and intestinal tissues (**Supplemental Figure 1d**)

as previously reported[38]. A closer investigation into the temperature-dependent differences in

lifespan revealed marked upregulation of *sqst-1* mRNA at higher temperature in these strains

(from ~5-fold in wild-type, up to ~75 fold in SQST-1::GFP over-expressing animals) (**Figure 1g**).

In order to measure the active trafficking of SQST-1 into the lysosome as a guide for selective

autophagy, we developed a tandem reporter system expressing SQST-1 fused to both GFP and

RFP[7] (GFP signal is quenched in low pH environments). We confirmed that the conversion of

tandem SQST-1 into an RFP-only signal is reduced in *lgg-1-* or *lgg-2*-silenced animals

(**Supplemental Figure 1e**). and autophagy-deficient *atg-18* mutants (**Supplemental Figure 1f**).

While relatively modest at 20°C (**Figure 1h-i**), increasing temperature up to 30°C for 24 hours

significantly enhanced the conversion to the RFP-only SQST-1 signal in the SQST-1 tandem

reporter, suggesting that selective autophagy is induced with heat stress. Substantial

temperature-dependent intestinal and neuronal accumulation of SQST-1 occurred during aging

and over-expressing SQST-1 at 25°C was also accompanied by higher levels of total ubiquitinated

proteins (**Figure 1a-f inset and Supplemental Figure 1g-h**). While over-expressing SQST-1 was

detrimental at 25°C, so was the loss of *sqst-1* (**Supplemental Figure 1i-j**), suggesting that SQST-

1 function has a temperature-specific role in the lifespan of wild-type animals. Expression and

steady-state levels of SQST-1 likely need to be tightly coordinated with cargo targeting and more

importantly, with the rate of protein degradation systems (i.e. autophagy and proteasome).

Altogether, we found that solely increasing SQST-1 is detrimental for lifespan at 25°C and

inconsequential at 20°C, indicating a temperature-dependent effect on SQST-1 dynamics in

aging.

Proteins that bind lipid droplets modulate SQST-1 dynamics

In order to better understand how SQST-1 is regulated, we opted for an unbiased genome-wide

RNAi screening approach in an integrated SQST-1 over-expressing strain (*psqst-1::sqst-1::gfp*)

at 25°C during development. SQST-1 modulators were subsequently validated by gene silencing

in a separate over-expressing strain (*psqst-1::sqst-1::rfp*) during development and adulthood.

Interestingly, the vast majority of modifiers led to the intestinal accumulation of fluorescently

tagged SQST-1 in both development and adulthood; no modifiers were identified that consistently

caused decreased levels of SQST-1 (**Supplemental Table 1**). A substantial portion of genetic

modifiers clustered in ribosomal-related proteins, including proteins coding for small and large

subunits, suggesting that ribosomal assembly and function substantially impact SQST-1

dynamics (**Figure 2a**, **Supplemental Table 1**). This is in line with studies in *C. elegans*[2] and

killifish[39], highlighting the age-related instability of several ribosomal proteins and consequent

ribosomal mis-assembly. Loss of ribosome subunit RPL-43 was also found to increase SQST-1

accumulation during development[40]. Since SQSTM1 was also recently reported to associated

with lipid droplets in macrophages[41], we wondered if SQST-1 might also associate with lipid

droplets. We co-expressed SQST-1::RFP and the lipid resident protein DHS-3 fused to GFP [42] in

nematodes and we found that close to 50% of SQST-1::RFP localizes to lipid droplets when

autophagy is inhibited (**Figure 2b and Supplemental Figure 2a**). Thus, we investigated whether

SQST-1 modulators may also be related to lipid droplet metabolism. We compared our SQST-1

modulators to proteins that have been found to bind lipid droplets[43] and found significant overlap

($p < 5.745e-60$) (**Figure 2c**). Interestingly, several proteins prone to age-dependent aggregation[2]

also modulated SQST-1 accumulation ($p < 6.030e-10$) (**Figure 2c**). Overall, 17 SQST-1

modulators have the propensity to both aggregate with age and bind lipid droplets (**Figure 2c,**

Supplemental Table 2). Several ribosomal subunits (*rps* and *rpl* genes) and mRNA-related

proteins (*inf-1* and *pab-1*) emerged, along with the HSP70 chaperone family member *hsp-*

1/HSPA8, indicating that ribosomal assembly, translation initiation, and protein folding are

important regulators of SQST-1 dynamics (**Figure 2d**). SQST-1::RFP accumulation upon

silencing of these modifiers was mostly gonadal and intestinal (**Figure 2d, Supplemental Figure**

2b). Interestingly, the expression of *sqst-1* was differentially regulated by the 17 SQST-1

modulators whereas silencing large ribosomal proteins generally increased *sqst-1* expression

(**Supplemental Figure 2b**). This highlights that *sqst-1* expression is potentially sensitive to

ribosomal protein stoichiometry. Here, our comparative analysis suggests that age-related SQST-

1 accumulation may be partly attributed to altered interactions between lipid droplets and

progressively unstable protein complexes like the ribosome.

Elevated lipid droplets reduce SQST-1 accumulation and extend lifespan

Intestinal lipid droplet accumulation is a striking feature of several long-lived nematodes[30-32]. Since

SQST-1 accumulated on lipid droplets when autophagy was inhibited (**Figure 2b**), and several

SQST-1 modifiers bind lipid droplets (**Figure 2c**), we speculated that lipid droplets may underlie

the ability of long-lived animals to stabilize or facilitate clearance of potentially aggregating

proteins and maintain proteome stability during aging. In order to specifically stimulate intestinal

lipid droplet accumulation and expansion, we silenced the conserved cytosolic lipase *atgl-1/ATGL*

[44] during adulthood to attenuate lipid droplet breakdown, which led to larger lipid stores thereby

mimicking a key feature of several long-lived animals (**Figure 3a, inset**). Silencing *atgl-1* resulted

in a significant lifespan extension in wild-type animals (12-28%), indicating that lipid droplet

accumulation is sufficient to mediate longevity (**Figure 3a-b**). Silencing ATGL-1 activator *lid-1* or

hormone-sensitive lipase orthologue *hosl-1*[45] also led to significant reduction in SQST-1

accumulation (**Supplemental Figure 2c**). These observations pointed to a possible lipid droplet-

mediated cytoprotective mechanism[46]. Indeed, silencing *atgl-1* in SQST-1::RFP over-expressing

animals led to a substantial increase in lifespan (**Figure 3c)** accompanied by a marked decrease

in SQST-1 accumulation (**Figure 3d**). Similar observations were made in animals over-expressing

SQST-1::GFP (**Supplemental Figure 2d**), suggesting that the lipid droplet-mediated effects

impacts SQST-1-mediated selective autophagy.

To determine whether accumulation of lipids is able to improve the lifespan of proteostatically-

impaired mutants, we silenced *atgl-1* in three well-established short-lived mutants, including *daf-*

16(mu86), *hlh-30(tm1978),* and *hsf-1(sy441)*[47]. Notably, the expression of these transcription

factors was increased at 25°C compared to 20°C, suggesting that their function may be stimulated

by heat (**Supplemental Figure 2e**). Silencing *atgl-1* extended the lifespan of *daf-16* and *hlh-30*

mutants, but not the lifespan of *hsf-1* mutants (**Figure 3e-g**), indicating that lipid droplet-mediated

lifespan extension may require the expression of key chaperones regulated by HSF-1/HSF1, such

as heat shock protein HSP-1, which binds lipid droplets and modulates SQST-1 levels (**Figure**

2d). Interestingly, lipid droplets may harbor chaperones with roles in aggregate clearance[48].

Notably, autophagic activity, as measured by the tandem mCherry::GFP::LGG-1 reporter[7],

showed that *atgl-1* silencing increases the conversion of autophagosomes (GFP+mCherry -

yellow puncta) into autolysosomes (mCherry only - red puncta) (**Figure 3h, Supplemental Figure**

2f). Accordingly, the lifespan of autophagy-deficient *atg-7* mutants was not increased by silencing

atgl-1 (**Supplemental Figure 2g**).

As our lifespan analyses highlighted novel proteostatic and longevity roles for lipid droplets, we

reasoned that long-lived animals with large lipid stores should accumulate less intestinal SQST-

1. Strikingly, *daf-2(e1370)* expressing SQST-1::GFP (**Figure 3i**) or SQST-1::RFP (**Supplemental**

Figure 2h) accumulated negligible intestinal SQST-1 during aging (GFP signal in mid-section was

entirely gonadal, see wild-type comparison in **Figure 1b**). In addition, SQST-1 over-expression

did not significantly affect the long lifespan of *daf-2* animals (**Figure 3i**). Loss of *sqst-1* did not

affect the lifespan of *daf-2* animals (**Supplemental Figure 2i**), as previously shown[16], highlighting

that SQST-1 function becomes less important at 25°C for the lifespan of organisms with relatively

stable proteomes and high lipid stores. The extent of heat-induced increase in SQST-1-mediated

selective autophagy was also attenuated in *daf-2* animals (**Supplemental Figure 2j**) compared

to wild-type (**Figure 1h**), suggesting that elevated lipid droplets may buffer the need for SQST-1

function during heat stress. Silencing *atgl-1* in *daf-2* animals enhanced their intestinal lipid stores

and further extended their lifespan (**Figure 3j**), indicating that elevated lipid droplet accumulation

can also extend lifespan in animals with enhanced proteostasis. Altogether, our data present an

important and previously unrecognized role for lipid droplets in SQST-1 dynamics and longevity.

Silencing atgl-1 elicits limited changes in gene expression

Lipid droplets have been recently shown to coordinate transcriptional programs by sequestering

factors that modulate transcription[49]. Since HSF-1 was required for lifespan extension by *atgl-1*

silencing, we hypothesized that the lipid droplet increase associated with *atgl-1* silencing might

affect the expression of HSF-1-regulated targets. Using RNA sequencing (RNA-seq), we found

that enhancing lipid stores by silencing *atgl-1* in WT or *daf-2* animals had limited effect on global

transcription. Silencing *atgl-1* resulted in 13 overlapping differentially expressed genes (DEG) in

wild-type animals and *daf-2* mutants of which 7 genes are down-regulated in both cases

(**Supplemental Figure 3a-b**). The mRNA levels of *sqst-1* remained unchanged by *atgl-1* silencing

(**Supplemental Figure 3c**). Altering *atgl-1* levels (silencing or over-expression) led to differential

expression of 50 overlapping genes, with no discernable transcription factor signature

(**Supplemental Figure 3d-e**). In addition, expression of *sqst-1* was unchanged in wild-type

animals with low or high levels of *atgl-1* (**Supplemental Figure 3e**). Overall, we concluded that

transcriptional regulation may not contribute significantly to the proteostatic-enhancing and

lifespan-extending effects of lipid droplets. Therefore, lipid droplets may impact proteostasis via

a more direct mechanism on the proteome itself.

Lipid droplets enhance proteostasis by modulating the accumulation of ubiquitinated proteins

The emerging connection between lipid droplet stores and proteome stability led us to investigate

how lipid droplet loss affects proteostasis and SQST-1 abundance. First, we tested whether lipid

droplet depletion affects lifespan using ATGL-1::GFP over-expressing animals[50]. Unlike at 20°C

(**Supplemental Table 3**)[51], we found that over-expressing ATGL-1 was detrimental to lifespan at

25°C (**Figure 4a**) and led to a significant increase in SQST-1 intestinal accumulation

accompanied by lower lipid stores (**Figure 4b**), suggesting that loss of lipid droplet stores can

interfere with SQST-1 dynamics. When autophagy or proteasome function was reduced by

silencing autophagosome protein *lgg-1* or proteasome subunit *rpn-6.1*, respectively, ubiquitinated

protein levels were higher in animals with lower lipid stores, suggesting that lipid droplets may

buffer the proteome and facilitate the processing of unstable or misfolded proteins (**Figure 4c**).

Strikingly, analyzing lipid droplets revealed preferential accumulation of ubiquitinated proteins in

lipid droplet-enriched fractions (**Figure 4d, Supplemental Figure 4a**), indicating that lipid droplets

contain a significant amount of proteins bound for degradation. Lipid droplet-associated

accumulation of ubiquitinated proteins was increased when autophagic or proteasomal

degradation was reduced (**Figure 4d**), suggesting that lipid droplets have the capacity to harbor

many degradation cargoes, which may become particularly relevant for proteostasis when

autophagic and proteasomal systems are failing during aging.

The association and function of ATGL-1 with lipid droplets was recently found to be antagonized

by the AAA-ATPase CDC-48/VCP[52]. CDC-48/VCP is best known for its role in ER-associated

degradation machinery[53], but it has been associated with other functions including endocytosis[54]

and more recently autophagy itself[55]. CDC-48/VCP helps unfold unstable, ubiquitinated, and

proteasomal degradation-bound proteins in concert with heat shock protein HSP-70, a key

regulated target of HSF-1[56]. Thus, we assayed the effect of the loss of CDC-48 on SQST-1 and

ATGL-1 levels using fluorescent reporters. Silencing *cdc-48.2* increased the levels of ATGL-

1::GFP (**Supplemental Figure 4b)** and led to the accumulation of SQST-1::RFP (**Supplemental**

Figure 4c). In mammalian cells, loss of proteasome function drives SQSTM1 to divert cargo to

selective autophagy[57], but stimulating proteasomal processing by over-expressing of VCP can

reduce SQSTM1 accumulation[58]. Here, we reasoned that CDC-48 functions related to processing

ubiquitinated targets and autophagy may underlie the ability of nematodes with elevated lipid

droplets to attenuate SQST-1 accumulation and modulate lifespan. Accordingly, silencing *atgl-1*

in *cdc-48.1* or *cdc-48.2* mutants failed to significantly extend lifespan (**Supplemental Figure 4d,**

Supplemental Table 4), indicating that lipid droplet-mediated lifespan extension requires

functional CDC-48.

As lipid droplet size increases, its surface also increases, and it is possible that the capacity of

this organelle to bind and stabilize proteins may also increase as well. Enhancing lipid droplet

stores by silencing *atgl-1* reduced the overall accumulation of ubiquitinated proteins, in particular

in the lower solubility (5% SDS soluble) fraction (**Figure 4e**), suggesting that typically insoluble

proteins are less likely to be ubiquitinated when lipid droplets abound. Notably, the types of

proteins that aggregate with age differ between wild-type and long-lived *daf-2* mutants, as the

latter tends to accumulate aggregating proteins that are less hydrophobic than those aggregating

in wild-type animals[4]. Reducing lipid droplet stores genetically by silencing lipogenic genes *sbp-*

1(SREBP2), *lpin-1(LIPN1)* or *fasn-1(FASN)* enhanced SQST-1 accumulation in wild-type animals

(**Supplemental Figure 4e**). Similarly, loss of lipid droplet stores by silencing *lpin-1* in *daf-2*

resulted in increased SQST-1 accumulation and overall protein ubiquitination (**Figure 4f-g**). As

proteostasis failure is a feature of neurodegeneration, we tested the effect of increasing lipid

droplet levels in proteotoxic contexts. Animals intestinally-expressing an aggregating poly-

glutamine protein fused to YFP (Q44::YFP)[59] had reduced numbers of aggregates when *atgl-1*

was silenced **(Figure 4h)**. Reducing the expression of *atgl-1* in a muscle-expressing proteotoxic

amyloid protein Aβ-42 resulted in a marked protection against aggregation-associated paralysis[60]

(**Figure 4i**). Altogether, our data demonstrate that lipid droplets are important for proteostasis and

contribute to lifespan by facilitating autophagy and stabilizing the proteome.

DISCUSSION

Decades of aging research has uncovered key longevity-regulating pathways in *C. elegans*[47], yet

the role of lipid droplets in the lifespan of several established long-lived nematodes has remained

unresolved. Here, while studying the regulation of the selective autophagy receptor SQST-1, we

unexpectedly uncovered a direct role for lipid droplets in proteostasis and lifespan. Several SQST-

1 modulators associate with lipid droplets and aggregate with age, including ribosomal,

translation-related, and folding-related proteins[43]. Ribosomal assembly dysfunction due to loss in

subunit stoichiometry may burden the autophagy machinery, and contribute to age-related

proteotoxicity[39]. Importantly, lipid droplet accumulation, by silencing cytosolic triacylglycerol lipase

atgl-1/ATGL, mitigates the progressive age-related SQST-1 accumulation, a benefit recapitulated

in long-lived, lipid droplet-rich *daf-2* mutant animals. Lipid droplets are cytoprotective as they

prevent aberrant SQST-1 accumulation, in part via the functioning of AAA-ATPase CDC-48/VCP

which processes ubiquitinated proteins for proteasomal degradation[61]. Strikingly, we found that

the majority of soluble ubiquitinated proteins in nematodes was associated with lipid droplets,

suggesting that unstable proteins bound for degradation may be stabilized by the surface of lipid

droplets. Our findings also provide support to the importance of the emerging lipid-droplet-

mediated protein degradation process[62,63] in longevity and aging.   Altogether, our study

strengthens the emerging concept that lipid droplets serve as a buffer for proteostasis[46] by

stabilizing proteomes and coordinating protein degradation machineries.

Understanding the regulation of SQST-1/SQSTM1 during aging and in different disease contexts

is important in order to determine the validity of stimulating selective autophagy as a therapeutic

strategy to improve proteostasis in age-related diseases[64]. Here, we find that over-expressing

SQST-1 had no effect on the lifespan of wild-type animals at 20°C[16] whereas elevated SQST-1

level at 25°C was detrimental. In hindsight, one may have predicted that over-expressing a

ubiquitin-binding protein with a propensity for oligomerization[65] might not be necessarily

advantageous, particularly in organisms nearing their protein solubility limit[2,3] and displaying

overall proteome instability during aging[5]. SQST-1 over-expression was primarily visible in

neuronal, gonadal and intestinal cells and accumulated in a temperature-dependent manner and

progressively during aging. The inability of stressed and aging animals to process ubiquitinated

cargos bound for proteasomal or autophagic degradation may lead to a gradual build-up of SQST-

1. Accordingly, SQST-1 over-expression may over-sensitize the animals to unstable,

ubiquitination-prone proteins, posing an additional challenge to the autophagic machinery. This

is exemplified by recent evidence that reducing SQST-1 specifically in the neurons of a nematode

model of ALS is protective[66], possibly since mutant fused in sarcoma (FUS) expression promotes

SQST-1 accumulation, which exacerbates proteotoxicity and neurodegeneration. Notably,

SQSTM1 can phase separate when interacting with ubiquitinated cargoes[67], but the impact of the

formation of these condensates on overall proteostasis is unclear.

Lipid droplet accumulation is a prominent and under-explored phenotype of several long-lived

nematodes, including well-established models such as germline-less *glp-1* animals, protein

translation *rsks-1* mutants, insulin/IGF-1 receptor *daf-2* mutants[68] as well as in long-lived animals

with reduced nuclear export[69]. In addition, lifespan extension by inhibiting intestinal lipid secretion

mediated by vitellogenins results in lipid redistribution to intestinal lipid droplet stores[29], a process

believed to generate precursors for lysosomally-derived lipid signals[28]. While enhanced lysosomal

lipolysis and lipophagy may stimulate longevity-associated lipid signaling, overactive cytosolic

lipolysis decreases the ability of cells to maintain a stable proteome, which may burden the

selective autophagy machinery. Our findings support that lipid droplets are much more than just

lipid storage organelles[46] and can serve as interfaces to enable protein stabilization and

processing.

Rationing of intestinal lipid stores was originally evoked as a potential mechanism to provide long-

term energy for dauer larvae that arrest feeding[35]. Here, we propose a role for lipid stores during

adulthood that focuses on their capacity to buffer proteostasis during aging in concert with heat

shock chaperones, thereby unburdening protein degradation systems (**Figure 5**). ATGL-1 over-

expression leads to lipid store depletion [51], but, more importantly, it also removes the protection

conferred by lipid droplets and renders animals vulnerable to proteotoxic stress associated with

reduced proteasomal function, heat and aging. Thus, regulating the activity of ATGL-1 via

transcription[70], post-translational modification[71] or alteration in CDC48/VCP activity[52] provide a

mechanism by which cells modify protein homeostasis. Overall, our data also show that the lipid

droplet-mediated decrease in protein ubiquitination and the enhancement in lifespan require the

function of the AAA-ATPase CDC-48/VCP, a ubiquitin protein processing enzyme and autophagy

modulator[55].

Lipid droplet expansion may belong to an arsenal of proteostatic measures that lipid-storing cells

and possibly neurons can employ to prevent protein misfolding and aggregation. While lipid

droplet accumulation in immune cells such as macrophages[72] and microglia[73] may be

inflammatory and contribute to aging, lipid droplets may confer some post-mitotic, differentiated

cells with added protection against protein aggregation[48] and mitochondrial lipid overload[74]. For

example, lipid droplet accumulation in Alzheimer's disease pathogenesis[75] may highlight an effort

from neuronal cells to improve proteome stability and resilience against progressive

proteotoxicity. Notably, drastic inhibition of triacylglycerol lipolysis becomes detrimental to

neurons[76] as larger lipid droplets are less easily broken down[50]. However, a recent study showed

that lipid droplet accumulation protected hyper-activated neurons from cell death[77], a

phenomenon potentially relevant in Alzheimer's disease[78]. Thus, moderate accumulation of lipid

droplets may be more desirable to provide protection against age-related proteotoxicity. In

agreement, our data show marked protection against paralysis in a proteostatic Alzheimer's

disease model by attenuating cytosolic lipolysis.

Enhancing proteostasis by stimulating the autophagy process has emerged as an attractive

strategy to mitigate several age-related diseases with pathological proteostatic decline such as

neurodegenerative diseases [79]. However, our study indicates that stimulating selective autophagy

by specifically increasing the expression of one selective autophagy receptor, SQST-1/SQSTM1,

is not sufficient to improve proteostasis and may exacerbate age-related proteostatic collapse.

Overall, our study suggests that therapeutic improvement in proteostasis may benefit from a

combinatorial approach in which the whole autophagy/lysosomal machinery is stimulated

concomitantly with a modest reduction in lipid droplet breakdown. Such an approach may prevent

pathogenic burdening of the autophagy/lysosome pathway with cargoes that could have

otherwise been stabilized by lipid droplets or routed to the proteasome.

METHODS

C. elegans **strain maintenance**

Nematodes were maintained at 20°C on agar NGM plates seeded with OP50 *E. coli* unless

otherwise noted, as previously described[80]. Synchronized populations were prepared using a

sodium hypochlorite solution to collect eggs as previously described[81]. Supplemental Tables 5

and 6 contains the list of strains used in this study. HT115 *E. coli* and RNAi clones from the

Ahringer library (Source Bioscience) were used for RNAi experiments[82].

Transgenic strain construction

DNA constructs for the plasmids pLAP26 (*psqst-1::sqst-1::rfp::unc-54 3'UTR*) and pLAP29 (*psqst-*

1::sqst-1::gfp::rfp::unc-54 3'UTR) were assembled using HiFi cloning and were injected into the

germline of Day 1 adults (pLAP26 was co-injected with the pLAP7 (*myo-2::gfp::unc-54*)).

Transgenic progeny was UV-irradiated for extrachromosomal array integration and selected for

100% transmission rate and backcrossed at least four times to wildtype N2. Details about strain

construction are provided in Supplemental Table 6.

Genome-wide RNAi screen

Approximately 50 synchronized transgenic eggs (*psqst-1::SQST-1::GFP*) were transferred onto

plates seeded with RNAi against all genes from the Ahringer library (Source BioScience), which

represents about 86% of the predicted genes in the *C. elegans* genome[82]. Nematodes were

developed at 25°C and changes in GFP intensity or expression pattern were monitored on Day 1

of adulthood. As a negative control, bacteria expressing the L4440 backbone (i.e. empty vector)

expressed in HT115 *E. coli* were used. The identity of each RNAi was not known to the scorer

and fluorescence intensity was scored using a V20 dissection fluorescence microscope (Zeiss).

Modifiers from the genome-wide RNAi screen were sequenced (GENEWIZ), validated three times

and corroborated using transgenic animals expressing *SQST-1::RFP*. RNAi against the genes

with the strongest effect on GFP expression were subsequently validated by silencing in

adulthood only. Details are provided in Supplemental Tables 1 and 2.

Lifespan analyses

Eggs obtained from bleaching were transferred onto OP50 *E. coli.* bacteria-seeded agar plates.

All lifespan analyses were carried out at 25°C starting at Day 1 of adulthood after development at

20°C, unless otherwise noted. For gene knockdown experiments, worms were transferred to

control bacteria expressing the empty vector (L4440) or expressing corresponding RNAi at day

one of adulthood. Viability was scored every 1 to 3 days, as previously described[83]. Survival

curves and statistical analysis were generated using the Stata 15.0 software (StataCorp). Details

are provided in Supplemental Tables 3 and 4.

Comparative analyses

Knocked-down genes with the strongest effect on SQST-1::GFP levels were analyzed for gene

ontology using WormCat[84] to identify which pathways were enriched. RNAi against genes that

overlapped with the SQST-1 modulators, lipid droplet proteome[43], and insoluble proteins[2] were

then tested in adult-only experiments. Statistical significance of the overlap between any two

groups of genes was calculated by hypergeometric probability (nemates.org, Lund laboratory,

University of Kentucky). For adult-only RNAi validation, eggs from wild-type and SQST-1

transgenic animals were developed on OP50 *E coli.* bacteria at 20°C and transferred to

corresponding RNAi plates at Day 1 of adulthood and grown at 25°C for an additional 72 hours.

Adults were transferred away from their progeny and bright field and fluorescent images were

taken daily.

Imaging and analyses

Worms were visualized using a Zeiss Discovery V20 fluorescence dissecting microscope (Zeiss,

White Plains, NY). Worms were immobilized with 0.1% sodium azide in M9 solution (42 mM

Na_2HPO_4 22 mM KH_2PO_4 86 mM NaCl, 1 mM $MgSO_4$-$7H_2O$) on agar plates and images were

taken via Zen imaging software, using consistent parameters (magnification and exposure)

within each experiment. For quantification of fluorescent signal, total fluorescence of each worm

in each image was analyzed using ImageJ software and averaged for mean total fluorescence.

Confocal microscopy images were obtained on an Olympus FV3000 inverted confocal laser

scanning microscope (Leduc Bioimaging Facility, Brown University). Quantification of

autophagosomes marked by GFP and mCherry showing yellow puncta and autolysosomes

marked with only mCherry as GFP is quenched in acidic pH showing up as red puncta in the

LGG-1 tandem reporter strain was carried out by color thresholding composite images for yellow

or red puncta followed by counting the thresholded puncta using particle analysis on ImageJ.

The overlap between lipid droplet protein (DHS-3) and tagged SQST-1 in the over-expressing

strain was also performed using color thresholding followed by particle analysis functions in

ImageJ (NIH).

Lipid staining

Worms were collected and washed twice with M9 solution and then fixed with 60% isopropanol

for 30 minutes. After fixation, worms were stained overnight on a rocker with freshly prepared

60% Oil Red O (stock solution of 0.5% ORO in isopropanol diluted with water, equilibrated

overnight on a rocker, and gravity-filtered). The following day, worms were washed with TBS-T

(50 mM Tris base, 150 mM Tris HCl, 1.5 M NaCl, and 0.05% Tween-20) and imaged using a

Zeiss Discovery V20 fluorescence dissecting microscope.

Gene expression analysis

RNA was extracted from approximately 3,000 Day 1 worms and cDNA was prepared as

previously described [8]. Gene expression levels were measured in biological triplicates using iTaq

Universal SYBR Green Supermix (BIO-RAD), and a Roche LightCycler 96 (Indianapolis, IN).

Expression was normalized using 2 housekeeping genes, *act-1* and *cyn-1* and statistical analyses

were performed using GraphPad Prism 7 (GraphPad Software)*.* See Supplemental Table 7 for

qPCR primers details.

RNAseq sample preparation and analysis

WT or *daf-2* mutant worms were developed on OP50 at 20°C and transferred to corresponding

RNAi plates at Day 1 of adulthood and grown at 25°C for an additional 96 hours. Worms over-

expressing ATGL-1::GFP were developed on OP50 at 20°C and grown at 25°C for an additional

96 hours. Adults were transferred away from their progeny daily. RNA was extracted and cDNA

was prepared as previously described[8]. RNA quality was confirmed by BioAnalyzer (Genomics

Core, Brown University), and *atgl-1* silencing was confirmed by qPCR before submitting samples

for RNAseq analysis by GENEWIZ as previously described[8].

Lipid droplet fractionation

Lipid droplets were isolated by ultracentrifugation from 24,000 Day 5 animals grown at 25°C on

RNAi for 96 hours. Worms were homogenized with a ball-bearing homogenizer in MSB buffer

(250mM sucrose and 10mM Tris-HCl pH 7.4) containing protease inhibitor (Roche). The lysate

was cleared of cuticle debris and unlysed worms by centrifuging at 500 x g for 5 minutes. 10% of

the cleared lysate was reserved as "Input". Nuclei and non-lipid droplet organelles were pelleted

out of the cytosol by an ultracentrifugation spin at 45,000 RPM in a TLA100.1 rotor for 30 minutes

at 4°C. The lipid droplet layer at the top of the centrifugation tube was aspirated, and

contaminating organelles were removed by an additional centrifugation of the lipid droplet fraction.

The final volume of lipid droplet and cytosol fractions were kept consistent across conditions.

Immunoblotting

Protein lysates were collected from worms using RIPA buffer (50 mM Tris-HCl, 250 mM sucrose,

1 mM EDTA, and Roche protease inhibitor tablet, with 1% or 5% SDS) and a handheld

homogenizer, and cleared lysate protein concentrations were quantified using the DC Protein

Assay kit (BIO-RAD). Equal amounts of total protein (10 μg) or comparative volumes for lipid

droplet fractions (Input volume is 10% of lipid droplet and cytosol fractions) were separated by

SDS-PAGE on a 4-15% Tris-Glycine gel and transferred to nitrocellulose. The membrane was

briefly stained with Ponceau S to confirm even transfer, rinsed with TBS-T until clear, blocked

with 5% nonfat dry milk in TBS-T, and immunoblotted with anti-Ubiquitin (Invitrogen MAI-10035),

anti-GFP (Santa Cruz SC-8334), and anti-Actin (Milllipore, MAB1501R). The membrane was

developed using SuperSignal West Femto Maximum Sensitivity Substrate (ThermoFisher) and

imaged on a ChemiDoc Imaging System (BIO-RAD).

DATA AVAILABILITY STATEMENT

465 RNA sequencing data are accessible on NCBI GEO (Accession number: GEO204953).

CONTRIBUTIONS

AVK, JM, WMP and JAL performed lifespan analyses and imaging as well as RNAi sequencing,

validation, and analysis. AVK, JM, JAL and JRJ validated the whole-genome RNAi screening. JM

performed the RNA sequencing and lipid droplet analyses. WMP conducted lipid staining and

AVK performed the confocal microscopy imaging. DIR performed RNAi sequencing and

conducted qPCR analyses. CN, RP and JLA performed lifespan analysis repeats. SQW and WMP

developed SQST-1 over-expressing and tandem reporter strains. LRL designed the experiments,

conducted initial RNAi screening, imaging, qPCR and lifespan analyses, and wrote the

manuscript. All co-authors edited the manuscript.

ACKNOWLEDGEMENTS

We are grateful for the technical support provided by Erin McConnell, Rachel Tam and Tuong

Tran. We thank the Malene Hansen laboratory (SBPMDI) and the Caenorhabditis Genetics

Center (U. Minnesota, P40 OD010440) for providing transgenic and mutant strains and the

Andrew Dillin laboratory (UC Berkeley/HHMI) for sharing expression plasmids. This work was

funded by grants from the National Institute of Health (R00 AG042494, R01 AG051810 and R21

AG068922) and a Glenn Award for Research in Biological Mechanisms of Aging from the Glenn

Foundation for Medical Research to LRL.

REFERENCES

1   Koga, H., Kaushik, S. & Cuervo, A. M. Protein homeostasis and aging: The importance of

exquisite quality control. *Ageing Res Rev* **10**, 205-215, doi:10.1016/j.arr.2010.02.001

(2011).

2   Reis-Rodrigues, P. *et al.* Proteomic analysis of age-dependent changes in protein

solubility identifies genes that modulate lifespan. *Aging Cell* **11**, 120-127,

doi:10.1111/j.1474-9726.2011.00765.x (2012).

3   Sui, X. *et al.* Widespread remodeling of proteome solubility in response to different protein

homeostasis stresses. *Proc Natl Acad Sci U S A* **117**, 2422-2431,

doi:10.1073/pnas.1912897117 (2020).

4   Walther, D. M. *et al.* Widespread Proteome Remodeling and Aggregation in Aging C.

elegans. *Cell* **161**, 919-932, doi:10.1016/j.cell.2015.03.032 (2015).

5   Ben-Zvi, A., Miller, E. A. & Morimoto, R. I. Collapse of proteostasis represents an early

molecular event in Caenorhabditis elegans aging. *Proc Natl Acad Sci U S A* **106**, 14914-

14919, doi:0902882106 [pii] 10.1073/pnas.0902882106 (2009).

6   Vilchez, D. *et al.* RPN-6 determines C. elegans longevity under proteotoxic stress

conditions. *Nature* **489**, 263-268, doi:nature11315 [pii] 10.1038/nature11315 (2012).

7   Chang, J. T., Kumsta, C., Hellman, A. B., Adams, L. M. & Hansen, M. Spatiotemporal

regulation of autophagy during Caenorhabditis elegans aging. *eLife* **6**,

doi:10.7554/eLife.18459 (2017).

8   Lapierre, L. R. *et al.* The TFEB orthologue HLH-30 regulates autophagy and modulates

longevity in Caenorhabditis elegans. *Nat Commun* **4**, 2267, doi:ncomms3267 [pii]

10.1038/ncomms3267 (2013).

9   Morley, J. F. & Morimoto, R. I. Regulation of longevity in Caenorhabditis elegans by heat

shock factor and molecular chaperones. *Mol Biol Cell* **15**, 657-664, doi:10.1091/mbc.E03-

07-0532 E03-07-0532 [pii] (2004).

10   Tiku, V. *et al.* Small nucleoli are a cellular hallmark of longevity. *Nat Commun* **8**, 16083,

doi:10.1038/ncomms16083 (2017).

11   Visscher, M. *et al.* Proteome-wide Changes in Protein Turnover Rates in C. elegans

Models of Longevity and Age-Related Disease. *Cell Rep* **16**, 3041-3051,

doi:10.1016/j.celrep.2016.08.025 (2016).

12   Dhondt, I. *et al.* FOXO/DAF-16 Activation Slows Down Turnover of the Majority of Proteins

in C. elegans. *Cell Rep* **16**, 3028-3040, doi:10.1016/j.celrep.2016.07.088 (2016).

13   Kirkin, V. & Rogov, V. V. A Diversity of Selective Autophagy Receptors Determines the

Specificity of the Autophagy Pathway. *Mol Cell* **76**, 268-285,

doi:10.1016/j.molcel.2019.09.005 (2019).

14   Pankiv, S. *et al.* p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of

ubiquitinated protein aggregates by autophagy. *J Biol Chem* **282**, 24131-24145,

doi:M702824200 [pii] 10.1074/jbc.M702824200 (2007).

15   Aparicio, R., Hansen, M., Walker, D. W. & Kumsta, C. The selective autophagy receptor

SQSTM1/p62 improves lifespan and proteostasis in an evolutionarily conserved manner.

Autophagy **16**, 772-774, doi:10.1080/15548627.2020.1725404 (2020).

16   Kumsta, C. *et al.* The autophagy receptor p62/SQST-1 promotes proteostasis and

longevity in C. elegans by inducing autophagy. *Nat Commun* **10**, 5648,

doi:10.1038/s41467-019-13540-4 (2019).

17   Aparicio, R., Rana, A. & Walker, D. W. Upregulation of the Autophagy Adaptor

p62/SQSTM1 Prolongs Health and Lifespan in Middle-Aged Drosophila. *Cell Rep* **28**,

1029-1040 e1025, doi:10.1016/j.celrep.2019.06.070 (2019).

18   Palikaras, K., Lionaki, E. & Tavernarakis, N. Coordination of mitophagy and mitochondrial

biogenesis during ageing in C. elegans. *Nature* **521**, 525-528, doi:10.1038/nature14300

(2015).

19   Johansen, T. & Lamark, T. Selective autophagy mediated by autophagic adapter proteins.

538   *Autophagy* **7**, 279-296, doi:14487 [pii] (2011).

539   20   Zhang, B. *et al.* Brain-gut communications via distinct neuroendocrine signals

540   bidirectionally regulate longevity in C. elegans. *Genes Dev* **32**, 258-270,

541   doi:10.1101/gad.309625.117 (2018).

542   21   Antebi, A. Regulation of longevity by the reproductive system. *Exp Gerontol* **48**, 596-602,

543   doi:10.1016/j.exger.2012.09.009 (2013).

544   22   Gelino, S. *et al.* Intestinal Autophagy Improves Healthspan and Longevity in C. elegans

545   during Dietary Restriction. *PLoS Genet* **12**, e1006135, doi:10.1371/journal.pgen.1006135

546   (2016).

547   23   Lapierre, L. R., Gelino, S., Melendez, A. & Hansen, M. Autophagy and lipid metabolism

548   coordinately modulate life span in germline-less C. elegans. *Curr Biol* **21**, 1507-1514,

549   doi:S0960-9822(11)00844-X [pii] 10.1016/j.cub.2011.07.042 (2011).

550   24   Wang, M. C., O'Rourke, E. J. & Ruvkun, G. Fat metabolism links germline stem cells and

551   longevity in C. elegans. *Science* **322**, 957-960, doi:322/5903/957 [pii]

552   10.1126/science.1162011 (2008).

553   25   Lapierre, L. R., Melendez, A. & Hansen, M. Autophagy links lipid metabolism to longevity

554   in C. elegans. *Autophagy* **8**, 144-146, doi:18722 [pii] 10.4161/auto.8.1.18722 (2012).

555   26   Chapin, H. C., Okada, M., Merz, A. J. & Miller, D. L. Tissue-specific autophagy responses

556   to aging and stress in C. elegans. *Aging (Albany NY)* **7**, 419-434 (2015).

557   27   Ramachandran, P. V. *et al.* Lysosomal Signaling Promotes Longevity by Adjusting

558   Mitochondrial Activity. *Dev Cell* **48**, 685-696 e685, doi:10.1016/j.devcel.2018.12.022

559   (2019).

560   28   Folick, A. *et al.* Aging. Lysosomal signaling molecules regulate longevity in Caenorhabditis

561   elegans. *Science* **347**, 83-86, doi:10.1126/science.1258857 (2015).

29   Seah, N. E. *et al.* Autophagy-mediated longevity is modulated by lipoprotein biogenesis.

Autophagy **12**, 261-272, doi:10.1080/15548627.2015.1127464 (2016).

30   O'Rourke, E. J., Soukas, A. A., Carr, C. E. & Ruvkun, G. C. elegans major fats are stored

in vesicles distinct from lysosome-related organelles. *Cell Metab* **10**, 430-435, doi:S1550-

4131(09)00301-5 [pii] 10.1016/j.cmet.2009.10.002 (2009).

31   Lapierre, L. R. *et al.* Autophagy genes are required for normal lipid levels in C. elegans.

Autophagy **9**, 278-286, doi:22930 [pii] 10.4161/auto.22930 (2013).

32   Brooks, K. K., Liang, B. & Watts, J. L. The influence of bacterial diet on fat storage in C.

elegans. *PLoS One* **4**, e7545, doi:10.1371/journal.pone.0007545 (2009).

33   Perez, C. L. & Van Gilst, M. R. A 13C isotope labeling strategy reveals the influence of

insulin signaling on lipogenesis in C. elegans. *Cell Metab* **8**, 266-274, doi:S1550-

4131(08)00248-9 [pii] 10.1016/j.cmet.2008.08.007 (2008).

34   Kirkwood, T. B. & Holliday, R. The evolution of ageing and longevity. *Proc R Soc Lond B*

Biol Sci **205**, 531-546 (1979).

35   Narbonne, P. & Roy, R. Caenorhabditis elegans dauers need LKB1/AMPK to ration lipid

reserves and ensure long-term survival. *Nature* **457**, 210-214, doi:nature07536 [pii]

10.1038/nature07536 (2009).

36   Miller, H. *et al.* Genetic interaction with temperature is an important determinant of

nematode longevity. *Aging Cell* **16**, 1425-1429, doi:10.1111/acel.12658 (2017).

37   Lapierre, L. R., Kumsta, C., Sandri, M., Ballabio, A. & Hansen, M. Transcriptional and

epigenetic regulation of autophagy in aging. *Autophagy* **11**, 867-880,

doi:10.1080/15548627.2015.1034410 (2015).

38   Dupuy, D. *et al.* Genome-scale analysis of in vivo spatiotemporal promoter activity in

Caenorhabditis elegans. *Nat Biotechnol* **25**, 663-668, doi:10.1038/nbt1305 (2007).

39   Kelmer Sacramento, E. *et al.* Reduced proteasome activity in the aging brain results in
ribosome stoichiometry loss and aggregation. *Mol Syst Biol* **16**, e9596,
doi:10.15252/msb.20209596 (2020).

40   Guo, B. *et al.* Genome-wide screen identifies signaling pathways that regulate autophagy
during Caenorhabditis elegans development. *EMBO Rep* **15**, 705-713,
doi:10.1002/embr.201338310 (2014).

41   Robichaud, S. *et al.* Identification of novel lipid droplet factors that regulate lipophagy and
cholesterol efflux in macrophage foam cells. *Autophagy* **17**, 3671-3689,
doi:10.1080/15548627.2021.1886839 (2021).

42   Zhang, P. *et al.* Proteomic study and marker protein identification of Caenorhabditis
elegans lipid droplets. *Mol Cell Proteomics* **11**, 317-328, doi:10.1074/mcp.M111.016345
(2012).

43   Vrablik, T. L., Petyuk, V. A., Larson, E. M., Smith, R. D. & Watts, J. L. Lipidomic and
proteomic analysis of Caenorhabditis elegans lipid droplets and identification of ACS-4 as
a lipid droplet-associated protein. *Biochim Biophys Acta* **1851**, 1337-1345,
doi:10.1016/j.bbalip.2015.06.004 (2015).

44   Zechner, R., Kienesberger, P. C., Haemmerle, G., Zimmermann, R. & Lass, A. Adipose
triglyceride lipase and the lipolytic catabolism of cellular fat stores. *J Lipid Res* **50**, 3-21,
doi:10.1194/jlr.R800031-JLR200 (2009).

45   Lee, J. H. *et al.* Lipid droplet protein LID-1 mediates ATGL-1-dependent lipolysis during
fasting in Caenorhabditis elegans. *Mol Cell Biol* **34**, 4165-4176, doi:10.1128/MCB.00722-
14 (2014).

46   Olzmann, J. A. & Carvalho, P. Dynamics and functions of lipid droplets. *Nat Rev Mol Cell*
Biol **20**, 137-155, doi:10.1038/s41580-018-0085-z (2019).

47   Denzel, M. S., Lapierre, L. R. & Mack, H. I. D. Emerging topics in C. elegans aging

research: Transcriptional regulation, stress response and epigenetics. *Mech Ageing Dev*

177, 4-21, doi:10.1016/j.mad.2018.08.001 (2019).

48   Moldavski, O. *et al.* Lipid Droplets Are Essential for Efficient Clearance of Cytosolic

Inclusion Bodies. *Dev Cell* **33**, 603-610, doi:10.1016/j.devcel.2015.04.015 (2015).

49   Mejhert, N. *et al.* Partitioning of MLX-Family Transcription Factors to Lipid Droplets

Regulates Metabolic Gene Expression. *Mol Cell* **77**, 1251-1264 e1259,

doi:10.1016/j.molcel.2020.01.014 (2020).

50   Zhang, S. O. *et al.* Genetic and dietary regulation of lipid droplet expansion in

Caenorhabditis elegans. *Proc Natl Acad Sci U S A* **107**, 4640-4645,

doi:10.1073/pnas.0912308107 (2010).

51   Zaarur, N. *et al.* ATGL-1 mediates the effect of dietary restriction and the insulin/IGF-1

signaling pathway on longevity in C. elegans. *Mol Metab* **27**, 75-82,

doi:10.1016/j.molmet.2019.07.001 (2019).

52   Olzmann, J. A., Richter, C. M. & Kopito, R. R. Spatial regulation of UBXD8 and p97/VCP

controls ATGL-mediated lipid droplet turnover. *Proc Natl Acad Sci U S A* **110**, 1345-1350,

doi:10.1073/pnas.1213738110 (2013).

53   Wu, X. & Rapoport, T. A. Mechanistic insights into ER-associated protein degradation.

Curr Opin Cell Biol **53**, 22-28, doi:10.1016/j.ceb.2018.04.004 (2018).

54   Bug, M. & Meyer, H. Expanding into new markets--VCP/p97 in endocytosis and

autophagy. *J Struct Biol* **179**, 78-82, doi:10.1016/j.jsb.2012.03.003 (2012).

55   Hill, S. M. *et al.* VCP/p97 regulates Beclin-1-dependent autophagy initiation. *Nat Chem*

Biol **17**, 448-455, doi:10.1038/s41589-020-00726-x (2021).

56   Vembar, S. S. & Brodsky, J. L. One step at a time: endoplasmic reticulum-associated

degradation. *Nat Rev Mol Cell Biol* **9**, 944-957, doi:10.1038/nrm2546 (2008).

57   Demishtein, A. *et al.* SQSTM1/p62-mediated autophagy compensates for loss of
proteasome polyubiquitin recruiting capacity. *Autophagy* **13**, 1697-1708,
doi:10.1080/15548627.2017.1356549 (2017).

58   Korolchuk, V. I., Mansilla, A., Menzies, F. M. & Rubinsztein, D. C. Autophagy inhibition
compromises degradation of ubiquitin-proteasome pathway substrates. *Mol Cell* **33**, 517-
527, doi:10.1016/j.molcel.2009.01.021 (2009).

59   Mohri-Shiomi, A. & Garsin, D. A. Insulin signaling and the heat shock response modulate
protein homeostasis in the Caenorhabditis elegans intestine during infection. *J Biol Chem*
283, 194-201, doi:10.1074/jbc.M707956200 (2008).

60   McColl, G. *et al.* Utility of an improved model of amyloid-beta (Abeta(1)(-)(4)(2)) toxicity in
Caenorhabditis elegans for drug screening for Alzheimer's disease. *Mol Neurodegener* **7**,
57, doi:10.1186/1750-1326-7-57 (2012).

61   Franz, A., Ackermann, L. & Hoppe, T. Create and preserve: proteostasis in development
and aging is governed by Cdc48/p97/VCP. *Biochim Biophys Acta* **1843**, 205-215,
doi:10.1016/j.bbamcr.2013.03.031 (2014).

62   Roberts, M. A. & Olzmann, J. A. Protein Quality Control and Lipid Droplet Metabolism.
Annu Rev Cell Dev Biol **36**, 115-139, doi:10.1146/annurev-cellbio-031320-101827 (2020).

63   Bersuker, K. & Olzmann, J. A. Establishing the lipid droplet proteome: Mechanisms of lipid
droplet protein targeting and degradation. *Biochim Biophys Acta Mol Cell Biol Lipids* **1862**,
1166-1177, doi:10.1016/j.bbalip.2017.06.006 (2017).

64   Galluzzi, L., Bravo-San Pedro, J. M., Levine, B., Green, D. R. & Kroemer, G.
Pharmacological modulation of autophagy: therapeutic potential and persisting obstacles.
Nat Rev Drug Discov **16**, 487-511, doi:10.1038/nrd.2017.22 (2017).

65   Zaffagnini, G. *et al.* Phasing out the bad-How SQSTM1/p62 sequesters ubiquitinated
proteins for degradation by autophagy. *Autophagy* **14**, 1280-1282,
doi:10.1080/15548627.2018.1462079 (2018).

66 Baskoylu, S. N. *et al.* Disrupted autophagy and neuronal dysfunction in C. elegans knockin

models of FUS amyotrophic lateral sclerosis. *Cell Rep* **38**, 110195,

doi:10.1016/j.celrep.2021.110195 (2022).

67 Zaffagnini, G. *et al.* p62 filaments capture and present ubiquitinated cargos for autophagy.

EMBO J **37**, doi:10.15252/embj.201798308 (2018).

68 Hansen, M., Flatt, T. & Aguilaniu, H. Reproduction, fat metabolism, and life span: what is

the connection? *Cell Metab* **17**, 10-19, doi:10.1016/j.cmet.2012.12.003 (2013).

69 Silvestrini, M. J. *et al.* Nuclear Export Inhibition Enhances HLH-30/TFEB Activity,

Autophagy, and Lifespan. *Cell Rep* **23**, 1915-1921, doi:10.1016/j.celrep.2018.04.063

(2018).

70 Littlejohn, N. K., Seban, N., Liu, C. C. & Srinivasan, S. A feedback loop governs the

relationship between lipid metabolism and longevity. *eLife* **9**, doi:10.7554/eLife.58815

(2020).

71 Ding, L. *et al.* Peroxisomal beta-oxidation acts as a sensor for intracellular fatty acids and

regulates lipolysis. *Nat Metab* **3**, 1648-1661, doi:10.1038/s42255-021-00489-2 (2021).

72 Schmitz, G. & Grandl, M. Lipid homeostasis in macrophages - implications for

atherosclerosis. *Rev Physiol Biochem Pharmacol* **160**, 93-125,

doi:10.1007/112_2008_802 (2008).

73 Marschallinger, J. *et al.* Lipid-droplet-accumulating microglia represent a dysfunctional

and proinflammatory state in the aging brain. *Nat Neurosci* **23**, 194-208,

doi:10.1038/s41593-019-0566-1 (2020).

74 Nguyen, T. B. *et al.* DGAT1-Dependent Lipid Droplet Biogenesis Protects Mitochondrial

Function during Starvation-Induced Autophagy. *Dev Cell* **42**, 9-21 e25,

doi:10.1016/j.devcel.2017.06.003 (2017).

75 Di Paolo, G. & Kim, T. W. Linking lipids to Alzheimer's disease: cholesterol and beyond.

Nat Rev Neurosci **12**, 284-296, doi:10.1038/nrn3012 (2011).

76 Inloes, J. M. *et al.* The hereditary spastic paraplegia-related enzyme DDHD2 is a principal

brain triglyceride lipase. *Proc Natl Acad Sci U S A* **111**, 14924-14929,

doi:10.1073/pnas.1413706111 (2014).

77 Yang, L. *et al.* Neuronal lipolysis participates in PUFA-mediated neural function and

neurodegeneration. *EMBO Rep* **21**, e50214, doi:10.15252/embr.202050214 (2020).

78 Nuriel, T. *et al.* Neuronal hyperactivity due to loss of inhibitory tone in APOE4 mice lacking

Alzheimer's disease-like pathology. *Nat Commun* **8**, 1464, doi:10.1038/s41467-017-

01444-0 (2017).

79 Labbadia, J. & Morimoto, R. I. The biology of proteostasis in aging and disease. *Annu Rev*

Biochem **84**, 435-464, doi:10.1146/annurev-biochem-060614-033955 (2015).

80 Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71-94 (1974).

81 Stiernagle, T. Maintenance of C. elegans. *WormBook*, 1-11,

doi:10.1895/wormbook.1.101.1 (2006).

82 Kamath, R. S. *et al.* Systematic functional analysis of the Caenorhabditis elegans genome

using RNAi. *Nature* **421**, 231-237 (2003).

83 Hansen, M., Hsu, A. L., Dillin, A. & Kenyon, C. New genes tied to endocrine, metabolic,

and dietary regulation of lifespan from a Caenorhabditis elegans genomic RNAi screen.

PLoS Genet **1**, 119-128 (2005).

84 Holdorf, A. D. *et al.* WormCat: An Online Tool for Annotation and Visualization of

Caenorhabditis elegans Genome-Scale Data. *Genetics* **214**, 279-294,

doi:10.1534/genetics.119.302919 (2020).

Figure 1

Figure 1. SQST-1 dynamics and lifespan modulation is temperature-dependent in *C. elegans*
a-f. Lifespan analysis of wild-type animals and animals over-expressing SQST-1 fused to RFP, GFP or GFP::RFP fed OP50 *E. coli*, developmentally raised at 20°C and then grown at 25°C (**a-c**) or 20°C (**d-f**) during adulthood (n=100). Insets include corresponding representative images of transgenic animals at Day 5 of adulthood, and quantified fluorescence, ±SD *t-test ****p<0.001*. Nt: Non transgenic. **g.** *sqst-1* mRNA levels in wild-type and transgenic animals were quantified by qPCR. Biological triplicates ±SD *t-test *p<0.05, **p<0.01, ***p<0.001* **h.** Levels of GFP and RFP were measured in transgenic tandem SQST-1::GFP::RFP animals (leveraging the pH sensitivity of GFP to visualize SQST-1 accumulation in autolysosome) after incubating Day 1 animals at 20°C, 25°C or 30°C for 24 hours. Average of 10 worms per image. n=5 images per condition ±SD *ANOVA *p<0.05, **p<0.01, ***p<0.001.* Difference in signal between GFP and RFP is due to the low pH sensitivity of GFP. Thus, RFP-only signal represents SQST-1 protein in autolysosomes. **i.** Representative images of transgenic animals (quantified in h.) incubated at 20°C, 25°C or 30°C for 24 hours. Details about lifespan analyses and repeats are available in Supplemental Table 4, Mantel-Cox log-rank. n.s.: not significant, **p<0.01,****p<0.001*.

Figure 2

a

Ribosome: subunit
Transcription: chromatin structure: histone
Ribosome: EIF
Lysosome: Vacuolar ATPase
mRNA functions: processing: snRNP
Cytoskeleton: actin function: actin
Non-coding RNA: tRNA: production
Trafficking: ER/Golgi: COP I
Metabolism: mitochondria: complex V

Legend
10^{-40}
10^{-20}
10^{-10}
10^{-5}
0.001
0.05
NS

b

DHS-3::GFP | SQST-1::RFP | Merge | High Magnification

Control
lgg-1 RNAi

50 µm | 5 µm

Co-localized / Not co-localized
% of total DHS-3::GFP signal
Control / *lgg-1* RNAi Distal Intestine

Co-localized / Not co-localized
% of total SQST-1::RFP signal
Control / *lgg-1* RNAi Distal Intestine

c

SQST-1 modulators (Total = 250)
Lipid droplet proteome (Total = 333)
Insoluble proteins (Total = 201)

51
17
1
44

SQST-1 modulators that aggregate with age and bind lipid droplets

inf-1	*rps-11*
rpl-17	*rps-14*
rpl-19	***rps-16***
rpl-23	*rps-23*
rpl-31	*rps-28*
rpl-7	*rps-3*
rpl-9	*rps-5*
rps-0	***hsp-1***
	pab-1

d

Control | *hsp-1* RNAi | *inf-1* RNAi | *pab-1* RNAi | *rpl-7* RNAi | *rps-16* RNAi

SQST-1::RFP

SQST-1::RFP
Fluorescence (AU)
**** **** **** **** *
Control, hsp-1, inf-1, pab-1, rpl-7, rps-16
RNAi

% of total signal
Head / Gonad / Intestine
Control, hsp-1, inf-1, pab-1, rpl-7, rps-16

Figure 2. SQST-1 regulators include proteins that bind lipid droplets and that aggregate with age

a. Pathway enrichment of genes that regulate SQST-1 levels using WormCat[77]. **b.** SQST-1::RFP accumulates after autophagy is inhibited by *lgg-1* silencing for 3 days during adulthood. Representative confocal microscopy image of the distal intestine of animals expressing both SQST-1::RFP and the lipid droplet-resident protein DHS-3 fused to GFP, subjected to *lgg-1* silencing for 3 days during adulthood. The co-localization between DHS-3::GFP and SQST-1::GFP is quantified. **c.** Overlap of SQST-1 regulators with age-dependent aggregating proteins[2] and lipid droplet-associated proteins[40]. **d.** Representative images of gene silencing of a set of overlapping SQST-1 regulators during adulthood for 3 days in animals over-expressing SQST-1::RFP. Fluorescence and tissue distribution quantified ±SD *ANOVA *p<0.05,****p<0.001*. Details about SQST-1 regulators are available in Supplemental Tables 1 and 2.

Figure 3

Figure 3. Silencing *atgl-1* extends lifespan and reduces SQST-1 accumulation in *C. elegans*
a. Lifespan analysis of wild-type animals and **c.** animals over-expressing SQST-1::RFP raised at 20°C on OP50 *E. coli* and then grown at 25°C during adulthood on control bacteria or bacteria expressing dsRNA against *atgl-1* (n=100). Corresponding images of Day 5 animals stained with Oil-Red-O demonstrating intestinal lipid store accretion under *atgl-1* RNAi (blue borders). **b.** Levels of *atgl-1* mRNA measured by qPCR showing efficient silencing after 3 days on *atgl-1* RNAi (Black: Control RNAi, Blue: *atgl-1* RNAi). Biological triplicates *t*-test *p<0.05* **d.** Representative fluorescence images taken after Day 1 animals over-expressing SQST-1::RFP and DHS-3::GFP were fed control bacteria or bacteria expressing dsRNA against *atgl-1* for 5 days at 25°C. **e-g.** Lifespan analysis of *daf-16(mu86)*, *hlh-30(tm1978),* and *hsf-1(sy441)* mutants raised at 20°C on OP50 *E. coli* and then grown at 25°C during adulthood on control bacteria or bacteria expressing dsRNA against *atgl-1* (n=100). Corresponding images of Day 5 animals stained with ORO demonstrating intestinal lipid store accretion under *atgl-1* RNAi (blue borders). **h.** Levels of autophagosomes and autolysosomes were measured in animals expressing the tandem reporter mCherry::GFP::LGG-1[7] after feeding Day 1 animals with control bacteria or bacteria expressing dsRNA against *atgl-1* for 2 days at 25°C. n=10 per condition ±SD *t*-test #*p<0.06* **i.** Lifespan analysis of *daf-2(e1370)* and *daf-2;SQST-1::GFP* animals raised at 20°C and then grown at 25°C during adulthood on OP50 *E. coli* (with representative image of Day 5 animals, comparative image of wild-type animals in Figure 1b). **j.** *daf-2(e1370)* animals were raised at 20°C on OP50 *E. coli* and then grown at 25°C during adulthood on control bacteria or bacteria expressing dsRNA against *atgl-1* (n=100). Corresponding images of Day 5 animals stained with ORO demonstrating intestinal lipid store accretion under *atgl-1* RNAi (blue borders). Details about lifespan analyses and repeats are available in Supplemental Tables 3 and 4, Mantel-Cox log-rank. n.s.: not significant, **p<0.01,***p<0.001*..

Figure 4

Figure 4. Lipid droplets modulate protein ubiquitination levels and enhance proteostasis
a. Lifespan analysis of wild-type and transgenic animals over-expressing ATGL-1::GFP developmentally raised at 20°C and then grown at 25°C during adulthood on OP50 *E. coli* (n=100). **b.** Levels of RFP signal and lipid droplets from Day 3 animals over-expressing SQST-1::RFP in wild-type or transgenic ATGL-1::GFP over-expressing background. **c.** Wild-type and ATGL-1::GFP over-expressing animals were raised at 20°C and then grown at 25°C during adulthood on control bacteria (Ctl) or bacteria expressing dsRNA against autophagy gene *lgg-1* or proteasome subunit gene *rpn-6.1* for 2 days. Levels of ubiquitinated proteins and actin were visualized by immunAnimals expressing lipid droplet-resident protein DHS-3 fused to GFP were raised at 20°C and then grown at 25°C during adulthood on control bacteria or bacteria expressing RNAi against *lgg-1* or *rpn-6* for 4 days. Levels of ubiquitinated proteins and DHS-3::GFP were visualized by immunoblotting from total input (I), cytosol (C) and lipid droplet (LD) fractions (comparative % loaded between fractions, i.e. 10%). oblotting. Biological replicates shown. **d. e.** Day 1 wild-type animals were fed control bacteria or bacteria expressing dsRNA against *rpn-6* or *atgl-1* for 3 days and ubiquitination levels were detected by immunoblotting. **f.** Levels of RFP signal and lipid droplets in Day 7 *daf-2* animals over-expressing SQST-1::RFP and fed control bacteria or bacteria expressing dsRNA against *lpin-1* during adulthood. **g.** Day 1 *daf-2* animals were fed control bacteria or bacteria expressing dsRNA against *atgl-1, rpn-6* or *lpin-1* for 5 days at 25°C and ubiquitination levels were detected by immunoblotting. **h.** Nematodes expressing intestinally Q44::YFP were grown at at 20°C on OP50 *E. coli* and fed control bacteria or bacteria expressing dsRNA against *atgl-1* at Day 1 of adulthood for 4 days at 25°C, followed by quantification of intestinal aggregates. ±SD *t*-test *$p<0.05$. **i.** Nematodes expressing heat-inducible human Aβ-42 were grown at 20°C on OP50 *E. coli* and fed control bacteria or bacteria expressing dsRNA against *atgl-1* at Day 1 of adulthood for 2 days at 25°C. Paralysis was scored thereafter. Triplicates of n=100 each, ±SD *t*-test *$p<0.05$, **$p<0.01$.

Figure 5. Lipid droplets buffer proteostasis by harboring proteins that challenge protein degradation pathways. When lipid stores are low, ubiquitinated proteins accumulate and eventually overwhelm the proteasomal system and selective autophagy receptors such as SQST-1. High lipid droplet content increases the capacity of cells to stabilize proteins (in conjunction with CDC-48 and HSF-1-regulated chaperones, such as HSP-1) that would otherwise accumulate and aggregate. As proteostasis declines with age, lipid droplets can facilitate autophagic degradation and stabilize the proteome, which prevents the burdening of selective autophagy.