Molecular reconstruction of recurrent evolutionary switching
in olfactory receptor specificity

Lucia L. Prieto-Godino[1,2,*], Hayden R. Schmidt[1] and Richard Benton[1,*]

[1]Center for Integrative Genomics
Faculty of Biology and Medicine
Génopode Building
University of Lausanne
CH-1015
Lausanne
Switzerland

[2]The Francis Crick Institute
London
NW1 1BF
United Kingdom

[*]Corresponding authors:
lucia.prietogodino@crick.ac.uk
Richard.Benton@unil.ch

Abstract

Olfactory receptor repertoires exhibit remarkable functional diversity, but how these proteins have evolved is poorly understood. Through analysis of extant and ancestrally-reconstructed drosophilid olfactory receptors from the Ionotropic Receptor (IR) family, we investigated evolution of two organic acid-sensing receptors, IR75a and IR75b. Despite their low amino acid identity, we identify a common "hotspot" in their ligand-binding pocket that has a major effect on changing the specificity of both IRs, as well as at least two distinct functional transitions in IR75a during evolution. Ligand-docking into IR models predicts that the hotspot does not contact odor molecules, suggesting that this residue indirectly influences ligand/receptor interactions. Moreover, we show that odor specificity is refined by changes in additional, receptor-specific sites, including those outside the ligand-binding pocket. Our work reveals how a core, common determinant of ligand-tuning acts within epistatic and allosteric networks of substitutions to lead to functional evolution of olfactory receptors.

Introduction

Amongst the senses, olfaction is particularly flexible over evolutionary time, enabling animals to adapt their recognition of the vast, ever-changing universe of volatile chemicals in the environment (Bargmann, 2006; Ramdya and Benton, 2010). This flexibility is reflected in the evolution of large, divergent families of olfactory receptors with different odor tuning properties. Several functional surveys of receptor repertoires in vertebrates (e.g., humans and the house mouse *Mus musculus*) and insects (e.g., the vinegar fly *Drosophila melanogaster* and the malaria mosquito *Anopheles gambiae*) have identified ligands for many receptors (Carey et al., 2010; Hallem and Carlson, 2006; Saito et al., 2009). Moreover, comparative sequence and functional analyses of orthologous receptors across species have started to identify amino acid differences that can explain species-specific receptor tuning properties (Adipietro et al., 2012; Auer et al., 2020; Butterwick et al., 2018; Del Mármol et al., 2021; Leary et al., 2012; Mainland et al., 2013; Prieto-Godino et al., 2017; Yang et al., 2017). However, the molecular basis of functional changes in receptors over evolutionary timescales – and whether common principles in this process exist between different receptors – remains unclear.

A powerful model to study olfactory receptor evolution is the Ionotropic Receptor (IR) repertoire, a protostomian chemosensory subfamily of ionotropic glutamate receptors (iGluRs) (Benton et al., 2009; Croset et al., 2010; Ni, 2021; Rytz et al., 2013). Although IRs and iGluRs have limited amino acid sequence identity, their overall conserved (predicted) secondary and tertiary structural organization suggests that the chemosensory receptors share many mechanistic similarities with their iGluR ancestors (Abuin et al., 2011). The best-characterized IRs are predicted to be hetero-tetramers formed of two subunits of a conserved co-receptor and two subunits of a "tuning" receptor (Abuin et al., 2019, 2011). The latter are more variable in sequence both within and between species, particularly in the extracellular ligand-binding domain (LBD), consistent with their diverse odor-recognition properties.

Functional characterization of IRs in different drosophilid species has
revealed orthologous receptors that have distinct odor recognition properties
(Prieto-Godino et al., 2017, 2016), notably IR75a, a receptor that likely originated
in the Neodipteran ancestor (>200 million years ago) and its paralog IR75b, which
arose through duplication of *Ir75a* in the Drosophilidae ancestor (~60-70 million
years ago) (Croset et al., 2010; Prieto-Godino et al., 2017). In *D. melanogaster* and
D. simulans, two cosmopolitan species that feed on a wide range of fermented fruit,
these receptors exhibit different sensitivity towards carboxylic acids: IR75a is tuned
predominantly to acetic acid, while IR75b responds maximally to butyric acid
(Prieto-Godino et al., 2017, 2016; Silbering et al., 2011). By contrast, in the closely-
related island endemic *D. sechellia*, which feeds and breeds exclusively on the ripe
noni fruit of the *Morinda citrifolia* shrub (Figure 1A), IR75a and IR75b preferentially
respond to butyric acid and hexanoic acid, respectively (Prieto-Godino et al., 2017,
2016). These differences are likely to be ecologically significant: for example, acetic
acid is a key product of microbial fermentation of vegetal material and regulates
numerous behaviors in *D. melanogaster* including attraction (Becher et al., 2010)
sexual receptivity (Gorter et al., 2016) and oviposition (Joseph et al., 2009; Kim et
al., 2018). Hexanoic acid is a dominant component of noni fruit and elicits attractive
behaviors in *D. sechellia* (Amlou et al., 1998; Dekker et al., 2006; Prieto-Godino et
al., 2017). In this work, we combine comparative *in vivo* functional analyses of
these receptors across the drosophilid phylogeny, with ancestral sequence
reconstruction, site-directed mutagenesis and protein modelling to investigate their
evolution.
116
117 **Results**
118
119 **Evolution of olfactory responses of IR75a across the drosophilid phylogeny**
120
The distinct responses of *D. sechellia* IR75a (*Dsec*IR75a) compared to orthologs
in its two generalist cousins (Prieto-Godino et al., 2016) suggested that acetic acid-
sensing was the ancestral function of IR75a. We tested this hypothesis by
measuring odor-evoked responses of IR75a-expressing olfactory sensory neurons
(OSNs) across the drosophilid phylogeny, representing >40 million years
divergence time (Figure 1A-C, see Material and methods). As stimuli we used a
panel of linear carboxylic acids spanning from one- to six-carbon chains (hereafter
abbreviated to C1-C6, where acetic acid is C2 and butyric acid is C4). All tested
species within the *melanogaster/obscura* group (except for *D. sechellia*) displayed
strongest responses to C2, similar to *D. melanogaster* and *D. simulans* (Figure 1B-
C). Unexpectedly, the responses of more divergent species were more similar to
those of *D. sechellia*, exhibiting strongest responses to C4 (Figure 1B-C).
To simplify data visualization and discern in an unbiased way which odors
contribute maximally to differential tuning of Ir75a neurons across species, we
performed principal component analysis (PCA) on their response profiles. The first
principal component (PC1) explains 67.5% of the variance in the data, mostly
capturing the inverse variation between C2 and C4 (Figure 1D). When plotting the
C2 and C4 responses against each other, Ir75a neurons of different species
segregated into two clusters with either high responses to C2 and low to C4 or vice
versa (Figure 1E). The clustering matched well the phylogeny with the exception of

D. sechellia, which grouped together with *D. willistoni*, *D. mojavensis* and *D. virilis*
(Figure 1A and 1E). These observations suggested a new model in which the
ancestral drosophilid IR75a was predominantly a C4 sensor that evolved to
become a C2 sensor in the last common ancestor of the melanogaster and obscura
groups, before reverting to an ancestral-like state in *D. sechellia* (Figure 2A).
146
147   **Functional analysis of ancestrally-reconstructed IR75a**
148
To test this hypothesis, we "resurrected" the ancestral IR75a receptors at the
ancestral drosophilid and *melanogaster/obscura* group phylogenetic nodes by
inferring their sequence via maximum likelihood (Randall et al., 2016) from the
sequence of orthologs from sixteen extant species (Figure 2A and Figure 2 – figure
supplement 1 and Materials and methods). We synthesized genes encoding the
inferred ancestral proteins – termed here IR75a^{Dros} and IR75a$^{mel-obs}$, respectively –
and integrated these into a common genomic location to avoid differential positional
influence on their expression. These transgenes were expressed individually in the
D. melanogaster "IR decoder neuron", an OSN that lack the endogenous tuning
receptor subunit but expresses the IR8a co-receptor (Abuin et al., 2011; Grosjean
et al., 2011; Prieto-Godino et al., 2016). When expressed in this system, the tuning
curves of *D. melanogaster* IR75a (*Dmel*IR75a) and *D. sechellia* IR75a (*Dsec*IR75a)
recapitulate the response profile of these receptors expressed in their endogenous
neurons, with strongest responses to C2 and C4, respectively (Figure 2B-C),
consistent with previous observations (Prieto-Godino et al., 2016). As predicted,
IR75a$^{mel-obs}$ responded similarly to *Dmel*IR75a (Figure 2B-C). Importantly, the more
ancient, resurrected receptor, IR75a^{Dros}, had a tuning profile that was almost
identical to *Dsec*IR75a, with maximal responses to C4 (Figure 2B-C). These results
indicate that IR75a has switched tuning profile at least twice during its evolutionary
history, from C4- to C2-sensing and, in *D. sechellia*, back again.
169
170   **Identifying the molecular basis of the functional evolution of IR75a**
171
To determine the molecular basis of the functional evolution of IR75a, we aligned
the sequences of IR75a$^{mel-obs}$ and IR75a^{Dros}. 114 positions in these proteins exhibit
different amino acids (82% identity), of which 45 are located within the LBD (Figure
2 – figure supplement 1). As this level of divergence precluded straightforward
experimental determination of the relevant sites, we first approached the problem
by focusing on the more recent "reverse" transition of C2-sensing to C4-sensing
that occurred on the branch leading to *Dsec*IR75a. We had previously narrowed
down this tuning switch to three amino acid positions within the internal pocket of
the bilobed (S1-S2) LBD (Prieto-Godino et al., 2016) (Figure 3A). Simultaneous
substitution of these sites in *Dmel*IR75a with the residues found in *Dsec*IR75a (i.e.,
T289S, Q536K and F538L) produced a receptor that faithfully recapitulated the
response properties of *Dsec*IR75a when assessed in the IR decoder neuron
(Figure 3B), as previously described (Prieto-Godino et al., 2016). Furthermore,
reverse amino acid substitutions in *Dsec*IR75a (S289T, K536Q and L538L)
conferred response properties characteristic of *Dmel*IR75a (Figure 3B). These
results indicate that the change in tuning is encoded entirely within these three
sites. Thus, there are only $2^3 = 8$ variants to transit between *Dmel*IR75a and

*Dsec*IR75a (including the wild-type sequences) – compared to $2^{114} = 2 \times 10^{34}$
possible variants between IR75a^{Dros} and IR75a$^{mel-obs}$ – offering an excellent
opportunity to study the functional evolution of an olfactory receptor.
We generated versions of *Dmel*IR75a in which each of these three sites
were substituted individually as well the three possible double substitutions. All
single amino acid changes had an impact on receptor responses, shifting tuning
towards C4 (and C3) to different extents, while still retaining some sensitivity to C2
(Figure 3C). Double substitutions showed further shifts towards the *Dsec*IR75a
tuning profiles but to varying degrees: *Dmel*IR75a^{T289S,Q536K} is a broadly-tuned
receptor that responds maximally (albeit weakly) to C2, C3 and C4,
*Dmel*IR75a^{Q536K,F538L} displays maximum responses to both C3 and C4, and
*Dmel*IR75a^{T289S,F538L} responds maximally to C4 alone (Figure 3C). To move beyond
simple descriptions of the effects of these mutations, we used visualizations and
analyses that give insights into the evolutionary landscape of this receptor, as
described below.
204
205 **Mapping the evolutionary landscape of IR75a**
206
Evolutionary landscapes map genotypes onto a quantitative phenotype to illustrate
evolutionary change as a "navigation" process. The roughness of the landscape
determines how accessible each of all possible paths are, and therefore the
likelihood that evolution proceeds through each path (Aguilar-Rodríguez et al.,
2017; Wright, 1932). We reasoned this approach was useful to understand the
evolution of IR75a as the lack of informative intraspecific genetic diversity within
these receptor genes in drosophilid species (data not shown) precluded the direct
identification of evolutionary intermediates.
Responses properties of a receptor are multidimensional, where each odor
defines a dimension. However, PCA of the responses of all *Dmel*IR75a receptor
mutants to all odors revealed that PC1 explains most of the variance (54.5%) and
captures the inverse variation between C2 and C4 – similar to the IR75a neuron
response PCA across the drosophilid phylogeny (Figure 1D) – and, to a lesser
extent, the co-variation of C5 with C4 (Figure 3D). We therefore plotted the
phenotypic data for each single, double and triple *Dmel*IR75a mutant in the C2
versus C4 space alone (Figure 3E) and in the PC1 space (Figure 3F),
acknowledging these simplifications represents only part of the functional changes
(albeit potentially the most important).
The C2 vs C4 plot enables visualization of all possible evolutionary paths by
joining the stepwise mutations from *Dmel*IR75a to *Dmel*IR75a^{T289S,Q536K,F538L}. Most
receptor variants are plotted close to the straight line (i.e., the shortest path) that
joins the initial and final state, with the *Dmel*IR75a^{F538L} single mutant showing the
largest individual "step" along this line (Figure 3E). If we assume that all steps along
this line would be favoured by selection, this observation suggests the hypothesis
that the F538L change might have been the first one of the three substitutions to
occur, as *de novo* mutations with larger effect are typically substituted first followed
by those with smaller effect (Holder and Bull, 2001; Orr, 2005). An alternative
hypothesis is that T289S variant already existed, perhaps as standing variation in
the population: T289S has little phenotypic consequence by itself (Figure 3C,E),
but it greatly augments the effect of the F538L substitution (Figure 3C,E). In this

case, T289S and F538L would have reached fixation together in the same genetic
background because their combined effect allows for a large adaptive leap from
one peak (C2) to the other (C4).The Q536K change alone has an intermediate
phenotype, but we suspect is unlikely to have been the first to occur, as
combination with either other amino acid change leads to a less specific receptor
(*Dmel*IR75a^{Q536K,F538L}, lying to the top-right of the line), or a receptor with overall
weak sensitivity (*Dmel*IR75a^{T289S,Q536K}, lying to the bottom-left of the line) (Figure
3C,E).
To formalize this analysis, we calculated epistasis (*i.e.*, non-additive effects
of different mutations), which can constrain the available evolutionary paths
(Aguilar-Rodríguez et al., 2017; Phillips, 2008). Using the projection of our
physiological responses onto PC1 as our phenotypic readout, we calculated for
each of the combinations of mutants (*i.e.*, single with single, or single with double)
whether the sum of the effects of each of the starting substitutions on the wild-type
background was significantly different from the effect of introducing both mutations
simultaneously (see Materials and methods). Notably, negative magnitude
epistasis was observed in all combinations of Q536K and F538L (whether T289S
was also present the background or not) (Figure 3F, red lines). This observation
supports our previous hypothesis that whether F538L or T289S was the first
mutation to occur, the second one would be the other of these two, because Q536K
incurs in negative epistasis when combined with F538L and it does not lead to a
monotonic increase in responses when combined with T289S (Figure 3F).
Furthermore, this visualization highlights the significant effect on tuning of the
F538L substitution, as well as the minimal (statistically non-significant) effect of the
T289S substitution, supporting its possible segregation in the population without
deleterious effects. It remains unclear what, if any, is the function of Q536K; it is
possible it was fixed by drift or affects some other aspect of receptor activity not
analyzed here.
265
266 **A hotspot for evolution of IR odor tuning**
267
We next asked whether the knowledge of the molecular basis of the C2→C4 tuning
change on the *D. sechellia* branch offers insight into the ancestral switch from C4
to C2 in the last common ancestor of the *melanogaster/obscura* group. Strikingly,
examination of the identity of residues aligned with *Dmel*IR75a F538 in orthologs
across the drosophilid phylogeny revealed a perfect correspondence between the
identity of this position and the best agonist for the receptor: all IR75a orthologues
of species responding most strongly to C2 have an F, while those that respond to
C4, have an L (Figure 4A and Figure 2 – figure supplement 1). Such
correspondence was not seen for amino acid identities at position 289 and 536: for
example, *Dwil*IR75a has a Q at the position equivalent to Q536 of *Dmel*IR75a
(Figure 2 – figure supplement 1), but this species' Ir75a neurons respond to C4,
like *D. sechellia* (Figure 1C).
These observations suggest that position 538 (or equivalent in orthologous
sequences of slightly different lengths) has been a "hotspot" for odor response
evolution, changing from L to F in the *melanogaster/obscura* ancestor and then
changing back on the *D. sechellia* branch. The reversion of amino acid identity in
*Dsec*IR75a is not due to an inverse mutation in the corresponding DNA sequence:

a C→T mutation in codon position one led to the L to F substitution in the
melanogaster/obscura ancestor, while in *D. sechellia* a C→A mutation in codon
position three led, convergently, to restoration of the L-encoding codon (Figure 4A).
The important contribution of a single amino acid in IR75a in determining the
specificity for shorter- or longer-chain acids was reminiscent of our observation of
the evolution of the paralogous receptor IR75b: the difference in tuning of
*Dmel*IR75b and *Dsec*IR75b to C4 and C6, respectively, is determined in large part
by a T523S substitution in the LBD (Prieto-Godino et al., 2017). IR75b and IR75a
exhibit only 38% amino acid identity. However, alignment of these receptors
revealed that position 523 in *Dmel*IR75b corresponds precisely to the 538 hotspot
in IR75a (Figure 4B). Thus, evolution of novel specificities in two different receptors
– which diverged from a common ancestor >60 million years ago (Prieto-Godino et
al., 2017) – is specified by changes in the same site within their LBDs.
To gain insights into the molecular mechanisms underlying the role of the
hotspot in determining the functional properties of IR75a, we performed docking
analysis of ligands in a protein homology model of the *Dmel*IR75a LBD (Prieto-
Godino et al., 2016). The top-ranked pose of C2 (or C4) in this model predicts an
ionic bond of the carboxyl group of this acid with R297 (Figure 4C and Figure 4 –
figure supplement 1). This arginine residue is conserved in all acid-sensing IRs
(Benton et al., 2009), and is essential for odor-evoked activity of IR84a (Abuin et
al., 2011); consistently, a similar interaction was predicted in a model of the IR84a
LBD with its ligand phenylacetic acid (Figure 4 – figure supplement 1). Moreover,
this interaction is equivalent to that of the α-carboxyl group of the glutamate ligand
with a conserved arginine in iGluRs (Figure 4C and figure 4 – figure supplement
1), suggesting a conserved mode of ligand-recognition in IRs and iGluRs. By
contrast, the *Dmel*IR75a hotspot residue F538 is not predicted to interact with
ligand molecules, as it is >4 Å away from any of the top-ranked ligand poses (Figure
4C and figure 4 – figure supplement 1). These observations suggest that the
hotspot is not directly involved ligand binding; how changes in the hotspot might
alter odor response properties of IRs is considered in the Discussion.
315
316   **Functional interactions between the IR75b hotspot and surface residues of**
317   **the LBD**
318
Although mutation of the hotspot in *Dmel*IR75b is sufficient to confer novel
responses to C5 and C6, this single change did not fully recapitulate the response
profile of *Dsec*IR75b, as *Dmel*IR75b^{T523S} retained sensitivity to shorter chain acids
(Figure 5A). Compared to *Dmel*IR75b and *D. simulans* IR75b (*Dsim*IR75b),
*Dsec*IR75b contain three additional changes within the LBD pocket region (P473S,
G492S and A520T; like the hotspot, all are within the S2 lobe), but in our previous
work none of these appeared to contribute substantially – individually or together –
to the refinement of specificity for C6 (Prieto-Godino et al., 2017). The *Dsec*IR75b
LBD contains six additional derived residues located in S1 (Prieto-Godino et al.,
2017), leading us to test whether these sites contribute to the observed changes in
odor responses.
Replacement of the entire S1 lobe in *Dmel*IR75b with that of *Dsec*IR75b
generated a receptor, *Dmel*IR75b^{DsecS1}, with significantly increased responses to
C6 and lower (albeit not statistically significant; Figure 5 source data 1) sensitivity

to C3, when compared to *Dmel*IR75b (Figure 5A). Addition of the hotspot
substitution (T523S) produced a receptor (*Dmel*IR75b^{DsecS1,T523S}) that is more
similar in response profile to *Dsec*IR75b than either *Dmel*IR75b^{DsecS1} or
*Dmel*IR75b^{T523S} (Figure 5A). However, this receptor still has robust sensitivity to
C4, like *Dmel*IR75b but unlike *Dsec*IR75b (Figure 5A). Further incorporation of the
three additional substitutions of residues in the S2 domain (Prieto-Godino et al.,
2017), generated a receptor (*Dmel*IR75b^{DsecS1,4mutS2} that, in terms of specificity, is
indistinguishable from *Dsec*IR75b (Figure 5A). However, this receptor has overall
reduced sensitivity for all acids when compared with *Dsec*IR75b, and its response
to C6 is significantly smaller than that of *Dsec*IR75b (Figure 5A). Despite this
overall reduced sensitivity, *Dmel*IR75b^{DsecS1,4mutS2} still has a significantly increased
sensitivity to C6 when compared with *Dmel*IR75b (Figure 5A).
To visualize how the S1 residues might impact receptor function, we
mapped the position of the six derived changes within this lobe onto a protein
homology model of *Dmel*IR75b (Figure 5B). All of these are located on the surface
of the LBD and therefore unlikely to contact ligands directly. IRs and iGluRs are
thought to exhibit the same global structure and stoichiometry (Abuin et al., 2019,
2011), prompting us to align the modelled *Dsec*IR75b LBD onto a homotetrameric
iGluR structure (PDB 3KG2) (Figure 5C). This analysis did not suggest particularly
close proximity of most of these residues to the interaction interface between
subunits (Figure 5C).
354
355   **Discussion**
356
As species adapt to new ecological niches, olfactory receptors evolve to define new
relationships between external signals and internal neural representations.
Population genetic analyses of *D. melanogaster* strains isolated from diverse global
habitats reveal that olfactory receptors (and other chemosensory protein families)
display some of the strongest genomic signatures of recent selection (Arguello et
al., 2016), suggesting these proteins act as "first responders" in local adaptation to
new environments. In this work we have used comparative sequence and
functional analyses across the well-defined phylogeny of the *Drosophila* genus to
study how members of the organic-acid sensing clade of IRs have changed over
evolutionary timescales.
Our most important finding is the discovery of a "hotspot" residue in these
IRs, whose mutation had a major effect on the odor specificity of one receptor
(IR75a) at two different timepoints during species diversification, as well as on the
tuning change of a distinct receptor (IR75b). For IR75a, our data support a model
in which the ancestral drosophilid IR75a was – contrary to previous assumptions –
a C4 sensor that switched, through mutation at the hotspot (and other sites), to C2
sensing in the *melanogaster/obscura* ancestor before "reverting" to C4-sensing in
D. sechellia. The ancestral function of IR75b is still unclear: we have not been able
to unambiguously identify Ir75b neurons across the drosophilid phylogeny, as they
are not easily distinguishable from those expressing a related receptor, IR75c
(Prieto-Godino et al., 2017). However, like IR75a, IR75b has adapted through
hotspot substitution along the lineage leading to *D. sechellia*. Neither the identity
nor even the chemical class of amino acids occupying the hotspot are conserved
in these receptors (phenylalanine (F, aromatic) or leucine (L, hydrophobic) in

IR75a; threonine (T) or serine (S) (both polar) in IR75b). These observations
indicate that the position of the hotspot in these IRs' LBDs, and not its identity *per*
se, must explain its central role in defining odor-response properties in different
receptors. Studies of other families of olfactory receptors in invertebrates and
vertebrates have revealed enormous inter- and intra-specific sequence variation,
some of which have been linked to differences in odor tuning (Adipietro et al., 2012;
Auer et al., 2020; Block, 2018; Butterwick et al., 2018; Del Mármol et al., 2021;
Leary et al., 2012; Mainland et al., 2013; Prieto-Godino et al., 2017; Yang et al.,
2017). Within this molecular and function diversity, it will be of interest to examine
whether analogous hotspots exist, revealing favored (or constrained) mechanisms
through which evolution selects for new odor-detection properties.
The mechanistic role of the hotspot remains unknown. Our ligand docking
into an IR75a LBD model suggest this residue does not contact ligands directly,
located at a distance of >4 Å away. While we acknowledge the caveats of
interpreting ligand-docking into protein models, our predictions are consistent with
experimentally-determined glutamate/iGluRs interactions, as well as our previous
structure-function analysis of IR84a (Abuin et al., 2011). IR75a orthologs across
the drosophilid phylogeny conserve all three main agonist-binding residues
characteristic of iGluRs (R297, T456, E507 in *Dmel*IR75a) (Benton et al., 2009;
Mayer, 2006), suggesting that the core contacts between odors and this receptor
are unchanged, and that their mutation might be more likely to lead to a non-
functional protein. By contrast, mutation of more distant residues that could, for
example, (de)stabilize ligand-induced LBD conformational changes to alter the
coupling between ligand binding and channel gating, may be a more subtle way of
modifying the tuning profile of these receptors, without the risk of drastic loss of
function. Future determination of the mechanistic impact of hotspot mutations will
be an important priority, which will likely require experimentally-determined
structures of odor-free and odor-bound IR LBDs. Such knowledge may also inform
our understanding of the mechanism of ligand-induced gating in iGluRs, where the
equivalent position (Y732 (Figure 5C)) is thought to have a function in ligand-gating
although its precise role is unclear (Armstrong and Gouaux, 2000; Mamonova et
al., 2008).
While the hotspot is clearly important, its contribution to modification of
tuning properties is shaped by additional changes in these receptors. For IR75a,
there are two additional substitutions within the ligand-binding pocket, while for
IR75b, one or more residues located on the external surface of the LBD are
relevant. In both cases, the functional consequences of combining these
substitutions with that of the hotspot are not easily predicted from their individual
impact, revealing complex epistatic interactions. Moreover, the distance of these
additional sites from the predicted odor-binding site, suggest the existence of
allosteric effects of certain sites on odor/receptor interactions. These results are of
interest in light of molecular evolutionary analyses of olfactory receptor repertoires,
which have identified numerous residues under positive selection (or relaxed
purifying selection) in different receptors – implying a contribution to functional
divergence – in regions far from the predicted ligand-binding pocket (Arguello et
al., 2016; Chen et al., 2010; Gardiner et al., 2009; Smadja et al., 2009; Steiger et
al., 2010). Together such observations argue that the evolution of olfactory receptor
specificity does not simply arise by alterations in direct interactions of receptors

with odor ligands, but rather can emerge from a complex network of interactions of amino acid substitutions – with major or minor effects – both near and far from the ligand-binding pocket.

One clear limitation of our study is the restriction of our profiling to a set of linear carboxylic acids at a single concentration. While these ligands are found in nature and are the best agonists for these receptors among many screened odors (Silbering et al., 2011), they necessarily only give a partial insight into the functional changes of individual receptors. The C4 to C6 switch of IR75b in *D. sechellia* is likely related to high abundance of C6 in its sole host fruit (noni). For IR75a the ecological framework is less clear: we speculate that the C4 to C2 switch in the *melanogaster*/*obscura* ancestor may be related to the use of host fruits with acetic-acid producing bacteria, which may then have been no longer relevant for *D. sechellia* which preferentially feeds upon ripe (non-fermenting) noni fruit (which contains much more C4 than C2 (Auer et al., 2020; Farine et al., 1996)). Future expansion of the odor profiling of these species' receptors will be essential to understand the pressures in the natural world that have selected for olfactory receptor proteins with new chemical recognition properties. Such knowledge, together with mechanistic insight into ligand/receptor interactions, may be useful to re-engineer the ligand-binding specificities of receptors (for example, as chemogenetic tools (Fukabori et al., 2020)) or design pharmacological manipulators to control the olfactory-guided behaviors of pest insects.

Acknowledgements

We thank Roman Arguello, Jean Hausser, Michael Shahandeh and other members of the Benton laboratory for discussions and comments on the manuscript. We thank Tom Baden for help with Igor analysis and Liliane Abuin and Steeve Cruchet for technical assistance. H.R.S. was supported by an EMBO Long-Term Fellowship (ALTF 940-2019) and a Helen Hay Whitney Foundation Fellowship. L.L.P.-G.'s laboratory was supported by a European Research Council (ERC) Starting Investigator Grant (802531), and the Francis Crick Institute, which receives its core funding from Cancer Research UK (FC001594), the UK Medical Research Council (FC001594), and the Wellcome Trust (FC001594). Work in R.B.'s laboratory was supported by ERC Consolidator and Advanced Grants (615094 and 833548, respectively), an HFSP Young Investigator Award (RGY0073/2011), the Swiss National Science Foundation Nano-Tera Envirobot project (20NA21_143082). For the purpose of Open Access, the authors have applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

Author contributions

L.L.P.-G. and R.B. conceived the project. L.L.P.-G. designed and performed all experiments and analyses, except for the ligand docking studies, which were performed by H.R.S. All authors contributed to interpretation of results. L.L.P.-G. and R.B. wrote the paper, with contributions from H.R.S.

Materials and methods

Drosophila strains and culture

Flies were maintained at 25ºC in 12 h light:12 h dark conditions. We used the following published *D. melanogaster* strains: *Ir84a^{Gal4}* (Grosjean et al., 2011), *UAS-DmelIr75a, UAS-DsecIr75a, UAS-DmelIr75a^{T289S,Q536K,F538L}* (Prieto-Godino et al., 2016). Other drosophilid species were obtained from the *Drosophila* Species Stock Center: *D. sechellia* (14021-0248.25), *D. simulans* (14021-0251.195), *D. yakuba* (14021-0261.01), *D. erecta* (14021-0224.01), *D. ananassae* (14024-0371.13), *D. pseudoobscura* (14011-0121.94), *D. willistoni (*14030-0811.24), *D. mojavensis* (15081-1352.22), *D. virilis* (15010-1051.87).

Molecular biology

cDNAs of *D. melanogaster* and *D. sechellia Ir75a* and *Ir75b* were previously described (Prieto-Godino et al., 2017, 2016). Site-directed mutagenesis was performed using standard procedures, and mutant cDNAs were subcloned into *pUAST attB* for transgenesis of *D. melanogaster* using the phiC31 site-specific integration system (landing site attP40) by BestGene and Genetic Services Inc. All transgenes were sequence-verified both before and after transformation.

Ancestral protein reconstruction

The 16 "modern" sequences of IR75a shown in Figure 2 – figure supplement 1 were used to computationally infer the ancestral sequences at all of the nodes of the tree using the known phylogeny. Ancestral sequences were calculated using the bio-informatic tool FastML (Ashkenazy et al., 2012) with rate variation modelled as a gamma distribution. The marginal posterior probability for most of the amino acids of all of the reconstructed nodes was above 0.8. For the hot-spot mutation the marginal posterior probability was 1. To ascertain the robustness of the inferences made by FastML, we also inferred ancestral sequences of the nodes using a different bioinformatic tool, codeml within the PAML package. The inferred sequences had a global identity of 96.5% and were fully identical at the three amino acid substitutions under study. DNA sequences encoding the IR75a^{Dros} and the IR75a^{obs-mel} sequences predicted by FastML were synthesized by Eurofins-Genomics, subcloned into *pUAST attB* and transformed into flies as described above.

In vivo electrophysiology

Single sensillum extracellular recordings were performed essentially as described in (Benton and Dahanukar, 2011; Prieto-Godino et al., 2017). For all stimuli, 10 µl odor (10% v/v in water) were use, and presented to the animal in a 1 s pulse. ac2 sensilla were identified in different species by targeting sensilla at antennal locations where ac2 are found in *D. melanogaster*, and using pyridine as a diagnostic odor, which activates the ac2 neuron expressing IR41a and appears to be conserved across species (Figure 1B). CAS numbers and sources of odors are

as follows: formic acid (64-18-6), acetic acid (64-19-7), propionic acid (79-09-4),
butyric acid (107-92-6), pentanoic acid (109-52-4), hexanoic acid (142-62-1),
pyridine (110-86-1) (diagnostic for ac2), octanol (111-87-5) (diagnostic for ac3).
Odor-evoked responses were calculated by summing the activity of all OSNs in a
sensillum to a given stimulus, as reliable spike-sorting is not possible and other
neurons housed in the ac2 or ac4 sensilla do not respond to acidic odors (Silbering
et al., 2011). We counted the number of spikes in a 500 ms window at stimulus
delivery (200 ms after stimulus onset due to a delay introduced by the air path),
subtracted the number of spikes in a 500 ms window 2 s before stimulus delivery,
and doubled the result to obtain spikes/s. To calculate solvent-corrected responses
(as shown in the figures), we subtracted from the response to each diluted odor,
the response obtained when stimulating with the corresponding solvent (water for
all odors except for pyridine and octanol, which were dissolved in paraffin oil (8012-
95-1). A maximum of three sensilla were tested per animal, and individual
genotypes were measured, in an interleaved fashion, on multiple independent
days. Normalized responses were calculated by dividing solvent-corrected
responses of a given sensillum to each odor by the maximal response of that
sensillum, such that each sensillum always had one odor whose normalized
response was 1. A pre-requisite for PCA analysis is to input z-scored responses;
these were calculated with the in-built function of MATLAB, which works according
to the definition of z-scoring, i.e., across each sensillum recording, each response
was subtracted from the mean response of all sensillar responses and divided by
the standard deviation.
545
546 **Statistical analysis**
547
Principal component analysis was carried with in-built MATLAB functions.
Statistical analyses were carried out with in-built functions in R Studio or Igor. For
all statistical tests, a Shapiro test for normally was first performed. If both samples
being compared were normally distributed, a t-test was performed; if one of the
samples was not normally distributed, a Wilcoxon test was run. When performing
multiple comparisons, p-values were corrected using the Bonferroni method.
Epistasis was calculated by determining whether the effects of each individual
mutation were added linearly by comparing the observed responses of the
combination of two genetic manipulations with the expected distribution of
individual mutations were added linearly. Briefly, to generate expected distributions
for the linear combination of two mutations, we subtracted the mean responses of
the initial receptor from the two "intermediate" receptors, this provides the "effect"
of each mutation(s) individually, we then we took 1000 random samples from the
"effects" of each of the two mutations and summed them to generate a distribution
of the expected responses if the two effects of each mutation added linearly. To
determine whether two mutations interacted epistatically, we statistically compared
the expected distribution with the actual distribution (the double manipulation
combination) using a Wilcoxon test. If the result of this test was $p < 0.05$ after
correction for multiple comparisons using Bonferroni method, the two mutations
were considered to interact epistatically.
568
569

Ligand docking

AutoDock Vina version 1.1.2 (Trott and Olson, 2009) was used to dock ligands into homology models of the LBDs of *Dmel*IR75a and *Dmel*IR84a (K205-N354-L451-W583) (Prieto-Godino et al., 2016) . The crystal structure of *Rnor*GluA2 bound to glutamate (PDB accession 1FTJ) was used to optimize the grid conditions and docking procedure for IRs. A12×12×12 Å grid was placed along the axis between the CZ atom in a conserved arginine residue (R96 in *Rnor*GluA2, R100 in *Dmel*IR75a) and the CD atom in a conserved glutamate residue (E193 in *Rnor*GluA2, E197 in the IR75a, this glutamate is not conserved in *Dmel*IR84a). The center of the grid was placed closer to the arginine than to the glutamate, such that if a straight line was drawn from the arginine to the glutamate, the grid center was one-third of the way along this line. For *Dmel*IR84a, the conserved glutamate is not present, and the position of the grid was manually determined using the conserved arginine (R317) as a guide and the grids for RnGluA2 and IR75a as reference. Docking was performed using an exhaustiveness of 8, with the default settings for AutoDock Vina; these were sufficient to accurately predict the position of the co-crystallized glutamate in RnGluA2 (Figure 4 – figure supplement 1). Ligand poses were inspected manually, but in all cases the top-ranked pose was judged to be the most plausible. Structure visualization was performed using PyMol version 2.3.3. The *Dsec*IR75b LBD model (Prieto-Godino et al., 2017) was aligned to the GluA2 homotetramer using the "align" command in PyMol.

Figure Legends

Figure 1. Evolution of olfactory responses of Ir75a neurons across the
drosophilid phylogeny.
(**A**) Phylogeny of analyzed drosophilid species and their known ecological niches
(Markow, 2015).
(**B**) Representative traces of extracellular recordings of neuronal responses in
antennal coeloconic 2 (ac2) sensilla (schematized top left) to the indicated odors in
different drosophilid species. ac2 sensilla were identified based on their
morphology, their location on the antennal surface – either near the entry to the
sacculus or in the distal posterior part intermingled with ac3 sensilla (Silbering et
al., 2011) – their pattern of basal firing, and the conserved responses of the Ir41a
neuron towards pyridine and the lack of response to octanol (which is detected by
the Or35a neuron in ac3 sensilla (Yao et al., 2005)).
(**C**) Heatmap of electrophysiological solvent-corrected responses (see Materials
and methods) of ac2 sensilla of all species in (**A**) to a series of linear carboxylic
acids; the color-scale is on the right. Each rectangle represents the responses
measured in a single sensillum.
(**D**) Loadings of the first principal component of all responses shown in (**C**).
(**E**) Responses of ac2 sensilla for all species plotted as z-scored responses (see
Material and methods) of C2 against z-scored responses to C4. This plotting
reveals two clear clusters with species responding maximally to C2 (orange) or C4
(blue).

Figure 1 Source data 1. Data for Figure 1 showing the solvent-corrected
spikes/second responses of ac2 sensilla for each of the species (Figure 1C). The
same responses z-scored (Figure 1E) and the results of the PCA analysis, showing
loadings and variance explained by each PC (Figure 1D). Here and in subsequent
source data files "NaN" ("not a number") indicates cases where data was not
available, usually due to the electrode coming out of the sensillum before the end
of the series of stimulations.

Figure 2. Ancestral sequence reconstruction of IR75a.
(**A**) Phylogeny of the protein sequences of drosophilid IR75a orthologs used for the
reconstruction of ancestral nodes (IR75a^{Dros} and IR75a$^{mel-obs}$) (see Figure 2 – figure
supplement 1). The orange and blue dots indicate maximal responses of the
corresponding neurons to C2 and C4, respectively (Figure 1C). The branches of
the tree have been similarly color-coded according to predicted receptor responses
based on parsimony.
(**B**) Representative traces of extracellular recordings of neuronal responses to the
indicated odors of receptors expressed in the IR decoder neuron (see text).
Genotypes are of the form: *UAS-xxx/UAS-xxx;Ir84a^{Gal4}/Ir84a^{Gal4}*, here and in all
subsequent figures.
(**C**) Quantification of responses of the indicated receptors expressed in the IR
decoder neuron. In this and subsequent similar figure panels, the top row shows
barplots of responses normalized by maximal response, and the bottom row shows
individual datapoints, mean and SEM of raw solvent-corrected responses to odor
stimuli.

641
Figure 2 – figure supplement 1. Alignment of IR75a orthologs.
Multiple sequence alignment of IR75a orthologs used for the ancestral protein
reconstruction. The ancestral versions that were functionally tested, IR75a^{Dros} and
IR75a$^{obs-mel}$, are also shown. Dark blue indicates identical residues, light blue
indicates similar residues, grey indicates dissimilar residues. The red lines indicate
the spans of S1 and S2 lobes of the LBD. The amino acids colored in red are those
positions that were estimated by FASTML (see Materials and methods) to have a
marginal posterior probability < 0.7.
650
651 **Figure 2 Source data 1.** Data for Figure 1C, responses in spikes/s of ac2 sensilla
652 from each of the genotypes, and normalized with respect the maximal response of
653 each sensilla.
654
Figure 3. Mapping the evolutionary landscape of IR75a.
(**A**) Top: cartoon of the domain organization of IR75a; bottom: protein homology
model of the *Dmel*IR75a LBD (adapted from (Prieto-Godino et al., 2016)). The three
functionally-important amino acid positions that differ between *Dmel*IR75a and
*Dsec*IR75a are highlighted in red.
(**B-C**) Quantification of responses of the indicated receptor versions expressed in
the IR decoder neuron.
(**D**) Loadings of the first principal component of all responses shown in (**C**).
(**E**) Responses of each of the indicated receptor versions plotted in the C2 vs C4
space.
(**F**) Visualization of epistasis and accessible mutational pathways. Each of the
individual, double and triple mutations are plotted in the PC1 axis (error bars are
standard error of the mean). Possible evolutionary paths join these with lines. Solid
lines indicate when the path joins two points that significantly increase PC1 value
(i.e. increased responses to C4 and C5 and decreased responses to C2); dashed
lines denote paths that, while accessible, do not lead to significantly increased PC1
values (see Source Data, for statistical values). Red lines indicate the cases where
two mutations interact epistatically when combined, *i.e.*, the combination of the two
mutations is not equal to the expected response if their effects added linearly (see
Materials and methods for details and Source Data for statistical values).
675
676 **Figure 3 Source data 1.** First tab: Data for Figure 3B and 3C, responses in
677 spikes/second of ac2 sensilla from each of the genotypes indicated, and
678 normalized with respect the maximal response of each sensilla. Second tab:
679 Results from the PCA analysis. Third tab: P-values resulting from the statistical
680 analysis in Figure 3F.
681
Figure 4. A hotspot for tuning IR sensitivity.
(**A**) Phylogeny of species' IR75a receptors used for the reconstruction of ancestral
nodes. The inferred identity of hotspot mutation for key nodes is shown as well as
for each of the extant receptors at the end of each leaf, together with the measured
(solid circle) or predicted (empty circle) sensitivity. On the right, the nucleotides
encoding the hotspot amino acid position in IR75a across species. *Drosophila*
species abbreviations (where not presented in Figure 1): *Dmir* (*miranda*), *Dper*

(*persimilis*), *Dame* (*americana*), *Dbuz* (*buzzatii*), *Dgri* (*grimshawi*), *Dalb*
(*albomicans*).
(**B**) Top: Protein sequence alignment of *Dmel*IR75a and *Dmel*IR75b. Dark blue
indicates identical residues, light blue indicates similar residues, grey indicates
dissimilar residues. The red lines indicate the spans of S1 and S2 lobes of the LBD.
The red box indicates the common amino acid position ("hotspot"). Bottom:
separate alignments of the hotspot region in IR75a and IR75b for *D. melanogaster*
and *D. sechellia* proteins showing that while the residue position is conserved, the
identity of the amino acids is different for these two receptors.
(**C**) Left: homology model of *Dmel*IR75a LBD (blue) with docked ligands; the insets
show the top-ranked poses for C2 (cyan) and C4 (purple) in the LBD pocket. Right:
crystal structure of the *Rattus norvegicus* (*Rnor*) iGluR GluA2 LBD (gray) in
complex with glutamate (yellow) (PDB accession 1FTJ). The hotspot residue is
shown in orange in *Dmel*IR75a, and the equivalent residue in *Rnor*GluA2 in
magenta. Amino acid numbering corresponds to that of the full-length sequence.
704
Figure 4 – figure supplement 1. Ligand docking in IRs and iGluRs.
Top-three docked poses for the indicated ligands in the *Dmel*IR75a and *Dmel*IR84a
LBD homology models, and the *Rnor*GluA2/glutamate crystal structure (PDB
1FTJ), ranked by AutoDock Vina docking score. In *Dmel*IR75a, all three poses of
C2 occupy a similar space in the binding pocket, contacting the conserved arginine
R297 and >4 Å from the hotspot residue 538 (orange). Similarly, in *Dmel*IR84a, all
three poses of phenylacetic acid occupy a similar position in the binding pocket and
make contact with the conserved arginine R317, which is essential for ligand-
evoked currents from IR84a *in vivo* (Abuin et al., 2011). In *Rnor*GluA2, the docked
poses of docked glutamate (green), particularly the top-ranked pose, are similar to
the positioning of co-crystallized glutamate (yellow). Amino acid numbering
corresponds to that of the full-length sequences.
717
Figure 5. Epistasis in the ligand binding domain of IR75b.
(**A**) Quantification of responses of the indicated wildtype and mutated IR75b
variants expressed in the IR decoder neuron. Note that the *Dmel*IR75b[DsecS1] variant
includes the six derived changes between *D. sechellia* and *D. simulans/D.*
melanogaster as well as a seventh amino acid substitution common to *D. sechellia*
and *D. simulans* receptors.
(**B**) Protein homology model of the *Dsec*IR75b LBD (adapted from (Prieto-Godino
et al., 2017)). The S1 and S2 lobes are colored pale and dark green, respectively.
Surface-located amino acids that differ between *Dmel*IR75b and *Dsec*IR75b in the
S1 domain are depicted in magenta; the hotspot (position 523) and other residues
in the ligand-binding pocket in the S2 domain are depicted in red.
(**C**) Homology model of the *Dmel*IR75b LBD (colored as in (**B**)) aligned to the full-
length homotetrameric crystal structure of *Rnor*GluA2 (PDB accession 3KG2),
which is colored by chain in faded gray, red, blue, and yellow. The *Rnor*GluA2
amino-terminal domain (which is not present in most IRs (Rytz et al., 2013)) has
been removed for clarity.
734
735 **Figure 5 Source data 1.** First tab: Data for Figure 5A, responses in spikes/second
736 of ac2 sensilla from each of the genotypes indicated, and normalized with respect

737 to the maximal response of each sensilla. Second tab: P-values resulting from the
738 statistical analysis, as discussed in the text.
739
740 **Supplementary file 1. Nucleotide sequences encoding the reconstructed**
741 **ancestral IR75a orthologs.**
742

743 **References**
744

Abuin L, Bargeton B, Ulbrich MH, Isacoff EY, Kellenberger S, Benton R. 2011.
Functional Architecture of Olfactory Ionotropic Glutamate Receptors. *Neuron*.
Abuin L, Prieto-Godino LL, Pan H, Gutierrez C, Huang L, Jin R, Benton R. 2019.
In vivo assembly and trafficking of olfactory Ionotropic Receptors. *BMC Biol*
17. doi:10.1186/s12915-019-0651-7
Adipietro KA, Mainland JD, Matsunami H. 2012. Functional evolution of
Mammalian odorant receptors. *PLoS Genet* **8**:e1002821.
doi:10.1371/journal.pgen.1002821
Aguilar-Rodríguez J, Payne JL, Wagner A. 2017. A thousand empirical adaptive
landscapes and their navigability. *Nat Ecol Evol*. doi:10.1038/s41559-016-
0045
Amlou M, Moreteau B, David JR. 1998. Genetic analysis of Drosophila sechellia
specialization: Oviposition behavior toward the major aliphatic acids of its
host plant. *Behav Genet*. doi:10.1023/A:1021689312582
Arguello JR, Cardoso-Moreira M, Grenier JK, Gottipati S, Clark AG, Benton R.
2016. Extensive local adaptation within the chemosensory system following
Drosophila melanogaster's global expansion. *Nat Commun* **7**:11855.
doi:10.1038/ncomms11855
Armstrong N, Gouaux E. 2000. Mechanisms for Activation and Antagonismof an
AMPA-Sensitive Glutamate Receptor: Crystal Structures of the GluR2 Ligand
Binding Core. *Neuron*.
Ashkenazy H, Penn O, Doron-Faigenboim A, Cohen O, Cannarozzi G, Zomer O,
Pupko T. 2012. FastML: A web server for probabilistic reconstruction of
ancestral sequences. *Nucleic Acids Res*. doi:10.1093/nar/gks498
Auer TO, Khallaf MA, Silbering AF, Zappia G, Ellis K, Álvarez-Ocaña R, Arguello
JR, Hansson BS, Jefferis GSXE, Caron SJC, Knaden M, Benton R. 2020.
Olfactory receptor and circuit evolution promote host specialization. *Nature*.
doi:10.1038/s41586-020-2073-7
Bargmann CI. 2006. Comparative chemosensation from receptors to ecology.
Nature. doi:10.1038/nature05402
Becher PG, Bengtsson M, Hansson BS, Witzgall P. 2010. Flying the fly: Long-
range flight behavior of drosophila melanogaster to attractive odors. *J Chem
Ecol*. doi:10.1007/s10886-010-9794-2
Benton R, Dahanukar A. 2011. Electrophysiological recording from Drosophila
olfactory Sensilla. *Cold Spring Harb Protoc*. doi:10.1101/pdb.prot5630
Benton R, Vannice KS, Gomez-Diaz C, Vosshall LB. 2009. Variant ionotropic
glutamate receptors as chemosensory receptors in Drosophila. *Cell*.
Block E. 2018. Molecular Basis of Mammalian Odor Discrimination: A Status
Report. *J Agric Food Chem*. doi:10.1021/acs.jafc.8b04471

Butterwick JA, del Mármol J, Kim KH, Kahlson MA, Rogow JA, Walz T, Ruta V.
2018. Cryo-EM structure of the insect olfactory receptor Orco. *Nature*.
doi:10.1038/s41586-018-0420-8
Carey AF, Wang G, Su C-YY, Zwiebel LJ, Carlson JR. 2010. Odorant reception in
the malaria mosquito Anopheles gambiae. *Nature* 1–7.
doi:10.1038/nature08834
Chen M, Peng ZG, He SP. 2010. Olfactory receptor gene family evolution in
stickleback and medaka fishes. *Sci China Life Sci*. doi:10.1007/s11427-010-
0025-4
Croset V, Rytz R, Cummins SF, Budd A, Brawand D, Kaessmann H, Gibson TJ,
Benton R. 2010. Ancient protostome origin of chemosensory ionotropic
glutamate receptors and the evolution of insect taste and olfaction. *PLoS*
Genet.
Dekker T, Ibba I, Siju KP, Stensmyr MC, Hansson BS. 2006. Olfactory shifts
parallel superspecialism for toxic fruit in Drosophila melanogaster sibling, D.
sechellia. *Curr Biol* **16**:101–109. doi:10.1016/j.cub.2005.11.075
Del Mármol J, Yedlin M, Ruta V. 2021. The structural basis of odorant recognition
in insect olfactory receptors. *bioRxiv*.
Farine J-PP, Legal L, Moreteau B, Le Quere jean-LL. 1996. Volatile components
of ripe fruits of Morinda citrifolia and their effects on Drosophila.
Phytochemistry 1–6. doi:10.1016/0031-9422(95)00455-6
Fukabori R, Iguchi Y, Kato S, Takahashi K, Eifuku S, Tsuji S, Hazama A,
Uchigashima M, Watanabe M, Mizuma H, Cui Y, Onoe H, Hikishima K,
Yasoshima Y, Osanai M, Inagaki R, Fukunaga K, Nishijo T, Momiyama T,
Benton R, Kobayashi K. 2020. Enhanced retrieval of taste associative
memory by chemogenetic activation of locus coeruleus norepinephrine
neurons. *J Neurosci*. doi:10.1523/JNEUROSCI.1720-20.2020
Gardiner A, Butlin RK, Jordan WC, Ritchie MG. 2009. Sites of evolutionary
divergence differ between olfactory and gustatory receptors of Drosophila.
Biol Lett. doi:10.1098/rsbl.2008.0723
Gorter JA, Jagadeesh S, Gahr C, Boonekamp JJ, Levine JD, Billeter J-CC. 2016.
The nutritional and hedonic value of food modulate sexual receptivity in
Drosophila melanogaster females. *Sci Rep* **6**:19441. doi:10.1038/srep19441
Grosjean Y, Rytz R, Farine J-P, Abuin L, Cortot J, Jefferis GSXE, Benton R.
2011. An olfactory receptor for food-derived odours promotes male courtship
in Drosophila. *Nature*.
Hallem EA, Carlson JR. 2006. Coding of Odors by a Receptor Repertoire. *Cell*
125:143–160. doi:10.1016/j.cell.2006.01.050
Holder KK, Bull JJ. 2001. Profiles of adaptation in two similar viruses. *Genetics*.
Joseph RM, Devineni A V., King IFGG, Heberlein U. 2009. Oviposition preference
for and positional avoidance of acetic acid provide a model for competing
behavioral drives in Drosophila. *Proc Natl Acad Sci U S A* **106**:11352–11357.
doi:10.1073/pnas.0901419106
Kim G, Huang JH, McMullen JG, Newell PD, Douglas AE. 2018. Physiological
responses of insects to microbial fermentation products: Insights from the
interactions between Drosophila and acetic acid. *J Insect Physiol*.
doi:10.1016/j.jinsphys.2017.05.005
Leary GP, Allen JE, Bunger PL, Luginbill JB, Linn CE, Macallister IE, Kavanaugh

MP, Wanner KW. 2012. Single mutation to a sex pheromone receptor
provides adaptive specificity between closely related moth species. *Proc Natl*
Acad Sci U S A **109**:14081–14086. doi:10.1073/pnas.1204661109

Mainland JD, Keller A, Li YR, Zhou T, Trimmer C, Snyder LL, Moberly AH,
Adipietro KA, Liu WLL, Zhuang H, Zhan S, Lee SS, Lin A, Matsunami H.
2013. The missense of smell: Functional variability in the human odorant
receptor repertoire. *Nat Neurosci* **17**:114–120. doi:10.1038/nn.3598

Mamonova T, Speranskiy K, Kurnikova M. 2008. Interplay between structural
rigidity and electrostatic interactions in the ligand binding domain of GluR2.
Proteins Struct Funct Genet **73**:656–671. doi:10.1002/prot.22090

Markow TA. 2015. The secret lives of Drosophila flies. *Elife* **4**.
doi:10.7554/eLife.06793

Mayer ML. 2006. Glutamate receptors at atomic resolution. *Nature*.
doi:10.1038/nature04709

Ni L. 2021. The Structure and Function of Ionotropic Receptors in Drosophila.
Front Mol Neurosci. doi:10.3389/fnmol.2020.638839

Orr HA. 2005. The genetic theory of adaptation: A brief history. *Nat Rev Genet*.
doi:10.1038/nrg1523

Phillips PC. 2008. Epistasis - The essential role of gene interactions in the
structure and evolution of genetic systems. *Nat Rev Genet*.
doi:10.1038/nrg2452

Prieto-Godino LL, Rytz R, Bargeton B, Abuin L, Arguello JR, Peraro MD, Benton
R. 2016. Olfactory receptor pseudo-pseudogenes. *Nature* **539**.
doi:10.1038/nature19824

Prieto-Godino LL, Rytz R, Cruchet S, Bargeton B, Abuin L, Silbering AF, Ruta V,
Dal Peraro M, Benton R. 2017. Evolution of Acid-Sensing Olfactory Circuits
in Drosophilids. *Neuron* **93**. doi:10.1016/j.neuron.2016.12.024

Ramdya P, Benton R. 2010. Evolving olfactory systems on the fly. *Trends Genet*.

Randall RN, Radford CE, Roof KA, Natarajan DK, Gaucher EA. 2016. An
experimental phylogeny to benchmark ancestral sequence reconstruction.
Nat Commun. doi:10.1038/ncomms12847

Rytz R, Croset V, Benton R. 2013. Ionotropic Receptors (IRs): Chemosensory
ionotropic glutamate receptors in Drosophila and beyond. *Insect Biochem*
Mol Biol.

Saito H, Chi Q, Zhuang H, Matsunami H, Mainland JD. 2009. Odor coding by a
mammalian receptor repertoire. *Sci Signal*. doi:10.1126/scisignal.2000016

Silbering AF, Rytz R, Grosjean Y, Abuin L, Ramdya P, Jefferis GSXE, Benton R.
2011. Complementary function and integrated wiring of the evolutionarily
distinct Drosophila olfactory subsystems. *J Neurosci*.

Smadja C, Shi P, Butlin RK, Robertson HM. 2009. Large gene family expansions
and adaptive evolution for odorant and gustatory receptors in the pea aphid,
Acyrthosiphon pisum. *Mol Biol Evol*. doi:10.1093/molbev/msp116

Steiger SS, Fidler AE, Mueller JC, Kempenaers B. 2010. Evidence for adaptive
evolution of olfactory receptor genes in 9 bird species. *J Hered*.
doi:10.1093/jhered/esp105

Trott O, Olson AJ. 2009. AutoDock Vina: Improving the speed and accuracy of
docking with a new scoring function, efficient optimization, and
multithreading. *J Comput Chem*. doi:10.1002/jcc.21334

Wright S. 1932. The roles of mutation, inbreeding, crossbreeding and selection in
evolution. *Sixth Int Congr Genet*.
Yang K, Huang LQ, Ning C, Wang CZ. 2017. Two single-point mutations shift the
ligand selectivity of a pheromone receptor between two closely related moth
species. *Elife*. doi:10.7554/eLife.29100
Yao CA, Ignell R, Carlson JR. 2005. Chemosensory coding by neurons in the
coeloconic sensilla of the Drosophila antenna. *J Neurosci* **25**:8359–8367.
doi:10.1523/JNEUROSCI.2432-05.2005
888

*Dmel*IR75a LBD
homology model
+
C2

*Dmel*IR84a LBD
homology model
+
phenylacetic acid

*Rnor*GluA2 LBD/glutamate
crystal structure
+
docked glutamate