Community level individual-based plant-pollinator networks unveil pollen flow dynamics and plant reproductive success

Alfonso Allen-Perkins[1,2,✉], **María Hurtado**[3], **David García-Callejas**[1,3], **Oscar Godoy**[3], **and Ignasi Bartomeus**[1]

[1]Departamento de Ecología Integrativa, Estación Biológica de Doñana, EBD-CSIC, Avda. Amrico Vespucio 26, Seville E-41092, Spain.
[2]Departamento de Ingeniería Eléctrica, Electrónica, Automática y Física Aplicada, ETSIDI, Universidad Politécnica de Madrid, 28040 Madrid, Spain.
[3]Departamento de Biología, Instituto Universitario de Investigación Marina (INMAR), Universidad de Cádiz, Puerto Real, E-11510, Spain.

Ecological networks are a widely used tool to understand the dynamics of ecological communities in which plants interact with their pollinator counterparts. However, while most mutualistic networks have been defined at the species level, ecological processes, such as pollination, take place at the individual level. This recognition has led to the development of individual-based networks, yet current approaches only account for individuals of a single plant species due to conceptual and mathematical limitations. Here, we introduce a multilayer framework designed to depict the conspecific and heterospecific pollen flows mediated by floral visitors among plant individuals belonging to different species. Pollen transfer is modeled as a transport-like system, where an ensemble of conspecific plant-pollinator "circuits" are coupled through pollinators. With this physical conceptualization of ecological processes, we investigate how the reproductive success of plant individuals is affected by the overall dynamics of the whole multilayer network (macrostructure), as well as by their local position within the network (mesostructure). To illustrate this multiscale analysis, we apply it to a dataset of nine well-resolved individual plant-pollinator interaction networks from annual plant grasslands. Our results show that the resulting individual-based networks are highly modular, with insect visitors effectively connecting individuals of the same and different plant species. We also obtain empirical evidence that network structure is critical for modulating individual plant reproduction. In particular, the mesoscale level is the best descriptor of plant reproductive success, as it integrates the net effect of local heterospecific and conspecific interactions on seed production of a given individual. We provide a simple, but robust set of metrics to scale down network ecology to functioning properties at the individual level, where most ecological processes take place, hence moving forward the description and interpretation of multitrophic communities across scales.

Multilayer networks | Plant-pollinator networks | Pollination | Ecosystem function | Centrality | Map equation

Correspondence: *alfonso.allen.perkins@gmail.com*

Introduction

Network approaches have been increasingly used in ecology to interpret the complex structure of interactions among species (Bascompte *et al.*, 2003; Martinez, 1991; Thebault & Fontaine, 2010). By understanding ecological communities as networks in which species are linked through pairwise interactions, ecologists have shown that network architecture strongly influences species dynamics (Bascompte & Jordano, 2013; Brose *et al.*, 2006), with quantifiable effects on community properties such as local stability (Allesina & Tang, 2012), robustness to extinctions (Memmott *et al.*, 2004), or the impact of invasive species or climate change on local assemblages (Memmott *et al.*, 2007). However, despite the fast-paced information gain on ecological network architecture, we are still lacking a solid foundation linking network architecture with higher-level properties, such as different functional effects of ecological communities (Magrach *et al.*, 2019; Poisot *et al.*, 2013; Thompson *et al.*, 2012).

To quantify ecological network structure and its links to ecological functioning, most approaches are based on a species-level perspective, yet this view ignores the fact that several ecological processes take place at the individual level and depend on individual variation (Bolnick *et al.*, 2011; Dall *et al.*, 2012; Herrera, 2017; Sih *et al.*, 2012; Wolf & Weissing, 2012). For example, pollination is a key function mediated by insects that transport pollen from one plant individual to another one. Collapsing all plant individuals from a given species into a single average (i.e. node) assumes there is no intraspecific variation in traits such as floral display, plant height, or flower morphology, and ultimately on pollinator visitation rates. This simplification clashes with several empirical examples showing that within species floral dissimilarities can result in different assemblages of pollinators (Arroyo-Correa *et al.*, 2020; Gómez *et al.*, 2020). Pollinators can discern and respond to intraspe-

cific flower variation (Herrera, 2017), influencing the vegetative (e.g., growth rate, carbon assimilation) and reproductive (e.g., fecundity) performance of individuals (Herrera, 2017). However, the study of individual-based networks is in its infancy (Guimarães, 2020) due to theoretical and methodological reasons. Empirically, the sampling effort required to resolve all species interactions into individual ones is labor-intensive, which partly explains the scarcity of empirical studies of individual-based networks, either for one species (Araújo *et al.*, 2008; Gómez & Perfectti, 2012; Gómez *et al.*, 2011; Miguel *et al.*, 2018; Tur *et al.*, 2013) or more species (Dupont *et al.*, 2014, 2010). But a more concerning issue is that we lack robust methods able to depict the complexity of integrating individual nodes belonging to multiple species. As a consequence, most studies on individual-based networks focus on the individuals of one species (usually a plant species) linked by its floral visitors (Gómez & Perfectti, 2012; Gómez *et al.*, 2011) or frugivore species (Miguel *et al.*, 2018), ignoring other individuals of the plant community that belong to other species.

Here, we present a set of tools derived from multilayer networks to understand (i) the general patterns of interactions between conspecific and heterospecific plant individuals belonging to multiple species, and (ii) how that structure relates to individual plant fecundity measured as seed production. Multilayer networks are networks with nodes and links distributed across layers. Each layer represents aspects or features of the nodes or the links that belong to the layer, and the links that connect nodes in different layers (i.e. inter-layer edges) provide information on the processes or interdependencies operating between such layers (see the reviews of Boccaletti *et al.* (2014) and Kivela *et al.* (2014) for further details about the multilayer paradigm and its formal definitions). In ecology, multilayer networks typically depict species interactions through time and space, or across types of interactions (García-Callejas *et al.*, 2017; Hutchinson *et al.*, 2018; Pilosof *et al.*, 2017). Here we take advantage of the flexibility of multilayer formalism to conceptualize individual-based pollination networks as transport-like systems (Aleta *et al.*, 2017; Domenico *et al.*, 2014), and model pollen transfer mediated by floral visitors as an ensemble of conspecific plant-pollinator circuits (or layers) coupled through pollinator species. Specifically, conspecific plant individuals and their floral visitor' species are the nodes of each layer, whereas inter-layer links account for both interspecific movements of insects and phenological overlap of plant species (Fig. 1). This analogy allows us to study in detail the structure of interactions at increasing scales: 1) micro-scale, by analyzing node roles through properties such as their degree centrality or strength (Gómez & Perfectti, 2012; Gómez, 2019); 2) meso-scale, through the study of local interaction patterns, or motifs (Delmas *et al.*, 2018; Milo *et al.*, 2002; Simmons *et al.*, 2018); and 3) macro-scale,

i.e. structural properties of the overall community from a flow-based perspective (Farage *et al.*, 2021; Rosvall *et al.*, 2009). Integrating different scales in a common formalism allows us to explore the relative importance of scale for individual plant reproduction. In particular, it has been recently suggested that meso-scale processes may be particularly important for analyzing the functional consequences of community structure because, at that scale, the local patterns of realised direct interactions can reveal the different mechanisms through which individual- and species-based nodes can indirectly influence each other (Simmons *et al.*, 2018). In order to explore in detail the meso-scale of pollen transport networks, here we use network three-node motifs or triplets (i.e., subgraphs representing patterns of interactions between three nodes) that contain two plant individual nodes. To take into account indirect interactions with conspecific and heterospecific partners, respectively, we introduce the concept of homospecific and heterospecific motifs, that is, motifs in which both plant individuals belong to the same species and motifs with two heterospecific individuals, respectively (see Figs. 1(a), (b) and (e)). This distinction is introduced to highlight the conspecific and heterospecific pollen sources of each plant individual, due to the insect movement and pollination.

We illustrate our approach by analyzing nine well-resolved individual-based plant-pollinator networks from a Mediterranean grassland ecosystem. Our general aim is to assess which network structure explains the reproductive success of plant individuals. We hypothesize that being a central node in the network increases seed production if this position maximizes the conspecific pollen flow. Conversely, it can also entail a decrease in reproductive success if pollen-flows mediated by insects increase conspecific pollen loss and heterospecific pollen deposition on stigmas as it has been previously showed experimentally (Morales & Traveset, 2009; Moreira-Hernández & Muchhala, 2019). Specifically, we aim to: (i) Describe the structure of the system at three community scales: micro-scale, through the use of node-level network properties; meso-scale, through the use of homospecific and heterospecific network motifs; and macro-scale, through the use of multilayer structure and modularity; and (ii) evaluate the relative importance of ecological structure at different scales for the reproductive success of plant individuals.

Material and methods

Data collection. We conducted our observational study at Caracoles Ranch, a 2680 ha salty grassland located within Doñana NP, SW Spain (37°04'01.0" N, 6°19'16.2" W). The climate is Mediterranean with mild winters and average 50-year annual rainfall of 550–570 mm with high interannual oscillations. Soils are sodic saline (electric conductivity > 4 dS/m and pH < 8.5) and annual vegetation dominates the grassland with no

perennial species present (Lanuza *et al.*, 2018). Along a 1 km long × 800 m wide area we established in 2015 nine square plots of side 8.5 m distributed in three subareas: three plots were in the upper part, three in the middle, and three in the lower part. The average distance between these three locations was 300 m and the average distance between plots within each location was 40 m. Each plot was divided into 36 square subplots of side 1 m with aisles of 0.5 m in between to allow access to subplots where measurements were taken (324 subplots in total).

During spring 2020, we sampled the annual plant vegetation and their floral visitors across the nine established plots. Specifically, we observed 23 co-occurring annual plant species (Supplementary material Appendix 1 table A1). We surveyed weekly (weather permitting) the number of floral visits to each plant species within each subplot from February to June, that is, covering all species phenology (see Supplementary material Appendix 2). Visits were only considered when the floral visitor touched the reproductive organs of the plant. All subplots within a plot were simultaneously sampled for 30 min each week by walking slowly between the subplots. Plot survey was randomised between weeks to avoid sampling effects. All floral visitors were either identified during the survey or they were net-collected for their posterior identification at the lab. Overall, this procedure rendered approximately 54 hours of floral visitors sampled over 19 weeks. This amount of sampling has proved enough to quantitatively characterize plant-pollinator interactions (e.g. Albor *et al.* (2019)). Floral visitors were identified to the species (42.34%), morphospecies (41.96%) or family (15.70%) level (Supplementary material Appendix 3 table A3). In addition, we also assigned each species to one of the following eleven functional groups: small beetles, big beetles, butterflies, flies, flower beetles, house flies, hoverflies, humbleflies, small flies, solitary bees, and wasps (Supplementary material Appendix 3 table A3). While some of these floral visitor types (like solitary bees) act as true mutualists, others (such as flower beetles) may rob plant nectar or pollen, and can even damage flowers (Morris *et al.*, 2003). Insects only visited the flowers of 10 of the 23 species present (Supplementary material Appendix 2 Fig. A2.1), and most of these visits were concentrated on three Asteraceae species, namely: *Chamaemelum fuscatum*, *Leontodon maroccanus* and *Pulicaria paludosa*.

To assess plant reproductive success, we sampled at least one individual plant species present per subplot and plot (Supplementary material Appendix 1 table A1), and we counted in the field the number of fruits per individual plant species. We also collected one fruit per individual per subplot and plot from the three most attractive species for insects (*C. fuscatum*, *L. maroccanus* and *P. paludosa*) which represent 95.65% of all insect visits and, thus, are the focus of the reproductive success analysis (see below). We cleaned the collected fruits and counted

the number of viable seeds. Then, the seed production of a given focal individual was estimated as the number of fruits collected for that species multiplied by its number of seeds per fruit. For 49 species-subplot combinations (out of 1,402 combinations, 3.50%) for which we could not obtain seed production information, we used the mean number of seeds per species of their plot.

Multilayer assembly. We built nine multilayer networks (one per plot) to describe (i) the pollen flow between individual plants mediated by insect visitors (i.e. we assume that floral visitors connecting two plant individuals are transferring pollen), and (ii) the effect of flowering phenology on such flow. We define each subplot and species as an individual plant node, for which we can associate their visitation patterns and reproductive success. To model the ensemble of conspecific plants-insects circuits that are coupled through insect species, we chose a node-colored approach, i.e., we consider that there are different types of nodes, defined by "colors", and nodes having similar "colors" are grouped in the same layer (see (Kivela *et al.*, 2014)). Here, "colors" represent the different plant species sampled. If an insect species visits several plant species, we include in the network one insect node per plant species and assign to each insect node the "colors" of the plant visited (see Figs. 1(a) and (e) for an example). Following this approach, the layers of each network (or plot multilayer) only contain nodes that represent (i) conspecific plant individuals (individual-based nodes) and (ii) their floral visitors (species-based nodes). We defined the intralayer links (or intralinks) of the multilayers, that is, the connections within a given layer that represent interactions between the individuals of a given plant species with its insect visitors. Firstly, for each plant-species $\mathcal{P}$, we built a weighted, undirected bipartite graph, denoted here as *visitation* or *interaction network* (VN-$\mathcal{P}$), from the individual plant–insect interaction matrix of each plot (see left-hand side of Fig. 1(c) for an example). Here the weight of links represents the total amount of visits during the sampling period that the focal individual i of species $\mathcal{P}$ recieved from an insect species j, and such amount is given by all the recorded visits of j to $\mathcal{P}$ in i's subplot. Secondly, assuming that, in a given visitation network VN-$\mathcal{P}$, the larger the weight of a connection $w_{ij}^{\mathcal{P}}$ between nodes i and j, the more likely that the interaction will transfer pollen between such individuals, we derived a directed *flow network* from the undirected visitation graph by assigning the following value to the strength of the link from node i to j:

$$f_{ij}^{\mathcal{P}} = \frac{\varepsilon_{ij} w_{ij}^{\mathcal{P}}}{\sum_{s \in V(VN-\mathcal{P})} w_{is}^{\mathcal{P}}} \tag{1}$$

where $f_{ij}^{\mathcal{P}}$ represents the fraction of pollen grains allocated in i that flows to j in the layer $\mathcal{P}$, ε_{ij} is the efficiency of the exchanges from node i to j, V(VN-$\mathcal{P}$) denotes the nodes in the visitation network of $\mathcal{P}$, and $w_{ij}^{\mathcal{P}} = 0$ if i and j are disconnected in VN-$\mathcal{P}$ (see Fig. 1(c)

for an example). Notice that (i) both $f_{ij}^{\mathcal{P}}$ and ε_{ij} take values between 0 and 1, and (ii) $f_{ij}^{\mathcal{P}}$ and ε_{ij} can be different from $f_{ji}^{\mathcal{P}}$ and ε_{ji}, respectively (i.e., flow networks are directed). For the sake of simplicity, hereafter we assume $\varepsilon_{ij} = 1$ for every pair i and j, meaning that all exchanges are equally efficient regardless of the species and individuals involved. We also assume that pollen flow between two plant individuals does not decay with distance, that is, visitor species effectively link plant individuals at the plot level. These two assumptions are justified in our study system because we do not know pollinator efficiency for different species, and we consider that all the insects can move within the plot easily. Nevertheless, this approach can be easily extended to impose efficiencies per guild, if known, and/or pollen flow decay with distance. Here, since $\varepsilon_{ij} = 1$, the interactions $f_{ij}^{\mathcal{P}}$ are the intralayer connections of layer $\mathcal{P}$.

Finally, to couple the different layers and, thus, assemble the multilayer network, we introduced the interlayer links (or interlinks), i.e., interconnections between insects of the same species that are present in several plant layers. We used these interlinks to model how easily the pollen flows from one plant species to another due to the phenological overlap between them and the heterospecific movement of insects. To do so, we proceeded as follows: if a given insect visitor x was present in layers $\mathcal{P}$ and $\mathcal{L}$, we created one interlink from node x in layer $\mathcal{P}$ to x in layer $\mathcal{L}$ with strength $f_x^{\mathcal{P} \to \mathcal{L}}$, and another interlayer connection from $\mathcal{L}$ to $\mathcal{P}$ with strength $f_x^{\mathcal{L} \to \mathcal{P}}$, where $\mathcal{P}$ and $\mathcal{L}$ denote respectively two plant species, and $f_x^{\mathcal{P} \to \mathcal{L}}$ is equal to the phenological overlap of $\mathcal{P}$ and $\mathcal{L}$ divided by the total duration of $\mathcal{P}$'s phenology (see Fig. 1(d) for an example of these calculations). In each plot, we estimated phenologies for each plant species from its weekly record of insect visits (see further details in Supplementary Materials Appendix 2). According to previous definitions, interlayer edges are directed (i.e., $f_x^{\mathcal{P} \to \mathcal{L}}$ and $f_x^{\mathcal{L} \to \mathcal{P}}$ can be different), and their strengths take values between 0 and 1 (see Fig. 1(d) for an example).

Network metrics. Once we created the multilayer arrangements, we assessed how their structures shape the pollen flow that reaches each individual plant. To do so, we estimated a series of network metrics related to network macro- and microstructural patterns, and we complemented it with mesostrucutural descriptors. For the macrostructure, we calculated (i) the modular partitions that better describe network pollen flows. For the microstructure, we obtained (ii) the within-layer and among-layer centralities, which are related with the individual's probability of receiving pollen and the effect of interlinks in such probability, respectively. Below we explain in detail each of these metrics. Finally, we characterized mesostructure through (iii) the 3-node motif characterizations, denoted as homospecific motifs and heterospecific motifs, which incorporates additional de-

tails on the direct and indirect interactions among plant individuals.

Modularity. Network modularity measures how closely connected nodes are divided into modules or compartments (Delmas *et al.*, 2018). Since our multilayers encode pollen flows, we defined modules as non-overlapping groups of nodes that capture flows within and across layers for a relatively long time, rather than clusters with a high internal density of links (i.e. (Domenico *et al.*, 2015)). For that reason, we used Infomap and its random walk approach (R package "infomapecology", (Farage *et al.*, 2021)) to unveil the module organization of our multilayers. Specifically, Infomap uses the map equation to measure the minimum amount of bits (or code length, L) that are needed to describe the movement of a random walker in and between the modules of a given network partition M (Farage *et al.*, 2021; Rosvall *et al.*, 2009). Then, the algorithm finds the partition that requires the least amount of information to describe modular flows.

For each (observed) multilayer network, we estimated the optimal module partition and tested the significance of the total number of modules m and of the code length L by using two null models that preserve the observed phenology (see Appendix 6 in Supplementary materials). Finally, based on the optimal module partitions, in Supplementary materials, we included further calculations of specialization roles, that characterize how the nodes connect the macro-scale modules (see Appendix 5, (Dormann & Strauss, 2013; Guimerà & Amaral, 2005; Olesen *et al.*, 2007; Watts *et al.*, 2016).

Centrality metrics. Centrality is a measure of how "important" a node is in a network. Since such "importance" depends on the structural features or dynamic behaviour we are interested in, there are several ways of being central and, thus, different means of scoring centrality (Delmas *et al.*, 2018; Gómez, 2019). Since our multilayer arrangements are directed and usually contain isolated pairs of plant individuals and insects, we selected traditional centrality metrics that (i) are well-defined in directed and disconnected networks, and (ii) are relevant to characterize the role of nodes from a pollen flow perspective, namely: the in- and out-strength, and the PageRank centrality. In addition, we developed a new metric to further assess the effect of interlinks on the PageRank, the among-layer centrality ratio, denoted as R. This set of metrics include both local measures (i.e., those that quantify the immediate influence between nodes, like the in- and out-strength) and global ones (i.e., those that account for the structure of the entire network, although they are defined at the node level, as the PageRank and the among-layer centrality ratio).

At the local level, the in- and out-strength of a node (i.e., sum of the weights of its in-links and out-links, respectively) are key from a pollen diffusion perspective (mod-

Fig. 1. (a) An example of a visitation network for a 6×6 plot with two plant species and two insect visitors, respectively. For the sake of readability, strengths $w_{ij}^{\mathcal{P}}$ have been omitted. (b) In this panel we highlight examples of homospecific (black solid lines) and heterospecific (black dotted lines) motifs, respectively, in Fig. 1(a). (c) An example of a visitation network and the resulting flow arrangement for plant species 1 (green) in Fig. 1(a). (d) An example of the estimation of interlinks for plant species 1 and 2 (green and orange, respectively) in Fig. 1(a). (e) Resulting multilayer network for the arrangement in Fig. 1(a). Continuous double-arrows represent directed intralayer links, whereas dashed ones show directed interlinks. Shaded areas highlight the multilayer analogs of the motifs in Fig. 1(b). As can be seen, layers are made up of conspecific focal individual plants and their interactions with their insect visitors, and floral visitors connect layers, depending on their movements and on the phenological overlap among plant species. Thus, if a given layer is isolated (i.e., it has no positive interlinks), its plant individuals only receive conspecific pollen mediated by insects; otherwise floral visitors will transport conspecific and heterospecific pollen among layers of plant species.

eled with random-walks): the larger the in-strength of a node (hereafter *within-layer centrality*), the larger its

probability is of receiving pollen grains from its direct neighbors. Instead, the probability that one node trans-

fers its load of pollen to its neighbors is proportional to its out-strength. At the global level, we calculated the PageRank index, whose value represents the probability of finding a grain of pollen (that diffuses on the system) in a given node, after a long time. Consequently, nodes with larger PageRank are likely to mediate and capture larger amounts of pollen flows than those with smaller values. Hereafter we will refer to the PageRank index as the *pollen arrival probability*. Finally, we assessed the effect of interlinks on the pollen arrival probability by estimating the *among-layer centrality ratio*, which is given by $R = MC/LC$, where MC and LC denote the pollen arrival probabilities obtained when the layers are coupled and isolated, respectively. Thus, if R is smaller than 1, the probability of receiving pollen is reduced when plant species coupling is considered, and vice versa. Notice that, in both cases, pollen flows are loaded with heterospecific pollen, depending on the phenologies of co-occurring plants. To calculate the centrality metrics of our multilayer networks we used the igraph package in R (Csardi & Nepusz, 2006).

Motif analysis. To complement the macro- and microdescriptions of the multilayer structures, we incorporate additional details on the local architecture of direct and indirect interactions among nodes (i.e., individual or species) by decomposing the networks into motifs, the basic building blocks of communities (Milo *et al.*, 2002). Motifs are well developed for food webs (Cirtwill *et al.*, 2018), mutualistic (Simmons *et al.*, 2020, 2018), and competitive networks (Godoy *et al.*, 2017), but have been seldom applied to multilayer networks. In our case, we analyzed the number of undirected triplets present in our visitation networks, that is, the pattern of connections of undirected path graphs with length 2 (see examples in Fig. 1(b)). However, even such simple threenode motifs can be differentiated according to the layers involved (see examples in Fig. 1(e)). Considering motifs containing one or two layers is interesting because it perfectly maps onto the effects of indirect conspecific and heterospecific pollen flows on focal individual's per capita seed production. In addition, it moves forward the bipartite motif descriptions from species-based networks, in which all the subgraph nodes are heterospecific (see Simmons *et al.* (2020, 2018)), to individual-based arrangements with conspecific and heterospecific individuals.

To conduct our motif analyses, we introduced a novel motif categorization according to the plant species involved: If all the plant individuals belong to the same species, the motif was referred to as homospecific motif; otherwise, the motif was classified as heterospecific (see Fig. 1(b)). According to previous studies on species-based mutualistic bipartite networks, such motifs can represent indirect competitive (Mitchell *et al.*, 2009; Ye *et al.*, 2013) or facilitative interactions (Carvalheiro *et al.*, 2014; Ghazoul, 2006). For example, in Fig. 1(a) plant individuals may be involved in exploita-

tive competition for finite pollinator resources, or interference competition through interspecific pollen deposition (Flanagan *et al.*, 2010; Mitchell *et al.*, 2009; Ye *et al.*, 2013). Conversely, facilitative effects may occur when the presence of one focal individual increases pollinator visits to a coflowering individual.

To incorporate species' phenological overlap in our motif analysis, we calculated both types of motifs from the bipartite networks that arose weekly in a given plot (see Fig. 1(a) for an example of such networks). Thus, heterospecific motifs only appear when different coflowering species that shared insect visitors are present in a given week. Then, the total amount of homospecific and of heterospecific motifs for a given focal individual and plot were estimated by pooling the weekly number of motifs in which that individual plant was present. Notice that, according to the previous methodology, we assumed that all floral visitors can easily reach all the subplots in a given plot, as we did when assembling the multilayer networks. Finally, we tested the significance of the homospecific and heterospecific motifs per focal individual and plot by using a null model that preserves the experimental phenology (see Appendix 8). A guide to calculating all the presented multilayer metrics is included in the next URL: `https://github.com/RadicalCommEcol/Multi_motifs`

Reproductive success analysis. One key aspect of this multilayer approach is that we can study the impact of direct and indirect interactions between plants and floral visitors at different structural scales on individual plant reproduction success (i.e. seed production). We explore that relationship by fitting generalized mixed models (GMMs) to the individuals of the three species that attracted more insect visitors (*C. fuscatum*, *L. maroccanus*, and *P. paludosa*, an early, middle, and late phenology species, respectively). Since our response variable (individual seed production) represents count data and we detected overdispersion issues and excess of zeroes in our exploratory analysis, we used negative binomial models in our inferences with a log link function and a constant zero-inflated term (see Eq. 2). As explanatory variables, all models included (i) microstructure descriptors, that is, the plant individuals' within-layer centrality (i.e. in-strenght) and among-layer centrality ratio (i.e. the pollen arrival probabilities when the layers are coupled over the pollen arrival probabilities when layers are isolated); and (ii) mesostructure descriptors, i.e. the number of homospecific and heterospecific motifs. We also used plot as a random intercept to account for multiple individuals of the same plant species measured at each plot. Thus, the model for each plant species can be fully specified as follows:

$$Seeds_{ij} \sim 0 \text{ with probabilty } \varphi_{ij} \text{ (Process1),} \qquad \textbf{(2)}$$

$$Seeds_{ij} \sim NB(\mu_{ij}, \alpha) \text{ with probabilty } 1 - \varphi_{ij} \text{ (Process2),}$$

$$E(Seeds_{ij}) = \mu_{ij}(1 - \varphi_{ij}),$$

$$Var\left(Seeds_{ij}\right) = \mu_{ij}\left(1 - \varphi_{ij}\right)\left(1 + \mu_{ij}\left(\varphi_{ij} + 1/\theta\right)\right),$$

$$\log\left(\mu_{ij}\right) = homospecific\ motifs_{ij}$$

$$+\ heterospecific\ motifs_{ij}$$

$$+\ within\text{-}layer\ centrality_{ij}$$

$$+\ among\text{-}layer\ centrality\ ratio_{ij}$$

$$+\ Plot_i,$$

$$Plot_i \sim N(0, \sigma_u),$$

$$\text{logit}\left(\varphi_{ij}\right) = ZI_0,$$

where $Seeds_{ij}$ is the seed production of the jth individual in Plot i for $i = 1, \cdots, 9$; $Plot_i$ is the random intercept, which is assumed to be normally distributed with mean 0 and variance σ_u^2; and represents the overdispersion parameter of the negative binomial. In addition, notice that, for each observation, there are two possible data generation processes and the result of a Bernoulli trial determines which process is used. For observation ij, Process 1 is selected with probability φ_{ij} and Process 2 with probability $1 - \varphi_{ij}$, where φ_{ij} is constant and depends on the constant term ZI_0. Process 1 generates only zero counts, whereas Process 2 generates counts from a negative binomial model.

To keep the regression variables on similar scales, all numeric explanatory variables were centered and scaled during the analysis. Finally, note that our models contain both individuals that received insect visits and individuals that did not. Network metrics for the latter group were estimated by assuming that their corresponding nodes were isolated from each other, as well as from the corresponding multilayers. For that reason, their within-layer centralities and their amount of homospecific and heterospecific motifs were set to zero, respectively, and the among-layer centrality ratio was set to one.

Our analyses were conducted in R, with the package "glmmTMB" (Brooks *et al.*, 2017). To avoid collinearity among explanatory variables, we checked their variance inflation factors with the R-package "performance" (Lüdecke *et al.*, 2020). We also checked model assumptions with the R-package "DHARMa" (Hartig, 2020).

Results

We recorded 1,794 insect visits in our system from February to June 2020. The visits were distributed heterogeneously among the functional groups of floral visitors, plant species, and plots, respectively (Supplementary materials Appendix 4 Fig. A4.1). Flower beetles (395 observations), solitary bees (280), and small flies (135) were the most common functional groups of floral visitors observed, and most interactions involved *L. maroccanus* (1,337 observations) and, to a lesser extent, *C. fuscatum* (268) or *P. paludosa* (111). Small flies and flower beetles were the most abundant insects visiting *C. fuscatum*, while small beetles and flower beetles were the most abundant for *L. maroccanus*, and solitary bees for *P. paludosa*.

Community structure. Plot multilayers obtained from the above field observations contain on average 4.4 ± 2.13 plant species (layers), 71.89 ± 37.34 nodes (of which 49.00 ± 25.29 are plant individuals and 22.89 ± 12.09 insect nodes), 176.89 ± 89.23 intralinks, and 16.00 ± 14.21 interlinks. By using Informap, we found on average 10.78 modules per plot, being (3, 16) the corresponding 95% confidence interval (CI). One third of the modules contains at least two plant species (Fig. 2), Supplementary materials Appendix 4 Fig. A4.3), which shows that insect visitors are effectively linking plant layers. Overall, the modules contain few plant individuals and most of them belong to the same species when the cluster contains several plant species. The average size of modules is 6.67 nodes with CI (2, 36) and the average number of plant individuals per module is 4.55 with CI (1, 26). As expected, the larger the plot richness and/or the longer phenologies, the larger the number of modules. Our results also suggest that modules reflect phenological constraints. For instance, *C. fuscatum* plant individuals in plot 2 do not share modules with those of *P. paludosa* (Fig. 2) because their respective phenologies do not overlap (Supplementary material Appendix 2 Fig. A2.1).

Results for insect species, which represent on average one third of the nodes per module (2.12 insect nodes per module with CI (1, 6)), are very similar to those of plant individuals (Supplementary materials Appendix 4 Fig. A4.3). Additional details on module features can be found in Supplementary materials (see Appendix 4). To assess the significance of the modules obtained, we compared the observed values for map equation ($L_{Observed}$) and the number of modules ($m_{Observed}$) in each plot with the null distribution obtained from our null models (see details in Appendix 6). According to our results, the optimal partitions of our multilayers provide a significantly better description of the pollen flows within the whole system (or exhibit more modular regularities, i.e., $L_{Observed}$ is smaller) than those estimated for random graphs, although the number of modules is not significantly different from that of null models in most plots (see Appendix 6). In other words, while the number of modules obtained is consistent with a null expectation, their composition is significantly different from it.

Centralities. Overall, within-layer centrality (i.e., in-strength) of plant individuals correlates with the total amount of visits that a focal individual received (Spearman's $\rho = 0.34$). Furthermore, we also found a generally strong, positive, and significant correlation between the within-layer centrality values and pollen arrival probabilities (Supplementary materials Appendix 7 Fig. A7.1). This is expected because the asymptotic probability of finding a grain of pollen in a given node depends on the probability of receiving such grain from the direct neighbors (which is mediated by the within-layer centrality of the node)

Pollen arrival probability is also positively correlated

Fig. 2. Number of plant individuals per module, plant species and plot.

Fig. 3. Dependence of plant individuals' pollen arrival probability (or PageRank index) on plot and species. Mean values are shown with red diamonds. Points show results for individual plants. Dashed lines show the theoretical pollen arrival probability if all the nodes had the same probability.

with the total amount of visits that a focal individual received (Spearman's $\rho = 0.47$). In Fig. 3 we show pollen arrival probability results per plant species and plot. Remarkably, the variance within a given plant species is larger than the one among species. In addition, we can see that, as expected, the larger the total number of plant individuals per plot, the smaller the average pollen arrival probability (Spearman's $\rho = -0.98$) and the theoretical pollen arrival probability, which is calculated as if all the nodes would have the same arrival probability (dashed lines in Fig. 3).

Finally, we see a moderate positive correlation between the among-layer centrality ratio (R) and the total amount of visits (Spearman's $\rho = 0.26$). According to our results, considering interlinks in our communities induces, on average, a loss of centrality in most plant individuals (i.e., $R < 1$). That is, on average most plant individuals receive much less conspecific pollen mediated by insect visitors, when the multilayer is coupled. Only *L. maroccanus* and *S. laciniata* plant individuals gain centrality (on average) when the plant layers are connected (Supplementary materials Appendix 7 Fig. A7.2).

Fig. 4. Dependence of the amount of homospecific and heterospecific motifs per focal individual plant species. Points show results for individual plants. Dashed lines locate those plant individuals whose numbers of homospecific and heterospecific motifs are equal.

Indeed, differences among plant species are significant in each plot (Kruskal-Wallis rank test: $p < 0.05$).

Homospecific and heterospecific motifs. In our communities, the number of homospecific motifs is highly correlated with the total number of visits (Spearman's $\rho = 0.73$), whereas the number of heterospecific motifs is not (Spearman's $\rho = 0.11$). Homospecific motifs are more abundant than heterospecific ones in *C. fuscatum* and *L. maroccanus* individuals, whereas both metrics are comparable in most focal individuals of *P. paludosa*, and, in the case of other plant species, heterospecific motifs predominate over homospecific motifs (Fig. 4). As expected, those plant species with the largest amount of visits and the smallest diversity of floral visitors (such as *C. fuscatum* and *L. maroccanus*) accumulate the largest amount of homospecific motifs in each plot (Supplementary materials Appendix 4 Fig. A4.1). That is, *C. fuscatum* and *L. maroccanus* individuals tend to share more weekly visits of the same insects within their focal individuals than visits with heterospecific individuals. Indeed, we found generally significant differences in the average number of homospecific motifs among plant species in each plot (Kruskal-Wallis rank test: $p < 0.05$). Regarding heterospecific motifs, both extremely common (i.e. *L. maroccanus*) or rare plant species show the largest amounts of such motifs. This is due to the fact that the latter species share all insect visitors with the former, along the sampling weeks (Supplementary material Appendix 2 Fig. A2.2).

Reproductive success analysis. Results for the GLMMs of the three most attractive species for the insect visitors (*C. fuscatum*, *L. maroccanus* and *P. paludosa*) show that, overall, our mesostructure descriptors (i.e., the total amount of homospecific and heterospecific motifs) exert a stronger influence on the seed production of the plant individuals considered than the microstructure metrics (i.e., the within-layer centrality and pollen

competition ratio) (see table 1 and Fig. 5).

Table 1. Results for the GLMMs of *C. fuscatum* (a), *L. maroccanus* (b), *P. paludosa* (c) showing the effect on plant reproductive success (seeds per individual plant). The term ZI_0 describes the zero inflation probability φ_{ij}, given by $\mathrm{logit}(\varphi_{ij}) = ZI_0$ (see Eq. 2 in section "Reproductive success analysis"). Important estimates (i.e. those that are significant at 10%) are highlighted in bold.

(a) *C. fuscatum*: the model contains 160 plant individuals. Conditional $R^2 = 0.319$, marginal $R^2 = 0.052$, with $\sigma_u^2 = 0.4821$ and $\theta = 1.09$.

Seeds per individual plant	Estimate	Std. Error	z value
(Intercept)	**5.0489**	**0.2653**	**19.03**
Homospecific motifs	**0.2847**	**0.1398**	**2.04**
Heterospecific motifs	**-0.1432**	**0.0822**	**-1.74**
Within-layer centrality	0.1379	0.0946	1.46
Among-layer centrality ratio	**0.0268**	**0.1049**	**0.26**
ZI₀	**-3.7841**	**0.6388**	**-5.92**

(b) *L. maroccanus*: the model contains 293 plant individuals. Conditional $R^2 = 0.304$, marginal $R^2 = 0.065$, with $\sigma_u^2 = 0.2503$ and $\theta = 2.37$.

Seeds per individual plant	Estimate	Std. Error	z value
(Intercept)	**5.7114**	**0.1717**	**33.26**
Homospecific motifs	**0.2323**	**0.0760**	**3.06**
Heterospecific motifs	-0.0455	0.0433	-1.05
Within-layer centrality	-0.0115	0.0424	-0.27
Among-layer centrality ratio	**0.0913**	**0.0550**	**1.66**
ZI₀	**-20.99**	**2111.82**	**-0.01**

(c) *P. paludosa*: the model contains 146 plant individuals. Conditional $R^2 = 0.190$, marginal $R^2 = 0.068$, with $\sigma_u^2 = 0.1373$ and $\theta = 0.834$.

Seeds per individual plant	Estimate	Std. Error	z value
(Intercept)	**5.6456**	**0.1666**	**33.89**
Homospecific motifs	**-0.1977**	**0.1804**	**-1.10**
Heterospecific motifs	**0.3118**	**0.1798**	**1.73**
Within-layer centrality	-0.0309	0.1101	-0.28
Among-layer centrality ratio	**-0.1662**	**0.1254**	**-1.33**
ZI$_0$	**-2.4673**	**0.3263**	**-7.56**

Results for *C. fuscatum* and *L. maroccanus* suggest that the larger the amount of homospecific (heterospecific) motifs, the larger the increase (decrease) in seed production. Besides, for both plant species gaining centrality leads to an increase in seed production. However *C. fuscatum* individuals with higher seed production are those showing larger values of within-layer centrality, regardless of the coupling between the layers, whereas *L. maroccanus* individuals mainly benefit from being central when their among-layer centrality ratio grows (that is, if their PageRank increases when the corresponding multilayers are coupled). These different ways of benefiting from centrality may reflect their different phenological overlaps: *L. maroccanus*' phenology overlaps completely with those of the remaining plant species, except *C. fuscatum*, whose overlap is only partial (Supplementary material Appendix 2 Fig. A2.1). Indeed, *C. fuscatum* individuals barely share one of their main visitors (small flies) with other plant species (Supplementary material Appendix 2 Fig. A2.2).

Regarding *P. paludosa* individuals, the results of our meso-descriptors contrast with those for *C. fuscatum* and *L. maroccanus*. For *P. paludosa*, the larger the amount of homospecific (heterospecific) motifs, the larger the decrease (increase) in seed production. Furthermore, being central when the multilayers are coupled (among-layer-centrality ratio) also impacts negatively on *P. paludosa*'s seed production. These opposed results compared to the other two Asteraceae species can reflect that, unlike *C. fuscatum* individuals, *P. paludosa* individuals share all their main visitors (i.e., solitary bees and house flies) with *L. maroccanus* (Supplementary material Appendix 2 Fig. A2.2).

Discussion

Our approach shows that multilayer representations of individual-based insect-plant visitation networks allow a deep description of the pollen flow processes and their connections with plant reproductive success. To achieve this, we modeled the connections between plant individuals mediated by floral visitors in each community as a transport-like system, in which there is an ensemble of conspecific plants-insects circuits (or layers) that are coupled through insect species that transfer pollen. Here layers contain conspecific focal individual plants and their interactions (intralinks) with their insect visitors, and those visitors connect layers (through the interlinks), depending on their movements and on the phenological overlap among plant species. Below, we first discuss implications for describing community structure, second, we highlight how to link these metrics to other processes such as pollination, and finally, we describe potential caveats and improvements.

Describing community-level individual-based networks. In species-rich plant communities, depicting individuals of each species in their own layer and layers linked via shared pollinators can rapidly help to visualize the within-species and among-species dynamics. For example, our nine individual plant-pollinator networks contain few layers that are coupled by a small number of directed interlinks, suggesting that plant species have dissimilar phenologies and share few species of floral visitors. In contrast, more clustered structures, where interlinks are abundant, may describe potential pollinator competition among plant species (Pauw, 2013). However, even the small number of interlinks observed in our study system are key determinants of the network structure from a flow perspective. Firstly, this is because, at the macrostructural level, interlinks shape the modular organization of the different individuals in clusters. The identity of the individuals of these modules is likely determined by the pollen flow dynamics. In our case, despite modules being dominated by single species, we observe modules that contain up to four different plant species. Indeed, those insect species that channel interlinks turn out to be network and module hubs of our arrangements. Second, at the micro-level, interlinks alter the plant individuals' probability of receiving pollen flow, with ramifications for the heterospecific pollen movement between plant individuals (Lanuza *et al.*, 2021). In general, pollen arrival probability of most plant individuals in our study drops when multilayers are coupled with the exception of several *L. maroccanus* individuals. Finally, the meso-scale descriptors also reflect the number of interlinks observed, such that in our case, and as expected, a paucity of interlinks translates into the overabundance of homospecific motifs for those plant species with the largest amount of visits and smallest diversity of floral visitors (such as *C. fuscatum* and *L. maroccanus*).

Linking network structure to network functioning. Beyond the description of network patterns, our individual-based approach directly maps into ecological processes and functions such as pollination. In fact, we found that the best descriptors of plant reproductive success are found at the meso-scale, since those descriptors integrate all the conspecific and heterospecific interactions of a given individual. This result agrees with recent

Fig. 5. Conditional plots for plant explanatory variables (R-package "visreg" (Breheny & Burchett, 2017)): Homospecific motifs (first row), heterospecific motifs (second row), within-layer centrality (third row), and among-layer centrality ratio (fourth row). Notice that, for the conditioning, the other variables are set to their median.

suggestions that the meso-scale captures important information from the local topology of the network, without adding too much complexity (Simmons *et al.*, 2018). In our models for *C. fuscatum*, *L. maroccanus* and *P. paludosa*, the total amount of homospecific motifs, depicting the amount and origin of floral visitors, showed significant relations with the seed production of all those species. We observed that homospecific motifs exerted a positive effect on the reproductive success of *C. fusca-tum* and *L. maroccanus* individuals, which is expected

if homospecific motifs quantify the number of conspe-cific visits. In addition, the number of heterospecific motifs showed the opposite trend, indicating that vis-its coming from heterospecifics do not contribute, and can even harm reproductive success (Ashman & Schoen, 1996). Surprisingly, we found the opposite trend for *P. paludosa*. This species is the less abundant of the three Asteraceae species studied in detail and while the main visitor of the other two species are small flies and beetles, its main visitors are highly mobile solitary bees (*Andrena*

and *Lassioglossum species*) which act as network hubs, and which are also strongly coupled with *L. maroccanus*. Indeed, our field observations show that *P. paludosa* may act as a magnet species (Lopezaraiza–Mikel *et al.*, 2007) because bees visited *L. maroccanus* mainly in plots with the presence of *P. paludosa*. Bees are efficient pollinators that can carry different pollen species on different body parts due to differences in flower morphology and stigma position (Armbruster *et al.*, 1994). This species-specific pollen placement may contribute to preventing high levels of heterospecific pollen deposition. Hence, the relationship between heterospecific motifs and reproductive success may depend on pollinator foraging behaviour. Harder to explain is the negative relationship of homospecific motifs with reproductive success. Bearing in mind that this relationship is comparatively weak (Fig. 5) and thus we cannot rule out the possibility of a spurious trend, a possibility is that the low number of *P. paludosa* homospecific motifs are strongly coupled with other plant species. So its homospecific motifs do not form closed loops, but are part of longer chains that basically distribute pollen flows to non-target individuals as suggested by the fact that 52% of the 25 modules with *P. paludosa* individuals also contain focal individuals of *L. maroccanus*, and only 20% of those modules are dominated by *P. paludosa* (i.e., the percentage of plant individuals of *P. paludosa* is larger than 50%), whereas 83.33% of the 18 modules with *C. fuscatum* are dominated by *C. fuscatum* focals, and, similarly, 84.37% of the 64 modules with *L. maroccanus* are dominated by the latter.

Despite the weaker effects of individual plants within-layer centrality, we also observed a general positive effect similar to the results found by Gómez & Perfectti (2012). In addition, among-layer centrality also seems to be important for *L. maroccanus*. This finding suggests that those *L. maroccanus* individuals that gain centrality when the multilayer is coupled (i.e., those that attract more insects) increase their chance of intercrossing with a higher number of conspecifics via longer loops or pollen carryover, which occurs when pollen collected from a plant is dispersed not only to the next plant but also to subsequent plants visited by a pollinator (Karron *et al.*, 1995). However, following the reasoning in Gómez & Perfectti (2012), since the main visitors of *L. maroccanus* are small beetles and flower beetles that usually feed on pollen and spend long periods of time on each flower they visit, the positive trend we found is apparently counter-intuitive. Despite *Brassicogethes sp.* and *Psilothrix viridicoerulea*, the most abundant floral visitor species, are described as poor pollinators (Bartomeus *et al.*, 2008), adults in flowers have pollen on their bodies and can disperse pollen at both within-field and landscape scales (Ramsay *et al.*, 2003), and its high abundance may be contributing to the relative importance we observed, as predicted by the mass ratio hypothesis (Grime, 1998). Besides, our model did

not account for the positive effect that selfing induced by insect movement may be exerting in *L. maroccanus*. Measuring pollinator efficiency (King *et al.*, 2013) may be a further step needed to understand the full picture.

Caveats and future directions. Collecting data to model individual-based networks at the community level is a daunting task, but as more examples emerge, we will be able to document generalities and validate untested assumptions of our framework. The most critical assumption is that individuals linked by the same pollinator species have larger probabilities of mating (Arroyo-Correa *et al.*, 2020; Gómez *et al.*, 2020). While reasonable at the scales investigated, pollinator behaviour (Devaux *et al.*, 2014) and morphology are expected to further modulate this relationship. Following individual pollinators to test for floral constancy (e.g. Jakobsson *et al.* (2008)) or analyzing pollen loads on pollinators bodies (Bosch *et al.*, 2009) deserves future attention to test this assumption. Once collected this information, it can be readily incorporated in our framework as it allows including assessments about different pollinator efficiencies or a distance decay function between nodes. In our framework, we have used only undirected motifs with three nodes to test our pollen flow approach. However, as pointed out by Simmons *et al.* (2018), it may be important to consider larger motifs in order to adequately capture meso-scale structure. Nevertheless, although more complex structures can be computed, the challenge remains in linking specific motifs to a clear ecological process. Thus, future work extending the downscaling of motifs to individual-level by considering quadruplets and other motifs of conspecific and heterospecific individuals with more than three nodes can be relevant to gain further insight into the functional consequences of community structure. Besides, since we have focused on binary interactions to perform our motif calculations (that is, to define a motif we were only interested in the presence or absence of interactions and not in the interaction strengths), another future research direction is to include quantitative information on interaction weights. Incorporating that extra layer of information may be relevant to identify how indirect interactions affect final ecosystem functions. However, despite being a promising direction for future investigation, quantitative bipartite motif analysis is not yet fully developed (see examples in Simmons *et al.* (2018) and further work is needed to conceptualize it and interpret their results (Simmons *et al.*, 2020).

Finally, It is important to note that all empirical networks may be biased by incomplete sampling, for instance, failing to detect the interactions of rare species (Blüthgen, 2010; Blüthgen *et al.*, 2008). This uncertainty affects the community detection presented in this work, as well as the network metrics we discussed so far. For instance, the map equation assumes complete data and, when networks are sparse and may contain missing links, the optimal solution is distorted (Smil-

janić *et al.*, 2020). To overcome these obstacles, further investigation should consider the use of improved flow-based community detection algorithms in development (Smiljanić *et al.*, 2020; Young *et al.*, 2020, 2019).

Overall, we provide a solid set of flexible tools to further explore individual-based networks. Downscaling interaction network ecology to the individual level may be important to map functions and process occurring as such scale, and to better capture intraspecific variability. Further research on individual-based community-wide networks would enable us to link network theory to processes determining the emergent properties of interactions at multiple scales.

ACKNOWLEDGEMENTS

The authors thank Doñana National Park for granting access to their facilities (permit 2019/14). We also acknowledge Francisco P. Molina (Seville, Spain) and Álvaro Pérez-Gómez (Cádiz, Spain) for helping with insect classification, and Pedro Jordano Lab for fruitful discussion. A.A.-P. and I.B. were funded through the 2017-2018 Belmont Forum and BiodivERsA joint call for research proposals, under the BiodivScen ERA-Net COFUND programme, and with the funding organisations AEI, NWO, ECCyT and NSF. O.G. acknowledges financial support provided by the Spanish Ministry of Economy and Competitiveness (MINECO) and by the European Social Fund through the Ramón y Cajal Program (RYC-2017- 23666). This study was partly funded by the Spanish Ministry of Science and Innovation (MICINN) and the European Social fund through the MeDiNaS project (Mechanisms of maintenance of species diversity in variable environments, RTI2018-098888-A-I00). M.H. acknowledges financial support provided by Spanish Ministry of Science, Innovation and Universities (Grant No. PRE2019-088280).

References

Albor, C., García-Franco, J. G., Parra-Tabla, V., Díaz-Castelazo, C. & Arceo-Gómez, G. (2019). Taxonomic and functional diversity of the co-flowering community differentially affect cakile edentula pollination at different spatial scales. *Journal of Ecology*, 107, 2167–2181.

Aleta, A., Meloni, S. & Moreno, Y. (2017). A multilayer perspective for the analysis of urban transportation systems. *Scientific Reports*, 7.

Allesina, S. & Tang, S. (2012). Stability criteria for complex ecosystems. *Nature*, 483, 205–208.

Araújo, M. S., es, P. R. G., Svanbäck, R., Pinheiro, A., Guimaraẽs, P., dos Reis, S. F. & Bolnick, D. I. (2008). Network analysis reveals contrasting effects of intraspecific competition on individual vs. population diets. *Ecology*, 89, 1981–1993.

Armbruster, W. S., Edwards, M. E. & Debevec, E. M. (1994). Floral character displacement generates assemblage structure of western australian triggerplants (stylidium). *Ecology*, 75, 315–329.

Arroyo-Correa, B., Bartomeus, I. & Jordano, P. (2020). Phenotypic and microsite plant individual variation determine the pollination network structure and its functional consequences.

Ashman, T.-L. & Schoen, D. J. (1996). Floral longevity: Fitness consequences and resource costs. In: *Floral Biology*. Springer US, pp. 112–139. URL `https://doi.org/10.1007/978-1-4613-1165-2_5`.

Bartomeus, I., Bosch, J. & Vilà, M. (2008). High invasive pollen transfer, yet low deposition on native stigmas in a carpobrotus-invaded community. *Annals of Botany*, 102, 417–424.

Bascompte, J. & Jordano, P. (2013). *Mutualistic Networks*. Princeton University Press. URL `https://doi.org/10.1515/9781400848720`.

Bascompte, J., Jordano, P., Melian, C. J. & Olesen, J. M. (2003). The nested assembly of plant-animal mutualistic networks. *Proceedings of the National Academy of Sciences*, 100, 9383–9387.

Blüthgen, N. (2010). Why network analysis is often disconnected from community ecology: A critique and an ecologist's guide. *Basic and Applied Ecology*, 11, 185–195.

Blüthgen, N., Fründ, J., Vázquez, D. P. & Menzel, F. (2008). WHAT DO INTERACTION NETWORK METRICS TELL US ABOUT SPECIALIZATION AND BIOLOGICAL TRAITS. *Ecology*, 89, 3387–3399.

Boccaletti, S., Bianconi, G., Criado, R., del Genio, C., Gómez-Gardeñes, J., Romance, M., Sendiña-Nadal, I., Wang, Z. & Zanin, M. (2014). The structure and dynamics of multilayer networks. *Physics Reports*, 544, 1–122.

Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., Rudolf, V. H., Schreiber, S. J., Urban, M. C. & Vasseur, D. A. (2011). Why intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution*, 26, 183–192.

Bosch, J., González, A. M. M., Rodrigo, A. & Navarro, D. (2009). Plant-pollinator networks: adding the pollinator's perspective. *Ecology Letters*, 12, 409–419.

Breheny, P. & Burchett, W. (2017). Visualization of regression models using visreg. *The R Journal*, 9, 56–71.

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Maechler, M. & Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal*, 9, 378–400.

Brose, U., Williams, R. J. & Martinez, N. D. (2006). Allometric scaling enhances stability in complex food webs. *Ecology Letters*, 9, 1228–1236.

Carvalheiro, L. G., Biesmeijer, J. C., Benadi, G., Fründ, J., Stang, M., Bartomeus, I., Kaiser-Bunbury, C. N., Baude, M., Gomes, S. I. F., Merckx, V., Baldock, K. C. R., Bennett, A. T. D., Boada, R., Bommarco, R., Cartar, R., Chacoff, N., Dänhardt, J., Dicks, L. V., Dormann, C. F., Ekroos, J., Henson, K. S., Holzschuh,

A., Junker, R. R., Lopezaraiza-Mikel, M., Memmott, J., Montero-Castaño, A., Nelson, I. L., Petanidou, T., Power, E. F., Rundlöf, M., Smith, H. G., Stout, J. C., Temitope, K., Tscharntke, T., Tscheulin, T., Vilà, M. & Kunin, W. E. (2014). The potential for indirect effects between co-flowering plants via shared pollinators depends on resource abundance, accessibility and relatedness. *Ecology Letters*, 17, 1389–1399.

Cirtwill, A. R., Riva, G. V. D., Gaiarsa, M. P., Bimler, M. D., Cagua, E. F., Coux, C. & Dehling, D. M. (2018). A review of species role concepts in food webs. *Food Webs*, 16, e00093.

Csardi, G. & Nepusz, T. (2006). The igraph software package for complex network research. *InterJournal*, Complex Systems, 1695.

Dall, S. R. X., Bell, A. M., Bolnick, D. I. & Ratnieks, F. L. W. (2012). An evolutionary ecology of individual differences. *Ecology Letters*, 15, 1189–1198.

Delmas, E., Besson, M., Brice, M.-H., Burkle, L. A., Riva, G. V. D., Fortin, M.-J., Gravel, D., Guimarães, P. R., Hembry, D. H., Newman, E. A., Olesen, J. M., Pires, M. M., Yeakel, J. D. & Poisot, T. (2018). Analysing ecological networks of species interactions. *Biological Reviews*, 94, 16–36.

Devaux, C., Lepers, C. & Porcher, E. (2014). Constraints imposed by pollinator behaviour on the ecology and evolution of plant mating systems. *Journal of Evolutionary Biology*, 27, 1413–1430.

Domenico, M. D., Lancichinetti, A., Arenas, A. & Rosvall, M. (2015). Identifying modular flows on multilayer networks reveals highly overlapping organization in interconnected systems. *Physical Review X*, 5.

Domenico, M. D., Sole-Ribalta, A., Gomez, S. & Arenas, A. (2014). Navigability of interconnected networks under random failures. *Proceedings of the National Academy of Sciences*, 111, 8351–8356.

Dormann, C. F. & Strauss, R. (2013). A method for detecting modules in quantitative bipartite networks. *Methods in Ecology and Evolution*, 5, 90–98.

Dupont, Y. L., Trøjelsgaard, K., Hagen, M., Henriksen, M. V., Olesen, J. M., Pedersen, N. M. E. & Kissling, W. D. (2014). Spatial structure of an individual-based plant-pollinator network. *Oikos*, 123, 1301–1310.

Dupont, Y. L., Trøjelsgaard, K. & Olesen, J. M. (2010). Scaling down from species to individuals: a flower-visitation network between individual honeybees and thistle plants. *Oikos*, 120, 170–177.

Farage, C., Edler, D., Eklöf, A., Rosvall, M. & Pilosof, S. (2021). Identifying flow modules in ecological networks using infomap. *Methods in Ecology and Evolution*.

Flanagan, R. J., Mitchell, R. J. & Karron, J. D. (2010). Increased relative abundance of an invasive competitor for pollination, lythrum salicaria, reduces seed number in mimulus ringens. *Oecologia*, 164, 445–454.

García-Callejas, D., Molowny-Horas, R. & Araújo, M. B. (2017). Multiple interactions networks: towards more realistic descriptions of the web of life. *Oikos*, 127, 5–22.

Ghazoul, J. (2006). Floral diversity and the facilitation of pollination. *Journal of Ecology*, 94, 295–304.

Godoy, O., Stouffer, D. B., Kraft, N. J. B. & Levine, J. M. (2017). Intransitivity is infrequent and fails to promote annual plant coexistence without pairwise niche differences. *Ecology*, 98, 1193–1200.

Gómez, J. M. & Perfectti, F. (2012). Fitness consequences of centrality in mutualistic individual-based networks. *Proceedings of the Royal Society B: Biological Sciences*, 279, 1754–1760.

Gómez, J. M., Perfectti, F., Armas, C., Narbona, E., González-Megías, A., Navarro, L., DeSoto, L. & Torices, R. (2020). Within-individual phenotypic plasticity in flowers fosters pollination niche shift. *Nature Communications*, 11.

Gómez, J. M., Perfectti, F. & Jordano, P. (2011). The functional consequences of mutualistic network architecture. *PLoS ONE*, 6, e16143.

Gómez, S. (2019). Centrality in networks: Finding the most important nodes. In: *Business and Consumer Analytics: New Ideas*. Springer International Publishing, pp. 401–433. URL `https://doi.org/10.1007/978-3-030-06222-4_8`.

Grime, J. P. (1998). Benefits of plant diversity to ecosystems: immediate, filter and founder effects. *Journal of Ecology*, 86, 902–910.

Guimarães, P. R. (2020). The structure of ecological networks across levels of organization. *Annual Review of Ecology, Evolution, and Systematics*, 51, 433–460.

Guimerà, R. & Amaral, L. A. N. (2005). Functional cartography of complex metabolic networks. *Nature*, 433, 895–900.

Hartig, F. (2020). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models*. URL `https://CRAN.R-project.org/package=DHARMa`. R package version 0.3.0.

Herrera, C. M. (2017). The ecology of subindividual variability in plants: patterns, processes, and prospects. *Web Ecology*, 17, 51–64.

Hutchinson, M. C., Mora, B. B., Pilosof, S., Barner, A. K., Kéfi, S., Thébault, E., Jordano, P. & Stouffer, D. B. (2018). Seeing the forest for the trees: Putting multilayer networks to work for community ecology. *Functional Ecology*, 33, 206–217.

Jakobsson, A., Padrón, B. & Traveset, A. (2008). Pollen transfer from invasive carpobrotus spp. to natives – a study of pollinator behaviour and reproduction success. *Biological Conservation*, 141, 136–145.

Karron, J. D., Tucker, R., Thumser, N. N. & Reinartz, J. A. (1995). Comparison of pollinator flight movements and gene dispersal patterns in mimulus ringens. *Heredity*, 75, 612–617.

King, C., Ballantyne, G. & Willmer, P. G. (2013). Why flower visitation is a poor proxy for pollination: measuring single-visit pollen deposition, with implications

for pollination networks and conservation. *Methods in Ecology and Evolution*, 4, 811–818.

Kivela, M., Arenas, A., Barthelemy, M., Gleeson, J. P., Moreno, Y. & Porter, M. A. (2014). Multilayer networks. *Journal of Complex Networks*, 2, 203–271.

Lanuza, J. B., Bartomeus, I., Ashman, T.-L., Bible, G. & Rader, R. (2021). Recipient and donor characteristics govern the hierarchical structure of heterospecific pollen competition networks. *Journal of Ecology.*

Lanuza, J. B., Bartomeus, I. & Godoy, O. (2018). Opposing effects of floral visitors and soil conditions on the determinants of competitive outcomes maintain species diversity in heterogeneous landscapes. *Ecology Letters*, 21, 865–874.

Lopezaraiza–Mikel, M. E., Hayes, R. B., Whalley, M. R. & Memmott, J. (2007). The impact of an alien plant on a native plant–pollinator network: an experimental approach. *Ecology Letters*, 10, 539–550.

Lüdecke, D., Makowski, D., Waggoner, P. & Patil, I. (2020). performance: Assessment of regression models performance. *CRAN.*

Magrach, A., Molina, F. P. & Bartomeus, I. (2019). Niche complementarity among pollinators increases community-level plant reproductive success.

Martinez, N. D. (1991). Artifacts or attributes? effects of resolution on the little rock lake food web. *Ecological Monographs*, 61, 367–392.

Memmott, J., Craze, P. G., Waser, N. M. & Price, M. V. (2007). Global warming and the disruption of plant?pollinator interactions. *Ecology Letters*, 10, 710–717.

Memmott, J., Waser, N. M. & Price, M. V. (2004). Tolerance of pollination networks to species extinctions. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 271, 2605–2611.

Miguel, M. F., Jordano, P., Tabeni, S. & Campos, C. M. (2018). Context-dependency and anthropogenic effects on individual plant-frugivore networks. *Oikos*, 127, 1045–1059.

Milo, R., Shen-Orr, S., Itzkovitz, S., Kashtan, N., Chklovskii, D. & Alon, U. (2002). Network motifs: Simple building blocks of complex networks. *Science*, 298, 824–827.

Mitchell, R. J., Flanagan, R. J., Brown, B. J., Waser, N. M. & Karron, J. D. (2009). New frontiers in competition for pollination. *Annals of Botany*, 103, 1403–1413.

Morales, C. L. & Traveset, A. (2009). A meta-analysis of impacts of alien vs. native plants on pollinator visitation and reproductive success of co-flowering native plants. *Ecology Letters*, 12, 716–728.

Moreira-Hernández, J. I. & Muchhala, N. (2019). Importance of pollinator-mediated interspecific pollen transfer for angiosperm evolution. *Annual Review of Ecology, Evolution, and Systematics*, 50, 191–217.

Morris, W. F., Bronstein, J. L. & Wilson, W. G. (2003). Three-way coexistence in obligate mutualist-exploiter interactions: The potential role of competition. *The American Naturalist*, 161, 860–875.

Olesen, J. M., Bascompte, J., Dupont, Y. L. & Jordano, P. (2007). The modularity of pollination networks. *Proceedings of the National Academy of Sciences*, 104, 19891–19896.

Pauw, A. (2013). Can pollination niches facilitate plant coexistence? *Trends in Ecology & Evolution*, 28, 30–37.

Pilosof, S., Porter, M. A., Pascual, M. & Kéfi, S. (2017). The multilayer nature of ecological networks. *Nature Ecology & Evolution*, 1.

Poisot, T., Mouquet, N. & Gravel, D. (2013). Trophic complementarity drives the biodiversity-ecosystem functioning relationship in food webs. *Ecology Letters*, 16, 853–861.

Ramsay, G., Thompson, C. & Squire, G. (2003). Quantifying landscape-scale gene flow in oilseed rape. department for environment, food and rural affairs.

Rosvall, M., Axelsson, D. & Bergstrom, C. T. (2009). The map equation. *The European Physical Journal Special Topics*, 178, 13–23.

Sih, A., Cote, J., Evans, M., Fogarty, S. & Pruitt, J. (2012). Ecological implications of behavioural syndromes. *Ecology Letters*, 15, 278–289.

Simmons, B. I., Beckerman, A. P., Hansen, K., Maruyama, P. K., Televantos, C., Vizentin-Bugoni, J. & Dalsgaard, B. (2020). Niche and neutral processes leave distinct structural imprints on indirect interactions in mutualistic networks. *Functional Ecology*, 35, 753–763.

Simmons, B. I., Cirtwill, A. R., Baker, N. J., Wauchope, H. S., Dicks, L. V., Stouffer, D. B. & Sutherland, W. J. (2018). Motifs in bipartite ecological networks: uncovering indirect interactions. *Oikos*, 128, 154–170.

Smiljanić, J., Edler, D. & Rosvall, M. (2020). Mapping flows on sparse networks with missing links. *Physical Review E*, 102.

Thebault, E. & Fontaine, C. (2010). Stability of ecological communities and the architecture of mutualistic and trophic networks. *Science*, 329, 853–856.

Thompson, R. M., Brose, U., Dunne, J. A., Hall, R. O., Hladyz, S., Kitching, R. L., Martinez, N. D., Rantala, H., Romanuk, T. N., Stouffer, D. B. & Tylianakis, J. M. (2012). Food webs: reconciling the structure and function of biodiversity. *Trends in Ecology & Evolution*, 27, 689–697.

Tur, C., Vigalondo, B., Trøjelsgaard, K., Olesen, J. M. & Traveset, A. (2013). Downscaling pollen-transport networks to the level of individuals. *Journal of Animal Ecology*, 83, 306–317.

Watts, S., Dormann, C. F., González, A. M. M. & Ollerton, J. (2016). The influence of floral traits on specialization and modularity of plant–pollinator networks in a biodiversity hotspot in the peruvian andes. *Annals of Botany*, 118, 415–429.

Wolf, M. & Weissing, F. J. (2012). Animal personali-

ties: consequences for ecology and evolution. *Trends in Ecology & Evolution*, 27, 452–461.

Ye, Z.-M., Dai, W.-K., Jin, X.-F., Gituru, R. W., Wang, Q.-F. & Yang, C.-F. (2013). Competition and facilitation among plants for pollination: can pollinator abundance shift the plant–plant interactions? *Plant Ecology*, 215, 3–13.

Young, J.-G., Cantwell, G. T. & Newman, M. E. J. (2020). Bayesian inference of network structure from unreliable data. *Journal of Complex Networks*, 8.

Young, J.-G., Valdovinos, F. S. & Newman, M. E. J. (2019). Reconstruction of plant–pollinator networks from observational data.

Appendix 1: List plant species observed in Caracoles (2020)

Table A1. List of plant species observed in Caracoles Ranch with floral visitors from a total of 23 species surveyed in the community. Taxonomic family of each species is provided. Sample size represents the total number of individuals sampled for reproductive success for each focal species, and it is correlated with their natural abundance observed at the study site.

Species	Family	Abundance	Sample size
Beta macrocarpa	Amaranthaceae	1,992	250
Centaurium tenuiflorum	Gentianaceae	1,946	190
Chamaemelum fuscatum	Asteraceae	1,347	151
Chamaemelum mixtum	Asteraceae	143	14
Leontodon maroccanus	Asteraceae	9,090	269
Melilotus sulcatus	Fabaceae	1,100	53
Pulicaria paludosa	Asteraceae	1,418	137
Scorzonera laciniata	Asteraceae	791	117
Sonchus asper	Asteraceae	1,012	106
Spergularia rubra	Caryophyllaceae	2,318	58

Appendix 2: Plant phenologies

Plant phenologies were extracted from our records of insect visits. Detailed weekly surveys of floral visitors during the flowering season showed that insects only visited the flowers of 10 of the species, and most of those visits were concentrated on only three of them (*Chamaemelum fuscatum*, *Leontodon maroccanus* and *Pulicaria paludosa*). As can be seen in Fig. A2.1, the phenologies of the plant species that received insect visits are varied, being able to differentiate early, medium and late phenologies. The earlier species is *C. fuscatum*; the medium species are *L. maroccanus*, *Spergularia rubra*, *Scorzonera laciniata*, *Sonchus asper*, *Chamaemelum mixtum*, *Melilotus sulcatus* and *Beta macrocarpa*; and the late species are *P. paludosa*, *Centaurium tenuiflorum*, and in 2020 a second flowering peak of *L. maroccanus*.

Fig. A2.1. Total amount of insect visits received by plant species per week in 2020.

To estimate the duration of plant phenologies (in weeks), we assumed that, if a plant species received insect visits in weeks i and j (for $i \neq j$), then such plant species were also visited during the period between i and j. For instance, B. macrocarpa's phenology lasts 6 weeks (from the 17th week to the 22nd one).

Finally, in Fig. A2.2 we disaggregate the phenological overlap among plant species by using the functional groups of flower visitors.

Fig. A2.2. Total amount of insect visits received by plant species and functional group of visitors per week in 2020.

Appendix 3: List of functional groups, families and species of Caracoles (2020)

Table A3. Number of visits observed in Caracoles Ranch by functional group, family and identifier.

Functional groups, **Family**, ID	Visits
Big beetles	**5**
Cerambycidae	**2**
Cerambycidae	2
Meloidae	**3**
Lagorina sericea	3
Butterflies	**13**
Geometridae	**2**
Geometridae	2
Lasiocampidae	**5**
Lasiocampa trifolii	5
Nymphalidae	**2**
Vanessa cardui	2
Pieridae	**4**
Euchloe crameri	1
Pieris brassicae	3
Flies	**1**
Diptera	**1**
Other diptera	1
Flower beetles	**395**
Chrysomelidae	**6**
Cryptocephalus sp.	6
Elateridae	**13**
Elateridae	13
Melyridae	**345**
Malachius bipustulatus	9
Melyridae	1

Psilothrix viridicoerulea	335
Mordellidae	**16**
Mordellidae	16
Oedemeridae	**15**
Oedemeridae	15
House flies	**82**
Calliphoridae	**6**
Calliphoridae	6
Muscidae	**44**
Musca sp.	44
Sarcophagidae	**23**
Sarcophaga sp.	23
Tachinidae	**9**
Cylindromyia sp.	9
Hoverflies	**62**
Syrphidae	**62**
Episyrphus balteatus	14
Eristalis sp.	3
Eupeodes corollae	1
Lomatia sp.	9
Sphaerophoria scripta	32
Syrphidae	3
Humbleflies	**57**
Bombyliidae	**57**
Anastoechus sp.	44
Bombilus major	13
Small beetles	**756**
Anthicidae	**10**
Anthicidae	10
Chrysomelidae	**2**
Cassida sp.	1
Phaedon sp.	1
Curculionidae	**1**
Curculionidae	1
Nitidulidae	**743**
Brassicogethes sp.	743
Small flies	**135**
Bibionidae	**4**
Dilophus sp.	4
Empididae	**2**
Empis sp.	2
Stratiomyidae	**6**
Nemotelus sp.	6
Ulidiidae	**123**
Ulidiidae	123
Solitary bees	**280**
Andrenidae	**136**
Andrena argentata	4
Andrena humilis	76
Andrena sp.	56
Apidae	**4**
Eucera sp.	4
Halictidae	**126**
Lasioglossum immunitum	4
Lasioglossum malachurum	111
Lasioglossum sp.	11
Megachilidae	**14**

Osmia ligurica	14
Wasp	**8**
Braconidae	**8**
Braconidae	8
Total	**1,794**

Appendix 4: Characterization of community structure

As can be seen in Fig. A4.1, the main visitors by functional groups were small beetles (756), followed by pollenivorous flower beetles (395), solitary bees (280) and small flies (135). The remaining visits are distributed between house flies (82), hoverflies (62), humbleflies (57), butterflies (13), wasps (8), big beetles (5) and flies(1). Regarding plants, during 2020, only ten focal species received insect visits in Caracoles, namely: *L. maroccanus* (1,337), *C. fuscatum* (268), *P. paludosa* (111), *C. tenuiflorum* (23), *M. sulcatus* (15), *B. macrocarpa* (14), *C. mixtum* (14), *S. laciniata* (7), *S. asper* (3), and *S. rubra* (2).

Fig. A4.1. Number of insect visits per functional group, plant species and plot.

Fig. A4.2. Number of individuals with visits per plant species.
Table A4. Characterization of the modules in each plot. Values in parenthesis show the results with 95% confidence intervals.

Plot	Plant individuals with visitors	Plant visitors	Number of modules	Plant species per module	Visitors per module	Plant individuals per module
1	49	23	10	1.20 (1, 3)	2.00 (1, 8)	4.90 (1, 28)
2	66	30	11	1.55 (1, 3)	1.91 (1, 8)	6.00 (1, 29)
3	66	30	14	1.64 (1, 3)	1.50 (1, 4)	4.71 (1, 13)
4	26	11	7	1.14 (1, 2)	1.43 (1, 2)	3.71 (1, 14)
5	19	8	6	1 1	1.33 (1, 3)	3.17 (1, 10)
6	14	7	3	1 1	2.33 (1, 4)	4.67 (1, 10)
7	43	22	14	1.29 (1, 3)	1.21 (1, 2)	3.07 (1, 10)
8	84	41	16	1.81 (1, 4)	1.56 (1, 5)	5.25 (1, 26)
9	74	34	16	1.50 (1, 3)	1.50 (1, 3)	4.62 (1, 21)

Regarding insect nodes per module, the average amount of them is 2.12 with CI (1, 6), and the amount of different species and morphospecies per module is 1.57 with CI (1, 7.92). Overall, there are 64 modules with a single insect species (or morphospecies), 24 with two different species, 4 modules with three, 2 modules with four, 1 with 5, and 2 modules with 8. Additional details on modules features can be found in table A4.

Fig. A4.3. Number of insect visitors per module, functional group and plot.

Fig. A4.4. Distribution of number of plant species, insect visitors species, and plant individuals per modules.

Fig. A4.5. Distribution of number of plant species per module and plot.

Fig. A4.6. Distribution of insect visitors species per module and plot.

Fig. A4.7. Distribution of number of plant individuals per module and plot.

Appendix 5: Specialization roles of plant individuals and floral visitors

Plant individuals and insect species within a module may exert a different impact on the pollen flow that they trap. For instance, depending on their centrality metrics, nodes may hoard the pollen influx and/or outflux. Besides, in the case of insect species, those that connect several plant species and modules can be responsible for distributing high loads of heterospecific pollen. To assess the roles of plant individuals and insect species in each multilayer, we classified them in the following categories: (i) peripherals, i.e., vertices linked to very few nodes that usually are within their module; (ii) module hubs, i.e., highly connected vertices linked to many nodes within their own module; (iii) connectors, i.e., nodes that link several modules; and (iv) network hubs, i.e., nodes acting as both connectors and module hubs (Olesen *et al.*, 2007). To do so, we calculated their weighted standard among module connectivity and within module degree, denoted by c and z, respectively (Dormann & Strauss, 2013; Olesen *et al.*, 2007). In unweighted and undirected networks, the former represents the level to which a given node is linked to other modules, and the later is its standardized number of links to other species in the same module (Guimerà & Amaral, 2005). Since our multilayers are weighted and directed, we must distinguish between the links that arrive at a node (or in-links) and those that depart from it (or out-links). For that reason, we adapted the previous definitions and computed two values of c (namely: in- and out-c) and two of z (in- and out-z) by using in- and out-strength of each node, which correspond to the sum of the weights of its in-links and the one for its out-links, respectively. From the in-links (out-links) perspective, we assigned the following roles to nodes: peripherals [low in-(out-)c and low in-(out-)z], module hubs [low in-(out-)c and high in-(out-)z], connectors [high in-(out-)c and low in-(out-)z], and network hubs [high in-(out-)c and high in-(out-)z].

To define the thresholds for in/out-c- and in/out-z-values, we followed Watts *et al.* (2016) and used the 95% quantiles of such metrics obtained from a null model for our original multilayer networks. Specifically we generated an ensemble of 100 null multilayers for each plot, respectively, and we calculated in/out-c- and in/out-z-values for each randomized multilayer. Our null model preserves the total number of observed interactions per insect visitor and assumes that the weight of interlinks are those of the observed multilayers. We estimated the thresholds for in/out-c and in/out-z over all the plots (i.e., over the results for 900 null multilayers), and their values are 1.00/1.00 and 1.93/1.62, respectively.

By using those thresholds, overall, around 30 - 40% of the plant and insect nodes were peripherals (with all their in- and out-links outside their own module), 40 - 60% were connectors among modules, 2.5 - 10% of insect species acted as module hubs, and 2.5 - 10% of insect species were network hubs. Since insect species channel all the links between conspecific and heterospecific plant individuals (see Fig. 1(e)), the observed absence of plant hubs is expected. In addition, our results also confirmed that most network and module hubs belong to the main insect visitors of the most visited plant species, namely: flower beetles, humbleflies, small beetles, bees. Specific details are as follows. From the in-link (or pollen influx to nodes) perspective, we found that 37.87% of all plant individuals and 29.61% of all floral visitors were peripherals (with c greater than 0, i.e., they all have in-links outside their own module), 58.50% of plant individuals and 50.00% of insect visitors were connectors among modules, 9.71% of insect visitors were module hubs, whereas 2.91% of insects were network hubs (Figs. A5.1 and A5.2). Influx roles for the remaining plant individuals (3.63%) and insect visitors (7.77%) can not be computed because the standard deviation of their respective within-layer centralities is equal to zero, and consequently their within module in-strength (i.e. in-z), which is defined in multiples of the within-layer centralities' standard deviations Dormann & Strauss (2013); Olesen *et al.* (2007)(Olesen et al. 2007, Dormann et al. 2009), can not be properly assessed. Influx network hubs were *Lasioglossum malachurum* (bee) in *P. paludosa* (plot 8), Ulidiidae (small flies) in *C. fuscatum* (plots 1, 2 and 3), *Psilothrix viridicoerulea* (flower beetles) in *L. maroccanus* layers (plot 6), and *Bombylius sp.* (humbleflies) in *C. tenuiflorum* (plot 2), whereas module hubs were *Anastoechus sp.* (humbleflies) in *L. maroccanus* (plot 3), *Andrena humilis* (solitary bees) in *L. maroccanus* (plot 7), *Andrena sp.* (solitary bees) in *C. fuscatum* (plot 9), *Brassicogethes sp.* (small beetles) in *L. maroccanus* (all plots), *Lasioglossum malachurum* (bee) in *P. paludosa* (plot 3), *Psilothrix viridicoerulea* (flower beetles) in *C. fuscatum* (plot 9) and *L. maroccanus* (plots 3, 4, 5, 7, and 8), and Ulidiidae (small flies) in *C. fuscatum* (plot 8). Results for the out-links (or pollen outflux from nodes) show that 37.87% of all plant individuals and 34.47% of all floral visitors were peripherals (with c greater than 0), 51.25% of plant individuals and 37.86% of insect visitors were connectors among modules, 2.91% of insect visitors were module hubs, whereas only 0.45% of plant individuals and 10.68% of insects were network hubs (see Figs. A5.3 and A5.4). Outflux roles for 10.88% of plant individuals and 14.08% of insect visitors are omitted. Outflux network hubs were *Anastoechus sp.* (humbleflies) in layer *C. tenuiflorum* (plot 2), *Andrena humilis* (solitary bees) in *S. laciniata* (plot 8), *Bombilus major* (humbleflies) in *P. paludosa* (plot 3), *Brassicogethes sp.* (Small beetles) in *C. fuscatum* (plots 3 and 9), *P. paludosa* (plot 8) and *S. laciniata* (in plot 8 and 9, respectively); *Empis sp.* (small flies) in *C. fuscatum* (plot 1); *Lasioglossum malachurum* (bees) in *C. tenuiflorum* (2 and 3), *M. sulcatus* (plot 8), and *P. paludosa* (1, 4, 8, and 9); Mordellidae (flower beetles) in *S. asper* (plot 9); *Musca sp.* (house flies) in *P. paludosa* (plot 8); *Psilothrix viridicoerulea* (flower beetles) in *C. fuscatum* (plot 3) and *P. paludosa* layers (plot 8); *Sphaerophoria scripta* (hoverflies) in *M. sulcatus* (plot 7); and Ulidiidae (Small flies) in *C. fuscatum* (plot 2), whereas module hubs are *Anastoechus sp.* (humbleflies) in

P. paludosa (plot 3); *Bombylius major* (humbleflies) in *P. paludosa* (plot 2); *Brassicogethes sp.* (small beetles) in *L. maroccanus* (plot 9); *Lasioglossum malachurum* (solitary bees) in *P. paludosa* (plot 3); and *Psilothrix viridicoerulea* (flower beetles) in *C. fuscatum* and *S. asper* (both in plot 9).

Fig. A5.1. Roles performed by plant individuals in the plot multilayers according to within module in-strength (in-z) and among module in-strength (in-c). Lines at in-z = 1.93 and in-c = 1.00 define species roles.

Fig. A5.2. Roles performed by floral visitors in the plot multilayers according to within module in-strength (in-z) and among module in-strength (in-c). Lines at in-z = 1.93 and in-c = 1.00 define species roles.

Fig. A5.3. Roles performed by plant individuals in the plot multilayers according to within module out-strength (out-z) and among module out-strength (out-c). Lines at out-z = 1.62 and out-c = 1.00 define species roles.

Fig. A5.4. Roles performed by floral visitors in the plot multilayers according to within module out-strength (out-z) and among module out-strength (out-c). Lines at out-z = 1.62 and out-c = 1.00 define species roles.

Appendix 6: Specialization roles of plant individuals and floral visitors

After estimating the map equation (in bits, $L_{Observed}$) and the number modules ($m_{Observed}$) for the multilayer of a given plot, we want to know whether these values are significant compared with a random expectation. To contrast the significance of such measures, we proposed two null models. In our first null model (Null model 1), the total number of observed interactions per plant species and per insect visitor are preserved in each layer. That is, only intra-layer connections with plant individuals are randomly reshuffled in this null model. Thus, the experimental phenology of each plot and the inter-links that codify it are equal to those of observed networks. Regarding our second null model (Null model 2), it preserves the total number of observed interactions per insect visitor and assumes that the weight of interlinks are those of the observed multilayers.

To derive a null distribution of map equation and number of modules for each null model, we generated an ensemble of 500 null multilayers for each plot, respectively. For each randomized multilayer, we have calculated the map equation ($L_{Simulated}$) and number of modules ($m_{Simulated}$). Then, we estimated the confidence interval for each null distribution and tested whether the observed values of our metrics belong to such intervals (non-significant

result) or not (significant result). Our results are summarized in tables A6.1 and A6.2, and Figs. A6.1, A6.2, A6.3 and A6.4. As can be seen, in each plot $L_{Observed}$ is significantly smaller than the results for randomized multilayer with a confidence level of 95%. Consequently, observed network flows tend to be more constrained within modules than in any of the random multilayers. On the other hand, we found no significant differences between $m_{Observed}$ and the number of modules of randomized multilayers with a confidence level of 95%.

Table A6.1. Observed map equation ($L_{Observed}$) and number of modules ($m_{Observed}$) for each plot, and the corresponding confidence intervals (C.I.) for the simulated map equation ($L_{Simulated}$) and simulated number of modules ($m_{Simulated}$), when the null model 1 is considered. Null distributions for $L_{Simulated}$ and $m_{Simulated}$ are shown in Figs.A6.1 and A6.2. Bold values denote significant results.

NULL MODEL 1

Plot	$L_{Observed}$	C. I. for $L_{Simulated}$	$m_{Observed}$	C. I. for $m_{Simulated}$
1	**3.709**	(3.908, 4.126)	**10**	(4, 8)
2	**3.971**	(4.367, 4.553)	11	(8, 11)
3	**3.696**	(3.943, 4.297)	14	(11, 16)
4	**3.259**	(3.678, 4.141)	7	(4, 8)
5	**3.126**	(3.175, 3.538)	6	(2, 6)
6	**2.997**	(3.093, 3.611)	3	(3, 5)
7	**3.680**	(3.851, 4.189)	**14**	(7, 11)
8	**3.539**	(4.019, 4.315)	16	(14, 19)
9	**3.701**	(3.940, 4.230)	16	(13, 19)

Fig. A6.1. Null distribution of map equation (in bits) obtained from 500 simulations for each plot. Each panel represents the results for a plot. The respective observed values are marked with vertical blue dashed lines.

Table A6.2. Observed map equation ($L_{Observed}$) and number of modules ($m_{Observed}$) for each plot, and the corresponding confidence intervals (C.I.) for the simulated map equation ($L_{Simulated}$) and simulated number of modules ($m_{Simulated}$), when the null model 2 is considered. Null distributions for $L_{Simulated}$ and $m_{Simulated}$ are shown in Figs. A6.3 and A6.4. Bold values denote significant results.

NULL MODEL 2

Plot	$L_{Observed}$	C. I. for $L_{Simulated}$	$m_{Observed}$	C. I. for $m_{Simulated}$
1	**3.709**	(4.590, 4.755)	**10**	(11, 16)
2	**3.971**	(4.940, 5.111)	**11**	(15, 19)
3	**3.696**	(4.744, 4.955)	14	(14, 19)
4	**3.259**	(3.716, 4.192)	7	(4, 8)
5	**3.126**	(3.481, 3.937)	6	(2, 7)
6	**2.997**	(3.376, 3.980)	3	(2, 6)
7	**3.680**	(4.512, 4.743)	14	(10, 16)
8	**3.539**	(4.733, 4.961)	**16**	(17, 23)
9	**3.701**	(4.658, 4.856)	16	(16, 22)

Fig. A6.2. Null distribution of number of modules obtained from 500 simulations for each plot. Each panel represents the results for a plot. The respective observed values are marked with vertical blue dashed lines.

Fig. A6.3. Null distribution of map equation (in bits) obtained from 500 simulations for each plot by using Null model 2. Each panel represents the results for a plot. The respective observed values are marked with vertical blue dashed lines.

Fig. A6.4. Null distribution of number of modules obtained from 500 simulations for each plot by using Null model 2. Each panel represents the results for a plot. The respective observed values are marked with vertical blue dashed lines.

Appendix 7: Supplementary figures for centrality measures

Fig. A7.1. Dependence of (multilayer) pollen arrival probabilities (PageRank) on within-layer centrality (in-strength) and plot. Linear regression lines are displayed in blue, with their corresponding coefficient of determination (R) and p-values.

Fig. A7.2. Dependence of the among-layer centrality ratio on plot and plant species. Mean values are shown with red diamonds.

Appendix 8: Supplementary figures for centrality measures

After calculating the total amount of homospecific and heterospecific motifs for those plant individuals that received insect visits, we contrasted the significance of such values by comparing them with a random expectation. To do so, we proposed a null model in which the total number of observed visits per plant species and per insect visitor are preserved in each plot and week. In that model, we randomly reshuffled the insect visits received by one plant species among those plant individuals that were present in a certain plot, during a given week. Thus, the experimental phenology observed in each plot was preserved.

To derive a null distribution of the total amount of homospecific and heterospecific motifs per focal individual, we generated an ensemble of 500 null networks for each plot and week. For each randomized realization, we calculated the number of homospecific and heterospecific motifs per plant individual, insect visitor, and week; and we pooled the results for plant individuals. Then, since null distributions are not usually Gaussian (see Fig. A8.1 for an example), we estimated their confidence intervals for our metrics and tested whether the observed values belong to such intervals (non-significant result) or not (significant result). Our findings are summarized in Figs. A8.2 and A8.3. According to such figures, 46.71% (82.09%) of plant individuals have a total amount of homospecific (heterospecific) motifs that is significantly different from the values obtained for randomized systems, with a confidence level of 95%. In the case of homospecific motifs, it is possible to see that, as expected, plant species with few (many) plant individuals tend to exhibit a higher (lower) proportion of significant results, specifically significantly larger (smaller) number of homospecific motifs. Remarkably, in the case of *C. fuscatum*, we observe that most of its homospecific triples are non-significant. This is due to the reduced phenological overlap among *C. fuscatum* and other plant species (see Fig. A2.1). Finally, it is worth mentioning that 97.27% of individual plants with significant heterospecific motifs exhibit larger values of such motifs than the randomized arrangements.

Fig. A8.1. Null distribution of homospecific motifs for the focal individual of *L. maroccanus* located in the plot 1 and subplot A3, obtained from 500 simulations. The observed value is marked with a vertical blue dashed line, and it is significantly smaller.

Fig. A8.2. Number of plant individuals with significant and non-significant homospecific motifs per plot and plant species.

Fig. A8.3. Number of plant individuals with significant and non-significant heterospecific motifs per plot and plant species.