High nucleotide substitution rates associated with retrotransposon proliferation drive

dynamic secretome evolution in smut pathogens

Depotter JRL[a], Ökmen B[a], Ebert MK[a], Beckers J[a], Kruse J[bc], Thines M[bc], Doehlemann G[a]*

[a]University of Cologne, CEPLAS, Institute for Plant Sciences, Cologne, Germany

[b]Senckenberg Biodiversity and Climate Research Centre (BiK-F), Frankfurt a. M., Germany.

[c]Institute of Ecology, Evolution and Diversity, Goethe University Frankfurt, Frankfurt a. M.,

Germany.

*Corresponding author

E-mail: g.doehlemann@uni-koeln.de (GD)

Key words: Ustilago, transposable element, genome expansion, DNA polymerase, mating-

type locus.

Abstract

Transposable elements (TEs) play a pivotal role in shaping diversity in eukaryotic genomes.

The covered smut pathogen on barley, *Ustilago hordei*, encountered a recent genome

expansion. Using long reads, we assembled genomes of 6 *U. hordei* strains and 3 sister

species, to study this genome expansion. We found that larger genome sizes can mainly be

attributed to long terminal repeat retrotransposons (LTR-RTs) of the *Copia* and *Gypsy*

superfamilies. From the studied smuts, LTR-RTs proliferated the most recently and to the

furthest extent in the *U. hordei* genome, in which they make up for 19.5% of the genome.

Interestingly, the extent of TE proliferation in different smut species is positively correlated

to the mating-type locus size, which is largest in *U. hordei* with up to ~560 kb. TE

transposition within the mating-type loci and their flanking regions are mating-type specific,

which is likely due to the very low recombination activity in this region. Furthermore, LTR-

RT proliferation was found to be associated with higher nucleotide substitution levels, as

genes in genome regions that are rich in dynamic LTR-RTs display higher nucleotide

substitution levels. The high nucleotide substitution rate particularly affected the evolution of

genes encoding secreted proteins as substitutions more frequently led to amino acid

alterations. The mechanism behind this increase in nucleotide substitution rate remains

elusive, but seems not to be a consequence of the repeat-induced point mutation (RIP)

mechanism, as genes and LTR-RTs did not display typical RIP substitutions.

INTRODUCTION

Transposable elements (TEs) play a pivotal role in the genome evolution of eukaryotic organisms, including fungi (1). Fungal genomes can vary considerably in size, which is often determined by the extent of TE proliferation (2, 3). On one side of the spectrum, Microsporidia, a diverse group of obligate intracellular parasitic fungi, contain members with extremely small genomes down to 2.3Mb that lack TEs (4, 5). In contrast, rust plant pathogens from the order Pucciniales contain members with genome sizes that are among the largest in the fungal kingdom (6, 7). For instance, the wheat stripe rust pathogen *Puccinia striiformis* f.sp. *tritici* has an estimated genome size of 135 Mb, which more than half consists of TE sequences (8). Mutations caused by TE transposition predominantly have a neutral or negative impact, but in particular cases they can also improve fungal fitness (3). For plant pathogenic fungi, TE transposition can be a source of mutagenesis to evade host immunity and/or lead to an optimized host interaction (9). TEs can also passively contribute to mutagenesis, as their transpositions increase homologous genome sequences that are prone to ectopic recombination (10, 11). Pathogens evolve by host jumps, radiation and subsequent arms races with their hosts (12), in which the latter attempts to detect pathogen ingress through the recognition of so-called invasion patterns (13). One invasion pattern that is typically detected are effectors, i.e. secreted proteins that facilitate host colonization (14). To quickly adapt to effector-triggered immunity and yet continue host symbiosis, effector genes often reside in genome regions that facilitate mutagenesis (12, 15), such as those rich in TEs (16). TE-rich genome regions may not only encounter higher mutation rates, but may also have a higher chance to fix mutations due to their functionally more accessory nature (17).

TEs are a diverse group of mobile nucleotide sequences that are categorized into two classes (1). Class I comprises retrotransposons that transpose through the reverse transcription of their messenger RNA (mRNA). Class II are DNA transposons that transpose

without mRNA intermediate. TEs are then further classified based on their sequence structure

(18). Retrotransposons with direct repeats at each end of their sequence are long terminal

repeat retrotransposons (LTR-RTs) (19). LTR-RTs can encode the structural and enzymatic

machinery for autonomous transposition. However, LTR-RT may lose this ability through

mutagenesis, but still be able to transpose using proteins of other LTR-RTs (20). These non-

autonomous LTR-RT derivates can be classified into large retrotransposon derivative

(LARD) and terminal-repeat retrotransposon in miniature (TRIM) depending on their size.

LTR-RTs can then be further classified into superfamilies including *Copia* and *Gypsy*, which

differ in the order of their reverse transcriptase and their integrase domain (18). A major type

of DNA transposons are terminal inverted repeats (TIRs). Terminal repeats are recognized by

the transposase enzyme for excision. Non-autonomous TIR derivatives that cannot encode

their own transposase are classified as miniature inverted TEs (MITEs) (21).

TE activity can have detrimental outcomes for the hosting genome. Consequently,

organisms have evolved several protection mechanisms to prevent uncontrolled TE

proliferation (22). TE activity can be epigenetically suppressed through DNA methylation,

histone modification or RNA interference (RNAi). Epigenetic TE suppression is reversible

and, thus, TEs can be re-activated, which typically occurs under particular stress conditions

(23). Alternatively, TEs can be irreversibly de-activated through mutagenesis (24). In fungi,

repeat-induced point mutation (RIP) is a meiosis-associated defence mechanism that

generates G to A and C to T transitions in duplicated sequences (25). RIP was first observed

in the ascomycete *Neurospora crassa* and has hitherto been experimentally verified in a

limited number of other ascomycetes (26–31). In basidiomycetes, RIP has never been

experimentally verified, but biases towards RIP substitution in repetitive DNA have been

reported (32–34).

Smut fungi are a diverse group of plant pathogenic basidiomycetes of which many

infect monocot plants, in particular grasses (35). Generally, these pathogens have small

genomes compared to other plant pathogens (36, 37). The corn smut species *Ustilago maydis*

and *Sporisorium reilianum* are closely related and have genome sizes of 19.8 Mb and 18.4

Mb, respectively (38, 39). This is partly due to their low level of repetitive sequences,

including TEs. In total, only 2.1 and 0.5% of the genome consists of TEs for the *U. maydis*

and *S. reilianum*, respectively (40). *Ustilago hordei* and *U. brachipodii-distachyi* are two

related smut fungi and have assembled genomes of 21.15 Mb and 20.5 Mb, respectively (33,

40). These larger genome assembly sizes correlate to their higher TE content, which is 11.8%

and 14.3 for *U. hordei* and *U. brachipodii-distachyi*, respectively (40). The assembled

genome of *U. brachipodii-distachyi* is originally published under the species name *U.*

bromivora (Rabe et al. 2016). *U. brachipodii-distachyi* infect members from the tribe

Brachipodieae, whereas *U. bromivora* affects bromes from the supertribe Triticodae (41, 42).

Considering the host specific nature of smut pathogens, we prefer to refer to this assembly as

U. brachipodii-distachyi instead of *U. bromivora*, as the assembled strain infects

Brachypodium species (40, 41).

Mating in grass-parasitic smut fungi is tetrapolar in *U. maydis* and *S. reilianum*,

whereas *U. hordei* and *U. brachipodii-distachyi* have a bipolar mating system. In the bipolar

system, there is one mating-type locus where recombination is suppressed (40, 43, 44). This

locus is flanked by the *a* locus, which contains pheromone/receptor genes, and the *b* locus,

which encodes the two homeodomain proteins bEast and bWest (45, 46). In the bipolar

mating-type system, there are two mating-type alleles, *MAT-1* and *MAT-2*, which are in *U.*

hordei ~500kb and ~430kb in size, respectively (44). A large fraction of the mating-type loci

consists of repetitive sequences, i.e. ~45% repeats for *U. hordei* (33, 47). In contrast, the

tetrapolar smuts, *U. maydis* and *S. reilianum*, have there *a* and *b* loci on different

chromosomes causing them to segregate independently during meiosis (39, 43).

Recently, the complete genome of the reference *U. hordei* strain Uh364 was re-

sequenced using the long-read PacBio technology and, instead of the previous 21.15 Mb

assembly (33), a 27.1 Mb assembly was obtained (48). Thus, the *U. hordei* genome

underwent a recent genome expansion as it is more than 30% larger than that of its relative *U.*

brachipodii-distachyi, which genome is assembled on chromosome level and is 20.5 Mb in

size (40). This finding triggered us to study the *U. hordei* genome more in depth and use

recently developed long-read sequencing technology to unravel how its genome recently

expanded.

Results

LTR-RT proliferation drives *U. hordei* genome size

To study the expansion in genome size of *U. hordei*, we sequenced 6 *U. hordei* strains of
different geographic origin (Figure S1). To obtain a high-quality assembly with large contigs
and high base calling confidence, we used paired-end Illumina reads in combination with
long reads obtained with the Oxford Nanopore MinION device. In total, 3.2-4.5 Gb of filtered
paired-end reads and 1.5-6.5 Gb of filtered Nanopore reads were used for assembly. The
resulting assemblies were composed of 23-46 contigs and ranged from 25.8-27.2 Mb in size
(Table 1). In total, 98.6-99.0% of the basidiomycete Benchmarking Universal Single-Copy
Orthologs (BUSCOs) were present in the *U. hordei* assemblies (Table 1) (49) which is
similar to the quality of the *U. maydis* reference genome assembly (98,8% of the BUSCOs)
(38). Strain Uh805 was assembled into 23 contigs that are homologous to the 23
chromosomes of *U. brachipodii-distachyi* indicating that inter-chromosomal rearrangements
did not occur during the divergence of these two species (40). Repetitive sequences in the
assemblies were identified and classified into TE orders (Table 1). In general, class I TE
sequences (retrotransposons) are over 6 times more abundant than class II TE sequences
(DNA transposons) in *U. hordei*. More than 90% of the class I TEs consist of LTR-RTs,
which is a total sequence amount ranging between 4,326 and 5,272 kb (Figure 1A, Table 1).
For the class II TE sequences, more than 75% are TIRs with a total sequence amount ranging
from 589 to 667 kb (Table 1). The number of LTR-RT sequences is positively correlated to
the assembly size (r = 0.94, p-value = 0.0051), whereas the number of TIR TEs is not (r = -
0.25, p-value = 0.63). Moreover, using strain Uh805 as a reference, 56-79% of the differences
in assembly size with other strains can be attributed to differences in LTR-RT content. In
conclusion, the variation in genome size between *U. hordei* strains can largely be attributed to
intraspecific differences in LTR-RT proliferation and retention.

Table 1. Transposable elements annotation in various smut genome assemblies.

Species	U. hordei						U. nuda	U. brachipodii-distachyi[40]	U. tritici	U. loliicola	U. maydis[38]	S. reilianum[50,51]
Strain	Uh359	Uh805	Uh811	Uh818	Uh1273	Uh1278	DE_29490	UB2112	Ut_3	K530	521	SRS1_H2-8
Assembly size (Mb)	27.0	25.8	26.2	26.2	27.2	26.6	21.4	20.4	20.4	20.8	19.7	18.5
Contigs	46	23	26	25	37	27	31	23	32	41	27	23
BUSCOs (%)	98.9	98.9	98.9	98.9	98.6	99.0	98.9	99.1	98.9	98.8	98.8	98.5
Total repeats (%)	38.2	35.3	36.4	36.5	38.9	36.0	22.6	17.0	16.4	8.9	4.6	3.6
Class I TEs (kb)§	5,625	4,611	4,985	4,897	5,663	4,940	1,786	672	739	102	199	5
LTR (kb)	5,272	4,326	4,615	4,549	5,208	4,607	1,537	463	462	9	185	5
LARD (kb)	126	54	213	136	233	104	106	39	43	65	11	0
TRIM (kb)	4	16	11	11	11	12	4	1	4	0	0	0
DIRS (kb)	163	117	59	117	97	122	98	135	119	0	0	0
LINE (kb)	8	0	0	5	22	7	9	4	71	27	3	0
SINE (kb)	0	1	0	0	1	2	1	0	1	1	0	0
Class II TEs (kb)§	781	746	708	791	731	692	395	473	285	482	5	103
TIR (kb)	604	619	589	667	612	622	342	418	242	322	5	100
MITE (kb)	12	13	11	18	13	12	11	9	5	88	0	2
Crypton (kb)	31	30	34	34	23	30	42	45	38	0	0	0
Maverick (kb)	38	11	22	21	31	15	0	0	0	0	0	0
Helitron (kb)	92	73	47	45	51	3	0	0	0	72	0	0

§ Only repetitive sequences that were larger than 500 bp were classified

The *U. hordei* secretome is activated upon plant colonization, whereas LTR-RTs are generally inactive

To guide gene annotation and to study gene/LTR-RT expression, RNA was extracted and sequenced from *U. hordei* grown in liquid medium and in barley leaves at 3 days post infection (dpi). RNA 150bp paired-end reads were obtained through the Illumina NovaSeq6000 System. In total, 5.1-8.4 and 36.0-45.2 Gb of raw reads were obtained for the samples grown in liquid medium and *in planta*, respectively. Genes were annotated with BRAKER with the strain Uh803 assembly as reference using RNA-Seq data and protein homology (53). In total, 7704 genes were predicted of which 396 were predicted to encode for proteins with a signal peptide and without a transmembrane domain, hereafter referred to as secreted proteins. RNA-seq reads were mapped to the Uh803 genome assembly and quantified for gene and LTR-RT loci. In total, 6229 of the 7704 (81%) predicted gene loci were expressed in either of the two *U. hordei* growth conditions, whereas only 19 of the 904 (2%) LTR-RTs displayed expression (more than one count per million of expressed sequence in at least two of the six samples) (Figure 1B). Moreover, only 7 of the expressed LTR-RTs displayed expression in more than half of their sequence in at least one of the samples. Of these 7, there was one *Copia* and one *Gypsy* LTR-RT that can be autonomous, as functional domains for aspartyl protease, reverse transcriptase and integrase could be identified. In conclusion, almost all LTR-RTs were inactive in the two tested environmental conditions. For the genes, 978 (16%) were upregulated *in planta*, whereas 1178 (19%) were downregulated (Figure 1B). Up- and downregulated genes were screened for Gene Ontology (GO) term enrichments, to see which biological processes are affected upon plant colonization. In total, 33 and 23 GO terms were enriched in *in planta* up- and down-regulated genes, respectively (Table S1). Generally, processes associated with the fungal membrane, including transmembrane transport were upregulated *in planta*, whereas cytoplasmic

processes associated with ribosome, mitochondrial membrane and proteasome were

downregulated (Table S1). In correspondence with these results, 214 of the 978 genes

upregulated *in planta* encode secreted proteins. Thus, 59% (214/360) of the expressed genes

that encode secreted proteins were upregulated *in planta*, which is a significant enrichment

(Fisher exact test, *p*-value = 1.26e-86) (Figure 1B). In contrast, only 10% of the genes

encoding a secreted protein were downregulated. Of these downregulated genes, 35% (13/37)

were predicted to have a carbohydrate-active (CAZyme) function, whereas this is 17%

(36/214) for *in planta* upregulated secretome genes, which is significantly less (Fisher exact

test, *p*-value = 0.014). In conclusion, the *U. hordei* transmembrane transport system and

secreted proteins are strongly activated upon plant colonization, whereas hardly any LTR-

RTs display expression.

The majority of the mating-type locus sequences are mating-type specific

Several plant pathogens have genomes compartmentalized in in a way that facilitate fast

effector repertoire evolution (12), such as repeat-rich regions (16). To test if TE-rich regions

are the cradle for the evolution of new pathogenicity effectors in *U. hordei,* the location of

secreted proteins and abundance of repeats and presence/absence polymorphisms were

plotted across the *U. hordei* genome (Figure 2). Strain Uh805 was used as a reference as it

contains 23 contigs that represent separate chromosomes. In these analyses, we considered

windows of 10 kb upstream of 5'-ends and 10kb downstream of 3'-ends of *U. hordei* genes

and determined the fraction of repeats and presence/absence polymorphisms within these 20

kb windows surrounding genes. Upregulated genes encoding secreted proteins, which are

promising effector candidates (15, 54), have a median of 13% of repetitive sequences in their

flanking region. This is not significantly different from other expressed genes that have a

median of 17% (t-test, *p*-value = 0.32). Thus, in contrast to some other filamentous plant

pathogens, *in planta* upregulated genes encoding secreted proteins are not especially

associated with repeat-rich genome regions in *U. hordei*. For presence/absence

polymorphisms, we calculated the fraction of the window that is absent in at least one of the

five other *U. hordei* strains. Here, upregulated genes encoding secreted proteins and other

expressed genes were also not significantly different from each other (t-test, *p*-value = 0.36,

both median = 0). In conclusion, *U. hordei* effector candidates, i.e. *in planta* upregulated

genes encoding secreted proteins, are not enriched in repeat-rich genome regions that

encounter frequent presence/absence polymorphisms.

When we plotted the *U. hordei* chromosomes, our attention was drawn to a ±816 kb

genome region on chromosome 1 between genome coordinates roughly at 950-1,766 kb,

which in only 27% of the region had a one-to-one best alignment to strain Uh1278, whereas

this ranged from 95% to 100% with other strains (Figure 2-3A). Moreover, this region is

repeat rich as it consists of 76% repeats and 36% of classified LTR-RTs, which is around

double the genome wide average of 35% and 17%, respectively (Table1). The *MAT-1* locus,

the region between and including the *a* and *b* loci, is embedded in this region and is 560 kb in

size, consisting of 77% of repetitive sequences (Table S2). Uh1278 is the only sequenced

MAT-2 U. hordei strain in this study (Tables S2). The low number of one-to-one best

alignment to strain Uh1278, indicates that most sequences within the mating-type locus and

its flanking regions are mating-type specific (Figure 3A). Similarly to *MAT-1*, *MAT-2* resides

in a genome region of ±829 kb that has a high repeat (77%) and LTR-RT (35%) content

(Figure 3a). The sequenced *MAT-2* locus is 472 kb, which is smaller than any of the

sequenced *MAT-1* loci that ranged from 536 to 564 kb (Table S2). Homologous

recombination is suppressed in the mating-type region (44). Consequently, any mutation

within these regions is by definition mating-type specific. The repetitive nature and the large

size of the mating-type loci indicates that many mating-type specific sequences likely

originate from TE transpositions (Figure 3B).

LTR-RT proliferation in smuts with a bipolar mating-type system

As LTR-RT played a predominant role in the genome expansion of *U. hordei*, we also

studied the impact of TE dynamics on the genome evolution of *U. hordei* sister species. We

sequenced genomes of *Ustilago loliicola*, *Ustilago nuda* and *Ustilago tritici*, which are smut

species that are close relatives of *U. hordei* and *U. brachipodii-distachyi* (41, 55). Using the

PacBio Sequel system we obtained a total of 6.3-9.7 Gb of raw long reads for the strains

listed in Table 1. Assemblies of 20.8, 21.4 and 20.4 Mb were obtained in 41, 31 and 32

contigs for *U. loliicola, U. nuda* and *U. tritici*, respectively. In total, 98.8-98.9% of the

BUSCOs were present in the assemblies (Table 1) (49). A phylogenetic tree was constructed

based on 1,667 BUSCO proteins including the newly sequenced species as well as *U.*

brachipodii-distachyi, *U. maydis* and *S. reilianum* (Figure 4A) (38, 40, 50, 51). *U. hordei, U.*

nuda, *U. brachipodii-distachyi* and *U. tritici* cluster together (HNBT-cluster hereafter) with

U. loliicola being the closest outgroup species. Within the cluster, *U. hordei* diverged the

most recently from *U. nuda*, which also infects *Hordeum* species (Figure 4A). Synteny

between the different contigs was also investigated and the ancestral gene order

reconstructed. Intriguingly, *S. reilianum* and *U. loliicola* do not have inter-chromosomal

rearrangement in comparison to their reconstructed last common ancestor (Figure 4A). The

U. maydis genome has one inter-chromosomal rearrangement with respect to its last common

ancestor with *S. reilianum.* HNBT-cluster species share one inter-chromosomal

rearrangement that occurred after their divergence from *U. loliicola* (Figure 4A). As a

consequence of this rearrangement, the *a* and *b* mating-type loci came to reside on the same

chromosome in linked to each other, resulting in the mating-type polarity switch from

tetrapolar to bipolar (43). This inter-chromosomal rearrangement is the only one observed in

the assemblies of *U. brachipodii-distachyi, U. nuda* and *U. hordei*, whereas the *U. tritici*

assembly has one additional inter-chromosomal rearrangement. In conclusion, HNBT-cluster

species have a bipolar mating system due to an inter-chromosomal rearrangement that

occurred after their divergence from *U. loliicola*.

To see if mating-type polarity relates to TE evolutionary dynamics, repeats were

annotated and classified in the TE orders for the different genomes in a similar fashion to the

U. hordei strains. In total, 22.6% of the *U. nuda* assembly consists of repetitive sequences,

which is 13-16% less than *U. hordei* from which *U. nuda* recently diverged (Table 1, Figure

4A). *Ustilago brachipodii-distachyi* (17.0 %) and *U. tritici* (16.4%) have similar levels of

repetitive sequences in their genomes (Table 1). The *U. loliicola*, *U. maydis* and *S. reilianum*

assemblies, which outgroup the HNBT-cluster, consist 8.9, 4.6 and 2.7% of repeats,

respectively. Thus, species of the HNBT-cluster underwent an expansion of repeat content in

their genome (Figure 4A). Interestingly, the repeat and the LTR-RT content of the HNBT-

cluster species genomes positively correlates to the size of their mating-type loci ($r = 0.98$, p-

value $= 0.02$, using strain Uh805 for *U. hordei*), which ranges from 190 kb for *U.*

brachipodii-distachyi to 560 kb for *U. hordei* (Figure 4A, Table S2). This confirms our

hypothesis that the size of the *MAT* locus is determined by TE proliferation and the insertion

of TEs in the *MAT* locus and its flanking regions (Figure 3B).This increase and interspecific

differences in repeat content between HNBT-cluster species can largely be attributed to

differences in LTR-RT content that ranged from 462 kb in *U. tritici* to 4,326 kb in *U. hordei*

(Uh805) (Table 1). In contrast, *U. loliicola, U. maydis* and *S. reilianum* contain 9, 185 and 5

kb of LTR-RT sequences, respectively (Table 1). Thus, LTR-RTs proliferated after the

divergence of *U. loliicola* and the HNBT-cluster, which is also the moment when the mating-

type polarity switched from tetrapolar to bipolar.

High local nucleotide substitution levels are associated with LTR-RT proliferation

Although HNBT-cluster species collectively encountered a LTR-RT proliferation, there are large differences within the cluster as *U. hordei* has more than 9 times the amount of LTR-RT sequences than *U. tritici* (Table 1). To study intraspecific differences, the nucleotide sequence identity distribution of the best reciprocal paralogous and orthologous LTR-RT sequences was calculated (Figure 4B). This was on the one hand done for the recently diverged species with the highest LTR-RT content, *U. hordei* and *U. nuda,* and on the other hand for *U. brachipodii-distachyi* and *U. tritici*. The distribution of the paralogous LTR-RTs in *U. brachipodii-distachyi* and *U. tritici* displayed two maxima at 82-83% and at 88-90% (Figure 4B). The maximum of the orthologous LTR-RTs between *U. brachipodii-distachyi* and *U. tritici* was at 93%. Thus, LTR-RTs in *U. brachipodii-distachyi* and *U. tritici* mainly proliferated before the last common ancestor of the HNBT species (Figure 4A). Orthologous LTR-RTs between *U. hordei* and *U. nuda* displayed a maximum at 89%, whereas for paralogous LTR-RTs a maximum at 93% was present for *U. nuda* and two maxima at 94 and 97% for *U. hordei* (Figure 4B). Thus, in contrast to *U. brachipodii-distachyi* and *U. tritici*, LTR-RTs continued to proliferated after the HNBT last common ancestor.

As TE-active genome regions have been associated with distinct nucleotide substitution regimes in fungi (10, 56), we studied if different extents of LTR-RT proliferation is associated with different nucleotide substitution regimes. Gene nucleotide divergence in *U. brachipodii-distachyi*/*U. tritici* and *U. hordei*/*U. nuda* was tested for clustering, where a cluster was defined to comprise at least 10 of a fraction of the most diverged orthologs in a window of 20 genes. Orthologs are defined to encode proteins with at least 80% identity and 80% overlap. Using this definition, the fraction of clustering increases with increasing fraction of the most divergent orthologs, until around 25% where the clustering plateaus

(Figure 4C). We further compared the clustering between species by using 20% of the most

divergent genes as its central within the 15 and 25% range where the clustering progresses in

a more or less constant linear relationship with the percentage of the most diverged orthologs

(Figure 4C). These clusters using 20% most diverged genes are hereafter referred to as

Diverged Gene Clusters (DGCs). For *U. brachipodii-distachyi*/*U. tritici* 24% of the 20%

most diverged orthologs reside in DGCs, which is at least two times less than for ortholog

comparison involving *U. hordei* and/or *U. nuda* (48-65%) (Figure 4C). Thus, gene nucleotide

sequence divergence occurs more clustered in *U. hordei* and *U. nuda* than in *U. brachipodii-*

distachyi and *U. tritici*, which might be linked to the more recent LTR-RT proliferation of *U.*

hordei and *U. nuda* (Figure 4A-B). In total, 89% and 78% of the *U. brachipodii-distachyi*

DGC genes are the same using *U. hordei* (1,346/1,507) and *U. nuda* (1,346/1,718) orthologs,

respectively (Figure S2). Thus, this indicates that clustered higher nucleotide substitution

rates occurred to a similar extent in *U. hordei* and *U. nuda* since their last common ancestor

and mainly affect the same genome regions.

310   We then looked if DGCs are distinct in gene density, intraspecific presence/absence

polymorphisms or LTR-RT content from other genome regions (Figure S3). For this, genome

windows of 20 kb were used and the DGCs were constructed based on the 20% most

diverged *U. hordei* genes with *U. brachipodii-distachyi* (Figure 2). We compared these two

species as (1) LTR-RT proliferated to the furthest extent in *U. hordei* and (2) *U. brachipodii-*

distachyi is its closest relative where LTR-RT did not extensively proliferate after the HNBT

last common ancestor (Figure 4). The median 20 kb window in *U. hordei* DGCs consist 46%

of expressed gene sequence. This is significantly less than the median of 51% for genome

regions outside DGCs (t-test, *p*-value = 0.025) (Figure S3). The fraction of Uh803 sequences

that is absent in at least one of the other sequenced *U. hordei* strains is not significantly

different within and outside DGCs (t-test, *p*-value = 0.12) (Figure S3). The fraction of

sequences within DGCs that is annotated as LTR-RT has a median of 21%, whereas this is

0% and significantly less for other genome regions (t-test, *p*-value = 2.95e-5) (Figure S3). In

conclusion, DGCs are generally gene poorer and LTR-RT richer than other genome regions.

To test if higher substitution levels are associated with LTR-RT proliferation, we

looked if interspecific LTR-RT variation especially occurs in DGCs. For genes that have a

single ortholog in every species of the HNBT-cluster, a window of 10 kb upstream of its 5'-

end and 10 kb downstream of its 3'-end was constructed, and the fraction of this 20 kb

window that consists of LTR-RT was determined. Then, the absolute difference of this

fraction for every *U. hordei* gene was calculated with its corresponding ortholog of one of the

HNBT-cluster species. These differences were then compared for genes that reside in DGCs

and genes that do not. The median differences in LTR-RT content of *U. hordei* genes in

DGCs were 8, 9 and 6% using the orthologs of *U. nuda*, *U. brachipodii-distachyi* and *U.*

tritici, respectively (Figure 5). In contrast, genes that do not reside in DGCs have a median

change of 0% independent of the species that was used for orthologs (Figure 5). This

difference in LTR-RT alterations is significant (Welch's t-test; *U. nuda*: *p*-value = 5.63e-13,

U. brachipodii-distachyi: *p*-value = 8.52e-23 and *U. tritici*: *p*-value = 4.92e-15). The same

phenomenon was also observed for *U. brachipodii-distachyi* DGC genes when compared to

U. hordei orthologs (Figure 5). Genes inside DGCs had a median LTR-RT alteration of 8%,

whereas this was 0% for genes outside DGCs (Welch's t-test, *p*-value = 1.17e-20). For the *U.*

nuda and *U. tritici* orthologs, medians of 0% occurred inside and outside *U. brachipodii-*

distachyi DGCs (Figure 5). Conceivably, the difference in LTR-RT content of *U.*

brachipodii-distachyi with *U. nuda* and *U. tritici* were too small to find significance (Table

1). In conclusion, differences in LTR-RT content between *U. hordei* and the other species

occur more frequently in genome regions were genes encountered a higher nucleotide

substitution level.

LTR-RTs do not display typical RIP substitutions

RIP is an genome defence mechanism that induces substitution levels in repetitive nucleotide sequences. However, RIP leaks to genome regions neighbouring these repeats, which can affect coding regions flanking TEs (57). To test if LTR-RTs are affected by RIP in HNBT-cluster species, we constructed ancestor LTR-RT sequences of LTR-RT families, using the convention that TE family members share at least 80% sequence identity in at least 80% of their sequence with one other family member (18). To facilitate the sequence alignment and ancestor sequence construction, we only took a subset of the LTR-RTs and excluded the terminal repetitive sequences (more details in Material and Methods). In total, ancestors of 13 TE-LTR families were reconstructed using 252 LTR-RT sequences (Figure 6A). The largest three families contained 65, 56 and 31 LTR-RTs respectively and contained only *U. hordei* and *U. nuda* members. The ancestor of the two largest families were from the *Copia* superfamily, whereas the third largest belong to *Gypsy* and contained a chromodomain at the C-terminal region of the integrase, so-called chromovirus (58). The substitutions that occurred between this subset of LTR-RTs and their respective ancestor are on average 84% transitions (Figure 6B). RIP targets repeats through C→T and G→A substitutions. On average, 19% of the encountered LTR-RT substitutions since their ancestor are T→C, which is significantly smaller than the fraction of 24% for the typical RIP substitution C→T (t-test, *p*-value = 1.10e-23). Conversely, the typical RIP substitution G→A occurred with an average of 19% less frequently, than A→G substitutions (average 22%) (Figure 6B) (t-test, *p*-value = 5.27e-7). In conclusion, there is no clear enrichment for C→T and G→A substitutions between the LTR-RTs and their reconstructed ancestor.

High local nucleotide substitution levels especially impact secretome evolution

To see if higher nucleotide substitution levels affect a particular group of genes more than others, we tested if genes residing in DGCs are enriched for particular GO terms. Also here we opted to choose *U. hordei* and *U. brachipodii-distachyi* for comparison due to their respective extensive and lack of LTR-RT proliferation after the HNBT last common ancestor (Figure 4). No single GO term was enriched in *U. hordei* DGC genes indicating that a higher local nucleotide substitution rate does not affect particular gene types more than others. Also, DGC genes were not more likely to encode secreted proteins (Fisher exact test, *p*-value = 0.80 Although high nucleotide substitution levels affect all type of genes, the consequences for the evolution of particular genes might be different. Presence/absence polymorphisms were compared within and outside DGCs. In DGCs, 27% of the *U. hordei* genes do not have an *U. brachipodii-distachyi* ortholog that encode for a protein with at least 80% identity and 80% overlap. Outside DGCs, 23 % of the *U. hordei* genes do not have such orthologs, which is significantly less than inside DGCs (Fisher exact test, *p*-value = 0.00047). Intriguingly, this difference is pronounced for genes that encode secreted proteins. Inside DGCs, 44% of the secretome genes are *U. hordei* specific, whereas this is 27% outside DGCs (Fisher exact test, *p*-value = 0.0038). Thus, the increased level of presence/absence polymorphisms in DGCs affects especially genes encoding secreted proteins.

We also studied the nucleotide substitution types. Substitutions were determined for *U. hordei* and *U. brachipodii-distachyi* orthologs (at least 35% identity and 40% overlap) that are single copy genes (using the same 35% and 40% cut-offs), as paralogs can impact the nucleotide substitution regime (59). In general, transitions had medians ranging from 0.016 to 0.019 transitions per codon outside DGCs, whereas transversions ranged from 0.0018 to 0.0056 transitions per codon (Figure 7A). Thus, similar to the LTR-RT substitutions, transitions occur more frequently than transversion in gene coding regions. Within DGCs,

median transitions and transversion ranged from 0.021 to 0.025 and from 0.0037 to 0.024

substitutions per codon, respectively (Figure 7A). All type of substitutions were significantly

higher for DGCs genes compared to genes outside DGCs (Welch's t-test, p-value < 0.001)

(Figure 7A). Intriguingly, the number of CG substitutions for DGC genes had a median that

was 4.26 times higher than for genes outside DGCs, whereas this ranged from 1.29 to 2.03

for other substitution types (Figure 7A). Although *U. hordei* encountered more recent and a

higher extant of LTR-RT proliferation than *U. brachipodii-distachyi* (Figure 4B, Table 1),

C→T, T→C, A→G and G→A substitutions between these two species were all higher inside

DGCs compared to outside (Figure 7B). Thus, the higher nucleotide substitution rate in

DGCs does not display typical RIP substitutions patterns, rather all substitution types are

increased.

Nucleotide substitutions in coding regions can be synonymous or nonsynonymous in

nature. Outside *U. hordei* DGCs, genes contain a median of 0.04 ds, whereas this is 0.06

within DGCs, which is a significant increase (Welch's t-test, p-value = 2.60e-78) (Figure

7C). Similarly, the median dn is 0.01 outside and 0.02 inside DGCs, respectively (Welch's t-

test, p-value = 5.85e-35) (Figure 7D). Within DGCs, the median fraction of dn and ds (dn/ds)

is 0.35 for genes that encode a secreted protein, whereas this is significantly less for other

genes that have a median of 0.30 (Welch's t-test, p-value = 0.031) (Figure 7E). In conclusion,

the higher nucleotide substitution levels in DGCs do more frequently lead to amino acid

changes in the secreted proteins than in proteins that are not secreted.

Discussion

Nucleotide substitution rates are unevenly distributed across genomes and can be influenced

by numerous factors, including neighbouring nucleotides, recombination frequencies and TE

activity (60–62). Nucleotide divergence in *U. hordei* and *U. nuda* occurs more clustered in

their genomes compared to *U. brachipodii-distachyi* and *U. tritici* (Figure 4C). These former

two smut species encountered a more recent and higher extent of LTR-RT proliferation

compared to the latter two (Figure 4B). LTR-RT proliferation appears to be associated with

higher nucleotide substitution rates, as *U. hordei* genome regions with high nucleotide

substitution levels are enriched for dynamic LTR-RTs (Figure 5). However, the exact

mechanism by which this occurs remains uncertain. The occurrence of genome regions with

higher nucleotide substitution rates is likely not a consequence of RIP, as LTR-RTs do not

more frequently encounter the typical RIP substitutions (Figure 6). Moreover, the higher local

nucleotide substitution levels can be observed for all substitution types (Figure 7A). The

associated higher nucleotide substitution levels in repeat-rich genome regions has been

previously found in other eukaryotes (63). It has been proposed that repeat-rich genome

regions are replicated by more error-prone polymerases. More specifically, it has been

suggested that DNA polymerase errors may be introduced during TE insertion. In rice, higher

nucleotide substitutions levels occur close to LTR-RT insertion sites (62). LTR-RT insertion

causes cuts in the host DNA that is then ligated by the host (64, 65). Wicker et al. (2016)

suggest that cut host DNA might become a target for 3'->5' exonuclease resulting in a

segment of single-stranded DNA (62). The complementary strand of this stretch of DNA

would then be synthesized by a replication complex with lower DNA polymerase fidelity and

mismatch repair. Conversely, higher substitution rates can also be the cause rather the

consequence why LTR-RT proliferation is observed in genome regions with higher

nucleotide substitution levels. Genome regions with pre-existing higher substitution regimes

may be more prone to TE fixation, as the more frequent substitutions increase the chance of

TE inactivation. Inactive TEs are generally less harmful for the host and may therefore have

lower chances to be selected against. However, the less clustered occurrence of high

nucleotide substitution levels in the LTR-RT poorer species *U. brachipodii-distachyi* and *U.*

tritici indicates that LTR-RT proliferation is the driver of clustered high nucleotide

substitutions rather than the consequence (Figure 4C).

Local higher levels of nucleotide substitutions impact the evolution of the genes that

reside in the affected genome regions. Generally, more nonsynonymous substitutions can be

observed (Figure 7D), but it is especially the genes that are encoding secreted proteins that

encounter amino acid alterations (Figure 7E). Conceivably, the generally more accessory

nature of secretome genes can make them more prone for amino acid changes (66). Although

the effect of nucleotide substitutions affected a particular gene type more than others,

enrichments for particular gene types could not be found. Also, genome regions with higher

nucleotide substitution levels, which are gene poorer and LTR-RT richer (Figure S3), are not

enriched for secretome genes, which include effector candidates. Thus, the model that

effector candidates often reside in accessory, repeat-rich genome regions that facilitates fast

adaptation, so-called two speed genome model (17, 37), does not hold for the DGCs. More

generally, repeat content and intraspecific presence/absence polymorphisms, features that

have been associated with the two-speed genome model, were not more frequently found in

the proximity of effector candidate genes compared to other genes (Figure 2) (56, 67). The

specificity and the universality of the two-speed genome model for filamentous plant

pathogens has recently been contested (68, 69). More plant pathogens have been reported

where effector candidates do not especially reside in gene-poor/repeat-rich regions, such as

the leaf spot pathogen *Ramularia collo-cygni* on barley, the earlier mentioned *P. striiformis* f.

sp. *tritici* and the barley powdery mildew pathogen *Blumeria gramin*is f. sp. *hordei* (70–72).

LTR-RT proliferation is the main factor that led to the *U. hordei* genome expansion

(Table 1). The expansion occurred especially in the mating-type locus that increased almost

three times in size in comparison to *U. brachipodii-distachyi* (Table S2). The lack of the

purifying recombination ability in this genome region can be the reason why LTR-RTs

especially accumulated in the mating-type and flanking genome regions (33, 44).

Conceivably, this process is reinforced by the increasing presence of repetitive sequences as

the transposition into a repeat-rich genome region is less likely to have a severe fitness cost

than a transposition in repeat-poor region. Furthermore, the co-occurrence of the LTR-RTs

proliferation in the HNBT branch and the switch in mating-type organization from tetra- to

bipolar may indicate that mating-type polarity switch impacts LTR-RT proliferation (73). In

case of biallelic *a* and *b* loci, the switch from a tetra-to bipolarity results in a basidiospore

compatibility change from 25 to 50%. Consequently, it takes a tetrapolar smut on average

longer to find a mating type than a bipolar smut. This longer time might increase the

opportunity to mate with spores from a different offspring and, thus, increase outcrossing.

The higher outcrossing rate for tetra- compared to bipolar smuts is even more pronounced

when multiallelism exists for the *a* and *b* loci (74). Multiallelism increased the compatibility

on population level, whereas compatibility within the same offspring remains 25 and 50% for

tetra- and bipolar smuts, respectively. Lower levels of outcrossing reduce the purifying

recombination ability of smuts, which may be the reason why LTRs could proliferate to a

further extent in bipolar smuts (33, 73).

TEs are import drivers of genome evolution as they directly cause mutagenesis

through their transpositions and indirectly increase the change of non-homologous

recombination due to their repetitive nature (11). LTR-RT proliferation in *U. hordei* indicates

that TE activity also influence local nucleotide substitution regimes and increase the

substitution levels in the genome regions where they insert. Consequently, genes in the

proximity of these insertion sites evolve faster (Figure 7D). Fast gene evolution may be

advantageous under stressful condition, when TEs are typically more active or change their

activity (23, 75).

Material and Methods

Genome sequencing and assembly

Genomic DNA from all smut species was isolated using a MasterPure™ Complete DNA&RNA Purification Kit (Epicentre®, Illumina®, Madison, Wisconsin, USA) according to the manufacturer's instructions. Long *U. hordei* reads were obtained with the Oxford Nanopore MinION device. The genomes of six *U. hordei* strains were sequenced: Uh359, Uh805, Uh811, Uh818, Uh121 and Uh122 (9). The library was prepared according the Oxford Nanopore Technology (ONT) protocol for native barcoding genomic DNA (EXP-NBD104 and SQK-LSK109). Three *U. hordei* strains were multiplexed for every run. The prepared library was loaded on an R9.4.1 Flow Cell. ONT reads were base-called, filtered (default value) and barcodes were trimmed with the Guppy Baseballing Software v3.5.1 of ONT. Paired-end *U. hordei* 150 bp reads were obtained with the Illumina HiSeq 4000 device. Library preparation (500bp insert size) and sequencing were performed by the BGI Group (Beijing, China). Paired-end *U. hordei* reads were filtered using Trimmomatic v0.39 with the settings "LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:100", only reads that remained paired after filtering were used in the assembly (76). An initial assembly was obtained by using the "ONT assembly and Illumina polishing pipeline" (https://github.com/nanoporetech/ont-assembly-polish). The assembly was further upgraded using the FinisherSC script (77) Mitochondrial contigs were remove from the assembly and was not used for any analysis. Additionally, small contigs were removed that contained a paired-end read coverage lower than 50% of the genome-wide average.

Long *U. loliicola, U. nuda* and *U. tritici* reads were obtained through Single Molecular Real-Time (SMRT) Sequencing using the PacBio Sequel system. *U. loliicola* K530 was isolated by Julia Kruse from *Lolium perenne* in Untermerzbach, Germany. *U. nuda* DE_29490 and *U. tritici* Ut_3 were both isolated by Karl-Josef Müller in Darzau, Germany.

The initial assembly was obtained using the Canu assembler and was further upgraded with

the FinisherSC script (77, 78). Mitochondrial contigs were remove from the assembly and

were not used for further analysis.

The quality of genome assemblies was assessed by screening the presences of

BUSCOs using the BUSCO software version 5.0.0 with the database "basidiomycota_odb10"

(49).

Transposable element annotation and classification

The smut genome assemblies were scavenged for repetitive sequences in order to construct a

repeat library for repeat annotation. Helitron TEs were identified using the EAHelitron script

(79). LTR-RTs were identified using LTRharvest (80). Miniature inverted-repeat TEs were

identified with MITE Tracker (81). Short interspersed nuclear elements were identified with

the SINE-scan tool (82). Finally, RepeatModeler (v1.0.11) was also used for *de novo* repeat

identification. These repeats were than combined with the repeat library from RepBase

(release20170127) (83). The CD-HIT-EST tool under default settings was used to remove

redundancy in the constructed library (84). RepeatMasker (v4.0.9) was then used to annotate

the repeats to specific genome locations. The annotated repeat sequences were filtered on size

and only sequences larger than 500bp were retained. Furthermore, repeats that were nested or

had more than 50% overlap with other repeats were removed from the library. In case two

repeats had reciprocally 50% overlap was the longest repeat retained. Repeats were classified

into different TE orders using the PASTEC tool using PiRATE-Galaxy (85, 86).

U. hordei **RNA sequencing and expression analysis**

Total RNA from *U. hordei* strain 4857-4 strain grown axenically and *in planta* was extracted

for three biological replicates. For the axenic samples, *U. hordei* was grown in YEPS light

(0.4% yeast extract, 0.4% peptone, and 2% saccharose) liquid medium at 22°C with 200 rpm

shaking till OD:1.0. For the *in planta* samples, Golden Promise barley cultivar was grown in

a greenhouse at 70% relative humidity, at 22°C during the day and the night; with a light/dark

regime of 15/9 hrs and 100 Watt/m^2 supplemental light when the sunlight influx intensity was

less than 150 Watt/m^2. Barley plants were infected with *U. hordei* through needle injection as

previously described (87) and samples were harvested 3dpi. Here, the third leaves of the *U.*

hordei infected barley plants were collected by cutting 1 cm below the injection needle sites.

Leaf samples were then frozen in liquid nitrogen and grinded using a mortar and pestle under

constant liquid nitrogen. The total RNA was isolated by using the TRIzol® extraction method

(Invitrogen; Karlsruhe, Germany) according to the manufacturer's instructions. Subsequently,

total RNA samples were treated with Turbo DNA-Free™ Kit (Ambion/Applied Biosystems;

Darmstadt, Germany) to remove any DNA contamination according to the manufacturer's

instructions. Total RNA was then sent to for library preparation and sequencing to Novogene

(Beijing, China). Libraries (250-300 bp insert size) were loaded on Illumina NovaSeq6000

System for 150bp paired-end sequencing using a S4 flowcell.

The reads were filtered using the Trinity software (v2.9.1) option trimmomatic under

the standard settings (88). The reads were then mapped to the reference genome using Bowtie

2 (v2.3.5.1) with the first 15 nucleotides on the 5'-end of the reads being trimmed due to

inferior quality (89). The reads were mapped onto a combined file of the *U. hordei* strain

Uh114 genome assembly and the *Hordeum vulgare* (IBSC_v2) (90) genome assembly. Reads

were counted to the *U. hordei* loci using the R package Rsubread (v1.34.7) (91). Here the

default minimum mapping quality score of 0 was used, to include reads that would have

multiple best mapping locations. For the gene loci, reads were counted that were mapped to

the predicted coding regions. For the LTR-RT loci, reads were only counted that mapped

within LTR-RT loci, excluding the reads that mapped onto the 10% of either edge of the

locus. Significant differential expression of a locus was calculated using the R package edgeR

(v3.26.8) (92). Significance of differential expression was calculated using t-tests relative to a

threshold of log2 fold change of 1 with Benjamini-Hochberg correction. The cut-off of 0.05

was used.

Gene annotation

U. hordei genomes were annotated using the BRAKER v2.1.4 pipeline with RNA-Seq and

protein supported training with the options "--softmasking" and "--fungus" enabled (53).

RNA-seq reads from *U. hordei* grown in axenic culture and *in planta* (all replicates) were

mapped to the assemblies using TopHat v2.1.1 (93). Protein predictions from numerous

Ustilaginales species were used to guide the annotation, i.e. *Anthracocystis flocculosa*,

Melanopsichium pennsylvanicum, *Moesziomyces antarcticus*, *S. reilianum*, *U. brachipodii-*

distachyi, *U. hordei*, *U. maydis* (33, 38–40, 94–96). *U. nuda* and *U. tritici* genomes were also

annotated with the BRAKER v2.1.4 pipeline, but no RNA-seq data was used to guide the

annotation. The option "--fungus" was enabled and the previously published protein files of

the following species were used for protein supported training: *M. pennsylvanicum*, *S.*

reilianum, *U. brachipodii-distachyi* and *U. maydis* (38–40, 94). Our annotation of *U. hordei*

Uh805 was also included to train the annotation software. The *U. brachipodii-distachyi*

genome was previously annotated and this annotation was used for analysis (40). Predicted

genes that included an internal stop codon or did not start with a methionine were removed.

Secreted proteins are proteins with a predicted signal peptide using SignalP version

5.0 (97) and the absence of a transmembrane domain predicted with TMHMM2.0c in the

protein sequence excluding the signal peptide (98). Gene Ontology (GO) terms were

annotated to the *U. hordei* strain Uh114 protein prediction using InterProScan (v5.42-78.0)

(99). Significance of GO term enrichments in a subset of genes were calculated with a Fisher

exact test with the alternative hypothesis being one-sided (greater). The significance values of

the multiple enrichments were corrected according Benjamini and Hochberg (100).

Carbohydrate-Active enzymes (CAZymes) were annotated using the dbCAN2 meta server

(101, 102). A protein was considered a CAZyme if at least two of the three tools

(HMMER,DIAMOND and Hotpep) predicted a CAZyme function.

Comparative genomic analyses.

Phylogenetic trees were constructed based on BUSCOs from the database

"basidiomycota_odb10" that are present without paralog in all members of the tree (49). For

every gene, the encoded protein sequences were aligned using MAFFT (v7.464) option "--

auto" (103). These aligned protein sequences were then concatenated for every species and

used for tree construction using RAxML (v8.2.11) with substitution model

"PROTGAMMAWAG" and 100 bootstraps (104). Here, protein sequences that were present

in least than 60% of the tree members were excluded for tree construction.

Synteny block between the smut genome assemblies of were identified with SynChro

with DeltaRBH = 3 (105, 106). The genome assembly of the epiphytic yeast *Moesziomyces*

bullatus ex *Albugo* was included in this analysis to use as an outgroup (107). The ancestral

chromosome gene order was constructed with AnChro with Delta´ = 3 and Delta´´ = 3 (106,

108). Inter-chromosomal rearrangements, i.e. translocations of two blocks, were identified

with ReChro Delta = 10 (106, 108). No inter-chromosomal rearrangements in *U. nuda* could

be automatically detected by ReChro. Here, the inter-chromosomal rearrangement that lead to

a mating-type polarity switch was manually determined.

The absent/present polymorphisms between the *U. hordei* strains were determined

with NUCmer (version 3.1) from the MUMmer package with the option "--maxmatch" (109).

From the same package, delta-filter with the option "-1" was used to find the one-to-one

alignments. A BED format file was created for the genome regions of the reference genome

that had a one-to-one alignment to the query genome. The BED format files of the different

query genomes with the same reference genome were then combined and the coverage for

every genome region was determined with the software bedtools (v.2.29.2) to determine the

percentage of presence a particular genome region in the other *U. hordei* strains (110).

Bedtools was also used to divide the genome in windows for which particular genome feature

were calculated, such as gene density, repeat content and presence/absence polymorphisms.

LTR-RT evolution

To know the sequence identity distribution, the best orthologous and paralogous LTR-RTs

were identified using blastn (v2.2.31+) (111). LTR-RTs that did not belong to an LTR-RT

family of multiple members, were excluded from the analysis. Members of the same LTR-RT

family share at least 80% sequence identity in at least 80% of their sequence with at least one

other member (18). Orthologous or paralogous LTR-RTs that have reciprocally the highest

bit-score were used for analysis. The nucleotide identity distribution of these orthologous and

paralogous LTR-RTs was constructed using Gaussian Kernel Density Estimation with a

kernel bandwidth of 1.5.

To reconstruct the ancestor LTR-RTs, a subset of the HNBT-cluster LTR-RTs were

used. LTR-RTs were included that were larger than 3 kb and smaller than 15 kb.

Furthermore, repetitive sequences within the LTR-RT (>50 bp) were indicated using blastn

(v2.2.31+) and removed from the sequence (111). The region between the repeats were then

used for ancestor construction if this region was larger than 500 bp. Here, bedtools (v.2.29.2)

function "getfasta" was used (110). Open reading frames (ORFs) and there encoding amino

acid sequence of were determined with esl-translate (-l 50) as part of the Easel (v0.46)

package. Functional domain within these amino acid sequences were determined with

pfam_scan.pl (-e_seq 0.01) using the Pfam database version 32.0 (112). Only sequences were

included in the ancestor construction if they had at least 3 different Pfam domains from the

following domains: PF00078, PF00665, PF03732, PF07727, PF08284, PF13975, PF13976,

PF14223, PF17917, PF17919 and PF17921. All of these predicted Pfam domain had to

located on the same nucleotide strand in order to be used for ancestor construction. These

sequences were then grouped in families, according to the definition that family members

share at least 80% sequence identity in at least 80% of their sequence with at least one other

member (18). Ancestors were constructed for families with at least four members, using

prank (v.170427) with the options "-showall" and "-F". For every family was the last

common ancestor used that could be classified in *Copia* or *Gypsy* using the tool LTRclassifier

(113). If LTRclassifier was unable to classify the LTR-RT, a more recent ancestor was used

as a reference. Nucleotide substitutions between LTR-RTs and their constructed ancestor

were then determined after they were aligned using MAFFT (v7.464) options "--auto" (103).

Gene divergence

One-to-one orthologs and homologs between *U. hordei* strains were found using the SiLiX

(v.1.2.10-p1) software (114). Homolog groups consisting of two members, each one of a

different strain/species, were considered one-to-one homologs. Fractions of the most

divergent orthologs were determined using the sequence identity of the hit with the best bit

score using blastn (v2.2.31+) (111). DGCs were identified with CROC with a minimum

number of 10 genes that can be found for a cluster using a gene window of 20 (115). Here,

the Benjamini and Hochberg method was used to correct for multiple testing. To extract the

genes within DGCs the software bedtools (v.2.29.2) was used (110). Also genes that did not

have an orthologs were considered in the DGC analyses. To overlap of DGC genes was

presented in venn diagrams that were constructed using InteractiVenn (116). Nucleotide

substitutions for orthologs were identified after orthologs were aligned using MAFFT

(v7.464) options "--auto" (103). Synonymous and nonsynonymous substitutions between

orthologs were identified using SNAP (117).

Data accession

676 Raw RNAseq reads and genome assemblies are deposited at NCBI under the BioProject

677 PRJNA698760.

Acknowledgements

This work has been supported by the Alexander von Humboldt Foundation, the European

Research Council (ERC☐2017☐COG 771035, conVIRgens), and the Cluster of Excellence

on Plant Sciences (CEPLAS; Germany's Excellence Strategy–EXC☐2048/1 – Project ID:

390686111) and the University of Cologne. We also thank the Regional Computing Centre of

Cologne (RRZK) for access to the Cologne High Efficient Operating Platform for Science

(CHEOPS). We thank Guus Bakkeren for sharing the *U. hordei* strains with us and critically

reading the manuscript. We thank Karl-Josef Müller for the isolation and sharing of the *U.*

nuda and *U. tritici* strain with us.

References

1. Bourque G, Burns KH, Gehring M, Gorbunova V, Seluanov A, Hammell M, Imbeault
M, Izsvák Z, Levin HL, Macfarlan TS, Mager DL, Feschotte C. 2018. Ten things you
should know about transposable elements. Genome Biol 19:199.
2. Stajich JE. 2017. Fungal genomes and instights into the evolution of the kingdom, p.
619–633. *In* The Fungal Kingdom.
3. Stukenbrock EH, Croll D. 2014. The evolving fungal genome. Fungal Biol Rev 28:1-
12.
4. Katinka MD, Duprat S, Cornillott E, Méténler G, Thomarat F, Prensier G, Barbe V,
Peyretaillade E, Brottier P, Wincker P, Delbac F, El Alaoul H, Peyret P, Saurin W,

699   Gouy M, Weissenbach J. 2001. Genome sequence and gene compaction of the
700   eukaryote parasite *Encephalitozoon cuniculi*. Nature 414:450-453.

701 5.   Corradi N, Pombert JF, Farinelli L, Didier ES, Keeling PJ. 2010. The complete
702   sequence of the smallest known nuclear genome from the microsporidian
703   *Encephalitozoon intestinalis*. Nat Commun 1:77.

704 6.   Ramos AP, Tavares S, Tavares D, Silva MDC, Loureiro J, Talhinhas P. 2015. Flow
705   cytometry reveals that the rust fungus, *Uromyces bidentis* (Pucciniales), possesses the
706   largest fungal genome reported-2489Mbp. Mol Plant Pathol 16:1006-1010.

707 7.   Tavares S, Ramos AP, Pires AS, Azinheira HG, Caldeirinha P, Link T, Abranches R,
708   Silva M do C, Voegele RT, Loureiro J, Talhinhas P. 2014. Genome size analyses of
709   Pucciniales reveal the largest fungal genomes. Front Plant Sci 5:422.

710 8.   Cuomo CA, Bakkeren G, Khalil HB, Panwar V, Joly D, Linning R, Sakthikumar S,
711   Song X, Adiconis X, Fan L, Goldberg JM, Levin JZ, Young S, Zeng Q, Anikster Y,
712   Bruce M, Wang M, Yin C, McCallum B, Szabo LJ, Hulbert S, Chen X, Fellers JP.
713   2017. Comparative analysis highlights variable genome content of wheat rusts and
714   divergence of the mating loci. G3 7:361–376.

715 9.   Ali S, Laurie JD, Linning R, Cervantes-Chávez JA, Gaudet D, Bakkeren G. 2014. An
716   immunity-triggering effector from the barley smut fungus *Ustilago hordei* resides in an
717   Ustilaginaceae-specific cluster bearing signs of transposable element-assisted
718   evolution. PLoS Pathog 10:e1004223.

719 10. Faino L, Seidl MF, Shi-Kunne X, Pauper M, van den Berg GCM, Wittenberg AHJ,
720   Thomma BPHJ. 2016. Transposons passively and actively contribute to evolution of
721   the two-speed genome of a fungal pathogen. Genome Res 26:1091–1100.

722 11. Seidl MF, Thomma BPHJ. 2014. Sex or no sex: Evolutionary adaptation occurs
723   regardless. BioEssays 36:335–345.

724 12. Thines M. 2019. An evolutionary framework for host shifts – jumping ships for
725   survival. New Phytol 224:605-617.

726 13. Cook DE, Mesarich CH, Thomma BPHJ. 2015. Understanding plant immunity as a
727   surveillance system to detect invasion. Annu Rev Phytopathol 53:541–563.

728 14. Rouxel T, Balesdent MH. 2017. Life, death and rebirth of avirulence effectors in a
729   fungal pathogen of Brassica crops, *Leptosphaeria maculans*. New Phytol 214:526-532.

730 15. Depotter JRL, Doehlemann G. 2020. Target the core: durable plant resistance against
731   filamentous plant pathogens through effector recognition. Pest Manag Sci 76:426-431.

732 16. Dong S, Raffaele S, Kamoun S. 2015. The two-speed genomes of filamentous
733   pathogens: waltz with plants. Curr Opin Genet Dev 35:57–65.

734 17. Croll D, McDonald BA. 2012. The accessory genome as a cradle for adaptive
735   evolution in pathogens. PLoS Pathog 8:e1002608.

736 18. Wicker T, Sabot F, Hua-Van A, Bennetzen JL, Capy P, Chalhoub B, Flavell A, Leroy
737   P, Morgante M, Panaud O, Paux E, SanMiguel P, Schulman AH. 2007. A unified
738   classification system for eukaryotic transposable elements. Nat Rev Genet 8:973-982.

739 19. Finnegan DJ. 2012. Retrotransposons. Curr Biol 22:R432-R437.

740 20. Havecker ER, Gao X, Voytas DF. 2004. The diversity of LTR retrotransposons.
741   Genome Biol 5:225.

21. Jiang N, Feschotte C, Zhang X, Wessler SR. 2004. Using rice to understand the origin

and amplification of miniature inverted repeat transposable elements (MITEs). Curr
Opin Plant Biol 7:115-119.

22.   Slotkin RK, Martienssen R. 2007. Transposable elements and the epigenetic regulation
of the genome. Nat Rev Genet 8:272–285.

23.   Fouché S, Badet T, Oggenfuss U, Plissonneau C, Francisco CS, Croll D. 2020. Stress-
driven transposable element de-repression dynamics and virulence evolution in a
fungal pathogen. Mol Biol Evol 37:221-239.

24.   Kinsey JA, Garrett-Engele PW, Cambareri EB, Selker EU. 1994. The *Neurospora*
transposon Tad is sensitive to repeat-induced point mutation (RIP). Genetics 138:657-
664.

25.   Selker EU, Stevens JN. 1985. DNA methylation at asymmetric sites is associated with
numerous transition mutations. Proc Natl Acad Sci U S A 82:8114-8118.

26.   Selker EU, Cambareri EB, Jensen BC, Haack KR. 1987. Rearrangement of duplicated
DNA in specialized cells of *Neurospora*. Cell 51:741-752.

27.   Graïa F, Lespinet O, Rimbault B, Dequard-Chablat M, Coppin E, Picard M. 2001.
Genome quality control: RIP (repeat-induced point mutation) comes to *Podospora*.
Mol Microbiol 40:586-595.

28.   Ikeda KI, Nakayashiki H, Kataoka T, Tamba H, Hashimoto Y, Tosa Y, Mayama S.
2002. Repeat-induced point mutation (RIP) in *Magnaporthe grisea*: implications for its
sexual cycle in the natural field context. Mol Microbiol 45:1355-1364.

29.   Cuomo CA, Güldener U, Xu J-R, Trail F, Turgeon BG, Di Pietro A, Walton JD, Ma L-
J, Baker SE, Rep M, Adam G, Antoniw J, Baldwin T, Calvo S, Chang Y-L, Harris LJ,
Hilburn K, Kennell JC, Kroken S, Magnuson JK, Muehlbauer G, Münsterkötter M,
Nelson D, Donnell KO, Ouellet T. 2007. The *Fusarium graminearum* genome reveals
a link between localized polymorphism and pathogen specialization. Science
317:1400–1402.

30.   Coleman JJ, Rounsley SD, Rodriguez-Carres M, Kuo A, Wasmann CC, Grimwood J,
Schmutz J, Taga M, White GJ, Zhou S, Schwartz DC, Freitag M, Ma LJ, Danchin
EGJ, Henrissat B, Coutinho PM, Nelson DR, Straney D, Napoli CA, Barker BM,
Gribskov M, Rep M, Kroken S, Molnár I, Rensing C, Kennell JC, Zamora J, Farman
ML, Selker EU, Salamov A, Shapiro H, Pangilinan J, Lindquist E, Lamers C,
Grigoriev I V., Geiser DM, Covert SF, Temporini E, Vanetten HD. 2009. The genome
of *Nectria haematococca*: contribution of supernumerary chromosomes to gene
expansion. PLoS Genet 5:e1000618.

31.   Idnurm A, Howlett BJ. 2003. Analysis of loss of pathogenicity mutants reveals that
repeat-induced point mutations can occur in the Dothideomycete *Leptosphaeria*
maculans. Fungal Genet Biol 39:31-37.

32.   Horns F, Petit E, Yockteng R, Hood ME. 2012. Patterns of repeat-induced point
mutation in transposable elements of basidiomycete fungi. Genome Biol Evol 4:240-
247.

33.   Laurie JD, Ali S, Linning R, Mannhaupt G, Wong P, Guldener U, Munsterkotter M,
Moore R, Kahmann R, Bakkeren G, Schirawski J. 2012. Genome comparison of barley
and maize smut fungi reveals targeted loss of RNA silencing components and species-
specific presence of transposable elements. Plant Cell 24:1733–1745.

34.   Hane JK, Anderson JP, Williams AH, Sperschneider J, Singh KB. 2014. Genome
788   sequencing and comparative genomics of the broad host-range pathogen *Rhizoctonia*
789   *solani* AG8. PLoS Genet 10:e1004281.
790   35.   Zuo W, Odiekmen B, Depotter JRL, Ebert MK, Redkar A, Misas Villamil J,
791   Doehlemann G. 2019. Molecular interactions between smut fungi and their host plants.
792   Annu Rev Phytopathol 57:411–430.
793   36.   Möller M, Stukenbrock EH. 2017. Evolution and genome architecture in fungal plant
794   pathogens. Nat Rev Microbiol 15:756–771.
795   37.   Raffaele S, Kamoun S. 2012. Genome evolution in filamentous plant pathogens: why
796   bigger can be better. Nat Rev Microbiol 10:417–430.
797   38.   Kämper J, Kahmann R, Bölker M, Ma LJ, Brefort T, Saville BJ, Banuett F, Kronstad
798   JW, Gold SE, Müller O, Perlin MH, Wösten HAB, De Vries R, Ruiz-Herrera J,
799   Reynaga-Peña CG, Snetselaar K, McCann M, Pérez-Martín J, Feldbrügge M, Basse
800   CW, Steinberg G, Ibeas JI, Holloman W, Guzman P, Farman M, Stajich JE,
801   Sentandreu R, González-Prieto JM, Kennell JC, Molina L, Schirawski J, Mendoza-
802   Mendoza A, Greilinger D, Münch K, Rössel N, Scherer M, Vraneš M, Ladendorf O,
803   Vincon V, Fuchs U, Sandrock B, Meng S, Ho ECH, Cahill MJ, Boyce KJ, Klose J,
804   Klosterman SJ, Deelstra HJ, Ortiz-Castellanos L, Li W, Sanchez-Alonso P, Schreier
805   PH, Häuser-Hahn I, Vaupel M, Koopmann E, Friedrich G, Voss H, Schlüter T,
806   Margolis J, Platt D, Swimmer C, Gnirke A, Chen F, Vysotskaia V, Mannhaupt G,
807   Güldener U, Münsterkötter M, Haase D, Oesterheld M, Mewes HW, Mauceli EW,
808   DeCaprio D, Wade CM, Butler J, Young S, Jaffe DB, Calvo S, Nusbaum C, Galagan J,
809   Birren BW. 2006. Insights from the genome of the biotrophic fungal plant pathogen
810   *Ustilago maydis*. Nature 444:97–101.
811   39.   Schirawski J, Mannhaupt G, Münch K, Brefort T, Schipper K, Doehlemann G, Di
812   Stasio M, Rössel N, Mendoza-Mendoza A, Pester D, Müller O, Winterberg B, Meyer
813   E, Ghareeb H, Wollenberg T, Münsterkötter M, Wong P, Walter M, Stukenbrock E,
814   Güldener U, Kahmann R. 2010. Pathogenicity determinants in smut fungi revealed by
815   genome comparison. Science 330:1546-1548.
816   40.   Rabe F, Bosch J, Stirnberg A, Guse T, Bauer L, Seitner D, Rabanal FA, Czedik-
817   eysenberg A, Uhse S, Bindics J, Genenncher B, Navarrete F, Kellner R, Ekker H,
818   Kumlehn J, Vogel JP, Gordon SP, Walter MC, Marcel TC, Mu M, Sieber CMK,
819   Mannhaupt G, Gu U, Kahmann R, Djamei A. 2016. A complete toolset for the study of
820   *Ustilago bromivora* and *Brachypodium* sp. as a fungal-temperate grass pathosystem.
821   Elife 5:e20522.
822   41.   Kruse J, Dietrich W, Zimmermann H, Klenke F, Richter U, Richter H, Thines M.
823   2018. *Ustilago* species causing leaf-stripe smut revisited. IMA Fungus 9:49–73.
824   42.   Maire RCJ. 1919. Une ustilaginale nouvelle de la flore nord-Africaine. Bull la Société
825   d'Histoire Nat l'Afrique du Nord 10:46–47.
826   43.   Bakkeren G, Kronstad JW. 1994. Linkage of mating-type loci distinguishes bipolar
827   from tetrapolar mating in basidiomycetous smut fungi. Proc Natl Acad Sci U S A
828   91:7085-7089.
829   44.   Lee N, Bakkeren G, Wong K, Sherwood JE, Kronstad JW. 1999. The mating-type and
830   pathogenicity locus of the fungus *Ustilago hordei* spans a 500-kb region. Proc Natl

Acad Sci U S A 96:15026-15031.

45. Gillissen B, Bergemann J, Sandmann C, Schroeer B, Bölker M, Kahmann R. 1992. A
two-component regulatory system for self/non-self recognition in *Ustilago maydis*.
Cell 68:647-657.

46. Raudaskoski M, Kothe E. 2010. Basidiomycete mating type genes and pheromone
signaling. Eukaryot Cell 9:847-859.

47. Bakkeren G, Jiang G, Warren RL, Butterfield Y, Shin H, Chiu R, Linning R, Schein J,
Lee N, Hu G, Kupfer DM, Tang Y, Roe BA, Jones S, Marra M, Kronstad JW. 2006.
Mating factor linkage and genome evolution in basidiomycetous pathogens of cereals.
Fungal Genet Biol 43:655-666.

48. Yadav V, Sun S, Billmyre RB, Thimmappa BC, Shea T, Lintner R, Bakkeren G,
Cuomo CA, Heitman J, Sanyal K. 2018. RNAi is a critical determinant of centromere
evolution in closely related fungi. Proc Natl Acad Sci U S A 115:3108-3113.

49. Seppey M, Manni M, Zdobnov EM. 2019. BUSCO: Assessing genome assembly and
annotation completeness, p. 227-245. *In* Methods in Molecular Biology. Humana Press
Inc.

50. Schweizer G, Munch K, Mannhaupt G, Schirawski J, Kahmann R, Dutheil J. 2018.
Positively selected effector genes and their contribution to virulence in the smut fungus
Sporisorium reilianum. Genome Biol Evol 10:629–645.

51. Zuther K, Kahnt J, Utermark J, Imkampe J, Uhse S, Schirawski J. 2012. Host
specificity of *Sporisorium reilianum* is tightly linked to generation of the phytoalexin
luteolinidin by *Sorghum bicolor*. Mol Plant-Microbe Interact 25:1230–1237.

52. Zhang H, Meltzer P, Davis S. 2013. RCircos: an R package for Circos 2D track plots.
BMC Bioinformatics 14:244.

53. Hoff KJ, Lomsadze A, Borodovsky M, Stanke M. 2019. Whole-genome annotation
with BRAKER, p. 65–95. *In* Methods in Molecular Biology. Humana Press Inc.

54. Depotter JRL, Zuo W, Hansen M, Zhang B, Xu M, Doehlemann G. 2021. Effectors
with different gears: divergence of *Ustilago maydis* effector genes is associated with
their temporal expression pattern during plant infection. J Fungi 7:16.

55. Wang Q-M, Begerow D, Groenewald M, Liu X-Z, Theelen B, Bai F-Y, Boekhout T.
2015. Multigene phylogeny and taxonomic revision of yeasts and related fungi in the
Ustilaginomycotina. Stud Mycol 81:55–83.

56. Depotter JRL, Shi-Kunne X, Missonnier H, Liu T, Faino L, van den Berg GCM, Wood
TA, Zhang B, Jacques A, Seidl MF, Thomma BPHJ. 2019. Dynamic virulence-related
regions of the plant pathogenic fungus *Verticillium dahliae* display enhanced sequence
conservation. Mol Ecol 28:3482-3495.

57. Van de Wouw AP, Cozijnsen AJ, Hane JK, Brunner PC, McDonald BA, Oliver RP,
Howlett BJ. 2010. Evolution of linked avirulence effectors in *Leptosphaeria maculans*
is affected by genomic environment and exposure to resistance genes in host plants.
PLoS Pathog 6:e1001180.

58. Gorinšek B, Gubenšek F, Kordiš D. 2004. Evolutionary genomics of chromoviruses in
eukaryotes. Mol Biol Evol 21:781-798.

59. Wagner A. 2002. Selection and gene duplication: a view from the genome. Genome
Biol 3:reviews1012.1.

60. Lercher MJ, Hurst LD. 2002. Human SNP variability and mutation rate are higher in
regions of high recombination. Trends Genet 18:337.340.
61. Hwang DG, Green P. 2004. Bayesian Markov chain Monte Carlo sequence analysis
reveals varying neutral substitution patterns in mammalian evolution. Proc Natl Acad
Sci U S A 101:13994-14001.
62. Wicker T, Yu Y, Haberer G, Mayer KFX, Marri PR, Rounsley S, Chen M, Zuccolo A,
Panaud O, Wing RA, Roffler S. 2016. DNA transposon activity is associated with
increased mutation rates in genes of rice and other grasses. Nat Commun 7:12790.
63. McDonald MJ, Wang WC, Huang H Da, Leu JY. 2011. Clusters of nucleotide
substitutions and insertion/deletion mutations are associated with repeat sequences.
PLoS Biol 9:e1000622.
64. Lee GE, Mauro E, Parissi V, Shin CG, Lesbats P. 2019. Structural insights on
retroviral DNA integration: learning from foamy viruses. Viruses 21:249-256.
65. Lesbats P, Engelman AN, Cherepanov P. 2016. Retroviral DNA integration. Chem
Rev 116:12730-12757.
66. Badet T, Croll D. 2020. The rise and fall of genes: origins and functions of plant
pathogen pangenomes. Curr Opin Plant Biol 56:65-73.
67. Haas BJ, Kamoun S, Zody MC, Jiang RHY, Handsaker RE, Cano LM, Grabherr M,
Kodira CD, Raffaele S, Torto-Alalibo T, Bozkurt TO, Ah-Fong AM V, Alvarado L,
Anderson VL, Armstrong MR, Avrova A, Baxter L, Beynon J, Boevink PC, Bollmann
SR, Bos JIB, Bulone V, Cai G, Cakir C, Carrington JC, Chawner M, Conti L,
Costanzo S, Ewan R, Fahlgren N, Fischbach M a, Fugelstad J, Gilroy EM, Gnerre S,
Green PJ, Grenville-Briggs LJ, Griffith J, Grünwald NJ, Horn K, Horner NR, Hu C-H,
Huitema E, Jeong D-H, Jones AME, Jones JDG, Jones RW, Karlsson EK, Kunjeti SG,
Lamour K, Liu Z, Ma L, Maclean D, Chibucos MC, McDonald H, McWalters J,
Meijer HJG, Morgan W, Morris PF, Munro C a, O'Neill K, Ospina-Giraldo M, Pinzón
A, Pritchard L, Ramsahoye B, Ren Q, Restrepo S, Roy S, Sadanandom A, Savidor A,
Schornack S, Schwartz DC, Schumann UD, Schwessinger B, Seyer L, Sharpe T, Silvar
C, Song J, Studholme DJ, Sykes S, Thines M, van de Vondervoort PJI, Phuntumart V,
Wawra S, Weide R, Win J, Young C, Zhou S, Fry W, Meyers BC, van West P,
Ristaino J, Govers F, Birch PRJ, Whisson SC, Judelson HS, Nusbaum C. 2009.
Genome sequence and analysis of the Irish potato famine pathogen *Phytophthora
infestans*. Nature 461:393–398.
68. Torres DE, Oggenfuss U, Croll D, Seidl MF. 2020. Genome evolution in fungal plant
pathogens: looking beyond the two-speed genome model. Fungal Biol Rev 3:136-143.
69. Frantzeskakis L, Kusch S, Panstruga R. 2019. The need for speed :
compartmentalized genome evolution in filamentous phytopathogens. Mol Plant
Pathol 20:3–7.
70. Stam R, Münsterkötter M, Pophaly SD, Fokkens L, Sghyer H, Güldener U,
Hückelhoven R, Hess M. 2018. A new reference genome shows the one-speed genome
structure of the barley pathogen *Ramularia collo-cygni*. Genome Biol Evol 10:3243-
3249.
71. Frantzeskakis L, Kracher B, Kusch S, Yoshikawa-Maekawa M, Bauer S, Pedersen C,
Spanu PD, Maekawa T, Schulze-Lefert P, Panstruga R. 2018. Signatures of host

specialization and a recent transposable element burst in the dynamic one-speed
genome of the fungal barley powdery mildew pathogen. BMC Genomics 19:381.

72. Schwessinger B, Sperschneider J, Cuddy WS, Garnica DP, Miller ME, Taylor JM, Dodds PN, Figueroa M, Park RF, Rathjen P. 2018. A near-complete haplotype-phased genome of the dikaryotic. MBio 9:e02275-17.

73. Laurie JD, Linning R, Wong P, Bakkeren G. 2013. Do TE activity and counteracting genome defenses, RNAi and methylation, shape the sex lives of smut fungi? Plant Signal Behav 8:e23853.

74. Coelho MA, Bakkeren G, Sun S, Hood ME, Giraud T. 2018. Fungal Sex: The Basidiomycota, p. 147–175. *In* Heitman, J, Howlett, BJ, Crous, PW, Stukenbrock, EH, James, TY, Gow, NAR (eds.), The Fungal Kingdom. American Society of Microbiology, Washington DC.

75. Horváth V, Merenciano M, González J. 2017. Revisiting the relationship between transposable elements and the eukaryotic stress response. Trends Genet 33:832-841.

76. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114-2120.

77. Lam KK, Labutti K, Khalak A, Lam K, Tse D. 2015. FinisherSC: a repeat-aware tool for upgrading de-novo assembly using long reads. Bioinformatics 31:3207–3209.

78. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu□: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res 27:722–736.

79. Hu K, Xu K, Wen J, Yi B, Shen J, Ma C, Fu T, Ouyang Y, Tu J. 2019. Helitron distribution in Brassicaceae and whole genome Helitron density as a character for distinguishing plant species. BMC Bioinformatics 20:354.

80. Ellinghaus D, Kurtz S, Willhoeft U. 2008. LTRharvest, an efficient and flexible software for de novo detection of LTR retrotransposons. BMC Bioinformatics 9:18.

81. Crescente JM, Zavallo D, Helguera M, Vanzetti LS. 2018. MITE Tracker: an accurate approach to identify miniature inverted-repeat transposable elements in large genomes. BMC Bioinformatics 19:348.

82. Mao H, Wang H. 2017. SINE-scan: An efficient tool to discover short interspersed nuclear elements (SINEs) in large-scale genomic datasets. Bioinformatics 33:743-745.

83. Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in eukaryotic genomes. Mob DNA 6:11.

84. Li W, Godzik A. 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. Bioinformatics 22:1658-1659.

85. Hoede C, Arnoux S, Moisset M, Chaumier T, Inizan O, Jamilloux V, Quesneville H. 2014. PASTEC: An automatic transposable element classification tool. PLoS One 9:e91929.

86. Berthelier J, Casse N, Daccord N, Jamilloux V, Saint-Jean B, Carrier G. 2018. A transposable element annotation pipeline and expression analysis reveal potentially active elements in the microalga *Tisochrysis lutea*. BMC Genomics 19:378.

87. Ökmen B, Mathow D, Hof A, Lahrmann U, Aßmann D, Doehlemann G. 2018. Mining the effector repertoire of the biotrophic fungal pathogen *Ustilago hordei* during host and non-host infection. Mol Plant Pathol 19:2603–2622.

88. Grabherr MG., Brian J. Haas, Moran Yassour Joshua Z. Levin, Dawn A. Thompson,
Ido Amit, Xian Adiconis, Lin Fan, Raktima Raychowdhury, Qiandong Zeng, Zehua
Chen, Evan Mauceli, Nir Hacohen, Andreas Gnirke, Nicholas Rhind, Federica di
Palma, Bruce W. N, Friedman and AR. 2013. Trinity: reconstructing a full-length
transcriptome without a genome from RNA-Seq data. Nat Biotechnol 29:644–652.
89. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat
Methods 9:357–359.
90. Mascher M, Gundlach H, Himmelbach A, Beier S, Twardziok SO, Wicker T, Radchuk
V, Dockter C, Hedley PE, Russell J, Bayer M, Ramsay L, Liu H, Haberer G, Zhang
XQ, Zhang Q, Barrero RA, Li L, Taudien S, Groth M, Felder M, Hastie A, Šimková
H, Stanková H, Vrána J, Chan S, Munõz-Amatriaín M, Ounit R, Wanamaker S, Bolser
D, Colmsee C, Schmutzer T, Aliyeva-Schnorr L, Grasso S, Tanskanen J, Chailyan A,
Sampath D, Heavens D, Clissold L, Cao S, Chapman B, Dai F, Han Y, Li H, Li X, Lin
C, McCooke JK, Tan C, Wang P, Wang S, Yin S, Zhou G, Poland JA, Bellgard MI,
Borisjuk L, Houben A, Doleael J, Ayling S, Lonardi S, Kersey P, Langridge P,
Muehlbauer GJ, Clark MD, Caccamo M, Schulman AH, Mayer KFX, Platzer M, Close
TJ, Scholz U, Hansson M, Zhang G, Braumann I, Spannagl M, Li C, Waugh R, Stein
N. 2017. A chromosome conformation capture ordered sequence of the barley genome.
Nature 544:427-433.
91. Liao Y, Smyth GK, Shi W. 2019. The R package Rsubread is easier, faster, cheaper
and better for alignment and quantification of RNA sequencing reads. Nucleic Acids
Res 47:e47.
92. Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: A Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26:139-
140.
93. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. 2013. TopHat2 :
accurate alignment of transcriptomes in the presence of insertions , deletions and gene
fusions. Genome Biol 14:R36.
94. Sharma R, Mishra B, Runge F, Thines M. 2014. Gene loss rather than gene gain is
associated with a host jump from monocots to dicots in the smut fungus
Melanopsichium pennsylvanicum. Genome Biol Evol 6:2034–2049.
95. Lefebvre F, Joly DL, Labbe C, Teichmann B, Linning R, Belzile F, Bakkeren G,
Belanger RR. 2013. The transition from a phytopathogenic smut ancestor to an
anamorphic biocontrol agent deciphered by comparative whole-genome analysis. Plant
Cell 25:1946–1959.
96. Morita T, Koike H, Koyama Y, Hagiwara H, Ito E, Fukuoka T, Imura T, Machida M,
Kitamoto D. 2013. Genome sequence of the basidiomycetous yeast *Pseudozyma*
antarctica T-34 , a producer of the glycolipid biosurfactants. Genome Announc
1:e00064-13.
97. Juan J, Armenteros A, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S,
Heijne G Von, Nielsen H. 2019. SignalP 5.0 improves signal peptide predictions using
deep neural networks. Nat Biotechnol 37:420–423.
98. Krogh A, Larsson B, Von Heijne G, Sonnhammer ELL. 2001. Predicting
transmembrane protein topology with a hidden Markov model: application to complete

genomes. J Mol Biol 305:567-580.

99.   Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J,
1009   Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M,
1010   Yong SY, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein function
1011   classification. Bioinformatics 30:1236–1240.

1012   100.   Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate □: a practical and
1013   powerful approach to multiple testing. J R Stat Soc B 57:289–300.

1014   101.   Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. 2012. DbCAN: a web resource for
1015   automated carbohydrate-active enzyme annotation. Nucleic Acids Res 40:W445-451.

1016   102.   Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y.
1017   2018. DbCAN2: A meta server for automated carbohydrate-active enzyme annotation.
1018   Nucleic Acids Res 46:W95-W101.

1019   103.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version
1020   7: Improvements in performance and usability. Mol Biol Evol 30:772–780.

1021   104.   Stamatakis A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-
1022   analysis of large phylogenies. Bioinformatics 30:1312–1313.

1023   105.   Drillon G, Carbone A, Fischer G. 2014. SynChro: a fast and easy tool to reconstruct
1024   and visualize synteny blocks along eukaryotic chromosomes. PLoS One 9:e92621.

1025   106.   Drillon G, Carbone A, Fischer G. 2013. Combinatorics of chromosomal
1026   rearrangements based on synteny blocks and synteny packs. J Log Comput 23:815-
1027   838.

1028   107.   Eitzen K, Sengupta P, Kroll S, Kemen E, Doehlemann G. 2021. A fungal member of
1029   the *Arabidopsis thaliana* phyllosphere antagonizes *Albugo laibachii* via a GH25
1030   lysozyme. Elife 10:e65306.

1031   108.   Vakirlis N, Sarilar V, Drillon G, Fleiss A, Agier N, Meyniel JP, Blanpain L, Carbone
1032   A, Devillers H, Dubois K, Gillet-Markowska A, Graziani S, Huu-Vang N, Poirel M,
1033   Reisser C, Schott J, Schacherer J, Lafontaine I, Llorente B, Neuvéglise C, Fischer G.
1034   2016. Reconstruction of ancestral chromosome architecture and gene repertoire reveals
1035   principles of genome evolution in a model yeast genus. Genome Res 26:918-932.

1036   109.   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL.
1037   2004. Versatile and open software for comparing large genomes. Genome Biol 5:R12.

1038   110.   Quinlan AR, Hall IM. 2010. BEDTools: A flexible suite of utilities for comparing
1039   genomic features. Bioinformatics 26:841–842.

1040   111.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment
1041   search tool. J Mol Biol 215:403–410.

1042   112.   El-Gebali S, Mistry J, Bateman A, Eddy SR, Luciani A, Potter SC, Qureshi M,
1043   Richardson LJ, Salazar GA, Smart A, Sonnhammer ELL, Hirsh L, Paladin L, Piovesan
1044   D, Tosatto SCE, Finn RD. 2019. The Pfam protein families database in 2019. Nucleic
1045   Acids Res 47:D427-D432.

1046   113.   Monat C, Tando N, Tranchant-Dubreuil C, Sabot F. 2016. LTRclassifier: a website for
1047   fast structural LTR retrotransposons classification in plants. Mob Genet Elements
1048   6:e1241050.

1049   114.   Miele V, Penel S, Duret L. 2011. Ultra-fast sequence clustering from similarity
1050   networks with SiLiX. BMC Bioinformatics 12:116.

1051   115.   Pignatelli M, Serras F, Moya A, Guigó R, Corominas M. 2009. CROC: finding
1052   chromosomal clusters in eukaryotic genomes. Bioinformatics 25:1552–1553.
1053   116.   Heberle H, Meirelles VG, da Silva FR, Telles GP, Minghim R. 2015. InteractiVenn: a
1054   web-based tool for the analysis of sets through Venn diagrams. BMC Bioinformatics
1055   16:169.
1056   117.   Korber B. 2000. HIV signature and sequence variation analysis., p. 55–72. *In* Rodrigo,
1057   AG, Learn, GH (eds.), Computational analysis of HIV molecular sequences. Kluwer
1058   Academic Publishers, Dordrecht.
1059

1060

1061

Figure 1: Long terminal repeat retrotransposons (LTR-RTs) recently proliferated in *Ustilago hordei* and are generally transcriptionally inactive. (A) The chromosomes of *U. hordei* strain Uh805 are depicted. The numbering of the chromosomes corresponds to the numbering of the previously published *U. bromivora* assembly (40). The grey lines on the chromosome bars indicate LTR-RTs. The best homolog for every LTR-RT is connected with ribbons in the centre of the circos plot (52). The colour of the ribbon indicates the sequence identity between the homologous region of the two LTR-RTs. **(B)** Differential expression of gene and LTR-RT *U. hordei* loci between growth in liquid culture medium and *in planta*. The numbers between brackets indicate how many genes and LTR-RTs have been annotated in the genome. The significance of differential expression was calculated using *t*-tests relative to a threshold of log2-fold-change of 1 with Benjamini-Hochberg correction. A *p*-value cut-off of 0.05 was used.

The genome architecture of *Ustilago hordei*

Figure 2: The *Ustilago hordei* genome architecture. The chromosomes of *U. hordei* strain Uh805 are depicted. The numbering of the chromosomes corresponds to the numbering of the previously published *U. brachipodii-distachyi* assembly (40). Repeat content and percentage presence/absence polymorphisms were calculated for 20 kb windows. The presence of Uh805 genome regions in the five other *U. hordei* strains is shown (Table 1). The location of effector gene candidates, i.e. *in planta* upregulated genes encoding secreted proteins, is depicted. Also, the location of the Divergent Gene Clusters (DGCs) is plotted for the 20% most divergent orthologs with *Ustilago brachipodii-distachyi*.

Figure 3: The chromosome 1 architecture of *MAT-1* and *MAT-2 Ustilago hordei*. (A) As reference, the *MAT-1* strain Uh805 and the *MAT-2* strain Uh1278 were used. Repeat content and percentage presence/absence polymorphisms were calculated for 20 kb windows for the reference strains. The presence-absence was determined between the *MAT-1* and *MAT-2* reference strains, in addition to between the *MAT-1* strains Uh805 and Uh811. The *MAT*, *a* and *b* loci are indicated. The orange square includes the mating-type loci and indicates the genome region where the repeat content is high and the sequences are generally mating-type species. **(B)** Model that explains the mating-type specificity of the sequences within and flanking the mating-type loci. The absence of recombination within the mating-type loci and its flanking regions makes that transpositions within these regions are mating-type specific.

Figure 4: Genome evolution of the HNBT-cluster species. (**A**) Phylogenetic relationship between smut pathogens based on Benchmarking Universal Single-Copy Orthologs (BUSCOs). Phylogenetic relationship between newly and previously sequences smut species with the *Ustilago maydis/Sporisorium reilianum* branch was as an outgroup. In total, 1,667 BUSCOs were used for tree construction. For *U. hordei*, strain Uh805 was used in the tree. The robustness of the trees was assessed using 100 bootstrap replicates. The colours of the contigs indicate the synteny with the ancestor contigs. The blue section of the circles indicate the repeat fraction that is present in the genome assembles and within the mating-type loci for species with a bipolar mating-types (**B**) Long terminal repeat retrotransposon (LTR-RT) expansion in the HNBT-cluster. Nucleotide sequence identity distribution of best reciprocal paralogous and orthologous LTR-RT sequences. Squares on the lines display maxima with the corresponding sequence identity value. (**C**) Extent of clustering of most diverged orthologs between different species. *Ustilago hordei* and *Ustilago brachipodii-distachyi* were used as reference genomes to calculated the clustering of the x% most divergent (y-axis) HNBT-cluster orthologs. The degree of clustering is defined as the fraction of the x% most diverged genes that reside with at least 10 together in a window of 20 genes. Significant differences were determined for 20% of the most divergent orthologs with the Fisher exact test for *p*-value < 0.05 and indicated with a capital letter (A-E).

Figure 5: Interspecific long terminal repeat retrotransposon (LTR-RT) alterations in windows around genes within and outside Diverged Gene Clusters (DGCs). *Ustilago hordei* and *Ustilago brachipodii-distachyi* were used as reference genomes and DGCs were determined based on the 20% most diverged genes compared to orthologs from the other HNBT-cluster species. Absolute differences in LTR-RT content between orthologs were determined for a window 10 kb upstream of the 5'-end and 10kb downstream of the 3'-end of the genes. Significant differences were calculated with the Welch's t-test. *** : *p*-value < 0.001 and n.s. : no significant difference.

Figure 6: Long terminal repeat retrotransposon (LTR-RT) substitutions. (**A**) In total, 252 LTR-RTs are included in the phylogenetic tree and their species of origin is indicated by their colour name. LTR-RTs families are indicated by the circle segments in different colour. For every family, an ancestor sequence was reconstructed and indicated on the phylogenetic tree by a circle. The superfamily and the functional domains are indicated for the ancestor sequences. (**B**) The substitutions encountered by LTR-RT since their reconstructed ancestor. For transitions, differences between in substitution frequencies were tested with a t-test. *** : *p*-value < 0.001.

Figure 7: Properties of Diverged Gene Clusters (DGCs). Difference if frequencies of (**A|B**) substitution type, (**C**) synonymous and (**D**) nonsynonymous substitutions between genes inside and outside DGCs. (**E**) Difference in fraction of relative nonsynonymous and synonymous levels of genes within DGCs that encode and do not encode a secreted protein. DGCs were defined for *U. hordei* by using the 20% most diverged orthologs with *U. brachipodii-distachyi*. Outliers were not depicted for the boxplots. Significant differences were indicated with "*" for *p*-value < 0.05 and "***" for *p*-value < 0.001. Significant differences were calculated with the Welch's t-test.

Figure S1 Phylogenetic relationship between *Ustilago hordei* lineages based on Benchmarking Universal Single-Copy Orthologs (BUSCOs). Phylogenetic relationship between the *U. hordei* strains with *U. nuda* as an outgroup. In total, 1,692 BUSCOs were used for tree construction. The robustness of the trees was assessed using 100 bootstrap replicates.

Figure S2: Overlap Diverged Gene Cluster (DGC) genes using orthologs of different species. *Ustilago hordei* and *Ustilago brachipodii-distachyi* were used as reference genomes to determine the clustering of the 20% most diverged genes using orthologs of different species.

Figure S3: Gene density, presence/absence polymorphisms and long terminal repeat retrotransposons (LTR-RTs) of Diverged Gene Clusters (DGCs). DGCs were determined based on the 20% most diverged *Ustilago hordei* genes with *Ustilago brachipodii-distachyi* orthologs. The various features were calculated over 20kb windows. Outliers were not depicted for the boxplots. Significant differences were calculated with the t-test. * : *p*-value < 0.05 , *** : *p*-value < 0.001 and n.s. : no significant difference.