The genetic population structure of Lake Tanganyika's *Lates* species flock, an endemic radiation of pelagic top predators

Jessica A. Rick*[1], Julian Junker*[2,3], Ismael A. Kimirei[4], Emmanuel A. Sweke[4,5], Julieth B. Mosille[4], Christian Dinkel[2], Salome Mwaiko[2,3], Ole Seehausen[2,3], Catherine E. Wagner[1]

* these authors contributed equally to this work

[1] Department of Botany and Program in Ecology, University of Wyoming, 1000 E University Dr., Laramie, Wyoming 82072 USA

[2] EAWAG Swiss Federal Institute of Aquatic Science and Technology, CH-6047 Kastanienbaum, Switzerland

[3] Division of Aquatic Ecology & Evolution, Institute of Ecology & Evolution, University of Bern, CH-3012 Bern, Switzerland

[4] Tanzania Fisheries Research Institute (TAFIRI), Dar es Salaam, Tanzania

[5] Deep Sea Fishing Authority (DSFA), Zanzibar, Tanzania

Corresponding authors: Jessica Rick (jrick@uwyo.edu) and Catherine E. Wagner (Catherine.Wagner@uwyo.edu)

Running title: Population genetics of *Lates* fishes

Abstract

Understanding genetic connectivity plays a crucial role in species conservation decisions,
and genetic connectivity is an important component of modern fisheries management. In this
study, we investigated the population genetics of four endemic *Lates* species of Lake
Tanganyika (*Lates stappersii*, *L. microlepis*, *L. mariae* and *L. angustifrons*) using reduced-
representation genomic sequencing methods. We find the four species to be strongly
differentiated from one another (mean interspecific F_{ST} = 0.665), with no evidence for
contemporary admixture. We also find evidence for strong genetic structure within *L. mariae*,
with the majority of individuals from the most southern sampling site forming a genetic group
that is distinct from the individuals at other sampling sites. We find evidence for much weaker
structure within the other three species (*L. stappersii, L. microlepis*, and *L. angustifrons*). Our
ability to detect this weak structure despite small and unbalanced sample sizes and
imprecise geographic sampling locations suggests the possibility for further structure
undetected in our study. We call for further research into the origins of the genetic
differentiation in these four species—particularly that of *L. mariae*—which may be important
for conservation and management of this culturally and economically important clade of fish.

***Keywords*:** fishery management, genomic differentiation, Lates angustifrons, Lates mariae,
Lates microlepis, Lates stappersii

Introduction

Life history traits—including reproduction, growth, and dispersal—strongly influence gene flow and the genetic structure of populations (Ellegren & Galtier 2016; Manier & Arnold 2006; Stearns 1992). Differences in life history traits can also explain why sympatric and ecologically similar taxa may differ in their genetic population structure (Peterman *et al.* 2015; Young *et al.* 2015). Understanding how life history traits like dispersal, reproduction, and growth, shape gene flow is therefore crucial to improving our understanding of local adaptation, speciation, persistence of populations, migration, and the spatial scale of effective management (Sunday *et al.* 2014).

In teleost fishes, life history traits such as spawning location, philopatry, timing of spawning, and duration of larval stage, have been shown to shape population structure (Olsen *et al.* 2011; Pettersson *et al.* 2019; Young *et al.* 2015). Anadromous salmonids are well-known for displaying homing behavior that leads to genetic distinction between populations that use different spawning grounds at different times, dividing populations into many genetic units despite the populations spending most of their lives together in the ocean (Wenburg *et al.* 1998; Brannon *et al.* 2004). Divergence between populations caused by temporal differences in the timing of spawning has also been observed in Atlantic herrings (Martinez Barrio *et al.* 2016; Pettersson *et al.* 2019). In the oceans, coral reef fishes which spawn on or near the substrate, which exhibit parental care strategies, and which lack planktonic larvae, generally have significantly greater population structure than pelagic spawners or substrate spawners with planktonic larvae (Riginos *et al.* 2014). Additionally, a shorter larval phase among pelagic spawners can lead to a more structured population, because this leads to a shorter larval dispersal distance (Selkoe & Toonen 2011).

Understanding the genetic population structure of species is essential for conservation and sustainable fisheries management (Reiss *et al.* 2009; Bernatchez 2016; Supple & Shapiro 2018). Disregarding population structure and managing a population complex as a single population can lead to overfishing or even extinction of the more vulnerable genetic groups (Reiss *et al.* 2009; Sterner 2007). In species with cryptic diversity, unintentional overfishing of distinct populations can lead to a loss of genetic diversity in the species as a whole. Stock diversity can contribute to the resilience of a fishery in the face of environmental fluctuations (Schindler *et al.* 2015) and the loss of differentiated populations can result in major losses in the consistency of fisheries yields (Hutchinson 2008; Schindler *et al.* 2010). The loss of population diversity can also have dramatic effects on ecosystem services and species persistence; understanding the extent to which population structure exists is crucial for predicting these consequences (Schindler *et al.* 2015; Schindler *et al.* 2010; Therkildsen *et al.* 2013). It is therefore essential to understand the extent and nature of population genetic

structure and differential adaptation in fishes that are important food and economic
resources.

The pelagic fish community and the fishery of Lake Tanganyika

Lake Tanganyika is the second largest inland fishery on the continent of Africa (Van der
Knaap *et al.* 2014). During the last few decades, harvest records from Lake Tanganyika's
fishery have indicated a general decline in population sizes (Kimirei *et al.* 2008; Van der
Knaap 2013; Van der Knaap *et al.* 2014; van Zwieten *et al.* 2002). This decline has resulted
from a combination of an increase in the number of fishermen and vessels on Lake
Tanganyika, changes in fishing practices (e.g., the use of beach seining, which targets fish in
their in-shore nursery habitats) (Kimirei *et al.* 2008; Petit & Shipton 2012; Van der Knaap
2013; Van der Knaap *et al.* 2014; van Zwieten *et al.* 2002), and warming lake surface
temperatures (Cohen *et al.* 2016; O'Reilly *et al.* 2003; Sayers *et al.* 2020). Decreases in fish
abundance are likely also linked to reduced productivity in the lake caused by stronger water
column stratification due to climate change (O'Reilly *et al.* 2003; Verburg *et al.* 2003). In
addition, anthropogenic activity (e.g., land use changes, pollution, and increased
sedimentation) impact both the water quality of Lake Tanganyika directly and the lake's
productivity via climate feedbacks (Ogutu-Ohwayo *et al.* 2016; Ivory *et al.* 2021; McGlue *et
al.* 2021). Local fishermen similarly acknowledge spatial and temporal changes in fish
abundance that are related to the ecological conditions of the lake (Bulengela *et al.* 2020; De
Keyzer *et al.* 2020). Consequently, there is increasing recognition of the need to develop
sustainable management strategies for the lake's pelagic fish stocks (Kimirei *et al.* 2008;
Mölsä *et al.* 1999; Mölsä *et al.* 2002; Van der Knaap 2013; Van der Knaap *et al.* 2014; van
Zwieten *et al.* 2002).

Lake Tanganyika's pelagic zone is relatively low in species diversity compared to that of
other large African lakes: it consists predominantly of six endemic fish species belonging to
two families (Coulter 1991). These include a monophyletic pair of mainly planktivorous
clupeids (*Stolothrissa tanganicae*, *Limnothrissa miodon*) and four mainly piscivorous latids of
the genus *Lates* (*L. stappersii*, *L. mariae*, *L. microlepis*, *L. angustifrons*). While all four *Lates*
species likely have planktivorous larvae (van Zwieten *et al.* 2016), only *L. stappersii* remains
pelagic throughout its life (Ellis 1978). The other three species move inshore to littoral weed
beds as juveniles (>3cm) before *L. microlepis* returns to the pelagic, *L. mariae* transitions to
the deepwater benthic zone, and *L. angustifrons* can be found throughout the water column
as adults (Coulter 1991). Of the four *Lates* species, *L. stappersii* is the most abundant and is
an important component of the current fishery of Lake Tanganyika (Coulter 1976, 1991;
Kimirei *et al.* 2008; Mannini 1998a, b; Mölsä *et al.* 1999; Mölsä *et al.* 2002; Munyandero

2002; Sarvala *et al.* 2002; Van der Knaap 2013; Van der Knaap *et al.* 2014; van Zwieten *et*
al. 2002).

The fish of Lake Tanganyika are influenced by a complex interplay of bottom-up control via
different mixing regimes and upwelling rates (Bergamino *et al.* 2010; O'Reilly *et al.* 2003;
Stenuite *et al.* 2007; Verburg & Hecky 2009; Verburg *et al.* 2003), and top-down control via
predators and fishing (Kimirei *et al.* 2008; Mannini 1998a; Mannini *et al.* 1996; Munyandero
2002; Van der Knaap 2013; Van der Knaap *et al.* 2014; van Zwieten *et al.* 2002). If spatially
heterogeneous environmental factors, such as mixing regimes and nutrient supply, are
consistent over generations and coupled with limited gene flow between spatially segregated
populations, divergent selection on life history traits (e.g., spawning phenology,
developmental timing and recruitment success) of fish in different parts of the lake could
develop and persist. Spatial environmental variation in Lake Tanganyika, in combination with
its large geographic extent, might therefore generate intraspecific genetic differentiation
among populations of pelagic fish, as has been observed in pelagic cichlids in both Lake
Tanganyika and Lake Malawi (Genner *et al.* 2010; Koblmüller *et al.* 2019). In these cichlids,
strong spawning site fidelity (Genner *et al.* 2010) and differences in foraging behavior
(Koblmüller *et al.* 2019) influence spatial genetic structure. In contrast, recent studies have
found that the two pelagic clupeids in Lake Tanganyika show no evidence of spatial genetic
structure (Junker *et al.* 2020; De Keyzer *et al.* 2019; Kmentová *et al.* 2020). Intraspecific
spatial heterogeneity in spawning times (Coulter 1976; Ellis 1978), dominant diet items
(Mannini *et al.* 1999), sizes at maturity (Mannini et al. 1996), and catch rates (Coulter 1991),
along the length of Lake Tanganyika support the possibility that gene flow may be reduced
among fish in different parts of the lake. Indeed, in *L. stappersii*, a study based on RAPD
markers found individuals from the Kigoma region to be genetically differentiated from the
rest of the sites sampled in this species (Kuusipalo 1999), with some additional differentiation
observed among individuals collected during the dry versus rainy.

Here, we use reduced-representation genomic data to test the prediction that differences in
life history traits among the four Lake Tanganyika-endemic *Lates* species result in
differences among the species in their intraspecific genetic connectivity and population
structure. We first assessed whether phenotypic identification coincides with genetically
based species boundaries and then examined the extent to which each of the four species is
genetically structured, as well as how the genetic structure observed in each species
corresponds to its known life history and the environment. Based on known differences in life
history among the four species, we hypothesized that we would find little genetic population
structure in the entirely pelagic *L. stappersii* and mainly pelagic *L. microlepis*. In contrast, we
expected the highest genetic differentiation in populations of the predominantly benthic *L.*
mariae. Finally, we hypothesized that *L. angustifrons* would show intermediate amounts of

structure since there is no evidence for large scale movements of this species, even though
they are found throughout the water column.

Material and Methods

Study system and sampling

Between 2001 and 2019, we collected tissue samples (fin clips) from *L. stappersii, L. mariae,*
L. microlepis, and *L. angustifrons* at 9 general locations along the Tanzanian shore of Lake
Tanganyika, which together span the ~490 km shoreline from the northern to southern border
(Fig. 1; Table S2). All fish collections were made in partnership with researchers of the
Tanzanian Fisheries Research Institute (TAFIRI) in Kigoma, Tanzania; most of the fish were
obtained opportunistically from fishermen, and therefore were already dead when we
received them. Given fuel costs and limitations of most outboard motors used by fishermen
in this region, fishermen generally fish within a 20km radius of their landing sites (personal
observation), but there remains inherent uncertainty in the exact location and depth at which
these fish were collected. Fish were identified in the field by sampling personnel following
United Nations Food and Agricultural Organization (FAO) identification guidelines (Eccles
1992) and guidance from Tanzanian researchers and local fishermen, and genetic methods
were later used to confirm species identities for each fish. The majority of fish were collected
during two targeted sampling campaigns in 2017 and 2018, the first of which occurred at the
end of the dry season (September-October 2017), while the second occurred just following
the rainy season (April-May 2018). For fish collected from 2016-2019, we recorded length
and weight, and took standardized pictures of each fish. For the few live fish that were
caught as part of other research, we took cuvette photographs of the live fish and
subsequently euthanized the fish with an overdose of MS222. We then took fin clips for
genetic analysis from all fish. Specimens smaller than 600mm were preserved in
formaldehyde and archived in the collections at TAFIRI (Kigoma, Tanzania), EAWAG
(Kastanienbaum, Switzerland), the University of Wyoming Museum of Vertebrates (Laramie,
WY, USA), or the Cornell University Museum of Vertebrates (Ithaca, NY, USA).

Genomic sequencing

We extracted DNA from fin clips using DNeasy Blood and Tissue Kits (Qiagen, Inc.) following
the standard protocol, with the addition of an RNAse A incubation step. We then prepared
genomic libraries for a combination of genotyping-by-sequencing (GBS, for *L. stappersii, L.*
mariae, L. microlepis, L. angustifrons) and restriction site associated DNA sequencing (RAD,
for *L. stappersii*).

We prepared GBS libraries following protocols outlined in Parchman *et al.* (2012), using MseI
and EcoRI restriction enzymes. Following fragmentation via restriction enzyme digestion,
fragmented DNA was barcoded by ligating short, unique individual-identifying DNA fragments
(barcodes) to each individual's fragmented DNA. These barcoded fragments were then
amplified by PCR and pooled for sequencing. We ran two replicate PCRs for each individual
and pooled the final PCR products into two libraries. The prepared libraries were size-
selected for 200-350bp fragments using Blue Pippin (Sage Science, MA). One library was
sequenced on one lane of Illumina HiSeq4000 (150bp single-end) at the University of Texas
at Austin's Genome Sequencing and Analysis Facility (UT GSAF; Austin, TX) and the other
library was sequenced on one lane of Illumina HiSeq4000 (150bp, single-end) at the
University of Oregon's Genomics and Cell Characterization Core Facility (Eugene, OR). Each
library contained 192 individuals, and select individuals were duplicated across libraries to
check for library compatibility.

The RAD libraries containing *L. stappersii* were prepared for sequencing following the
protocol by Baird *et al.* (2008) with the following modifications: we used between 400ng and
1000ng genomic DNA per sample and digested with *SbfI* overnight. We tagged each
individual using P1 adapters (synthesized by Microsynth) with custom six to eight base pair
barcodes and multiplexed 67 barcoded individuals per library, resulting in a total of three
RAD libraries. The libraries were sheared using an S220 series Adaptive Focused Acoustic
(AFA) ultra-sonicator (Covaris) with the manufacturer's settings for a 400 bp mean fragment
size. We size-selected for fragments between 300 and 700bp using a sageElf (Sage
Scientific Electrophoretic Lateral Fractionator; Sage Science, Beverly, MA). The enrichment
step was done in 6 aliquots with a total volume of 200 µl. Volumes were combined prior to
the final size selection (again 300-700bp) step using the sageELF. Sequencing was done by
the Lausanne Genomic Technologies sequencing facilities (University of Lausanne,
Switzerland). All three RAD libraries were single-end sequenced on one lane each of the
Illumina HiSeq2000 (100bp SE).

Sequence data preparation

We filtered raw sequencing reads from each library by first removing the reads derived from
the PhiX genome and other common contaminants (e.g. excess barcodes, primers, and
adapters) using bowtie2 (v2.0.0, Langmead & Salzberg 2012). We filtered reads for an intact
enzyme restriction site (*SbfI* for RAD libraries, MseI and EcoRI for GBS libraries), de-
multiplexed the fastq files, and matched sequence barcodes to individual fish using custom
perl and bash scripts.

For the GBS libraries, the reads were mapped to the chromosome-level *Lates calcarifer*
reference genome (v3, NCBI Genome Assembly GCA_001640805.1; Vij *et al.* 2016) using
bwa mem (v0.7.17; Li & Durbin 2009) with default settings. Following alignment, we excluded
any individual with < 50,000 reads or less than 60% of raw reads assembled to the reference
genome. We identified variable sites (i.e. single nucleotide polymorphisms; SNPs) in the
assembly using SAMtools mpileup (v1.8; Li *et al.* 2009) and bcftools (v1.8; Li *et al.* 2009). In
calling variable sites, we omitted indels and kept only high-quality biallelic variant sites
(QUAL > 20 and GQ > 9). We then filtered SNPs by minor allele frequency and amount of
missing data using vcftools (Danecek *et al.* 2011), additionally only calling genotypes with a
minimum read depth of 5. We then selected only SNPs with a minor allele frequency greater
than 0.01 that were present in at least 50% of individuals. Finally, we thinned each data set
to retain one SNP per locus (-thin 90). We assessed the distribution of missing data across
individuals and mean read depth across sites using vcftools (Danecek *et al.* 2011),
additionally using whoa (v0.0.2.999, Anderson 2018) in R (v4.0.4, R Core Team 2021) to
assess heterozygote miscall rate at low read depth sites to ensure that our read depth filter
was stringent enough.

For the RAD libraries, we barcode-trimmed reads down to 84 nucleotides using
process_radtags from Stacks (Catchen *et al.* 2013). The FASTX-toolkit v.0.0.13
(http://hannonlab.cshl.edu/fastx_toolkit/) was used for quality filtering. In a first step, we kept
only reads with all base quality scores greater than 10; in a second step, we removed all
reads with more than 5% of the bases with quality score below 30. The RAD library was then
mapped to the *Lates calcarifer* reference genome (Vij et al. 2016) and filtered, following the
same steps as for the GBS libraries above (MAF > 0.01, missing data < 0.5, one SNP per
locus, read depth > 5). There was no overlap of individuals included in the RAD and GBS
sequencing libraries.

After filtering variants, we calculated the distribution of minor allele reads at heterozygous
sites for each individual in both the RAD and GBS libraries, using a custom bash script
(available at http://github.com/jessicarick/lates-popgen). If there is an excess of sequencing
errors or contamination for an individual, then we would expect the ratio of minor allele to
major allele reads to be significantly less than 1:1. Thus, we plotted the distribution of minor
allele reads at heterozygous sites for each individual and removed individuals with a
deficiency in minor allele reads across the majority of heterozygous sites. After removing
these individuals, we then re-called genotypes and re-filtered the SNP data set to produce
our final SNP data set.

Species assignment

Not all samples could be identified to species in the field. Some fish were caught in deep
water and brought quickly to the surface by the fishermen, making them difficult to identify
due to barotrauma. Furthermore, juveniles of the large species *L. microlepis*, *L. mariae* and
L. angustifrons can be difficult to distinguish. We used three different lines of evidence to
confirm species identification and ensure that morphological identification matched genetic
assignment: (1) phenotypic identification based on FAO species descriptions; (2) individual
ancestry assignment using a model-based clustering approach; and (3) inference of
monophyletic groups in phylogenetic analysis. The details for the latter two of these steps are
described below.

With the GBS data, we first visualized clusters in our data using principal component analysis
using EMU (Meisner *et al.* 2021). EMU uses an iterative method to infer population structure
in the presence of missing data and is therefore able to infer population structure even for
datasets with high, uneven, or non-random proportions of missing data, which tend to
produce bias in other PCA methods (Meisner *et al.* 2021). We used the EM acceleration
method in EMU, using 10 eigenvectors for optimization (-e 10) with a maximum of 100
optimization iterations and keeping all eigenvectors as output. We then inferred species
groups using the model-based genetic clustering program entropy (Gompert *et al.* 2014;
Shastry *et al.* 2021). Entropy infers population structure from multilocus SNP data by
assigning individuals partially or completely to groups using Bayesian estimation from
genotype likelihoods. In taking uncertainty about individual genotypes into account via
genotype likelihoods, the model integrates outcomes over genotype uncertainty. We
converted our VCF containing sites with <50% missing data to the mpgl genotype likelihood
format using the vcf2mpgl R script (available at http://github.com/jessicarick/lates-popgen;
adapted from https://bitbucket.org/buerklelab/mixedploidy-entropy/src, v2.0). We then ran
entropy for K=4 clusters, running three independent MCMC chains of 80,000 total steps,
discarding the first 10,000 steps as burn-in, and retaining every 10^{th} value (thin=10), resulting
in 7000 samples from the posterior distribution of each chain. Our MCMC chains were run in
parallel using GNU parallel (Tange 2011). We checked MCMC chains for mixing and
convergence of parameter estimates by plotting a trace of the MCMC steps. We then
visualized assignments in R and assigned individuals to species based on group assignment
probabilities (q).

We further corroborated species identities using maximum likelihood phylogenetic inference
methods in RAxML (v8.1.17; Stamatakis 2014). We concatenated all SNPs, removed
invariant sites using the raxml_ascbias python script (v1.0, from
https://github.com/btmartin721/raxml_ascbias), and used the Lewis correction for invariant
sites with the ASC_GTRGAMMA model of molecular evolution (Lewis 2001) within RAxML to

infer the maximum likelihood phylogeny, including *L. calcarifer* (Vij et al. 2016) as the

outgroup. We used 100 rapid bootstraps to estimate confidence in our maximum likelihood

tree. We then visually identified monophyletic groups on our maximum likelihood tree and

used these groupings to further verify species identities from entropy and PCA clusters.

Genetic diversity

We calculated genetic diversity for each species in ANGSD (v0.931, Korneliussen *et al.*

2014). From the BAM alignment files, we first calculated the site allele frequency likelihoods

based on individual genotype likelihoods (option -doSaf 1) using the samtools model (option -

GL 1), with major and minor alleles inferred from genotype likelihoods (option -doMajorMinor

1) and allele frequencies estimated according to the major allele (option -doMaf 2). We

filtered sites for a minimum read depth of 10, minimum mapping quality of 20, and minimum

quality (*q*-score) of 20. From the site allele frequency spectra, we then calculated the

maximum-likelihood estimate of the folded site frequency spectra (SFS) using the ANGSD

realSFS program (with option -fold 1). The folded SFSs were then used to calculate genome-

wide genetic diversity (Watterson's θ), using the ANGSD thetaStat program (Korneliussen *et*

al. 2013).

Population genetic structure and isolation-by-distance

For each species, we then re-called variants to create a species-specific set of SNPs from

the GBS data. As with the species combined GBS data, we identified variable sites using

SAMtools mpileup and bcftools, omitting indels and keeping only high-quality biallelic variant

sites (QUAL > 20 and GQ > 9). We then filtered SNPs by minor allele frequency (> 0.01) and

amount of missing data (<50% missing) using vcftools (Danecek et al. 2011), additionally

only calling genotypes with a minimum read depth of 5. Finally, we thinned to one SNP per

locus (-thin 90).

For each species, we then performed principal component analysis (PCA) using EMU

(Meisner *et al.* 2021), after using PLINK (v1.90b6.9, Purcell *et al.* 2007; Chang *et al.* 2015) to

convert our VCF into BED format. EMU is able to infer population structure even for datasets

with high, uneven, or non-random rates of missing data, which tend to produce bias in other

PCA methods (Meisner *et al.* 2021). We again used the EM acceleration method, using 10

eigenvectors for optimization (-e 10) with a maximum of 100 optimization iterations and

keeping all eigenvectors as output. We also performed PCA in EMU on the RAD dataset for

L. stappersii. Individuals that had been duplicated among libraries were kept separate

through these initial analyses, and then their reads were combined and variants were re-

called for each species after ensuring that the duplicates clustered near to one another in
both species identification and intraspecific PCA analyses.

We estimated heterozygosity at each site using the dartR package in R (v1.9.9, Gruber *et al.*
2018). We then used the Reich-Patterson F_{ST} estimator (Reich *et al.* 2009), which is robust
even with small sample sizes, to calculate genetic divergence between sampling sites within
each species. We calculated Reich-Patterson F_{ST} estimates using a custom function in R
(available at http://github.com/jessicarick/reich-fst), using functions from the vcfR (v1.12,
Knaus & Grunwald 2017) and dartR packages for importing and manipulating our VCF. We
then tested for isolation-by-distance (IBD) using a Mantel test (mantel.randtest from ade4 in
R; v1.7-16, Dray & Dufour 2007) to calculate the correlation between the natural logarithm of
Euclidean geographic distance and standardized genetic distance ($F_{ST}/1-F_{ST}$) for all pairwise
combinations of sampling sites within each species. We chose to pool these tests by
sampling site, as we do not have more specific locations for where most of the fish were
caught.

To test for intraspecific population subdivision more formally, we again used entropy. For
each species, we ran entropy for K=1 to K=6 to infer the most likely number of ancestral
groups within each species, and to assign individuals to these clusters. We ran three
independent MCMC chains of 100,000 total steps, discarding the first 10,000 steps as burn-
in, and retaining every 10^{th} value (thin=10), resulting in 9000 samples from the posterior
distribution of each chain. We checked MCMC chains for mixing and convergence of
parameter estimates by plotting a trace of the MCMC steps. We then calculated the deviance
information criterion (DIC) for each value of K and used the model with the lowest DIC as our
best for modeling the variation observed in our data. From our entropy results, we assigned
individuals to intraspecific groups using a cutoff of q > 0.6 at each value of K. We then
calculated the mean of Reich-Patterson F_{ST} estimates between these K groups to understand
how genetically distinct these groups are from one another. We also ran entropy at K=2 and
K=3 using datasets with a more stringent missing data filter (miss=0.9) to test whether group
assignments were contingent on the choice of bioinformatic filters. We also calculated Reich-
Patterson F_{ST} estimates between each pair of individuals within each species in R, to
facilitate comparisons that do not rely on group assignments. With the group assignments
and individual pairwise F_{ST} estimates, we statistically tested and graphically visualized
relationships between these genetic measures and sampling year, sampling time of year, fish
standard length, and juvenile status, to examine hypotheses about heterogeneity in
spawning location and timing.

Results

The GBS libraries yielded an average of 245 million reads across 200 individuals. The RAD libraries containing the *L. stappersii* individuals yielded an average of 271 million reads, including 2.5% - 8.6% bacteriophage PhiX genomic DNA. On average, the mapping rate for *L. stappersii* individuals' GBS reads to the *L. calcalifer* reference genome was 89.8% (RAD, 89.6%), and mapping rates averaged 91.6% for *L. microlepis*, 91.3% for *L. mariae* and 88.7% for *L. angustifrons* individuals.

These mapping rates resulted in 1,463,702 unfiltered variable sites in *L. stappersii*, 1,286,810 in *L. microlepis*, 961,153 in *L. mariae* and 764,456 in *L. angustifrons*. After filtering for missing data (< 50%), minor allele frequency (MAF > 0.01), read depth (minDP > 5), and keeping only one SNP per locus (-thin 90), our species-specific GBS data sets contained 38,160 SNPs from 58 *L. stappersii* individuals (RAD, 16,802 SNPs from 63 different individuals), 33,429 SNPs from 34 *L. microlepis* samples, 30,431 SNPs from 38 *L. mariae* samples and 19,519 SNPs from 20 *L. angustifrons* samples. The data set with all four species contained 4,999,192 unfiltered and 44,823 filtered SNPs. None of the four species exhibited heterozygote excess that would be expected if many sites were erroneously called as heterozygous, and estimates of miscall rates are low for all read depths > 2 (Fig. S1). The majority of individuals who were lost were filtered out due to an imbalance in the reads supporting major and minor alleles at heterozygous loci. In our final dataset, the mean read depth was 15.2 for *L. stappersii*, 23.1 for *L. microlepis*, 20.9 for *L. mariae*, and 25.8 for *L. angustifrons* (Fig. S2).

Species assignment

The species-combined SNP dataset for GBS data contained 200 individuals, of which we removed 50 individuals due to low numbers of reads (n=2), low assembly rate to the reference genome (n=4), or highly unbalanced read ratios at heterozygous sites (n=44), which is indicative of contamination. The remaining individuals all had > 50,000 reads that aligned to the *L. calcarifer* reference genome. The SNP alignment for RAxML contained 48,776 SNPs after removing invariant sites. The rooted maximum likelihood phylogeny (rooted on *L. calcarifer*; Fig. 2A) shows that each of the four species is reciprocally monophyletic. In addition, we infer a sister taxon relationship between *L. mariae* and *L. angustifrons,* and a sister taxon relationship between *L. stappersii* and *L. microlepis* (conflicting with the best supported topology in Koblmüller *et al.* 2021); however, short internode distances between these splits and low bootstrap support for the initial split suggest that all three speciation events may have occurred within a short period, and more investigation of the divergence history of the clade is warranted but not the focus of this

current work. The PCA based on the GBS data was consistent with the species affinities

indicated by monophyletic groups in RAxML (Fig. 2). The first genetic PC axis (27.3%)

separates *L. stappersii* from the remaining three species. The second PC axis (12.4%) then

separates *L. mariae* and *L. microlepis* from one another, and the third axis (4.8%) separates

L. angustifrons from the other three species. Using entropy at K=4, we found each of the four

species to make a distinct genetic cluster (Fig. 2C). Based on these combined analyses, 58

individuals were assigned to *L. stappersii*, 34 to *L. microlepis*, 38 to *L. mariae*, and 20 to *L.*

angustifrons. While the genetic approaches revealed consistent species identifications, we

had phenotypically misidentified one *L. microlepis*, one *L. mariae* and three *L. angustifrons*

(Table S1).

Population structure

All four species had low but similar levels of genetic diversity (overall θ_W = 0.00222), and

genetic diversity was elevated in *L. stappersii* (θ_W = 0.00230) compared to the three other

species (*L. microlepis*, θ_W = 0.00094; *L. mariae*, θ_W = 0.00092; *L. anguistifrons*, θ_W = 0.00064;

Fig. 2A). Genetic diversity for the RAD dataset in *L. stappersii* was similar to that estimated

for the GBS data (RAD θ_W = 0.00223). Reflecting the distinctiveness seen in the all-species

PCA and phylogeny, F_{ST} values are high between species (mean Reich-Patterson F_{ST} =

0.607; Table S3). None of the four species shows a significant relationship between genetic

and geographic distance, suggesting an overall lack of isolation by distance (Fig. S3; all

Mantel test *p*-values > 0.2).

The PCA for *L. stappersii* based on GBS data (Fig. 3A) did not reveal any clear genetic

clusters. We similarly did not see any clustering in the RAD data for *L. stappersii* (Fig. S4). In

entropy, K=1 was the best fit for the GBS data according to DIC (Fig. 5A); however, at K=2,

some individuals from Kigoma, North Mahale and South Mahale form a distinct genetic

cluster, and at K=3, groups appear that co-occur at all sampling sites. These groups at K=2

and K=3 were largely but not completely stable (i.e., the same individuals were clustered

together) at a higher threshold for missing data (miss=0.5 vs. miss=0.9 had 18 individuals

switching groups at K=2 and 10 individuals at K=3; Fig. S7, S8). This weak genetic structure

(between-cluster K=2 Reich-Patterson F_{ST} = 0.00205, 95% bootstrap CI: 0.00084-0.00346,

Table S4) was also partly reflected in F_{ST} values between sampling sites, where F_{ST} values of

comparisons between individuals of South Mahale or North Mahale with samples from other

sites were elevated, albeit small ($\leq$ 0.012, Fig. 4A); however, only the F_{ST} values between

North Mahale and each of Mpinbwe and Kagunga had 95% bootstrap confidence intervals

not overlapping zero (Fig. S9A). The RAD data showed similar weak structure at K=2 (Fig.

S5; between-cluster F_{ST} = 0.00146, 95% bootstrap CI: 0.00087-0.00207), but these groups

were found in sympatry at all five sampling sites, similar to the groups at K=3 for the GBS
data. At K=3, F_{ST} values between groups for the RAD data had bootstrap confidence
intervals overlapping zero (Table S3).

The PCA for *L. microlepis* did not reveal clear genetic clusters but it did highlight several
individuals from the sampling sites Kigoma, North Mahale, and South Mahale, that were
differentiated from the other samples on PC1 and PC2 (Fig. 3B). These differentiated
individuals were not those with high amounts of missing data (Fig. S10B). All of the sampling
sites had low differentiation (mean F_{ST} = 0.00471; Fig. 4B) but all pairwise comparisons had
95% confidence intervals that did not overlap zero except for the Kirando-Kigoma and
Kirando-South Mahale comparisons (Fig. S9B). In entropy, K=1 was found to be the best fit
using DIC (Fig. 5B); at K=2, four individuals collected in Kigoma are differentiated from the
others, albeit with low genetic differentiation and a between-group F_{ST} estimate that overlaps
zero (Reich-Patterson F_{ST} = 0.00074, 95% CI: -0.00389 - 0.00549; Table S4). At K=3, the
additional group is found at all four sampling sites; the differentiation between the two largest
groups is low, but significant (Reich-Patterson F_{ST} = 0.0030, 95% CI: 0.00155 - 0.00474),
while the F_{ST} estimates between each of these groups and the smaller group identified at
K=2 had 95% bootstrap confidence intervals overlapping zero. These groups at K=2 and K=3
were largely stable at a higher threshold for missing data (miss=0.5 vs. miss=0.9 had 4
individuals switching groups at K=2 and 8 individuals at K=3; Fig. S7, S8).

The PCA for *L. mariae* clusters three individuals from Kasanga separate from all other
individuals on the first PC axis (explaining 5.28% of the variance), while the second PC axis
separates out some individuals from Kigoma, Wampembe, and Kirando from the others (Fig.
3C). The differentiation among individuals on both PC1 and PC2 does not correlate with the
individuals' amount of missing data (Fig. S10C). In entropy, K=2 has the lowest DIC
estimate, where the three individuals from Kasanga form a distinct cluster (Reich-Patterson
F_{ST} between entropy-identified groups = 0.119, 95% bootstrap CI: 0.113-0.127; Fig 4C and
Table S4). At K=3, most of the other sampling sites have individuals assigned to both of the
non-Kasanga clusters, although these two non-Kasanga clusters have very low
differentiation (F_{ST} = 0.00359, 95% bootstrap CI: 0.00180-0.00623) compared to that
between the Kasanga cluster and each of the other two clusters (mean F_{ST} = 0.120). These
groups at K=2 and K=3 were largely stable at a higher threshold for missing data (miss=0.5
vs. miss=0.9 had no individuals switching groups at K=2 and 15 individuals at K=3; Fig. S7,
S8). The F_{ST} estimates between all fish sampled from Kasanga and other sampling sites are
all significantly different from zero (Fig. S9C). All other between-site F_{ST} estimates had 95%
bootstrap confidence intervals overlapping zero.

For *L. angustifrons*, the PCA did not reveal any genetic structure (Fig. 3D) and the clustering
model with K=1 is the best fit for our data (Fig. 5D). Clusters identified at higher levels of K
had low differentiation (Table S4; K=2, F_{ST} = 0.00498; K=3, mean F_{ST} = 0.0228), and were
found co-occurring at most sampling sites (Fig. 5D). These groups at K=2 and K=3 were
largely but not completely stable at a higher threshold for missing data (miss=0.5 vs.
miss=0.9 had 4 individuals switching groups at K=2 and 7 individuals at K=3; Fig. S7, S8). In
addition, only three pairwise F_{ST} estimates between sampling sites were significant:
Kagunga-North Mahale, Kigoma-Kirando, and North Mahale-Wampembe (Fig. S9D);
however, even these sites had low differentiation (mean F_{ST} between sites = 0.0132), and it
should be noted that North Mahale only has one fish sampled, so these estimates may not
be indicative of patterns at the site as a whole.

We find no relationship between size difference and extent of differentiation (pairwise F_{ST} or
genetic cluster membership) in *L. mariae, L. microlepis,* or *L. angustifrons*. In *L. stappersii*, a
weak positive relationship exists between the difference in standard length and the extent of
genetic differentiation (F_{ST}) between individuals, such that individuals that are more different
in size are also more differentiated genetically (Fig. S12). In *L. microlepis*, we found no
relationship between genetic cluster membership and juvenile status at K=2 or K=3 when
examining only the fish collected in Kigoma, the only site with large enough numbers for
doing this analysis. We additionally find no systematic difference between F_{ST} estimates
between only juvenile fish and those estimated between only adult fish for *L. microlepis* or *L.*
mariae (Fig. S13, S14). When only looking at fish collected at the same site (Fig. S14), we
find that juvenile-adult comparisons had higher F_{ST} values in *L. mariae* than adult-adult
comparisons (two-tailed t-test: t = 2.3484, df = 28, p-value = 0.02615), but did not differ from
juvenile-juvenile comparisons (two-tailed t-test: t = -1.7863, df = 31.781, p-value = 0.0836),
and juvenile-juvenile and adult-adult comparisons were not significantly different from one
another (two-tailed t-test: t = 1.9658, df = 151, p-value = 0.05116). In *L. angustifrons*, F_{ST}
values among juveniles at the same site were lower than those between juveniles and adults
at the same site (two-tailed t-test: t = -3.8334, df = 16.646, p-value = 0.001376) or between
adults at the same site (two-tailed t-test: t = 2.2254, df = 27.295, p-value = 0.0345), while F_{ST}
estimates did not differ significantly between juvenile-juvenile and adult-adult comparisons
(two-tailed t-test: t = -1.6493, df = 14.084, p-value = 0.1212). In *L. microlepis*, F_{ST} estimates
for fish from the same site were marginally higher in juvenile-juvenile pairs than in adult-adult
pairs (two-tailed t-test: t = 2.2525, df = 19.005, p-value = 0.03631), but did not differ between
each of these categories and juvenile-adult comparisons (p-value > 0.07). We do not find any
evidence for juveniles collected at the same time of year to be more closely related than
juveniles collected during different parts of the year in *L. microlepis* (two-tailed t-test: t = -

1.2229, df = 378.64, p-value = 0.2221; Fig. S15), and do not have enough data to analyze
this relationship in the other three species.

Discussion

Differences in life history characteristics can have significant influence on variation in
population genetic structure among taxa, even in closely related species. In this study, we
analyzed population genetic structure within the four endemic pelagic top predators of Lake
Tanganyika, *Lates spp.*, using reduced-representation genomic sequencing data sets. From
our samples collected at discrete sampling sites along the length of the lake, we find two key
patterns: first, we find a strongly differentiated genetic group in *Lates mariae* at the most
southern sampling site, Kasanga; second, we find no additional pattern of isolation-by-
distance or geographic population structure within any of the four *Lates* species. We find
weak evidence for weakly differentiated genetic groups that exist together at the majority of
sites in all four species. While numerous explanations could account for a lack of geographic
population structure and the existence of sympatric weakly differentiated groups, more
research is needed to understand mechanisms causing the population structure identified in
these species. Our results have potential implications for understanding the conservation and
management of these fishes, with implications for this regionally important fishery.

Strong genetic differentiation in southern Lates mariae

In *L. mariae*, the majority of samples from the most southern site, Kasanga, are strongly
genetically differentiated from all other samples. The magnitude of F_{ST} estimates between
these distinct individuals and the other *L. mariae* individuals (F_{ST} ~0.135), suggests
unexpectedly strong intraspecific genetic structure given a lack of obvious geographic
barriers. Our sampling does not include *L.* microlepis or *L. angustifrons* individuals from this
furthest south site, so we cannot use our current data to determine whether this pattern is
also reflected in the other two large *Lates* species.

It is plausible that the differentiation in *L. mariae* may be driven by limnological patterns that
differ between the northern and southern portions of the lake. In particular, the southern part
of the lake experiences intense upwelling of cold, nutrient rich waters from the deep during
the dry season (Verburg *et al.* 2011), which mixes with oxygenated water from the surface
and consequently has high nutrient concentrations and primary productivity rates, which in
turn fuels zooplankton growth (Plisnier *et al.* 2009; Loiselle *et al.* 2014). In contrast, the
northern part of the lake has a more stable, persistently stratified, clear water column
(Plisnier *et al.* 1999; Bergamino *et al.* 2010; Mziray *et al.* 2018). As a result, there is
asynchronous variability between environments in the north and south, as well as high

environmental variability over the course of the year in the south (i.e, higher primary
productivity in the dry season compared to the rainy season). This spatial variability affects
zooplankton community composition, with shrimps and calanoid copepods prevailing in the
south (Kurki *et al.* 1999), while cyclopoid copepods and medusa dominate in the north (Kurki
et al. 1999; Cirhuza and Plisnier 2016). Increases in primary productivity generally lead to
high densities of zooplankton, as well as *Stolothrissa* and *Limnothrissa* sardines (Mulimbwa
et al. 2014), as the blooms seem to positively affect sardine spawning and recruitment,
although recruitment pulses sometimes occur during times of poor nutrient conditions.
Mannini *et al.* (1999) found corresponding spatial variability in diet for *L. stappersii*, but
spatial variability in diet has not been documented for *L. mariae* or the other large-bodied
species. While adult *L. mariae* do not depend on the pelagic sardines as a primary food
source (Coulter 1976), it is possible that such temporal nutrient variability leads to variability
in the main *L. mariae* prey (benthic cichlids) as well.

Such a spatially and temporally variable environment might lead to local differential selection
in the southern part of the lake, and this selection combined with restricted movement of
individuals between this and other populations of *L. mariae* could lead to the genetic
differentiation in individuals from Kasanga. *L. mariae* abundance is greater in the south than
in the north (Coulter 1970; personal communication), which suggests that spatial variation in
environmental conditions may underlie differences in fitness or population growth. A variable
environment may also lead to within-site differentiation among individuals caught in Kasanga
if the two groups were to adapt to this variability in different ways.

Another possibility is that one group is resident at Kasanga, while the other disperses
throughout the lake. While all *L. mariae* individuals sampled from Kasanga were mature and
L. mariae is believed to be the least vagile of the four *Lates* species (Coulter 1976), it is not
clear how vagile these individuals typically are; it is possible that some individuals remain
resident while others migrate, as is the case in Atlantic cod (Kirubakaran *et al.* 2016; Berg et
al. 2017). To investigate these possibilities, future studies should expand sampling in the
southern portion of the lake, including sampling various life stages of this and other *Lates* to
understand the genetic composition of the southern populations and any important ecological
implications of genetic differentiation.

It is also possible that the differentiation in the most southern part of Lake Tanganyika in *L.*
mariae is related to the lake's bathymetry and differences in the depth of the oxygenated
layer, as suggested by Coulter (1976), who observed asynchronous changes in local
abundance between *L. mariae* populations in the southwest versus southeast arms of the
lake. From catch rates, Coulter (1970) observed that southern populations of at least two of
the *Lates* species (*L. angusitfrons* and *L. mariae*) were likely distinct, as they were heavily

exploited in the southeast arm with no apparent effect on catches in the southwest arm.
These southern reaches of the lake are relatively shallow, and the oxygenated layer extends
up to 200m deep, which allows for an abundance of oxygenated habitat for benthic species
(Coulter 1976).

Despite the life history differences between pelagic cichlids and the *Lates* species, the
genetic patterns observed in *L. mariae* are similar to those observed previously in the
benthopelagic cichlids. For the four benthopelagic *Diplotaxodon* species of Lake Malawi with
genetically structured populations, research by Genner *et al.* (2010) indicated spawning site
fidelity drives population structure. In contrast, Koblmüller *et al.* (2019) hypothesized that
preying upon benthic cichlids—which requires shorter-distance movements than preying on
pelagic prey—can explain the genetic structure in the *Bathybatini* of Lake Tanganyika.
Similar to the *Bathybatini, L. mariae* hunts benthic cichlids and is described as relatively
resident (Coulter 1976). However, our results did not recover any evidence for significant
isolation-by-distance in *L. mariae*, suggesting that the genetic structure that we observe is
likely not the result of dispersal limitation. Interestingly, females of *L. mariae* likely aggregate
over specific spawning grounds (Coulter 1991), similar to *Diplotaxon*. If samples of *L. mariae*
were taken from spawning grounds, it would be possible to infer whether the divergence we
observe is related to strong spawning site fidelity and differentiation between these stocks.
However, the sampling methods used in our study (i.e., obtaining fish from fishermen) limit
our knowledge of exact catch location. In addition, these sampling methods mean that a
range of ages and reproductive maturities of fish were included in our study. To our
knowledge, the location of spawning grounds has not been documented for any of the four
Lates species. In addition, it is unclear how the pelagic phase of eggs, larvae, and small
juveniles (<2.5cm), of *L. mariae* influences dispersal in the species. Spawning pelagically
(Riginos *et al.* 2014) and having a long pelagic larval stage (Selkoe & Toonen 2011; Shanks
2009) generally facilitate gene flow and reduce population structure in marine taxa, although
these patterns are complicated by specific behavior exhibited by both adults and larvae (Bay
et al. 2006; Riginos *et al.* 2011). Future studies in *L. mariae* should sample spawning fish
and work to understand how genetic divergence relates to spawning site location and timing.

No pattern of isolation-by-distance or geographic population structure

Although the samples spanned the ~550 km Tanzanian shoreline of Lake Tanganyika, none
of the genetic differences between populations relates to geographic distance, suggesting no
isolation-by-distance in any of the four species. In addition, admixture models with only a
single panmictic population fit our data the best in all species except for *L. mariae* (Fig. 5).
This suggests that individuals of all four species are mobile enough that gene flow is not
restricted among our sampling sites, other than the distinct Kasanga population.

Deviations from isolation-by-distance are common among marine animals, where
geographically proximate populations often have greater population differentiation than
populations further apart, a phenomenon sometimes referred to as "chaotic patchiness"
(Johnson and Black 1984). This phenomenon has been attributed to larval dispersal via
ocean currents (White et al. 2010), and it is possible that currents within Lake Tanganyika
may similarly disperse *Lates* eggs and larvae prior to their movement to inshore habitats.
While it is presumed that all four *Lates* species spawn pelagically (Ellis 1978; Coulter 1991)
and the larvae of the larger three species are planktonic until ~3cm in size (Coulter 1976),
spawning behavior, spawning location within the lake, and the length of time that larvae may
be carried around by lake currents remain undocumented for all four species. Lake currents
vary throughout the year (Plisnier *et al.* 1999; Verburg *et al.* 2011), and therefore
understanding the timing and location of reproduction for all four species will be important to
understanding the role that these currents may play in dispersing planktonic eggs and larvae.

While geographically widespread populations are not particularly surprising given the large
body sizes and long-distance swimming capabilities of these fishes, it conflicts with evidence
of a resident lifestyle in *L. mariae* (Chapman 1976; Coulter 1976, 1991; Muller *et al.* 2001)
and evidence for some spatial structure in the Kigoma area in a previous population genetic
study of *L. stappersii* (Kuusipalo 1999). It should be noted that our sampling is unbalanced
among sites and lacking in the most southern reaches of the lake for *L. angustifrons* and *L.*
microlepis. It is therefore possible that more (albeit still likely weak) genetic structure exists
than we have been able to detect with available samples. That said, Koblmüller *et al.* (2019)
similarly found no evidence of genetic population structure in two eupelagic bathybatine
cichlid species of Lake Tanganyika, Koblmüller *et al.* (2015) found no evidence for population
structure in the pelagic cichlid *Boulengerochromis microlepis*, and De Keyzer *et al.* (2019)
and Junker *et al.* (2020) found no population structure in either of the pelagic sardine
species. There are many life history differences between the sardines, cichlids, and *Lates*
spp. that occur sympatrically in the open waters of Lake Tanganyika, but these results
together suggest that spatial environmental heterogeneity along the length of the lake is not
sufficient to restrict gene flow and produce geographically isolated or clinal populations in
these diverse species.

Sympatric, weakly-differentiated groups exist in all four species

While our admixture models suggest panmixia as the best model in *L. stappersii, L.*
microlepis, and *L. angustifrons*, and two distinct groups in *L. mariae*, we find sympatric
differentiation in all four *Lates* species at higher K values. Model DIC values suggest that
these higher K values are not the best fit for our data, and differentiation is weak (mean F_{ST} =
0.032; Table S4), but individual assignments do not behave as would be expected for truly

panmictic populations. In our data, we see individuals with high assignment confidence to
one group or another (i.e., each bar is mostly a single color in Fig. 5 and Fig. S6); if the
population were truly panmictic, then we would expect to see all individuals showing genetic
contributions from all groups at higher than optimal values of K. For that reason, we believe
these weakly differentiated groups may be biologically relevant, albeit weakly differentiated.
These groups do not appear to correspond to sex or any other phenotypic character that we
recorded. There are several potential explanations for this consistent structure, of which we
will discuss the evidence for three: (a) spatially segregated spawning sites with spawning
site-based differentiation, where adults move throughout the lake but return to the same
spawning grounds to breed; (b) temporally segregated spawning groups; or (c) historical
allopatric groups that have become sympatric yet partially retained their genetic
distinctiveness.

Genetically distinct groups that occur in sympatry must have at least partial reproductive
isolation, and in fish without clear sexual selection, this often occurs intraspecifically as a
segregation of spawning either in space or time. If there is site fidelity to regionally
differentiated spawning grounds but between-region movement outside of spawning, then
this could create patterns of differentiation such as those that we observed. If spatially
segregated spawning sites exist, then the adults that are sampled at a single site may have
migrated from a more distant site (unless they are collected while spawning, in which case
we would expect the genetics to match with their geographic location), but we would expect
littoral juveniles to exhibit a stronger pattern of geographic-based differentiation or isolation
by distance. In addition, we would expect individual differentiation among juveniles collected
at the same site to be lower than differentiation among adults collected at the same site, or
differentiation between juveniles and adults collected at the same site. We did not find this to
be the case in any of the four species (Fig. S14), with the caveat that our sample sizes for
these comparisons are low and our data may exhibit batch effects due to our sampling
strategies. We did find stronger differentiation in *L. angustifrons* and *L. mariae* between
juveniles and adults collected at the same site than among juveniles collected at the same
site; however, the mean F_{ST} for adult-adult comparisons was not larger than that of juvenile-
juvenile comparisons, so the evidence supporting this hypothesis is mixed.

Genetic differentiation related to temporal stratification in spawning behavior—such as
having a population of "dry season" spawners and a population of "rainy season" spawners—
could also explain the existence of sympatric groups. Coulter (1976) suggested that the four
species likely spawn continuously, but with peaks in number of ripe females from August to
November/December, and potentially other smaller peaks in March/April. In addition,
observations of well-differentiated size distributions of young *Lates* found in littoral weed
beds suggests periodicity in spawning (Coulter 1976). Unfortunately, we do not have data on

the spawning status of all of the fish collected in this study, which would facilitate
investigating this hypothesis. However, we were able to test whether juveniles collected at a
similar time of year were more closely related than those collected at a different time of year,
regardless of sampling site. We do not find any evidence for juveniles collected at the same
time of year to be more closely related than juveniles collected during different parts of the
year in *L. microlepis* (Fig. S15); however, our sampling is not conducive to being able to
rigorously investigate this relationship, as some sites were only sampled in one year or at
one time of year. Additionally, only a few juveniles were collected opportunistically from each
species, as they are too small to be targeted by fishermen. Investigating the possibility of
discrete spawning groups will be an important next step in understanding the weak genetic
structure present in these species, and will require more targeted sampling.

A third potential explanation for the intraspecific structure could lie in the geologic history of
Lake Tanganyika. With an estimated age of 9-12 million years, Lake Tanganyika is one of
the oldest lakes in the world and has been influenced by extensive lake level fluctuations
(Cohen *et al.* 1993). Based on the bathymetry of the lake, it is thought that Lake Tanganyika
was divided into three paleo-lakes during periods of extremely low water levels (Cohen *et al.*
1997). Such environmental fluctuations may have led to oscillations between sympatry or
parapatry and allopatry for fish populations within the lake, which have been invoked as a
"species pump" (Greenwood 1981; Sedano *et al.* 2010) contributing to diversity in some
clades in the littoral (Janzen & Etienne 2017; Sturmbauer *et al.* 2016) and benthic or pelagic
(Duftner *et al.* 2005; Koblmüller *et al.* 2005) zones of the lake. While it has recently been
suggested that the origin of the Lake Tanganyika *Lates* radiation is much younger than other
endemic fish radiations in the lake (Koblmüller *et al.* 2021), it remains unclear whether
speciation in this clade was similarly triggered by these environmental fluctuations—as well
as whether more recent fluctuations could explain our contemporary observations of weak
genetic structure. Such fluctuations could also lead to persistent genetic structure within
species, for example, if spawning site fidelity within separate basins is maintained after the
basins reconnect. However, we would expect strong philopatry to produce much stronger
genetic differentiation than observed here.

Interestingly, *L. stappersii* in this study which were caught during the dry season in the south
of the lake show distinct carbon isotope signals from samples caught in the north (Ehrenfels
et al. 2021). These differences in carbon isotope ratios correlate with different rates of
primary productivity in the northern and the southern basin during the dry season (Loiselle *et*
al. 2014), and with differences in prey composition in different areas of the lake (Plisner *et al.*
2009; Kurki *et al.* 1999). This is evidence for at least partly resident populations of *L.*
stappersii, which stay in the northern or southern basin during the dry season for long
enough to incorporate the different isotopic signals of these basins. Such movement patterns

may reduce gene flow between *L. stappersii* populations from the basins; however, these
isotopic groups do not correspond with the genetic groups found at K=2 here (Ehrenfels *et al*,
2021)*.*

It is also important to keep the limitations of these data in mind. Admixture analyses using
programs such as STRUCTURE (Pritchard *et al.* 2000) are biased in cases of unbalanced
sampling among sites, such that differentiation is often overestimated at sites with the most
individuals and underestimated at sites with few individuals (e.g., Puechmaille 2016; Wang
2017). While it is unclear whether this is also true for entropy analyses, it is possible that a
similar issue influences our results. For example, we find multiple groups at the Kigoma site
in *L. microlepis* at K=2, which is also the site with the most individuals (Fig. 5B). Furthermore,
our assignment of individual locations to fisherman landing sites or markets is less precise
than if we had exact locations where the fish were caught. While we only have general
knowledge of how far fishermen travel from the landing site to fish during the day, it is
possible that fish collected at the same sampling site were collected up to ~40 km away from
one another. In addition, our samples collected in the Kigoma fish markets may have come
from farther away from those collected at landing sites (Petit & Shipton 2012). These
limitations may influence our ability to detect geographic trends in our data.

Recommendations for conservation and sustainable management of Lake Tanganika's Lates

We find evidence for weak differentiation within sampling sites in all four species. We
furthermore find clear evidence for a genetically distinct group within *L. mariae* at our most
southern sampling site, in agreement with previous studies suggesting that this species is the
most resident of all four *Lates* species (Coulter 1976, 1991). Intriguingly, the weak
intraspecific structure in all four species is not linked to isolation-by-distance, but rather
involves differentiation between some individuals at multiple sites, suggesting the existence
of a mechanism for maintaining differentiation while these distinct genetic groups are in
sympatry. In contrast, the low F_{ST} estimates between samples from distant sites and lack of
isolation-by-distance suggest that these cies disperse over large distances, indicating a need
for management of these species on at a lake-wide scale. Furthermore, it is likely that more
extensive sampling of these little-studied species may uncover important additional diversity.

To maintain intraspecific genetic variation, viable population sizes, and possible adaptive
differences between genetic groups within the *Lates* species, more research is needed on
the location and timing of spawning for the four species and whether restricting catches in
certain areas and periods of the year could help the larger-bodied species recover from low
population sizes. Harvest restrictions have been implemented in regions around Lake
Tanganyika where community conservation areas are active (Kimirei & Sweke 2018) but are
not in place throughout the lake. Such management practices are performed successfully in

the cod fishery of Denmark (albeit after collapse of this fishery and extinction of many of the
original stocks; Dahle *et al.* 2018), and in the salmon fishery of Bristol Bay (Schindler *et al.*
2015; Schindler *et al.* 2010). However, implementation of such a strategy requires detailed
evidence of spawning ground locations and timing of spawning, as well as for fishermen to
be able to distinguish between the genetically distinct stocks. Therefore, studies are needed
to shed more light on the phenotypic, ecological, and phenological characteristics of the
genetic groups within each species. In addition, studies are needed to examine whether fish
from different spawning grounds form the distinct genetic clusters that we identified in this
study. Genomic analyses, although helpful in generating hypotheses about the natural
history of these species, must be paired with an in-depth understanding of natural history,
including integration of local knowledge, for managing these fish stocks. A sustainable
management approach informed by these combined data is crucial to protect the inshore
nursery habitats and ensure recruitment of the species in Lake Tanganyika.

Funding

This work was supported by the Swiss National Science Foundation (grant CR23I2-166589
to OS), the University of Wyoming (start-up funding to CEW), the National Science
Foundation (grant DEB-1556963 to CEW), the National Institute of General Medical Sciences
of the National Institutes of Health (Institutional Development Award grant P20GM103432),
and the American Genetics Association (Ecological, Evolutionary, and Conservation
Genomics Research Award to JAR).

Acknowledgements

Computing was accomplished with an allocation from the University of Wyoming's Advanced
Research Computing Center, on its Teton Intel x86_64 cluster
(https://doi.org/10.15786/M2FY47) and the Genetic Diversity Center (GDC) of ETH Zürich.
Special thanks go to Benedikt Ehrenfels for coordinating field campaigns, Mupape Mukuli for
facilitating logistics during fieldwork, and to the crew of the MV Maman Benita for fieldwork
assistance. We thank the whole team at the Tanzanian Fisheries Research Institute for their
support. A special thanks goes to Mary Kishe for her support during fieldwork permission
processes and to the Tanzanian Commission for Science and Technology (COSTECH) for
the permits to do this research. Live fish samples were collected under the approved
University of Wyoming IACUC protocol #20160602CW00241-01. We appreciate The Nature
Conservancy's Tuungane Project for providing funding and logistical support, with special
thanks going to Colin Apse and Peter Limbu. We also thank the members of the Wagner Lab

at the University of Wyoming and the FishEc group at EAWAG for discussions about the
analyses and manuscript. We also thank the Associate Editor and four anonymous reviewers
for the valuable comments that greatly improved the quality of this manuscript.

Data Availability

We have deposited the primary data underlying these analyses as follows:

- Sampling locations, morphological data, and filtered VCF files have been archived on
Zenodo (http://doi.org/10.5281/zenodo.5216259)
- Fastq files for each individual have been archived in NCBI SRA PRJNA776855, and
will be released upon publication
- Scripts for data analysis are available on Github (https://github.com/jessicarick/lates-
popgen), and the version at publication will be archived on Zenodo
(http://doi.org/10.5281/zenodo.5216259)

References

Anderson, EC (2018) whoa: Where's my Heterozygotes? Observations on genotyping
Accuracy. Version 0.0.2.999. *Zenodo*, doi: 10.5281/zenodo.2401764
Baird NA, Etter PD, Atwood TS*, et al.* (2008) Rapid SNP discovery and genetic mapping
using sequenced RAD markers. *PLoS One* **3**.
Bay LK, Crozier RH, Caley MJ (2006) The relationship between population genetic structure
and pelagic larval duration in coral reef fishes on the Great Barrier Reef. *Marine*
Biology **149**, 1247-1256.
Berg PR, Star B, Pampoulie C*, et al.* (2017) Trans-oceanic genomic divergence of Atlantic
cod ecotypes is associated with large inversions. *Heredity* **119**, 418-428.
Bergamino N, Horion S, Stenuite S*, et al.* (2010) Spatio-temporal dynamics of phytoplankton
and primary production in Lake Tanganyika using a MODIS based bio-optical time
series. *Remote Sensing of Environment* **114**, 772-780.
Bernatchez L (2016) On the maintenance of genetic variation and adaptation to
environmental change: considerations from population genomics in fishes. *Journal of*
Fish Biology **89**, 2519-2556.
Brannon EL, Powell MS, Quinn TP, Talbot A (2004) Population Structure of Columbia River
Basin Chinook Salmon and Steelhead Trout. *Reviews in Fis heries Science* **12**, 99-
232.
Bulengela G, Onyango P, Brehm J, *et al.* (2020) "Bring fishermen at the center": the value of
local knowledge for understanding fisheries resources and climate-related changes in
Lake Tanganyika. *Environment, Development and Sustainability* **22**, 5621-5649.
Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis
tool set for population genomics. *Mol Ecol* **22**, 3124-3140.
Chang CC, Chow CC, Tellier LCAM, *et al.* (2015) Second-generation PLINK: rising to the
challenge of larger and richer datasets. *Gigascience* **4**, s13742-015-0047-8.
Chapman DW (1976) Acoustic Estimates of Pelagic Ichthyomass in Lake Tanganyika with an
Inexpensive Echo Sounder. *Transactions of the American Fisheries Society* **105**, 581-
587.
Cirhuza DM, Plisnier PD (2016) Composition and seasonal variations in abundance of
Copepod (Crustacea) populations from the northern part of Lake Tanganyika. *Aquatic*
Ecosystem Health & Management **19**, 401-410.
Cohen AS, Gergurich EL, Kraemer BM*, et al.* (2016) Climate warming reduces fish
production and benthic habitat in Lake Tanganyika, one of the most biodiverse
freshwater ecosystems. *Proc Natl Acad Sci U S A* **113**, 9563-9568.
Cohen AS, Lezzar K-E, Tiercelin J-J, Soreghan M (1997) New palaeogeographic and lake-
level reconstructions of Lake Tanganyika: implications for tectonic, climatic and
biological evolution in a rift lake. *Basin Research* **9**, 107-132.
Cohen AS, Soreghan MJ, Scholz CA (1993) Estimating the age of formation of lakes: an
example from Lake Tanganyika, East African Rift system. *Geology* **21**, 511-514.
Coulter GW (1970) Population changes within a group of fish species in Lake Tanganyika
following their exploitation. *Journal of Fish Biology* **2**, 329-353.
Coulter GW (1976) The biology of Lates species (Nile perch) in Lake Tanganyika, and the
status of the pelagic fishery for Lates species and Luciolates stappersii (Blgr.).
Journal of Fish Biology **9**, 235-259.
Coulter GW (1991) *Lake Tanganyika and its Life* British Museum (Natural History) Cromwell
Road, London SW7 5BD & Oxford University Press, Walton Street, Oxford OX2 6DP.
Dahle G, Johansen T, Westgaard J-I, Aglen A, Glover KA (2018) Genetic management of
mixed-stock fisheries "real-time": The case of the largest remaining cod fishery
operating in the Atlantic in 2007–2017. *Fisheries Research* **205**, 77-85.
Danecek P, Auton A, Abecasis G*, et al.* (2011) The variant call format and VCFtools.
Bioinformatics **27**, 2156-2158.
De Keyzer ELR, De Corte Z, Van Steenberge M, *et al.* (2019) First genomic study on Lake
Tanganyika sprat *Stolothrissa tanganicae*: a lack of population structure calls for

integrated management of this important fisheries target species. *BMC Ecology and*
Evoluiton **19**, 6.
De Keyzer ELR, Mulungula PM, Lufungula GA, *et al.* (2020) Local perceptions on the state of
the pelagic fisheries and fisheries management in Uvira, Lake Tanganyika, DR
Congo. *Journal of Great Lakes Research* **46**, 1740-1753.
Dray S, Dufour A-B (2007) The ade4 Package: Implementing the Duality Diagram for
Ecologists. *2007* **22**, 20.
Duftner N, Koblmüller S, Sturmbauer C (2005) Evolutionary relationships of the
Limnochromini, a tribe of benthic deepwater cichlid fish endemic to Lake Tanganyika,
East Africa. *Journal of Molecular Evolution* **60**, 277-289.
Eccles DH (1992) Field guide to the freshwater fishes of Tanzania. In: *species identification*
sheets for fishery purposes (ed. FAO), p. 145. FAO, Rome.
Ehrenfels B, Junker J, Namutebi D, *et al.* (2021) Isotopic signatures induced by upwelling tag
regional fish populations in Lake Tanganyika, *bioRxiv*, doi:
10.1101/2021.10.05.463178.
Ellegren H, Galtier N (2016) Determinants of genetic diversity. *Nature Reviews Genetics* **17**,
422-433.
Ellis CMA (1978) Biology of Luciolates-Stappersi in Lake Tanganyika (Burundi). *Transactions*
of the American Fisheries Society **107**, 557-566.
Genner MJ, Nichols P, Shaw PW*, et al.* (2010) Population structure on breeding grounds of
Lake Malawi's 'twilight zone' cichlid fishes. *Journal of Biogeography* **37**, 258-269.
Gompert Z, Lucas LK, Buerkle CA*, et al.* (2014) Admixture and the organization of genetic
diversity in a butterfly species complex revealed through common and rare genetic
variants. *Mol Ecol* **23**, 4555-4573.
Greenwood, PH (1981) Species-flocks and explosive evolution. *Chance, change and*
challenge- the evolving biosphere, 61-74.
Gruber B, Unmack PJ, Berry OF, Georges A (2018) dartr: An r package to facilitate analysis
of SNP data generated from reduced representation genome sequencing. *Molecular*
Ecology Resources **18**, 691-699.
Hutchinson WF (2008) The dangers of ignoring stock complexity in fishery management: the
case of the North Sea cod. *Biol Lett* **4**, 693-695.
Ivory SJ, McGlue MM, Peterman C, *et al* (2021) Climate, vegetation, and weathering across
space and time in Lake Tanganyika (tropical eastern Africa). *Quaternary Science*
Advances **3**, 10023.
Janzen T, Etienne RS (2017) Inferring the role of habitat dynamics in driving diversification:
evidence for a species pump in Lake Tanganyika cichlids.
Johnson MS, Black R (1984) Pattern beneath the chaos: The effect of recruitment on genetic
patchiness in an intertidal limpet. *Evolution* **38**, 1371-1383.
Junker J, Rick JA, McIntyre PB*, et al.* (2020) Structural genomic variation leads to genetic
differentiation in Lake Tanganyika's sardines. *Mol Ecol* **29**, 3277-3298.
Kimirei IA, Mgaya YD, Chande AI (2008) Changes in species composition and abundance of
commercially important pelagic fish species in Kigoma area, Lake Tanganyika,
Tanzania. *Aquatic Ecosystem Health and Management* **11**, 29-35.
Kimirei IA, Sweke EA (2018) Community-based conservation areas as a fisheries
management tool in the East African Great Lakes: an example from Tanzania. In: 34[th]
SIL Congress, 19-24 August 2018. Nanjing, China. pp. 230 Book of abstracts.
Kirubakaran TG, Grove H, Kent MP*, et al.* (2016) Two adjacent inversions maintain genomic
differentiation between migratory and stationary ecotypes of Atlantic cod. *Mol Ecol*
25, 2130-2143.
Kmentová N, Koblmüller S, Van Steenberge M, *et al.* (2020) Weak population structure and
recent demographic expansion of the monogenean parasite *Kapentagyrus* spp.
infecting clupeid fishes of Lake Tanganyika, East Africa. *International Journal for*
Parasitology **50**, 471-486.
Knaus BJ, Grünwald NJ (2017) VCFR: a package to manipulate and visualize variant call
format data in R. *Mol Ecol Resources* **17**, 44-53.

Koblmüller S, Duftner N, Katongo C, *et al.* (2005) Ancient divergence in bathypelagic Lake
Tanganyika deepwater cichlids: Mitochondrial phylogeny of the tribe Bathybatini.
Journal of Molecular Evolution **60**, 297-314.
Koblmüller S, Odhiambo EA, Sinyinza D, *et al.* (2015) Big fish, little divergence:
phylogeography of Lake Tanganyika's giant cichlid, *Boulengerochromis microlepis*.
Hydrobiologia **748**, 29-38.
Koblmüller S, Zangl L, Börger C*, et al.* (2019) Only true pelagics mix: comparative
phylogeography of deepwater bathybatine cichlids from Lake Tanganyika.
Hydrobiologia **832**, 93-103.
Koblmüller S, Schöggl CA, Lorber CJ, *et al.* (2021) African lates perches (Teleostei, Latidae,
Lates): Paraphyly of Nile perch and recent colonization of Lake Tanganyika.
Molecular Phylogenetics and Evolution **160**, 107141.
Korneliussen TS, Albrechtsen A, Nielsen R (2014) ANGSD: Analysis of Next Generation
Sequencing Data. *BMC Bioinformatics* **15**, 356.
Korneliussen TS, Moltke I, Albrechtsen A, Nielsen R (2013) Calculation of Tajima's D and
other neutrality test statistics from low depth next-generation sequencing data. *BMC*
Bioinformatics **14**, 289.
Kurki H, Mannini P, Vuorinen I, *et al.* (1999) Macrozooplankton communities in Lake
Tanganyika indicate food chain differences between the northern part and the main
basins. *Hydrobiologia* **407**, 123-129.
Kuusipalo L (1999) Genetic differentiation of endemic nile perch *Lates stappersi*
(*Centropomidae*, *Pisces*) populations in Lake Tanganyika suggested by RAPD
markers. In: Lindqvist O.V., Mölsä H., Salonen K., Sarvala J. (eds) *From Limnology to*
Fisheries: Lake Tanganyika and Other Large Lakes. Developments in Hydrobiology,
vol 141. Springer, Dordrecht.
Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**,
357-359.
Lewis PO (2001) A Likelihood Approach to Estimating Phylogeny from Discrete
Morphological Character Data. *Systematic Biology* **50**, 913-925.
Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics (Oxford, England)* **25**, 1754-1760.
Li H, Handsaker B, Wysoker A*, et al.* (2009) The Sequence Alignment/Map format and
SAMtools. *Bioinformatics* **25**, 2078-2079.
Loiselle S, Cózar A, Adgo E, *et al.* (2014) Decadal Trends and Common Dynamics of the
Bio-Optical and Thermal Characteristics of the African Great Lakes. *PLoS One* **9**,
e93656.
Manier MK, Arnold SJ (2006) Ecological correlates of population genetic structure: a
comparative approach using a vertebrate metacommunity. *Proceedings of the Royal*
Society B: Biological Sciences **273**, 3001-3009.
Mannini P (1998a) *Ecology of the Pelagic Fish Resources of Lake Tanganyika*, University of
Hull.
Mannini P, Aro E, Katonda I*, et al.* (1996) Pelagic fish stocks of Lake Tanganyika: biology
and exploitation. FAO/FINNIDA Research for the Management of the Fisheries of
Lake Tanganyika. In: *GCP/RAF/271/FIN—TD/53 (En)*, p. 60p.
Mannini P, Katonda I, Kissaka B, Verburg P (1999) Feeding ecology of Lates stappersii in
Lake Tanganyika. *Hydrobiologia* **407**, 131-139.
Martinez Barrio A, Lamichhaney S, Fan G*, et al.* (2016) The genetic basis for ecological
adaptation of the Atlantic herring revealed by genome sequencing. *Elife* **5**.
McGlue MM, Ivory SJ, Stone JR, *et al.* (2020) Solar irradiance and ENSO affect food security
in Lake Tanganyika, a major African inland fishery. *Science Advances* **6**, eabb2191.
Meisner J, Liu S, Huang M, Albrechtsen A (2021) Large-scale inference of population
structure in presence of missingness using PCA. *Bioinformatics*, btab027.
Mölsä H, Reynolds JE, Coenen EJ, Lindqvist OV (1999) Fisheries research towards resource
management on Lake Tanganyika. *Hydrobiologia* **407**, 1-24.

Mölsä H, Sarvala J, Badende S*, et al.* (2002) Ecosystem monitoring in the development of
sustainable fisheries in Lake Tanganyika. *Aquatic Ecosystem Health and*
Management **5**, 267-281.
Mulimbwa N, Raeymaekers JAM, Sarvala J (2014) Seasonal changes in the pelagic catch of
two clupeid zooplanktivores in relation to the abundance of copepod zooplankton in
the northern end of Lake Tanganyika. *Aquatic Ecosystem Health & Management* **17**,
25-33.
Muller P, Li XP, Niyogi KK (2001) Non-photochemical quenching. A response to excess light
energy. *Plant Physiol* **125**, 1558-1566.
Munyandero J (2002) The lake Tanganyika clupeid and latid fishery system: Indicators and
problems inherent in assessments and management. *African Study Monographs* **23**,
117-145.
Mziray P, Kimirei IA, Staehr PA, *et al.* (2018) Seasonal patterns of thermal stratification and
primary production in the northern parts of Lake Tanganyika. *Journal of Great Lakes*
Research **44**, 1209–1220.
O'Reilly CM, Alin SR, Piisnier PD, Cohen AS, McKee BA (2003) Climate change decreases
aquatic ecosystem productivity of Lake Tanganyika, Africa. *Nature* **424**, 766-768.
Ogutu-Ohwayo R, Natugonza V, Musinguzi L, *et al.* (2016) Implications of climate variability
and change for African lake ecosystems, fisheries productivity, and livelihoods.
Journal of Great Lakes Research **42**, 498-510.
Olsen JB, Crane PA, Flannery BG*, et al.* (2011) Comparative landscape genetic analysis of
three Pacific salmon species from subarctic North America. *Conservation Genetics*
12, 223-241.
Parchman TL, Gompert Z, Mudge J*, et al.* (2012) Genome-wide association genetics of an
adaptive trait in lodgepole pine. *Mol Ecol* **21**, 2991-3005.
Peterman WE, Anderson TL, Ousterhout BH*, et al.* (2015) Differential dispersal shapes
population structure and patterns of genetic differentiation in two sympatric pond
breeding salamanders. *Conservation Genetics* **16**, 59-69.
Petit P, Shipton T (2012) IUU Fishing on Lake Tanganyika. *Programme for the*
Implementation of a Regional Fisheries Strategy for the Eastern and Southern Africa
and Indian Ocean Region.
Pettersson ME, Rochus CM, Han F*, et al.* (2019) A chromosome-level assembly of the
Atlantic herring genome-detection of a supergene and other signals of selection.
Genome Research **29**, 1919-1928.
Plisnier PD, Chitamwebwa D, Mwape L, *et al.* (1999) Limnological annual cycle inferred from
physical-chemical fluctuations at three stations of Lake Tanganyika. *Hydrobiologia*
407, 45-58.
Plisnier PD, Mgana H, Kimirei I, *et al.* (2009) Limnological variability and pelagic fish
abundance (Stolothrissa tanganicae and Lates stappersii) in Lake Tanganyika.
Hydrobiologia **625**,117-134.
Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using
multilocus genotype data. *Genetics* **155**: 945-959.
Puechmaille SJ (2016) The program STRUCTURE does not reliably recover the correct
population structure when sampling is uneven: subsampling and new estimators
alleviate the problem. *Molecular Ecology Resources* **16**, 608-627.
Purcell S, Neale B, Todd-Brown K, *et al.* (2007) PLINK: a toolset for whole-genome
association and population-based linkage analysis. *American Journal of Human*
Genetics, 81: 559-575.
R Core Team (2021) R: A Language and Environment for Statistical Computing. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.
Reich D, Thangaraj K, Patterson N, Price AL, Singh L (2009) Reconstructing Indian
population history. *Nature* **461**, 489-494.
Reiss H, Hoarau G, Dickey-Collas M, Wolff WJ (2009) Genetic population structure of marine
fish: mismatch between biological and fisheries management units. *Fish and*
Fisheries **10**, 361-395.

Riginos C, Buckley YM, Blomberg SP*, et al.* (2014) Dispersal Capacity Predicts Both
Population Genetic Structure and Species Richness in Reef Fishes. *The American*
Naturalist **184**, 52-64.
Riginos C, Doublas KE, Jin Y, *et al.* (2011) Effects of geography and life history traits on
genetic differentiation in benthic marine fishes. *Ecography* **34**, 566-575.
Sarvala J, Tarvainen M, Salonen K, Mölsä H (2002) Pelagic food web as the basis of
fisheries in Lake Tanganyika: A bioenergetic modeling analysis. *Aquatic Ecosystem*
Health and Management **5**, 283-292.
Sayers M, Bosse K, Fahnenstiel G, Shuchman R (2020) Carbon fixation trends in eleven of
the world's largest lakes: 2003-2018. *Water* **12**, 3500.
Schindler DE, Armstrong JB, Reed TE (2015) The portfolio concept in ecology and evolution.
Frontiers in Ecology and the Environment **13**, 257-263.
Schindler DE, Hilborn R, Chasco B*, et al.* (2010) Population diversity and the portfolio effect
in an exploited species. *Nature* **465**, 609-612.
Sedano RE, Burns KJ, Riddle B (2010) Are the Northern Andes a species pump for
Neotropical birds? Phylogenetics and biogeography of a clade of Neotropical
tanagers (Aves: Thraupini). *Journal of Biogeography* **37**, 325-343.
Selkoe KA, Toonen RJ (2011) Marine connectivity: a new look at pelagic larval duration and
genetic metrics of dispersal. *Marine Ecology Progress Series* **436**, 291-305.
Shanks (2009) Pelagic larval duration and dispersal distance revisited. *The Biological*
Bulletin **216**, 373-385.
Shastry, V, Adams PE, Lindtke D, *et al.* (2021) Model-based genotype ancestry estimation
for potential hybrids with mixed-ploidy. *Mol Ecol Resources* **21**: 1434-1451.
Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**, 1312-1313.
Stearns SC (1992) *The evolution of life histories* Oxford Univ. Press, New York.
Stenuite S, Pirlot S, Hardy MA*, et al.* (2007) Phytoplankton production and growth rate in
Lake Tanganyika: evidence of a decline in primary productivity in recent decades.
Freshwater Biology **52**, 2226-2239.
Sterner T (2007) Unobserved diversity, depletion and irreversibility The importance of
subpopulations for management of cod stocks. *Ecological Economics* **61**, 566-574.
Sturmbauer C, Börger C, Van Steenberge M, Koblmüller S (2016) A separate lowstand lake
at the northern edge of Lake Tanganyika? Evidence from phylogeographic patterns in
the cichlid genus Tropheus. *Hydrobiologia* **791**, 51-68.
Sunday JM, Popovic I, Palen WJ, Foreman MG, Hart MW (2014) Ocean circulation model
predicts high genetic structure observed in a long-lived pelagic developer. *Mol Ecol*
23, 5036-5047.
Supple MA, Shapiro B (2018) Conservation of biodiversity in the genomics era. *Genome*
Biology **19**, 131.
Tange O (2011) GNU Parallel - The Command-Line Power Tool. ;login: The USENIX
Magazine, **36**:42-47.
[dataset] Temasek Life Sciences Laboratory (2016). *Lates calcarifer* isolate:ASB-BC8
Genome sequencing and assembly. National Center for Biotechnology Information
Sequence Read Archive, Assembly GCA_001640805.1.
Therkildsen NO, Hemmer-Hansen J, Als TD*, et al.* (2013) Microevolution in time and space:
SNP analysis of historical DNA reveals dynamic signatures of selection in Atlantic
cod. *Mol Ecol* **22**, 2424-2440.
Van der Knaap M (2013) Comparative analysis of fisheries restoration and public
participation in Lake Victoria and Lake Tanganyika. *Aquatic Ecosystem Health and*
Management **16**, 279-287.
Van der Knaap M, Katonda KI, De Graaf GJ (2014) Lake Tanganyika fisheries frame survey
analysis: Assessment of the options for management of the fisheries of Lake
Tanganyika. *Aquatic Ecosystem Health and Management* **17**, 4-13.
van Zwieten PAM, Roest FC, Machiels MAM, Van Densen WLT (2002) Effects of inter-
annual variability, seasonality and persistence on the perception of long-term trends

in catch rates of the industrial pelagic purse-seine fishery of northern Lake
Tanganyika (Burundi). *Fisheries Research* **54**, 329-348.
van Zwieten PAM, Kolding J, Plank MJ*, et al.* (2016) The Nile perch invasion in Lake Victoria:
cause or consequence of the haplochromine decline? *Canadian Journal of Fisheries*
and Aquatic Sciences **73**, 622-643.
Verburg P, Antenucci JP, Hecky RE (2011) Differential cooling drives large-scale convective
circulation in Lake Tanganyika. *Limnology and Oceanography* **56**, 910-926.
Verburg P, Hecky RE (2009) The physics of the warming of Lake Tanganyika by climate
change. *Limnology and Oceanography* **54**, 2418-2430.
Verburg P, Hecky RE, Kling H (2003) Ecological consequences of a century of warming in
Lake Tanganyika. *Science* **301**, 505-507.
Vij S, Kuhl H, Kuznetsova IS*, et al.* (2016) Chromosomal-Level Assembly of the Asian
Seabass Genome Using Long Sequence Reads and Multi-layered Scaffolding. *PLOS*
Genetics **12**, e1005954-e1005954.
Wang, J (2017) The computer program STRUCTURE for assigning individuals to
populations: easy to use but easier to misuse. *Molecular Ecology Resources* **17**, 981-
990.
Wenburg JK, Bentzen P, Foote CJ (1998) Microsatellite analysis of genetic population
structure in an endangered salmonid: the coastal cutthroat trout (Oncorhynchus clarki
clarki). *Molecular Ecology* **7**, 733-749.
White JW, Botsford LW, Hastings A, Largier JL (2010) Population persistence in marine
reserve networks: incorporating spatial heterogeneities in larval dispersal. *Marine*
Ecology Progress Series **398**, 49-67.
Young EF, Belchier M, Hauser L*, et al.* (2015) Oceanography and life history predict
contrasting genetic population structure in two Antarctic fish species. *Evolutionary*
Applications **8**, 486-509.
1100

1101

Figure Captions

Figure 1. Sampling site locations and years in which fish were collected at each location (shaded boxes). Circle sizes on the map are proportional to the number of samples in the final genotyping-by-sequencing dataset for all species combined, with a minimum of 1 and maximum of 45 samples collected per site across all years. Dashed lines indicate the general boundaries of the north, central, and southern basin boundaries within the lake. Additional sampling information is available in Table S1 and Table S2.

Figure 2. Relationships among the four endemic *Lates* species in Lake Tanganyika based on the GBS data set, as demonstrated by their (a) phylogenetic relationships, (b) relationships using principal component analysis, (c) genetic clustering using entropy at K=4. The phylogeny has been rooted using *L. calcarifer* as an outgroup and nodal bootstrap support is indicated for the main nodes separating the species. In (a), genetic diversity of the given species, as measured using Watterson's Theta (θ_W) in ANGSD, is indicated. Percentages in (b) indicate that amount of variance explained by the given principal component axis (axes 1-4 shown). In (c) each vertical bar represents an individual, and the colors represent the proportion of ancestry assigned to each of the four species. Individuals are colored by *a posteriori* genetic assignment of species, rather than phenotypic identification.

Figure 3. Principal component analysis of individual species based on the GBS data set, with colors indicating sampling locations, ordered from north to south. For each species, the plot shows the location of each individual on the two most dominant PCA axes, with percent variance explained in parentheses.

Figure 4. Heatmaps visualizing estimates of differentiation (Reich-Patterson F_{ST}) between sampling locations above the diagonal, with observed heterozygosities (H_o) along the diagonal (dark gray backgrounds) for the genotyping-by-sequencing dataset only. The red numbers across the top of each heatmap indicate sample size for the given species at the given sampling site. For F_{ST} values, light gray indicates no differentiation (i.e., bootstrap confidence intervals overlap 0), yellow indicates low differentiation, and red indicates relative high levels of differentiation.

Figure 5. Intraspecific population structure inferred from the GBS data set and entropy analyses at K=2 and K=3 for (A) *L. stappersii*, (B) *L. microlepis*, (C) *L. mariae*, and (D) *L. angustifrons*. Maps show the distribution of individuals among sampling locations and inset plots show discriminant information content (DIC) values for each value of K, with the lowest DIC value for each species indicated with a larger black dot. In colored barplots, each bar corresponds to one individual and colors indicate inferred ancestry from each of the K groups. Plots for values of K=4 to K=6 can be found in Supplementary Fig. S6.

NORTH ● Kagunga ● Kigoma ● North Mahale ● South Mahale ● Ikola ● Mpinbwe ● Kirando ● Wampembe ● Kasanga SOUTH

(A) *Lates stappersii*

PC2 (2.46%)

PC1 (2.48%)

(B) *Lates microlepis*

PC2 (4.1%)

PC1 (4.59%)

(C) *Lates mariae*

PC2 (3.45%)

PC1 (5.28%)

(D) *Lates angustifrons*

PC2 (6.62%)

PC1 (7.98%)

● Kagunga ● Kigoma ● North Mahale ● South Mahale ● Ikola ● Mpinbwe ● Kirando ● Wampembe ● Kasanga

(A) *Lates stappersii*

(B) *Lates microlepis*

(C) *Lates mariae*

(D) *Lates angustifrons*

(A) *Lates stappersii* — Kagunga, Kigoma, North Mahale, South Mahale, Mpimbwe, Kirando, Wampembe, Kasanga

(B) *Lates microlepis* — Kigoma, North Mahale, South Mahale, Kirando

(C) *Lates mariae* — Kagunga, Kigoma, Ikola, Kirando, Wampembe, Kasanga

(D) *Lates angustifrons* — Kagunga, Kigoma, North Mahale, South Mahale, Kirando, Wampembe