The genetic population structure of Lake Tanganyika's *Lates* species flock, an endemic radiation of pelagic top predators

Jessica A. Rick*[1], Julian Junker*[2,3], Ismael A. Kimirei[4], Emmanuel A. Sweke[4,5], Julieth B. Mosille[4], Christian Dinkel[2], Salome Mwaiko[2,3], Ole Seehausen[2,3], Catherine E. Wagner[1]

* denotes equal contribution

[1] Department of Botany and Program in Ecology, University of Wyoming, 1000 E University Dr., Laramie, Wyoming 82072 USA

[2] EAWAG Swiss Federal Institute of Aquatic Science and Technology, CH-6047 Kastanienbaum, Switzerland

[3] Division of Aquatic Ecology & Evolution, Institute of Ecology & Evolution, University of Bern, CH-3012 Bern, Switzerland

[4] Tanzania Fisheries Research Institute (TAFIRI), Dar es Salaam, Tanzania

[5] Deep Sea Fishing Authority (DSFA), Zanzibar, Tanzania

Corresponding authors: Jessica Rick (jrick@uwyo.edu) and Catherine E. Wagner (Catherine.Wagner@uwyo.edu)

Running title: Population genetics of *Lates* fishes

Abstract

Life history traits are important in shaping gene flow within species and can thus determine
whether a species exhibits genetic homogeneity or population structure across its range.
Understanding genetic connectivity plays a crucial role in species conservation decisions,
and genetic connectivity is an important component of modern fisheries management in
fishes exploited for human consumption. In this study, we investigated the population
genetics of four endemic *Lates* species of Lake Tanganyika (*Lates stappersii*, *L. microlepis*,
L. mariae and *L. angustifrons*), using reduced-representation genomic sequencing methods.
We find the four species to be strongly differentiated from one another (mean interspecific
F_{ST} = 0.665), with no evidence for contemporary admixture. We also find evidence for high
levels of genetic structure within *L. mariae*, with the majority of individuals from the most
southern sampling site forming a genetic group distinct from the individuals at other sampling
sites. We find evidence for much weaker structure within the other three species, *L.
stappersii, L. microlepis*, and *L. angustifrons*, despite small and unbalanced sample sizes
and imprecise geographic sampling locations, suggesting the possibility for further structure
undetected in our study. We call for further research into the origins of the genetic
differentiation that we observe in these four species, particularly that of *L. mariae*, which may
be important for the conservation and management of this species.

***Keywords*:** Lates angustifrons, Lates mariae, Lates microlepis, Lates stappersii, population
genetics, genomic differentiation

Introduction

Life history traits, including reproduction, growth and dispersal, strongly influence gene flow and the genetic structure of populations (Ellegren & Galtier 2016; Manier & Arnold 2006; Stearns 1992). Differences in life history traits can also explain why sympatric and ecologically similar taxa may differ in their genetic population structure (Peterman *et al.* 2015; Young *et al.* 2015). Understanding how life history traits like dispersal, reproduction, and growth shape gene flow is therefore crucial to improving our understanding of local adaptation, speciation, persistence of populations, migration, and the spatial scale of effective management (Sunday *et al.* 2014).

In teleost fishes, traits such as spawning location, philopatry, timing of spawning, and duration of larval stage, have been shown to shape population structure (Olsen *et al.* 2011; Pettersson *et al.* 2019; Young *et al.* 2015). Anadromous salmonids are well-known for displaying homing behavior that leads to genetic distinction between populations that use different spawning grounds at different times, dividing populations into many genetic units despite the populations spending most of their lives together in the ocean (Wenburg *et al.* 1998; Brannon *et al.* 2004). Divergence between populations caused by temporal differences in the timing of spawning has also been observed in Atlantic herrings (Martinez Barrio *et al.* 2016; Pettersson *et al.* 2019). In the oceans, coral reef fishes which spawn on or near the substrate, which exhibit parental care strategies, and which lack planktonic larvae, generally have significantly greater population structure than pelagic spawners or substrate spawners with planktonic larvae (Riginos *et al.* 2014). Additionally, a shorter larval phase among pelagic spawners can lead to a more structured population, because this leads to a shorter larval dispersal distance (Selkoe & Toonen 2011).

Understanding the genetic population structure of species is essential for conservation and sustainable fisheries management (Reiss *et al.* 2009; Bernatchez 2016; Supple & Shapiro 2018). Disregarding population structure and managing a population complex as a single population can lead to overfishing or even extinction of the more vulnerable genetic groups (Reiss *et al.* 2009; Sterner 2007). In species with cryptic diversity, unintentional overfishing of distinct populations can lead to a loss of genetic diversity in the species as a whole. Stock diversity can contribute to the resilience of a fishery to environmental fluctuations (Schindler *et al.* 2015) and the loss of differentiated populations can compound to major losses in the consistency of fisheries yields (Hutchinson 2008; Schindler *et al.* 2010). The loss of population diversity can also have drastic effects on ecosystem services and species persistence, and understanding the extent to which population structure exists is crucial for predicting these consequences (Schindler *et al.* 2015; Schindler *et al.* 2010; Therkildsen *et al.* 2013). It is therefore essential to understand the extent and nature of population genetic

structure and differential adaptation in fishes that are important food and economic
resources.

The pelagic fish community and the fishery of Lake Tanganyika

The pelagic fish of Lake Tanganyika are one such fishery. Lake Tanganyika is the second
largest inland fishery on the continent of Africa (Van der Knaap *et al.* 2014). During the last
few decades, fish catches in Lake Tanganyika's fishery have exhibited a general downward
trend (Kimirei *et al.* 2008; Van der Knaap 2013; Van der Knaap *et al.* 2014; van Zwieten *et al.*
2002). This decline has resulted from a combination of an increase in the number of
fishermen and vessels on Lake Tanganyika, changes in fishing practices (e.g., the use of
beach seining, which targets fish in their in-shore nursery habitats) (Kimirei *et al.* 2008; Petit
& Shipton 2012; Van der Knaap 2013; Van der Knaap *et al.* 2014; van Zwieten *et al.* 2002),
and warming lake surface temperatures (Cohen *et al.* 2016; O'Reilly *et al.* 2003; Sayers *et al.*
2020). Decreases in fish abundance are likely also linked to reduced productivity in the lake
caused by stronger water column stratification due to climate change (O'Reilly *et al.* 2003;
Verburg *et al.* 2003). In addition, anthropogenic activity (e.g., land use changes, pollution,
and increased sedimentation) impact both the water quality of Lake Tanganyika directly and
the lake's productivity via climate feedbacks (Ogutu-Ohwayo *et al.* 2016; Ivory *et al.* 2021;
McGlue *et al.* 2021). Local fishermen similarly acknowledge spatial and temporal changes in
fish abundance related to the ecological conditions of the lake (Bulengela *et al.* 2020; De
Keyzer *et al.* 2020). Consequently, there is increasing recognition of the need to develop
sustainable management strategies for the lake's pelagic fish stocks (Kimirei *et al.* 2008;
Mölsä *et al.* 1999; Mölsä *et al.* 2002; Van der Knaap 2013; Van der Knaap *et al.* 2014; van
Zwieten *et al.* 2002).

Lake Tanganyika's pelagic zone is relatively low in species diversity compared to that of
other large African lakes: it consists predominantly of six endemic fish species belonging to
two families (Coulter 1991). These include a monophyletic pair of mainly planktivorous
clupeids (*Stolothrissa tanganicae*, *Limnothrissa miodon*) and four mainly piscivorous latids of
the genus *Lates* (*L. stappersii*, *L. mariae*, *L. microlepis*, *L. angustifrons*). While all four *Lates*
species likely have planktivorous larvae (van Zwieten *et al.* 2016), only *L. stappersii* remains
pelagic throughout its life (Ellis 1978). The other three species move inshore to littoral weed
beds as juveniles (>3cm) before *L. microlepis* returns to the pelagic, *L. mariae* transitions to
the deepwater benthic zone, and *L. angustifrons* can be found throughout the water column
as adults (Coulter 1991). Of the four *Lates* species, *L. stappersii* is the most abundant and an
important component of the current fishery of Lake Tanganyika (Coulter 1976, 1991; Kimirei
et al. 2008; Mannini 1998a, b; Mölsä *et al.* 1999; Mölsä *et al.* 2002; Munyandero 2002;

Sarvala *et al.* 2002; Van der Knaap 2013; Van der Knaap *et al.* 2014; van Zwieten *et al.*
2002).

The fish of Lake Tanganyika are influenced by a complex interplay of bottom-up control via
changes in the supply of nutrient-rich deep waters, driven by different mixing regimes and
upwelling rates (Bergamino *et al.* 2010; O'Reilly *et al.* 2003; Stenuite *et al.* 2007; Verburg &
Hecky 2009; Verburg *et al.* 2003), and top-down control via predators and fishing (Kimirei *et*
al. 2008; Mannini 1998a; Mannini *et al.* 1996; Munyandero 2002; Van der Knaap 2013; Van
der Knaap *et al.* 2014; van Zwieten *et al.* 2002). If spatially heterogeneous environmental
factors, such as mixing regimes and nutrient supply, are consistent over generations and
coupled with limited gene flow between spatially segregated populations, divergent selection
on life history traits (e.g. spawning phenology, developmental timing and recruitment
success) of fish in different parts of the lake could develop and persist. Spatial environmental
variation in Lake Tanganyika, in combination with its large geographic extent, might therefore
generate intraspecific genetic differentiation among populations of pelagic fish, as has been
observed in pelagic cichlids in both Lake Tanganyika and Lake Malawi (Genner *et al.* 2010;
Koblmüller *et al.* 2019). In these cichlids, strong spawning site fidelity (Genner *et al.* 2010)
and differences in foraging behavior (Koblmüller *et al.* 2019) influence spatial genetic
structure. In contrast, recent studies have found that the two pelagic clupeids in Lake
Tanganyika show no evidence of spatial genetic structure (Junker *et al.* 2020; De Keyzer *et*
al. 2019; Kmentová *et al.* 2020). Intraspecific spatial heterogeneity in spawning times
(Coulter 1976; Ellis 1978), dominant diet items (Mannini *et al.* 1999), sizes at maturity
(Mannini et al. 1996), and catch rates (Coulter 1991), along the length of Lake Tanganyika
support the possibility that gene flow may be reduced among fish in different parts of the
lake. Indeed, in *L. stappersii*, a study based on RAPD markers found individuals from the
Kigoma region to be genetically differentiated from the rest of the sites sampled (Kuusipalo
1999), with some additional differentiation observed among individuals collected during the
dry versus rainy season.

Here, we test these predictions using genomic data collected from each of the four *Lates*
species from across their distribution within Lake Tanganyika. We use reduced-
representation genomic sequencing datasets from four endemic *Lates* species—*L.*
stappersii, L. mariae, L. microlepis, and *L. angustifrons*—to examine their genetic
connectivity. We first assessed whether phenotypic identification coincides with genetically
based species boundaries and then examined the extent to which each of the four species is
genetically structured, as well as how the genetic structure observed in each species
corresponds to its known life history and the environment. Based on known differences in life
history among the four species, we hypothesized that we would find little genetic population
structure in the entirely pelagic *L. stappersii* and mainly pelagic *L. microlepis*. In contrast, we

expected the highest genetic differentiation in populations of the predominantly benthic *L.*
mariae. Finally, we hypothesized that *L. angustifrons* would show intermediate amounts of
structure since there is no evidence for large scale movements of this species, even though
they are found throughout the water column.

Material and Methods

Study system and sampling

Between 2001 and 2019, we collected tissue samples (fin clips) from *L. stappersii, L. mariae,*
L. microlepis, and *L. angustifrons* at 9 general locations along the Tanzanian shore of Lake
Tanganyika, which together span the ~490 km shoreline from the northern to southern border
(Fig 1). All fish collections were made in partnership with researchers of the Tanzanian
Fisheries Research Institute (TAFIRI) in Kigoma, Tanzania; most of the fish were obtained
opportunistically from fishermen, and therefore were already dead when we received them.
Fishermen generally fish within a ~20 km radius from their landing sites, but there remains
inherent uncertainty in the exact location and depth at which these fish were collected. Fish
were identified in the field by sampling personnel following United Nations Food and
Agricultural Organization (FAO) identification guidelines (Eccles 1992) and guidance from
Tanzanian researchers and local fishermen, and genetic methods were later used to confirm
species identities for each fish. The majority of fish were collected during two targeted
sampling campaigns in 2017 and 2018, the first of which occurred at the end of the dry
season (September-October 2017), while the second occurred just following the rainy
season (April-May 2018). For fish collected from 2016-2019, we recorded length and weight,
and took standardized pictures of each fish. For the few live fish that were caught as part of
other research, we took cuvette photographs of the live fish and subsequently euthanized the
fish with an overdose of MS222. We then took fin clips for genetic analysis from all fish.
Specimens smaller than 600mm were preserved in formaldehyde and archived in the
collections at TAFIRI (Kigoma, Tanzania), EAWAG (Kastanienbaum, Switzerland), the
University of Wyoming Museum of Vertebrates (Laramie, WY, USA), or the Cornell University
Museum of Vertebrates (Ithaca, NY, USA).

Figure 1. Sampling site locations and years in which fish were collected at each location (shaded boxes). Circle sizes on the map are proportional to the number of samples in the final dataset for all species combined, with a minimum of 1 and maximum of 45 samples collected per site across all years.

Genomic sequencing

We extracted DNA from fin clips using DNeasy Blood and Tissue Kits (Qiagen, Inc.) following the standard protocol, with the addition of an RNAse A incubation step. We then prepared genomic libraries for a combination of genotyping-by-sequencing (GBS, for *L. stappersii, L. mariae, L. microlepis, L. angustifrons*) and restriction site associated DNA sequencing (RAD, for *L. stappersii*).

We prepared GBS libraries following protocols outlined in Parchman *et al.* (2012), using MseI and EcoRI restriction enzymes. Following fragmentation via restriction enzyme digestion, fragmented DNA was barcoded by ligating short, unique individual-identifying DNA fragments (barcodes) to each individual's fragmented DNA. These barcoded fragments were then amplified by PCR and pooled for sequencing. We ran two replicate PCRs for each individual and pooled the final PCR products into two libraries. The prepared libraries were size-selected for 200-350bp fragments using Blue Pippin (Sage Science, MA). One library was sequenced on one lane of Illumina HiSeq4000 (150bp single-end) at the University of Texas

at Austin's Genome Sequencing and Analysis Facility (UT GSAF; Austin, TX) and the other
library was sequenced on one lane of Illumina HiSeq4000 (150bp, single-end) at the
University of Oregon's Genomics and Cell Characterization Core Facility (Eugene, OR). Each
library contained 192 individuals, and select individuals were duplicated across libraries to
check for library compatibility.

The RAD libraries containing the *L. stappersii* were prepared for sequencing following the
protocol by Baird *et al.* (2008) with the following modifications: we used between 400ng and
1000ng genomic DNA per sample and digested with *SbfI* overnight. We tagged each
individual using P1 adapters (synthesized by Microsynth) with custom six to eight base pair
barcodes and multiplexed 67 barcoded individuals per library. The libraries were sheared
using an S220 series Adaptive Focused Acoustic (AFA) ultra-sonicator (Covaris) with the
manufacturer's settings for a 400 bp mean fragment size. We size-selected for fragments
between 300 and 700bp using a sageElf (Sage Scientific Electrophoretic Lateral
Fractionator; Sage Science, Beverly, MA). The enrichment step was done in 6 aliquots with a
total volume of 200 µl. Volumes were combined prior to the final size selection (again 300-
700bp) step using the sageELF. Sequencing was done by the Lausanne Genomic
Technologies sequencing facilities (University of Lausanne, Switzerland). All three libraries
were single-end sequenced on one lane each of the Illumina HiSeq2000 (100bp SE).

Sequence data preparation

We filtered raw sequencing reads from each library by first removing the reads derived from
the PhiX genome and other common contaminants (e.g. excess barcodes, primers, and
adapters) using bowtie2 (v2.0.0, Langmead & Salzberg 2012). We filtered reads for an intact
enzyme restriction site (*SbfI* for RAD libraries, MseI and EcoRI for GBS libraries), de-
multiplexed the fastq files, and matched sequence barcodes to individual fish using custom
perl and bash scripts.

For the GBS libraries, the reads were mapped to the chromosome-level *Lates calcarifer*
reference genome (v3, NCBI Genome Assembly GCA_001640805.1; Vij *et al.* 2016) using
bwa mem (v0.7.17; Li & Durbin 2009) with default settings. Following alignment, we excluded
any individual with < 50,000 reads or less than 60% of raw reads assembled to the reference
genome. We identified variable sites (i.e. single nucleotide polymorphisms; SNPs) in the
assembly using SAMtools mpileup (v1.8; Li *et al.* 2009) and bcftools (v1.8; Li *et al.* 2009). In
calling variable sites, we omitted indels and kept only high-quality biallelic variant sites
(QUAL > 20 and GQ > 9). We then filtered SNPs by minor allele frequency and amount of

missing data using vcftools (Danecek *et al.* 2011), additionally only calling genotypes with a
minimum read depth of 5. We then selected only SNPs with a minor allele frequency greater
than 0.01 that were present in at least 50% of individuals. Finally, we thinned each data set
to retain one SNP per locus (-thin 90). We assessed the distribution of missing data across
individuals and mean read depth across sites using vcftools (Danecek *et al.* 2011),
additionally using whoa (v0.0.2.999, Anderson 2018) in R (v4.0.4, R Core Team 2021) to
assess heterozygote miscall rate at low read depth sites to ensure that our read depth filter
was stringent enough.

For the RAD libraries, we barcode-trimmed reads down to 84 nucleotides using
process_radtags from Stacks (Catchen *et al.* 2013). The FASTX-toolkit v.0.0.13
(http://hannonlab.cshl.edu/fastx_toolkit/) was used for quality filtering. In a first step, we kept
only reads with all base quality scores greater than 10; in a second step, we removed all
reads with more than 5% of the bases with quality score below 30. The RAD library was then
mapped to the *Lates calcarifer* reference genome (Vij et al. 2016) and filtered, following the
same steps as for the GBS libraries above (MAF > 0.01, missing data < 0.5, one SNP per
locus, read depth > 5). There was no overlap of individuals included in the RAD and GBS
sequencing libraries.

After filtering variants, we calculated the distribution of minor allele reads at heterozygous
sites for each individual in both the RAD and GBS libraries, using a custom bash script
(available at http://github.com/jessicarick/lates-popgen). If there is an excess of sequencing
errors or contamination for an individual, then we would expect the ratio of minor allele to
major allele reads to be significantly less than 1:1. Thus, we plotted the distribution of minor
allele reads at heterozygous sites for each individual and removed individuals with a
deficiency in minor allele reads across the majority of heterozygous sites. After removing
these individuals, we then re-called genotypes and re-filtered the SNP data set to produce
our final SNP data set.

Species assignment

Not all samples could be identified to species in the field. Some fish were caught deep and
brought quickly to the surface by the fishermen, making them difficult to identify.
Furthermore, juveniles of the large species *L. microlepis*, *L. mariae* and *L. angustifrons* can
be difficult to distinguish. We used three different lines of evidence to confirm species
identification and ensure that morphological identification matched genetic assignment: (1)
phenotypic identification based on FAO species descriptions; (2) individual ancestry

assignment using a model-based clustering approach; and (3) inference of monophyletic
groups in phylogenetic analysis. The details for each of these steps are described below.

With the GBS data, we first visualized clusters in our data using principal component analysis
using EMU (Meisner *et al.* 2021). EMU uses an iterative method to infer population structure
in the presence of missing data and is therefore able to infer population structure even for
datasets with high, uneven, or non-random proportions of missing data, which tend to
produce bias in other PCA methods (Meisner *et al.* 2021). We used the EM acceleration
method in EMU, using 10 eigenvectors for optimization (-e 10) with a maximum of 100
optimization iterations and keeping all eigenvectors as output. We then inferred species
groups using the model-based genetic clustering program entropy (Gompert *et al.* 2014;
Shastry *et al.* 2021). Entropy infers population structure from multilocus SNP data by
assigning individuals partially or completely to groups using Bayesian estimation from
genotype likelihoods. In taking uncertainty about individual genotypes into account via
genotype likelihoods, the model integrates outcomes over genotype uncertainty. We
converted our VCF containing sites with <50% missing data to the mpgl genotype likelihood
format using the vcf2mpgl R script (available at http://github.com/jessicarick/lates-popgen;
adapted from https://bitbucket.org/buerklelab/mixedploidy-entropy/src, v2.0). We then ran
entropy for K=4 clusters, running three independent MCMC chains of 80,000 total steps,
discarding the first 10,000 steps as burn-in, and retaining every 10^{th} value (thin=10), resulting
in 7000 samples from the posterior distribution of each chain. Our MCMC chains were run in
parallel using GNU parallel (Tange 2011). We checked MCMC chains for mixing and
convergence of parameter estimates by plotting a trace of the MCMC steps. We then
visualized assignments in R and assigned individuals to species based on group assignment
probabilities (q).

We further corroborated species identities using maximum likelihood phylogenetic inference
methods in RAxML (v8.1.17; Stamatakis 2014). We concatenated all SNPs, removed
invariant sites using the raxml_ascbias python script (v1.0, from
https://github.com/btmartin721/raxml_ascbias), and used the Lewis correction for invariant
sites with the ASC_GTRGAMMA model of molecular evolution (Lewis 2001) within RAxML to
infer the maximum likelihood phylogeny, including *L. calcarifer* (Vij et al. 2016) as the
outgroup. We used 100 rapid bootstraps to estimate confidence in our maximum likelihood
tree. We then visually identified monophyletic groups on our maximum likelihood tree and
used these groupings to further verify species identities from entropy and PCA clusters.

Genetic diversity

We calculated genetic diversity for each species in ANGSD (v0.931, Korneliussen *et al.*
2014). From the BAM alignment files, we first calculated the site allele frequency likelihoods
based on individual genotype likelihoods (option -doSaf 1) using the samtools model (option -
GL 1), with major and minor alleles inferred from genotype likelihoods (option -doMajorMinor
1) and allele frequencies estimated according to the major allele (option -doMaf 2). We
filtered sites for a minimum read depth of 10, minimum mapping quality of 20, and minimum
quality (*q*-score) of 20. From the site allele frequency spectra, we then calculated the
maximum-likelihood estimate of the folded site frequency spectra (SFS) using the ANGSD
realSFS program (with option -fold 1). The folded SFS were then used to calculate genome-
wide genetic diversity (Watterson's θ), using the ANGSD thetaStat program (Korneliussen *et*
al. 2013).

Population genetic structure and isolation-by-distance

For each species, we then re-called variants to create a species-specific set of SNPs from
the GBS data. As with the species combined GBS data, we identified variable sites using
SAMtools mpileup and bcftools, omitting indels and keeping only high-quality biallelic variant
sites (QUAL > 20 and GQ > 9). We then filtered SNPs by minor allele frequency (> 0.01) and
amount of missing data (<50% missing) using vcftools (Danecek et al. 2011), additionally
only calling genotypes with a minimum read depth of 5. Finally, we thinned to one SNP per
locus (-thin 90).

For each species, we then performed principal component analysis (PCA) using EMU
(Meisner *et al.* 2021), after using PLINK (v1.90b6.9, Purcell *et al.* 2007; Chang *et al.* 2015) to
convert our VCF into BED format. EMU is able to infer population structure even for datasets
with high, uneven, or non-random rates of missing data, which tend to produce bias in other
PCA methods (Meisner *et al.* 2021). We again used the EM acceleration method, using 10
eigenvectors for optimization (-e 10) with a maximum of 100 optimization iterations and
keeping all eigenvectors as output. We also performed PCA in EMU on the RAD dataset for
L. stappersii.

We estimated heterozygosity at each site using the dartR package in R (v1.9.9, Gruber *et al.*
2018). We then used the Reich-Patterson F_{ST} estimator (Reich *et al.* 2009), which is
consistent even with small sample sizes, to calculate genetic divergence between sampling
sites within each species. We calculated Reich-Patterson F_{ST} estimates using a custom
function in R (available at http://github.com/jessicarick/reich-fst), using functions from the
vcfR (v1.12, Knaus & Grunwald 2017) and dartR packages for importing and manipulating

our VCF. We then tested for isolation-by-distance (IBD) using a Mantel test (mantel.randtest

from ade4 in R; v1.7-16, Dray & Dufour 2007) to calculate the correlation between the natural

logarithm of Euclidean geographic distance and standardized genetic distance ($F_{ST}/1-F_{ST}$) for

all pairwise combinations of sampling sites within each species. We chose to pool these tests

by sampling site, as we do not have more specific locations for where most of the fish were

caught.

To test for intraspecific population subdivision more formally, we again used entropy. For

each species, we ran entropy for K=1 to K=6 to infer the most likely number of ancestral

groups within each species, and to assign individuals to these clusters. We ran three

independent MCMC chains of 100,000 total steps, discarding the first 10,000 steps as burn-

in, and retaining every 10^{th} value (thin=10), resulting in 9000 samples from the posterior

distribution of each chain. We checked MCMC chains for mixing and convergence of

parameter estimates by plotting a trace of the MCMC steps. We then calculated the deviance

information criterion (DIC) for each value of K and used the model with the lowest DIC as our

best for modeling the variation observed in our data. From our entropy results, we assigned

individuals to intraspecific groups using a cutoff of q > 0.6 at each value of K. We then

calculated the mean of Reich-Patterson F_{ST} estimates between these K groups to understand

how genetically distinct these groups are from one another. We also ran entropy at K=2 and

K=3 using datasets with a more stringent missing data filter (miss=0.9) to test whether group

assignments were contingent on the choice of bioinformatic filters. We also calculated Reich-

Paterson F_{ST} estimates between each pair of individuals within each species in R, to facilitate

comparisons that do not rely on group assignments. With the group assignments and

individual pairwise F_{ST} estimates, we statistically tested and graphically visualized

relationships between these genetic measures and sampling year, sampling time of year, fish

standard length, and juvenile status, to examine hypotheses about heterogeneity in

spawning location and timing.

Results

The GBS libraries yielded an average of 245 million reads across 200 individuals. The RAD

libraries containing the *L. stappersii* individuals yielded an average of 271 million reads,

including 2.5% - 8.6% bacteriophage PhiX genomic DNA. On average, the mapping rate for

L. stappersii individuals' GBS reads to the *L. calcalifer* reference genome was 89.8% (RAD,

89.6%), and mapping rates averaged 91.6% for *L. microlepis*, 91.3% for *L. mariae* and

88.7% for *L. angustifrons* individuals.

These mapping rates resulted in 1,463,702 unfiltered variable sites in *L. stappersii*,
1,286,810 in *L. microlepis*, 961,153 in *L. mariae* and 764,456 in *L. angustifrons*. After filtering
for missing data (< 50%), minor allele frequency (MAF > 0.01), read depth (minDP > 5), and
keeping only one SNP per locus (-thin 90), our species-specific GBS data sets contained
38,160 SNPs from 58 *L. stappersii* individuals (RAD, 16,802 SNPs from 63 different
individuals), 33,429 SNPs from 34 *L. microlepis* samples, 30,431 SNPs from 38 *L. mariae*
samples and 19,519 SNPs from 20 *L. angustifrons* samples. The data set with all four
species contained 4,999,192 unfiltered and 44,823 filtered SNPs. None of the four species
exhibited heterozygote excess that would be expected if many sites were erroneously called
as heterozygous, and estimates of miscall rates are low for all read depths > 2 (Fig. S1). The
majority of individuals who were lost were filtered out due to an imbalance in the reads
supporting major and minor alleles at heterozygous loci. In our final dataset, the mean read
depth was 15.2 for *L. stappersii*, 23.1 for *L. microlepis*, 20.9 for *L. mariae*, and 25.8 for *L.*
angustifrons (Fig. S2).

Species assignment

The species-combined SNP dataset for GBS data contained 200 individuals, of which we
removed 50 individuals due to low numbers of reads (n=2), low assembly rate to the
reference genome (n=4), or highly unbalanced read ratios at heterozygous sites (n=44),
which is indicative of contamination. The remaining individuals all had > 50,000 reads that
aligned to the *L. calcarifer* reference genome. The SNP alignment for RAxML contained
48,776 SNPs after removing invariant sites. The rooted maximum likelihood phylogeny
(rooted on *L. calcarifer*; Fig. 2A) shows that each of the four species is reciprocally
monophyletic. In addition, we infer a sister taxon relationship between *L. mariae* and *L.*
angustifrons, and a sister taxon relationship between *L. stappersii* and *L. microlepis*
(conflicting with the best supported topology in Koblmüller *et al.* 2021); however, short
internode distances between these splits and low bootstrap support for the initial split
suggest that all three speciation events may have occurred within a short period, and more
investigation of the divergence history of the clade is warranted but not the focus of this
current work. The PCA based on the GBS data was consistent with the species affinities
indicated by monophyletic groups in RAxML (Fig. 2). The first genetic PC axis (27.3%)
separates *L. stappersii* from the remaining three species. The second PC axis (12.4%) then
separates *L. mariae* and *L. microlepis* from one another, and the third axis (4.8%) separates
L. angustifrons from the other three species. Using entropy at K=4, we found each of the four
species to make a distinct genetic cluster (Fig. 2C). Based on these combined analyses, 58
individuals were assigned to *L. stappersii*, 34 to *L. microlepis*, 38 to *L. mariae*, and 20 to *L.*

angustifrons. While the genetic approaches revealed consistent species identifications, we
had phenotypically misidentified one *L. microlepis*, one *L. mariae* and three *L. angustifrons*
(Table S1).

Figure 2. Relationships among the four endemic *Lates* species in Lake Tanganyika based on the GBS
data set, as demonstrated by their (a) phylogenetic relationships, (b) relationships using principal
component analysis, (c) genetic clustering using entropy at K=4. The phylogeny has been rooted
using *L. calcarifer* as an outgroup and nodal bootstrap support is indicated for the main nodes
separating the species. In (a), genetic diversity of the given species, as measured using Watterson's
Theta (θ_W) in ANGSD, is indicated. Percentages in (b) indicate that amount of variance explained by
the given principal component axis (axes 1-4 shown). In (c) each vertical bar represents an individual,
and the colors represent the proportion of ancestry assigned to each of the four species. Individuals
are colored by *a posteriori* genetic assignment of species, rather than phenotypic identification.

Population structure

All four species had low but similar levels of genetic diversity (overall θ_W = 0.00222), and
genetic diversity was elevated in *L. stappersii* (θ_W = 0.00230) compared to the three other
species (*L. microlepis*, θ_W = 0.00094; *L. mariae*, θ_W = 0.00092; *L. anguistifrons*, θ_W = 0.00064;
Fig 2A). Genetic diversity for the RAD dataset in *L. stappersii* was similar to that estimated
for the GBS data (RAD θ_W = 0.00223). Reflecting the distinctiveness seen in the all-species
PCA and phylogeny, F_{ST} values are high between species (mean Reich-Patterson F_{ST} =
0.607; Table S2). None of the four species shows a significant relationship between genetic
and geographic distance, suggesting an overall lack of isolation by distance (Fig. S3; all
Mantel test *p*-values > 0.2).

The PCA for *L. stappersii* based on GBS data (Fig 3A) did not reveal any clear genetic
clusters. We similarly did not see any clustering in the RAD data for *L. stappersii* (Fig. S4). In

entropy, K=1 was the best fit for the GBS data according to DIC (Fig. 5A); however, at K=2, some individuals from Kigoma, North Mahale and South Mahale form a distinct genetic cluster, and at K=3, groups appear that co-occur at all sampling sites. These groups at K=2 and K=3 were largely but not completely stable (i.e., the same individuals were clustered together) at a higher threshold for missing data (miss=0.5 vs. miss=0.9 had 18 individuals switching groups at K=2 and 10 individuals at K=3; Fig. S7, S8). This weak genetic structure (between-cluster K=2 Reich-Patterson F_{ST} = 0.00205, 95% bootstrap CI: 0.00084-0.00346, Table S3) was also partly reflected in F_{ST} values between sampling sites, where F_{ST} values of comparisons between individuals of South Mahale or North Mahale with samples from other sites were elevated, albeit small ($\leq$ 0.012, Fig. 4A); however, only the F_{ST} values between North Mahale and each of Mpinbwe and Kagunga had 95% bootstrap confidence intervals not overlapping zero (Fig. S9A). The RAD data showed similar weak structure at K=2 (Fig. S5; between-cluster F_{ST} = 0.00146, 95% bootstrap CI: 0.00087-0.00207), but these groups were found sympatrically at all five sampling sites, similar to the groups at K=3 for the GBS data. At K=3, F_{ST} values between groups for the RAD data had bootstrap confidence intervals overlapping zero (Table S2).

Figure 3. Principal component analysis of individual species based on the GBS data set, with colors indicating sampling locations, ordered from north to south. For each species, the plot shows the location of each individual on the two most dominant PCA axes, with percent variance explained in parentheses.

Figure 4. Heatmaps visualizing estimates of differentiation (Reich-Patterson F_{ST}) between sampling locations above the diagonal, with observed heterozygosities (H_o) along the diagonal (dark gray backgrounds). The red numbers across the top of each heatmap indicate sample size for the given species at the given sampling site. For F_{ST} values, light gray indicates no differentiation (i.e., bootstrap confidence intervals overlap 0), yellow indicates low differentiation, and red indicates relative high levels of differentiation.

The PCA for *L. microlepis* did not reveal clear genetic clusters but it did highlight several individuals from the sampling sites Kigoma, North Mahale, and South Mahale, that were differentiated from the other samples on PC1 and PC2 (Fig. 3B). These differentiated individuals were not those with high amounts of missing data (Fig. S10B). All of the sampling sites had low differentiation (mean F_{ST} = 0.00471; Fig. 4B) but all pairwise comparisons had 95% confidence intervals that did not overlap zero except for the Kirando-Kigoma and Kirando-South Mahale comparisons (Fig. S9B). In entropy, K=1 was found to be the best fit using DIC (Fig. 5B); at K=2, four individuals collected in Kigoma are differentiated from the others, albeit with low genetic differentiation and a between-group F_{ST} estimate that overlaps zero (Reich-Patterson F_{ST} = 0.00074, 95% CI: -0.00389 - 0.00549; Table S3). At K=3, the additional group is found at all four sampling sites; the differentiation between the two largest groups is low, but significant (Reich-Patterson F_{ST} = 0.0030, 95% CI: 0.00155 - 0.00474), while the F_{ST} estimates between each of these groups and the smaller group identified at

K=2 had 95% bootstrap confidence intervals overlapping zero. These groups at K=2 and K=3

were largely stable at a higher threshold for missing data (miss=0.5 vs. miss=0.9 had 4

individuals switching groups at K=2 and 8 individuals at K=3; Fig. S7, S8).

The PCA for *L. mariae* clusters three individuals from Kasanga separate from all other

individuals on the first PC axis (explaining 5.28% of the variance), while the second PC axis

separates out some individuals from Kigoma, Wampembe, and Kirando from the others (Fig.

3C). The differentiation among individuals on both PC1 and PC2 does not correlate with the

individuals' amount of missing data (Fig. S10C). In entropy, K=2 has the lowest DIC

estimate, where the three individuals from Kasanga form a distinct cluster (Reich-Patterson

F_{ST} between entropy-identified groups = 0.119, 95% bootstrap CI: 0.113-0.127; Fig 4C and

Table S3). At K=3, most of the other sampling sites have individuals assigned to both of the

non-Kasanga clusters, although these two non-Kasanga clusters have very low

differentiation (F_{ST} = 0.00359, 95% bootstrap CI: 0.00180-0.00623) compared to that

between the Kasanga cluster and each of the other two clusters (mean F_{ST} = 0.120). These

groups at K=2 and K=3 were largely stable at a higher threshold for missing data (miss=0.5

vs. miss=0.9 had no individuals switching groups at K=2 and 15 individuals at K=3; Fig. S7,

S8). The F_{ST} estimates between all fish sampled from Kasanga and other sampling sites are

all significantly different from zero (Fig. S9C). All other between-site F_{ST} estimates had 95%

bootstrap confidence intervals overlapping zero.

For *L. angustifrons*, the PCA did not reveal any genetic structure (Fig. 3D) and the clustering

model with K=1 is the best fit for our data (Fig. 5D). Clusters identified at higher levels of K

had low differentiation (Table S3; K=2, F_{ST} = 0.00498; K=3, mean F_{ST} = 0.0228), and were

found co-occurring at most sampling sites (Fig. 5D). These groups at K=2 and K=3 were

largely but not completely stable at a higher threshold for missing data (miss=0.5 vs.

miss=0.9 had 4 individuals switching groups at K=2 and 7 individuals at K=3; Fig. S7, S8). In

addition, only three pairwise F_{ST} estimates between sampling sites were significant:

Kagunga-North Mahale, Kigoma-Kirando, and North Mahale-Wampembe (Fig. S9D);

however, even these sites had low differentiation (mean F_{ST} between sites = 0.0132), and it

should be noted that North Mahale only has one fish sampled, so these estimates may not

be indicative of patterns at the site as a whole.

Figure 5. Intraspecific population structure inferred from the GBS data set and entropy analyses at K=2 and K=3 for (A) *L. stappersii*, (B) *L. microlepis*, (C) *L. mariae*, and (D) *L. angustifrons*. Maps show the distribution of individuals among sampling locations and inset plots show discriminant information content (DIC) values for each value of K, with the lowest DIC value for each species indicated with a larger black dot. In colored barplots, each bar corresponds to one individual and colors indicate inferred ancestry from each of the K groups. Plots for values of K=4 to K=6 can be found in Supplementary Fig. S6.

We find no relationship between size difference and extent of differentiation (pairwise F_{ST} or
genetic cluster membership) in *L. mariae, L. microlepis,* or *L. angustifrons*. In *L. stappersii*, a
weak positive relationship exists between the difference in standard length and the extent of
genetic differentiation (F_{ST}) between individuals, such that individuals that are more different
in size are also more differentiated genetically (Fig. S12). In *L. microlepis*, we found no
relationship between genetic cluster membership and juvenile status at K=2 or K=3 when
examining only the fish collected in Kigoma, the only site with large enough numbers for
doing this analysis. We additionally find no systematic difference between F_{ST} estimates
between only juvenile fish and those estimated between only adult fish for *L. microlepis* or *L.*
mariae (Fig. S13, S14). When only looking at fish collected at the same site (Fig. S14), we
find that juvenile-adult comparisons had higher F_{ST} values in *L. mariae* than adult-adult
comparisons (two-tailed t-test: t = 2.3484, df = 28, p-value = 0.02615), but did not differ from
juvenile-juvenile comparisons (two-tailed t-test: t = -1.7863, df = 31.781, p-value = 0.0836),
and juvenile-juvenile and adult-adult comparisons were not significantly different from one
another (two-tailed t-test: t = 1.9658, df = 151, p-value = 0.05116). In *L. angustifrons*, F_{ST}
values among juveniles at the same site were lower than those between juveniles and adults
at the same site (two-tailed t-test: t = -3.8334, df = 16.646, p-value = 0.001376) or between
adults at the same site (two-tailed t-test: t = 2.2254, df = 27.295, p-value = 0.0345), while F_{ST}
estimates did not differ significantly between juvenile-juvenile and adult-adult comparisons
(two-tailed t-test: t = -1.6493, df = 14.084, p-value = 0.1212). In *L. microlepis*, F_{ST} estimates
for fish from the same site were marginally higher in juvenile-juvenile pairs than in adult-adult
pairs (two-tailed t-test: t = 2.2525, df = 19.005, p-value = 0.03631), but did not differ between
each of these categories and juvenile-adult comparisons (p-value > 0.07). We do not find any
evidence for juveniles collected at the same time of year to be more closely related than
juveniles collected during different parts of the year in *L. microlepis* (two-tailed t-test: t = -
1.2229, df = 378.64, p-value = 0.2221; Fig. S15), and do not have enough data to analyze
this relationship in the other three species.

Discussion

Differences in life history characteristics can have significant influence on differences in
population genetic structure among taxa, even in closely related species. In this study, we
analyzed population genetic structure within the four endemic pelagic top predators of Lake
Tanganyika, *Lates spp.*, using reduced-representation genomic sequencing data sets. From
our samples collected at discrete sampling sites along the length of the lake, we find three

main patterns: first, we find a strongly differentiated genetic group in *Lates mariae* at the

most southern sampling site, Kasanga; second, we find no additional pattern of isolation-by-

distance or geographic population structure within any of the four *Lates* species; and third,

we find weak evidence for weakly differentiated genetic groups that exist in sympatry at the

majority of sites in all four species. While numerous explanations could account for a lack of

geographic population structure and the existence of sympatric weakly differentiated groups,

more research is needed to understand mechanisms causing the population structure

identified in the different species. These results have potential implications for understanding

the conservation and management of these fishes, with implications for this regionally

important fishery.

Strong genetic differentiation in southern Lates mariae

In *L. mariae*, the majority of samples from the most southern site, Kasanga, are genetically

differentiated from all other samples. The magnitude of F_{ST} estimates between these distinct

individuals and the other *L. mariae* individuals (F_{ST} ~0.135), suggests unexpectedly strong

intraspecific genetic structure given a lack of obvious geographic barriers. Unfortunately, our

sampling does not include individuals from this furthest south site for *L.* microlepis or *L.*

angustifrons, so we cannot determine whether this pattern is also reflected in the other two

large *Lates* species.

It is plausible that the differentiation in *L.* mariae may be driven by known limnological

patterns that differ between the northern and southern portions of the lake. In particular, the

southern part of the lake experiences intense upwelling of cold, nutrient rich waters from the

deep during the dry season (Verburg *et al.* 2011), which mixes with oxygenated water from

the surface and consequently has high nutrient concentrations and primary productivity rates,

which in turn fuels zooplankton growth (Plisnier *et al.* 2009; Loiselle *et al.* 2014). In contrast,

the northern part of the lake has a more stable, persistently stratified, and clear water column

(Plisnier *et al.* 1999; Bergamino *et al.* 2010; Mziray *et al.* 2018). As a result, there is

asynchronous variability between environments in the north and south (i.e., higher primary

productivity in the south than in the north of the lake), as well as high environmental

variability over the course of the year in the south (i.e, higher primary productivity in the dry

season compared to the rainy season). Spatial variability in conditions affects zooplankton

community composition, with shrimps and calanoid copepods prevailing in the south (Kurki *et*

al. 1999), while cyclopoid copepods and medusa dominate in the north (Kurki *et al.* 1999;

Cirhuza and Plisnier 2016). Increases in primary productivity generally lead to high densities

of zooplankton, as well as *Stolothrissa* and *Limnothrissa* sardines (Mulimbwa *et al.* 2014), as

the blooms seem to positively affect sardine spawning and recruitment, although recruitment

pulses sometimes occur during times of poor nutrient conditions as well. Mannini *et al.*
(1999) found corresponding spatial variability in diet for *L. stappersii*, but spatial variability in
diet has not been documented for *L. mariae* or the other large-bodied species. While adult *L.*
mariae do not depend on the pelagic sardines as a primary food source (Coulter 1976), it is
possible that such temporal nutrient variability leads to variability in *L. mariae* prey (benthic
cichlids) as well.

Such a spatially and temporally variable environment might lead to local differential selection
in the southern part of the lake, and this selection combined with restricted movement of
individuals between this and other populations of *L. mariae* could lead to the genetic
differentiation in individuals from Kasanga. *L. mariae* abundance is greater in the south than
in the north (Coulter 1970; I. Kimerei, E. Sweke, personal observation), which suggests that
spatial variation in environmental conditions may underlie differences in fitness or population
growth. A variable environment may also lead to within-site differentiation among individuals
caught in Kasanga if the two groups were to adapt to this variability in different ways. Another
possibility is that one group is adapted to remaining in Kasanga, while the other disperses
further throughout the lake. While all *L. mariae* individuals sampled from Kasanga were
mature and *L. mariae* is believed to be the least vagile of the four *Lates* species (Coulter
1976), it is not clear how vagile these individuals typically are; it is possible that some
individuals remain resident while others migrate, as is the case in Atlantic cod (Kirubakaran
et al. 2016; Berg et al. 2017). To investigate these possibilities, future studies should expand
sampling in the southern portion of the lake, including sampling various life stages of this and
other *Lates* to understand the genetic composition of the southern populations and any
important ecological implications of genetic differentiation.

It is also possible that the differentiation in the most southern part of Lake Tanganyika in *L.*
mariae is related to the lake's bathymetry and differences in the depth of the oxygenated
layer, as suggested by Coulter (1976), who observed asynchronous changes in local
abundance between *L. mariae* populations in the southwest versus southeast arms of the
lake. From catch rates, Coulter (1970) observed that southern populations of at least two of
the *Lates* species, *L. angusitfrons* and *L. mariae*, were likely distinct, as they were heavily
exploited in the southeast arm with no apparent effect on catches in the southwest arm.
These southern reaches of the lake are relatively shallow, and the oxygenated layer extends
up to 200m deep, which allows for a large area of oxygenated bottom to be available to
benthic species (Coulter 1976).

Despite the life history differences between pelagic cichlids and the *Lates* species, the
genetic patterns observed in *L. mariae* are similar to those observed previously in the
benthopelagic cichlids. For the four benthopelagic *Diplotaxodon* species of Lake Malawi with

genetically structured populations, research by Genner *et al.* (2010) indicated spawning site
fidelity drives population structure. In contrast, Koblmüller *et al.* (2019) hypothesized that
preying upon benthic cichlids—which requires shorter-distance movements than preying on
pelagic prey—can explain the genetic structure in the *Bathybatini* of Lake Tanganyika.
Similar to the *Bathybatini, L. mariae* hunts benthic cichlids and is described as relatively
resident (Coulter 1976). However, our results did not recover any evidence for significant
isolation-by-distance in *L. mariae*, suggesting that the genetic structure that we observe is
likely not the result of dispersal limitation. However, females of *L. mariae* do aggregate over
specific spawning grounds (Coulter 1991), similar to *Diplotaxon*. If samples of *L. mariae* were
taken over spawning grounds, it is possible that results would reveal clearly distinct
populations as would be expected from strong spawning site fidelity and differentiation
between these stocks. However, the sampling methods used in our study (i.e., obtaining fish
from fishermen) limit our knowledge of whether fish were collected over spawning grounds.
In addition, these sampling methods mean that a range of ages and reproductive maturities
of fish were included in our study. To our knowledge, the location of spawning grounds has
not been documented for any of the four *Lates* species. In addition, it is unclear how the
pelagic phase of eggs, larvae, and small juveniles (<2.5cm), of *L. mariae* influences dispersal
in the species, as spawning pelagically (Riginos *et al.* 2014) and having a long pelagic larval
stage (Selkoe & Toonen 2011; Shanks 2009) generally facilitate gene flow and reduces
population structure in marine taxa (but see Bay *et al.* 2006; Riginos *et al.* 2011), although
these patterns are complicated by specific behavior exhibited by both adults and larvae.
Future studies in *L. mariae* should sample spawning fish and work to understand the location
of spawning grounds to further assess whether the intraspecific divergences that we observe
could arise from spawning site fidelity.

No pattern of isolation-by-distance or geographic population structure

Although the samples spanned the ~550 km Tanzanian shoreline of Lake Tanganyika, none
of the genetic differences between populations relates to geographic distance, suggesting no
isolation-by-distance in any of the four species. In addition, admixture models with only a
single panmictic population fit our data the best in all species except for *L. mariae* (Fig. 5).
This suggests that individuals of all four species are mobile enough that gene flow is not
restricted among our sampling sites, other than the distinct Kasanga population.

Deviations from isolation-by-distance are common among marine animals, where
geographically proximate populations often have greater population differentiation than
populations further apart, a phenomenon sometimes referred to as "chaotic patchiness"
(Johnson and Black 1984). This phenomenon has been attributed to larval dispersal via

ocean currents (White et al. 2010), and it is possible that currents within Lake Tanganyika
may similarly disperse *Lates* eggs and larvae prior to their movement to inshore habitats.
While it is presumed that all four *Lates* species spawn pelagically (Ellis 1978; Coulter 1991)
and the larvae of the larger three species are planktonic until ~3cm in size (Coulter 1976),
spawning behavior, spawning location within the lake, and the length of time that larvae may
be carried around by lake currents remain unknown for all four species. Lake currents vary
throughout the year (Plisnier *et al.* 1999; Verburg *et al.* 2011), and therefore understanding
the timing and location of reproduction for all four species will be important to understanding
the role that these currents may play in dispersing planktonic eggs and larvae.

While geographically widespread populations are not particularly surprising given the large
body sizes and long-distance swimming capabilities of these fishes, including evidence for
long-distance movements in *L. stappersii* and *L. microlepis*, it conflicts with evidence of a
resident lifestyle in *L. mariae* (Chapman 1976; Coulter 1976, 1991; Muller *et al.* 2001) and
evidence for some spatial structure in the Kigoma area in a previous population genetic study
of *L. stappersii* (Kuusipalo 1999). It should also be noted that our sampling is unbalanced
among sites and lacking in the most southern reaches of the lake for *L. angustifrons* and *L.*
microlepis. It is therefore possible that more (albeit still likely weak) genetic structure exists
than we have been able to detect with available samples. That being said, Koblmüller *et al.*
(2019) similarly found no evidence of genetic population structure in two eupelagic
bathybatine cichlid species of Lake Tanganyika, Koblmüller *et al.* (2015) found no evidence
for poulation structure in the pelagic cichlid *Boulengerochromis microlepis*, and De Keyzer *et*
al. (2019) and Junker *et al.* (2020) found no population structure in either of the pelagic
sardine species. There are many life history differences between the sardines, cichlids, and
Lates that occur sympatrically in the open waters of Lake Tanganyika, but these results
together suggest that spatial environmental heterogeneity along the length of the lake is not
sufficient to restrict gene flow and produce geographically isolated or clinal populations in
these diverse species.

Sympatric, weakly-differentiated groups exist in all four species

While our admixture models suggest panmixia as the best model in *L. stappersii, L.*
microlepis, and *L. angustifrons*, and two distinct groups in *L. mariae*, we find groups with
restricted gene flow in all four *Lates* species that co-occur at the same sampling sites at
higher levels of K. Model DIC values suggest that these higher levels of division are not the
best fit for our data and differentiation is weak between groups (mean F_{ST} = 0.032; Table S3),
but individual assignments do not behave as we would expect if we were to increase the
number of groups in a truly panmictic populations. In our data, we see individuals being

assigned with high confidence to one group or another (i.e., each bar is mostly a single color
in Fig. 5 and Fig. S6); if the population were truly panmictic, then we would expect to see all
individuals showing genetic contributions from all groups. For that reason, we believe these
weakly differentiated groups may be biologically relevant, albeit weakly differentiated. These
groups do not appear to correspond to sex or any other phenotypic character that we
recorded. There are several potential explanations for this consistent structure, of which we
will explore the evidence for three: (a) spatially segregated spawning sites and philopatry in
these species, where adults move throughout the lake but return to the same spawning
grounds to breed; (b) temporally segregated spawning groups within these species; or (c)
historical allopatric groups that have become sympatric yet partially retained their genetic
distinctiveness.

Genetically distinct groups that occur in sympatry must have at least partial reproductive
isolation, and in fish without clear sexual selection, this often occurs intraspecifically as a
segregation of spawning either in space or time. If there is site fidelity to regionally
differentiated spawning grounds but between-region movement outside of spawning, then
this could create patterns of differentiation such as those that we observe. If spatially
segregated spawning sites exist, then the adults that are sampled at a single site may have
migrated from a more distant site (unless they are collected while spawning, in which case
we would expect the genetics to match with their geographic location), but we would expect
littoral juveniles to exhibit a stronger pattern of geographic-based differentiation or isolation
by distance. In addition, we would expect juveniles collected at the same site to be more
closely related than adults collected at the same site, or juveniles and adults collected at the
same site. We did not find this to be the case in any of the four species (Fig. S14), with the
caveat that our sample sizes for these comparisons is low and our data may exhibit batch
effects due to our sampling strategies. We did find stronger differentiation in *L. angustifrons*
and *L. mariae* between juveniles and adults collected at the same site than between
juveniles and juveniles collected at the same site; however, the mean F_{ST} for adult-adult
comparisons was not larger than that of juvenile-juvenile comparisons, so the evidence
supporting this hypothesis is mixed. However, again, our sample size in each of these
categories for each sampling site is small, and therefore future studies would be helpful for
quantifying support for this hypothesis.

Genetic differentiation related to temporal stratification in spawning behavior—such as
having a population of "dry season" spawners and a population of "rainy season" spawners—
could also explain the existence of sympatric groups. Coulter (1976) suggested that the four
species likely spawn continuously, but with peaks in number of ripe females from August to
November/December, and potentially other smaller peaks in March/April. In addition,
observations of well-differentiated size distributions of young *Lates* found in littoral weed

beds suggests periodicity in spawning (Coulter 1976). Unfortunately, we do not have data on
the spawning status of the fish collected in this study, which would facilitate investigating this
hypothesis. However, we were able to test whether juveniles collected at a similar time of
year were more closely related than those collected at a different time of year, regardless of
sampling site. We do not find any evidence for juveniles collected at the same time of year to
be more closely related than juveniles collected during different parts of the year in *L.*
microlepis (Fig. S15); however, our sampling is not conducive to being able to rigorously
investigate this relationship, as some sites were only sampled in one year or at one time of
year. Additionally, only a few juveniles were collected opportunistically from each species, as
they are too small to be targeted by fishermen. Investigating the possibility of discrete
spawning groups will be an important next step in understanding the weak genetic structure
present in these species, and will require more targeted sampling.

Another third potential explanation for this consistent structure could lie in the geologic
history of Lake Tanganyika. With an estimated age of 9 -12 million years, Lake Tanganyika is
one of the oldest lakes in the world and has been influenced by extensive lake level
fluctuations (Cohen *et al.* 1993). Based on the bathymetry of the lake, it is thought that Lake
Tanganyika was divided into three paleo-lakes during periods of extremely low water levels
(Cohen *et al.* 1997). Environmental fluctuations such as these may have led to oscillations
between sympatry or parapatry and allopatry for fish populations within the lake, which have
been invoked as a "species pump" (Greenwood 1981; Sedano *et al.* 2010) contributing to
diversity in some clades in the littoral (Janzen & Etienne 2017; Sturmbauer *et al.* 2016) and
benthic or pelagic (Duftner *et al.* 2005; Koblmüller *et al.* 2005) zones of the lake. While it has
recently been suggested that the origin of the Lake Tanganyika *Lates* radiation is much
younger than other endemic fish radiations in the lake (Koblmüller *et al.* 2021), it remains
unclear whether speciation in this clade was similarly triggered by these environmental
fluctuations—as well as whether more recent fluctuations could explain our contemporary
observations of weak genetic structure. Such fluctuations could also lead to persistent
genetic structure within species, for example, if spawning site fidelity within separate basins
is maintained after the basins reconnect. However, we would expect strong philopatry to
produce much stronger genetic differentiation than observed here.

Interestingly, *L. stappersii* in this study which were caught during the dry season in the south
of the lake show distinct carbon isotope signals from samples caught in the north (Ehrenfels
et al, *in prep*). These differences in carbon isotope ratios correlate with different rates of
primary productivity in the northern and the southern basin during the dry season (Loiselle *et*
al. 2014), and with differences in prey composition in different areas of the lake (Plisner *et al.*
2009; Kurki *et al.* 1999). This is evidence for at least partly resident populations of *L.*
stappersii, which stay in the northern or southern basin during the dry season for long

enough to incorporate the different isotopic signals of these basins. Such movement patterns
may reduce gene flow between *L. stappersii* populations from the basins; however, these
isotopic groups do not seem to correspond to the genetic groups found at K=2 here
(Ehrenfels *et al, in prep).*

With these potential explanations in mind, it is important to keep the limitations of our data in
mind. Admixture analyses using programs such as STRUCTURE (Pritchard *et al.* 2000) have
been shown to be biased with unbalanced sampling among sites, such that differentiation is
often overestimated at sites with the most individuals and underestimated at sites with few
individuals (e.g., Puechmaille 2016; Wang 2017). While it is unclear whether this is true for
entropy analyses as well, it is important to interpret our analyses with this in mind. For
example, we find multiple groups at the Kigoma site in *L. microlepis* at K=2, which is also the
site with the most samples present in the data set (Fig. 5B). Furthermore, our assignment of
individual locations to fisherman landing sites or markets is less precise than if we had exact
locations where the fish were caught. While we only have general knowledge of how far
fishermen generally travel from the landing site to fish during the day, it is possible that fish
collected at the same sampling site were collected up to ~40 km away from one another. In
addition, our samples collected in the Kigoma fish markets may have come from farther away
from those collected at the landing sites (Petit & Shipton 2012). These limitations influence
our ability to detect geographic trends in our data; however, we believe that these limitations
would only have minor effects on the inferences discussed here.

Recommendations for conservation and sustainable management of Lake Tanganika's Lates

We find evidence for unexpected weak yet genetically distinct and sympatric groups within all
four species. We furthermore find evidence for genetically distinct group within *L. mariae* at
our most southern sampling site, in agreement with previous studies suggesting that this
species is the most resident of all four *Lates* species (Coulter 1976, 1991). Intriguingly, this
structure is not linked to isolation-by-distance, but rather involves differentiation between
some individuals at multiple sites, suggesting the existence of a mechanism for maintaining
differentiation while these distinct genetic groups are in sympatry. In contrast, the low F_{ST}
estimates between samples from distant sites and lack of isolation-by-distance suggest that
these species do disperse over large distances, suggesting a need for management of these
species on at a lake-wide scale. Furthermore, it is likely that more extensive sampling of
these little-studied species may uncover important additional diversity.

To maintain intraspecific genetic variation, viable population sizes, and possible adaptive
differences between genetic groups within the *Lates* species, more research is needed on

the location and timing of spawning for the four species and whether restricting catches in
certain areas and periods of the year could help the larger-bodied species to recover from
low population sizes. Harvest restrictions have been implemented in regions around Lake
Tanganyika where community conservation areas are active (Kimirei & Sweke 2018) but are
not in place throughout the lake. Such management practices are performed successfully in
the cod fishery of Denmark (albeit after collapse of this fishery and extinction of many of the
original stocks; Dahle *et al.* 2018), and in the salmon fishery of Bristol Bay (Schindler *et al.*
2015; Schindler *et al.* 2010). However, implementation of such a strategy requires detailed
evidence of spawning ground locations and timing of spawning, as well as for fishermen to
be able to distinguish between the genetically distinct stocks. Therefore, studies are needed
to shed more light on the phenotypic, ecological, and phenological characteristics of the
genetic groups within each species. In addition, studies are needed to examine whether fish
from different spawning grounds form the distinct genetic clusters that we identified in this
study. Genomic analyses, although helpful in generating hypotheses about the natural
history of these species, will need to be paired with an in-depth understanding of natural
history, including integration of local knowledge, for managing these fish stocks. A
sustainable management approach is crucial to protect the inshore nursery habitats and
ensure recruitment of the species in Lake Tanganyika.

Funding

This work was supported by the Swiss National Science Foundation (grant CR23I2-166589
to OS), the University of Wyoming (start-up funding to CEW), the National Science
Foundation (grant DEB-1556963 to CEW), the National Institute of General Medical Sciences
of the National Institutes of Health (Institutional Development Award grant P20GM103432),
and the American Genetics Association (Ecological, Evolutionary, and Conservation
Genomics Research Award to JAR).

Acknowledgements

Computing was accomplished with an allocation from the University of Wyoming's Advanced
Research Computing Center, on its Teton Intel x86_64 cluster
(https://doi.org/10.15786/M2FY47) and the Genetic Diversity Center (GDC) of ETH Zürich.
Special thanks go to Benedikt Ehrenfels for coordinating field campaigns, Mupape Mukuli for
facilitating logistics during fieldwork, and to the crew of the MV Maman Benita for fieldwork
assistance. We thank the whole team at the Tanzanian Fisheries Research Institute for their
support. A special thanks goes to Mary Kishe for her support during fieldwork permission

processes and to the Tanzanian Commission for Science and Technology (COSTECH) for
the permits to do this research. Live fish samples were collected under the approved
University of Wyoming IACUC protocol #20160602CW00241-01. We appreciate The Nature
Conservancy's Tuungane Project for providing funding and logistical support, with special
thanks going to Colin Apse and Peter Limbu. Thanks to members of the Wagner lab at the
University of Wyoming and the FishEc group at EAWAG for discussions about the analyses
and manuscript. We also thank the Associate Editor and four anonymous reviewers for the
valuable comments that greatly improved the quality of this manuscript.

Data Availability

We have deposited the primary data underlying these analyses as follows:

- Sampling locations, morphological data, filtered VCF files have been archived on
Zenodo (http://doi.org/10.5281/zenodo.5216259)
- Fastq files for each individual have been archived in NCBI SRA XXXXX
- Scripts for data analysis are available on Github (https://github.com/jessicarick/lates-
popgen), and the version at publication will be archived on Zenodo

References

Anderson, EC (2018) whoa: Where's my Heterozygotes? Observations on genotyping Accuracy. Version 0.0.2.999. *Zenodo*, doi: 10.5281/zenodo.2401764

Baird NA, Etter PD, Atwood TS, *et al.* (2008) Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLoS One* **3**.

Bay LK, Crozier RH, Caley MJ (2006) The relationship between population genetic structure and pelagic larval duration in coral reef fishes on the Great Barrier Reef. *Marine Biology* **149**, 1247-1256.

Berg PR, Star B, Pampoulie C, *et al.* (2017) Trans-oceanic genomic divergence of Atlantic cod ecotypes is associated with large inversions. *Heredity* **119**, 418-428.

Bergamino N, Horion S, Stenuite S, *et al.* (2010) Spatio-temporal dynamics of phytoplankton and primary production in Lake Tanganyika using a MODIS based bio-optical time series. *Remote Sensing of Environment* **114**, 772-780.

Bernatchez L (2016) On the maintenance of genetic variation and adaptation to environmental change: considerations from population genomics in fishes. *Journal of Fish Biology* **89**, 2519-2556.

Brannon EL, Powell MS, Quinn TP, Talbot A (2004) Population Structure of Columbia River Basin Chinook Salmon and Steelhead Trout. *Reviews in Fis heries Science* **12**, 99-232.

Bulengela G, Onyango P, Brehm J, *et al.* (2020) "Bring fishermen at the center": the value of local knowledge for understanding fisheries resources and climate-related changes in Lake Tanganyika. *Environment, Development and Sustainability* **22**, 5621-5649.

Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis tool set for population genomics. *Mol Ecol* **22**, 3124-3140.

Chang CC, Chow CC, Tellier LCAM, *et al.* (2015) Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, s13742-015-0047-8.

Chapman DW (1976) Acoustic Estimates of Pelagic Ichthyomass in Lake Tanganyika with an Inexpensive Echo Sounder. *Transactions of the American Fisheries Society* **105**, 581-587.

Cirhuza DM, Plisnier PD (2016) Composition and seasonal variations in abundance of Copepod (Crustacea) populations from the northern part of Lake Tanganyika. *Aquatic Ecosystem Health & Management* **19**, 401-410.

Cohen AS, Gergurich EL, Kraemer BM, *et al.* (2016) Climate warming reduces fish production and benthic habitat in Lake Tanganyika, one of the most biodiverse freshwater ecosystems. *Proc Natl Acad Sci U S A* **113**, 9563-9568.

Cohen AS, Lezzar K-E, Tiercelin J-J, Soreghan M (1997) New palaeogeographic and lake-level reconstructions of Lake Tanganyika: implications for tectonic, climatic and biological evolution in a rift lake. *Basin Research* **9**, 107-132.

Cohen AS, Soreghan MJ, Scholz CA (1993) Estimating the age of formation of lakes: an example from Lake Tanganyika, East African Rift system. *Geology* **21**, 511-514.

Coulter GW (1970) Population changes within a group of fish species in Lake Tanganyika following their exploitation. *Journal of Fish Biology* **2**, 329-353.

Coulter GW (1976) The biology of Lates species (Nile perch) in Lake Tanganyika, and the status of the pelagic fishery for Lates species and Luciolates stappersii (Blgr.). *Journal of Fish Biology* **9**, 235-259.

Coulter GW (1991) *Lake Tanganyika and its Life* British Museum (Natural History) Cromwell Road, London SW7 5BD & Oxford University Press, Walton Street, Oxford OX2 6DP.

Dahle G, Johansen T, Westgaard J-I, Aglen A, Glover KA (2018) Genetic management of mixed-stock fisheries "real-time": The case of the largest remaining cod fishery operating in the Atlantic in 2007–2017. *Fisheries Research* **205**, 77-85.

Danecek P, Auton A, Abecasis G, *et al.* (2011) The variant call format and VCFtools. *Bioinformatics* **27**, 2156-2158.

De Keyzer ELR, De Corte Z, Van Steenberge M, *et al.* (2019) First genomic study on Lake Tanganyika sprat *Stolothrissa tanganicae*: a lack of population structure calls for

integrated management of this important fisheries target species. *BMC Ecology and*
Evoluiton **19**, 6.
De Keyzer ELR, Mulungula PM, Lufungula GA, *et al.* (2020) Local perceptions on the state of
the pelagic fisheries and fisheries management in Uvira, Lake Tanganyika, DR
Congo. *Journal of Great Lakes Research* **46**, 1740-1753.
Dray S, Dufour A-B (2007) The ade4 Package: Implementing the Duality Diagram for
Ecologists. *2007* **22**, 20.
Duftner N, Koblmüller S, Sturmbauer C (2005) Evolutionary relationships of the
Limnochromini, a tribe of benthic deepwater cichlid fish endemic to Lake Tanganyika,
East Africa. *Journal of Molecular Evolution* **60**, 277-289.
Eccles DH (1992) Field guide to the freshwater fishes of Tanzania. In: *species identification*
sheets for fishery purposes (ed. FAO), p. 145. FAO, Rome.
Ellegren H, Galtier N (2016) Determinants of genetic diversity. *Nature Reviews Genetics* **17**,
422-433.
Ellis CMA (1978) Biology of Luciolates-Stappersi in Lake Tanganyika (Burundi). *Transactions*
of the American Fisheries Society **107**, 557-566.
Genner MJ, Nichols P, Shaw PW*, et al.* (2010) Population structure on breeding grounds of
Lake Malawi's 'twilight zone' cichlid fishes. *Journal of Biogeography* **37**, 258-269.
Gompert Z, Lucas LK, Buerkle CA*, et al.* (2014) Admixture and the organization of genetic
diversity in a butterfly species complex revealed through common and rare genetic
variants. *Mol Ecol* **23**, 4555-4573.
Greenwood, PH (1981) Species-flocks and explosive evolution. *Chance, change and*
challenge- the evolving biosphere, 61-74.
Gruber B, Unmack PJ, Berry OF, Georges A (2018) dartr: An r package to facilitate analysis
of SNP data generated from reduced representation genome sequencing. *Molecular*
Ecology Resources **18**, 691-699.
Hutchinson WF (2008) The dangers of ignoring stock complexity in fishery management: the
case of the North Sea cod. *Biol Lett* **4**, 693-695.
Ivory SJ, McGlue MM, Peterman C, *et al* (2021) Climate, vegetation, and weathering across
space and time in Lake Tanganyika (tropical eastern Africa). *Quaterny Science*
Advances **3**, 10023.
Janzen T, Etienne RS (2017) Inferring the role of habitat dynamics in driving diversification:
evidence for a species pump in Lake Tanganyika cichlids.
Johnson MS, Black R (1984) Pattern beneath the chaos: The effect of recruitment on genetic
patchiness in an intertidal limpet. *Evolution* **38**, 1371-1383.
Junker J, Rick JA, McIntyre PB*, et al.* (2020) Structural genomic variation leads to genetic
differentiation in Lake Tanganyika's sardines. *Mol Ecol* **29**, 3277-3298.
Kimirei IA, Mgaya YD, Chande AI (2008) Changes in species composition and abundance of
commercially important pelagic fish species in Kigoma area, Lake Tanganyika,
Tanzania. *Aquatic Ecosystem Health and Management* **11**, 29-35.
Kimirei IA, Sweke EA (2018) Community-based conservation areas as a fisheries
management tool in the East African Great Lakes: an example from Tanzania. In: 34[th]
SIL Congress, 19-24 August 2018. Nanjing, China. pp. 230 Book of abstracts.
Kirubakaran TG, Grove H, Kent MP*, et al.* (2016) Two adjacent inversions maintain genomic
differentiation between migratory and stationary ecotypes of Atlantic cod. *Mol Ecol*
25, 2130-2143.
Kmentová N, Koblmüller S, Van Steenberge M, *et al.* (2020) Weak population structure and
recent demographic expansion of the monogenean parasite *Kapentagyrus* spp.
infecting clupeid fishes of Lake Tanganyika, East Africa. *International Journal for*
Parasitology **50**, 471-486.
Knaus BJ, Grünwald NJ (2017) VCFR: a package to manipulate and visualize variant call
format data in R. *Mol Ecol Resources* **17**, 44-53.
Koblmüller S, Duftner N, Katongo C, *et al.* (2005) Ancient divergence in bathypelagic Lake
Tanganyika deepwater cichlids: Mitochondrial phylogeny of the tribe Bathybatini.
Journal of Molecular Evolution **60**, 297-314.

Koblmüller S, Odhiambo EA, Sinyinza D, *et al.* (2015) Big fish, little divergence: phylogeography of Lake Tanganyika's giant cichlid, *Boulengerochromis microlepis*. *Hydrobiologia* **748**, 29-38.

Koblmüller S, Zangl L, Börger C*, et al.* (2019) Only true pelagics mix: comparative phylogeography of deepwater bathybatine cichlids from Lake Tanganyika. *Hydrobiologia* **832**, 93-103.

Koblmüller S, Schöggl CA, Lorber CJ, *et al.* (2021) African lates perches (Teleostei, Latidae, *Lates*): Paraphyly of Nile perch and recent colonization of Lake Tanganyika. *Molecular Phylogenetics and Evolution* **160**, 107141.

Korneliussen TS, Albrechtsen A, Nielsen R (2014) ANGSD: Analysis of Next Generation Sequencing Data. *BMC Bioinformatics* **15**, 356.

Korneliussen TS, Moltke I, Albrechtsen A, Nielsen R (2013) Calculation of Tajima's D and other neutrality test statistics from low depth next-generation sequencing data. *BMC Bioinformatics* **14**, 289.

Kurki H, Mannini P, Vuorinen I, *et al.* (1999) Macrozooplankton communities in Lake Tanganyika indicate food chain differences between the northern part and the main basins. *Hydrobiologia* **407**, 123-129.

Kuusipalo L (1999) Genetic differentiation of endemic nile perch *Lates stappersi* (*Centropomidae*, *Pisces*) populations in Lake Tanganyika suggested by RAPD markers. In: Lindqvist O.V., Mölsä H., Salonen K., Sarvala J. (eds) *From Limnology to Fisheries: Lake Tanganyika and Other Large Lakes*. Developments in Hydrobiology, vol 141. Springer, Dordrecht.

Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**, 357-359.

Lewis PO (2001) A Likelihood Approach to Estimating Phylogeny from Discrete Morphological Character Data. *Systematic Biology* **50**, 913-925.

Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics (Oxford, England)* **25**, 1754-1760.

Li H, Handsaker B, Wysoker A*, et al.* (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079.

Loiselle S, Cózar A, Adgo E, *et al.* (2014) Decadal Trends and Common Dynamics of the Bio-Optical and Thermal Characteristics of the African Great Lakes. *PLoS One* **9**, e93656.

Manier MK, Arnold SJ (2006) Ecological correlates of population genetic structure: a comparative approach using a vertebrate metacommunity. *Proceedings of the Royal Society B: Biological Sciences* **273**, 3001-3009.

Mannini P (1998a) *Ecology of the Pelagic Fish Resources of Lake Tanganyika*, University of Hull.

Mannini P, Aro E, Katonda I*, et al.* (1996) Pelagic fish stocks of Lake Tanganyika: biology and exploitation. FAO/FINNIDA Research for the Management of the Fisheries of Lake Tanganyika. In: *GCP/RAF/271/FIN—TD/53 (En)*, p. 60p.

Mannini P, Katonda I, Kissaka B, Verburg P (1999) Feeding ecology of Lates stappersii in Lake Tanganyika. *Hydrobiologia* **407**, 131-139.

Martinez Barrio A, Lamichhaney S, Fan G*, et al.* (2016) The genetic basis for ecological adaptation of the Atlantic herring revealed by genome sequencing. *Elife* **5**.

McGlue MM, Ivory SJ, Stone JR, *et al.* (2020) Solar irradiance and ENSO affect food security in Lake Tanganyika, a major African inland fishery. *Science Advances* **6**, eabb2191.

Meisner J, Liu S, Huang M, Albrechtsen A (2021) Large-scale inference of population structure in presence of missingness using PCA. *Bioinformatics*, btab027.

Mölsä H, Reynolds JE, Coenen EJ, Lindqvist OV (1999) Fisheries research towards resource management on Lake Tanganyika. *Hydrobiologia* **407**, 1-24.

Mölsä H, Sarvala J, Badende S*, et al.* (2002) Ecosystem monitoring in the development of sustainable fisheries in Lake Tanganyika. *Aquatic Ecosystem Health and Management* **5**, 267-281.

Mulimbwa N, Raeymaekers JAM, Sarvala J (2014) Seasonal changes in the pelagic catch of two clupeid zooplanktivores in relation to the abundance of copepod zooplankton in

the northern end of Lake Tanganyika. *Aquatic Ecosystem Health & Management* **17**, 25-33.

Muller P, Li XP, Niyogi KK (2001) Non-photochemical quenching. A response to excess light energy. *Plant Physiol* **125**, 1558-1566.

Munyandero J (2002) The lake Tanganyika clupeid and latid fishery system: Indicators and problems inherent in assessments and management. *African Study Monographs* **23**, 117-145.

Mziray P, Kimirei IA, Staehr PA, *et al.* (2018) Seasonal patterns of thermal stratification and primary production in the northern parts of Lake Tanganyika. *Journal of Great Lakes Research* **44**, 1209–1220.

O'Reilly CM, Alin SR, Piisnier PD, Cohen AS, McKee BA (2003) Climate change decreases aquatic ecosystem productivity of Lake Tanganyika, Africa. *Nature* **424**, 766-768.

Ogutu-Ohwayo R, Natugonza V, Musinguzi L, *et al.* (2016) Implications of climate variability and change for African lake ecosystems, fisheries productivity, and livelihoods. *Journal of Great Lakes Research* **42**, 498-510.

Olsen JB, Crane PA, Flannery BG*, et al.* (2011) Comparative landscape genetic analysis of three Pacific salmon species from subarctic North America. *Conservation Genetics* **12**, 223-241.

Parchman TL, Gompert Z, Mudge J*, et al.* (2012) Genome-wide association genetics of an adaptive trait in lodgepole pine. *Mol Ecol* **21**, 2991-3005.

Peterman WE, Anderson TL, Ousterhout BH*, et al.* (2015) Differential dispersal shapes population structure and patterns of genetic differentiation in two sympatric pond breeding salamanders. *Conservation Genetics* **16**, 59-69.

Petit P, Shipton T (2012) IUU Fishing on Lake Tanganyika. *Programme for the Implementation of a Regional Fisheries Strategy for the Eastern and Southern Africa and Indian Ocean Region.*

Pettersson ME, Rochus CM, Han F*, et al.* (2019) A chromosome-level assembly of the Atlantic herring genome-detection of a supergene and other signals of selection. *Genome Research* **29**, 1919-1928.

Plisnier PD, Chitamwebwa D, Mwape L, *et al.* (1999) Limnological annual cycle inferred from physical-chemical fluctuations at three stations of Lake Tanganyika. *Hydrobiologia* **407**, 45-58.

Plisnier PD, Mgana H, Kimirei I, *et al.* (2009) Limnological variability and pelagic fish abundance (Stolothrissa tanganicae and Lates stappersii) in Lake Tanganyika. *Hydrobiologia* **625**,117-134.

Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* **155**: 945-959.

Puechmaille SJ (2016) The program STRUCTURE does not reliably recover the correct population structure when sampling is uneven: subsampling and new estimators alleviate the problem. *Molecular Ecology Resources* **16**, 608-627.

Purcell S, Neale B, Todd-Brown K, *et al.* (2007) PLINK: a toolset for whole-genome association and population-based linkage analysis. *American Journal of Human Genetics*, 81: 559-575.

R Core Team (2021) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Reich D, Thangaraj K, Patterson N, Price AL, Singh L (2009) Reconstructing Indian population history. *Nature* **461**, 489-494.

Reiss H, Hoarau G, Dickey-Collas M, Wolff WJ (2009) Genetic population structure of marine fish: mismatch between biological and fisheries management units. *Fish and Fisheries* **10**, 361-395.

Riginos C, Buckley YM, Blomberg SP*, et al.* (2014) Dispersal Capacity Predicts Both Population Genetic Structure and Species Richness in Reef Fishes. *The American Naturalist* **184**, 52-64.

Riginos C, Doublas KE, Jin Y, *et al.* (2011) Effects of geography and life history traits on genetic differentiation in benthic marine fishes. *Ecography* **34**, 566-575.

Sarvala J, Tarvainen M, Salonen K, Mölsä H (2002) Pelagic food web as the basis of
fisheries in Lake Tanganyika: A bioenergetic modeling analysis. *Aquatic Ecosystem*
Health and Management **5**, 283-292.
Sayers M, Bosse K, Fahnenstiel G, Shuchman R (2020) Carbon fixation trends in eleven of
the world's largest lakes: 2003-2018. *Water* **12**, 3500.
Schindler DE, Armstrong JB, Reed TE (2015) The portfolio concept in ecology and evolution.
Frontiers in Ecology and the Environment **13**, 257-263.
Schindler DE, Hilborn R, Chasco B*, et al.* (2010) Population diversity and the portfolio effect
in an exploited species. *Nature* **465**, 609-612.
Sedano RE, Burns KJ, Riddle B (2010) Are the Northern Andes a species pump for
Neotropical birds? Phylogenetics and biogeography of a clade of Neotropical
tanagers (Aves: Thraupini). *Journal of Biogeography* **37**, 325-343.
Selkoe KA, Toonen RJ (2011) Marine connectivity: a new look at pelagic larval duration and
genetic metrics of dispersal. *Marine Ecology Progress Series* **436**, 291-305.
Shanks (2009) Pelagic larval duration and dispersal distance revisited. *The Biological*
Bulletin **216**, 373-385.
Shastry, V, Adams PE, Lindtke D, *et al.* (2021) Model-based genotype ancestry estimation
for potential hybrids with mixed-ploidy. *Mol Ecol Resources* **21**: 1434-1451.
Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**, 1312-1313.
Stearns SC (1992) *The evolution of life histories* Oxford Univ. Press, New York.
Stenuite S, Pirlot S, Hardy MA*, et al.* (2007) Phytoplankton production and growth rate in
Lake Tanganyika: evidence of a decline in primary productivity in recent decades.
Freshwater Biology **52**, 2226-2239.
Sterner T (2007) Unobserved diversity, depletion and irreversibility The importance of
subpopulations for management of cod stocks. *Ecological Economics* **61**, 566-574.
Sturmbauer C, Börger C, Van Steenberge M, Koblmüller S (2016) A separate lowstand lake
at the northern edge of Lake Tanganyika? Evidence from phylogeographic patterns in
the cichlid genus Tropheus. *Hydrobiologia* **791**, 51-68.
Sunday JM, Popovic I, Palen WJ, Foreman MG, Hart MW (2014) Ocean circulation model
predicts high genetic structure observed in a long-lived pelagic developer. *Mol Ecol*
23, 5036-5047.
Supple MA, Shapiro B (2018) Conservation of biodiversity in the genomics era. *Genome*
Biology **19**, 131.
Tange O (2011) GNU Parallel - The Command-Line Power Tool. ;login: The USENIX
Magazine, **36**:42-47.
[dataset] Temasek Life Sciences Laboratory (2016). *Lates calcarifer* isolate:ASB-BC8
Genome sequencing and assembly. National Center for Biotechnology Information
Sequence Read Archive, Assembly GCA_001640805.1.
Therkildsen NO, Hemmer-Hansen J, Als TD*, et al.* (2013) Microevolution in time and space:
SNP analysis of historical DNA reveals dynamic signatures of selection in Atlantic
cod. *Mol Ecol* **22**, 2424-2440.
Van der Knaap M (2013) Comparative analysis of fisheries restoration and public
participation in Lake Victoria and Lake Tanganyika. *Aquatic Ecosystem Health and*
Management **16**, 279-287.
Van der Knaap M, Katonda KI, De Graaf GJ (2014) Lake Tanganyika fisheries frame survey
analysis: Assessment of the options for management of the fisheries of Lake
Tanganyika. *Aquatic Ecosystem Health and Management* **17**, 4-13.
van Zwieten PAM, Roest FC, Machiels MAM, Van Densen WLT (2002) Effects of inter-
annual variability, seasonality and persistence on the perception of long-term trends
in catch rates of the industrial pelagic purse-seine fishery of northern Lake
Tanganyika (Burundi). *Fisheries Research* **54**, 329-348.
van Zwieten PAM, Kolding J, Plank MJ*, et al.* (2016) The Nile perch invasion in Lake Victoria:
cause or consequence of the haplochromine decline? *Canadian Journal of Fisheries*
and Aquatic Sciences **73**, 622-643.

Verburg P, Antenucci JP, Hecky RE (2011) Differential cooling drives large-scale convective
circulation in Lake Tanganyika. *Limnology and Oceanography* **56**, 910-926.
Verburg P, Hecky RE (2009) The physics of the warming of Lake Tanganyika by climate
change. *Limnology and Oceanography* **54**, 2418-2430.
Verburg P, Hecky RE, Kling H (2003) Ecological consequences of a century of warming in
Lake Tanganyika. *Science* **301**, 505-507.
Vij S, Kuhl H, Kuznetsova IS*, et al.* (2016) Chromosomal-Level Assembly of the Asian
Seabass Genome Using Long Sequence Reads and Multi-layered Scaffolding. *PLOS*
Genetics **12**, e1005954-e1005954.
Wang, J (2017) The computer program STRUCTURE for assigning individuals to
populations: easy to use but easier to misuse. *Molecular Ecology Resources* **17**, 981-
990.
Wenburg JK, Bentzen P, Foote CJ (1998) Microsatellite analysis of genetic population
structure in an endangered salmonid: the coastal cutthroat trout (Oncorhynchus clarki
clarki). *Molecular Ecology* **7**, 733-749.
White JW, Botsford LW, Hastings A, Largier JL (2010) Population persistence in marine
reserve networks: incorporating spatial heterogeneities in larval dispersal. *Marine*
Ecology Progress Series **398**, 49-67.
Young EF, Belchier M, Hauser L*, et al.* (2015) Oceanography and life history predict
contrasting genetic population structure in two Antarctic fish species. *Evolutionary*
Applications **8**, 486-509.
1153