A divergent protein kinase A in the human pathogen *Leishmania* is associated with

cell cortex microtubules and controls cell shape

Fischer Weinberger, R.[1], Bachmaier, S.[2], Ober, V.[2] Githure, G.B.[2], Dandugudumula, R.[1],

Phan, I.Q.[3,4], Almoznino, M.[1], Polatoglou, E.[2], Tsigankov, P.[1], Nitzan Koren, R[1].,

Myler, P.J.[3,4,5], Boshart, M.[2] and Zilberstein, D.[1*]

[1]Faculty of Biology, Technion - Israel Institute of Technology, Haifa 32000, Haifa, Israel

[2]Faculty of Biology, Genetics, Ludwig-Maximilian University Munich, Großhaderner
Straße 2-4, 82152 Martinsried, Germany

[3]Seattle Structural Genomics Center for Infectious Disease, Seattle, WA, USA

[4]Center for Global Infectious Disease Research, Seattle Children's Research Institute,
307 Westlake Ave N, Seattle, WA 98109-5219, USA.

[5]Department of Pediatrics, Department of Biomedical Informatics & Medical Education,
and Department of Global Health, University of Washington, Seattle, WA 98195, USA

*Corresponding author email: danz@technion.ac.il

Summary

Parasitic protozoa of the genus *Leishmania* cycle between the phagolysosome of mammalian macrophages, where they reside as rounded intracellular amastigotes, and the midgut of female sand flies, which they colonize as elongated extracellular promastigotes. Previous studies indicated that protein kinase A (PKA) plays an important role in the initial steps of promastigote development into amastigotes. Here, we describe a novel regulatory subunit of PKA (which we have named PKAR3) that is unique to *Leishmania* and most (but not all) other Kinetoplastea. PKAR3 is localized to subpellicular microtubules (SPMT) in the cell cortex, where it recruits a specific catalytic subunit (PKAC3). Promastigotes of *PKAR3* or *PKAC3* null mutants lose their elongated shape and are round but remain flagellated. Truncation of an N-terminal formin homology-like domain of PKAR3 results in its detachment from the SPMT, also leading to rounded promastigotes. Thus, the tethering of PKAC3 kinase activity via PKAR3 at the cell cortex is essential for maintenance of the elongated shape of promastigotes. This role of PKAR3 is reminiscent of PKARIβ and PKARIIβ binding to microtubules of mammalian neurons, which is essential for the elongation of dendrites and axons, respectively. Interestingly, PKAR3 does not bind cAMP but nucleoside analogs with a very high affinity similar to the PKAR1 isoform of *Trypanosoma*. We propose that these early diverged protists have re-purposed PKA for a novel signaling pathway that spatiotemporally controls microtubule remodeling and cell shape *via* PKA activity.

Keywords: differentiation, protein kinase A, signaling, microtubule cortex, morphogenesis, *Leishmania*, parasitism

Introduction

Leishmania donovani is a (trypanosomatid) protozoan parasite that causes kala azar, a fatal form of visceral leishmaniasis in humans (1). These organisms cycle between the midgut of female sand flies, where they reside as flagellated extracellular promastigotes, and the phagolysosomes of mammalian macrophages, where they live as aflagellate amastigotes (2). Infective metacyclic promastigotes are introduced into the host during the sand fly blood meal; after which they are phagocytosed by resident macrophages near the bite site (3,4). Once inside the host phagolysosome, promastigotes encounter two physical cues that distinguish this environment from that of the vector gut: acidic pH (~5.5) and elevated temperature (~37°C). Promastigotes process these cues into a signal that initiates development into amastigotes (5-7).

Among the earliest events during L. donovani promastigote-to-amastigote differentiation are changes in the phosphorylation profile of many proteins (8). Significantly, while protein kinase A (PKA)-specific phosphorylation is common in promastigotes, most of these phosphoproteins are dephosphorylated within a few minutes after initiation of differentiation into amastigotes (7). This suggests that regulation of PKA activity is important for regulation of Leishmania development. PKA-mediated signaling pathways are ubiquitous in eukaryotic cells, being implicated in growth control, development, and metabolism. The canonical PKA regulatory system in vertebrates involves assembly of catalytic (PKAC) and regulatory (PKAR) subunits into an inactive heterotetrameric complex. Dimerization is mediated by a dimerization and docking (D/D) domain in the N-terminal portion of PKAR (9). When cAMP binds the cyclic nucleotide-binding domains (cNBDs) near the C-terminal of each PKAR, the resultant conformational change triggers dissociation of the PKAC/R complex and release of catalytically active PKAC subunits. Compartmentation of PKA activity to different cellular microdomains is

achieved (at least in higher eukaryotes) through interaction of the D/D domain of PKAR with a

diverse family of A-kinase anchoring proteins (AKAPs) that vary according to biological context

(10). However, PKARs from many lower eukaryotes (including trypanosomatids) appear to lack

the D/D domain (11,12) and they lack obvious AKAPs (13).

Most eukaryote genomes (with the notable exception of plants and algae) encode one or

more PKAC subunits (13). Trypanosomatids (including *L. donovani*) are no exception, having

two subunits (PKAC1 and C2) that are almost identical and a third (C3) that is more divergent

(14-16). Phosphoproteomic analyses of *L. donovani* revealed that all three PKAC subunits

contain at least two phosphorylation sites (17). One, within the kinase active site loop, matches

the canonical PKA phosphorylation motif (RXXpS/pT) and is phosphorylated only in

promastigotes. The second is located within a canonical ERK1/ERK2 substrate motif (pSP) close

to the C-terminus and while PKAC1/C2 is phosphorylated at this site in promastigotes, PKAC3

is phosphorylated only after exposure to the differentiation signal (8). In other eukaryotes,

phosphorylation of both a threonine (T_{197}) in the activation loop and a serine (S_{338}) near the C-

terminus is necessary for full activation of PKAC, suggesting that PKAC1/C2 are active in

L. donovani promastigotes, while PKAC3 is transiently activated only upon receiving the

differentiation signal.

Most eukaryotes also possess one or more PKAR subunits, whose number and evolution

are more dynamic than for PKAC (13). All trypanosomatids have a regulatory subunit (PKAR1)

that has been shown to localize to the flagellum of *Trypanosoma brucei* (18)(19)(20) and

L. donovani, where it is expressed only in the promastigote stage (7,21). However, in contrast to

higher eukaryotes, trypanosomatid PKAR1 appears to be monomeric and unable to bind cAMP

(11,12).

Interestingly, *L. donovani* (along with other *Leishmania* species and *Trypanosoma cruzi*,

but not *T. brucei*) have a second PKAR-like subunit (14), containing at least 10 sites that are

phosphorylated or dephosphorylated during differentiation, implying that it may have a

regulatory role (8). Here, we show that this protein (which we have renamed PKAR3) forms a

holoenzyme complex with PKAC3 and anchors it to subpellicular microtubules at the cell cortex

via a formin FH2-like domain at the N-terminus of PKAR3. Promastigotes of *L. donovani* null

mutants lacking either PKAR3 or PKAC3 are predominantly rounded compared to wild type

(WT) cells, suggesting that tethering of PKA kinase activity to microtubules is necessary for

maintenance of elongated cell shape. Thus, this study ties together an evolutionary divergent

PKA localized to the microtubule cytoskeleton of *Leishmania* with developmental

morphogenesis.

Results

Leishmania PKA regulatory subunits are evolutionary divergent.

Previous analyses of the PKAR subunits from a wide range of organisms revealed that they have undergone dynamic and divergent evolution during eukaryote history (13). We used BLAST searches to identify PKAR orthologues in 15 different genera and two unclassified trypanosomatids from the Kinetoplastea (**Table S1**). *Leishmania* and most other trypanosomatids contain two paralogues (PKAR1 and PKAR3) that lack the canonical dimerization and docking (D/D) domain found in the N-terminal region of the PKARs of many (but not all) eukaryotes (**Fig. 1A**). Instead, PKAR1 has an RNI-like domain containing Leucine Rich Repeats (LRRs) that may mediate different protein-protein interactions but not homodimerization (11,2). PKAR3 has a long N-terminal region without obvious domain profile signatures. Construction of a phylogenetic tree (**Fig. S1**) revealed that PKAR1 and PKAR3 represent two separate branches distinct from that containing the canonical PKAR(s) found in most other eukaryotes.

Cytoskeletal localization of PKAR3 to cortical subpellicular microtubules.

As shown in **Fig. S2**, polyclonal antibodies raised in rabbits against recombinant PKAR3 reacted with a 72 kDa protein in wild-type (WT) cells. This protein was absent in null mutants (Δ*pkar3*), but reappeared after ectopic expression of full-length *PKAR3* in the null mutant (Δ*pkar3::PKAR3$_{FL}$*). Confocal microscopy using the antibody against PKAR3 revealed its localization to the cell cortex in *Leishmania* promastigotes (**Fig. 1B**). The cell cortex contains a rigid cytoskeleton made of stable subpellicular microtubules (SPMT) that are composed largely of β-tubulin (22)(23)(24). Previous studies have shown that non-ionic detergents can solubilize the trypanosomatid surface membrane, while leaving the SPMT intact (25-27). Trématent of

L. donovani promastigotes with the zwitterionic detergent n-dodecyl β-D-maltoside (DDM,

which is neutral at pH 7) resulted in enrichment of β-tubulin in the DDM-insoluble fraction (**Fig.**

1B and **1C**). In contrast, cytosolic HSP83 (28) is absent from the DDM-insoluble fraction. The

presence of PKAR3 in the DDM-insoluble pellet suggests that it is associated with the SPMT.

Binding of a PKA regulatory subunit (PKARIβ) to stable microtubules has been

documented in neuronal axons of mammalian cells (29), and is mediated by interaction of the

dimerization and docking (D/D) domain of PKARIβ with a microtubule associated protein 2

(MAP2) (30). The *Leishmania* genome does not contain an obvious orthologue of MAP2 and

PKAR3 lacks the canonical D/D domain, suggesting its localization to the SPTM is mediated by

a different mechanism. The N-terminal 50 amino acids of PKAR3 is relatively well-conserved

and analysis using HHpred (31,32) revealed structural similarity to three Protein Data Bank

(PDB) entries containing a formin homology (FH2) domain (**Fig. S3**). Formins are a diverse

family of proteins that participate in polymerization of actin and assembly of microtubule

cytoskeletons in many eukaryotes (33), leading us to speculate that this domain might be

responsible for binding of PKAR3 to the subpellicular microtubules. To test this hypothesis, we

generated null mutants of *L. donovani* expressing an ectopic version of *PKAR3* lacking the N-

terminal 90 amino acids (*Δpkar3::PKAR3*$_{ΔN90}$). Western blot analysis confirmed that both the

full length and truncated proteins are present in whole cell lysates (**Fig. 2A**). However, only full

length PKAR3 remains bound to the DDM-insoluble pellet extracted from these cells, the

truncated protein was present only in the DDM supernatant, suggesting that it no longer binds the

SPMT. Fluorescence microscopy using antibodies against PKAR3 confirmed that the truncated

protein no longer localized to the cortex, being spread throughout the promastigote cell body

(**Fig. 2B**). Conversely, when a plasmid containing the N-terminal 90 amino acids of PKAR3

(with a C-terminal HA-tag) was ectopically expressed in the *Δpkar3* mutant

(Δpkar3::PKAR3$_{N90}$*),* the fusion protein was enriched in the DDM-insoluble pellet (**Fig. 2C**).

Taken together, these results indicate that binding of PKAR3 to the microtubule cytoskeleton is

mediated through the N-terminus containing the FH2-like domain.

To further characterize the association of PKAR3 with the SPMT, we used fluorescence

resonance energy transfer (FRET) to determine if they are located within close proximity in

individual promastigotes (**Fig. 3**). The high value of corrected acceptor FRET intensity observed

when PKAR3 and β-tubulin are co-expressed shows that they co-localize within the SPMT (**Fig.**

3A). The low corrected acceptor FRET intensity in the *Δpkar3* mutant validates the approach

(**Fig. 3B**). Similarly, we saw low corrected acceptor FRET intensity between β-tubulin and an

amino acid transporter (34) located in the cell membrane (**Fig. 3C**). In contrast to the finding

above that N-terminal truncated PKAR3 was not retained in the SPMT pellet after DDM

treatment, there was significant corrected acceptor FRET detected in *Δpkar3::PKAR3*$_{\Delta N90}$ cells

(**Fig. 3D**). However, the number of pixels qualified as "interacting" was ~30% less than obtained

with PKAR3$_{FL}$ (**Fig. 3E**). As the concentration of ectopically expressed PKAR3$_{FL}$ and

PKAR3$_{\Delta N90}$ proteins was high in these cells (**Fig. 2A**), other regions of PKAR3 may also interact

with microtubules (albeit more transiently), especially in the absence of detergent. The FRET

analyses support our hypothesis that PKAR3 is attached to the SPMT *via* a direct or indirect

interaction with β-tubulin.

PKAR3 associates with PKAC3 to form a holoenzyme complex.

Because antibodies against *L. donovani* PKA catalytic submits are not available, we

expressed a Ty1-tagged version of *L. donovani* PKAR3 (LdPKAR3-Ty1) under the control of a

tetracycline repressor in the *Trypanosoma brucei Δpkar1* cell line (7,12). As expected,

expression of LdPKAR3-Ty1 increased in the presence of tetracycline (**Fig. S4A**), with some

"leaky" background expression present in the absence of tetracycline. Although the majority of

LdPKAR3-Ty1 was found in the DDM-soluble supernatant after detergent fractionation of

T. brucei bloodstream form cells, the protein was also present in the DDM-insoluble pellet **(Fig.**

S4B), indicating that the association with the SPMT seen in *L. donovani* might be preserved after

ectopic expression in *T. brucei*. Affinity purification of LdPKAR3-Ty1 from the DDM

supernatant using anti-Ty1 antibody coupled to magnetic beads, followed by Western blotting

with antibodies against the PKA catalytic subunits, showed that LdPKAR3 formed a stable

complex with TbPKAC3 **(Fig. 4A),** but not with TbPKAC1/2 (**Fig. 4B**). Interestingly, up-

regulation of LdPKAR3-Ty1 expression was accompanied by increased TbPKAC3 expression,

possibly due to its stabilization in a complex with R3 (**Fig. S4A**). This result was confirmed by

co-expression of *L. donovani* PKAR3 with *L. donovani* PKAC3 in the *L. tarentolae* LEXSY

expression system (Jena Bioscience, Jena) and purification of the heteromeric complex by

tandem affinity chromatography using an N-terminal His$_6$-tag on PKAR3 and a Strep-tag on

PKAC3 (**Fig. 4C**). As a positive control, we co-purified the *L. donovani* PKAR1*/*C3 complex in

a parallel experiment (**Fig. 4D**). We conclude that PKAR3 and PKAC3 form a stable complex

similar to the PKAR1/PKAC complexes previously reported in *T. brucei* (12) and validate

PKAR3 as subunit of a PKA holoenzyme complex.

To further assess the interaction between PKAR3 and PKAC3 *in vivo*, HA-tagged

PKAC3 was ectopically expressed in *L. donovani* WT and *Δpkar3* promastigotes. Western blot

analysis of whole cell lysates showed that while PKAC3-HA was expressed at similar levels in

both the WT and null mutant; it was detected at high levels in the DDM-insoluble fraction from

WT promastigotes, but not in *Δpkar3* (**Fig. 5A**). These results were confirmed by proteomic

analysis of the DDM-insoluble fraction of WT and *Δpkar3* promastigotes, which showed that the

relative abundance of SPMT-associated PKAC3 was ~30-fold higher in WT than *Δpkar3*

(**Fig. 5B).** In contrast, actin (which is not associated with PKAR3) showed little difference in

relative abundance between samples.

We also employed FRET to confirm the molecular proximity of PKAR3 and PKAC3-HA

in *L. donovani* promastigotes. As shown in the left panel **of Fig. 5C**, the two proteins show a

high value of corrected acceptor FRET intensity. Promastigotes labeled with antibodies against

the donor (PKAR3) or acceptor (PKAC3-HA) alone, displayed no background FRET (**Fig. 5C,**

right panels). These results confirm PKAR3-mediated recruitment of PKAC3 to the SPMT in the

parasite cell cortex.

Structural analysis of the interaction between PKAR3 and PKAC3.

The mammalian PKAR and PKAC subunits have been extensively studied at the atomic

level, both alone and in complex (35)(36)(37) and an X-ray structure is available for *T. cruzi*

PKAR1 (12), allowing us to perform homology modeling based on these structures. Since the N-

terminal sequence of PKAR3 is divergent from human (and other) PKAR subunits and is

predicted to be largely low-complexity and disordered (except for the N-terminal 50 amino

acids), our models were built using only the C-terminal portion including the inhibitor/pseudo-

inhibitor sequence and putative nucleoside binding domains. The conserved residues in

mammalian cNBDs (R_{210} and R_{334} of PKARIα) have been replaced by N_{443} and A_{567} in PKAR3

(**Fig. 6A & B**) and E_{560} in the putative nucleoside binding pocket of latter would likely clash

with cAMP phosphate (**Fig. 6B**). We expressed the C-terminal portion of *L. donovani* PKAR3 in

E. coli, followed by affinity purification and refolding (**Fig. S5)** to produce ligand-free protein

for binding assays by isothermal titration calorimetry (ITC) to test whether it binds cAMP. These

experiments show a complete absence of cAMP binding under these conditions (**Fig. 6C**) and we

found high-affinity binding of the nucleoside analogues 7-cyano-7-deaza-inosine (7CN-7C-Ino,

also known as Jaspamycin) and Toyocamycin (Toyo), with dissociation constants of 1 nM and

3.8 nM, respectively (**Fig. 6D**), similar to those for the cAMP-independent PKAR1 of *T. brucei*

(12).

Electrostatic interaction due to salt-bridges between conserved amino acids have been

shown to be important for interaction between PKAR and PKAC subunits in higher eukaryotes

(38). The salt-bridge triad between R_{240}/D_{266} in bovine PKARIα and R_{194} in PKAC that stabilizes

this interaction appears to be conserved in *L. donovani* PKAR3 (R_{475}/E_{501}) and PKAC3 (R_{166})

(**Fig. 6E**). However, non-conservative replacement of the first residue (R_{95} in PKARIα to S_{324} in

PKAR3) in the inhibitor/pseudo-inhibitor site is predicted to disrupt interaction with negatively

charged and polar residues (T_{23}, E_{99}, E_{142} and Y_{302}) in PKAC3 (**Fig. 6F**). Furthermore, changes

from E_{143}/R_{239} in PKARIα to D_{373}/D_{473} in PKAR3 (**Fig. 6A**) likely prevents formation of a salt

bridge critical for cAMP-induced allosteric activation of PKARIα (38).

The PKAR3/C3 complex is necessary for maintaining the elongated shape of promastigotes.

Microscopic examination revealed that a significant number of cells in the *PKAR3* null

mutant (*Δpkar3*) population lost the normally elongated shape of wild type (WT) promastigotes

and became rounded (although still flagellated; **Fig. 7A**). Ectopic expression of full-length (FL),

PKAR3 largely restored the elongated phenotype, but expression of a truncated (ΔN90) version

did not. Similarly, *Δpkac3* promastigotes were more rounded than WT, while ectopic expression

of full-length *PKAC3* restored the elongated shape. To quantify these changes in shape, we

employed Image Streaming flow cytometry (39) to calculate the cell aspect ratio (width over

length) of 10,000 cells from each population. ImageStream® analysis identified three different

shape groups; elongated (aspect ratio of 0.1-0.4), rounded (aspect ratio of 0.6-1), and ovoid

(aspect ratio of 0.4-0.6)(**Fig. S6**). As expected, elongated cells formed the majority in

promastigotes (**Fig. S6A,** upper panel), while axenic amastigotes was largely rounded (**Fig. S6A,**

lower panel), with the relative numbers of ovoid cells (with an aspect ratio between elongated

and round) being similar in both promastigotes and amastigotes. In contrast, promastigotes from

the *Δpkar3 and Δpkac3* mutants showed a significantly different distribution of aspect ratio from

WT *L. donovani* (**Fig. 7B**). The majority (52%) of WT promastigotes were elongated with 13%

rounded; but only 22% of *Δpkar3* promastigotes are elongated, with the proportion of rounded

cells increasing to 56%. Ectopic expression of full-length *PKAR3* in the mutant cell-line

(*Δpkar3::FL)* partially restored the fraction of elongated cells to 37% and reduced the fraction of

rounded cells to 32%. However, mutants expressing the *PKAR3$_{ΔN90}$* add-back were similar to the

Δpkar3 parent, with 46% being rounded and only 17% elongated. *PKAC3* null mutants (Δ*pkac3*)

also showed a substantial increase in the proportion (53%) of rounded cells with only 14% being

elongated. Ectopic expression of full-length *PKAC3* in the *Δpkac3* mutant partially restored the

WT phenotype (30% rounded and 31% elongated). These results clearly indicate that both

PKAR3 and PKAC3 are critical for maintenance of an elongated cell morphology in *Leishmania*

promastigotes and that recruitment of PKAC3 to the SPMT by the N-terminal domain of PKAR3

is critical for maintaining normal morphogenesis in promastigotes.

Discussion

In this study, we have functionally characterized a novel regulatory subunit of protein kinase A (PKAR3) that recruits a specific catalytic subunit of PKA (PKAC3) to subpellicular microtubules (SPMT) in the cell cortex of *Leishmania*. This interaction is essential for maintaining the elongated shape of promastigotes, since mutants of *PKAR3* (Δ*pkar3*) become rounded (but remain flagellated), as do those lacking *PKAC3*. Furthermore, truncation of a formin homology (FH2)-like domain at the N-terminal end of PKAR3 resulted in detachment of PKAR3 from the SPMT, again leading to rounded promastigotes. We postulate that the PKAC3 phosphorylates one or more proteins in the cell cortex, allowing precise temporospatial regulation of microtubule remodeling throughout the parasite lifecycle.

In fission yeast, PKA activity triggered by glucose starvation promotes microtubule destabilization by phosphorylation of the microtubule rescue factor CLASP/Cls/Peg1, although this process apparently does not involve recruitment by PKAR (40). Microtubule-associated PKA has also been shown to play a critical role in elongation of axons and dendrites by mammalian neurons (41), with PKARIIβ being responsible for the former and PKARIβ for the latter (30). In this system, microtubule-associated protein 2 (MAP2) binds the D/D domain of the PKARs to link them to the microtubules of neurons (42). However, *Leishmania* lacks an obvious orthologue of MAP2, so it appears that the N-terminal FH2-like domain of PKAR3 compensates by recruiting the PKAR3/C3 complex to microtubules, although whether it binds directly or connects *via* a cryptic anchoring protein remains to be seen.

Homologues of PKAC and PKAR are present in most (but not all) eukaryotes (13), with the canonical PKAR containing an N-terminal D/D domain followed by a linker with the

inhibitory sequence and two C-terminal cAMP binding domains. The D/D domain typically

dimerizes to form a four-helix bundle that serves as a docking site for different A-kinase-

anchoring proteins (AKAPs) that allow the PKAR/C complex to be localized to specific sites in

the cell. However, the D/D domain has been lost several times during eukaryotic evolution,

including in trypanosomatids, where PKAR1 contains an RNI-like LRR domain and PKAR3

contains a long, mostly disordered, N-terminal region. Interestingly, *Paratrypanosoma confusum*

(a trypanosomatid parasite isolated from mosquitoes) has an additional paralogue of PKAR (see

Table S1) that is more closely related to those found in other eukaryotes (**Fig. S1**). All three

paralogues are also found in *Bodo saltans* (a free-living kinetoplastid), suggesting that PKAR1

and PKAR3 arose by gene duplication (and subsequent domain shuffling) early in the evolution

of the Kinetoplastea. Interestingly, PKAR3 has been lost in several trypanosomatid genera;

Phytomonas and *Vickermania* (monoxenous insect and/or plant pathogens), as well as four sub-

genera (*Duttonella*, *Herpetosoma*, *Nannomonas* and *Trypanozoon*) of *Trypanosoma (*extra-

cellular parasites of mammals*)*. Since many of these species appear to lack amastigotes, this

pattern is consistent with the hypothesis that PKAR3 plays an important role in regulating

changes in cell morphology during intracellular development.

It has been previously shown that sequence changes in the cNBDs of PKAR1 make it

refractory to cAMP binding (12). Here we show that PKAR3 is equally refractory to cAMP and

that two high-affinity ligands of PKAR1 (7-CN-7C-ino and toyocamycin) also bind PKAR3 with

affinities in the nanomolar range. However, attempts to dissociate the PKAR3-PKAC3 complex

using 7-CN-7-C-ino treatment were unsuccessful, leaving it open whether PKAR3 is a ligand-

controlled inhibitor of PKAC3. Indeed, the first loop of the pseudo-inhibitor domain of PKAR3

contains a serine instead of the arginine conserved in other PKARs, suggesting that it cannot

dock the PKAC3 active site. PKAC3 may remain catalytically active while complexed with

PKAR3, allowing maintenance of the elongated shape of promastigotes. Active PKAR/C

complexes have also been shown in higher mammals (30). It is possible that *Leishmania* (and

other Kinetoplastidae) utilize PKAR3 primarily as an anchoring or tethering subunit to localize a

specific PKA catalytic subunit close to its targets in the cell. In a similar way PKAR1 seems to

tether PKAC1 and 2 to the flagellum in *Leishmania* promastigotes and in *Trypanosoma* (7,18-

20). Whether the nucleoside ligands of PKAR1 and PKAR3 regulate kinase activity of PKAC1/2

and PKAC3, respectively, in the functional context of an upstream pathway remains to be

determined. Alternatively, the temporal control of kinase activity might operate via the catalytic

subunit and the role of PKAR3 be limited to the tethering function, i.e. the spatial control. This is

indeed an attractive speculation as it has been shown that PKAC3 mRNA (previously named c-

lpk2) is downregulated upon induced promastigote to amastigote differentiation with very fast

kinetics (15). This model is perfectly compatible with the observed rounding upon induction of

differentiation and the mutant phenotypes.

In conclusion, we have shown that *Leishmania* (and likely other trypanosomatid

parasites) have re-purposed PKA for a cAMP-independent mechanism to spatiotemporally

regulate microtubule remodeling and cell shape during parasite development upon host infection.

Acknowledgements

We thank the Smoler Proteomics Center at the Technion for all proteomic analyses. Funding for this work was provided by the USA-Israel Binational Science Foundation (BSF), grant number 2017030 to DZ and PJM, the German-Israel Foundation (GIF) grant number I-112-416.10-2017 to DZ and MB, contract HHSN272201700059C from the National Institute of Allergy and Infectious Diseases to PJM, and grant 16GW0281K from the Federal Ministry of Education and Science of Germany to MB.

Author contributions

RFW, SB, RD, MA, VO, GBG, EP, PT and RNK conducted experiments. IQP conducted structure modeling. PJM supervised these efforts and performed the phylogenetic analyses. MB supervised the work in *T. brucei*, *L. tarentolae* and the ITC experiments. DZ supervised the experiments in *L. donovani* and managed the overall project. He also wrote the manuscript together with MB, PJM and SB.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1: PKAR3 is anchored to the DDM-insoluble subpellicular microtubules (SPMT) at the cortex of *L. donovani* cells.

A) Schematic representation of selected *Homo sapiens* (Hsap), *Dictyostelium discoideum* (Ddis), *Saccharomyces cerevisiae* (Scer), *Plasmodium falciparum* (Pfal) and *Leishmania donovani* (Ldon) PKAR orthologues showing the dimerization and docking (D/D) domain (red), RNI-like LRR domain (green), (pseudo-)inhibitor site (brown), and nucleoside binding domains (blue). A lighter shade indicates a significant sequence difference from that of the canonical (human) PKAR. **B**) Indirect immunofluorescence of PKAR3 (upper row), β-tubulin (middle row) and HSP83 (bottom row) in intact promastigote cells (left column) or the DDM-insoluble pellet containing the SPMT (right column). Staining was carried out using antibodies against PKAR3, β-tubulin and HSP83. Fluorescence was detected using a Zeiss LSM 700 inverted confocal microscope, with the scale bar representing 5μm. **C**) Proteins from intact or DDM-extracted (supernatant and pellet) promastigotes were separated on 10% SDS-PAGE and the blotted proteins subjected to Western analysis using the antibodies indicated on the left side of the panels.

Figure 2: PKAR3 binds to the SPMT *via* an N-terminal FH2-like domain.

(**A**) Proteins were extracted from intact and DDM-treated (supernatant and pellet) *L. donovani* promastigotes of the wild type (WT), PKAR3 null mutant (*Δpkar3*), as well as *Δpkar3::FL* and *Δpkar3::ΔN90* cell lines were separated by 10% SDS-PAGE and subsequently subjected to Western blot analysis using anti-PKAR3 (upper row), anti-β tubulin (middle row) and anti-HSP83 (lower row). **B**) Indirect immunofluorescence of PKAR3 in *Δpkar3::ΔN90* promastigotes

using antibody against PKAR3. Fluorescence was detected using a Zeiss LSM 700 inverted

confocal microscope, with the scale bar representing 5μm. **C)** Proteins were extracted from intact

or DDM-treated promastigotes from the *Δpkar3::HA-N90* cell line and subjected to Western blot

analysis using antibodies against the HA-tag (upper row), β tubulin (middle row) and HSP83

(lower row).

Figure 3: PKAR3 associates with tubulin of the SPMT.

(**A**) Promastigotes of *L. donovani* WT were labeled with rabbit anti-PKAR3 and mouse anti-β

tubulin, followed by secondary antibody detection with goat anti-rabbit antibodies conjugated to

Alexa fluor 568 and goat anti-mouse antibodies conjugated to Alexa fluor 647. Fluorescence

Resonance Energy Transfer (FRET) emission was calculated using the Axio Vision program,

with the color scale on the left of each panel indicating the strength of FRET emission. (**B**) Null

mutants of *PKAR3* (*Δpkar3*) were labeled with the same antibodies as above. FRET analysis is

shown in the left panel with cell fluorescence shown in the right panel. (**C**) WT promastigotes

were labeled with rabbit antibodies against the AAP24 transporter and mouse antibodies against

β-tubulin, followed by secondary antibody detection with goat anti-rabbit antibodies conjugated

to Alexa fluor 568 and donkey anti-mouse antibodies conjugated to Alexa fluor 647. FRET

analysis is shown in the left panel and fluorescence in the right panel. (**D**) FRET analysis

between PKAR3 and β-tubulin in cells expressing *Δpkar3::ΔN90*, carried out as in panels A and

B. (**E**) Quantitation of pixels above the interaction threshold FRET value of PKAR3 and β-

tubulin as determined from *Δpkar3::FL* and *Δpkar3::ΔN90* add-back cell lines, as described

above. One-way ANOVA indicated that number of FRET pixels containing *Δpkar3::ΔN90/*β

tubulin were 30% less than *Δpkar3::FL/* β tubulin ($P<0.05$, n=3)

Figure 4: PKAR3 associates with PKAC3.

(**A**) Proteins from the DDM-soluble fractions of the *T. brucei* MiTat 1.2 *PKAR1* null mutant (7) parental cell line (Δ*tbpkar1*) and those expressing an tetracycline-regulatable version of *L. donovani PKAR3* containing a C-terminal Ty1-tag (*LdR3-Ty1*) were immunoprecipitated with mouse monoclonal antibodies against Ty1, followed by Western blotting with rabbit antibodies against *L. donovani* PKAR3 and *T. brucei* PKAC3. The following abbreviations denote the different fractions interrogated: IN (input), FT (flow-through), and E (elution). The sizes of the molecular weight marker are shown in kilodaltons (kDa). An unidentified protein that cross-reacted with anti-PKAC3 antibody is indicated by an asterisk. (**B**) The same samples were probed with antibodies against PKAR3 and PKAC1. (**C**) *L. donovani* PKAR3 and PKAC3 containing N-terminal His_6 and Strep tags, respectively, were co-expressed in *L. tarentolae* and soluble proteins affinity purified on tandem Ni-NTA and Strep-Tactin columns. Aliquots from the input (IN) and eluate (E1, E2) fractions were analyzed by Western blotting with rabbit antibodies against LdPKAR3, as well as mouse monoclonal antibodies against the His (Bio-Rad) and Strep (Qiagen) tags. (**D**) A similar experiment performed using tagged *L. donovani* PKAR1 and PKAC3 and probed with rabbit antibodies against TbPKAR1.

Figure 5: PKAC3 binding to the subpellicular microtubules is mediated by PKAR3.

A) *L. donovani* promastigote wild type (WT) and PKAR3 null mutants (*Δpkar3*) cell lines expressing HA-tagged PKAC3 were subjected to subpellicular microtubule enrichment using 0.5% DDM, and proteins probed by Western blot using antibodies against the HA-tag (upper row), PKAR3 (middle row) and β-tubulin (lower row). **B)** A gel slice containing proteins with a molecular mass range of 35-42 kDa was excised and subjected to mass-spectroscopy using an Orbitrap (Thermo, Ltd). The relative abundance of peptides from PKAC3, PKAR3, and actin

(ACT) are shown for two replicate experiments using WT (■, ■) and *Δpkar3* (■, ■) cell lines.

(**C**) The WT cell line expressing HA-tagged PKAC3 was labeled with rabbit anti-PKAR3 and

mouse anti-HA, followed by goat anti-rabbit antibodies conjugated to Alexa fluor 568 and goat

anti-mouse antibodies conjugated to Alexa fluor 647. The left panels show FRET emission color

and strength as calculated using the Axio Vision program, with the color scale ruler on the left

indicating the strength of interaction between the two labeled proteins. The two right panels

show control experiments using antibody against PKAR3 (donor) or HA-tagged PKAC3

(acceptor) alone, followed by the two secondary antibodies together as in the left panel.

Figure 6: Structural modeling of the PKAR3/C3 complex and ligand binding.

(**A**) Protein sequence alignment of *Bos taurus* PKARIα (Btau_PKARIα) with *L. donovani*

PKAR1 (Ldon_PKAR1) and PKAR3 (Ldon_PKAR3), indicating regions with known structural

importance. Amino acids that are conserved in all three sequences are highlighted in black. (**B**)

Structural modelling of the two nucleotide binding domains of Ldon_PKAR3 (pink) based on

those from Btau_PKARIα (tan), highlighting the most important changes in the conserved amino

acid sequence. (**C**) Binding isotherms from isothermal titration calorimetry (ITC) using refolded

ligand-free recombinant protein containing the C-terminal portion (residues 321-647) of

L. donovani PKAR3 titrated with cAMP. The graph gives the difference power (DP) between the

reference and sample cells upon ligand injection as a function of time. The measurement

corresponds to noise, excluding binding of cAMP. The results shown are from a representative

example of three independent replicates. (**D**) Similar experiments performed with 7CN-7C-

inosine (left) or Toyocamycin (right). The upper panels show the difference power (DP) between

the reference and sample cells after ligand injection as a function of time, while the lower panels

show the total heat exchange per mole of injectant (integrated peak areas from upper panel)

plotted against the molar ratio of ligand to protein. The K_D is the mean of three or more

independent replicates, while the graphs are from a single representative replicate. (**E**) Structural

model of the complex formed between Ldon_PKAR3 (pink) and Ldon_PKAC3 (blue) showing

amino acids predicted to be involved in the salt-bridges and electrostatic interactions downstream

of the pseudo-substrate. (**F**) Side chains of critical residues involved in the PKAR-PKAC

interaction for experimentally determined Btau_PKARIα/Cα [PDBID:2QCS] (left) and predicted

*Ldon*_PKAR3/C3 (right) complexes.

Figure 7: Promastigotes from PKAR3 or PKAC3 mutants are more rounded.

(**A**) Microscopic bright field pictures of cells from late-log cultures of *L. donovani* wild type

(WT), *PKAR3* null mutant (*Δpkar3)*, *PKAC3* null mutant (*Δpk*ac3), and their add-back

derivatives (*Δpkar3::FL*, *Δpkar3::ΔN90*, and *Δpkac3::FL*). (**B**) The percentage of elongated (■)

rounded (■) cells in the samples above as determined by image stream analysis. Ovoid cells

represented ~30% in all cell lines and are not included in the graphs. Asterisks indicate statistical

significance differences from WT of $p < 0.05$ (*) and $p < 0.01$ (**), respectively, in three replicate

experiments.

STAR Methods

Phylogenetic analysis of PKA regulatory subunits. The amino acid sequences of PKAR

orthologues were obtained from 19 kinetoplastid genomes representing 17 different genera and

two unclassified Trypanosomatidae (**Supplementary Table S1**), as well as 8 other eukaryotic

species (**Supplementary Table S2**). In several cases, tBlastn searches of unannotated genomes

were performed to identify the open reading frame(s) encoding PKAR proteins. Protein domains

within each sequence were identified using the InterProScan (43) module of Geneious Prime

(https://www.geneious.com) and/or the HHpred website of the MPI Bioinformatics Toolkit

(31,32). Protein sequences were aligned using the Clustal Omega (44) of Geneious and

phylogenetic trees constructed using the RAxML module implemented in Rapid Bootstrapping

mode with 100 trees using the GAMMA BLOSUM62 substitution matrix (45).

Cell culture. *L. donovani* MHOM/SD/00/1S (28) axenic promastigotes were grown in Earle's-

based medium 199 (M199; Biological Industries, Ltd) supplemented with 10% heat-inactivated

Fetal Bovine Serum (GIBCO, Ltd) and 1% Penicillin-Streptomycin solution (Biological

Industries, Ltd). Trypanosomes were cultivated at 37°C and 5% CO_2 in modified HMI-9 culture

medium (46) supplemented with 10% heat-inactivated fetal bovine serum. Cell density was kept

below 1×10^6 cells/ml by regular dilution.

Detergent enrichment of subpellicular microtubules. Axenic *L. donovani* promastigote (20 ml of

cells at 1×10^7 cells/ml) were washed twice with ice-cold phosphate buffered saline (PBS). The

final pellet was suspended to 200 µl PBS containing protease inhibitors mix (Roche cOmplete

protease inhibitor) and 0.5% n-dodecyl β-D-maltoside (DDM) in Eppendorf tubes. After a short

vortex, the tubes were placed on ice for 15 min, and subsequently centrifuged. The pellet was

washed twice with 1 ml of ice-cold PBS containing the protease inhibitors. For

immunofluorescence analysis, the pellet was suspended in 100 µl PBS. For Western blot

analysis, the pellet was suspended in 100 µl Laemmli buffer. *T. brucei* subpellicular

microtubules were enriched employing the same protocol used for *Leishmania*.

Generation of a polyclonal anti-PKAR3 antibody. The full-length CDS of R3 (encoded by

LinJ.34.2680) was PCR amplified from genomic DNA of *L. infantum* strain JPCM5 using

primers (**Table S2**) designed to introduce an N-terminal His$_6$ tag followed by a TEV protease

cleavage site and cloned *via Bam*HI and *Not*I restriction sites into pETDuet-1 (Novagen, Merck

Millipore). The protein was expressed in *E. coli* Rosetta and purified using Ni-NTA columns

followed by TEV protease cleavage. Recombinant R3 was injected to two rabbits with 5 mg/ml

protein each. Rabbits were boosted four times with 5 mg/ml of the recombinant R3. Injections

and bleedings were carried out as a contracted service (Sigma-Aldrich, Ltd).

Western blotting. Western blot analysis was done as described previously (47), using a 1:1,000

dilution of polyclonal rabbit anti-R3, anti-TbPKAR1 (7), anti-TbPKAC1/C2 (12) and anti-

TbSAXO (25). Rabbit antibodies against TbPKAC3 were used at 1:250 dilution (12). For

Western blot detection of proteins recombinantly expressed in *L. tarentolae*, mouse anti-His

(Bio-Rad) and mouse anti-Strep (Qiagen) were used. The secondary antibodies IRDye680LT

goat anti-rabbit (1:50000; LICOR) and IRDye800CW goat anti-mouse (1:10000; LICOR) were

used for detection with the Odyssey^{TM} CLx imaging system (LICOR). Rabbit antiserum against

L. donovani HSP83 (28) or β-tubulin (cell signalling, Ltd) were used as protein loading marker.

Gene knockouts. The *PKAR3* gene was deleted from the 1S-2D clone of *L. donovani* using

homologous recombination as described previously by Inbar et al. (34). PCR analyses of genomic

DNA was used to confirm the absence of the targeted gene (**Fig. S1A**). Polyclonal antibodies raised

in rabbits against recombinant PKAR3 reacted with a protein of the correct molecular mass

(72 kDa) in Western blots of wild type cells, but not *Δpkar3* mutants (**Fig. S1B**). The protein was

present (at slightly elevated levels) in add-back mutants (*Δpkar3::*FL) ectopically expressing full-

length R3. The *PKAC3* gene was deleted using the same strategy as above. PCR analyses of

L. donovani genomic DNA extracted from single colonies confirmed that both copies of the gene

are missing from *Δpkac3*, while the antibiotic resistance genes that were inserted to replace *PKAC3*

are in place (**Fig. S7**).

Immunofluorescence. Indirect immunofluorescence analysis was carried out following the

methods described by Inbar et al. (34). Briefly, late log phase promastigotes were washed twice

in phosphate buffered saline (PBS) and then fixed in 1% formaldehyde/PBS on a slide for 10 min

before permeabilization by exposure to 0.2% TritonX-100/PBS for 10 min. Cells were incubated

with blocking solution [5% (v/v) non-fat dried skimmed milk powder/PBST] for 30 min at room

temperature, incubated with rabbit anti-PKAR3 (this study), rabbit anti-promastigote membrane

proteins (28) or mice anti-HA tag antibodies (1:500 dilution) for one hour. Subsequently, slides

were washed three times with PBS-Tween and incubated with the fluorescent secondary

antibody in darkness for 30 min. Slides were washed three time with PBS-Tween and a drop of

DAPI Fluoromount G (Southern Biotech Ltd) was added. Slides were covered with slips, sealed,

and then kept in darkness. Subsequently, they were examined using a Zeiss LSM 700 inverted

confocal laser scanning microscope. Image processing was done using Zen lite software, Zeiss.

Fluorescence Resonance Energy Transfer (FRET). Late log phase *L. donovani* promastigotes

ectopically expressing PKAR3 and PKAC3-HA were fixed in PBS containing 4%

paraformaldehyde and then settled on slides. PKAR3 was labelled using polyclonal anti-PKAR3

antibody (1:500), followed by detection with goat anti-rabbit Alexa 568 (1:500). PKAC3 was

labelled with anti-HA antibody (1:1000, BioLegend, Ltd) and secondary goat anti-mouse Alexa

647 (1:500). Confocal microscopy was obtained using a Zeiss LSM 700 Inverted confocal laser

scanning microscope. An excitation wavelength of 578 nm and an emission wavelength of 640

nm and below were used for Alexa568, whereas an excitation wavelength of 651 nm and an

emission wavelength of 640 nm and above were used for Alexa647. FRET was assessed with

AxioVision FRET software, using the mathematical approach - Youvan's technique. Youvan's

technique (Fc) is the basis of all correction measurement techniques (out of 4 techniques) in

AxioVision FRET. It refers to corrected FRET values when three FRET filter sets are used. The

measured values are corrected for the background (bg) and for the crosstalk from the donor (don)

and the acceptor (acc). Calculation of the intensity values for FRET follows the formula: Fc =

(fretgv - bg) - cfdon (dongv - bg) - cfacc (accgv - bg), where gv is the fluorescence value in

every pixel and cf is a constant, which is calculated for each of the samples with only one

secondary antibody. Subsequently, the number of interacting molecules in the FRET images

were counted using the microscopic analysis software, IMARIS version 9.3.

Image streaming flow cytometry. Analysis was carried out using Amnis® ImageStream®X Mk

II)(48)(49)(50). The Image Streamer was calibrated by feeding it with late log phase axenic

promastigotes of wild type and mutants. Late log phase *L. donovani* promastigotes were washed

once using ice cold phosphate buffered saline (PBS). Subsequently, cells were fixed by

suspending them in PBS containing 4% paraformaldehyde. An aliquot of these cell suspension

was injected into the Imagestream Mark II. Each cell was subjected to a snapshot in bright field.

The data was then analyzed using IDEAS software, version 6.10, with compensation according

to the software standards. Cells were gated for single cells in focus. Further, cells were classified

based on shape (elongated, ovoid, round) by considering aspect ratio vs symmetry features.

Aspect ratio feature assesses the elongatedness of cells by height/width ratio. Symmetry 2

measures the tendency of the object to have single axis of elongation.

Inducible overexpression of PKAR3 in Trypanosoma brucei. The *PKAR3* CDS was PCR

amplified from genomic DNA of *L. donovani* promastigotes using primers R3_HindIII_fw and

R3_lr_Ty1_BamHI_rev (**Table S2**) to introduce a C-terminal Ty1 tag. The PCR product was

cloned into the *Hin*dIII and *Bam*HI sites of pHD615 (51), which was then linearized with *Not*I

for transfection of a previously described homozygous *T. brucei PKAR1* deletion mutant (7) that

additionally expresses a double tetracycline repressor (pHD1313; (52). Transfected cells were

selected with 2 µg/ml blasticidin (pHD615) and 2.5 µg/ml phleomycin (pHD1313). PKAR3

expression was induced with 1 µg/ml tetracycline.

Immunoprecipitation in T. brucei. Immunoprecipitation of PKAR3-Ty1 was carried out by

binding anti-Ty1 (53) to magnetic protein A beads (Dynabeads, Invitrogen) followed by a 2-hour

incubation with 1×10^8 trypanosomes lysed in lysis buffer (10 mM Tris/Cl pH 7.5; 150 mM

NaCl; 0.5 mM EDTA; 0.5% NP-40; Roche cOmplete protease inhibitor) for 30 min at 4°C.

Beads were washed 4x with lysis buffer and proteins were eluted by incubation with 50 µl 2×

Laemmli sample buffer for 5 min at 95°C.

Tandem affinity purification from Leishmania tarentolae. PKAR1 and *PKA*R3 were N-terminally

tagged with a hexa-histidine peptide by PCR amplification from *L. donovani* genomic DNA. The

PCR products were inserted in the pLEXSY_I-ble3® vector (Jena Bioscience). PKAC3 was N-

terminally strep-tagged and inserted into pLEXSY_I-neo3®. The vectors were then linearized

with *Swa*I and stably co-transfected for holoenzyme expression (PKAR1 and PKAC3 or PKAR3

and PKAC3) into the LEXSY T7-TR® cell line (Jena Bioscience). Protein co-expression, parasite

cultivation and protein purification were performed as described in (12) with a few

modifications. Briefly, co-expression was induced with 10 µg/ml tetracycline for 48 hours. The

cells were harvested by centrifugation (2000×g for 5 min), washed with PBS, and lysed in His

binding buffer (50 mM NaH_2PO_4 pH 8, 300 mM NaCl, 10 mM imidazole, 1% Triton-X, protease

inhibitor cocktail). The soluble fraction was loaded on to a gravity flow Ni-NTA column

(Thermo Fisher Scientific). The column was washed twice with His wash buffer (50 mM

NaH_2PO_4 pH 8, 300 mM NaCl, 20 mM imidazole) prior to elution of the kinase complex with

His elution buffer (50 mM NaH_2PO_4 pH 8, 300 mM NaCl, 250 mM imidazole). The Ni-NTA

eluate was loaded on to a gravity flow Strep-Tactin column (IBA), which was then washed with

Strep wash buffer (50 mM NaH_2PO_4 pH 8, 300 mM NaCl). Lastly, the kinase complexes were

eluted with Strep elution buffer (50 mM NaH_2PO_4 pH 8, 150 mM NaCl, 50 mM biotin). The

integrity of the co-expressed holoenzyme complexes was analysed by western blot.

Pulldown of subpellicular microtubules-bound proteins (proteomics). These experiments aimed

to determine the levels of PKAC3 binding to the subpellicular microtubules of *L. donovani* wild

type (WT) and *Δpkar3*. Microtubules of WT and *Δpkar3* promastigotes were DDM-enriched as

described above. Subsequently, proteins associated with this fraction were separated on 9% SDS-

PAGE and a slice containing proteins at the molecular mass range of 35-42 kDa was excised and

subsequently subjected to mass spectrometry. The proteins in the gel were reduced with 3 mM

DTT (60ºC for 30 min), modified with 10 mM iodoacetamide in 100 mM ammonium

bicarbonate (in the dark, room temperature for 30 min) and digested in 10% acetonitrile and 10

mM ammonium bicarbonate with modified trypsin (Promega) at a 1:10 enzyme-to-substrate

ratio, overnight at 37ºC. The resulted peptides were desalted using C18 tips (Homemade stage

tips) dried and resuspended in 0.1% formic acid**.** The peptides were resolved by reverse-phase

chromatography on 0.075 X 180-mm fused silica capillaries (J&W) packed with Reprosil

reversed phase material (Dr Maisch GmbH, Germany). The peptides were eluted with linear 60

minutes gradient of 5 to 28%, 15 minutes gradient of 28 to 95% and 15 minutes at 95%

acetonitrile with 0.1% formic acid in water at flow rates of 0.15 μl/min. Mass spectrometry was

performed by Q Exactive plus mass spectrometer (Thermo) in a positive mode using repetitively

full MS scan followed by collision induced dissociation (HCD) of the 10 most dominant ions

selected from the first MS scan. The mass spectrometry data was analyzed using Proteome

Discoverer 1.4 software with Sequest (Thermo) algorithm against *L. donovani* and *L. infantum*

proteomes from TriTryp database with mass tolerance of 20 ppm for the precursor masses and

0.05 Da for the fragment ions (TriTrypDB). Oxidation on Met, and Phosphorylation on Ser, Thr,

Tyr were accepted as variable modifications and Carbamidomethyl on Cys was accepted as static

modifications. Minimal peptide length was set to six amino acids and a maximum of two

miscleavages was allowed. Peptide- and protein-level false discovery rates (FDRs) were filtered

to 1% using the target-decoy strategy. Semi-quantitation was done by calculating the peak area

of each peptide based on its extracted ion currents (XICs) and the area of the protein that is the

average of the three most intense peptides from each protein. The mass spectrometry proteomics

data have been deposited to the ProteomeXchange Consortium via the PRIDE (54) partner

repository with the dataset identifier PXD025222.

Molecular modelling. PKAR3 homology models were built from each of the PDB

templates 2QCS, 5JR7 and 4MX3 with Chimera Modeller (55,56). Multiple sequence

alignments were produced with Clustal Omega (44). The PKAR3/C3 complex was predicted

by AlphaFold2 using MMseqs2 on ColabFold v1.5.2, with pdb70

templates, alphafold2_multimer_v3 weights for complex predictions using 20 recycles

and tolerance set to 0.5, followed by AMBER relaxation (57). Structure images were produced

with UCSF Chimera (58).

Ligand binding studies by isothermal titration calorimetry (ITC). Protein purification was

performed as described by (12) with the following modification: N-terminally truncated

LdPKAR3 (aa 321-647) was cloned into pET11_Sumo3 (NEB) with an N-terminal Sumo3-tag

and expressed in *E. coli* Rosetta (DE3). Purification of R3(321-647) by Ni-NTA affinity

chromatography was followed by SenP2 protease mediated cleavage of the N-terminal Sumo3-

tag during dialysis of the protein in 50mM HEPES pH 7.5 and 50mM NaCl. The cleaved protein

was either stored at -80°C or directly denatured and separated from prebound ligands as

described (12). For ligand binding studies by ITC the protein was further dialyzed against 50mM

Tris-Cl pH 8.5, 9.6mM NaCl, 0.4mM KCl, 2mM MgCl2, 2mM CaCl2, 0.5M Arginine, 0.75M

Guanidine-HCl, 0.4M Sucrose, 10mM DTT over night at 4°C and subsequently purified by gel

filtration chromatography on a Superdex Increase 10/300 GL column (GE healthcare). Refolded

and purified proteins were diluted to 10-20 µM in ITC buffer (50mM HEPES pH 7.5 50mM

NaCl and 1% DMSO). 100-200 µM 7-cyano-7-deaza-inosine (7CN-7C-ino) or Toyocamycin

(Toyo) was diluted in the same buffer. ITC measurements were carried out on a MicroCal

PEAQ-ITC (Malvern instrument). 2-4 µl of ligand were injected in a series of 13-19 injections

into the protein sample at 298K. The Differential Power (DP) between reference and sample cell

was maintained at 8-10 µcal s^{-1} in all experiments. Data analysis was performed with the

MicroCal PEAQ-ITC software applying a model with one binding site.

Bibliography

1. L. Maxfield, J. S. Crane, *Leishmaniasis* (StatPearls Publishing, 2020).

2. J. Alvar, *et al.*, Leishmaniasis worldwide and global estimates of its incidence. *PLoS One* **7**, e35671 (2012).

3. K. P. Chang, D. M. Dwyer, Multiplication of a human parasite (*Leishmania donovani*) in phagolysosomes of hamster macrophages in vitro. *Science* **193**, 678–680 (1976).

4. B. L. Herwaldt, Leishmaniasis. *Lancet* **354**, 1191–1199 (1999).

5. E. Barak, *et al.*, Differentiation of *Leishmania donovani* in host-free system: analysis of signal perception and response. *Mol Biochem Parasitol* **141**, 99–108 (2005).

6. D. Zilberstein, Lysosome Sensing Is a Key Mechanism in *Leishmania* Intracellular Development. *Front. Microbiol.* **12**, 667807 (2021).

7. S. Bachmaier, *et al.*, Protein kinase A signaling during bidirectional axenic differentiation in *Leishmania*. *Int J Parasitol* **46**, 75–82 (2016).

8. P. Tsigankov, *et al.*, Regulation dynamics of *Leishmania* differentiation: deconvoluting signals and identifying phosphorylation trends. *Mol Cell Proteomics* **13**, 1769–1786 (2014).

9. S. S. Taylor, R. Ilouz, P. Zhang, A. P. Kornev, Assembly of allosteric macromolecular switches: lessons from PKA. *Nat Rev Mol Cell Biol* **13**, 646–658 (2012).

10. W. Wong, J. D. Scott, AKAP signalling complexes: focal points in space and time. *Nat Rev Mol Cell Biol* **5**, 959–970 (2004).

11. J. Bubis, *et al.*, The gene product of a *Trypanosoma equiperdum* ortholog of the cAMP-dependent protein kinase regulatory subunit is a monomeric protein that is not capable of binding cyclic nucleotides. *Biochimie* **146**, 166–180 (2018).

12. S. Bachmaier, *et al.*, Nucleoside analogue activators of cyclic AMP-independent protein
kinase A of Trypanosoma. *Nat. Commun.* **10**, 1421–1435 (2019).

13. M. Peng, *et al.*, Spatial Organization in Protein Kinase A Signaling Emerged at the Base
of Animal Evolution. *J Proteome Res* **14**, 2976–2987 (2015).

14. S. Bachmaier, M. Boshart, "Kinetoplastid AGC kinases" in *Protein Phosphorylation in*
Eukaryotic Parasites: Potential for Chemotherapy, D. Doerig, C. S. Spaeth, M. Wiese, P.
M. Selzer, Eds. (Wiley-Blackwell, 2013), pp. 99–122.

15. M. M. Siman-Tov, A. Radi, M. Shapira, C. L. Jaffe, Cloning from *Leishmania major* of a
developmentally regulated gene, c-lpk2, for the catalytic subunit of the cAMP-dependent
protein kinase. *Mol. Biochem. Parasitol.* **77**, 201–215 (1996).

16. M. M. Siman-Tov, A. C. Ivens, C. L. Jaffe, Molecular cloning and characterization of two
new isoforms of the protein kinase A catalytic subunit from the human parasite
Leishmania. *Gene* **288**, 65–75 (2002).

17. P. Tsigankov, P. F. Gherardini, M. Helmer-Citterich, G. F. Spath, D. Zilberstein,
Phosphoproteomic analysis of differentiating *leishmania* parasites reveals a unique stage-
specific phosphorylation motif. *J Proteome Res* **12**, 3405–3412 (2013).

18. M. Oberholzer, *et al.*, Independent analysis of the flagellum surface and matrix proteomes
provides insight into flagellum signaling in mammalian-infectious *Trypanosoma brucei*.
Mol Cell Proteomics **10**, 1–14 (2011).

19. C. Fort, S. Bonnefoy, L. Kohl, P. Bastin, Intraflagellar transport is required for the
maintenance of the *Trypanosome* flagellum composition but not its length. *J. Cell Sci.*
129, 3026–41 (2016).

20. D. Chiasson, *et al.*, A unified multi-kingdom Golden Gate cloning platform. *Sci. Rep.* **9**,

10131 (2019).

21.   D. Rosenzweig, *et al.*, Retooling *Leishmania* metabolism: from sand fly gut to human

macrophage. *FASEB J* **22**, 590–602 (2008).

22.   K. Gull, The cytoskeleton of trypanosomatid parasites. *Annu Rev Microbiol* **53**, 629–655

(1999).

23.   L. Kohl, K. Gull, Molecular architecture of the *trypanosome* cytoskeleton. *Mol Biochem*

Parasitol **93**, 1–9 (1998).

24.   D. M. Dwyer, Isolation and partial characterization of surface membranes from

Leishmania donovani promastigotes. *J Protozool* **27**, 176–182 (1980).

25.   D. Dacheux, *et al.*, A MAP6-related protein is present in protozoa and is involved in

flagellum motility. *PLoS One* **7**, e31344 (2012).

26.   D. Robinson, P. Beattie, T. Sherwin, K. Gull, Microtubules, tubulin, and microtubule-

associated proteins of trypanosomes. *Methods Enzym.* **196**, 285–299 (1991).

27.   T. Sherwin, K. Gull, The cell division cycle of *Trypanosoma brucei brucei*: timing of

event markers and cytoskeletal modulations. *Philos Trans R Soc L. B Biol Sci* **323**, 573–

588 (1989).

28.   Y. Saar, *et al.*, Characterization of developmentally-regulated activities in axenic

amastigotes of Leishmania donovani. *Mol Biochem Parasitol* **95**, 9–20 (1998).

29.   M. Vicente-Manzanares, F. Sanchez-Madrid, Cell polarization: a comparative cell biology

and immunological view. *Dev.Immunol.* **7**, 51–65 (2000).

30.   R. Ilouz, *et al.*, Isoform-specific subcellular localization and function of protein kinase A

identified by mosaic imaging of mouse brain. *Elife* **6** (2017).

31.   L. Zimmermann, *et al.*, A Completely Reimplemented MPI Bioinformatics Toolkit with a

New HHpred Server at its Core. *J. Mol. Biol.* **430**, 2237–2243 (2018).

32. F. Gabler, *et al.*, Protein Sequence Analysis Using the MPI Bioinformatics Toolkit. *Curr.*

Protoc. Bioinforma. **72** (2020).

33. J. Faix, R. Grosse, Staying in Shape with Formins. *Dev. Cell* **10**, 693–706 (2006).

34. E. Inbar, D. Schlisselberg, M. Suter Grotemeyer, D. Rentsch, D. Zilberstein, A versatile

proline-alanine transporter in the unicellular pathogen *Leishmania donovani* regulates

amino acid homeostasis and osmotic stress responses. *Biochem J* **449**, 555–566 (2013).

35. J. G. Bruystens, *et al.*, Structure of a PKA RIalpha Recurrent Acrodysostosis Mutant

Explains Defective cAMP-Dependent Activation. *J Mol Biol* **428**, 4890–4904 (2016).

36. J. G. Bruystens, *et al.*, PKA RIα homodimer structure reveals an intermolecular interface

with implications for cooperative cAMP binding and Carney complex disease. *Structure*

22, 59–69 (2014).

37. C. Kim, C. Y. Cheng, S. A. Saldanha, S. S. Taylor, PKA-I holoenzyme structure reveals a

mechanism for cAMP-dependent activation. *Cell* **130**, 1032–1043 (2007).

38. E. P. Barros, *et al.*, Electrostatic Interactions as Mediators in the Allosteric Activation of

Protein Kinase A RIα. *Biochemistry* **56**, 1536–1545 (2017).

39. S. E. Headland, H. R. Jones, A. S. V. D'Sa, M. Perretti, L. V., Cutting-edge analysis of

extracellular microparticles using ImageStream(X) imaging flow cytometry. *Sci. Rep.* **4** ,

2537-2541 (2014).

40. M. Kelkar, S. G. Martin, ARTICLE PKA antagonizes CLASP-dependent microtubule

stabilization to re-localize Pom1 and buffer cell size upon glucose limitation (2015)

https:/doi.org/10.1038/ncomms9445 (December 16, 2022).

41. Y. A. Huang, *et al.*, Microtubule-associated type II protein kinase A is important for

neurite elongation. *PLoS One* **8**, e73890 (2013).

42. A. Harada, J. Teng, Y. Takei, K. Oguchi, N. Hirokawa, MAP2 is required for dendrite

elongation, PKA anchoring in dendrites, and proper PKA signal transduction. *J Cell Biol*

158, 541–549 (2002).

43. M. Blum, *et al.*, The InterPro protein families and domains database: 20 years on. *Nucleic*

Acids Res. **49**, D344–D354 (2021).

44. F. Sievers, D. G. Higgins, Clustal Omega for making accurate alignments of many protein

sequences. *Protein Sci* **27**, 135–145 (2018).

45. A. Stamatakis, RAxML version 8: A tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

46. E. Vassella, b Reuner, b Yutzy, M. Boshart, Differentiation of African Trypanosones is

controlled by density sensing mechanism which signals cell cycle arrest via the cAMP

pathway. *J. Cell Sci.* **110**, 2661–2671 (1997).

47. I. Darlyuk, *et al.*, Arginine homeostasis and transport in the human pathogen Leishmania

donovani. *J Biol Chem* **284**, 19800–19807 (2009).

48. Y. Lee, Q. Wang, I. Shuryak, D. J. Brenner, H. C. Turner, Development of a high-

throughput γ-H2AX assay based on imaging flow cytometry. *Radiat. Oncol.* **14**, 150

(2019).

49. M. Oliveira, *et al.*, First report of the in vitro antileishmanial properties of extremophile

plants from the Algarve Coast. *Nat Prod Res* **32**, 600–604 (2018).

50. M. A. Rodrigues, Automation of the *in vitro* micronucleus assay using the Imagestream®

imaging flow cytometer. *Cytom. Part A* **93**, 706–726 (2018).

51. S. Biebinger, L. Elizabeth Wirtz, P. Lorenz, C. Clayton, Vectors for inducible expression

of toxic gene products in bloodstream and procyclic *Trypanosoma brucei*. *Mol. Biochem.*

Parasitol. **85**, 99–112 (1997).

52. V. P. Alibu, L. Storm, S. Haile, C. Clayton, D. Horn, A doubly inducible system for RNA

interference and rapid RNAi plasmid construction in *Trypanosoma brucei*. *Mol. Biochem.*

Parasitol. **139**, 75–82 (2005).

53. P. Bastin, A. Bagherzadeh, K. R. Matthews, K. Gull, A novel epitope tag system to study

protein targeting and organelle biogenesis in *Trypanosoma brucei*. *Mol. Biochem.*

Parasitol. **77**, 235–239 (1996).

54. Y. Perez-Riverol, *et al.*, The PRIDE database and related tools and resources in 2019:

Improving support for quantification data. *Nucleic Acids Res.* **47**, D442–D450 (2019).

55. E. C. Meng, E. F. Pettersen, G. S. Couch, C. C. Huang, T. E. Ferrin, Tools for integrated

sequence-structure analysis with UCSF Chimera. *BMC Bioinformatics* **7**, 339 (2006).

56. B. Webb, A. Sali, Comparative Protein Structure Modeling Using MODELLER. *Curr*

Protoc Bioinforma. **47**, 5 6 1-32 (2014).

57. M. Mirdita, *et al.*, ColabFold: making protein folding accessible to all. *Nat. Methods* **19**,

679–682 (2022).

58. E. F. Pettersen, *et al.*, UCSF Chimera--a visualization system for exploratory research and

analysis. *J Comput Chem* **25**, 1605–1612 (2004).

Figure 1

A
Hsap_PKARIα
Scer_BCY1
Ddis_PKAR
Pfal_PKAR
Ldon_PKAR1
Ldon_PKAR3

B Intact promastigote DDM-insoluble
        cells   pellet

PKAR3

β-tubulin

HSP83

C

                                    DDM
                         Intact ⌐   ⌐
              KDa   cells   Pellet   Sup

PKAR3   75 →

β-tubulin   55 →

HSP83   83 →

Figure 2

Figure 3

A

Δpkar3::*FL*
anti-PKAR3 + anti-β-tubulin

B

Δpkar3
anti-PKAR3 + anti-β-tubulin

C

WT
anti-AAP24 + anti-β-tubulin

D

Δpkar3::*ΔN90*
α PKAR3+α β-tubulin

E

Figure 4

Figure 5

Figure 6

Figure 7