Article Summary Line: Transmission of SARS-CoV-2 between people and their pets is

common, but can be mitigated by measures to reduce human-animal contact.

Running Title: SARS-CoV-2 transmission between people and pets

Keywords: SARS-CoV-2; COVID-19; zoonoses; One Health; anthropozoonoses; household

transmission

Word count: 146 (abstract); 3,499 (text)

Household transmission of SARS-CoV-2 from humans to pets in Washington and Idaho:

burden and risk factors

Authors: Julianne Meisner, BVM&S MS[a*]; Timothy V. Baszler, DVM PhD[b,c], Kathryn H.

Kuehl, DVM[d], Vickie Ramirez, MA[a], Anna Baines, DVM[a], Lauren A. Frisbie, MPH[e], Eric T.

Lofgren, PhD[c], David M. DeAvila MS[b], Rebecca M. Wolking BS[b], Dan S. Bradway BS[b, c],

Hannah Wilson BS[b], Beth Lipton DVM[f], Vance Kawakami DVM[f], Peter M. Rabinowitz, MD

MPH[a]

Affiliations:

[a]University of Washington, Seattle, WA, USA

[b]Washington Animal Disease Diagnostic Laboratory, Washington State University, Pullman,

WA, USA

[c]Paul G. Allen School for Global Health, Washington State University, Pullman, WA, USA

[d]Department of Veterinary Clinical Sciences, Washington State University, Pullman, WA, USA

[e]Washington State Department of Health, Tumwater, WA, USA

[f] Public Health – Seattle & King County, Seattle, WA, USA

Abstract

SARS-CoV-2 is believed to have emerged from an animal reservoir; however, the frequency of

and risk factors for inter-species transmission remain unclear. We carried out a community-based

study of pets in households with one or more confirmed SARS-CoV-2 infection in humans.

Among 119 dogs and 57 cats with completed surveys, clinical signs consistent with SARS-CoV-

2 were reported in 20 dogs (21%) and 19 cats (39%). Out of 81 dogs and 32 cats sampled for

testing, 40% of dogs and 43% of cats were seropositive, and 5% of dogs and 8% of cats were

PCR positive; this discordance may be due to delays in sampling. Respondents commonly

reported close human-animal contact and willingness to take measures to prevent transmission to

their pets. Reported preventative measures showed a slightly protective trend for both illness

and seropositivity in pets, while sharing of beds and bowls had slight harmful effects.

Background

Coronaviruses infect multiple mammalian species, and SARS-CoV-2 virus, the etiological agent of COVID-19 infection, likely jumped to humans from a mammalian source [1]. While currently the virus is spreading person to person, the ACE2 receptor involved in SARS-CoV-2 transmission is present in multiple species and there are numerous reports of infections in pets [2–4]. Currently, 110 domestic cats and 95 domestic dogs in the USA have been reported by USDA-APHIS to have SARS-CoV-2 infection [5]. Workplace transmission of SARS-CoV-2 between humans and animals has also been documented, including in zoos (felids and non-human primates) and on mink farms [6,7]. This is consistent with previous reports of SARS-CoV-1 infecting cats and ferrets, and laboratory studies demonstrating experimental SARS-CoV-2 infection of non-human primates, ferrets, hamsters, and rabbits [8]. Less is known, however, about the frequency of and risk factors for SARS-CoV-2 transmission between humans and companion animals in a household setting. Furthermore, the natural history of SARS-CoV-2 infection in pets is poorly understood.

Given the close contact many people have with their pets and the intimate nature of their shared environment, exacerbated during periods of human quarantine or isolation, it is important to better understand the role of companion animals in community infection patterns, including contribution to virus evolution and emergence of novel strains. In light of evidence from mink farms that animal-origin variants may contain spike mutations and other changes that could affect clinical features of infection [9,10], recent evidence suggesting mouse origins of the omicron variant [11], and Hong Kong's recent decision to cull 2,000 hamsters after a pet shop worker was infected with the delta variant [12], ongoing monitoring of SARS-CoV-2

transmission between humans and animals in household and other human-animal contact settings

remains critical.

We report our findings from the COVID-19 and Pets Study (CAPS), an ongoing cross-

sectional community-based study of pets in households of persons with documented COVID-19

infection. The goal of the study is to describe the frequency of transmission between humans and

animals within a household, and to determine human, animal, and environmental risk factors for

that transmission, in a One Health framework.

Methods

The COHERE [13] and STROBE [14] statements were used to guide reporting of the

findings and the preparation of this manuscript.

Study population

We defined a household as one or more persons ages 18 or older, co-housing with at least

one pet that does not live solely outdoors. Pets were defined as dogs, cats, ferrets, and hamsters

based on prior research documenting experimental COVID-19 infection in these species [15,16].

We conducted this study in King, Snohomish, Yakima, Whitman, Pierce, Spokane, and

Benton counties in Washington, and Latah County in Idaho. This paper reports on sampling

conducted from April 2020 to September 2021.

Study design

CAPS is a cross-sectional study with individual- and household-level data collection.

Study participation involved two components, detailed below: an online survey followed by

animal sampling.

Recruitment and eligibility

Households were recruited through partnerships with other COVID-19 clinical trials and

community studies, social media, word of mouth, community partners, and by contact tracers

from Public Health—Seattle & King County during case investigation/contact tracing calls.

Individuals were screened for eligibility using the UW Research Electronic Data Capture

(REDCap) system [17], a HIPAA-compliant web tool for clinical research, with criteria

including county of residence, pet ownership, and one or more household member with

confirmed SARS-CoV-2 infection via PCR or antigen testing by a provider or laboratory.

Animals with known fearful and/or aggressive behavior were excluded, however, other animals

in the corresponding household were eligible.

Ethical approvals

This study and its protocols received ethical approval from the University of

Washington's Institutional Review Board STUDY00010585) and Office of Animal Welfare

(PROTO201600308: 4355-01). Informed consent was obtained via REDCap, or over the phone

with the study coordinator, after the nature and possible consequences of study involvement had

been explained. Once eligibility was confirmed and consent was obtained, individuals completed

the online survey.

Survey

A comprehensive survey was completed by a household member prior to scheduling of

the sampling visit. Surveys were completed online by the study participant using the REDCap

interface, or via phone with the study coordinator. Human items included COVID-19 symptom

onset, specific symptoms experienced and severity; comorbidities; vaccination status including

dates and type; and reported COVID-19-like illness of any other household members, including

those who did not have confirmatory testing. Animal items, stratified on individual animal,

included veterinary clinical variables, history of illness compatible with SARS-CoV-2 infection,

and contact between individual animals and individual members of the household.

Environmental items included type and size of home, type of flooring (carpet, wood, etc.), and

availability of outdoor space for pets to roam.

A second survey was completed verbally at the time of sampling on any changes in the

clinical status of human and animal household members since the REDCap survey was

completed, including new hospitalizations, symptoms, or COVID-19 diagnoses. Confirmation of

SARS-CoV-2 test date and positive result was also performed through review of test results by

the sampling team. Self-test results were not accepted.

<u>Animal sampling</u>

Sampling was performed by a team of two study personnel including at least one

veterinarian. In most cases sampling was conducted at the participant's home; however, several

animals were tested at veterinary hospitals. No chemical restraint was used, nor muzzles due to

biosafety concerns.

Species-appropriate restraint was employed using standard techniques to allow for

venipuncture and collection of 3 mL of blood into a labeled serum separator tube. Following

venipuncture, swab samples were collected from both rostral nares/nasal passage and the caudal

oropharynx and then placed into one Primestore Molecular Transport Medium (MTM) tube. A

separate fecal swab was collected from the rectum and placed into a separate Primestore MTM

tube. All participants received educational information from the field team about measures to

mitigate household COVID-19 transmission.

Swab and serum samples were transported on ice within 24 hours to the Washington

Animal Disease Diagnostic Laboratory (WADDL) for PCR and antibody testing.

<u>Testing</u>

SARS-CoV-2 RT-PCR

Respiratory or fecal swabs: RNA extraction and SARS-Cov-2 reverse transcriptase (RT)

real-time PCR was performed as described [18]. Following initial viral detection by PCR, three

dog samples and one cat sample were submitted to University of Minnesota Genomics Center

(Oakdale, MN 55128) for whole genome sequencing (WGS) [19]. A second cat sample was

submitted to the USDA National Veterinary Services Laboratories (NVSL) in Ames Iowa for

WGS. Mutational analysis was performed using the GISAID EpiFlu Database CoVsurver:

Mutation Analysis of hCoV-19 [20,21]. All five sequences were deposited into GISAID, with

accession numbers EPI_ISL_7845315, EPI_ISL_7845316, EPI_ISL_7845317,

EPI_ISL_7845318, and EPI_ISL_8897004. SARS-CoV-2 lineages were assigned using the

Phylogenetic Assignment of Named Global Outbreak LINeages (Pango lineage) tool [22,23].

SARS-CoV-2 Spike Protein Receptor Binding Domain (RBD) ELISA

WADDL developed canine and feline SARS-CoV-2 ELISA assays using recombinant

SARS-CoV-2 Spike Receptor Binding Domain protein as antigen (S-RBD). The recombinant

RBD was obtained from the UW Center for Emerging and Reemerging Infectious Disease

(CERID) laboratory of Dr. Wesley Van Voorhis through an institutional Material Transfer

Agreement. WADDL used an in-house standard operating procedure for indirect ELISA of

SARS-CoV-2 in 96-well format based on a previous publication in humans [24]. The major

components of the assay included: 1) rS-RBD coating of plates as target antigen (2ug/ml in

Sigma Carbonate-Bicarbonate Buffer); 2) 1:100 dilution of test sera (diluted in ChronBlock

ELISA Buffer-Chondrex Inc.); 3) anti dog IgG-HRP as linker (Southern BioTech goat anti-

canine IgG) and 4) Sigma (TMB) liquid substrate system to develop OD. Plates were blocked

with ChronBlock ELISA buffer per manufacturer's instructions, washing solution consisted of

PBS+0.1% Tween 20 (Sigma), and plates were read on a plate reader at 450 nM. Test sera were

run in triplicate and utilized at "*test OD*".

For the canine RBD ELISA, the negative controls consisted of sera from six pre-COVID

dogs, archived at WADDL and tested for canine Adenovirus (CAV), canine Distemper Virus

(CDV), canine Coronavirus (CCV), canine Parainfluenza (CPI), and canine Parvovirus (CPV)

IgG. All six samples had antibody presence of one or more of the tests performed, however no

sera reacted in the SARS-CoV-2 canine RBD ELISA. For the cat RBD ELISA, the negative

controls consisted of sera from three pre-COVID cats from WADDL archives, tested for feline

Coronavirus (FIP-FeCV) and feline Panleukopenia Virus (FPV)- IgG. Two of the three samples

had antibody presence of one or more of the tests performed (including 2 for FIP-FeCV);

however, neither reacted in the SARS-CoV-2 feline RBD ELISA. Negative controls were run in

triplicate and the mean was utilized as "*negative control OD*." A ratio of *test OD: negative*

control OD was used to determine the results. The positive cutoff of 2.0 *test OD: negative*

control OD ratio equated to the mean of negative controls + 3 standard deviations of the mean.

SARS-CoV-2 RBD ELISA was repeated three times for all samples, and the final results

were tabulated as a mean value obtained from the repeated testing. As no dog or cat in

Washington or Idaho had been confirmed to be SARS CoV-2 positive via serology prior to our

study, the first antibody positive case for each species and state was sent to the NVSL for

confirmation via virus neutralization (VN) assay in keeping with regulatory recommendations.

Both canine and feline SARS-CoV-2 RBD ELISA positive samples were confirmed at NVSL by

VN.

<u>Statistical analyses</u>

The primary aim of this study was to estimate the burden of household SARS-CoV-2

transmission from humans to their pets. Secondary aims included describing the nature of

human-animal contact within households and identifying risk factors for household transmission,

including human-animal contact. All analyses were conducted in R [25].

Outcome

Animal infection with SARS-CoV-2 was defined as an animal meeting one or more of

the following criteria: (1) SARS CoV-2 RBD ELISA seropositive status, (2) PCR positive status,

or (3) illness consistent with SARS-CoV-2 infection, hereafter referred to as "illness," defined as

participant answer of "yes" to the survey question: "Since the time of COVID

diagnosis/symptom onset in the household, has this animal had any new issues with difficulty

breathing, coughing and/or decreased interest in playing, walking, or eating?" Serostatus was

parameterized as ELISA ratio, log-transformed for the sake of interpretability; PCR positive

status and illness were parameterized as binary variables.

Regression models

Outcome was defined as an animal case of SARS-CoV-2 (definition above). Separate

regression models were fit for each outcome definition.

Household-level exposures for animal infection included residence in house versus

apartment or condominium (binary), home size in square feet (continuous), and the number of

human confirmed SARS-CoV-2 cases (continuous). Animal-level exposures for infection

included bedsharing with one or more human household members (binary), sharing bowls with

one or more household members (binary), and SARS-CoV-2 positive household members taking

precautions to prevent transmission to their pets following diagnosis, including not petting or

kissing the animal, staying in a different room, and having someone else feed and walk the

animal (binary). We also examined the association between canine seropositivity and illness

compatible with SARS-CoV-2 infection in the animal, and between seropositivity and time since

the animal was first exposed, defined as two days prior to the first date any household member

had symptoms of COVID-19 or tested positive, whichever was earlier.

We identified possible confounders *a priori* using a directed acyclic graph (DAG; Figure

1). The minimum sufficient adjustment set was defined, using this DAG and DAGitty.net,

separately for each exposure [26]. Animal species was explored as an effect modifier using a

multiplicative interaction term, and stratified results presented in all cases in which this

interaction term reached statistical significance ($p \leq 0.05$).

For each exposure of interest we implemented a generalized estimating equation (GEE)

approach with an exchangeable working correlation structure, household as the clustering

variable, and binomial models with a logit (binary outcomes) or Gaussian (continuous outcomes)

link, using the geepack package in R [27]. For regression of ELISA ratio on illness and time

since first exposure, we performed linear regression using the glm() function in R.

Results

<u>Recruitment</u>

In total, 107 eligible households enrolled and completed the survey. No households

currently living as unhoused enrolled. Two households corresponded to a single dog which was

moved from the participant's home to a family member's home immediately after the onset of

the participant's COVID-19 symptoms, leaving 105 households corresponding to 119 dogs and

57 cats available for analyses; no ferrets or hamsters enrolled or were sampled.

Sample collection is detailed in Figure 2. In total, 83 households corresponding to 100

dogs and 47 cats had a sampling visit conducted. Of these, six dogs and eight cats belonged to

households were not sampled due to temperament, leaving 94 dogs and 39 cats with PCR results,

while an additional 13 dogs and 9 cats were safe to restrain for swab (PCR) samples but not for

serum collection, leaving 81 dogs and 32 cats with serology results.

<u>Descriptive statistics</u>

Descriptive statistics are presented in Table 1. On average, at least six weeks (dogs) and

two weeks (cats) elapsed between the last human COVID-19 diagnosis in the household and

animal sampling. Of the 119 dogs and 57 cats with completed surveys, 20.4% (95% CI 12.9%,

29.7%) of dogs and 38.8% (95% CI 25.2%, 53.8%) of cats had reported illness. Of the 94 dogs

and 39 cats who were PCR tested, 5.3% (95% CI 1.8%, 12%) of dogs and 7.7% (95% CI 1.6%,

20.9%) of cats were positive; of the 81 dogs and 32 cats who had serum collected, 40.2% (95%

CI 29.6%, 51.7%) of dogs and 40.6% (95% CI 23.7%, 59.4%) of cats were seropositive.

Individual animal SARS-CoV-2 RBD ELISA results are shown in Figure 3 (dogs) and Figure 4

(cats). SARS-CoV-2 RBD ELISA test OD:negative control OD ratios in seropositive animals

ranged from 2.03 – 21.22 in dogs and from 3.01 – 30.35 in cats.

Five dog swabs (Cts 26.08 – 37.67) and 3 cats (Cts 27.03 – 39.97) were PCR positive on

nasal/oropharyngeal swabs; one of these dogs was also PCR positive from fecal swab (Ct 39.20).

Five PCR positive samples (2 cats and 3 dogs) had Cts sufficient for WGS (Ct<30): The earliest

cat sample (April 2021) that underwent WGS fell into Pango clade B.1.2. A later dog sample

sequenced as Delta sublineage B.1.617.2.103 (AY103), while the other three (2 cat, 1 dog)

samples sequenced as Delta sublineage B.1.617.2.25 (AY25). Of the five PCR positive dogs,

three were PCR positive prior to being seropositive and two were simultaneously PCR and

seropositive.

There were 11 households with two or more positive animals, and among multi-pet

households with at least one positive pet, mean prevalence (PCR or serology) was 91%. Out of

eight total PCR positive cases, all were detected after April 2021, when the first case of the Delta

variant was documented in Washington State.

Nearly one-third of dogs engaged in activities outside of the household during periods of

human isolation or quarantine. Over 50% of both cats and dogs resided in households whose

residents reported awareness of CDC guidelines to prevent human-animal transmission of

SARS-CoV-2, and 48 (41%) dogs and 17 (30%) cats resided in households which reported

taking precautions to prevent such transmission to household pet(s) following diagnosis. No cats

and only two dogs resided in a household in which an infected person was hospitalized for

COVID-19. Nearly all dogs (83%) and most cats (72%) had access to yards or gardens and were

allowed on furniture (86% of dogs and 100% of cats), and the majority were kissed by (75% of

dogs and 68% of cats) and shared beds (69% of dogs and 73% of cats) with human household

members. Almost all dogs' (91%) and cats' (95%) bowls were washed in the kitchen.

Regression models

Results of regression models are presented in Table 2 as prevalence odds ratios for the

binary outcome of illness, reflecting the cross-sectional design of this study, and as $\exp^{\beta}$ for the

outcome of ELISA ratio, which can be interpreted as the relative change (ratio scale) in ELISA

ratio for a one unit change in the exposure. As so few animals were PCR positive, we did not run

regression models for this outcome. With the exception of house size, which was adjusted for

house type as the minimum sufficient adjustment set was very small for this exposure,

confounders were not adjusted for due to concerns regarding overfitting arising from the small

sample size. Effect modification by species was found only for house type.

Dogs residing in houses on average had a 79% (95% CI 2%, 211%) higher ELISA ratio

than dogs residing in apartments or condos, while the inverse association was detected for cats

(49% lower mean ELISA ratio, 95% CI 75% lower, 3% higher) and for the outcome of illness in

both cats and dogs (48% lower prevalence odds, 95% CI 80% lower, 34% higher); this

association reached statistical significance for dogs only. No other effect estimates reached

statistical significance; however, there were positive trends across both outcome definitions for

bed sharing with humans, sharing bowls, and being indoor only; and a negative effect for

precautions taken to prevent SARS-CoV-2 transmission following diagnosis. We also found

ELISA ratio was positively associated with illness; however, we did not find evidence of an

272 effect of time since first exposure on ELISA ratio, nor of house square footage on either

273 outcome.

Discussion

275  We present the results of a cross-sectional, One Health study of SARS-CoV-2

276 transmission between people and their pets. The study results indicate that household

277 transmission of SARS-CoV-2 from humans to animals occurs frequently and infected animals

278 commonly display signs of illness. Notably, in 9 out of 11 households with multiple pets of

279 whom at least one tested positive (PCR or serology), all tested pets were positive. We

280 furthermore show that close human-animal contact is common among people and their pets in

281 this study population, that this contact appears to facilitate SARS-CoV-2 transmission, and that

282 pet owners are familiar with and willing to adopt measures to protect their pets from COVID-19.

283  There are several limitations to our approach. First, several weeks had elapsed from first

284 reported exposure to household sample collection from animals in most households, possibly

285 limiting our ability to detect viral shedding by PCR testing but strengthening our ability to detect

286 seroconversion. Second, while we assume transmission is from humans to pets, the cross-

287 sectional nature of this study precludes certainty regarding the direction of transmission.

288 Nevertheless, as SARS-CoV-2 is transmitted predominantly human-to-human, few cases of

289 SARS-CoV-2 have been documented in dogs and cats, and no cases have been documented to be

290 transmitted from dogs or cats to humans, we believe transmission in this study was exclusively

291 from humans to pets. Third, our study is subject to residual confounding due to inability to adjust

292 for confounders without risking over-fitting. We do not expect unmeasured or unadjusted

293 confounders to exert strong effects other than latent (and therefore difficult to measure and

model) constructs, such as socioeconomic status, strength of the human-animal bond, and level

of concern about zoonotic disease transmission. Finally, our definition of illness in pets is

simple, derived from a single survey item, and vulnerable to misclassification if these clinical

signs are due to other etiologies. This survey was created early in the COVID-19 pandemic,

although illness in pets is still not well-characterized.

We believe respondents misunderstood the question, "Is this animal indoor only vs.

indoor/outdoor" as 37% of dogs were reported to be indoor-only, however we believe this

variable retains its connection to degree of animal contact despite mismeasurement (i.e., a dog

labeled as "indoor only" likely spends more time in an indoor setting with humans than a dog

labeled as indoor/outdoor). We do not expect strong measurement error in any of the other

variables examined. As no gold-standard for canine anti-SARS-CoV-2 serology exists, validation

of our ELISA assay was limited to analytic validation and we could not reliably estimate

diagnostic sensitivity of our serological test; full diagnostic validation was not possible due to the

absence of sufficient gold-standard positive and negative samples, a limitation arising from the

status of SARS-CoV-2 as an emerging pathogen. However, all pre-COVID-19 samples evaluated

were negative, indicating specificity approaches 100%, and all samples sent to USDA-NVSL for

confirmatory PCR and serology testing had concordant results. While our primary aim—to

estimate the burden of human-animal SARS-CoV-2 transmission—was estimated with

reasonable precision, due to our small sample size variance was high for effect estimates

produced by our regression model. Finally, by nature of our recruitment methods and study

population, generalizability of our findings is likely limited to highly-educated, higher-income

individuals residing in urban and suburban communities.

Conclusions

These limitations aside, our study contributes important and novel findings to the

literature on cross-species transmission of SARS-CoV-2, with relevance to other zoonoses

transmitted in a household setting. Furthermore, we collected human, animal, and environmental

data, representing a true One Health approach to this critical research question. Finally, our

findings indicate households in this population are willing to adopt measures to protect their pets

from SARS-CoV-2 infection, and that these measures may be effective, indicating an

opportunity to prevent household transmission of zoonoses through health education and policy.

Acknowledgements

Data collection: Jessica Bell, DVM and Raelynn Farnsworth, DVM, Washington State

University College of Veterinary Medicine; Katherine Burr, DVM and Gemina Garland Lewis,

MS, Center for One Health Research, University of Washington.

Survey Review: J. Scott Weese, Ontario Veterinary College, University of Guelph.

Recombinant SARS-CoV-2 receptor binding domain source material: Dr. Wes Van Voorhis,

Center for Emerging and Re-emerging Infectious Diseases, University of Washington, Seattle,

WA, USA [28,29].

Funding:

The project was funded in part with funds from: 1) Department of Health and Human Services,

FDA Vet-LIRN Veterinary Diagnostic Laboratory Program grant #5U18FD006180-04, 2)

National Institute of Allergy and Infectious Diseases, National Institutes of Health, Department

of Health and Human Services, under Contract No.: HHSN272201700059C (WCV) and grant

Author Bio

Julianne Meisner is a post-doctoral fellow in Environmental and Occupational Health

Sciences at the University of Washington. Her research interests lie at the intersection of human

and animal health, and the application of rigorous epidemiologic methods to research at this

interface.

References

[1]   M. Konda, B. Dodda, V.M. Konala, S. Naramala, S. Adapa, Potential Zoonotic Origins of SARS-CoV-2 and Insights for Preventing Future Pandemics Through One Health Approach, Cureus. (2020). https://doi.org/10.7759/cureus.8932.

[2]   M. Kiros, H. Andualem, T. Kiros, W. Hailemichael, S. Getu, A. Geteneh, D. Alemu, W.E. Abegaz, COVID-19 pandemic: current knowledge about the role of pets and other animals in disease transmission. , Virol J. 17 (2020) 143.

[3]   A. Csiszar, F. Jakab, T. Valencak, Z. Lanszki, G. Tóth, G. Kemenesi, S. Tarantini, V. Fazekas-Pongor, Z. Ungvari, Companion animals likely do not spread COVID-19 but may get infected themselves, Geroscience. 42 (2020) 1229–1236.

[4]   T.S. Salajegheh, D.P. Magalhães, P. Rahimi, F. Shahabinejad, S. Dhakal, M.Y. Singh, A. Shehata, J. Lama, J. Klein, M. Safdar, M. Rahman, Filipiak KJ, A. Rodríguez-Morales, M. Sobur, F. Kabir, B. Vazir, L. Mboera, M. Caporale, M. Islam, J. Amuasi, R. Gharieb, P. Roncada, S. Musaad, B. Tilocca, M. Koohi, A. Taghipour, A. Sait, K. Subbaram, A. Jahandideh, P. Mortazavi, M. Abedini, D. Hokey, U. Hogan, M. Shaheen, A. Elaswad, M. Elhaig, M. Fawzy, Transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) to animals: an updated review, J Transl Med. 18 (2020) 358.

[5]   US Department of Agriculture, Animal and Plant Health Inspection Service, Https://Www.Aphis.Usda.Gov/Aphis/Dashboards/Tableau/Sars-Dashboard. (2022).

[6]   R.J. Molenaar, S. Vreman, R.W. Hakze-van der Honing, R. Zwart, J. de Rond, E. Weesendorp, L.A.M. Smit, M. Koopmans, R. Bouwstra, A. Stegeman, W.H.M. van der Poel, Clinical and Pathological Findings in SARS-CoV-2 Disease Outbreaks in Farmed Mink (Neovison vison), Veterinary Pathology. (2020). https://doi.org/10.1177/0300985820943535.

[7]   B.E.E. Martina, B.L. Haagmans, T. Kuiken, R.A.M. Fouchier, G.F. Rimmelzwaan, G. Van Amerongen, J.S.M. Peiris, W. Lim, A.D.M.E. Osterhaus, SARS virus infection of cats and ferrets, Nature. (2003). https://doi.org/10.1038/425915a.

[8]   American Veterinary Medical Association, SARS-CoV-2 in Animals, (n.d.). www.avma.org/resources-tools/animal-health-and-welfare/covid-19/sars-cov-2-animals-including-pets.

[9]   P. F, SARS-CoV-2 variants lacking ORF8 occurred in farmed mink and pangolin, Gene. 784 (2021) 145596. https://doi.org/10.1016/j.gene.2021.145596.

[10] S. Guo, K. Liu, J. Zheng, The Genetic Variant of SARS-CoV-2: Would it matter for Controlling the Devastating Pandemic?, International Journal of Biological Sciences. (2021). https://doi.org/10.7150/ijbs.59137.

[11] C. Wei, K. Shan, W. Wang, S. Zhang, Q. Huan, W. Qian, Evidence for a mouse origin of the SARS-CoV-2 Omicron variant, J Genet Genomics. 21 (2021) S1673-8527.

[12] de Guzman, Hong Kong Says Hamsters May Have Infected a Pet Shop Worker With COVID-19. Now They All Must Die, Https://Time.Com/6139876/Hong-Kong-Hamsters-Covid-19/. (2022). https://time.com/6139876/hong-kong-hamsters-covid-19/ (accessed January 23, 2022).

[13] M.F. Davis, S.C. Rankin, J.M. Schurer, S. Cole, L. Conti, P. Rabinowitz, G. Gray, L. Kahn, C. Machalaba, J. Mazet, M. Pappaioanou, J. Sargeant, A. Thompson, S. Weese, J. Zinnstag, Checklist for One Health Epidemiological Reporting of Evidence (COHERE), One Health. (2017). https://doi.org/10.1016/j.onehlt.2017.07.001.

[14] STROBE statement - Checklist of items that should be included in reports of
observational studies (© STROBE Initiative), International Journal of Public Health.
(2008). https://doi.org/10.1007/s00038-007-0239-9.
[15] M. Imai, K. Iwatsuki-Horimoto, M. Hatta, S. Loeber, P.J. Halfmann, N. Nakajima, T.
Watanabe, M. Ujie, K. Takahashi, M. Ito, S. Yamada, S. Fan, S. Chiba, M. Kuroda, L.
Guan, K. Takada, T. Armbrust, A. Balogh, Y. Furusawa, M. Okuda, H. Ueki, A.
Yasuhara, Y. Sakai-Tagawa, T.J.S. Lopes, M. Kiso, S. Yamayoshi, N. Kinoshita, N.
Ohmagari, S.I. Hattori, M. Takeda, H. Mitsuya, F. Krammer, T. Suzuki, Y. Kawaoka,
Syrian hamsters as a small animal model for SARS-CoV-2 infection and countermeasure
development, Proceedings of the National Academy of Sciences of the United States of
America. (2020). https://doi.org/10.1073/pnas.2009799117.
[16] J. Shi, Z. Wen, G. Zhong, H. Yang, C. Wang, B. Huang, R. Liu, X. He, L. Shuai, Z. Sun,
Y. Zhao, P. Liu, L. Liang, P. Cui, J. Wang, X. Zhang, Y. Guan, W. Tan, G. Wu, H. Chen,
Z. Bu, Z. Bu, Susceptibility of ferrets, cats, dogs, and other domesticated animals to
SARS-coronavirus 2, Science. (2020). https://doi.org/10.1126/science.abb7015.
[17] P.A. Harris, R. Taylor, B.L. Minor, V. Elliott, M. Fernandez, L. O'Neal, L. McLeod, G.
Delacqua, F. Delacqua, J. Kirby, S.N. Duda, The REDCap consortium: Building an
international community of software platform partners, Journal of Biomedical Informatics.
(2019). https://doi.org/10.1016/j.jbi.2019.103208.
[18] C. Eckstrand, T. Baldwin, K. Rood, M. Clayton, J. Lott, R. Wolking, D. Bradway, T.
Baszler, An outbreak of SARS-CoV-2 with high mortality in mink (Neovison vison) on
multiple Utah farms, PLoS Pathog. 17 (2021) e1009952.
[19] J. Plitnick, S. Griesemer, E. Lasek-Nesselquist, N. Singh, D. Lamson, K. St George,
Whole-Genome Sequencing of SARS-CoV-2: Assessment of the Ion Torrent AmpliSeq
Panel and Comparison with the Illumina MiSeq ARTIC Protocol, J Clin Microbiol. 59
(2021) e006492.
[20] GISAID, CoVsurver: Mutation Analysis of hCoV-19, Https://Www.Gisaid.Org/Epiflu-
Applications/Covsurver-Mutations-App. (2022).
[21] S. Khare, C. Gurry, L. Freitas, M. Schultz, G. Bach, A. Diallo, N. Akite, J. Ho, R. Lee, W.
Yeo, Curation Team GC, S. Maurer-Stroh, GISAID's Role in Pandemic Response, China
CDC Wkly. 3 (2021) 1049–1051.
[22] A. Rambaut, E. Holmes, Á. O'Toole, V. Hill, J. McCrone, C. Ruis, L. du Plessis, O.
Pybus, A dynamic nomenclature proposal for SARS-CoV-2 lineages to assist genomic
epidemiology., Nat Microbiol. 5 (2020) 1403–1407.
[23] Á. O'Toole, V. Hill, J. McCrone, E. Scher, A. Rambaut, Pangolin COVID-19 Lineage
Assigner, Https://Pangolin.Cog-Uk.Io/. (2022).
[24] I. Phan, S. Subramanian, D. Kim, M. Murphy, D. Pettie, L. Carter, I. Anishchenk, L.
Barrett, J. Craig, L. Tillery, R. Shek, W. Harrington, D. Koelle, A. Wald, D. Veesler, N.
King, J. Boonyaratanakornkit, N. Isoherranen, A. Greninger, K. Jerome, H. Chu, B.
Staker, L. Stewart, P. Myler, W. van Voorhis, In silico detection of SARS-CoV-2 specific
B-cell epitopes and validation in ELISA for serological diagnosis of COVID-19, Sci Rep.
11 (2021) 4290.
[25] R.F. for S. Computing., R Core Team 2020, R: A Language and Environment for
Statistical Computing. (2020).

[26]   J. Textor, B. van der Zander, M.S. Gilthorpe, M. Liśkiewicz, G.T. Ellison, Robust causal
435   inference using directed acyclic graphs: The R package "dagitty," International Journal of
436   Epidemiology. (2016). https://doi.org/10.1093/ije/dyw341.
437   [27]   S. Højsgaard, U. Halekoh, J. Yan, The R Package geepack for Generalized Estimating
438   Equations Journal of Statistical Software, CRAN. (2006).
439   [28]   A.C. Walls, Y.J. Park, M.A. Tortorici, A. Wall, A.T. McGuire, D. Veesler, Structure,
440   Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein, Cell. (2020).
441   https://doi.org/10.1016/j.cell.2020.02.058.
442   [29]   I.Q. Phan, S. Subramanian, D. Kim, M. Murphy, D. Pettie, L. Carter, I. Anishchenko, L.K.
443   Barrett, J. Craig, L. Tillery, R. Shek, W.E. Harrington, D.M. Koelle, A. Wald, D. Veesler,
444   N. King, J. Boonyaratanakornkit, N. Isoherranen, A.L. Greninger, K.R. Jerome, H. Chu,
445   B. Staker, L. Stewart, P.J. Myler, W.C. Van Voorhis, In silico detection of SARS-CoV-2
446   specific B-cell epitopes and validation in ELISA for serological diagnosis of COVID-19,
447   Scientific Reports. (2021). https://doi.org/10.1038/s41598-021-83730-y.
448
449

	n (%)	
	Dogs (N=119)	Cats (N=57)
Animal		
Illness consistent with SARS-CoV-2	20 (20%)	19 (39%)
Seropositive	33 (40%)	13 (41%)
PCR positive	5 (5%)	3 (8%)
ELISA ratio	3.9 (4.93)*	9.88 (12.51)*
Activity[a] during human quarantine	33 (28%)	7 (12%)
Respondent took precautions[b]	48 (41%)	17 (30%)
Age	6.05 (3.86)*	6.40 (4.50)*
Male	66 (56%)	28 (49%)
Respondent aware of CDC guidelines[c]	62 (53%)	29 (53%)
Time from first diagnosis[d] to sampling (days)	51.17 (60.64)*	29.28 (19.17)*
Time from last diagnosis[d] to sampling (days)	43.06 (69.44)*	15.16 (40.93)*
Humans		
Index case age	41.78 (13.24)*	47.91 (14.38)*
Index case male	34 (29%)	14 (25%)
Index case preexisting condition[e]	27 (23%)	18 (32%)
Index case was hospitalized	2 (2%)	0 (0%)
No. SARS-CoV-2 positive household members	1.78 (1.28)*	1.72 (1.13)*

No. household members with COVID-19-like symptoms[f]	0.27 (0.63)*	0.26 (0.55)*
No. household residents	3.43 (1.49)*	3.07 (1.28)*

Environment		
Reside in a house	91 (76%)	51 (89%)
Reside in an apartment or condominium	51 (24%)	6 (11%)
Square footage of housing	1856.32 (932.74)*	1980.88 (1095.15)*
Number of bedrooms	3.24 (1.4)*	3.19 (1.22)*
Number of floors	1.87 (0.82)*	1.84 (0.62)*
Access to outdoor space where pets can roam	99 (83%)	41 (72%)

Human-animal contact		
Bowls used by animals cleaned in the kitchen	108 (91%)	54 (95%)
Humans and animals share bowls	15 (13%)	8 (14%)
Humans wash hands before handling animals	15 (13%)	2 (4%)
Humans wash hands after handling animals	50 (42%)	12 (21%)
Animal bedshares with humans	81 (69%)	41 (73%)
Animal shares a bedroom but not a bed with humans	54 (46%)	19 (34%)
Animal is indoor-only	43 (37%)	35 (61%)
Animal sleeps outdoors	1 (1%)	5 (9%)
Humans pet the animal	117 (100%)	56 (100%)
Humans kiss the animal	88 (75%)	38 (68%)
Animal is allowed on furniture	101 (86%)	56 (100%)

Table 1: Descriptive statistics for 119 dogs and 57 cats corresponding to 105 households.

*mean (standard deviation). [a]Activity defined as going to a veterinary clinic or groomer, being walked off-leash, or visiting an off-leash park, dog park, kennel, or daycare facility. [b]Precautions to prevent human-animal SARS-CoV-2 transmission following diagnosis: not petting or kissing the animal, staying in a different room, and having someone else feed and walk the animal. [c]Guidelines to prevent human-animal SARS-CoV-2 transmission. [d]First diagnosis: earliest known, confirmed SARS-CoV-2 diagnosis in the household; final diagnosis: last known, confirmed SARS-CoV-2 diagnosis in the household. [e]Prexisting conditions: diabetes, kidney disease, heart disease, hypertension, immunosuppression. [f]Household members who had COVID-19-like symptoms but did not get tested.

	Illness consistent with SARS-CoV-2 [a]	ELISA ratio[a]
Exposure	POR (95% CI)	exp$^\beta$ (95% CI)
Indoor-only	1.63 (0.77, 3.45)	1.07 (0.61, 1.88)
House type[b]	0.52 (0.2, 1.34)	1.79 (1.02, 3.11) (dogs) 0.51 (0.25, 1.03) (cats)
House square footage	1 (1, 1)	1 (1, 1)
Share bowls[c]	1.29 (0.39, 4.25)	1.78 (1.07, 4.49)
Bedsharing	1.48 (0.66, 3.33)	1.16 (0.68, 1.95)
Took precautions[d]	0.71 (0.29, 1.75)	0.81 (0.48, 1.37)
No. SARS-CoV-2 infected humans	0.78 (0.54, 1.13)	1.18 (0.85, 1.64)
Illness consistent with SARS-CoV-2	-	1.09 (0.59, 2.01)
Time since first exposure (days)[e]	-	1 (1, 1)

Table 2: Regression model results. House size was adjusted for house type, but no other

models were not adjusted for confounders due to overfitting concerns. [a]Survey results available

for 119 dogs and 57 cats, serology results available for 81 dogs and 32 cats. [b]House versus

apartment or condominium. [c]Animals and humans share bowls. [d]Precautions taken to prevent

human-animal SARS-CoV-2 transmission following diagnosis: not petting or kissing the animal,

staying in a different room, and having someone else feed and walk the animal. [e]First exposure

defined as 2 days prior to first positive diagnosis in the household or onset of symptoms,

whichever was earlier. POR: prevalence odds ratio; 95% CI: 95% confidence interval.

Figure 1: Directed acyclic graph for human-animal SARS CoV2 transmission. Variables

outlined with a square are the exposures of interest, while outcome (approximated by serostatus,

PCR result, and illness in separate models) is outlined with a circle. HAB: human-animal bond;

SES: socioeconomic status; took precautions: SARS-CoV-2 positive household member(s) took

precautions to prevent transmission to pet; indoor-only: animal does not go outdoors; bedshare:

animal shares a bed with one or more household members.

Figure 2: Flowchart depicting serological and PCR sampling. Out of 119 dogs and 57 cats

corresponding to 105 households with completed surveys, PCR testing is complete for 94 dogs

and 39 cats, and serological testing is complete for 81 dogs and 32 cats. The remaining pets were

not sampled due to safety concerns.

Figure 3: SARS-CoV-2 RBD ELISA Serology data, cats. PCR testing is complete for 39 cats,

and serological testing is complete for 32 cats. The remaining pets were not sampled due to

safety concerns.

Figure 4: SARS-CoV-2 RBD ELISA Serology data, dogs. PCR testing is complete for 94

dogs, and serological testing is complete for 81 dogs. The remaining pets were not sampled due

to safety concerns.