Hot and cold spots of pest-induced US urban tree death, 2020-2050.

Authors

Emma J. Hudgins[1*], Frank H. Koch[2], Mark J. Ambrose[3], Brian Leung[1,4]

Affiliations

[1] Dept. of Biology, McGill University.

[2] USDA Forest Service, Southern Research Station.

[3] Dept. of Forestry and Environmental Resources, North Carolina State University.

[4] School of Environment, McGill University.

*Corresponding author, Stewart Biology Building, 1205 avenue Docteur Penfield, Montreal, Quebec, Canada, H3A 1B1, tel: +15142452054, emma.hudgins@mail.mcgill.ca

Short title: US urban pest impacts, 2020-2050

Keywords: Forest ecology, Economic impact, Scenario modelling, Invasive species, Invasive insects

Article Type: Letter

Words in abstract: 149

Words in main text: 4939

Words in text box: 431

Number of references: 34

Number of figures: 4

Number of tables: 1
Number of text boxes: 1

Statement of authorship: EJH and BL conceptualized the study. EJH performed the analyses and created the figures. EJH, BL, FHK and MJA wrote the paper. FHK and MJA provided insect and urban tree expertise and commented on realism of results. MJA manages the urban tree database.

Data accessibility statement: All data, code, and materials used in the analyses (except where privacy restrictions apply) are available at www.github.com/emmajhudgins/UStreedamage. A DOI will be provided at the time of acceptance via Zenodo.

Abstract

Urban trees are important nature-based solutions for future wellbeing and livability but are at

high risk of mortality from insect pests. Until now, the magnitudes and spatial distributions of

risks were unknown. Here, we combine new street tree models in ~30,000 United States (US)

communities with models of invasive insect spread and host tree death. We estimate that an

additional 1.4 million street trees will be killed from 2020 through 2050. Roughly 23% of

communities will experience 95% of all mortality, and 90% will be due to the emerald ash borer

(*Agrilus planipennis*), which is expected to kill virtually all ash trees (*Fraxinus* spp.) in >6000

communities. Mortality hotspots include Milwaukee, WI and Chicago, IL, while damage is lower

across the southern and central US. Asian wood borers of maple and oak trees pose the highest

future risk, where a new establishment could cost US$ 4.9B over thirty years.

Introduction

Previous analyses suggest that impacts associated with urban trees are likely to comprise the dominant share of economic damages caused by invasive alien forest insects (IAFIs) in the United States (US) (Aukema et al. 2011). Urban tree populations include highly susceptible species like ash (*Fraxinus* spp.) that are being decimated by emerald ash borer (EAB, *Agrilus planipennis*) (Kovacs et al. 2010). To eliminate the potential for injury or property damage due to dead trees, infested urban trees must be treated or removed (Fahrner et al. 2017). Moreover, the importance of urban forests is only expected to grow. While urbanization is already very high in the US (82% in 2018), it has not yet reached saturation (World Bank, http://data.worldbank.org, UN DESA, http://population.un.org). At the same time, there has been a push for urban 'greening' (i.e., increasing urban forest canopy). Urban trees perform many important ecosystem services, including lowering cooling costs (Norton et al. 2015), buffering against flooding, increasing air quality, carbon sequestration, improving citizens' mental and physical health outcomes, and creating important habitat (Van den Berg et al. 2010; Roy et al. 2012). The high tree mortality risk posed by IAFIs can greatly diminish these myriad benefits.

While IAFI life histories differ, they are known to be transported long distances by humans (Hulme 2009), potentially with similar drivers across entire dispersal pathways (Hudgins et al. 2017, 2020). Thus, the creation of a pathway-level damage estimate can provide insight into the benefit of limiting future spread via these pathways (e.g. through quarantines, highway checkpoints to limit firewood movement). Past estimates of IAFI damage have been important in providing support for phytosanitary measures such as ISPM15 (IPPC 2002), a wood packing material treatment protocol, whose adoption is growing worldwide (Leung et al. 2014). A

previous pathway-level estimate for the cumulative cost of all US IAFIs was performed a decade

ago and had substantial data limitations (Aukema et al. 2011). Since then, contemporary

advances allow direct estimates of spread for every IAFI species as well as host prevalence and

IAFI-induced mortality for every tree species in every community across the United States. This

allows not only the estimation of country-wide IAFI damages, but also IAFI and host-specific

damages and their spatial distribution. Further, we can examine the impact of tree mortality

dynamics on cost dynamics and derive better risk assessments of not-yet established pests, based

their functional traits and host distributions.

In this paper, we synthesized four subcomponent models of IAFI invasions: 1) a model of

spread for 57 IAFI species, 2) a model for the distribution of all urban street tree host genera

across all US communities, 3) a model of host mortality in response to IAFI-specific infestation

for all urban host tree species, and 4) the cost of removing and replacing dead trees, to provide

the best current estimate of the damage to street trees, including explicit estimates for all known

IAFIs across all major insect guilds.

Materials and Methods

We synthesized four subcomponent models of IAFI invasions (see conceptual diagram, Fig. S1).

IAFI dispersal forecasts

We modelled spread using the Semi-Generalized Dispersal Kernel (SDK, Hudgins et al. 2020).

This is a spatially explicit, negative exponential dispersal kernel model that can account for

additional spatial predictors in source and recipient sites. The SDK builds from the Generalized

Dispersal Kernel (GDK, Hudgins et al. 2017) as a starting point, using human population

density, forested land area and tree density in source and destination sites as moderators of

spread. The SDK combines up to three species-specific corrections for each species to maximize

predictive ability: 1) a species-specific intercept term, 2) information on an IAFI's likely initial

invasion location, and 3) niche-related limitations when evidenced in the literature. The SDK

was applied to all 57 IAFIs believed to cause some damage from Aukema et al. (2011) and

projected from 2020 to 2050 (Fig. S2).

Street tree models

Our fitting set consisted of 653 street tree databases for US communities where street tree

inventory data had been collected (Fig. S3, Koch et al. 2018). In two communities (Tinley Park

and, IL and Fort Wayne, IN), preventative cutting for EAB was conducted prior to the most

recent inventory and was therefore accounted for within our dataset. We modelled the abundance

and diameter at breast height (DBH) for trees within each genus, as treatment costs are

dependent on number and diameter of trees (Aukema et al. 2011). We split trees into three

diameter classes (small = 0-30cm, medium = 31-60cm, large >60cm). We first fit models for the

total tree abundance of all species by diameter class, and then used these total tree models to help

predict genus-specific tree abundance within each diameter class. Street tree inventory data are

not always reliably reported to the species level across municipalities, and some species are so

rare in street tree inventories that it would have been very difficult to develop robust species-

level models, so we limited our examination to the genus level. Since IAFIs may not be equally

impactful to all host tree species in a genus, we had to estimate the genus-level severity of each

IAFI species for each IAFI-host combination. We did so by estimating the species-level

breakdown of each genus based on their average relative proportions across our 653 inventoried

communities, and assuming this distribution was representative in other projected communities.

We modelled the total abundance of street trees in a community using boosted regression

trees (BRT, *gbm.step* within R package *dismo,* Hijmans et al. 2017, Appendix S1) relating the

logarithmically-scaled total tree abundance within a diameter class to community-specific

predictors, employing environmental variables from WORLDCLIM (Fick & Hijmans 2017) and

community characteristics used in Koch et al. (2018), and sourced largely from the National

Land Cover Database (NLCD, Homer et al. 2015), the US Census and the American Community

Survey (https://www.census.gov/data.html, Table S1).

Next, we estimated the abundance of street trees within each genus, using the same

climatic and demographic factors as the total tree abundance model as well as the total tree

abundance model output as predictors (Fig. S1). We considered two approaches: 1) zero-inflated

Poisson generalized additive models (GAMs), or 2) a two-step BRT approach. We compared

BRT and GAM models that were fit to all genera simultaneously (general BRT/GAM models

using genus-specific intercept terms) with models that were fit to each genus separately

(customized BRT/GAM models) (Fig. S1). We chose the model that produced the strongest

relationship for each genus using R^2 values that were relative to the 1:1 line (i.e, a normalized

mean squared error, R^2_{MSE}, Appendix S1).

We synthesized the previous two modelling steps, intersecting IAFI spread forecasts with

predicted tree distributions (using observed tree data where available), to create forecasts of tree

exposure, which we define as the sum of predicted density of each IAFI species, multiplied by

their predicted host tree abundance in each community.

Host mortality model

We examined the impacts of the three major feeding guilds of IAFIs (34): Foliage feeders

included insects that feed on leaf or needle tissue. Sap feeders included all species that consume

sap, including scale insects and gall-forming species. Borers included species that feed on

phloem, cambium, or xylem. Across insect guilds, the logic from Aukema et al. (2011) appeared

to hold: most species were innocuous, but a small number caused high mortality (Table S7). In

contrast, while several invasive pathogens were mentioned in Potter et al. (2019), pathogens are

only reliably reported when they produce noticeable impacts (Aukema et al. 2011). To avoid

mischaracterizing their impacts, we removed pathogens from the remainder of our analysis.

We ranked the severity of a given IAFI infestation on a particular host using a scale based

on observed long-term percent mortality (Table S7, defined in Potter et al. 2019). We fit a Beta

distribution to the frequency distribution of IAFIs in each of these categories using Stan

(Carpenter et al. 2017), a program and language for efficient Bayesian estimation. We used the

posterior mean as the expected mortality for an IAFI in each severity category, rather than the

simple midpoint of the range of each category.

We define the term 'mortality debt' as the period between an IAFI initiating damage

within a community and reaching its estimated asymptotic host mortality within that community

(Text Box 1). While we had estimates of the asymptotic proportional mortality of host trees

(Potter et al. 2019), we had no information on the rate by which trees reach this plateau. Previous

estimates have ranged from 5 to 100 years (Aukema et al. 2011, Pugh 2010), so we analyzed

three scenarios within this range (10, 50, 100 years). To account for what is currently known

about the mortality dynamics of IAFIs within each of the feeding guilds, we also examined

scenarios based on our most likely scenario of the duration of mortality debt across IAFI feeding

guilds. For simplicity, we assumed mortality increased by a constant fraction over time until

reaching its maximum and levelling off. For example, in the 50-year mortality debt scenario, if

an IAFI's maximum host mortality was defined as 90%, mortality would increase by 9% at each

5-year timestep for 10 timesteps until 90% mortality had been reached.

<u>Management costs</u>

As a final layer that allowed us to move from mortality estimates to cost estimates, we estimated

the cost of removing and replacing dead trees. We used this cost because we believe it to be the

minimum management response required, and because the extent and variability of preventive

behaviour would be much harder to estimate. However, we note that this cost does not account

for additional preventive cutting or any non-cutting management such as spraying or soil

drenching with pesticides. We assumed that cutting was a one-time 100% effective treatment

against IAFIs, or in other words, that newly planted trees were of different species and thus not

susceptible to the same IAFI species that killed the previous trees. We assumed a 2% discount

rate for future damages (Aukema et al. 2011) and that infestations were independent, or in other

words that invasion by one IAFI did not interfere with invasion by another. This is likely a fair

assumption, as there is minimal host sharing across IAFIs, and IAFI species each infest only a

small proportion of hosts at a given time interval, so there is minimal potential for species

interactions (Aukema et al. 2010). We assumed the same per-tree cost estimates for cutting and

replacing dead trees as in Aukema et al. (2011), where the cost of cutting increases nonlinearly

with size class. If we assume that street trees are always under the jurisdiction of local

governments, the cost of removal and replacement of each tree is US$450 for small trees,

US$600 for medium trees, and US$1200 for large trees (these costs jump to an estimated

US$600, US$800, and US$1500 for homeowners). We reported all costs incurred from 2020 to

2050 in 2019 US dollars based on a 2% discount rate relative to these baseline costs. Since these

baseline per-tree management costs came from a 2011 publication, we converted them to 2019

dollars via the consumer price index, which amounted to an inflation of 13.65% (World Bank,

https://data.worldbank.org), though we note that the present-day costs of per-tree removal may

have declined with advances in technology.

<u>Model synthesis</u>

Once all subcomponent models had been parameterized, we synthesized the street tree estimates,

IAFI spread estimates, host mortality estimates, and removal costs to produce overall cost

estimates (Fig. S1). We summed the damages from 2020 to 2050 to obtain a total discounted cost

for this 30-year window. We then obtained annualized costs by calculating an annuity over the

30-year time horizon using the following equation:

203
$$Annualized\ damage = D\frac{\sum_{time=min}^{max} Costs_{time}}{(1-(1+D)^{min-max})}$$
(1)

Where D is the discount rate (2%).

We assessed parameter uncertainty in proportional host mortality by sampling from our

posterior beta mortality distribution. We also used sensitivity analysis to explore the effect of

different mortality debt scenarios, including 1) our most likely scenario, 2) setting all guilds to

10, 50, or 100-year debts, and 3) varying each guild separately while holding the other two

guilds at their most likely scenario. While our host distribution models were based on standard

modelling approaches (e.g., GAM), our Bayesian formulations underlying the mortality

estimates were novel and needed to be tested theoretically, to ensure that parameters were

identifiable, and reproduced the correct behavior. See Appendix S2 for details of our theoretic

analyses.

<u>Potential impacts to non-street trees</u>

To provide a rough estimate of non-street tree impacts, we built a model for whole-community

trees (i.e., street + non-street trees) from the dataset of 56 communities where genus-level

estimates were reported, subtracted predicted street trees from this whole community estimate,

and apportioned the remaining trees into residential and non-residential trees based on their

average fractions across all sites where land type breakdowns were provided (32 municipalities).

Results

<u>Urban tree pest exposure</u>

Total tree abundance models were predictive with some outliers (Appendix S1, Fig. S4, small

trees: $R^2 = 0.78$, medium trees: $R^2 = 0.58$ large trees: $R^2 = 0.42$). Removing the outliers changed

the R^2 to 0.76 for small trees, 0.76 for medium trees, and 0.58 for large trees. Our genus-level

abundance models were strong but became slightly weaker for rare genus - size class

combinations (Fig. 1, overall R^2 for all genera of small trees: $R^2 = 0.93$, medium trees: $R^2 = 0.93$,

large trees: $R^2 = 0.92$). While relationships were variable across genera, the genera that were fit

most poorly did not make up a large proportion of predicted trees, and none were below $R^2 =$

0.25 (Fig. S5).

The optimal genus-level fitting approach differed across genera depending on diameter

class, prevalence of genera, and whether presence/absence or tree abundance was the response

variable (Table S2). Generally, rarer genera were better fit by global BRT and GAM models,

which utilized information from all other species while common species were better fit by

customized models (Fig. S6). According to our models, while subject to regional variation, the

population of street trees is mostly made up of maple (*Acer*) and oak *(Quercus)*, with substantial

ash (*Fraxinus,* Fig. S7).

Predicted street tree exposure (measured as the number of predicted susceptible trees in

Fig. 2a * IAFI relative propagule pressure in Fig. 2b, Hudgins et al. 2020) across all tree types

from 2020 to 2050 was generally high in the eastern US, and only sporadically high across the

western US (Fig. 2c). Predicted street tree exposure was highest among maples (*Acer* spp.*,*

25.6M predicted exposed trees)*,* oaks (*Quercus* spp.*,* 5.9M)*,* and pines (*Pinus* spp. 3.4M). The

greatest number of trees were predicted to become exposed to San Jose scale (*Quadraspidiotus*

perniciosus, 7.3M), Japanese beetle (*Popillia japonica*, 6.7M), and calico scale (*Eulecanium*

cerasorum, 6.4M). Among residential and community trees, exposure was greatest among

maples, oaks, and *Prunus* spp. (1.7B,1.1B, 707M, respectively), and the most frequently

predicted IAFI encounters were with the same three species (Japanese beetle, San Jose scale, and

calico scale).

Host tree mortality

The best-fitting mortality model indicated that most IAFIs fall in the low severity groups. Within

all severity groups, the majority of IAFIs were at the low end of severity (Fig. 3, full results in

Appendix S2). In our most likely mortality debt scenario (i.e., 10-year scenario for borers, 50-

year scenario for defoliators, 100-year scenario for sap feeders, Text Box 1), we estimated a

mortality level of 0.7-2.5% beyond expected natural mortality of street trees by 2050, where our

most likely scenario fell on the higher end of this range (Table 1). Predicted street tree death

varied by a factor of four based on the mortality debt scenario, with longer debts leading to lower

total mortality between now and 2050 (Table 1). This was because in longer mortality debt

scenarios, trees experience mortality in the years 2020-2050 from IAFIs that initially established

in their communities in 2000 (50yr debt) or 1950 (100yr), but our highest impact IAFI (EAB)

can only begin to cause mortality after 2002 in any scenario. Sensitivity was driven largely by

wood boring species, as demonstrated by the sensitivity of mortality estimates to their mortality

debt scenarios ("Vary Borers" row, Table 1). We also found that longer mortality debts led to a

smoother cost curve, or costs that do not vary much due to more consistent host mortality rates

(Fig. 4).

Spatially, future damages will be primarily borne in the Northeast and Midwest, driven

by EAB spread (Fig. 2d). We predict that EAB will reach asymptotic mortality in 6747 new

cities, which means that 98.98% of its preferred ash hosts will die. Thus, the mortality is

predicted to be concentrated in a 902,500km^2 zone encompassing many major Midwestern and

Northeastern cities (Fig. S10). This mortality is also predicted to result in a 98.8% loss of all ash

street trees within this zone. Over 230,000 ash street trees are predicted to have died before 2020,

and there are a further 69 cities where EAB is predicted to reach asymptotic mortality within 10

years of 2050 (i.e., 98.8% ash mortality by 2060). Due to the restricted range of forest ash

relative to urban ash, we predict that 68% of ash trees and 76% of communities containing street

ash will remain unexposed to EAB in 2060. Furthermore, at-risk ash trees are unequally

distributed. We projected the highest risk close to the leading edge of present-day EAB

distributions, particularly in areas predicted to have high ash densities. The top "mortality

hotspot cities", where projected additional mortality is in the range of 5,000-25,000 street trees,

include Milwaukee, WI, the Chicago Area (Chicago/Aurora/Naperville/Arlington Heights, IL),

Cleveland, OH, and Indianapolis, IN (Fig. 2d). Cities predicted to have high mortality outside of

the Midwest include New York, NY, Philadelphia, PA, and Seattle, WA – communities with

high numbers of street trees and high human population densities, which attract EAB propagules

within our spread model. The states most impacted by street tree mortality match these patterns,

where the highest mortality is predicted for Illinois, New York, and Wisconsin.

Cost estimates

We estimated annualized street tree costs across all guilds to be between US$29-33M per year in

our most likely scenario (mean = $30M, Table 1, Fig. S11). Roughly 90% of all costs across the

entire US were due to EAB-induced ash mortality. The total cost associated with street tree

mortality in the top ten hotspot cities was estimated at $50M from 2020 to 2050, with $13M in

Milwaukee, WI alone.

The ranking of feeding guild severity was relatively robust across mortality debt

scenarios, despite the potential for differences due to the interaction of IAFI-specific spread and

mortality debt dynamics. Costs were higher for longer mortality debt scenarios for borers,

peaked at intermediate debt for defoliators, and peaked at the longest debt for sap feeders. These

patterns were due to the relative rates of historical and contemporary range expansion of more

impactful IAFIs (i.e., high impact borers have more rapid recent range expansion, while

contemporary high impact defoliator expansion is slow compared to 50 years ago). Borers were

predicted to be the most damaging feeding guild ($8M-28M mean annualized street tree damages

across scenarios), and EAB was consistently the top threat. Defoliators were predicted to be the

second most damaging feeding guild in the next 30 years (means = $0.8M-$1.4M), despite

having more widespread hosts than wood borers, due to lower asymptotic mortality levels.

Defoliators had a 1-2 order of magnitude lower cost than wood-boring species, but again showed

consistency in which species were the top threats within the guild. Consistent with previous work

in Aukema et al. (2011), European gypsy moth had the highest cost of all defoliators, followed

by Japanese beetle and cherry bark tortrix (*Enarmonia formosana*). The sap-feeding group

accrued the lowest costs in the next 30 years due to their lower asymptotic mortality and rarer

street tree hosts (mean = $0.2M-1.1M). Hemlock woolly adelgid (*Adelges tsugae*) was the

highest impact sap feeder, followed by oystershell scale (*Lepidosaphes ulmi*) and elongate

hemlock scale (*Fiorinia externa*). Total costs were only notably sensitive to borer mortality debt

scenario misspecification (Table 1), which is promising, given our certainty of the shorter

scenario for EAB.

<u>Potential impacts to non-street trees</u>

Mean added mortality (i.e., above background rates) for residential and non-residential

community trees in the most likely scenario was 1.0% (13.3M residential and 72.1M non-

residential trees, Table S10). While recognizing that non-street tree management will likely be

more variable, to provide a rough estimate, we assumed that non-street trees would be managed

in the same way as street trees (i.e., removal and replacement of dead trees). In this scenario,

added mortality would incur an estimated annualized cost of $1.5B for non-residential trees and

$356M for residential trees. Further, a disproportionate amount of the total damages (91% of the

mortality to residential non-residential community trees) is expected to be felt in the hotspot

zone, with 12.1 million residential and 65.9 million non-residential community trees expected to

be killed. Given the relatively limited data, and the difference in potential management

behaviour for these trees, we caution against overinterpretation of these results.

<u>Novel IAFI risk forecast</u>

Our framework allowed us to identify the factors leading to the greatest impacts for IAFIs

already known to have established in the United States. We were able to identify the most

common urban host trees, the sites facing the greatest future IAFI propagule pressure, and the

IAFI-host combinations with the greatest mortality. However, this approach can also be

synthesized with IAFI entry scenarios to understand potential impacts of novel invasive IAFIs.

To illustrate the utility of this framework for prediction, we have provided a checklist of risk

factors in Table S11 and future spread simulations in Table S12 and Fig. S12. We show that

entry via a southern port (e.g., the Port of South Louisiana) would lead to the greatest number of

exposed trees. Further, an EAB-like borer of oak and maple trees could kill 6.1 million street

trees and cost $4.9B over the next 30 years.

Discussion

While previous analyses have indicated that urban trees are associated with the largest share of

economic damages due to IAFIs (Aukema et al. 2011, Paap et al. 2017, Kovacs et al. 2010), until

recently, data did not exist on the urban distribution of host trees (Koch et al. 2018), the spread

of IAFIs (Hudgins et al. 2017,2020), nor the mortality risk for hosts due to different IAFIs

(Potter et al. 2019). With these new models, it is now possible to forecast where and when IAFIs

will have the most damages across the US. Our analysis suggests an overall added mortality of

between 2.1-2.5% of all street trees, amounting to $US 30M per year in management costs.

However, the most interesting and potentially useful element was our ability to forecast hotspots

of future forest IAFI damages, including a 902,500km^2 region that we expect to experience

95.7% of all mortality, in large part due to a 98.8% loss of its ash street trees due to EAB. This

type of forecasting has been highlighted as a crucial step in prioritizing management funds

(McGeoch et al. 2016). These data can be used by municipal pest managers to anticipate future

costs, and may help motivate improved spread control programs that aim to identify the potential

source counties of future invasions and mitigate the worst anticipated impacts (complete forecast

available at http://github.com/emmajhudgins/UStreedamage).

Beyond present IAFI risks, our integrated model can also act as a risk assessment tool for

street tree mortality caused by novel IAFIs (Table S11-S12, Fig. S12). While ash trees are

assured to be dramatically affected by EAB over the next few decades, our models suggest oak

and maple to be the most common street tree genera nationwide. Further, while ash species are

being substituted with less susceptible tree species, maples and oaks continue to be widely

planted within our street tree inventories. Therefore, IAFIs with host species spanning these

genera should be of heightened concern. Secondly, the timescale and magnitude of the impacts

of wood borers (see also Aukema et al. 2011) make them the highest risk to street trees. We

integrated these two pieces of information with information on major ports of entry within the

US (American Association of Port Authorities 2015, http://aapa.com/), as well as our general

model of IAFI spread (Hudgins et al. 2020), to forecast the extent of exposed maple and oak

street trees from 2020-2050 (Fig. S12, Table S12). Our analyses show that entry via a southern

port would lead to the greatest number of exposed trees. Further, larger trade volumes between

the US and Asia compared to other regions (Sardain et al. 2019) suggest Asian natives will be

the most likely future established IAFIs. One potential candidate species fitting these criteria is

citrus longhorned beetle, which is an Asian wood borer thought to have many potential host

species within the United States, including ash, maple and oak (Haack et al. 2010). The lack of

more thorough regulation of live plant imports and strict implementation of current wood

treatment protocols such as ISPM15 (Lovett et al. 2016) increase the susceptibility of the US to

invasion and subsequent spread of this species and other potentially high-risk borers.

382 Our impact estimates vary substantially based on dynamics of host mortality following

383 initial IAFI invasion, especially because of variability in the duration and functional form of

384 mortality debt. Luckily, the guild (borers) and species (EAB) whose impact on total community

385 costs are most sensitive to correct specification of the mortality debt dynamics are the ones for

386 which we are most confident. Several publications have demonstrated near-complete decimation

387 of ash stands in the decade following EAB infestation (Kovacs et al. 2010, Knight et al. 2013,

388 Fei et al. 2019). Furthermore, since total tree mortality is asymptotically equivalent across all

389 mortality debt regimes, if other feeding guilds possessed 10-year mortality debt regimes, we

390 should have been able to detect a rapid die-off of their hosts as they spread, similarly to what we

391 found for EAB (albeit scaled by their maximum mortality rates). This is not the case in the

392 literature (Fei et al. 2019).

393 With our integrated model, we also estimated economic damages, which updates the

394 decade old Aukema et al. (2011) using recent advances (Koch et al. 2018, Potter et al 2019).

395 Surprisingly, the previous cost estimates were not that different at the country scale. The

396 previous cost estimate separated urban trees into residential and non-residential types, grouping

397 street trees in the latter. We estimate annualized costs for non-residential trees to be somewhat

398 lower than those in Aukema et al. (2011, $1.3B versus $2.0B in total "Local Government

399 expenditures"). This lower estimate is likely because of a lower rate of predicted ash exposure to

400 EAB (i.e., lower predicted ash abundance in areas of predicted EAB spread) in non-residential

401 areas. Interestingly, our estimate of residential tree costs is roughly one third that in Aukema et

402 al. (2011, $303M vs $1.1B in total "Household Expenditures"), again likely due to a (more

403 extreme) overestimate in the nationwide prevalence of residential ash trees in the previous

404 publication.

We predict that 75% of communities containing ash trees and 68% of all street ash will

remain untouched by EAB by 2060 because of the lack of forest ash beyond our forecasted

invasion extent, thus limiting exposure. Spatially, our results show lower threat in the western

US. This pattern is consistent with previous findings (Lovett et al. 2016) and can be explained by

the high impacts of EAB, European gypsy moth, and hemlock woolly adelgid, whose

distributions are projected to concentrate further east in the short term. However, some of the

highest-impact non-native pathogens have emerged in the western US and were not captured in

this analysis (Kinloch Jr. 2003, Rizzo & Garbelotto 2003). Western regions could also see high

future risks due to the polyphagous shot hole borer (*Euwallacea whitfordiodendrus)* and its

insect-disease complex with fusarium fungus (*Fusarium spp.*) (Coleman et al. 2019). This

complex has already established in California and has maple and oak trees among its many hosts.

While the substantial advances that emerged recently allowed us to develop a more fully

integrated model, we also identified data deficiencies which require additional research. A

relative quantification of additional sources of uncertainty is provided in Appendix S3. Further,

this cost estimate only examines the cutting of dead trees. The analysis fails to account for

preventative cutting prior to EAB arrival, to fully examine non-street tree management, and to

assess the impacts of IAFIs that have not yet established in the United States. Furthermore, our

analysis assumes a complete identification of 'high impact IAFIs'. Some presently established

IAFIs may not yet have been identified as 'high impact', either due to lags in their impact, and/or

lags in the detection of this impact (Coutts et al. 2018), but may become 'high impact' before

2050.

We have shown that the suite of known IAFIs have the potential to kill roughly a hundred

million additional urban trees in the US in the next 30 years. While these numbers themselves

are striking, reporting only a country-level impact estimate without IAFI species, tree, and

community-level resolution does little to inform management prioritizations. Here, we were able

to identify specific urban centers, IAFI species, and host tree genera associated with the vast

majority of these impacts. We predict that 90% of all street tree mortality within the next 30

years will be EAB-induced ash mortality, and that ~95% of all street tree mortality will be

concentrated in less than 25% of all communities. These estimates illustrate the gravity of IAFI

infestations for communities in the path of high impact invaders that are rich in susceptible hosts.

Moreover, we were able to use this framework to identify a checklist of biotic and

spatiotemporal risk factors for future high-impact street tree IAFIs.

Acknowledgments

EJH would like to thank her PhD supervisory committee members T. Jonathan Davies and

Patrick M. A. James for their invaluable comments, as well as the thoughtful comments and

questions from thesis external examiner Dominique Gravel and colleague Andrew Liebhold. EJH

also acknowledges the continual support and feedback from lab members Dat Nguyen, Abbie

Gail Jones, Charlotte Steeves, Shriram Varadarajan, and Lidia Della Venezia. This work was

supported by a NSERC CGS-D awarded to EJH. This research was funded, in part, through Cost

Share Agreements 17-CS-11330110-025, 18-CS-11330110-026, and 19-CS-11330110-027,

between the USDA Forest Service and North Carolina State University.

References

Aukema, J. E., Leung, B., Kovacs, K., Chivers, C., Britton, K. O., Englin, J., ... *et al.* (2011).
Economic impacts of non-native forest insects in the continental United States. *PLoS*
One, 6(9).
Aukema, J. E., McCullough, D. G., Von Holle, B., Liebhold, A. M., Britton, K., & Frankel, S. J.
(2010). Historical accumulation of nonindigenous forest pests in the continental United
States. *BioScience*, 60(11), 886-897.
Carpenter, B., Gelman, A., Hoffman, M. D., Lee, D., Goodrich, B., Betancourt, M., ... *et al*.
(2017). Stan: A probabilistic programming language. *J Stat. Soft.*, 76(1).

Coleman, T. W., Poloni, A. L., Chen, Y., Thu, P. Q., Li, Q., Sun, J., ... *et al.* (2019). Hardwood injury and mortality associated with two shot hole borers, Euwallacea spp., in the invaded region of southern California, USA, and the native region of Southeast Asia. *Ann. Forest Sc.i*, 76(3), 1-18.

Coutts, S. R., Helmstedt, K. J., & Bennett, J. R. (2018). Invasion lags: The stories we tell ourselves and our inability to infer process from pattern. *Divers. Distrib*, 24(2), 244-251.

Fahrner, S. J., Abrahamson, M., Venette, R. C., & Aukema, B. H. (2017). Strategic removal of host trees in isolated, satellite infestations of emerald ash borer can reduce population growth. *Urban For. Urban Green.*, 24, 184-194.

Fei, S., Morin, R. S., Oswalt, C. M., & Liebhold, A. M. (2019). Biomass losses resulting from insect and disease invasions in US forests. *Proc. Natl. Acad. Sci.* 116(35), 17371-17376.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1‑km spatial resolution climate surfaces for global land areas. *Int. J Climatol.*, 37(12), 4302-4315.

Haack, R. A., Hérard, F., Sun, J., & Turgeon, J. J. (2010). Managing invasive populations of Asian longhorned beetle and citrus longhorned beetle: a worldwide perspective. *Ann. Rev. Entomol.*, 55.

Hijmans, R. J., Phillips, S., Leathwick, J., Elith, J., & Hijmans, M. R. J. (2017). Package 'dismo'. *Circles*, 9(1), 1-68.

Hilbert, D. R., Roman, L. A., Koeser, A. K., Vogt, J., & van Doorn, N. S. (2019). Urban tree mortality: a literature review. *Arbor. Urban For.*, 45(5), 167-200.

Hochberg, M.E., & Weis, A.E. (2001). Ecology: bagging the lag. *Nature* 409, 992–93.

Homer, C.G., Dewitz, J.A., Yang, L., Jin, S., Danielson, P., Xian, G., … *et al*. (2015.) Completion of the 2011 National Land Cover Database for the conterminous United States – representing a decade of land cover change information. *Photogramm. Eng. Remote Sensing,* 81, 345–354.

Hudgins, E. J., Liebhold, A. M., & Leung, B. (2017). Predicting the spread of all invasive forest pests in the United States. *Ecol. Lett.*, 20(4), 426-435.

Hudgins, E. J., Liebhold, A. M., & Leung, B. (2020). Comparing generalized and customized spread models for nonnative forest pests. *Ecol. Appl.,* 30(1), e01988.

Hulme, P. E. (2009). Trade, transport and trouble: managing invasive species pathways in an era of globalization. *J Appl. Ecol.*, 46(1), 10-18.

IPPC (International Plant Protection Convention). 2002. International standards for phytosanitary measures. Rome, Italy: FAO. www.fao.org/docrep/009/a0450e/a0450e00.htm. Viewed 12 May 2020.

Kinloch Jr, B. B. (2003). White pine blister rust in North America: past and prognosis. *Phytopathol.*, 93(8), 1044-1047.

Knight, K. S., Brown, J. P., & Long, R. P. (2013). Factors affecting the survival of ash (*Fraxinus* spp.) trees infested by emerald ash borer (*Agrilus planipennis*). *Biol. Invasions*, 15(2), 371-383.

Koch, F. H., Ambrose, M. J., Yemshanov, D., Wiseman, P. E., & Cowett, F. D. (2018). Modeling urban distributions of host trees for invasive forest insects in the eastern and central USA: A three-step approach using field inventory data. *Forest Ecol. Manage.*, 417, 222-236.

Kovacs, K. F., Haight, R. G., McCullough, D. G., Mercader, R. J., Siegert, N. W., & Liebhold, A. M. (2010). Cost of potential emerald ash borer damage in US communities, 2009–2019. *Ecol. Econ.*, 69(3), 569-578.

Kovacs, K., Václavík, T., Haight, R. G., Pang, A., Cunniffe, N. J., Gilligan, C. A., &
Meentemeyer, R. K. (2011). Predicting the economic costs and property value losses
attributed to sudden oak death damage in California (2010–2020). *J Environ.*
Manage., 92(4), 1292-1302.
Leung, B., Springborn, M. R., Turner, J. A., & Brockerhoff, E. G. (2014). Pathway-level risk
analysis: the net present value of an invasive species policy in the US. *Front. Ecol.*
Environ., *12*(5), 273-279.

Liebhold, A.M., & Tobin, P.C. (2008). Population ecology of insect invasions and their
management. *Ann. Rev. Entomol.,* 53, 387–408.
Lovett, G. M., Weiss, M., Liebhold, A. M., Holmes, T. P., Leung, B., Lambert, K. F., ... &
Weldy, T. (2016). Nonnative forest insects and pathogens in the United States: Impacts
and policy options. *Ecol. Appl.*, *26*(5), 1437-1455.
McGeoch, M. A., Genovesi, P., Bellingham, P. J., Costello, M. J., McGrannachan, C., &
Sheppard, A. (2016). Prioritizing species, pathways, and sites to achieve conservation
targets for biological invasion. *Biol. Invasions*, 18(2), 299-314.
Norton, B. A., Coutts, A. M., Livesley, S. J., Harris, R. J., Hunter, A. M., & Williams, N. S.
(2015). Planning for cooler cities: A framework to prioritise green infrastructure to
mitigate high temperatures in urban landscapes. *Landsc. Urban Plan.*, 134, 127-138.
Paap, T., Burgess, T. I., & Wingfield, M. J. (2017). Urban trees: bridge-heads for forest pest
invasions and sentinels for early detection. *Biol. Invasions*, 19(12), 3515-3526.
Potter, K. M., Escanferla, M. E., Jetton, R. M., & Man, G. (2019). Important insect and disease
threats to United States tree species and geographic patterns of their potential impacts.
Forests, 10(4), 304.
Pugh, S.A. (2010). *Michigan's forest resources, 2008*. Research Note NRS-50. Newtown
Square, PA: U.S. Forest Service, Northern Research Station. 4 pp.
Rizzo, D. M., & Garbelotto, M. (2003). Sudden oak death: endangering California and Oregon
forest ecosystems. *Front. Ecol. Environ.*, 1(4), 197-204.
Roy, S., Byrne, J., & Pickering, C. (2012). A systematic quantitative review of urban tree
benefits, costs, and assessment methods across cities in different climatic zones. *Urban*
For. Urban Green., 11(4), 351-363.
Sardain, A., Sardain, E., & Leung, B. (2019). Global forecasts of shipping traffic and biological
invasions to 2050. *Nat. Sust.*, 2(4), 274-282.
Van den Berg, A. E., Maas, J., Verheij, R. A., & Groenewegen, P. P. (2010). Green space as a
buffer between stressful life events and health. *Soc. Sci. Med.*, 70(8), 1203-1210.

Text Boxes

Text Box 1. Mortality debt explained.

The joint impact of maximum mortality and mortality debt is best illustrated by a series of examples. Estimates of street tree natural mortality range around 2.4-2.6% per year (Hilbert et al. 2019). Within a 30-year window, this would amount to roughly 53% natural street tree mortality. Our model assumes that if IAFI enters site at the beginning of this window (2020), it first undergoes a 10-year time lag, and can then cause mortality in the final 20 years. The maximum level of mortality induced by a borer (EAB on several *Fraxinus* spp., Category H = 98.98%), would result in 98.98% additional mortality (mortality of remaining the trees that survived natural mortality) at the end of a 30-year window. This level of mortality would be clearly detectable above natural street tree mortality. Hemlock woolly adelgid has a similar maximum mortality to EAB (Category H on *Tsuga* spp.), but we have assumed that sap-feeder mortality takes 100 years to reach asymptotic levels. As such, by the end of a 30-year window, only (98.98%/100)*20 years = 19.80% of additional host trees would be killed. While defoliators have shorter mortality debts, they tend to cause lower mortality, making their impacts the least detectable above background mortality. For defoliators, the IAFI with the greatest damage on any host is the larch casebearer, (*Coleophora laricella* on *Larix laricina*, category E = 16.46%). Given a 50-year mortality debt for defoliators, the maximum mortality above background rates by 2050 is (16.46% / 50) * 20 years = 6.58%. While these estimates are much lower, many host trees of sap feeders and defoliators are very common, and this mortality could very well be inflating the perceived background mortality rates of these host trees measured in (Hilbert et al. 2019).

EAB is estimated to kill the majority of its susceptible hosts in the first decade following infestation (Kovacs et al. 2014), while maximum mortality is estimated to take closer to 100 years for hemlock woolly adelgid (Aukema et al. 2011), so we used the 10 and 100-year scenarios for borers and sap-feeders, respectively. A recent publication examining mortality rates in forested areas suggested that European gypsy moth has a mortality rate intermediate between borers and sap-feeders, so we set defoliators at 50-years (20). Once an IAFI was predicted to infest an area, we imposed a 10-year initial lag phase between IAFI arrival at a site and the initial onset of damage (Hochberg & Weiss 2001, Liebhold & Tobin 2008) and then began increasing the host mortality following our mortality debt scenario to the asymptotic level (defined by the host mortality model).

540
541

Figures

Figure 1. Fit of the genus-specific host tree models across all genera and size classes.

Figure 2. Model outputs for the first three subcomponent models, including **a.** predicted street tree abundance, **b.** predicted newly invaded sites of existing IAFIs, **c.** predicted street tree exposure levels (number of focal host tree + IAFI interactions) from 2020 to 2050, and finally **d.** Predicted total tree mortality from 2020 to 2050 in the most likely mortality debt scenario across space. The top seven most impacted cities or groups of nearby cities are shown in terms of total tree mortality 2020 to 2050 (A = Milwaukee, WI; B = Chicago/Aurora/Naperville/Arlington Heights, IL; C = New York, NY; D = Seattle, WA; E = Indianapolis, IN; F = Cleveland, OH; G = Philadelphia, PA).

Figure 3. Posterior distribution for the beta model of host mortality due to IAFIs within each severity category. 95% Bayesian credible intervals are shown in grey, and the posterior median is shown in black. Colored bins represent severity categories extended from Potter et al. *2019).

Figure 4. Depiction of the influence of mortality debt on temporal cost patterns. Predicted costs 2020 to 2050 for the 10 year (yellow), 50 year (teal), and 100 year (purple) mortality debt scenarios with a 10 year initial invasion lag. The most likely scenario predictions are shown as a dashed red line. Costs are presented in 5-year increments in accordance with the timestep length within our spread model.

562   **Figure 1**

Figure 2

 Figure 3

Figure 4

572   **Tables**
573
Table 1. Predicted annualized cost (in 2019 US dollars) and tree mortality across invasion
scenarios from 2020 to 2050 across all 57 IAFI species. "Most likely" indicates the scenario with
expert-elicited mortality debt durations by feeding guild, "Vary" scenarios hold all guilds but the
focal guild constant at their most likely scenario, and "All" fix all three guilds at a given
mortality debt duration. Mean mortality for most likely scenario = 2.3%, 1.38M trees, US$ 30M
annualized (US$ 679M over the next 30 years).
580

Mortality Debt Scenario	Annualized Cost (US$ millions)		Tree Mortality (Millions)		Percent Mortality	
	lower 95% CI	upper 95% CI	lower 95% CI	upper 95% CI	lower 95% CI	upper 95% CI
Most likely	28.5	33.2	1.29	1.54	2.1%	2.5%
Vary Borers	10.1	32.1	0.45	1.45	0.7%	2.4%
Vary Defoliators	28.1	32.6	1.28	1.48	2.1%	2.4%
Vary Sap-feeders	28.5	32.5	1.30	1.47	2.1%	2.4%
All 10	27.8	30.4	1.27	1.39	2.1%	2.3%
All 50	18.5	22.3	0.84	1.00	1.4%	1.7%
All 100	9.77	13.5	0.44	0.60	0.7%	1.0%