Fluoroquinolone-Resistant *Escherichia coli* Originating on Dairy Farms are an

Infrequent Cause of Bacteriuria in Humans Living in the Same Geographical

Region

Oliver Mounsey[1], Hannah Schubert[2], Jacqueline Findlay[1], Katy Morley[2], Emma

F. Puddy[1,2], Virginia C. Gould[1,2], Paul North[3], Karen E. Bowker[3], O. Martin

Williams[4], Philip B. Williams[4], David C. Barrett[2], Tristan A. Cogan[2], Katy M.

Turner[2], Alasdair P. MacGowan[3], Kristen K. Reyher[2], Matthew B. Avison[1]*

[1]School of Cellular & Molecular Medicine, University of Bristol, Bristol, UK

[2]Bristol Veterinary School, University of Bristol, Bristol, UK

[3]Department of Microbiology, Infection Sciences, Southmead Hospital, North

Bristol NHS Trust, Bristol, UK

[4]Bristol Royal Infirmary, University Hospitals Bristol and Weston NHS

Foundation Trust, Bristol, UK

Synopsis

Background

Our primary aim was to test whether cattle-associated fluoroquinolone-resistant (FQ-R) *Escherichia coli* found on dairy farms were a significant cause of bacteriuria in humans living in the same 50 x 50 km geographical region located in South West England. Another aim was to identify risk factors for the presence of FQ-R *E. coli* on dairy farms.

Methods

FQ-R *E. coli* were isolated during 2017-18 from 42 dairy farms and from community urine samples. Forty-two cattle and 489 human urinary isolates were subjected to WGS, allowing phylogenetic comparisons. Risk factors were identified using a Bayesian regularisation approach.

Results

Of 489 FQ-R human isolates, 255 were also 3rd generation cephalosporin-resistant (3GC-R), with strong genetic linkage between *aac(6')Ib-cr* and *bla*$_{CTX-M-15}$. We identified possible farm-to-human sharing for pairs of ST744 and ST162 isolates, but core genome SNP distances (71 and 63, respectively) were smaller in pairs of ST744 and ST162 isolates from different farms (7 and 3 SNPs, respectively). Total farm fluoroquinolone use showed a positive association with the odds of isolating FQ-R *E. coli* while total dry cow therapy use showed a negative association.

Conclusions

This work suggests that FQ-R *E. coli* found on dairy farms have a limited impact on community bacteriuria within the local human population, however, this appears

40   greater than observed for 3GC-R *E. coli* when studied in parallel. Reducing

41   fluoroquinolone use may reduce the on-farm prevalence of FQ-R *E. coli*, and this

42   reduction may be greater when dry cow therapy is targeted to the ecology of

43   resistant *E. coli* on the farm.

Introduction

Fluoroquinolones are an important class of antibacterial drugs, included in the World Health Organisation's list of 'Highest Priority Critically Important Antimicrobials'.[1] They are extensively used to treat infections in humans, companion animals, and farmed animals. These important medicines work by disrupting the activity of type II topoisomerases, causing the release of DNA that has double-strand breaks, leading to cell death.[2]

In addition to safety concerns about fluoroquinolone use in humans,[3] bacteria such as *Escherichia coli* are increasingly becoming fluoroquinolone-resistant (FQ-R).[4] Here, FQ-R is mainly caused by mutations altering the quinolone resistance-determining regions (QRDRs) of the primary target, DNA gyrase subunit A (GyrA), and the secondary target, DNA topoisomerase IV subunits (ParC) and, to a lesser extent, ParE.[5, 6] However, additional mechanisms can also contribute to FQ-R, including regulatory mutations leading to AcrAB-TolC efflux pump over-production and expression of various plasmid-mediated quinolone resistance (PMQR) genes. For example, the various *qnr* genes, *oqxAB* efflux pump genes, and *aac(6')Ib-cr*, which encodes a mutated aminoglycoside-modifying enzyme.[5, 7]

Because most FQ-R in *E. coli* results from multiple mutations in target enzymes and because PMQRs cannot confer FQ-R alone, the spread of FQ-R *E. coli* typically involves vertical dissemination of resistant clones. Most commonly in humans, the widespread proliferation of an FQ-R subset of the ST131 complex,[8] which is also frequently associated with 3rd generation cephalosporin resistance (3GC-R).[9] Additionally, the pandemic ST1193 group,[10, 11] which is only occasionally 3GC-R.[10, 12]

In secondary care, the fluoroquinolone ciprofloxacin is commonly used as an oral

switch following intravenous therapy for serious infections caused by Gram negative

bacteria.[13] Accordingly, FQ-R is an important threat to human health, and particularly

when found in urinary *E. coli* since a substantial proportion of sepsis has an urinary

origin and *E. coli* are the leading cause of urosepsis.[14] The use of fluoroquinolones in

primary care in the UK has reduced in recent years, primarily due to changes in

policies concerning prescribing for community urinary tract infections. For example,

in the 1.5 million population centred on the city of Bristol, we recently demonstrated a

24% fall in dispensing of ciprofloxacin, by far the most used fluoroquinolone in

primary care, between 2013 and 2016. We noted a commensurate reduction in FQ-R

in community-origin urinary *E. coli* in the same region.[15]

Fluoroquinolones have also been extensively used for the treatment of companion[16]

and farmed[17] animals. Usage has decreased in farmed animals in the UK over

recent years[18] and voluntary use restrictions were introduced in June 2018 that

mean fluoroquinolones are only used as a last resort, backed up by susceptibility

testing[19]. There is still the possibility, however, that FQ-R *E. coli* that have been

selected on farms might colonise humans and ultimately cause disease. There is

also strong evidence that antibacterial resistant (ABR) *E. coli* are shared between

companion animals and humans.[20] Hence, considering FQ-R in humans through the

lens of the One Health research framework may help obtain a wider picture of

selection and transmission. Our primary aim was to test the hypothesis that FQ-R *E.*

coli obtained in 2017-18 from dairy farms located within a 50 x 50 km region of South

West England were closely related to FQ-R human urinary *E. coli* collected in

parallel from the same region, suggestive of transmission. Whilst our similarly

motivated studies of 3GC-R *E. coli* showed no evidence of recent sharing of isolates

between dairy farms and humans,[21, 22] we considered that the clonal nature of FQ-R

E. coli might give a different outcome.

Materials and Methods

Isolates.

Collection of FQ-R *E. coli* from dairy farms in South West England has been

described previously.[23] Briefly, 4145 samples were collected from faecally

contaminated sites around 53 dairy farms (49 farms yielded FQ-R *E. coli*). Samples

were collected at monthly visits between January 2017 and December 2018. FQ-R

isolates were selected by plating processed samples onto Tryptone Bile X-

Glucuronide (TBX) agar (Sigma-Aldrich, Dorset, UK) containing 0.5 mg/L

ciprofloxacin, the EUCAST breakpoint.[24] One FQ-R *E. coli* colony was picked from

each plate if growth occurred. FQ-R urinary *E. coli* isolates were obtained from

routine urine microbiology at Severn Infection Partnership, Southmead Hospital.

Urine samples were submitted between Sept 2017 and Dec 2018 from 146 GPs

located throughout Bristol and including coverage in Gloucestershire, Somerset, and

Wiltshire.[15] Bacterial identification was carried out using BD CHROMagar Orientation

Medium (BD, GmbH, Heidelberg, Germany). Antibiotic susceptibilities were defined

by disc testing using ciprofloxacin and ceftriaxone as indicators of FQ-R and 3GC-R

as interpreted according to EUCAST guidelines.[24] FQ-R/3GC-R urinary isolates and

FQ-R but 3GC-susceptbile (FQ-R/3GC-S) isolates were all subcultured onto TBX

agar containing 0.5 mg/L ciprofloxacin to confirm FQ-R status.

Ethics

118 Farmers gave fully informed consent to participate in this study. Ethical approval was

119 obtained from the University of Bristol's Faculty of Health Sciences Research Ethics

120 Committee (ref 41562). Ethical approval was not needed for the retention of human

121 urinary isolates since these were anonymous and isolated for routine diagnostics.

PCR, WGS, and Phylogenetic Analysis

PMQR genes (*qnrA*, *qnrB*, *qnrC*, *qnrD*, *qnrS*, *aac(6')-Ib-cr*, *oqxAB*, and *qepA*) were

identified by multiplex PCR as previously described.[25] WGS was performed by

MicrobesNG (https://microbesng.uk/) as previously described.[9, 22] Resistance genes

and STs were assigned using ResFinder [26] and MLST 2.0 [27] on the Centre for

Genomic Epidemiology (http://www.genomicepidemiology.org/) platform.

Phylogenetic analysis was carried out using the Bioconda environment [28] on the

Cloud Infrastructure for Microbial Bioinformatics.[29] The reference sequences were

ST131 isolate EC958 (accession: HG941718), ST744 isolate EC590 (accession:

NZ_CP016182), and ST162 isolate W2-5 (accession: NZ_CP032989). Sequence

alignments were with Snippy (https://github.com/tseemann/snippy). Maximum

likelihood phylogenetic trees were constructed using RAxML, utilising the GTRCAT

model of rate heterogeneity and the software's autoMR and rapid bootstrap.[30, 31]

Trees were illustrated using Microreact.[32]

Risk Factor Analysis

The risk factors fell into four categories: farm management, antibacterial usage

(ABU), sample characteristics and meteorological. These categories, and the

methods of data collection for each, are described in our recent wider risk factor

study and its supplementary information.[23] All code can be found at

https://github.com/HannahSchubert1/OH-STAR-modelling-code. A random intercept

Bayesian model with farm as a random effect was fitted using R with the BRMS

(https://cran.r-project.org/web/packages/brms/index.html) package. The presence of

FQ-R *E. coli* within a sample was the dependent variable. The remaining variables

were all fixed effects and were split into two groups: "main effects" which were the

ABU variables (measured as milligram per population-corrected unit) hypothesised

to affect FQ-R *E. coli* presence, namely, total fluoroquinolone usage, novobiocin

usage, and total 3^{rd} and 4^{th} generation cephalosporin usage; and "regularised

effects" which were all remaining variables.

For the main effects, uninformative priors (normal distribution with a mean 0,

standard deviation 5) were used; for the regularised effects, a regularizing prior

(horseshoe prior with a single degree of freedom) [33] was used. The mean intercept

was also given an uninformative prior (normal distribution with mean 0, standard

deviation 5) while the standard deviation of the random effects was given the default

prior for BRMS (a half-Student-t distribution with 3 degrees of freedom and a scale

factor of 10). Four chains were sampled with 2000 iterations. The target acceptance

criterion was increased from the default 0.8 to 0.98 to decrease the chance of

sampling divergencies.

Results

Molecular epidemiology of FQ-R human urinary E. coli

A total of 489 FQ-R human urinary *E. coli* isolates were collected; 255 were also 3GC-R, 234 were 3GC-S. According to PMQR gene multiplex PCR, nineteen isolates harboured *qnr* genes (14 *qnrS*, 5 *qnrB*) and two carried *oqxAB*. Notably, 135 (52.9%) of the 3GC-R isolates but only 26 (11.1%) of 3GC-S isolates carried *aac(6')Ib-cr* (Chi square: 96.7, p<0.0001).

WGS was performed for 188 FQ-R urinary isolates; 90 were 3GC-R and 98 were 3CG-S. **Figure 1** shows a phylogenetic tree drawn based on core genome alignment for these isolates. As well as illustrating the dominance of ST131 (mainly 3GC-R, green markers) and ST1193 (mainly 3GC-S, orange markers) this tree also shows that clusters of isolates fell into ST744, ST162, ST69 and ST38 (**Figure 1**). A total of 22 different combinations of QRDR mutations were identified in the 188 sequenced isolates. However, a large majority (155/188) fell within three QRDR types, identified here a Types A, B and C (**Table 1**). Isolates within common FQ-R clonal groups ST131 and ST1193 were dominated by QRDR Types A and B, respectively, whilst QRDR Type C was found in isolates from 13 diverse STs.

WGS also confirmed the strong bias for *aac(6')Ib-cr* gene carriage, with 39 (43.3%) 3GC-R isolates being *aac(6')Ib-cr*-positive, versus only four (4.1%) of the 3GC-S isolates (Chi square: 16.5, p<0.0001). We hypothesised that physical linkage of *aac(6')Ib-cr* with an over-represented 3GC-R gene was driving this bias and **Table 2** reports that carriage of *aac(6')Ib-cr* was strongly associated with carriage of *bla*$_{CTX-M-15}$. We thus further hypothesised that this was due to physical linkage of *bla*$_{CTX-M-15}$

with the class 1 integron In*37*, carrying *aac(6')Ib-cr*. Even given the difficulties of

interpreting short-read WGS data, there was evidence in favour of this hypothesis.

Of 11 isolates where the genomic locations of these genes could be definitively

identified, in three, In*37* and *bla*$_{CTX-M-15}$ were both embedded (at separate locations)

in contigs surrounded by known chromosomal genes. In eight, In*37* and *bla*$_{CTX-M-15}$

were immediately adjacent on the same contig, sometimes surrounded by known

chromosomal genes, and sometimes not, suggesting a plasmid location.

3GC-R ST131 urinary *E. coli* isolates commonly carry *bla*$_{CTX-M-15}$ in this study

region.[9] Therefore, it was not surprising to find a positive association between

aac(6')Ib-cr carriage and ST131 and between *aac(6')Ib-cr* carriage and QRDR Type

A (**Table 2**). However, of 32 FQ-R/3GC-S ST131 *E. coli*, only four carried *aac(6')Ib-*

cr, confirming *aac(6')Ib-cr* was linked with *bla*$_{CTX-M-15}$ and not with ST131.

In contrast, ST1193 isolates in this study were found mainly to be 3GC-S (n=28,

versus n=6 for 3GC-R; **Figure 1**), and there was a significant negative association

between *aac(6')Ib-cr* carriage and the isolate being ST1193 (**Table 2**). There was

also a negative association between carriage of *bla*$_{CTX-M-27}$ and carriage of *aac(6')Ib-*

cr (**Table 2**). These associations reinforce the conclusion that In*37* and *bla*$_{CTX-M-15}$

are closely genetically linked in urinary *E. coli* in our study region. No other

association tested was found to be statistically significant.

Molecular epidemiology of FQ-R E. coli from dairy farms and influence on human

UTIs

We have recently reported a survey of resistant *E. coli* in faecal samples from 53

dairy farms in South West England.[23] Of 4145 faecal samples collected from

vicinities near animals, 263 were positive for FQ-R *E. coli*, representing 49 of 53

farms surveyed.[23] Of 42 farms located within a 50 x 50 km region, which included the

homes of the people providing the urinary samples discussed above, 245 FQ-R *E.*

coli isolates from this earlier study [23] were taken forward for further analyses,

representing one unique isolate from each positive sample. Multiplex PCRs showed

that *aac(6')Ib-cr* (found to be highly prevalent in the human urinary *E. coli* isolates)

was entirely absent from the farm isolates and only nine isolates carried a PMQR

gene (5 *qnrS*, 2 *qnrA* and 2 *qnrD)*.

WGS was performed for 42 FQ-R cattle-associated *E. coli* which were selected so

that at least one isolate represented each farm that was positive for FQ-R *E. coli*

within the 50 x 50 km region. Nine different STs were identified, with dominance of

ST744 (21/42 isolates), ST162 (8/42 isolates), and ST10 (5/42 isolates). ST162 and

ST10 have QRDR Type C, with three mutations as defined above (**Table 1**). ST744

also has these same three QRDR mutations, plus the addition of a distinctive *parC*

mutation causing A56T. Because ST744 and ST162 were also common among FQ-

R/3GC-S human isolates (**Figure 1**), we generated a phylogenetic tree to consider

relationships between the cattle FQ-R and human FQ-R/3GC-S *E. coli* collected in

parallel within our 50 x 50 km study area (**Figure 2**). This tree suggested overlapping

ST744 and ST162 isolates found on different farms, but, importantly, also among

human urinary isolates. Detailed trees generated using ST744 or ST162 reference

genomes (**Figure 3**) confirmed this. The closest relationship between two human

urinary isolates was 931 or 175 SNPs for ST744 and ST162, respectively, but the

closest relationship between a human and a cattle isolate was 71 or 63 SNPs,

respectively.

Fluoroquinolone use as a driver of FQ-R E. coli on dairy farms and the suppressive

effect of dry cow therapy

We next considered factors that might influence the prevalence of FQ-R *E. coli* on

farms. Our aim was to identify potential interventions that might reduce FQ-R *E. coli*

on farms, and so reduce any ongoing zoonotic threat. In our recent paper, we

showed that the odds of a sample being positive for FQ-R *E. coli* was significantly

greater if the sample came from the environment of heifer calves.[23] This is important

because dairy heifer calves are normally reared on the farm as replacement milking

cows, so factors associated with the increased carriage of FQ-R *E. coli* in calves

could have a long-term effect on the whole farm. Accordingly, we performed a new

risk factor analysis to identify management and ABU factors associated with the

odds of finding FQ-R *E. coli*-positive samples in the environment of heifer calves.

Of 631 samples collected from the environments of heifer calves, 103 (16.3%) were

positive for FQ-R *E. coli*. We identified two variables that were associated with

sample-level positivity for FQ-R *E. coli*. One variable - the total usage of

fluoroquinolones in the year the samples were collected - was positively associated

with finding FQ-R *E. coli* in a sample (odds ratio 2.39, 95% credible interval [1.01,

6.02]). Another variable - the percentage of cows within the herd dried off using any

antibacterial dry cow therapy (a preparation inserted into the teats of pregnant cows

thought to be at risk of developing mastitis between lactations, or with existing

subclinical infections) - was negatively associated with finding FQ-R *E. coli* in

samples (odds ratio 0.24, 95% credible interval [0.11, 0.50]). Full model outputs and

model checking is presented in Supplementary.

Discussion

We have shown in previous similarly powered work performed in parallel on the

same study farms that there was no evidence of recent sharing of 3GC-R *E. coli*

between farms and humans in the study region.[21, 22] This agrees with the findings of

similarly motivated studies from other groups.[34-36] However, when considering FQ-R

E. coli populations, we now report evidence for mixing of farm and human *E. coli*,

and that the shared *E. coli* clones can cause bacteriuria.

A core genome SNP distance of 30 or fewer is commonly seen in phylogenetic

analyses of Enterobacteriales isolates that are confirmed to be part of an acute

outbreak of foodborne illness [37] and hospital studies frequently set a cut-off of <100

SNPs to define an outbreak.[38] Finding FQ-R human/cattle isolates pairs differing by

71 (ST744) or 63 (ST162) SNPs is therefore suggestive of a situation where human

and cattle isolates in this region do intermingle. However, this observation should be

considered in the context of our finding that the closest isolates from two different

farms were only three and seven SNPs apart for ST162 and ST744, respectively

(**Figure 3**).

We conclude therefore, that whilst farm-related strains made up a small component

of the total FQ-R urinary *E. coli* in our survey, there is evidence for zoonotic

transmission to humans (**Figure 3**), which contrasts with studies considering 3GC-R

E. coli.[21, 22, 34-36] More work is needed to establish the exact routes of transmission,

but even the minor zoonotic potential of cattle-associated FQ-R *E. coli* identified here

should act as a stimulus to reduce the prevalence of such bacteria on farms. This

work also suggests potential ways to achieve that objective.

We have reported at a regional level that reducing fluoroquinolone use in primary

care was associated with a reduction in the proportion of *E. coli*-positive community

urine samples where the isolate was FQ-R.[15] Accordingly, it was interesting to find

that overall fluoroquinolone use at farm level was positively associated with the odds

of finding FQ-R *E. coli*-positive faecal samples in the environments around dairy

heifer calves. The implication is that reducing fluoroquinolone use on farms may well

reduce the prevalence of FQ-R *E. coli* in heifer calves, which, as they join the milking

herd, may lead to a general reduction of FQ-R *E. coli* on the farm.

Notably, in mid-2018, as our surveillance of resistant *E. coli* on study farms was

ending, the use of fluoroquinolones was effectively stopped, except in the very rare

instance where susceptibility testing confirmed that no other antibacterial treatment

option was available.[19] This Red Tractor farm assurance scheme regulation is

applicable to the vast majority of UK dairy farms.

Our final key finding was that dry cow therapy use may expedite the reduction of FQ-

R *E. coli* in heifer calves, which was unexpected. We hypothesised that the reason

for the association between increased usage of dry cow therapy on a farm and

reduced prevalence of FQ-R *E. coli* in the environments of heifer calves was that

relatively few FQ-R *E. coli* from these farms were cross-resistant to the antibacterial

ingredients of dry cow therapies. These antibacterials can be released in the

colostrum and first milk from treated cows. Since this colostrum is usually fed to

calves at birth,[39-41] it is plausible that calves receiving colostrum from treated cows

are protected from colonisation by FQ-R *E. coli* due to the dose of antibacterial

inadvertently received shortly after birth.

We calculated that 83% (in terms of weight of active ingredient) of the dry cow

therapy anti-Gram-negative antibacterials used on study farms during the period of

our project (2017-2018) were cephalosporins or cloxacillin (3.44 kg). Framycetin

(neomycin B) made up the remainder (0.69 kg). Notably, of the 42 FQ-R cattle *E. coli*

isolates subjected to WGS, only two (4.8%) were resistant to both cephalosporins

and cloxacillin, as inferred from WGS.

Whilst genetically inferred framycetin resistance (presence of an *aph* gene) was

more common among sequenced FQ-R cattle *E. coli* isolates (15/42, 36%), it was far

less common among FQ-R isolates than among CTX-M β-lactamase-positive 3GC-R

E. coli isolates from these same farms collected in parallel (109/135, 81% of

isolates).[22] Accordingly, there is evidence for suppression of FQ-R *E. coli* by dry cow

therapy use, irrespective of active agent, because the FQ-R bacteria found on these

farms are rarely resistant to the active agents used.

Much attention has been paid to reducing dry cow therapy use on dairy farms, as a

way of reducing total ABU.[39-41] We would not suggest a shift away from this

approach, because inappropriate use might increase the selection of zoonotic threat

of cattle *E. coli* that are resistant to other antibacterials. Indeed, we have already

shown a positive association between cefquinome dry cow therapy use on dairy

farms and *E. coli* carrying CTX-M type β-lactamases,[23] the most common cause of

3GC-R in *E. coli* isolated from community UTIs in our region.[9] Whilst use of

cefquinome in dry cow therapy is also effectively stopped under the 2018 Red

Tractor regulations,[19] it is certainly possible that switching to the use of first-

generation cephalosporins or cloxacillin dry cow therapy could maintain selection for

CTX-M producers, which are resistant to these agents. Importantly, however, 50% of

3GC-R on our study farms was caused by chromosomally encoded AmpC hyper-

production,[21, 22] an enzyme inhibited by cloxacillin.[42] We suggest, therefore, that a

better alternative to cefquinome dry cow therapy might be cloxacillin, if all else is

equal.

In contrast, a switch from cefquinome to framycetin-containing dry cow therapy might

be less favoured because, as we have shown previously, framycetin dry cow therapy

use co-selects CTX-M positive strains.[23] Furthermore, whilst we have identified that

framycetin resistance is less common in FQ-R $E.\ coli$ than in bla_{CTX-M} positive $E.\ coli$

on farms (36% versus 81%), we found framycetin resistance to be more common

than cloxacillin resistance in FQ-R $E.\ coli$ (36% versus 4.8%), supporting the use of

cloxacillin over framycetin or a first-generation cephalosporin as the best choice to

drive down FQ-R $E.\ coli$ whilst less strongly selecting for 3GC-R $E.\ coli$ (i.e. actively

selecting against the proportion caused by AmpC enzymes).

Overall, our One Health approach to investigating selection and transmission of

critically important ABR $E.\ coli$ within our study region highlights the rare but not

insignificant zoonotic potential for cattle-associated FQ-R $E.\ coli$ as a cause of

bacteriuria in humans. Our results also highlight that reducing fluoroquinolone use on

farms, whilst carefully selecting the most appropriate dry cow therapy active

ingredient to match the ecology of resistance found on a particular farm, should most

effectively reduce that zoonotic potential. Our work certainly demonstrates the

foresight of the recently introduced Red Tractor regulations designed to effectively

eliminate use of highest-priority critically important antibacterials on dairy farms in

the UK.[19]

Acknowledgements

Genome sequencing was provided by MicrobesNG (http://www.microbesng.uk). We wish to thank all the farmers and veterinary surgeons who participated in this study and the biomedical scientists and other laboratory staff at Severn Pathology, Southmead Hospital, Bristol, for assistance in collecting urinary isolates. We are grateful to Dr Rob Arbon, Jean Golding Institute, University of Bristol, for advice about the Bayesian modelling work.

Funding

This work was funded by grant NE/N01961X/1 to M.B.A., K.K.R., K.M.T., D.C.B., A.P.M. and T.A.C., grant MR/S004769/1 to M.B.A., K.K.R. and K.M.T., and grant BB/T004592/1 to K.K.R and M.B.A. from the Antimicrobial Resistance Cross Council Initiative supported by the seven United Kingdom research councils and the National Institute for Health Research. This work was also funded by grant MR/T005408/1 to P.B.W. and M.B.A. from the Medical Research Council.

Transparency declaration

D.C.B. was president of the British Cattle Veterinary Association 2018-19. Otherwise, the authors declare no competing interests. Farming and veterinary businesses who contributed data and permitted access for sample collection were not involved in the design of this study or in data analysis and were not involved in drafting the manuscript for publication.

Author Contributions

Conceived the Study: D.C.B., K.K.R., M.B.A.

Collection of Data: O.M., H.S., J.F. E.F.P., K.M. P.N., supervised by K.B., A.P.M., T.A.C., K.K.R., M.B.A.

Cleaning and Analysis of Data: O.M., H.S., E.F.P., V.C.G., supervised by O.M.W., P.B.W., K.M.T., K.K.R., M.B.A.

Initial Drafting of Manuscript: O.M., H.S., M.B.A.

Corrected and Approved Manuscript: All Authors.

References

1.	World Health Organisation. WHO Critically Important Antimicrobials for Human Medicine 6th revision. https://www.who.int/foodsafety/publications/antimicrobials-sixth/en/ (23rd February 2021, date last accessed).

2.	Aldred KJ, Kerns RJ, Osheroff N. Mechanism of quinolone action and resistance. *Biochemistry* 2014; **53**: 1565-74.

3.	Yu X, Jiang DS, Wang J et al. Fluoroquinolone Use and the Risk of Collagen-Associated Adverse Events: A Systematic Review and Meta-Analysis. *Drug Safety* 2019; **42**: 1025-33.

4.	Fasugba O, Gardner A, Mitchell BG et al. Ciprofloxacin resistance in community- and hospital-acquired *Escherichia coli* urinary tract infections: a systematic review and meta-analysis of observational studies. *BMC Infectious Diseases* 2015; **15**: 545.

5.	Webber M, Piddock LJ. Quinolone resistance in *Escherichia coli*. *Veterinary Research* 2001; **32**: 275-84.

6.	Park KS, Yang HS, Nam YS et al. Mutations in DNA Gyrase and Topoisomerase IV in Ciprofloxacin-Nonsusceptible Extended-Spectrum -Lactamase-Producing *Escherichia coli* and *Klebsiella pneumoniae*. *Clinical Laboratory* 2017; **63**: 535-41.

7.	Morgan-Linnell SK, Zechiedrich L. Contributions of the Combined Effects of Topoisomerase Mutations toward Fluoroquinolone Resistance in *Escherichia coli*. *Antimicrobial Agents and Chemotherapy* 2007; **51**: 4205-8.

8.   Stoesser N, Sheppard AE, Pankhurst L et al. Evolutionary History of the

Global Emergence of the *Escherichia coli* Epidemic Clone ST131. *mBio* 2016;

7: e02162-15.

9.   Findlay J, Gould VC, North P et al. Characterization of cefotaxime-resistant

urinary *Escherichia coli* from primary care in South-West England 2017–18.

Journal of Antimicrobial Chemotherapy 2019; **75**: 65-71.

10. Johnson TJ, Elnekave E, Miller EA et al. Phylogenomic Analysis of

Extraintestinal Pathogenic *Escherichia coli* Sequence Type 1193, an

Emerging Multidrug-Resistant Clonal Group. *Antimicrobial Agents and*

Chemotherapy 2019; **63**: e01913-18.

11. Tchesnokova V, Radey M, Chattopadhyay S et al. Pandemic fluoroquinolone

resistant *Escherichia coli* clone ST1193 emerged via simultaneous

homologous recombinations in 11 gene loci. *Proceedings of the National*

Academy of Sciences 2019; **116**: 14740-8.

12. Tchesnokova VL, Rechkina E, Larson L et al. Rapid and Extensive Expansion

in the United States of a New Multidrug-resistant *Escherichia coli* Clonal

Group, Sequence Type 1193. *Clinical Infectious Siseases* 2019; **68**: 334-7.

13. Kalra OP, Raizada A. Approach to a patient with urosepsis. *Journal of Global*

Infectious Diseases 2009; **1**: 57-63.

14. Tamma PD, Conley AT, Cosgrove SE et al. Association of 30-Day Mortality

With Oral Step-Down vs Continued Intravenous Therapy in Patients

Hospitalized With Enterobacteriaceae Bacteremia. *Journal of the American*

Medical Association Internal Medicine. 2019; **179**: 316-23.

15.   Hammond A, Stuijfzand B, Avison MB et al. Antimicrobial resistance

437   associations with national primary care antibiotic stewardship policy: Primary

438   care-based, multilevel analytic study. *PloS One* 2020; **15**: e0232903.

439   16.   Prescott JF, Boerlin P. Antimicrobial use in companion animals and Good

440   Stewardship Practice. *Veterinary Record* 2016; **179**: 486-8.

441   17.   Magouras I, Carmo LP, Stärk KDC et al. Antimicrobial Usage and -Resistance

442   in Livestock: Where Should We Focus? *Frontiers in Veterinary Science* 2017;

443   **4**: 148.

444   18.   Veterinary Medicines Directorate. Veterinary Antimicrobial Resistance and

445   Sales   Surveillance   2019.

446   https://www.gov.uk/government/publications/veterinary-antimicrobial-

447   resistance-and-sales-surveillance-2019 (22/03/21 2021, date last accessed).

448   19.   Assured   Food   Standards.   Red   Tractor   Farm   Standards.

449   https://assurance.redtractor.org.uk/standards   (22/03/21   2021,   date   last

450   accessed).

451   20.   Toombs-Ruane LJ, Benschop J, French NP et al. Carriage of extended-

452   spectrum beta-lactamase-and AmpC beta-lactamase-producing *Escherichia*

453   *coli* from humans and pets in the same households. *Applied and*

454   *Environmental Microbiology*. 2020. **86**: e01613-20.

455   21.   Alzayn M, Findlay J, Schubert H et al. Characterization of AmpC-

456   hyperproducing *Escherichia coli* from humans and dairy farms collected in

457   parallel in the same geographical region. *Journal of Antimicrobial*

458   *Chemotherapy* 2020; **75**: 2471-9.

459   22.   Findlay J, Mounsey O, Lee WWY et al. Molecular Epidemiology of

460   *Escherichia coli* Producing CTX-M and pAmpC β-Lactamases from Dairy

Farms Identifies a Dominant Plasmid Encoding CTX-M-32 but No Evidence

for Transmission to Humans in the Same Geographical Region. *Applied and*

Environmental Microbiology 2020; 87: e01842-20.

23.   Schubert H, Morley K, Puddy EF et al. Reduced Antibacterial Drug

Resistance and *bla*$_{CTX-M}$ β-Lactamase Gene Carriage in Cattle-Associated

Escherichia coli at Low Temperatures, at Sites Dominated by Older Animals,

and on Pastureland: Implications for Surveillance. *Applied and Environmental*

Microbiology 2021; **87**: e01468-20.

24.   EUCAST. Breakpoint tables for interpretation of MICs and zone diameters.

Version

8.1.http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoin

t_tables/v_8.1_Breakpoint_Tables.pdf.

25.   Ciesielczuk H, Hornsey M, Choi V et al. Development and evaluation of a

multiplex PCR for eight plasmid-mediated quinolone-resistance determinants.

Journal of Medical Microbiology 2013; **62**: 1823-7.

26.   Zankari E, Hasman H, Cosentino S et al. Identification of acquired

antimicrobial resistance genes. *Journal of Antimicrobial Chemotherapy* 2012;

67: 2640-4.

27.   Wirth T, Falush D, Lan R et al. Sex and virulence in *Escherichia coli*: an

evolutionary perspective. *Molecular Microbiology* 2006; **60**: 1136-51.

28.   Grüning B, Dale R, Sjödin A et al. Bioconda: sustainable and comprehensive

software distribution for the life sciences. *Nature Methods* 2018; **15**: 475-6.

29.   Connor TR, Loman NJ, Thompson S et al. CLIMB (the Cloud Infrastructure for

Microbial Bioinformatics): an online resource for the medical microbiology

community. *Microbial Genomics* 2016; **2**: e000086.

30.   Stamatakis A. RAxML-VI-HPC: maximum likelihood-based phylogenetic

analyses with thousands of taxa and mixed models. *Bioinformatics* 2006; **22**:

2688-90.

31.   Stamatakis A, Meier H, Ludwig T. RAxML-III: a fast program for maximum

likelihood-based inference of large phylogenetic trees. *Bioinformatics* 2004;

21: 456-63.

32.   Argimon S, Abudahab K, Goater RJ et al. Microreact: visualizing and sharing

data for genomic epidemiology and phylogeography. *Microbial Genomics*

2016; **2**: e000093.

33.   Carvalho CM, Polson NG, Scott JG. The horseshoe estimator for sparse

signals. *Biometrika* 2010; **97**: 465-80.

34.   Day MJ, Hopkins KL, Wareham DW et al. Extended-spectrum β-lactamase-

producing *Escherichia coli* in human-derived and foodchain-derived samples

from England, Wales, and Scotland: an epidemiological surveillance and

typing study. *Lancet Infectious Diseases* 2019; **19**: 1325-35.

35.   Ludden C, Raven KE, Jamrozy D et al. One Health Genomic Surveillance of

Escherichia coli Demonstrates Distinct Lineages and Mobile Genetic

Elements in Isolates from Humans versus Livestock. *mBio* 2019; **10**: e02693-

18.

36.   Montealegre MC, Talavera Rodríguez A, Roy S et al. High Genomic Diversity

and Heterogenous Origins of Pathogenic and Antibiotic-Resistant *Escherichia*

coli in Household Settings Represent a Challenge to Reducing Transmission

in Low-Income Settings. *mSphere* 2020; **5**: e00704-19.

37. Pightling AW, Pettengill JB, Luo Y et al.Interpreting Whole-Genome Sequence

Analyses of Foodborne Bacteria for Regulatory Applications and Outbreak

Investigations. Frontiers in Microbiology. 2018; **9**: 1482.

38. Constantinides B, Chau, KK, Quan TP et al. Genomic surveillance of

Escherichia coli and *Klebsiella* spp. in hospital sink drains and patients.

Microbial Genomics 2020; **6**: mgen000391.

39. Morgans LC, Bolt S, Bruno-McClung E et al. A participatory, farmer-led

approach to changing practices around antimicrobial use on UK farms.

Journal of Dairy Science 2021; **104**: 2212-30.

40. Rowe SM, Godden SM, Nydam DV et al. Randomized controlled non-

inferiority trial investigating the effect of 2 selective dry-cow therapy protocols

on antibiotic use at dry-off and dry period intramammary infection dynamics.

Journal of Dairy Science 2020; **103**: 6473-92.

41. Scherpenzeel CGM, den Uijl IEM, van Schaik G et al. Effect of different

scenarios for selective dry-cow therapy on udder health, antimicrobial usage,

and economics. *Journal of Dairy Science* 2016; **99**: 3753-64.

42. Tzouvelekis LS, Tzelepi E, Mentis AF et al. Identification of a novel plasmid-

mediated beta-lactamase with chromosomal cephalosporinase characteristics

from *Klebsiella pneumoniae*. *The Journal of Antimicrobial Chemotherapy*

1993; **31**: 645-54.

Tables

QRDR Type	Associated ST	Mutations relative to wildtype strain (K12)	Number of isolates (of 188)
Type A	ST131	*gyrA* S83L, *gyrA* D87N, *parC* S80I, *parC* E84V, *parE* I529L	90
Type B	ST1193	*gyrA* S83L, *gyrA* D87N, *parC* S80I, *parE* L416F	37
Type C	Thirteen STs	*gyrA* S83L, *gyrA* D87N, *parC* S80I	28

533 **Table 1.** Common QRDR mutation patterns identified within FQ-R human urinary *E.*

534 *coli.*

$aac(6')Ib\text{-}cr$ carriage versus:	Chi square	Association	p-value
ST131	6.2	Positive	0.013
ST1193	4.3	Negative	0.038
ST38	1.9	Positive	0.17
$bla_{CTX\text{-}M\text{-}15}$ carriage	47.8	Positive	<0.0001
$bla_{CTX\text{-}M\text{-}27}$ carriage	5.0	Negative	0.025
QRDR Type A	8.7	Positive	0.003
QRDR Type B	3.5	Negative	0.060
QRDR Type C	3.0	Negative	0.083

538 **Table 2.** Chi square analyses examining the association of $aac(6')Ib\text{-}cr$ carriage with

539 other different genotypic factors.

540 All comparisons had degrees of freedom = 1.

 Figures

Figure 1. Maximum likelihood phylogenetic tree showing FQ-R urinary isolates.
Green nodes were also 3GC-R, whilst yellow nodes were 3GC-S. Sequences were
aligned against an ST131 reference (grey node). STs have been indicated where
more than five isolates from that ST were present.

Figure 2. Maximum likelihood phylogenetic tree showing FQ-R, 3GC-S urinary
isolates (orange) and cattle isolates (blue). Sequences were aligned against an
ST131 reference (black node). Prevalent STs have been indicated.

Figure 3. Maximum likelihood phylogenetic tree showing (A) ST744 isolates and (B) ST162 isolates also present in Figure 2. 3GC-S urinary isolates (orange) and isolates collected from environments nearby to cattle housing (blue). Sequences were aligned against references of the same ST (black node).