An RNA-based feed-forward mechanism ensures motor switching in *oskar* mRNA transport

Imre Gáspár[1,2] *, Ly Jane Phea[1], Mark A. McClintock[3], Simone Heber[1], Simon L. Bullock[3] * and Anne Ephrussi[1] *

1 Developmental Biology Unit, European Molecular Biology Laboratory, 69117 Heidelberg, Germany

2 Current address: Turbine.ai, Budapest, Hungary

3 Division of Cell Biology, MRC Laboratory of Molecular Biology, Cambridge, United Kingdom

*Correspondence: imre.gaspar@turbine.ai, sbullock@mrc-lmb.cam.ac.uk, anne.ephrussi@embl.org

ABSTRACT

Regulated recruitment and activity of motor proteins is essential for intracellular transport of cargoes, including messenger ribonucleoprotein complexes (RNPs). Here we show that orchestration of *oskar* RNP transport in the *Drosophila* germline relies on the interplay of two double-stranded RNA binding proteins, Staufen and the dynein adaptor Egalitarian (Egl). We find that Staufen antagonizes Egl-mediated transport of *oskar* mRNA by dynein both *in vitro* and *in vivo*. Following delivery of nurse cell-synthesized *oskar* mRNA into the oocyte by dynein, recruitment of Staufen to the RNPs results in dissociation of Egl and a switch to kinesin-1-mediated translocation of the mRNA to its final destination at the posterior pole of the oocyte. We additionally show that Egl associates with *staufen* (*stau*) mRNA in the nurse cells, mediating its enrichment and translation in the ooplasm. Our observations identify a novel feed-forward mechanism, whereby dynein-dependent accumulation of *stau* mRNA, and thus protein, in the oocyte enables motor switching on *oskar* RNPs by downregulating dynein activity.

INTRODUCTION

Proper execution of the genetic program relies on regulatory mechanisms that act at multiple levels of gene expression, from transcription to post-translational events. In many cell types, including neurons, early embryos and oocytes, a key process that regulates genetic output at the subcellular level is RNA localization, which is often driven by cytoskeletal motors (St Johnston, 2005; Gaspar, 2011; Marchand *et al.* 2012; Glock *et al.* 2017; Mofatteh and Bullock, 2017). The kinesin-1 and cytoplasmic dynein-1 (dynein) motors, which move RNPs towards the plus end and minus end of microtubules, respectively, play a central role in mRNA trafficking in several systems (Gaspar, 2011; Mofatteh and Bullock, 2017).

A paradigm for the study of microtubule-based RNP transport is *oskar* mRNA, the protein product of which induces abdomen and germline formation in the embryo (Ephrussi and Lehmann, 1992). *oskar* mRNA is produced in the nurse cells of the *Drosophila* germline syncytium in the early stages of oogenesis and transported by dynein into the developing oocyte where microtubule minus ends are enriched (Ephrussi *et al.* 1991; Januschke *et al.*, 2002; Clark *et al.* 2007; Sanghavi *et al.*, 2013; Jambor *et al.*, 2014; Vazquez-Pianzola *et al.*, 2017). mRNA trafficking from the nurse cells to the oocyte is dependent on the Egl protein, which binds specialized stem-loop structures in localizing mRNAs and links them together with its coiled-coil binding partner Bicaudal-D (BicD) to dynein and the dynein activating complex dynactin (Navarro *et al.*, 2004; Dienstbier *et al.*, 2009; Dix *et al.*, 2013). *In vitro* reconstitution studies have shown that Egl and BicD also play a key role in switching on dynein motility (McClintock *et al.*, 2018; Sladewski *et al.*, 2018).

During mid-oogenesis, when the microtubule cytoskeleton is reorganized, most of Egl's mRNA targets localize to the anterior of the oocyte, where microtubule minus ends are now concentrated (Gaspar, 2011; Lasko, 2012). In contrast, *oskar* mRNA is transported to the posterior pole, where the microtubule plus ends are focused (Parton *et al.*, 2011); (Sanghavi *et al.*, 2013), by the Kinesin-1 heavy chain (Khc) (Brendza *et al.*, 2000; Palacios and St Johnston, 2002; Zimyanin *et al.*, 2008; Williams *et al.*, 2014). Khc is associated with nascent *oskar* RNPs in the nurse cell cytoplasm (Gáspár *et al.*, 2017), raising the question of how the

activity of the motor is coordinated with that of dynein during delivery of *oskar* mRNA from
the nurse cells to the oocyte posterior.

Another double-stranded RNA binding protein (dsRBP), Staufen, is necessary for the Khc-
mediated localization of *oskar* to the posterior pole (Ephrussi *et al.* 1991; St Johnston *et al.*
1991; Januschke *et al.*, 2002; Sanghavi *et al.*, 2013; Jambor *et al.*, 2014; Vazquez-Pianzola
et al., 2017; St Johnston *et al.*, 1992; Zimyanin *et al.*, 2008), as well as for production of
Oskar protein at this site (St Johnston *et al.* 1991; Ephrussi and Lehmann, 1992; Schuldt *et*
al., 1998; Micklem *et al.*, 2000). Staufen is also involved in trafficking of other mRNAs, such
as *bicoid* in the oocyte and *prospero* in dividing neuroblasts (Ferrandon *et al.*, 1994; Schuldt
et al., 1998). Mammalian orthologues of *Drosophila* Staufen, mStau1 and mStau2, play roles
in the bidirectional transport of *CaMKIIα* and *Rgs4* mRNAs in dendrites (Heraud-Farlow *et*
al., 2013; Bauer *et al.*, 2019) and *Xenopus* Staufen is a component of *Vg1* transport RNPs in
the oocyte (Yoon and Mowry, 2004). These observations indicate that Staufen's function as
a regulator of RNP transport is evolutionarily conserved. However, it is unclear how Staufen
proteins orchestrate mRNA trafficking processes.

Here, we show that Staufen antagonizes dynein-based transport of *oskar* RNPs within the
Drosophila oocyte and thus favors kinesin-1-driven translocation of the mRNA to the
posterior pole. Recombinant Staufen can inhibit dynein-based transport of *oskar* mRNPs that
are reconstituted *in vitro* with purified components, showing that it is a direct regulator of the
transport machinery. Staufen displaces Egl from *oskar* RNPs after they have reached the
oocyte. Staufen does not, however, perturb dynein and dynactin association with the RNPs,
revealing that proteins in addition to Egl contribute to motor linkage in this system. These
data suggest that Staufen interferes with dynein activation by promoting dissociation of Egl
from RNPs. We also provide evidence that Staufen's inhibition of dynein is restricted to the
oocyte by Egl- and dynein-mediated concentration of nurse cell-synthesized *stau* mRNA
within the ooplasm. Thus, Egl delivers its own negative regulator into the oocyte, constituting
a feed-forward regulatory mechanism that spatially and temporally controls dynein activity
and thus *oskar* mRNA localization during oogenesis.

RESULTS

Staufen suppresses minus end-directed transport of *oskar* RNPs in ooplasmic
extracts

oskar mRNA is localized at the posterior pole of the oocyte during mid-oogenesis (Ephrussi
et al. 1991; Kim-Ha *et al.* 1991). In *stau* mutant oocytes, enrichment of *oskar* mRNA at the
posterior pole is greatly reduced, and a considerable amount of *oskar* remains at the anterior
margin (Fig 1A'; (Ephrussi *et al.* 1991; St Johnston *et al.* 1991). To assess the effect of *stau*
depletion on the balance of *oskar* directionality on microtubules, we evaluated the movement
of individual MCP-GFP-labeled *oskar-MS2* RNPs in control and *stau* RNAi ooplasmic
extracts by TIRF microscopy (Gaspar and Ephrussi, 2017; Gáspár *et al.*, 2017). This
analysis revealed that the normal ~ 2:1 dominance of microtubule plus end- versus minus

end-directed *oskar* RNP runs (Gaspar and Ephrussi, 2017; Gáspár *et al.*, 2017) was lost,
such that there was no overt bias in the directionality of motility (Fig 1B).
93
While analyzing the motility, we noticed that the relative GFP intensity of the motile fraction
of *oskar* RNPs in the *stau* RNAi extracts was lower than in the control RNAi condition,
indicating fewer mRNA molecules on average. This observation might explain the paucity of
oskar mRNA transport detected in live *stau* mutant oocytes when using the less sensitive
confocal live-cell imaging technique (compare Videos S1-S4; (Zimyanin *et al.*, 2008)).
However, loss of Staufen does not seem to affect the overall RNA content of *oskar* RNPs in
the oocyte as revealed by single-molecule fluorescent *in situ* hybridization (smFISH) (Fig
S5B), indicating a selective effect on the size of motile RNPs.
102
Since differences in cargo size and/or composition could confound the analysis, we
compared motility of *oskar* RNPs of similar sizes by stratifying the GFP intensity of RNPs
into 'relative' RNA units (see Methods and Fig 1C). This approach confirmed that motile
oskar RNPs contained fewer RNA molecules on average in the *stau* RNAi condition (Fig 1C)
and additionally showed that the loss of plus-end dominance could be observed for motile
RNPs with 1 or 2 relative units of *oskar* RNA, which make up the majority of the motile
population (Fig 1B). These data show that Staufen is needed for the plus-end dominance
that localizes *oskar* to the posterior pole.
111
The distribution of *oskar* mRNA and the directionality of RNP transport in *stau* mutants differ
from what is observed in other mutants affecting *oskar* localization: in *Tm1^gs* (Fig 1A" and
(Gáspár *et al.*, 2017) and *Khc^null* (Brendza *et al.*, 2000; Zimyanin *et al.*, 2008; Williams *et al.*,
2014; Gáspár *et al.*, 2017) mutants, *oskar* mRNA does not localize at the posterior of the
oocyte, and minus-end-directed transport predominates as a failure of Khc mediated
posterior-ward transport (Gáspár *et al.*, 2017). To explore the basis of the directionality
defect in *stau* mutants, we quantified the colocalization in egg chambers of functional,
endogenously tagged fluorescent Khc molecules and *oskar* mRNA detected by smFISH.
This method was previously used to show that Tm1 mutations reduce the association of
oskar RNPs with the motor (Gáspár *et al.*, 2017). When Staufen was depleted, there was no
reduction in Khc colocalization with *oskar* mRNA in either the nurse cells or the oocyte (Fig
1D), indicating that the observed loss of plus-end-directed bias (Fig 1B) is not due to
reduced association of Khc with *oskar* RNPs.
125
To assess whether there is a change in motor activity when Staufen is disrupted, we
analyzed the speed and run lengths of GFP-labeled *oskar-MS2* RNPs towards both
microtubule ends in *stau* RNAi and control ooplasmic extracts. The velocity of *oskar* RNP
movement appears to be influenced by their RNA content, as larger RNPs moved more
slowly in either direction (Fig 1E). When Staufen was depleted, there was a substantial and
statistically significant increase in minus-end-directed run lengths and velocities of RNPs
containing 1 or 2 relative units of *oskar* RNA (Fig 1D). In contrast, features of plus-end-
directed motion were generally not affected in the *stau* RNAi condition with the exception of
a modest, but significant, increase in plus end velocity for RNPs with 2 *oskar* RNA units (Fig
1E). These data indicate that the loss of plus-end-directed dominance of *oskar* RNPs when
Staufen is depleted is predominantly associated with hyperactivity of the minus end-directed
dynein transport machinery. Previous work with gain-of-function mutations in the dynein
activator BicD showed that increased dynein activity leads to ectopic anterior accumulation

of *oskar* RNA (Navarro *et al.*, 2004; Liu *et al.*, 2013), as is observed in *stau* mutant oocytes
(Fig 1A',(St Johnston *et al.*, 1991)). Altogether, these findings point to a role of Staufen in
suppression of minus end-directed transport of *oskar* RNA by dynein.
142
143
144   **Staufen antagonizes dynein-mediated *oskar* mRNA transport by purified components**
145
146   To test if Staufen has a direct inhibitory effect on minus-end directed *oskar* RNP transport,
147   we made use of an *in vitro* motility assay that involves reconstitution of a minimal RNP
148   assembled from purified Egl, BicD, dynactin, dynein and *in vitro* transcribed RNA
149   (McClintock *et al.*, 2018). Motility of this RNP along immobilized, fluorescent microtubules
150   can be visualized by TIRF microscopy using fluorophores on the mRNA and dynein.
151
In these experiments we used a ~529 nt fragment of the *oskar* 3'UTR that is sufficient for
nurse cell-to-oocyte transport (Jambor *et al.*, 2014). In the presence of Egl, BicD and
dynactin, *oskar* mRNA frequently underwent long-distance transport in association with
dynein (Fig. 2A and Fig S2A). No transport of *oskar* was detected when Egl and BicD were
omitted from the mix (Fig. S2A); in this condition, association of *oskar* mRNA with
microtubules or processive movement of microtubule-bound dynein complexes was almost
never observed (Fig. S2A-C). These observations corroborate the conclusion that Egl and
BicD link mRNAs to dynein-dynactin and activate movement of the motor along microtubules
(Dienstbier *et al.*, 2009; McClintock *et al.*, 2018; Sladewski *et al.*, 2018).
161
Addition of recombinant Staufen to the Egl, BicD, dynein and dynactin assembly mix
significantly reduced the number of *oskar* mRNA transport events (Fig. 2A and B). This
effect was associated with fewer mRNAs being recruited to the microtubules (Fig. 2C).
However, those mRNAs that interacted with the microtubule were just as likely to undergo
long-distance transport as those incubated without Staufen (Fig. 2D). The presence of
Staufen also reduced the number of processive dynein complexes and the proportion of
microtubule-associated dyneins that underwent processive movement (Fig. 2E-G). We
conclude from these experiments that Staufen directly inhibits transport of *oskar* by Egl,
BicD, dynactin and dynein and that this effect is associated with impaired activation of motor
movement.
172
Dynein movement during mRNA transport is switched on by the N-terminal coiled-coil region
of BicD, which recruits dynactin (and thereby its activating functions) to the motor complex
(Dienstbier *et al.*, 2009; McKenney *et al.*, 2010, 2014; Schlager *et al.*, 2014; Hoogenraad
and Akhmanova, 2016). These events are triggered by docking of RNA-bound Egl to the C-
terminal region of BicD, which frees the N-terminal region from an autoinhibitory interaction
(McClintock *et al.*, 2018; Sladewski *et al.*, 2018). To better resolve how dynein activity is
impaired by Staufen, we assessed its effect on dynein activation by the N-terminal coiled-coil
of the mouse BicD orthologue BicD2 (BicD2N), which is constitutively active in the absence
of cargo (Hoogenraad *et al.*, 2003; Dienstbier *et al.*, 2009; McKenney *et al.*, 2010, 2014;
Schlager *et al.*, 2014; Hoogenraad and Akhmanova, 2016). In the presence of dynactin and
BicD2N, Staufen partially reduced the number of processive dynein movements and the
proportion of processive dyneins (Fig. 2H, I and Fig. S3). However, Staufen's inhibitory effect
on the motor was proportionally much stronger in the presence of *oskar*, Egl, full-length BicD
and dynactin (Fig. 2H, I and Fig S3). These data suggest that Staufen inhibits both the

activation of dynein-dynactin motility by BicD proteins, as well as stimulation of this event by
Egl and RNA.
189
190
191 **A balance of Staufen and Egl is required for *oskar* mRNA localization *in vivo***
192
We next explored the interplay between Egl and Staufen function *in vivo* by manipulating the
dosage of these proteins and determining the effects on *oskar* mRNA localization in mid-
oogenesis. We evaluated mRNA distributions qualitatively (Fig 3) and confirmed our
conclusions through an established, unbiased statistical averaging method (Ghosh *et al.*,
2012; Gaspar *et al.*, 2014) (Fig S4). Similar to the situation in *stau* mutant egg-chambers
(Fig 3A, B and Figs S1A, B, E, S4A and B), UAS/Gal4-mediated overexpression of Egl in the
germline resulted in a strong *oskar* signal at the anterior of the oocyte cortex, and a weaker
enrichment of the mRNA at the posterior (Fig 3C and S4C). This finding corroborates recent
observations in Egl-overexpressing egg-chambers by Mohr *et al.* (Mohr *et al.*, 2021).
202
Altogether, these observations suggest that Staufen and Egl antagonize one another during
oskar mRNA localization in mid-oogenesis. To further test this idea, we overexpressed both
GFP-Staufen (using the maternal tubulin promoter (Micklem *et al.* 2000)) and Egl in the
germline and examined *oskar* localization. While GFP-Staufen overexpression in this
manner had no major effect on the localization of *oskar* (Fig S1F-I and (Heber *et al.*, 2019)),
it suppressed the ectopic anterior localization of *oskar* mRNA caused by elevated Egl levels
(Fig 3C, D and Fig S4C, D). This observation is consistent with mutual antagonism between
Egl and Staufen. Further supporting this notion, removal of a functional copy of *egl* partially
suppressed *oskar* mislocalization at the anterior of egg-chambers with reduced Staufen
levels (*stau* RNAi, *egl[1]/+*; Fig S3C-E). Furthermore, overexpression of Staufen suppressed
the ectopic anterior localization of *oskar* mRNA caused by two different dominant *BicD*
alleles (Fig 3E, F and Fig S4E-H). At least one of these mutations (*BicD[1]*) appears to
suppress the autoinhibited state of BicD in which the dynein and Egl binding sites are folded
back on one another (Liu *et al.*, 2013). Antagonism between BicD and Staufen is further
supported by the observation that reduced *stau* gene dosage enhances embryonic lethality
associated with the dominant *BicD* alleles (Navarro *et al.*, 2004). Collectively, these genetic
interaction experiments indicate a mutually antagonistic relationship between Staufen and
Egl, as well as the latter protein's binding partner, BicD, during the localization of *oskar*
mRNA in mid-oogenesis.
222
223 **Staufen dissociates Egl from *oskar* RNPs in the oocyte**
224
225 To shed light on how Staufen and Egl influence each other *in vivo*, we examined the
226 association of each protein with *oskar* mRNA using smFISH in combination with
227 transgenically-expressed, fluorescently-tagged proteins (Gáspár *et al.*, 2017; Heber *et al.*,
228 2019).
229
Distribution of transgenic Staufen-GFP, expressed by the *stau* promoter at close to
endogenous levels (Fig S1I) resembles that of the endogenous protein (St Johnston *et al.*
1991), and this molecule is functional during *oskar* localization (Fig S1D, E and G). In the
oocyte, Staufen-GFP and *oskar* mRNA frequently overlapped. In the nurse cells, there was
little overlap between Staufen-GFP and *oskar*, as previously reported (Little *et al.*, 2015),
and only low levels of Staufen-GFP signal were observed, regardless of the developmental

stage of the egg-chamber (Fig 4A, A' and B). In the oocyte, the fraction of Staufen-
associated *oskar* RNPs increased as a function of their *oskar* mRNA content (which was
determined by calibration of smFISH signals (see Materials and Methods and (Little *et al.*,
2015)); Fig S4A). This increase was substantially more pronounced during stage 9, when
there is a strong enrichment of *oskar* at the oocyte posterior, than in earlier stages of
oogenesis (Fig 4B). During stage 9, not only the likelihood of Staufen association with *oskar*
RNPs increased but also the relative levels of Staufen per RNP (Fig 4B'). We previously
observed the same behavior for overexpressed GFP-Staufen and mammalian GFP-Stau2
(Heber *et al.*, 2019). Notably, however, no or greatly reduced scaling of the Staufen signal
with *oskar* mRNA content of the RNPs was observed in the oocyte prior to the onset of *oskar*
localization to the posterior (Fig 4B'). This is likely due to limited availability of the protein, as
suggested by the relatively low fluorescent signal in early stage Staufen-GFP oocytes (Fig
S5G). Collectively, our analysis shows that Staufen preferentially associates with *oskar* RNA
in the oocyte and that this association is largely governed by the RNA content of the RNPs
and the enrichment of the pool of Staufen protein in the ooplasm.
251
During the same stages of oogenesis, Egl-GFP (which was mildly overexpressed with the α-
tubulin84B promoter (Fig S5I, (Dienstbier *et al.*, 2009)) was abundant in both the nurse cells
and the oocyte (Fig 4C and C'). This pattern matches the distribution of endogenous Egl
protein, as revealed by immunostaining (Mach and Lehmann, 1997). Colocalization analysis
showed that, in contrast to Staufen-GFP, Egl-GFP was already associated with *oskar* RNPs
in the nurse cells (Fig 4D) and maintained its association with *oskar* in the oocyte until the
onset of posterior localization. However, from stage 9 onward, we detected no significant
colocalization between Egl and RNPs containing four or more copies of *oskar* mRNA (Fig
4D). The amount of Egl signal associated with the RNPs increased proportionately with
oskar mRNA content during early oogenesis but reduced by about two-fold and became
independent of *oskar* mRNA copy number during stage 9 (Fig S5D'). As this loss of Egl from
oskar RNPs was an unexpected finding, we wondered if it was caused by the dilution of Egl-
GFP by endogenous, untagged Egl. However, this was not the case as the reduction of Egl
association with *oskar* RNPs was robust to changes in the dosage of the Egl-GFP transgene
or endogenous unlabeled Egl and was also observed when no unlabeled Egl (*egl^1/egl^2*
rescued by the Egl-GFP transgene) was present in the oocyte (Fig S5F'). Collectively, these
results indicate that whereas Staufen levels increase on *oskar* RNPs as posterior-ward
movement in the oocyte is initiated, there is a concomitant decrease in Egl levels on these
complexes.
271
The opposing trends of Staufen and Egl association with *oskar* RNPs in the nurse cells and
the oocyte (Fig 4B,B' and D,D'), together with the increasing concentrations of Staufen in the
ooplasm prior to the onset of *oskar* localization (Fig S5G), are consistent with Staufen
inhibiting Egl association with *oskar* RNPs in the oocyte. To test this notion, we analyzed the
association of Egl-GFP with *oskar* RNPs in *stau* mutants. When Staufen protein was absent
or greatly reduced, Egl remained associated with *oskar* RNPs in stage 9 oocytes, and the
relative amount of Egl signal that colocalized with the RNPs increased with *oskar* mRNA
content (Fig 4E, E' and Fig S5 E, E'). Conversely, when Staufen was mildly overexpressed
using an RFP-Staufen transgene (Zimyanin *et al.*, 2008), the amount of Egl associated with
oskar RNPs in the oocyte (in particular with those containing 4+ copies of RNA) decreased
(Fig 4F, F').
283

To test biochemically whether Staufen impairs the association of Egl with *oskar* RNPs *in*
vivo, we performed UV crosslinking and GFP immunoprecipitation followed by quantitative
RT-PCR on egg-chambers co-expressing Egl-GFP and either *stau* RNAi or a control RNAi in
the germline (Fig 4G, Fig S5H). Substantially more *oskar* mRNA was co-immunoprecipitated
with Egl-GFP from extracts of egg-chambers expressing *stau* RNAi compared to the control
(Fig 4G). We conclude from these experiments that Staufen antagonizes the association of
Egl with *oskar* mRNA *in vivo*.
291
292
293   **Aberrant anterior localization of *oskar* mRNA correlates with increased Egl**
294   **association**
295
As described above, BicD is a direct binding partner of Egl, and contacts the dynein and
dynactin complexes (Dienstbier *et al.*, 2009; Liu *et al.*, 2013; Vazquez-Pianzola *et al.*, 2017;
McClintock *et al.*, 2018; Sladewski *et al.*, 2018). To gain further insight into how the
association of Egl influences trafficking of *oskar* mRNA, we examined the effect of the two
gain-of-function *BicD* mutations in more detail. Both alleles caused an increase in the
association of Egl-GFP with mid- to large-sized (4-16 RNA copies) *oskar* RNPs in stage 9
oocytes when compared to the wild-type control (Fig 4E, E' and Fig S5F, F'). Although the
magnitude of the effect differed between the two *BicD* alleles, in both cases the amount of
Egl signal on the RNPs scaled with *oskar* RNA copy number up to 32 copies, in contrast with
the low amount of Egl detected on *oskar* RNPs in wild-type stage 9 egg-chambers (Fig 4E'
and Fig S5F'). Thus, there is a correlation between the ability of the BicD gain-of-function
alleles to misdirect *oskar* RNPs to the anterior of the oocyte and their stimulation of Egl
association with these structures in the oocyte cytoplasm.
309
We reasoned that the anterior localization of *oskar* mRNA in the oocyte observed when
untagged Egl was strongly overexpressed (Fig 3C and S4C) might be due to an aberrant,
concentration-dependent retention of the protein on the RNPs. To test this idea, we
introduced the moderately overexpressed Egl-GFP transgene (Dienstbier *et al.*, 2009) into
egg-chambers overexpressing untagged Egl under UAS-Gal4 control (Fig. S5I). We
observed numerous large RNP granules containing high amounts of Egl-GFP (which
presumably also contain unlabelled Egl) and *oskar* mRNA (Fig 5A and A') in the nurse cells
and at the anterior cortex of the oocyte. Such RNPs were seldom detected in the absence of
UAS/Gal4-mediated Egl overexpression (Fig 5A and E). Quantitative analysis revealed that
the relative Egl-GFP signal scaled with *oskar* RNA content in the RNPs when untagged Egl
was overexpressed (Fig 5C'). The effect was similar to that observed in the absence of
Staufen, as well as in egg-chambers with *BicD* gain-of-function alleles (Fig 4E' and S4F'). In
contrast, we found no significant change in Staufen association with *oskar* RNPs in stage 9
oocytes strongly overexpressing Egl (Fig 5D and D'). This finding indicates that Egl does not
overtly interfere with the loading of Staufen on *oskar* RNPs. However, substantially more
high copy number *oskar* RNPs (containing 17+ copies of *oskar*) were localized at the
anterior pole when Egl was strongly overexpressed than in the wild type (Fig 5E), consistent
with our earlier finding that Egl antagonizes Staufen's function in promoting posterior
localization of *oskar* mRNA.
329
Taken together, these data reveal that a shared feature of *oskar* mislocalization to the
anterior in the absence of Staufen, upon Egl overexpression, and in gain-of-function *BicD*

mutants is the failure to release Egl from the RNPs during mid-oogenesis. Our data indicate
that this release is mediated by Staufen, which interferes with the association of Egl with
oskar RNPs, especially those with high *oskar* mRNA content.

Dissociation of Egl does not prevent association of dynein with *oskar* RNPs

As described above, Egl functions as an adaptor molecule linking mRNA localization signals
to dynein-dynactin (Navarro *et al.*, 2004; Dienstbier *et al.*, 2009; Amrute-Nayak and Bullock,
2012). However, it has been proposed that Egl is not the only means to link the motor
complex to RNPs. In *ex vivo* experiments, both the *K10* and *hairy* mRNAs recruit some
dynein-dynactin complexes through Egl-BicD, and others through an Egl-BicD independent
mechanism, which presumably involved other RBP linkers or direct interactions between the
RNA and the motor complex (Bullock *et al.*, 2006; Amrute-Nayak and Bullock, 2012; Dix *et*
al., 2013; Soundararajan and Bullock, 2014). Only those dynein-dynactin complexes
interacting with Egl-BicD were capable of long-distance transport, explaining how these two
adaptors promote minus-end-directed trafficking of mRNAs. How Egl influences dynein's
association with RNPs has not, however, been examined *in vivo*.

To address this question, we quantified the colocalization of *oskar* mRNA (detected by
smFISH) with fluorescently tagged versions of p50/Dynamitin, a dynactin subunit, BicD and
Dynein heavy chain (Dhc) in the nurse cells and oocyte. We saw little difference in
association of each protein with *oskar* mRNA between the stage 9 nurse cells and oocyte
(Fig. 6A-D). This contrasts with the situation for Egl, which showed a relative reduction in
association with *oskar* in the stage 9 oocyte (Fig. 4D and D'). The fraction of *oskar* RNPs
associated with BicD and the dynein-dynactin subunits also did not change as a function of
RNA content in the oocyte (Fig 6D). Notably, the relative amount of these proteins
associated with *oskar* RNPs in the oocyte was approximately double that in the nurse cells
(Fig 6E). The amount of BicD and p50 associated with the RNPs in the oocyte was
independent of *oskar* copy number, whereas Dhc association increased as a function of
oskar mRNA content (Fig 6E). This result indicates that Dhc and BicD/p50 are not recruited
to *oskar* RNPs by identical mechanisms.

Increased levels of Egl on *oskar* RNPs in the absence of Staufen (Fig 4E, E' and S5E, E') or
in the presence of excess Egl (Fig 5D, D') could in principle result in increased recruitment of
the dynein motor. However, when we analyzed GFP-Dhc association with *oskar* RNPs, we
found no major difference between *stau* RNAi, Egl-overexpressing, and control oocytes (Fig
6F and F'). These data indicate that Egl is not obligatory for recruitment of dynein-dynactin to
a native cargo *in vivo*. Intriguingly, *stau* RNAi and Egl overexpressing oocytes also did not
display an increase in the association of BicD with *oskar* RNPs (Fig 6G and G'). Thus, there
appears to be an Egl-independent mechanism for recruiting BicD to these structures.

The loss of Egl from larger *oskar* RNPs during stage 9 suggests that dynein – although
present – is not fully active. This notion is consistent with our earlier observation in control
ooplasmic extracts that RNPs with a greater RNA content, and which therefore have less Egl
bound (Fig 1B and 4D and D'), engaged in minus-end-directed motion less frequently than
those with a lower RNA content: there was an approximately 4-fold plus-end dominance
(77.9% vs 22.1% of plus vs minus-end directed runs) for *oskar* RNPs with more than two

relative RNA units, versus an approximately 2-fold plus-end dominance (~65% vs ~35%) for
smaller RNPs (Fig 1B). In summary, in stage 9 oocytes, *oskar* RNPs lacking Egl maintain
their association with the dynein machinery but are defective in minus-end-directed motion.

379
380 **Staufen activity is controlled by Egl-mediated delivery of its mRNA into the oocyte**
381
As shown above, proper *oskar* localization to the posterior pole depends on availability of
Staufen protein in stage 9 oocytes. However, it is unclear how this spatio-temporal restriction
of Staufen is brought about. We noticed that only trace amounts of endogenous Staufen (Fig
7C and D) or Staufen-GFP (Fig S7C) accumulated in the early arrested oocyte primordium in
egl RNAi egg-chambers. This observation raised the possibility that Egl mediates the
enrichment of Staufen in the ooplasm of wild-type egg-chambers. The normal ooplasmic
accumulation of Staufen could in principle result from its transport in association with target
mRNAs from the nurse cells into the oocyte. However, this seems unlikely, as we detected
no association of Staufen-GFP with its main target, *oskar* mRNA, in the nurse cells (Fig 4A-
B'), and both Staufen and Staufen-GFP signal accumulated in the oocyte even in the
absence of *oskar* mRNA (Fig S7A-C). Alternatively, *stau* mRNA could be transported by Egl-
BicD-dynein-dynactin into the oocyte, leading to accumulation of Staufen protein at this site.
Supporting this notion, whereas *stau* mRNA and protein were strongly enriched in oocytes of
early-stage egg chambers expressing control RNAi, we observed little or no such enrichment
in egg-chambers in which *egl* was knocked down by RNAi (Fig 7B and D). Furthermore,
while *stau* mRNA did not display any specific localization pattern in the ooplasm of wild-type
oocytes at mid-oogenesis (Fig 7E), upon Egl overexpression the transcript was enriched at
the anterior pole of the oocyte (Fig 7E'). This observation suggests that, similar to other
oocyte-enriched transcripts, such as *oskar*, *gurken* and *bicoid* (Ephrussi *et al.*, 1991; Berleth
et al., 1988; Neuman-Silberberg and Schüpbach, 1993; Theurkauf *et al.*, 1993; Clark *et al.*
2007), *stau* mRNA is a target of Egl-dependent transport from the nurse cells by dynein.

403
404 Providing further support for this notion, colocalization analysis revealed that *stau* mRNA
405 and Egl-GFP were associated in the germline, and that this association was significantly
406 lower in the oocyte than in the nurse cells (Fig 7F). We also detected Staufen protein in
407 association with *stau* RNPs in stage 9 oocytes (Fig 7G), as we previously observed for *oskar*
408 RNPs (Fig 4).

409
Additional experiments provided evidence that the restriction of Staufen activity to the oocyte
is critical for proper RNA localization and oogenesis. In contrast to the direct expression of
GFP-Staufen from the maternal tubulin promoter, which results in nurse cell accumulation of
the protein only from mid-oogenesis onwards (Fig S7I), strong overexpression of GFP-
Staufen with highly active maternal Gal4 drivers results in high levels and aggregation of the
protein in the nurse cells of early egg-chambers (Fig S7E-H). Ectopic GFP-Staufen
expression in this manner resulted in failure of *oskar,* as well as *bicoid,* mRNA enrichment in
the oocyte (Fig S7E-F''). These egg-chambers also displayed oocyte polarity defects and
fragmentation of the karyosome (Fig S7G and H), which are hallmarks of defects in
localization of mRNA cargoes for dynein (Neuman-Silberberg and Schüpbach, 1993; Jenny
et al., 2006).
421

Taken together, these data indicate that the Egl-dependent transport of *stau* mRNA into the
oocyte by dynein underlies ooplasmic enrichment of Staufen protein, which in turn
inactivates the machinery for minus end-directed transport of *oskar* RNPs in the stage 9
oocytes. This constitutes a feed-forward mode of regulation, whereby the dynein transporter
system ensures the localized production of its inhibitor, leading to its own inactivation. This
predefined program thereby allows kinesin-1-dependent delivery of *oskar* mRNA to the
oocyte posterior.
429
430
431 **DISCUSSION**
432
The activity of motor molecules is essential for establishing proper localization of mRNA
molecules, which in turn underlies spatiotemporal restriction of protein function (St Johnston,
2005; Martin and Ephrussi, 2009; Mofatteh and Bullock, 2017; Abouward and Schiavo,
2021). The dynein and kinesin microtubule motors play key roles in positioning of mRNAs in
many systems, including by acting sequentially on the same RNP species (Baumann *et al.*,
2012; Gagnon *et al.*, 2013; Mofatteh and Bullock, 2017; Turner-Bridger *et al.*, 2018).
However, it is unclear how the opposing activities of these motors are coordinated during
RNP trafficking.
441
Here we use the tractable *Drosophila* egg-chamber to reveal a mechanism for
spatiotemporal control of dynein and kinesin-1 activity. Central to this system are two
dsRBPs, Staufen and Egl. Several genetic interaction studies have shown previously that
these proteins have opposing activities in the context of *oskar* mRNA localization in the
oocyte and anteroposterior patterning (Mohler and Wieschaus, 1986; Navarro *et al.*, 2004)
and both proteins were shown to form a complex with *oskar* (Laver *et al.*, 2013; Sanghavi *et
al.*, 2016). Recent work by Mohr and colleagues (Mohr *et al.*, 2021), which was published
while this manuscript was in preparation, further corroborated the ability of Egl to antagonize
localization of *oskar* to the posterior of the oocyte. Mohr *et al.* also investigated the potential
interplay of Staufen and Egl *in vitro* with RNA binding assays. It was shown that *in vitro* Egl
binds to a truncated version of stem-loop 2 in the *oskar* 3' UTR (SL2, (Jambor *et al.*, 2014)
that they termed the Transport and Anchoring Signal (TAS) (Mohr *et al.*, 2021). The TAS
partially overlaps with one of the Staufen Recognized Structures (Laver *et al.*, 2013) which
are important for Staufen binding *in vitro* and for proper *oskar* mRNA localization *in vivo*
(Mohr *et al.*, 2021). These data suggest that Staufen could antagonize Egl function by
interfering with binding of the latter protein to *oskar* mRNA. However, Mohr et al. did not
directly test this possibility. We found that knocking down Staufen increases the association
of Egl with the mRNA, demonstrating a role for Staufen in antagonizing association of Egl
with *oskar* RNA. The results of our *in vitro* motility assays are also consistent with this
scenario, as RNA binding to Egl is a prerequisite for full dynein activity within this minimal
RNP (McClintock *et al.*, 2018; Sladewski *et al.*, 2018). It is conceivable that the transport and
anchoring functions ascribed to TAS are two facets of the same underlying molecular
mechanism, as increased dynein activity in the absence of Staufen would drive enrichment
of *oskar* RNPs at microtubule minus ends that are nucleated at the anterior cortex
(Januschke *et al.*, 2006).
467
Our study additionally reveals through smFISH-based colocalization analysis in egg-
chambers how the interactions of Staufen and Egl with *oskar* mRNA are orchestrated in time

and space. When Staufen levels are low, such as in early oogenesis, amounts of Egl per
RNP scale with *oskar* RNA content. During stage 9, this scaling is lost in the oocyte and the
relative amount of Egl on *oskar* RNPs decreases as Staufen is recruited to *oskar*. Our data
indicate that Staufen-mediated displacement of Egl is a critical step in switching to kinesin-1-
based trafficking of *oskar* mRNA to the posterior pole. This switch is also likely to involve the
activities of exon junction complex components, which are required for posterior localization
of *oskar* (Newmark and Boswell, 1994; Hachet and Ephrussi, 2001, 2004; Mohr, Dillon and
Boswell, 2001; van Eeden *et al.*, 2001; Palacios *et al.*, 2004; Zimyanin *et al.*, 2008; Ghosh *et*
al., 2012). Our study provides a framework for understanding how the activities of these
factors are coordinated with that of Staufen.
480
481 Intriguingly, whilst an excess of Staufen can dissociate Egl from *oskar* RNPs, increasing Egl
482 concentration does not overtly affect Staufen recruitment with these structures. This
483 observation might be due to the five-to-one excess of SRSs over the TAS in each *oskar*
484 molecule (Mohr *et al.*, 2021), which could mask the loss of Staufen binding to the TAS-
485 proximal SRS.
486
We also found that dissociation of Egl from *oskar* RNPs in the stage 9 oocyte does not
detectably alter the amount of dynein on these structures. This work lends *in vivo* support to
the notion that additional proteins or RNA sequences can recruit dynein and dynactin in an
inactive state to RNPs (Bullock *et al.*, 2006; Amrute-Nayak and Bullock, 2012; Dix *et al.*,
2013; Soundararajan and Bullock, 2014). Presumably, Egl recruits only a small fraction of
the total number of dynein complexes on *oskar* RNPs or activates the motility of complexes
that are also linked to the RNA by other factors. Unexpectedly, we reveal that the bulk
association of BicD with *oskar* RNPs is also not dependent on Egl, pointing to an additional
mechanism for recruiting BicD, presumably in the autoinhibited state, that prevents dynein
activation (Hoogenraad *et al.*, 2003; Liu *et al.*, 2013). While the relative amount of the dynein
machinery associated with *oskar* RNPs is greater in the oocyte than the nurse cells, dynein's
association with *oskar* RNPs in the oocyte does not scale with RNA content. How the
amount of dynein recruited to these RNPs could be limited is unclear. However, this
mechanism could be a means to prevent sequestration of this multi-functional motor to a
very abundant cargo (~0.5-1 million copies of *oskar* mRNA per oocyte (Little *et al.*, 2015)).
502
Whilst our data build a strong case that a key function of Staufen in *oskar* mRNA localization
is to limit dynein activity by displacing Egl, several lines of evidence suggest that this is not
its only role in this context. We found that Staufen had a partial inhibitory effect on minus-
end-directed motility of purified dynein-dynactin complexes activated in an Egl-independent
manner by a constitutively active truncation of a BicD protein. Understanding the
mechanistic basis of this effect will be the goal of future studies. Moreover, Mohr *et al.* (Mohr
et al., 2021) found that the SRSs that are not proximal to the TAS in the sequence of *oskar*
also contribute to posterior localization of *oskar* mRNA in a Staufen-dependent manner.
Although it is possible that these elements are close enough to the binding site for Egl-BicD-
dynein-dynactin in the folded RNA molecule to interfere directly with the assembly or activity
of the complex, they could also regulate *oskar* mRNA distribution through an independent
mechanism. Other observations in our study hint at other roles of Staufen. We observed that
when Staufen is depleted motile *oskar* RNPs tend to have a lower RNA content, suggesting
an additional function of the protein in RNA oligomerization. Furthermore, whilst the
magnitude of the effect was much smaller than for minus-end-directed motion, there was

increased plus-end-directed velocity of a subset of *oskar* RNPs in *ex vivo* motility assays
when Staufen was disrupted. Although this observation could reflect dynein's ability to
promote kinesin-1 activity (Hancock, 2014), it is also possible that Staufen directly tunes the
activity of the plus end-directed motor.
522
An important question raised by our analysis of Staufen's effects on *oskar* mRNA transport is
how the timing and location of this process is controlled. We provide evidence that this key
aspect of *oskar* mRNA localization is based on another mRNA localization process in which
Egl, as part of s*tau* RNPs, is responsible for the enrichment of *stau* mRNA in the developing
oocyte. We propose this mechanism constitutes a feed-forward type of switch, whereby the
activity of the dynein-mediated transport machinery deploys its own negative regulator to a
distant location. The resultant increase in levels of ooplasmic Staufen levels in the ooplasm
prevents dynein-mediated transport of *oskar* to the anterior of the oocyte, while the almost
complete absence of the protein from the nurse cells is likely to be important for
uninterrupted transport of *oskar* RNPs into the oocyte by dynein during early- and mid-
oogenesis. Presumably, *stau* translation is suppressed during transit into the oocyte or the
protein is translated en route but only builds up to meaningful levels where the RNA is
concentrated in the oocyte. Staufen protein might also modulate its own localization in the
ooplasm by antagonizing the association of *stau* mRNA with Egl, and thereby its minus end-
directed motility. Consistent with this notion, the level of Egl on *stau* RNPs declines at a
similar stage to when Staufen inhibits the association of Egl with *oskar* RNPs. As Staufen
controls localization and translation of other mRNAs in the maturing oocyte (St Johnston *et*
al., 1991; Ferrandon *et al.*, 1994), mRNA-based regulation of the protein distribution in the
egg-chamber is likely to have functions that extend beyond orchestrating *oskar* mRNA
localization. Given the functional conservation of Staufen protein (Heber *et al.*, 2019) and the
observation that the mRNA encoding mStau2 is localized in dendrites of mammalian
neurons (Zappulo *et al.*, 2017), it is plausible that the feed-forward loop established by *stau*
RNA localization during *Drosophila* oogenesis is an evolutionarily conserved process that
controls RNA trafficking and protein expression in polarized cells.
547

METHODS

Fly strains

To knock down *stau* and *egl* RNA levels, we used the P{TRiP.GL01531} (FBal0281234) and the P{TRiP.HM05180} (FBal0240309) transgenic lines. A TRiP line against the *w* gene (P{TRiP.GL00094} - FBal0262579) was used as the negative control. The following mutant lines were used to disrupt or modify gene function: *stau*[D3] (FBal0016165) and *stau*[R9] (FBal0032815) alleles in heterozygous combination; *egl*[1] (FBal0003574) and *egl*[2] (FBal0003575); *BicD*[1] (FBal0001140) and *BicD*[2] (FBal0001141); *osk*[A87] (FBal0141009) in combination with *Df(3R)p-XT103* (FBab0002842) to remove *oskar* mRNA from egg-chambers. *w*[1118] (FBal0018186) was used as the wild-type control.

To overexpress Staufen, we used the αTub67C:GFPm6-Staufen (FBal0091177), αTub67C:RFP-Staufen (Zimyanin *et al.*, 2008) transgenic lines and a P{UASp-Staufen} transgene inserted onto the 3^rd chromosome (kind gift of F. Besse).

To overexpress unlabelled Egl protein, we used the P{UASp-egl.B} transgene (FBtp0022425) inserted on the X or the 3^rd chromosome. To label proteins of interest, we used the following fluorescently tagged reporter lines: P{Staufen-GFP} inserted on the 3^rd chromosome, expressing Staufen-GFP under control of the endogenous *stau* promoter (kind gift of F. Besse); P{tub-egl.EGFP} (FBtp0041299) inserted on the 3^rd chromosome, driving expression of Egl-GFP; P{UASp-BicD.eGFP} (FBtp0069955) and P{UAS-DCTN2-p50::GFP} (FBtp0016940) transgenic insertions to label BicD and Dynamitin, respectively. To drive the expression of the TRiP RNAi lines and other UASp constructs in the female germline, one copy of P{osk-Gal4} (FBtp0083699) inserted onto either the 2^nd or the 3^rd chromosome was used. For moderate overexpression of UASp-GFP-Staufen, we used one copy of P{matα4-GAL-VP16} (FBtp0009293) inserted onto the 2^nd chromosome.

The endogenous Dhc locus (*Dhc64C*) was tagged with the EmeraldGFP coding sequence to generate a GFP-Dhc expressing fly line according to protocols of the flyCRISPR website (Gratz *et al.*, 2013; O'Connor-Giles, Wildonger and Harrison, 2014) (https://flycrispr.org/). The locus was targeted using CRISPR/Cas9 and a guide RNA targeting the following sequence: 5' GAGTCACCCATGTCCCACAA. The introduced double-stranded break was repaired by homologous recombination using an in-frame EmeraldGFP coding sequence flanked by two ~ 700 bp long homology arms targeting around the Dhc translational initiation codon. F1 generation embryos were screened for GFP fluorescence to identify individuals with a modified genome. Flies homozygous for GFP-Dhc are viable and fertile.

All stocks were raised on normal cornmeal agar at 25 °C, and females were fed with wet yeast overnight before harvesting their ovaries.

***Ex vivo* motility assay of *oskar* RNPs**

The *ex vivo* motility analysis of *oskar* RNPs were carried out as described in (Gáspár *et al.*, 2017). Briefly, control and *stau* RNAi ovaries expressing mCherry-alpha-tubulin and *oskMS2*-GFP were dissected and transferred onto silanized coverslips in a drop of BRB80.

Several stage 9 egg-chambers were isolated and pulled under Voltalef 10S halocarbon oil
on the same coverslip. Under the oil, the nurse cells were removed by laceration using two
fine tungsten needles. The isolated oocyte still in a "sack" of follicle cells were spatially
separated from the remnants of nurse cell cytoplasm and with a gentle prick on the oocyte
anterior the ooplasm was released onto the coverslip by a continuous slow pulling on the
posterior of the oocyte-follicle "sack". Once several such preps were created, the ooplasmic
extracts were imaged using a Leica 7000 TIRF microscope with 100x oil (NA=1.4) objective
to visualize *oskMS2*-GFP and mCherry labeled microtubules. Time-lapse series were
collected and analyzed as described in (Gáspár *et al.*, 2017).
605

606   To analyze the relative mRNA content of *oskMS2*-GFP RNPs, a series of Gaussian
607   functions was fitted to the GFP signal intensity distribution in each time-lapse series
608   individually using the mixtools package in R (Benaglia *et al.*, 2009; R Core Team, 2014). The
609   smallest μ value of Gaussian fits was used to represent a single unit of RNA and each RNP
610   was normalized to this value.
611

613   **RNA co-immunoprecipitation (RIP) from ovarian lysate**
614

RIP was carried out as described in (Gáspár *et al.*, 2017). Briefly, ovaries from 50 flies were
dissected in BRB80 (80 mM PIPES, 1 mM $MgCl_2$, 1 mM EGTA, pH 6.8) and lysed in Pre-XL
buffer (20 mM Tris-Cl, 150 mM KCl, 4 mM $MgCl_2$, 2xPIC, 1 mM PMSF, pH 7.5;
supplemented with 40U of RiboLock RNase Inhibitor per 100µL lysate). Ovaries were
ground using a pestle and centrifuged for 1 min at 500 x g. The supernatant was extracted
and crosslinked at $0.3J/cm^2$. The lysate was equalized with 1 volume of Pre-XL buffer, 1
volume of RIPA (10 mM Tris/Cl pH 7.5; 150 mM NaCl; 0,5 mM EDTA; 0,1% SDS; 1% Triton
X-100; 1% Deoxycholate, 0.09% Na-Azide) buffer and 8 volumes of low salt buffer (20 mM
Tris-Cl, 150 mM KCl, 0.5 mM EDTA, 1 mM PMSF). GFP-Trap®_MA beads were washed
with low-salt buffer and blocked for 60 mins at room temperature in Casein Blocking Buffer
(Sigma-Aldrich) supplemented with 50 µg/mL Heparin. Lysate was incubated with beads for
90 minutes at 4°C. The beads were then washed six times with high salt buffer (20 mM Tris-
Cl, 500 mM NaCl, 1 mM EDTA, 0.1% SDS, 0.5 mM DTT, 1xPIC, 1mM PMSF) and twice with
PBT (PBS + 0.1% Triton). Endogenous RNA cross-linked to bait protein was recovered from
the beads using the QuickRNA Microprep Kit (Zymo Research). Complementary DNA was
synthesized using the SuperScript III First-Strand Synthesis kit (Invitrogen) and used as
template for PCR using *oskar* and *bicoid* primers.
632

633
634   **Single molecule fluorescent hybridization (smFISH)**
635

636   Single molecule FISH was carried out as described previously (Gáspár *et al.*, 2017; Gaspar,
637   Wippich and Ephrussi, 2017; Heber *et al.*, 2019). Briefly, ssDNA oligonucleotides
638   complementary to *oskar* and *stau* mRNAs (Table S1) were mixed and labeled with Atto532,
639   Atto565 or Atto633 according to the protocol described in Gaspar et al., 2017 (Gaspar,
640   Wippich and Ephrussi, 2017).
641

Drosophila ovaries expressing fluorescent reporter transgenes were dissected in 2% PFA,
0.05% Triton X-100 in PBS and fixed for 20 min. The fixative was removed by two 5 minute

washes in PBT (PBS □+□ 0.1% Triton X-100, pH 7.4). Ovaries were pre-hybridized in
200 □ µL 2 □× □ HYBEC (300 □ mM NaCl, 30 □ mM sodium citrate pH 7.0, 15% ethylene
carbonate, 1 □ mM EDTA, 50 □ µg per mL heparin, 100 □ µg per mL salmon sperm DNA, 1%
Triton X-100) for 10 □ min at 42 □ °C. Meanwhile, the 50 □ µL probe mixture (5 □ nM per
individual oligonucleotide) was prepared and pre-warmed to hybridization temperature. After
10 minutes of pre-hybridization, the probe mix was mixed into the pre-hybridization mixture.
After 2 □ hours of hybridization at 42 □ °C, the unbound probe molecules were washed out of
the specimen by two washes in pre-warmed HYBEC and a final wash in PBT at room
temperature. To boost signal intensity for the protein enrichment analysis of Staufen-GFP in
early egg-chambers, we incubated the ovaries in GFP-Booster (Chromotek) diluted 1:2000
in PBT for 1 hour, then washed the sample three times for 10 minutes in PBT. Ovaries were
mounted in Vectashield and processed for smFISH analysis.

656
657
658 **Microscopy**
659
660 Imaging of oocytes was carried out on a Leica TCS SP8 confocal laser scanning microscope
661 using a 20x dry (NA □= □ 0.75) objective for imaging the RNA distribution in the oocytes and a
662 63x oil immersion (NA □= □ 1.4) objective for analysis of RNA or Staufen-GFP protein
663 enrichment in early oocytes and for colocalization analysis between RNPs and the
664 fluorescently labeled Staufen and Egl proteins.

665
Imaging of reconstituted RNPs by TIRF microscopy was performed as described previously
(McClintock *et al.*, 2018) using a customized Nikon TIRF system controlled by Micro-
manager (Edelstein *et al.*, 2014), with the exception of data presented in Figures 2H and I,
and Figure S3. For these experiments, movies were acquired with 100-ms exposures at 4
frames/s for a total of 3 min on a Nikon Eclipse Ti2 inverted TIRF system equipped with a
Nikon 100x Apo TIRF oil immersion objective (NA = 1.49) and Photometrics Prime 95B
CMOS camera. TMR-dynein was imaged using the 561 nm laser line with a Chroma ET-561
nm Laser Bandpass Set (ET576lp and ET600/50m) and ET605/52m emission filter.

674
675
676 **RNA distribution analysis**
677
Analysis of *oskar* and *bicoid* mRNA distribution was carried out as described in (Gaspar *et*
al., 2014; Gáspár *et al.*, 2017; Heber *et al.*, 2019). Briefly, we manually defined the outlines
of the oocytes and their anteroposterior (AP) axis, using a custom-made ImageJ plug-in, the
smFISH signal was redistributed into a 100 □× □ 100 matrix. Each column of the matrix
represents the relative amount of signal found under 1% of the AP axis length, with anterior
on the left (column 1) and posterior on the right (column 100). The matrices from different
oocytes of the same genotype and stage were averaged to obtain an average view RNA
localization. The center-of-mass (relative to the geometric center of the oocyte) was
determined and compared statistically using a Kruskal–Wallis test followed by pairwise
Mann–Whitney U test against the control.

688
689
690 **RNA and protein enrichment analysis**
691

The boundaries of the somatic follicle cells, nurse cells and oocyte were manually defined, and the fluorescence intensity of smFISH or GFP signal in these compartments was measured by ImageJ. The extent of enrichment of RNA and protein in the somatic follicle cells and the oocyte was obtained by normalizing the measured fluorescence intensity values to the corresponding values obtained for the nurse cells.

Colocalization analysis between RNPs and fluorescent reporters

Analysis was carried out as described in (Gáspár *et al.*, 2017; Heber *et al.*, 2019). Briefly, images for colocalization analysis were deconvolved using Huygens Essential (https://svi.nl) and segmented using a custom ImageJ (Schneider, Rasband and Eliceiri, 2012) plug-in. Nearest-neighbor pairs between RNPs and fluorescent reporters were identified, their position and signal intensities were extracted from the nurse cell and oocyte compartments, excluding any nuclear areas in the field of view. Quantification of mRNA copy number per RNP and normalization of fluorescent reporter signal was carried out by fitting multiple Gaussian functions to the corresponding signal intensity distributions taken from the nurse cells using the mixtools package in R (Benaglia *et al.*, 2009; R Core Team, 2014). The μ value of Gaussian fit that described the largest portion of the distribution in the nurse cells (almost the lowest value of all fitted μ values) was taken as the signal intensity of one unit (for RNPs the intensity of a signal mRNA molecule). These unit values were used to normalize raw signal intensities. RNPs were clustered based on this normalized intensity under the following rule: [$2^i:2^{i+1}$), i $\in$ [0:8], i.e., 1, 2:3, 4:7, 8:15, etc. The observed nearest-neighbor colocalization frequencies were computed for each cluster and were compared to the expected colocalization frequencies (governed by the object-densities, determined in randomized object localizations). Similarly, the mean, normalized intensity of colocalizing fluorescent reporter molecules was calculated for each cluster. Correlation between RNA content and the normalized mean fluorescent reporter intensity was tested and compared using least-squares means analysis in R (Lenth, 2016). We typically analyzed 3000-80000 *oskar* RNPs imaged in 4-6 egg-chambers collected from 3-4 females per condition.

Statistical analysis

Statistical analyses were performed as indicated in the figure legends using R (R Core Team, 2014), RStudio (www.rstudio.com), and GraphPad Prism version 9.1.1 for MacOS (GraphPad Software, San Diego, California USA, www.graphpad.com). All graphs were plotted by the ggplot2 library in R (Wickham, 2016) and GraphPad Prism version 9.1.1 for MacOS (GraphPad Software, San Diego, California USA, www.graphpad.com).

Protein expression, purification, and fluorescent labeling

Recombinant dynein (human), Egl (*Drosophila*), BicD (*Drosophila*), and BicD2N (mouse) were expressed, purified, and fluorescently labeled as described previously (Hoang *et al.*, 2017; McClintock *et al.*, 2018). Briefly, complexes of the complete dynein holoenzyme,the coexpressed Egl/BicD complex, and BicD2N were expressed in *Sf*9 insect cells and purified by affinity chromatography using ZZ-tags on the Dynein heavy chain, Egl, or BicD2N,

respectively. Fluorescent labeling of the Dynein heavy chain and BicD2N with SNAP-Cell
TMR Star or SNAP-Surface Alexa Fluor 647 via N-terminal SNAPf tags was performed
either on-column prior to elution from the IgG affinity resin by cleavage of the ZZ epitope by
TEV protease (dynein) or in solution following TEV-cleavage (BicD2N). All complexes were
further purified by FPLC-based gel-filtration chromatography, using either TSKgel
G4000SWxl (dynein) or Superose 6 Increase 3.2/300 GL (Egl/BicD) columns. The dynactin
complex (porcine) was purified natively from pig brains as described previously (Schlager *et*
al., 2014; Urnavicius *et al.*, 2015). Extracts were clarified and fractionated by FPLC using SP
Sepharose (cation exchange, ~250ml bed volume), Mono Q 4.6/100 PE (anion exchange),
and TSKgel G4000SW (gel-filtration) columns.
750
Recombinant Staufen (*Drosophila*) was cloned as fusion protein with an N-terminal His_6-
rsEGFP2-tag and a C-terminal His_6-tag in pFastBacDual between the KpnI and HindIII sites
and expressed in *Sf*21 insect cells using the Bac-to-Bac expression system (Thermo Fisher).
The protein was purified by affinity chromatography on HisTrap FF in 1x PBS + 880 mM
NaCl, 400 mM arginine, 10 mM - 200 mM imidazole, 2 mM DTT and HiTrap Heparin HP in
40 mM Bis-Tris pH 7.5, 150 mM NaCl – 1000 mM NaCl, 40 mM arginine, 2 mM DTT and a
final gel-filtration on Superdex200 Increase in a final buffer of 40 mM Bis-Tris pH 7.5, 150
mM NaCl, 2 mM DTT.
759
760 In all cases, eluted protein was dispensed into single-use aliquots, flash-frozen in liquid
761 nitrogen, and stored at -80°C.
762
763
764 **RNA synthesis and purification**
765
The *oskar* RNA used for the reconstitution assay was uncapped and synthesized *in vitro*
using the MEGAscript T7 Transcription Kit (Ambion). The RNA was transcribed from a
linearized plasmid DNA template containing an ~529-nt region of the *osk* 3'UTR previously
defined as region 2+3 that includes the oocyte entry signal (OES) and promotes the
localization of reporter transcripts to the developing oocyte (Jambor *et al.*, 2014). For
fluorescent labeling of transcripts, Cy5 UTP or ChromaTide Alexa Fluor 488 UTP was
included in the transcription reaction with a 2-fold or 4-fold excess of unlabelled UTP,
yielding transcripts with an average of ~5 or ~14 fluorophores per RNA molecule,
respectively. Excess fluorescent UTP was removed using 2 successive rounds of Sephadex
G-50 desalting spin columns per transcription reaction. Transcripts were subsequently
purified by ammonium acetate/ethanol precipitation and resuspension in nuclease-free
water. RNA was dispensed into single-use aliquots and stored at -80°C.
778
779
780 *In vitro* **RNP reconstitution assays**
781
TIRF-based single-molecule reconstitution assays were performed as described previously
(McClintock *et al.*, 2018). Briefly, taxol-stabilized porcine microtubules were polymerized with
a mixture of unlabelled tubulin, HiLyte 488 tubulin and biotin-conjugated tubulin, and
immobilized in imaging flow chambers by streptavidin-based linkages to biotin-PEG
passivated cover slips. Assembly mixes containing relevant combinations of dynein,
dynactin, Egl, BicD, BicD2N, *oskar* RNA, and Staufen were then diluted to concentrations

suitable for imaging of single molecules (~2.5 nM for dynein), applied to the flow chamber and subsequently imaged by TIRF microscopy. For assays testing the effect of Staufen on RNA or dynein motility, the complexes were assembled for 45-60 minutes on ice with components diluted in GF150 buffer (25 mM HEPES pH 7.3, 150 mM KCl, 1 mM $MgCl_2$, 5 mM DTT, 0.1 mM MgATP, 10% glycerol) to concentrations of 100 nM TMR-dynein, 200 nM dynactin, 500 nM Egl/BicD or BicD2N, and 1 µM *oskar* RNA. This assembly mix was then diluted 1:1 with either ~4 µM rsEGFP2- Staufen or Staufen storage buffer (40 mM Bis-Tris pH 7.5, 150 mM NaCl, 2 mM DTT) and incubated for at least a further 15 minutes on ice. Just prior application to the imaging chamber, this mixture was further diluted 20-fold in Motility Buffer (30 mM HEPES pH 7.3, 50 mM KCl, 5 mM $MgSO_4$, 1 mM EGTA pH 7.3, 1 mM DTT, 0.5 mg ml^{-1} BSA, 1 mg ml^{-1} a-casein, 20 µM taxol) with added MgATP (2.5 mM final concentration) and 1x oxygen scavenging system (1.25 µM glucose oxidase, 140 nM catalase, 71 mM 2-mercaptoethanol, 25 mM glucose final concentrations) to reduce photobleaching. For assays testing the effect of Egl/BicD on RNA binding and motility on microtubules, the complexes were assayed as above except that primary assembly mixes consisted of 100 nM Alexa Fluor 647-dynein, 200 nM dynactin, 500 nM Egl/BicD or Egl/BicD storage buffer (GF150), and 1 µM *oskar* RNA, which was subsequently diluted 1:1 in Staufen storage buffer and processed as above. Binding and motility of RNA and dynein in the in vitro reconstitution assays were manually analyzed by kymograph in FIJI (Schindelin *et al.*, 2012) as described previously (McClintock *et al.*, 2018). Briefly, microtubule binding events were defined as those lasting for a minimum of 3 continuous frames of acquisition and processive events were defined as those that exhibited unidirectional movement of at least 5 pixels (525-550 nm depending on the microscope system) during this time, regardless of velocity.

ACKNOWLEDGEMENTS

We thank the EMBL Advanced Light Microscopy and Gene Core Facilities for their support. We acknowledge Alessandra Reversi (EMBL *Drosophila* Injection Service) for fly transgenesis and thank Florence Besse for *Drosophila* strains and Dierk Niessing for reagents for recombinant Staufen expression. Work in S.L.B.'s group is supported by the Medical Research Council, as part of United Kingdom Research and Innovation (also known as UK Research and Innovation) [MRC file reference number MC_U105178790]. For the purpose of the MRC open access policy, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising. M.A.M. is supported by a BBSRC project grant (BB/T00696X/1). L.-J.P. is supported by a FOR2333 grant of the Deutsche Forschungsgemeinschaft (Germany) to A.E. A.E. gratefully acknowledges the support of the European Molecular Biology Laboratory.

AUTHOR CONTRIBUTIONS

I.G. and M.A.M. designed experiments. I.G., L.J.P., M.A.M. and S.H. carried out the experiments and analyzed the data. S.L.B. and A.E. supervised the work. All authors discussed the data and contributed to manuscript preparation.

COMPETING INTERESTS STATEMENT

The authors declare that there are no competing interests.

 FIGURES

841
Figure 1: ***oskar* RNA localization and transport are impaired in *stau* mutants.**

(A-A") Endogenous *oskar* mRNA localization (shown in blue (low intensity)/yellow (high intensity)) in wild-type
(A), *stau* null (A') and *Tm1-I/C* null stage 9 oocytes (A") (anterior to the left and posterior to the right). Scale
bar represents 20 μm. In *stau* null oocytes, there is a pronounced anterior accumulation of *oskar* in addition to
the normal posterior localization; in Tm1-I/C mutant oocytes, *oskar* RNA is predominantly enriched at the
anterior cortex.

(B) Polarity distribution of *oskar*-MS2 RNP runs in control RNAi and *stau* RNAi ooplasmic extracts. Numbers
indicate the number of *oskar* RNP runs measured for all motile *oskar* RNPs and for RNPs categorized by
relative RNA content (Video S3 and S4).

(C) Relative mRNA content of motile *oskar-MS2* RNPs in control RNAi and *stau* RNAi ooplasmic extracts.
Pink, green and blue indicate the fraction of *oskar* RNPs with one, two, or more relative RNA units,
respectively (see Methods). Numbers show the number of *oskar* RNPs in that category.

(D) Association of Khc and *oskar* RNPs *in situ* in the nurse cells and the oocyte of control RNAi versus *stau*
RNAi samples.

(E-F) Speed (E) and run length (F) of motile *oskar-MS2* RNPs towards the plus or minus ends of microtubules
in control RNAi and *stau* RNAi ooplasmic extracts. RNPs are stratified by relative RNA content (D). Numbers
above the boxplots (E and F) indicate the differences in mean travel distance between *stau* and control RNAi
extracts. In *stau* RNAi extracts there were too few motile *oskar-MS2* RNPs with >2 RNAs to establish a
statistical trend (C,E-F). P-values of Fisher exact (B) or pairwise Mann-Whitney U-tests (D-F) are shown.

Figure 2. Staufen impairs *oskar* mRNA transport and dynein activation *in vitro*.

(A) Kymographs (time-distance plots) exemplifying the behavior of *oskar* RNA and dynein in the presence (filled circle) and absence (open circle) of Staufen; in both conditions, Egl, BicD and dynactin are also present but not fluorescently labeled. Minus end is to the left and plus end to the right.

(B-G) Quantification of motile properties of *oskar* RNA (B-D) and dynein (E-G) in the conditions shown in A. Charts show frequency of processive movements (B, E), total number of microtubule (MT) binding events (C, F) and fraction of microtubule-binding events that result in processive motility (D, G).

(H-I) Quantification of effect of Staufen on number of processive dynein complexes and fraction of dynein-microtubule binding events that result in processive motility for motor activated by dynactin, *oskar* RNA, and Egl/BicD (H) vs motor activated by dynactin and BicD2N (I). In H and I, values for conditions with Staufen were normalized to the corresponding condition without Staufen to obtain relative metrics. Plots show the mean ± standard deviation (SD) of values from individual microtubules (represented by black circles) derived from analysis of 586-1341 single RNA particles or 1247-2207 single dynein particles per condition. Statistical significance and P-values were determined with Mann-Whitney tests (B-G) or with Brown-Forsythe and Welch ANOVA tests (H and I).

879
Figure 3: Mutual interference of Staufen and Egl during *oskar* mRNA localization.
(A-F) Representative micrographs of the distribution of endogenous *oskar* mRNA (detected by smFISH; blue -
low intensity, yellow - high intensity, see scale in F) in stage 9 oocytes (anterior to the left, posterior to the right).
Ectopic anterior accumulation of *oskar* mRNA in oocytes lacking Staufen protein (B; see also Fig 1A'),
overexpressing Egl (C) or heterozygous for one copy of the dominant, hyperactive *BicD^1* allele (E). Anterior
accumulation of *oskar* is not observed in wild-type oocytes (A), or upon Staufen overexpression in Egl-
overexpressing oocytes (D) or *BicD^1/+* oocytes (F). Scale bar represents 20 μm.

887
Figure 4: Staufen interferes with association of Egl with *oskar* RNPs in the oocyte.

(A-A') Staufen-GFP (green) distribution in the nurse cell cytoplasm (left-hand images) and oocyte (right-hand
images) before (stage 6-7, A) and during (stage 8-9, A') the posterior localization of endogenous *oskar* mRNA
(magenta) in the oocyte of the same egg-chamber.

(B-B') Quantification of Staufen-GFP association with *oskar* RNPs in nurse cell cytoplasm and ooplasm at the
indicated stages as a function of RNA copy number.

(C-C') Distribution of Egl-GFP (green) in the nurse cell cytoplasm (left-hand images) and oocyte (right-hand
images) before (stage 7, C) and during (stage 9, C') the posterior localization of endogenous *oskar* mRNA
(magenta) in the oocyte of the same egg-chamber.

In A, A', C and C', nurse cell images are shown with different brightness/contrast settings for better
visualization of the signals. Cyan arrows point to examples of colocalization. Scale bars represent 10 μm.

(D-D') Quantification of association of Egl-GFP with *oskar* RNPs in nurse cell cytoplasm and ooplasm at the
indicated stages.

(E-F') Association of Egl-GFP with *oskar* RNPs in the stage 9 oocyte in the indicated genotypes. In A-B'
endogenous Staufen was absent (*stau^{R9}/stau^{D3}* background), whereas in C-F' endogenous, unlabeled Egl
was present.

Error bars represent 95% confidence intervals in B, B', D-F'. The size of the circles is proportional to the
relative abundance of each category of *oskar* RNPs within the overall population. Triangles indicate that the
fraction of GFP-positive *oskar* RNPs is not significantly different from zero ($p>0.01$, B, B', D-F'). In B', D', E'
and F', GFP protein intensities on *oskar* RNPs in the oocytes are normalized to GFP signal intensities in the
corresponding nurse cells

(G) Fold-enrichment of *oskar* mRNA precipitated from UV crosslinked Egl-GFP ovarian extracts relative to the
GFP control in control and *stau* RNAi. Different colors represent different experiments. Solid ($p<0.05$) or
dashed lines ($p>0.05$) connect paired data (pairwise Student's t-test was used to determine significance).
Empty triangles indicate non-significant enrichment ($p>0.05$) of *oskar* relative to the GFP control. Error bars
show SDn.

916
Figure 5: Effect of Egl overexpression on *oskar* RNP localization and composition.

(A-B') Images of *oskar* mRNA and Egl-GFP (A and A') or Staufen-GFP (B and B') in stage 9 egg-chambers
strongly overexpressing Egl in the germline (*oskar-Gal4>UAS-Egl*). In both genotypes, abnormally large
RNPs containing *oskar* mRNA are observed in the nurse cells and in the anterior region of the oocyte. These
RNPs frequently colocalize with Egl-GFP but rarely co-localize with Staufen-GFP. Scale bar represents 10
µm.

(C-D') Quantification of Egl-GFP and Staufen-GFP association with *oskar* RNPs as a function of RNA copy
number. Error bars represent 95% confidence intervals and the size of the circles is proportional to the
relative abundance of each category of *oskar* RNP within the overall population. Triangles indicate that the
fraction of GFP-positive *oskar* RNPs is not-significantly different from zero (p>0.01, C-D').

(E) Relative distribution of *oskar* RNPs grouped by RNA content along the first 30% of the anteroposterior
axis of stage 9 oocytes. In the wild-type control, less than 1% of large (65+ copies) *oskar* RNPs are close to
the oocyte anterior (first bin). When Egl is overexpressed, approximately 10% of large *oskar* RNPs are
present close to the oocyte anterior (first bin, arrow), while the rest of the oocyte has a distribution of *oskar*
RNPs similar to the control (Fig S5A).

933
Figure 6: Recruitment of the dynein machinery during the transport of *oskar* RNPs.

(A-C) Localization of GFP tagged versions (green) of BicD (A), p50/dynamitin (B) and Dhc (C) with respect to
oskar mRNA (magenta) in egg-chambers during mid-oogenesis. Scale bar represents 10 μm.

(D, E) Quantification of *oskar* RNP association with the indicated proteins in the nurse cells and the oocyte as a
function of RNA copy number.

(F, G) Quantification of *oskar* RNP association with GFP-Dhc in the oocytes of indicated genotypes.

Error bars represent 95% confidence intervals and the size of the circles is proportional to the relative abundance
of each category of *oskar* RNP within the overall population. Triangles indicate that the fraction of GFP-positive
oskar RNPs is not-significantly different from zero (p>0.01, D-G).

943
944

945
Figure 7: Egl promotes ooplasmic enrichment of Staufen mRNA and protein

(A-C') Distribution of endogenous *oskar* mRNA (A, A'; green), *stau* mRNA (B, B'; cyan) and Staufen protein
(C, C'; magenta) in early egg-chambers expressing control RNAi (A, B, C) or *egl* RNAi driven by *osk*-Gal4 (A',
B', C'). Oocytes of *egl* RNAi egg-chambers contain trace amounts of *oskar* mRNA, likely due to the action of
residual Egl protein.

(D) Quantification of oocyte enrichment of *stau* mRNA or Staufen protein relative to the sibling nurse cells.
Oocytes were identified through their enrichment of *oskar mRNA.* Enrichment of *stau* RNA or Staufen protein
in the somatic follicle cells, which do not express the shRNA, is used as a control. 19 control RNAi and 16 *egl*
RNAi samples were analyzed, respectively (also indicated on the panel). P-values of unpaired student's t-test
are shown.

(E, E') Localization of *stau* mRNA in wild-type and Egl-overexpressing stage 8 oocytes.

(F, G) Fraction of *stau* RNPs associating with Egl in the nurse cells and in the oocyte (F), and Staufen in the
oocyte (G). Scale bars represent 10 μm.

962
Figure S1: Expression and effect of the different Staufen transgenes on *oskar* mRNA localization.

(A) *oskar* RNA localization (shown in blue/yellow) in stage 9 *stau^{R9}/stau^{D3}* mutant oocytes. Scale bars
represent 20 µm.

(B-F) Mean (green) and variance (SD, magenta) of *oskar* RNA distribution in stage 9 and stage 10-11
oocytes. Numbers indicate the number of oocytes analyzed for each condition and scale bar represents 10%
of anteroposterior axis length. Anterior is to the left, posterior is to the right.

(G) Position of *oskar* center-of-mass along the AP axis in stage 9 and stage 10-11 oocytes for each condition.
0 is the geometric center of the oocyte, with the posterior pole located at 58%.

(H) Oskar protein expression in Staufen null oocytes coexpressing transgenic Staufen-GFP or GFP-Staufen.
Scale bars represent 20 µm.

(I) Western blot detection of Staufen in wild-type (lane 1), *stau^{R9}/stau^{D3}* (2), Staufen-GFP (3), GFP-Staufen
(4), *stau* RNAi (5) and control RNAi (6) ovarian lysates. Note the similar distribution of *oskar* in *stau^{R9}/stau^{D3}*
and in *stau* RNAi oocytes (C, D and G), despite residual Staufen expression in Staufen RNAi ovarian lysates
(I, lanes 5 and 6). The overexpressing GFP-Staufen (I, lane 4) transgene largely rescues *oskar*
mislocalization (F and G) and Oskar protein expression defects (H) observed in Staufen null mutants. The
Staufen-GFP transgene, expressed at low levels (I, lane 3), rescues *oskar* RNA localization at stage 9, but
the RNA is not maintained at the posterior at stages 10-11 (E and G), likely due to insufficient Oskar protein
expression at the posterior (H), which is essential for *oskar* mRNA anchoring at the oocyte posterior during
the later stages (Vanzo and Ephrussi, 2002).

Figure S2. Egl and BicD are required for formation of transport-competent dynein-RNA complexes *in vitro*.

(A) Example kymographs (time-distance plots) showing behavior of dynein and *oskar* RNA in the presence and absence of Egl and BicD; in each condition, dynactin is also present but not fluorescently labeled (filled and empty circles represent the presence and absence of indicated proteins, respectively). Egl and BicD were co-expressed and co-purified (see Methods).

990 (B and C) Charts showing the total number of microtubule (MT) binding events for *oskar* RNA (B) and number of
991 processive dynein complexes on microtubules under conditions shown in A. In B, values were corrected for non-
992 specific background binding of *oskar* RNA to the imaging surface as described in Methods. Mean ± SD of values
993 from individual microtubules (represented by black circles) are displayed. Statistical significance and P-values
994 were determined with Mann-Whitney tests.

995
996

Figure S3. Example kymographs (time-distance plots) showing the behavior of dynein activated by *oskar* RNA, Egl/BicD, and dynactin or by BicD2N and dynactin in the presence (filled circle) and absence (open circle) of Staufen. Quantification of these data is presented in Fig. 2H and I.

1004
Figure S4: Suppression of *oskar* mislocalization

(A-J) Average distribution of *oskar* mRNA (green) and variability of the distribution (SD, magenta) in stage 9
oocytes of the indicated genotypes. N indicates the number of oocytes analyzed. Scale bar represents 10% of
anteroposterior axis length. Anterior is to the left, posterior to the right. Arrowheads indicate the ectopic
localization of *oskar* mRNA at the anterolateral corners.

(K) Distribution of the observed *oskar* center-of-mass in stage 9 *stau* RNAi oocytes in the presence of two
(red) or one (yellow) functional copies of e*gl* as a function of oocyte size, used here as a proxy for
developmental stage. Solid lines show the best linear fits to the data. The equation and the square of the
goodness-of-fit (R^2) are indicated. Such moderate rescue was expected as *oskar* RNPs entering the oocyte
are thought to be associated with Egl. Note that there is no significant linear correlation between oocyte size
(developmental stage) and the position of *oskar* RNA center-of-mass in *stau* RNAi (red), indicating an *oskar*
mislocalization phenotype. There is a moderate correlation with a significant slope (underlined) when one
copy of e*gl* is removed (yellow), indicating progressive posterior localization of *oskar* mRNA at stage 9.

1018
Figure S5: Staufen and Egl association with *oskar* RNPs

(A, B) *oskar* mRNA content distribution in (A) wild-type nurse cells and oocytes at stages 5-7 (yellow) and 9
(blue) of oogenesis, and (B) in *stau* null (red) and dominant *BicD¹* mutants (purple). In the nurse cells, most *oskar*
RNPs contain 1-2 copies of the RNA, consistent with previous reports (Little et al., 2015). *oskar* RNP content
increases in the oocyte: at stages 5-7 most RNPs contain 4+ copies of the RNA, which decreases to >2 copies
during *oskar* posterior localization (stage 9; Little et al., 2015).

(C, D) Normalized Staufen-GFP (B') or Egl-GFP (D', E') signal intensity as a function of *oskar* mRNA content at
stages 5-7 (yellow) and stage 9 (blue). (C) Staufen-GFP signal intensity in the complete absence (yellow and
blue) and in the presence (red) of endogenous, unlabeled Staufen. Fitted linear models showing the correlation
between (C) Staufen-GFP and (D) Egl-GFP signal intensity and *oskar* mRNA copy number as solid lines and
equations (top - stages 5-7, bottom - stage 9). Underscored parameters of the models are significantly different
from zero (p<0.05). The slopes of the two fitted models are significantly different (p<0.0001, ANOVA).

(E-F') Association of Egl-GFP with *oskar* RNPs in oocytes expressing (E, E') *stau* RNAi (red), control RNAi
(brown), or with (F, F') *BicD¹* (purple) or *BicD²* (pink) alleles. (E) Note that knock-down of Staufen results in
similar retention of Egl on *oskar* RNPs as in the complete absence of Staufen protein (Fig 3E, E'). (D-F') Egg-
chambers expressed a single copy of Egl-GFP in the presence of two endogenous wild-type e*gl* alleles, except in
the case of the rescued e*gl* mutants (F, *egl¹/egl²,* green). Although when unlabelled Egl was absent (*egl¹/egl²,*
green), we observed a slightly elevated fraction of Egl positive RNPs, larger RNPs containing 16+ copies of *oskar*
mRNA displayed no significant association with Egl (F) and the relative amounts of Egl on *oskar* RNPs were
identical to what was observed in the presence of endogenous, unlabelled Egl (F', blue).

(G) Mean signal intensity of Staufen-GFP measured at multiple locations throughout developing oocytes. Size of
the oocytes (x-axis) is used as a proxy of developmental time and, along with morphological features, for staging
of the oocytes (shaded areas as indicated in the panel). (C-G) Error bars show 95% confidence interval of the
mean.

(H) Western blot of input lysates and eluates after RNA immunoprecipitation in the presence (lane 3) or the
absence (lane 4) of Staufen. Bait proteins - monomeric EGFP (lane 1), GFP-Staufen (lane 2) and Egl-GFP (lanes
3,4) - are detected by anti-GFP antibody. Anti-tubulin staining served to monitor potential contamination of the
eluates.

(I) Western blot showing endogenous Egl protein detected by anti-Egl antibody in the indicated genotypes.
Tubulin was used as a loading control.

In H and I, blue and yellow indicate low and high intensity of signal, respectively.

.
1051
1052
1053
1054

A

1055

1056 **Figure S6: Localization of *oskar* RNPs along the anteroposterior axis.**
1057 (A) Relative distribution of *oskar* RNPs grouped by RNA content along the anteroposterior axis in wild-type
1058 and in Egl overexpressing oocytes.

Figure S7: Localization of Staufen protein and *stau* mRNA in the egg-chamber

(A-B') Staufen protein expression detected by immunofluorescence (blue, A and A') or by the fluorescent reporter Staufen-GFP (B and B', green) in early egg-chambers in the presence (A and B) and complete absence of *oskar* mRNA (A' and B'). Note that lack of *oskar* from the oocyte blocks progression of oogenesis beyond stage 6 (Jenny et al., 2006).

(C) Enrichment of Staufen-GFP signal in early oocytes with one (green) or no (red) functional *oskar* alleles expressing *oskar* mRNA, and in oocytes expressing *egl* RNAi (gray) or control RNAi (brown). Enrichment is relative to the sibling nurse cells in the egg-chamber. Enrichment of Staufen-GFP in somatic follicle cells, which do not express the shRNA, serves as a control. P-values of pairwise Student's t-test are shown. Note that complete lack of *oskar* mRNA, an abundant binding partner of Staufen, has only a moderate effect on Staufen enrichment in the developing oocyte (also observed in A-B'), whereas knock-down of Egl almost completely abolishes Staufen ooplasmic accumulation.

(D) Quantification of the enrichment of *stau* mRNA in the oocyte in *egl* RNAi (gray), control RNAi (yellow) and Egl overexpressing oocytes (dark blue). Note that an excess of Egl has a minuscule effect on *stau* RNA accumulation in the oocyte, suggesting that in the wild-type, most of the *stau* mRNA expressed in the nurse cells is transported into the oocyte.

(E-F'') Early egg-chambers overexpressing GFP-Staufen (F, blue) under control of the *matTub*-Gal4 driver. Note that in control oocytes of a similar stage (E-E''), and in oocytes expressing low levels of GFP-Staufen (F, left egg-chamber), *oskar* mRNA (E'', F'', magenta) is enriched. Enrichment of *oskar* and *bicoid* (E', F', green) mRNAs is greatly reduced in oocytes expressing high levels of GFP-Staufen (F, right egg-chamber). This phenotype is reproducibly observed in >30 egg-chambers derived from three separate crosses.

(G-H') Egg-chambers in early and mid-oogenesis overexpressing GFP-Staufen (blue) under control of the *oskar*-Gal4 driver from the beginning of oogenesis. The vast majority of such egg-chambers fail to develop beyond stage 6, likely as a consequence of greatly reduced ooplasmic accumulation of *oskar* mRNA (green), which appears to be trapped in the nurse cells in large aggregates associated with GFP-Staufen (some examples are highlighted by arrowheads in panels F-H'). Such aggregates of endogenous Staufen are not observed in wild-type egg-chambers (E-E''). Similarly, no accumulation of *stau* mRNA in the oocyte is observed in these *oskar-Gal4>UAS-GFP-Staufen* oocytes (G' and H'), where *stau* mRNA levels are uniformly high in the germline (compare the signal in the follicle cell layer to that in the nurse cells and in the oocyte; see Figure 7). (H, H') In the oocytes occasionally escaping early developmental arrest, we invariably observed failure in nuclear migration from the posterior to the anterior, reflecting a defect in repolarization of the oocyte microtubule network (Januschke et al., 2006). Consequently, *oskar* mRNA remains in the center of these oocytes, which - although they complete oogenesis - fail to result in viable progeny. These phenotypes are reproducibly observed in >30 egg-chambers derived from three separate crosses.

(I) Expression of GFP-Staufen under the control of maternal tubulin promoter. GFP-Staufen (cyan) can hardly be detected in early egg chambers (highlighted by arrows), and the forming aggregates remain associated with the nurse cell nuclei until mid-oogenesis. Scale bars represent 5 μm.

1102 Video S1 and S2: *Ex vivo* imaging of *oskMS2*-GFP RNPs in extracts of control (Video S1) and *stau* RNAi
1103 (Video S4) stage 9 oocytes. *oskar* RNPs are shown in green, microtubule plus tips are labeled by EB1-
1104 mCherry (magenta). Note the frequent runs of dim *oskar* RNPs in the *stau* RNAi condition (Video S2)
1105
1106 Video S3 and S4: Live cell imaging of *oskMS2*-GFP RNPs in control (Video S3) and *stau* RNAi (Video S4)
1107 stage 9 oocytes. Anterior (left) and posterior (right) regions of the same oocyte are shown. *oskMS2*-GFP
1108 signal is rendered in blue-yellow to allow visual appreciation of dim (blue) and bright (yellow) *oskar* RNPs.
1109
1110
1111 Table S1: ssDNA oligos used to synthesize smFISH probes targeting the long 3' UTR of *stau* A/B isoforms or
1112 the 5' extended regions of *stau* A/C isoforms.
1113

REFERENCES

Abouward, R., and G. Schiavo. 2021. Walking the line: mechanisms underlying directional mRNA transport and localisation in neurons and beyond. Cell. Mol. Life Sci. 78:2665–2681. doi:10.1007/s00018-020-03724-3.

Amrute-Nayak, M., and S.L. Bullock. 2012. Single-molecule assays reveal that RNA localization signals regulate dynein-dynactin copy number on individual transcript cargoes. Nat. Cell Biol. 14:416–423. doi:10.1038/ncb2446.

Bauer, K.E., I. Segura, I. Gaspar, V. Scheuss, C. Illig, G. Ammer, S. Hutten, E. Basyuk, S.M. Fernández-Moya, J. Ehses, E. Bertrand, and M.A. Kiebler. 2019. Live cell imaging reveals 3'-UTR dependent mRNA sorting to synapses. Nat. Commun. 10:3178. doi:10.1038/s41467-019-11123-x.

Baumann, S., T. Pohlmann, M. Jungbluth, A. Brachmann, and M. Feldbrügge. 2012. Kinesin-3 and dynein mediate microtubule-dependent co-transport of mRNPs and endosomes. J. Cell Sci. 125:2740–2752. doi:10.1242/jcs.101212.

Benaglia, T., D. Chauveau, D.R. Hunter, and D. Young. 2009. mixtools□: An R package for Analyzing Finite Mixture Models. J. Stat. Softw. 32. doi:10.18637/jss.v032.i06.

Berleth, T., M. Burri, G. Thoma, D. Bopp, S. Richstein, G. Frigerio, M. Noll, and C. Nüsslein-Volhard. 1988. The role of localization of bicoid RNA in organizing the anterior pattern of the Drosophila embryo. EMBO J. 7:1749–1756. doi:10.1002/j.1460-2075.1988.tb03004.x.

Brendza, R.P., L.R. Serbus, J.B. Duffy, and W.M. Saxton. 2000. A function for kinesin I in the posterior transport of oskar mRNA and Staufen protein. Science. 289:2120–2122. doi:10.1126/science.289.5487.2120.

Bullock, S.L., A. Nicol, S.P. Gross, and D. Zicha. 2006. Guidance of bidirectional motor complexes by mRNA cargoes through control of dynein number and activity. Curr. Biol. 16:1447–1452. doi:10.1016/j.cub.2006.05.055.

Clark, A., C. Meignin, and I. Davis. 2007. A Dynein-dependent shortcut rapidly delivers axis determination transcripts into the Drosophila oocyte. Development. 134:1955–1965. doi:10.1242/dev.02832.

Dienstbier, M., F. Boehl, X. Li, and S.L. Bullock. 2009. Egalitarian is a selective RNA-binding protein linking mRNA localization signals to the dynein motor. Genes Dev. 23:1546–1558. doi:10.1101/gad.531009.

Dix, C.I., H.C. Soundararajan, N.S. Dzhindzhev, F. Begum, B. Suter, H. Ohkura, E. Stephens, and S.L. Bullock. 2013. Lissencephaly-1 promotes the recruitment of dynein and dynactin to transported mRNAs. J. Cell Biol. 202:479–494. doi:10.1083/jcb.201211052.

Edelstein, A.D., M.A. Tsuchida, N. Amodaj, H. Pinkard, R.D. Vale, and N. Stuurman. 2014. Advanced methods of microscope control using μManager software. J. Biol. Methods. 1. doi:10.14440/jbm.2014.36.

van Eeden, F.J., I.M. Palacios, M. Petronczki, M.J. Weston, and D. St Johnston. 2001. Barentsz is essential for the posterior localization of oskar mRNA and colocalizes with it to the posterior pole. J. Cell Biol. 154:511–523. doi:10.1083/jcb.200105056.

Ephrussi, A., L.K. Dickinson, and R. Lehmann. 1991. Oskar organizes the germ plasm and directs localization of the posterior determinant nanos. Cell. 66:37–50. doi:10.1016/0092-8674(91)90137-n.

Ephrussi, A., and R. Lehmann. 1992. Induction of germ cell formation by oskar. Nature. 358:387–392. doi:10.1038/358387a0.

Ferrandon, D., L. Elphick, C. Nüsslein-Volhard, and D. St Johnston. 1994. Staufen protein associates with the 3'UTR of bicoid mRNA to form particles that move in a microtubule-dependent manner. Cell. 79:1221–1232. doi:10.1016/0092-8674(94)90013-2.

Gagnon, J.A., J.A. Kreiling, E.A. Powrie, T.R. Wood, and K.L. Mowry. 2013. Directional transport is mediated by a Dynein-dependent step in an RNA localization pathway. PLoS Biol. 11:e1001551. doi:10.1371/journal.pbio.1001551.

Gaspar, I., and A. Ephrussi. 2017. Ex vivo Ooplasmic Extract from Developing Drosophila Oocytes for Quantitative TIRF Microscopy Analysis. Bio Protoc. 7. doi:10.21769/BioProtoc.2380.

Gáspár, I., V. Sysoev, A. Komissarov, and A. Ephrussi. 2017. An RNA-binding atypical tropomyosin recruits kinesin-1 dynamically to oskar mRNPs. EMBO J. 36:319–333. doi:10.15252/embj.201696038.

Gaspar, I., F. Wippich, and A. Ephrussi. 2017. Enzymatic production of single-molecule FISH and RNA capture probes. RNA. 23:1582–1591. doi:10.1261/rna.061184.117.

Gaspar, I., Y.V. Yu, S.L. Cotton, D.-H. Kim, A. Ephrussi, and M.A. Welte. 2014. Klar ensures thermal robustness of oskar localization by restraining RNP motility. J. Cell Biol. 206:199–215. doi:10.1083/jcb.201310010.

Gaspar, I. 2011. Microtubule-based motor-mediated mRNA localization in Drosophila oocytes and embryos. Biochem. Soc. Trans. 39:1197–1201. doi:10.1042/BST0391197.

Ghosh, S., V. Marchand, I. Gáspár, and A. Ephrussi. 2012. Control of RNP motility and localization by a splicing-dependent structure in oskar mRNA. Nat. Struct. Mol. Biol. 19:441–449. doi:10.1038/nsmb.2257.

Glock, C., M. Heumüller, and E.M. Schuman. 2017. mRNA transport & local translation in neurons. Curr. Opin. Neurobiol. 45:169–177. doi:10.1016/j.conb.2017.05.005.

Gratz, S.J., A.M. Cummings, J.N. Nguyen, D.C. Hamm, L.K. Donohue, M.M. Harrison, J. Wildonger, and K.M. O'Connor-Giles. 2013. Genome engineering of Drosophila with the CRISPR RNA-guided Cas9 nuclease. Genetics. 194:1029–1035. doi:10.1534/genetics.113.152710.

Hachet, O., and A. Ephrussi. 2001. Drosophila Y14 shuttles to the posterior of the oocyte
and is required for oskar mRNA transport. Curr. Biol. 11:1666–1674. doi:10.1016/s0960-
9822(01)00508-5.

Hachet, O., and A. Ephrussi. 2004. Splicing of oskar RNA in the nucleus is coupled to its
cytoplasmic localization. Nature. 428:959–963. doi:10.1038/nature02521.

Hancock, W.O. 2014. Bidirectional cargo transport: moving beyond tug of war. Nat. Rev.
Mol. Cell Biol. 15:615–628. doi:10.1038/nrm3853.

Heber, S., I. Gáspár, J.-N. Tants, J. Günther, S.M.F. Moya, R. Janowski, A. Ephrussi, M.
Sattler, and D. Niessing. 2019. Staufen2-mediated RNA recognition and localization requires
combinatorial action of multiple domains. Nat. Commun. 10:1659. doi:10.1038/s41467-019-
09655-3.

Heraud-Farlow, J.E., T. Sharangdhar, X. Li, P. Pfeifer, S. Tauber, D. Orozco, A.
Hörmann, S. Thomas, A. Bakosova, A.R. Farlow, D. Edbauer, H.D. Lipshitz, Q.D. Morris, M.
Bilban, M. Doyle, and M.A. Kiebler. 2013. Staufen2 regulates neuronal target RNAs. Cell
Rep. 5:1511–1518. doi:10.1016/j.celrep.2013.11.039.

Hoang, H.T., M.A. Schlager, A.P. Carter, and S.L. Bullock. 2017. DYNC1H1 mutations
associated with neurological diseases compromise processivity of dynein-dynactin-cargo
adaptor complexes. Proc Natl Acad Sci USA. 114:E1597–E1606.
doi:10.1073/pnas.1620141114.

Hoogenraad, C.C., and A. Akhmanova. 2016. Bicaudal D family of motor adaptors: linking
dynein motility to cargo binding. Trends Cell Biol. 26:327–340.
doi:10.1016/j.tcb.2016.01.001.

Hoogenraad, C.C., P. Wulf, N. Schiefermeier, T. Stepanova, N. Galjart, J.V. Small, F.
Grosveld, C.I. de Zeeuw, and A. Akhmanova. 2003. Bicaudal D induces selective dynein-
mediated microtubule minus end-directed transport. EMBO J. 22:6004–6015.
doi:10.1093/emboj/cdg592.

Jambor, H., S. Mueller, S.L. Bullock, and A. Ephrussi. 2014. A stem-loop structure directs
oskar mRNA to microtubule minus ends. RNA. 20:429–439. doi:10.1261/rna.041566.113.

Januschke, J., L. Gervais, S. Dass, J.A. Kaltschmidt, H. Lopez-Schier, D. St Johnston,
A.H. Brand, S. Roth, and A. Guichet. 2002. Polar transport in the Drosophila oocyte requires
Dynein and Kinesin I cooperation. Curr. Biol. 12:1971–1981. doi:10.1016/S0960-
9822(02)01302-7.

Januschke, J., L. Gervais, L. Gillet, G. Keryer, M. Bornens, and A. Guichet. 2006. The
centrosome-nucleus complex and microtubule organization in the Drosophila oocyte.
Development. 133:129–139. doi:10.1242/dev.02179.

Jenny, A., O. Hachet, P. Závorszky, A. Cyrklaff, M.D.J. Weston, D.S. Johnston, M.
Erdélyi, and A. Ephrussi. 2006. A translation-independent role of oskar RNA in early
Drosophila oogenesis. Development. 133:2827–2833. doi:10.1242/dev.02456.

Kim-Ha, J., J.L. Smith, and P.M. Macdonald. 1991. oskar mRNA is localized to the
posterior pole of the Drosophila oocyte. Cell. 66:23–35. doi:10.1016/0092-8674(91)90136-m.

Lasko, P. 2012. mRNA localization and translational control in Drosophila oogenesis.
Cold Spring Harb. Perspect. Biol. 4. doi:10.1101/cshperspect.a012294.

Laver, J.D., X. Li, K. Ancevicius, J.T. Westwood, C.A. Smibert, Q.D. Morris, and H.D.
Lipshitz. 2013. Genome-wide analysis of Staufen-associated mRNAs identifies secondary
structures that confer target specificity. Nucleic Acids Res. 41:9438–9460.
doi:10.1093/nar/gkt702.

Lenth, R.V. 2016. Least-Squares Means: the R package lsmeans. J. Stat. Softw. 69:1–33.
doi:10.18637/jss.v069.i01.

Little, S.C., K.S. Sinsimer, J.J. Lee, E.F. Wieschaus, and E.R. Gavis. 2015. Independent
and coordinate trafficking of single Drosophila germ plasm mRNAs. Nat. Cell Biol. 17:558–
568. doi:10.1038/ncb3143.

Liu, Y., H.K. Salter, A.N. Holding, C.M. Johnson, E. Stephens, P.J. Lukavsky, J.
Walshaw, and S.L. Bullock. 2013. Bicaudal-D uses a parallel, homodimeric coiled coil with
heterotypic registry to coordinate recruitment of cargos to dynein. Genes Dev. 27:1233–
1246. doi:10.1101/gad.212381.112.

Mach, J.M., and R. Lehmann. 1997. An Egalitarian-BicaudalD complex is essential for
oocyte specification and axis determination in Drosophila. Genes Dev. 11:423–435.
doi:10.1101/gad.11.4.423.

Marchand, V., I. Gaspar, and A. Ephrussi. 2012. An intracellular transmission control
protocol: assembly and transport of ribonucleoprotein complexes. Curr. Opin. Cell Biol.
24:202–210. doi:10.1016/j.ceb.2011.12.014.

Martin, K.C., and A. Ephrussi. 2009. mRNA localization: gene expression in the spatial
dimension. Cell. 136:719–730. doi:10.1016/j.cell.2009.01.044.

McClintock, M.A., C.I. Dix, C.M. Johnson, S.H. McLaughlin, R.J. Maizels, H.T. Hoang,
and S.L. Bullock. 2018. RNA-directed activation of cytoplasmic dynein-1 in reconstituted
transport RNPs. eLife. 7. doi:10.7554/eLife.36312.

McKenney, R.J., W. Huynh, M.E. Tanenbaum, G. Bhabha, and R.D. Vale. 2014.
Activation of cytoplasmic dynein motility by dynactin-cargo adapter complexes. Science.
345:337–341. doi:10.1126/science.1254198.

McKenney, R.J., M. Vershinin, A. Kunwar, R.B. Vallee, and S.P. Gross. 2010. LIS1 and
NudE induce a persistent dynein force-producing state. Cell. 141:304–314.
doi:10.1016/j.cell.2010.02.035.

Micklem, D.R., J. Adams, S. Grünert, and D. St Johnston. 2000. Distinct roles of two
conserved staufen domains in oskar mRNA localization and translation. EMBO J. 19:1366–
1377. doi:10.1093/emboj/19.6.1366.

Mofatteh, M., and S.L. Bullock. 2017. SnapShot: Subcellular mRNA Localization. Cell.
169:178-178.e1. doi:10.1016/j.cell.2017.03.004.

Mohler, J., and E.F. Wieschaus. 1986. Dominant maternal-effect mutations of Drosophila
melanogaster causing the production of double-abdomen embryos. Genetics. 112:803–822.
doi:10.1093/genetics/112.4.803.

Mohr, S., A. Kenny, S.T.Y. Lam, M.B. Morgan, C.A. Smibert, H.D. Lipshitz, and P.M.
Macdonald. 2021. Opposing roles for Egalitarian and Staufen in transport, anchoring and
localization of oskar mRNA in the Drosophila oocyte. PLoS Genet. 17:e1009500.
doi:10.1371/journal.pgen.1009500.

Mohr, S.E., S.T. Dillon, and R.E. Boswell. 2001. The RNA-binding protein Tsunagi
interacts with Mago Nashi to establish polarity and localize oskar mRNA during Drosophila
oogenesis. Genes Dev. 15:2886–2899. doi:10.1101/gad.927001.

Navarro, C., H. Puthalakath, J.M. Adams, A. Strasser, and R. Lehmann. 2004. Egalitarian
binds dynein light chain to establish oocyte polarity and maintain oocyte fate. Nat. Cell Biol.
6:427–435. doi:10.1038/ncb1122.

Neuman-Silberberg, F.S., and T. Schüpbach. 1993. The Drosophila dorsoventral
patterning gene gurken produces a dorsally localized RNA and encodes a TGFα-like protein.
Cell. 75:165–174. doi:10.1016/S0092-8674(05)80093-5.

Newmark, P.A., and R.E. Boswell. 1994. The mago nashi locus encodes an essential
product required for germ plasm assembly in Drosophila. Development. 120:1303–1313.

O'Connor-Giles, Wildonger, and Harrison. 2014. Generating targeting chiRNAs.
FlyCRISPR Web Site.

Palacios, I.M., D. Gatfield, D. St Johnston, and E. Izaurralde. 2004. An eIF4AIII-
containing complex required for mRNA localization and nonsense-mediated mRNA decay.
Nature. 427:753–757. doi:10.1038/nature02351.

Palacios, I.M., and D. St Johnston. 2002. Kinesin light chain-independent function of the
Kinesin heavy chain in cytoplasmic streaming and posterior localisation in the Drosophila
oocyte. Development. 129:5473–5485. doi:10.1242/dev.00119.

Parton, R.M., R.S. Hamilton, G. Ball, L. Yang, C.F. Cullen, W. Lu, H. Ohkura, and I.
Davis. 2011. A PAR-1-dependent orientation gradient of dynamic microtubules directs
posterior cargo transport in the Drosophila oocyte. J. Cell Biol. 194:121–135.
doi:10.1083/jcb.201103160.

R Core Team. 2014. R: A language and environment for statistical computing. R
Foundation for StatisticalComputing.

Sanghavi, P., S. Laxani, X. Li, S.L. Bullock, and G.B. Gonsalvez. 2013. Dynein associates
with oskar mRNPs and is required for their efficient net plus-end localization in Drosophila
oocytes. PLoS ONE. 8:e80605. doi:10.1371/journal.pone.0080605.

Sanghavi, P., G. Liu, R. Veeranan-Karmegam, C. Navarro, and G.B. Gonsalvez. 2016.
Multiple roles for Egalitarian in polarization of the Drosophila egg chamber. Genetics.
203:415–432. doi:10.1534/genetics.115.184622.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S.
Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D.J. White, V. Hartenstein, K.
Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-
image analysis. Nat. Methods. 9:676–682. doi:10.1038/nmeth.2019.

Schlager, M.A., H.T. Hoang, L. Urnavicius, S.L. Bullock, and A.P. Carter. 2014. In vitro
reconstitution of a highly processive recombinant human dynein complex. EMBO J.
33:1855–1868. doi:10.15252/embj.201488792.

Schneider, C.A., W.S. Rasband, and K.W. Eliceiri. 2012. NIH Image to ImageJ: 25 years
of image analysis. Nat. Methods. 9:671–675. doi:10.1038/nmeth.2089.

Schuldt, A.J., J.H. Adams, C.M. Davidson, D.R. Micklem, J. Haseloff, D. St Johnston, and
A.H. Brand. 1998. Miranda mediates asymmetric protein and RNA localization in the
developing nervous system. Genes Dev. 12:1847–1857. doi:10.1101/gad.12.12.1847.

Sladewski, T.E., N. Billington, M.Y. Ali, C.S. Bookwalter, H. Lu, E.B. Krementsova, T.A.
Schroer, and K.M. Trybus. 2018. Recruitment of two dyneins to an mRNA-dependent
Bicaudal D transport complex. eLife. 7. doi:10.7554/eLife.36306.

Soundararajan, H.C., and S.L. Bullock. 2014. The influence of dynein processivity control,
MAPs, and microtubule ends on directional movement of a localising mRNA. eLife.
3:e01596. doi:10.7554/eLife.01596.

St Johnston, D., D. Beuchle, and C. Nüsslein-Volhard. 1991. Staufen, a gene required to
localize maternal RNAs in the Drosophila egg. Cell. 66:51–63. doi:10.1016/0092-
8674(91)90138-o.

St Johnston, D., N.H. Brown, J.G. Gall, and M. Jantsch. 1992. A conserved double-
stranded RNA-binding domain. Proc Natl Acad Sci USA. 89:10979–10983.
doi:10.1073/pnas.89.22.10979.

St Johnston, D. 2005. Moving messages: the intracellular localization of mRNAs. Nat.
Rev. Mol. Cell Biol. 6:363–375. doi:10.1038/nrm1643.

Theurkauf, W.E., B.M. Alberts, Y.N. Jan, and T.A. Jongens. 1993. A central role for
microtubules in the differentiation of Drosophila oocytes. Development. 118:1169–1180.
doi:10.1242/dev.118.4.1169.

Turner-Bridger, B., M. Jakobs, L. Muresan, H.H.-W. Wong, K. Franze, W.A. Harris, and
C.E. Holt. 2018. Single-molecule analysis of endogenous β-actin mRNA trafficking reveals a
mechanism for compartmentalized mRNA localization in axons. Proc Natl Acad Sci USA.
115:E9697–E9706. doi:10.1073/pnas.1806189115.

Urnavicius, L., K. Zhang, A.G. Diamant, C. Motz, M.A. Schlager, M. Yu, N.A. Patel, C.V.
Robinson, and A.P. Carter. 2015. The structure of the dynactin complex and its interaction
with dynein. Science. 347:1441–1446. doi:10.1126/science.aaa4080.

Vazquez-Pianzola, P., B. Schaller, M. Colombo, D. Beuchle, S. Neuenschwander, A.
Marcil, R. Bruggmann, and B. Suter. 2017. The mRNA transportome of the BicD/Egl
transport machinery. RNA Biol. 14:73–89. doi:10.1080/15476286.2016.1251542.

Wickham, H. 2016. ggplot2 - Elegant Graphics for Data Analysis . Springer-Verlag New
York, New York, NY.

Williams, L.S., S. Ganguly, P. Loiseau, B.F. Ng, and I.M. Palacios. 2014. The auto-
inhibitory domain and ATP-independent microtubule-binding region of Kinesin heavy chain
are major functional domains for transport in the Drosophila germline. Development.
141:176–186. doi:10.1242/dev.097592.

Yoon, Y.J., and K.L. Mowry. 2004. Xenopus Staufen is a component of a
ribonucleoprotein complex containing Vg1 RNA and kinesin. Development. 131:3035–3045.
doi:10.1242/dev.01170.

Zappulo, A., D. van den Bruck, C. Ciolli Mattioli, V. Franke, K. Imami, E. McShane, M.
Moreno-Estelles, L. Calviello, A. Filipchyk, E. Peguero-Sanchez, T. Müller, A. Woehler, C.
Birchmeier, E. Merino, N. Rajewsky, U. Ohler, E.O. Mazzoni, M. Selbach, A. Akalin, and M.
Chekulaeva. 2017. RNA localization is a key determinant of neurite-enriched proteome. Nat.
Commun. 8:583. doi:10.1038/s41467-017-00690-6.

Zimyanin, V.L., K. Belaya, J. Pecreaux, M.J. Gilchrist, A. Clark, I. Davis, and D. St
Johnston. 2008. In vivo imaging of oskar mRNA transport reveals the mechanism of
posterior localization. Cell. 134:843–853. doi:10.1016/j.cell.2008.06.053.