A chromosome-level genome assembly of a maize elite breeding line Dan340

Yikun Zhao[1,5], Yuancong Wang[2,5], De Ma[3,5], Guang Feng[4], Yongxue Huo[1], Zhihao

Liu[1], Ling Zhou[2], Yunlong Zhang[1], Liwen Xu[1], Liang Wang[4], Han Zhao[2,*], Jiuran

Zhao[1,*], Fengge Wang[1,*]

[1] Maize Research Center, Beijing Academy of Agricultural and Forest Sciences (BAAFS)/Beijing

Key Laboratory of Maize DNA Fingerprinting and Molecular Breeding

[2] Provincial Key Laboratory of Agrobiology, Institute of Crop Germplasm and Biotechnology,

Jiangsu Academy of Agricultural Sciences

[3] Novogene Bioinformatics Institute

[4] Dandong Academy of Agricultural Sciences

[5] These authors contributed equally to this article.

* **Correspondence**: Han Zhao (zhaohan@jaas.ac.cn), Jiuran Zhao (maizezhao@126.com),

Fengge Wang (fenggewangmaize@126.com)

Key findings

1. The final assembly of Dan340 genome was 2,348.72 Mb, including 2,738 contigs

and 2,315 scaffolds with a N50 of 41.49 Mb and 215.35 Mb, respectively.

2. The percent of reads mapped to the reference genome was up to 97.48%.

3. The results showed that 96.84% of the plant single-copy orthologues were

complete. Complete single-copy and multi copy genes accounted for 87.36% and

9.48% of the genes, respectively. Taken together, these results indicated that our

Dan340 genome assembly presented high quality and coverage.

Abstract

The maize cultivar Dan340 is an excellent backbone inbred line of Luda Red Cob

Group with several desirable characters, such as disease resistance, lodging resistance,

high combining ability, wide adaptability and so on. In this study, we constructed a

high-quality chromosome-level reference genome for Dan340 by combining PacBio

long HiFi sequencing, Illumina short reads and chromosomal conformational capture

(Hi-C) sequencing reads. The final assembly of Dan340 genome was 2,348.72 Mb,

including 2,738 contigs and 2,315 scaffolds with a N50 of 41.49 Mb and 215.35 Mb,

respectively. The percent of high quality Illumina reads mapped to the reference

genome was up to 97.48%. The assembly of this genome will not only facilitate our

understanding about intraspecific genome diversity in maize, but also provides a

novel resource for maize breeding.

Background

Maize (*Zea mays* ssp. *mays* L.) is one of the most important crops grown worldwide

for food, forage, and biofuel, with an annual production of more than 1 billion tons

(Yang et al., 2017). With the rapid growth of human population and economic

demand, it has been predicted that maize will account for 45% of total cereal demand

by the year 2050 (Hubert et al., 2010). In addition, it is also an important model

organism for fundamental research in genetics and genomics (Hake and Ross-Ibarra,

2015).

Several reference genome of common maize inbred line in breeding have been

released since 2009 (Schnable et al., 2009; Hirsch et al., 2016; Jiao et al., 2017; Sun et

al., 2018; Li et al., 2019). The maize genome exhibits high levels of genetic diversity

among different inbred lines (Lu et al., 2015; Yang et al., 2017; Sun et al., 2018).

The maize cultivar Dan340 is an excellent backbone inbred line of Luda Red Cob

Group with several desirable characters, such as disease resistance, lodging resistance,

high combining ability, wide adaptability and so on (Zhang et al., 2018).

In the present study, we constructed a high-quality chromosome-level reference

genome for Dan340 by combining PacBio long HiFi sequencing, Illumina short reads

and chromosomal conformational capture (Hi-C) sequencing reads. The assembly of

this genome will not only facilitate our understanding about intraspecific genome

diversity in maize, but also provides a novel resource for maize breeding

improvement.

Plant material and DNA sequencing

Inbred line Dan340 was selected for genome sequencing and assembly because it is

an elite maize cultivar, which plays an important role in maize breeding and genetic

research. The plants were grown at 25°C in a greenhouse of Beijing Academy of

Agriculture and Forestry Sciences, Beijing, China. Fresh and tender leaves were

harvested from the best-growing individual and immediately frozen in liquid nitrogen,

followed by preservation at -80 °C in the laboratory prior to DNA extraction.

Genomic DNA was extracted from the leaf tissue of a single plant using the

DNAsecure Plant Kit (Tiangen Biotech Co., Ltd., Beijing, China). To ensure that

DNA extracts were useable for all types of genomic libraries, the quality and quantity

were evaluated using a NanoDrop 2000 spectrophotometer (NanoDrop Technologies,

Wilmington, DE, USA) and electrophoresis on a 0.8% agarose gel, respectively.

PacBio circular consensus sequencing (CCS) libraries were prepared using SMRTbell

Express Template Prep Kit 2.0 (Pacific Biosciences Ref. No. 101-685-400), following

the manufacturer's protocols, and subsequently sequenced on the PacBio sequel II

platform (Pacific Biosciences, Menlo Park, CA, USA). Consensus reads (CCS reads,

also termed HiFi reads) were generated using ccs software v.3.0.0

(https://github.com/pacificbiosciences/unanimity/) with the parameter '--min-passes 3

--min-rq 0.99'. The total CCS yield was 63.54 Gb, with a mean read length of 15.60

kb.

In addition, one Illumina paired-end sequencing library with an insert size of 350 bp

was generated using NEB Next Ultra DNA Library Prep Kit (NEB, USA) following

manufacturer's protocol, and subsequently sequenced using an Illumina HiSeq X Ten

platform (Illumina, San Diego, CA, USA) in Novogene Bioinformatics Institute,

Beijing, China. Approximately 80.66 Gb (~20X) of Illumina sequencing data were

obtained.

One Hi-C library was constructed using young leaves following the published

procedures with certain modifications (Belton et al., 2012). In brief, approximately

5-g leaf samples were cut into minute pieces and cross-linked by 4% formaldehyde

solution at room temperature in a vacuum for 30 min. Then, the sample was mixed

with excess 2.5 M glycine to quench the cross linking reaction for 5 min and then

placed on ice for 15 min. The cross-linked DNA was extracted and then digested with

MboI restriction enzyme. The sticky ends of the digested fragments were biotinylated

and proximity ligated to form ligation junctions that were enriched for and then

ultrasonically sheared to a size of 200-600 bp. The biotin-labelled DNA fragments

were pulled down and ligated with Illumina paired-end adapters and then amplified by

PCR to produce the Hi-C sequencing library. The library was sequenced using an

Illumina HiSeq X Ten platform (Illumina, San Diego, CA, USA).

Genome assembly

To obtain a high quality genome assembly of Dan340, we employed both PacBio HiFi

sequences and Illumina short-read, with scaffolding informed by chromosomal

conformation capture (Hi-C). The assembly was performed in a stepwise fashion.

First, de novo assembly of the long CCS reads generated from the PacBio SMRT

Sequencing was performed using Hifiasm (Cheng et al., 2021). In brief, we first used

all CCS clean reads to do error correction by performing all-vs-all pairwise alignment

between them. In this step, haplotype phasing was achieved using heterozygous SNPs

and most sequencing errors had been removed. Then, error-corrected reads were used

to construct phased assembly graph by determining if one overlap is among the reads

coming from different haplotypes. Next, based on the graph topological structures and

the phasing relationship among different haplotypes, Hifiasm produced a primary

contigs (p-contigs). Then, Pilon (Walker et al., 2014) was used to perform the second

round of error correction with short paired-end reads generated from Illumina Hiseq

Platforms. Subsequently, the Purge Haplotigs pipeline

(https://bitbucket.org/mroachawri/purge_haplotigs/overview) was used to remove

redundant sequences formed due to heterozygosity.

For Hi-C reads, to avoid reads with artificial bias, we removed the following type of

reads: (a) Reads with ≥ 10% unidentified nucleotides (N); (b) Reads with > 10 nt

aligned to the adapter, allowing ≤ 10% mismatches; (c) Reads with > 50% bases

having phred quality < 5. The filtered Hi-C reads were aligned against the contig

assemblies with BWA (version 0.7.8). Reads were excluded from subsequent analysis

if they did not align within 500 bp of a restriction site or did not uniquely map, and

the number of Hi-C read pairs linking each pair of scaffolds is tabulated. ALLHiC

v0.8.12 (Zhang et al., 2019) was used in simple diploid mode to scaffold the genome

and optimize the ordering and orientation of each clustered group, producing a

chromosomal level assembly. Juicebox Assembly Tools v1.9.8

(https://github.com/aidenlab/Juicebox) (Durand et al., 2016) was finally used to

visualize and manually correct the large-scale inversions and translocations to obtain

the final pseudo-chromosomes. The final assembly of Dan340 genome was 2,348.72

Mb, including 2,738 contigs and 2,315 scaffolds with a N50 of 41.49 Mb and 215.35

Mb (Table 1), respectively. A total of 91.30% sequences was anchored to 10

pseudo-chromosomes.

Evaluation of assembly quality

We assessed the quality of the assembly by two independent methods. First, the short

reads obtained from the Illumina sequencing data were aligned to the final assembly

with BWA version 0.7.8 (http://bio-bwa.sourceforge.net/) (Li and Durbin, 2010). Our

result showed that the percent of reads mapped to the reference genome was up to

97.48%. Second, the BUSCO (Benchmarking Universal Single-Copy Otheologs,

(Simao et al., 2015) database embryophyta_odb10 was used to evaluate the

completeness of gene regions, which including 1614 conserved core genes. The

results showed that 96.84% of the plant single-copy orthologues were complete.

Conclusions

1. We assembled the chromosome-level genome of maize elite inbred line Dan340

based on long CCS reads from the third-generation PacBio Sequel II sequencing

platform, with scaffolding informed by chromosomal conformation capture (Hi-C).

2. The final assembly of Dan340 genome was 2,348.72 Mb, including 2,738 contigs

and 2,315 scaffolds with a N50 of 41.49 Mb and 215.35 Mb, respectively.

References

Belton JM, McCord RP, Gibcus JH, Naumova N, Zhan Y, Dekker J. Hi-C: a

comprehensive technique to capture the conformation of genomes. Methods. 2012; 58,

268-276.

Cheng H, Concepcion GT, Feng X, Zhang H, Li H. (2021) Haplotype-resolved de

novo assembly using phased assembly graphs with hifiasm. Nat Methods. 18,

170-175.

Durand NC, Robinson JT, Shamim MS, Machol I, Mesirov JP, Lander ES, Aiden EL.

(2016) Juicebox Provides a Visualization System for Hi-C Contact Maps with

Unlimited Zoom. Cell Syst. 3, 99-101.

Hake S, Ross-Ibarra J. (2015) Genetic, evolutionary and plant breeding insights from

the domestication of maize. Elife. 4, e05861.

Hubert B, Rosegrant M, Boekel MA, Ortiz R. (2010). The future of food: scenarios

for 2050. Crop Sci. 50, 33-50.

Hirsch CN, Hirsch CD, Brohammer AB, Bowman MJ, Soifer I, Barad O, Shem-Tov

D, Baruch K, Lu F, Hernandez AG, Fields CJ, Wright CL, Koehler K, Springer NM,

Buckler E, Buell CR, de Leon N, Kaeppler SM, Childs KL, Mikel MA. (2016) Draft

Assembly of Elite Inbred Line PH207 Provides Insights into Genomic and

Transcriptome Diversity in Maize. Plant Cell. 28, 2700-2714.

Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, Campbell MS, Stein JC, Wei X,

Chin CS, Guill K, Regulski M, Kumari S, Olson A, Gent J, Schneider KL,

Wolfgruber TK, May MR, Springer NM, Antoniou E, McCombie WR, Presting GG,

McMullen M, Ross-Ibarra J, Dawe RK, Hastie A, Rank DR, Ware D. (2017)

Improved maize reference genome with single-molecule technologies. Nature. 546,

524-527.

Li C, Song W, Luo Y, Gao S, Zhang R, Shi Z, Wang X, Wang R, Wang F, Wang J,

Zhao Y, Su A, Wang S, Li X, Luo M, Wang S, Zhang Y, Ge J, Tan X, Yuan Y, Bi X,

He H, Yan J, Wang Y, Hu S, Zhao J. (2019) The HuangZaoSi Maize Genome

Provides Insights into Genomic Variation and Improvement History of Maize. Mol

Plant. 12, 402-409.

Li H, Durbin R. (2010) Fast and accurate long-read alignment with Burrows-Wheeler

transform. Bioinformatics. 26, 589-595.

Lu F, Romay MC, Glaubitz JC, Bradbury PJ, Elshire RJ, Wang T, Li Y, Li Y,

Semagn K, Zhang X, Hernandez AG, Mikel MA, Soifer I, Barad O, Buckler ES.

(2015) High-resolution genetic mapping of maize pan-genome sequence anchors. Nat

Commun. 6, 6914.

Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, Liang C, Zhang J,

Fulton L, Graves TA, Minx P, Reily AD, Courtney L, Kruchowski SS, Tomlinson C,

Strong C, Delehaunty K, Fronick C, Courtney B, Rock SM, Belter E, Du F, Kim K,

Abbott RM, Cotton M, Levy A, Marchetto P, Ochoa K, Jackson SM, Gillam B, Chen

W, Yan L, Higginbotham J, Cardenas M, Waligorski J, Applebaum E, Phelps L,

Falcone J, Kanchi K, Thane T, Scimone A, Thane N, Henke J, Wang T, Ruppert J,

Shah N, Rotter K, Hodges J, Ingenthron E, Cordes M, Kohlberg S, Sgro J, Delgado B,

Mead K, Chinwalla A, Leonard S, Crouse K, Collura K, Kudrna D, Currie J, He R,

Angelova A, Rajasekar S, Mueller T, Lomeli R, Scara G, Ko A, Delaney K, Wissotski

M, Lopez G, Campos D, Braidotti M, Ashley E, Golser W, Kim H, Lee S, Lin J,

Dujmic Z, Kim W, Talag J, Zuccolo A, Fan C, Sebastian A, Kramer M, Spiegel L,

Nascimento L, Zutavern T, Miller B, Ambroise C, Muller S, Spooner W, Narechania

A, Ren L, Wei S, Kumari S, Faga B, Levy MJ, McMahan L, Van Buren P, Vaughn

MW, Ying K, Yeh CT, Emrich SJ, Jia Y, Kalyanaraman A, Hsia AP, Barbazuk WB,

Baucom RS, Brutnell TP, Carpita NC, Chaparro C, Chia JM, Deragon JM, Estill JC,

Fu Y, Jeddeloh JA, Han Y, Lee H, Li P, Lisch DR, Liu S, Liu Z, Nagel DH, McCann

MC, SanMiguel P, Myers AM, Nettleton D, Nguyen J, Penning BW, Ponnala L,

Schneider KL, Schwartz DC, Sharma A, Soderlund C, Springer NM, Sun Q, Wang H,

Waterman M, Westerman R, Wolfgruber TK, Yang L, Yu Y, Zhang L, Zhou S, Zhu

Q, Bennetzen JL, Dawe RK, Jiang J, Jiang N, Presting GG, Wessler SR, Aluru S,

Martienssen RA, Clifton SW, McCombie WR, Wing RA, Wilson RK. (2009) The

B73 maize genome: complexity, diversity, and dynamics. Science. 326, 1112-1115.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. (2015)

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. Bioinformatics. 31, 3210-3212.

Sun S, Zhou Y, Chen J, Shi J, Zhao H, Zhao H, Song W, Zhang M, Cui Y, Dong X,

Liu H, Ma X, Jiao Y, Wang B, Wei X, Stein JC, Glaubitz JC, Lu F, Yu G, Liang C,

Fengler K, Li B, Rafalski A, Schnable PS, Ware DH, Buckler ES, Lai J. (2018)

Extensive intraspecific gene order and gene structural variations between Mo17 and

other maize genomes. Nat Genet. 50, 1289-1295.

AUGUSTUS: ab initio prediction of alternative transcripts. Nucleic Acids Res. 34,

W435-9.

Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA,

Zeng Q, Wortman J, Young SK, Earl AM. (2014) Pilon: an integrated tool for

comprehensive microbial variant detection and genome assembly improvement. PLoS

One. 9(11):e112963.

Yang N, Xu XW, Wang RR, Peng WL, Cai L, Song JM, Li W, Luo X, Niu L, Wang

Y, Jin M, Chen L, Luo J, Deng M, Wang L, Pan Q, Liu F, Jackson D, Yang X, Chen

LL, Yan J. (2017) Contributions of Zea mays subspecies mexicana haplotypes to

modern maize. Nat Commun. 8, 1874.

Zhang R, Xu G, Li J, Yan J, Li H, Yang X. (2018) Patterns of genomic variation in

Chinese maize inbred lines and implications for genetic improvement. Theor Appl

Genet. 131, 1207-1221.

Zhang X, Zhang S, Zhao Q, Ming R, Tang H. (2019) Assembly of allele-aware,

chromosomal-scale autopolyploid genomes based on Hi-C data. Nature Plants. 5,

833-845.

229
230
231   Table 1 Results of genome assembly assisted by Hi-C technology.

	Contig length(bp)	Scaffold length(bp)	Contig number	Scaffold number
Total	2,348,678,871	2,348,721,171	2,738	2,315
Max	97,718,682	343,826,844	−	−
Number >=2000	−	−	2,738	2,315
N50	41,489,332	215,353,106	20	5
N60	32,900,726	193,940,028	26	6
N70	22,291,114	184,974,134	35	7
N80	11,599,268	165,218,414	49	9
N90	3,788,017	153,467,472	80	10

232
233
234