Methylotroph Natural Product Identification by Inverse Stable Isotopic Labeling

Dale A. Cummings Jr., Alice I. Snelling, and Aaron W. Puri*

Department of Chemistry and the Henry Eyring Center for Cell and Genome Science,
University of Utah, Salt Lake City, UT 84112

Keywords: natural products, methylotroph, stable isotope, acyl-homoserine lactone, quorum sensing

*Corresponding author:
Aaron W. Puri
315 S 1400 E Rm 2020
Salt Lake City, UT 84112
USA
(801) 213-1408
a.puri@utah.edu

Abstract

Natural products are an essential source of bioactive compounds. Isotopic labeling is an effective way to identify natural products that incorporate a specific precursor; however, this approach is limited by the availability of isotopically-enriched precursors. We used an inverse stable isotopic labeling approach to identify natural products by growing bacteria on a ^{13}C-carbon source and then identifying ^{12}C-precursor incorporation by mass spectrometry. We applied this approach to methylotrophs, ecologically important bacteria predicted to have significant yet underexplored biosynthetic potential. We demonstrate this method identifies *N*-acyl homoserine lactone quorum sensing signals produced by diverse methylotrophs grown on three one-carbon compounds. We then apply this approach to simultaneously identify five uncharacterized signals produced by a methylotroph, and link these compounds to their synthases. We envision that this method can identify other classes of natural products synthesized by methylotrophs and other organisms that grow on inexpensive and readily available ^{13}C-carbon sources.

Introduction

Natural products are an important source of small molecule therapeutics (Newman and Cragg, 2020), agricultural compounds (Cantrell et al., 2012), and other bioactive metabolites. Precursor feeding studies are commonly used to determine the biosynthetic route to a compound of interest, and can also be used to identify previously undetected compounds produced by an organism (Gross et al., 2007; Kinnel et al., 2017; May et al., 2020; Schaefer et al., 2018). However, this approach can be limited by the synthetic or commercial availability of an isotopically enriched precursor. Alternatively, in an inverse stable isotopic labeling approach, a fully ^{13}C-labeled organism is fed a ^{12}C-biosynthetic precursor to identify natural products that incorporate this precursor. This incorporation can be detected by a negative shift in the mass-to-charge ratio (*m/z*) consistent with incorporation of all or part of the ^{12}C-precursor. With this approach researchers can use any precursor without having to synthesize or purchase a ^{13}C-labeled version. Inverse stable isotopic labeling has been previously used to help determine the biosynthetic origins of the cofactor pyrroloquinoline quinone (Kleef and Duine, 1988) and an isocyanide-containing antibiotic (Brady and Clardy, 2005). However, to our knowledge, this approach has not been used to systematically identify previously uncharacterized natural products.

Inverse stable isotopic labeling is particularly well-suited for bacteria that can grow on inexpensive and readily available ^{13}C carbon sources, such as methylotrophs. Methylotrophs are bacteria that grow on reduced compounds with no carbon-carbon bonds, such as methane, methanol, methylamine, and dimethyl sulfide (Chistoserdova et al., 2009). These organisms play important roles in biogeochemical cycling and bioremediation (Nisbet et al., 2014; Singh et al., 2010; Stein and Klotz, 2016), as well as plant-microbe interactions (Fedorov et al., 2011). Genomic analysis also predicts that methylotrophs possess the biosynthetic potential to produce myriad natural products that have not yet been identified (Puri, 2019). Identifying and characterizing these molecules can help scientists understand how these important organisms interact with each other and their environment, discover new drug leads, and help synthetic biologists optimize the desirable activities of methylotrophs using exogenous small molecules.

A benefit of working with methylotrophs is that ^{13}C-labeled versions of their one-carbon growth substrates are relatively inexpensive. For this reason, a methylotroph was used to create ^{13}C-labeled nucleic acids from ^{13}C-methanol for DNA structural studies by NMR spectroscopy (Batey et al., 1996). Stable isotope labeling experiments have also been repeatedly used to identify active methylotrophs in the environment (Kalyuzhnaya et al., 2008; Radajewski et al., 2000). We sought to take advantage of this benefit to aid in the discovery of methylotroph natural products.

As a proof of concept, we applied this method to identify quorum sensing (QS) signals produced by methylotrophs. QS is a form of chemical communication used by bacteria to regulate gene expression in a cell density-dependent manner (Papenfort and Bassler, 2016; Whiteley et al., 2017). In one well-characterized form of QS, Gram-negative proteobacteria use *N*-acyl-homoserine lactones (acyl-HSLs) produced by LuxI-family synthases. Acyl-HSL signals vary in their acyl chain, but all possess a common homoserine lactone derived from methionine via *S*-adenosyl-*L*-methionine (Moré et al., 1996; Schaefer et al., 1996) (Figure 1A). These signals are detected by LuxR-family receptors, which are also transcription factors that regulate gene expression upon signal binding. Methylotrophs often possess QS systems, and several strains have three or more annotated *luxI*-family synthase genes in their genome (Marx et al., 2012; Poonguzhali et al., 2007). The majority of these genes are predicted to encode CoA-utilizing enzymes known to produce noncanonical signals (Liao et al., 2018). However, relatively few methylotroph acyl-HSL signals have been characterized, and many cannot be predicted based on the amino acid sequences of their synthases (Puri, 2019).

Acyl-HSL quorum sensing signals are often detected using reporter assays, in which a strain is engineered so that the LuxR-family receptor activates the expression of a reporter gene upon signal binding (Thornhill and McLean, 2018). However, signal detection is dependent on the specificity of the receptor used, which can lead to false negatives. Reporter assays are often paired with thin layer chromatography to correlate detected signals with the retention factors of known standards, which can be

84 inaccurate. Liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS) can also be used
85 to identify acyl-HSLs based on known fragmentation patterns (Morin et al., 2003; Patel et al., 2016), but
86 access to these instruments may still limit routine analysis using this method, and fragmentation patterns
87 may differ for some acyl-HSLs. For example, the aryl-HSL *p*-coumaroyl-HSL does not produce the

Figure 1. (A) Acyl-HSL biosynthesis incorporates methionine. *L*-Met: *L*-methionine. MetK: *S*-adenosyl-*L*-methionine synthase. SAM: *S*-adenosyl-*L*-methionine. MTA: 5'-methylthioadenosine. (B) The inverse stable isotope labeling approach applied to acyl-HSL signals. Note that the acyl carbonyl carbon is not denoted as ^{12}C or ^{13}C as it can be derived from endogenous or exogenous sources and therefore may or may not be labeled when an organism is grown on a ^{13}C-carbon source.

88 characteristic *m/z* 102 HSL fragment (Schaefer et al., 2008).
89  Detection of methionine incorporation into the common HSL portion of acyl-HSLs is a more
90 generalizable method that has been successfully used to detect QS signals (Ahlgren et al., 2011; Eberhard
91 et al., 1991; Lindemann et al., 2011; Schaefer et al., 2018). ^{14}C-labeled methionine can be used in feeding
92 studies to indirectly identify signals by comparing radioactivity retention via HPLC to the retention times
93 (RTs) of known acyl-HSL standards. We sought to build off of this approach by using an inverse stable
94 isotopic labeling method with ^{12}C-methionine to identify QS signals produced by methylotrophs (Figure
95 1B). We show this method works with three ^{13}C-labeled one-carbon sources (methane, methanol, and
96 methylamine) in two diverse methylotrophs. We then apply the method to simultaneously identify five
97 QS signals produced by *Methylorubrum rhodinum* DSM2163 (Green and Ardley, 2018), including three
98 which could not be predicted based on sequence homology to characterized QS systems. Finally, we link
99 these signals to the synthase genes responsible for their production.
100
101 **Results**
102
103 *Inverse stable isotopic labeling identifies previously characterized acyl-HSL signals*
104
105  To verify the utility of the inverse stable isotopic labeling approach for identifying natural
106 products, we began by applying it to a methylotroph that produces acyl-HSL signals that have been

previously characterized. The methanol-oxidizing *Methylorubrum extorquens* PA1 produces a single
LuxI-family signal synthase, MlaI, with 100% amino acid sequence identity to the characterized version
found in *Methylorubrum extorquens* AM1 (Marx et al., 2012; Nieto Penalver et al., 2006). Vorholt and
coworkers previously demonstrated that MlaI produces two QS signals with unsaturated acyl chains: *N*-
(2-*trans*-7-*cis*-tetradecenoyl)-*L*-homoserine lactone (referred to as 2*E*,7*Z*-$C_{14:2}$-HSL) and *N*-(7-*cis*-
tetradecenoyl)-*L*-homoserine lactone (7*Z*-$C_{14:1}$-HSL) (Nieto Penalver et al., 2006) (Figure 2).
113

C$_8$-HSL

3-OH-C$_8$-HSL

3-oxo-C$_8$-HSL

3-OH-C$_{10}$-HSL

2*E*,7*Z*-C$_{14:2}$-HSL

7*Z*-C$_{14:1}$-HSL

Figure 2. Acyl-HSLs identified in this work. Stereochemistry was not determined and is inferred based on previous studies. All biologically produced acyl-HSLs that have been characterized have homoserine lactones with an *L* stereocenter. The hydroxyls in 3-OH-C$_8$-HSL and 3-OH-C$_{10}$-HSL are shown as 3*R* because their synthases are annotated as acyl carrier protein (ACP)-linked (KEGG Orthology term K13060) and these enzymes use acyl-ACPs from fatty acid biosynthesis (Hoang et al., 2002; Val and Cronan, 1998) where the stereoselective β-ketoacyl acyl carrier protein reductase FabG produces 3*R*-OH fatty acyl chains (Volpe and Vagelos, 1976).

114
In order to streamline the inverse stable isotopic labeling procedure for future screening of
diverse bacteria with potentially different growth rates, we chose to add the ^{12}C-methionine precursor at
the beginning of growth. We grew the *M. extorquens* PA1Δ*cel* strain CM2730 (referred to as PA1)
(Delaney et al., 2013) with ^{13}C-methanol and different concentrations of ^{12}C-methionine. We extracted the
supernatant with acidified ethyl acetate, analyzed the extract by LC-MS, and detected features using the
program MZmine2 (Pluskal et al., 2010). We identified two features with *m/z* values of 322 and 324,

which correspond to protonated versions of 2E,7Z-$C_{14:2}$-HSL and 7Z-$C_{14:1}$-HSL, respectively, that have
incorporated the four carbons of ^{12}C-methionine but are otherwise fully ^{13}C-labeled (Figure S1 and Table
1). When we cultured PA1 with ^{12}C-methanol as a control, we identified features with the same RTs but
with m/z values of 308 and 310, which correspond to the protonated ^{12}C-versions of 2E,7Z-$C_{14:2}$-HSL and
7Z-$C_{14:1}$-HSL, respectively. Together, these results show that the inverse labeling approach can be used to
correctly identify acyl-HSLs.
The methionine titration indicated that the optimal concentration of the ^{12}C-methionine precursor
was 0.5 mM when added at the beginning of growth (Figure S1). Higher-than-necessary label
concentrations may result in slowed growth or nonspecific incorporation of ^{12}C-precursor carbons due to
toxicity or methionine metabolism, respectively. We did not observe any fully ^{13}C-labeled acyl-HSLs
using our workflow upon addition of ^{12}C-methionine at any concentration we tested. This indicates that
for acyl-HSL natural products it is necessary to include a separate ^{13}C-carbon source condition with no
precursor as a benchmark to identify ^{12}C-enriched features in the ^{13}C-carbon source + ^{12}C-methionine
condition.
135

Table 1. Identification of acyl-HSL signals using the inverse stable isotopic labeling approach. Max paired m/z refers to the maximum feature m/z observed in that condition where the desired m/z difference of four units was observed compared to the other ^{13}C-condition. Note the RT differences between PA1 and DSM2163 for 2E,7Z-$C_{14:2}$-HSL and 7Z-$C_{14:1}$-HSL were resolved when smaller injection volumes were used after initial screening (see Figure S3).

Strain	Carbon source	^{12}C-carbon source		^{13}C-carbon source		^{13}C-carbon source + ^{12}C-methionine		# Pairs detected	Identified Acyl-HSL
		RT (min)	m/z	RT (min)	Max paired m/z	RT (min)	Max paired m/z		
M. extorquens PA1	Methanol	23.0	310	23.0	328	23.1	324	6	7Z-$C_{14:1}$-HSL
		22.7	308	22.7	326	22.8	322	5	2E,7Z-$C_{14:2}$-HSL
	Methylamine	23.1	310	23.1	327	23.2	323	5	7Z-$C_{14:1}$-HSL
		22.8	308	22.8	325	22.9	321	5	2E,7Z-$C_{14:2}$-HSL
M. tundripaludum 21/22	Methane	15.6	272	15.6	286	15.6	282	1	3-OH-C_{10}-HSL
M. rhodinum DSM2163	Methanol	22.9	308	22.9	326	22.8	322	5	2E,7Z-$C_{14:2}$-HSL
		23.2	310	23.2	328	23.1	324	5	7Z-$C_{14:1}$-HSL
		12.0	244	12.0	256	12.0	252	4	3-OH-C_8-HSL
		15.6	228	15.5	240	15.5	236	3	C_8-HSL
		12.7	242	12.7	254	12.7	250	1	3-oxo-C_8-HSL

In addition to detecting fully ^{13}C-labeled acyl-HSLs in the PA1 culture grown on ^{13}C-methanol
without addition of the ^{12}C-precursor, we also detected several features with the same RT but stepwise
decreases of one m/z unit. We also identified the same spectral pattern in the ^{13}C-methanol + ^{12}C-
methionine condition, but decreased by four m/z units, corresponding to ^{12}C-methionine incorporation
(Figure S2). Many methylotrophs in the alphaproteobacteria class, including *M. extorquens*, assimilate
reduced one-carbon compounds such as methanol using the serine cycle (Chistoserdova et al., 2009). In
the serine cycle, half of the carbon input is derived from carbon dioxide, which we did not label in our
system and could therefore result in incomplete ^{13}C-labeling of metabolites of interest. The fact that we do
observe fully ^{13}C-labeled acyl-HSLs is likely because ^{13}C-carbon dioxide is rapidly produced by the
bacterium's central metabolism. When we grew PA1 on the carbon source ^{13}C-methylamine, we did not
identify complete ^{13}C-labeling but this did not affect our ability to identify a decrease of four m/z units
corresponding with ^{12}C-methionine incorporation (Table 1) because the extent of acyl-HSL ^{13}C-labeling
was consistent between the ^{13}C-methylamine and ^{13}C-methylamine + ^{12}C-methionine conditions.
We wrote a Python script to identify pairs of features with matching RTs and a decrease of four
m/z units in the ^{13}C-carbon source + ^{12}C-methioine condition compared to the ^{13}C-carbon source condition
(see Methods). For the RTs corresponding to 2*E*,7*Z*-C$_{14:2}$-HSL and 7*Z*-C$_{14:1}$-HSL, our script identifies
several pairs of features with the desired m/z difference due to the aforementioned incomplete labeling
(Table 1). Notably, out of all the feature pairs identified in the PA1 culture extract, the two acyl-HSL
signals have the most identified pairs that correspond with a single feature at the same RT in the ^{12}C-
carbon source control (Table S1). We can therefore use this knowledge to prioritize identified features
that are more likely to be acyl-HSLs.
In order to determine if this method is more broadly applicable, we applied it to the methane-
oxidizing bacterium *Methylobacter tundripaludum* 21/22 (21/22), which was previously shown to
produce the QS signal *N*-(3-hydroxydecanoyl)-*L*-homoserine lactone (3-OH-C$_{10}$-HSL) (Puri et al., 2017)
(Figure 2). When we grew 21/22 with ^{13}C-methane, we identified a feature with an m/z value of 286,
corresponding to the protonated and fully ^{13}C-labeled 3-OH-C$_{10}$-HSL. We detected a feature with the
same RT and a decrease of four m/z units in the ^{13}C-methane + ^{12}C-methionine condition (Table 1),
indicating ^{12}C-methionine incorporation. These results show that this method can be broadly applied, as it
works with the methane-oxidizing gammaproteobacterium 21/22 as well as the methanol- and
methylamine-oxidizing alphaproteobacterium PA1.
167
168   *Simultaneous identification of five acyl-HSL signals produced by the methylotroph* M. rhodinum
169   *DSM2163*
170
Next, we applied the inverse stable isotopic labeling method to identify the signals produced by
the methanol-oxidizer *M. rhodinum* DSM2163 (DSM2163), which possesses three predicted LuxI-family
acyl-HSL synthases. A previous study used a reporter assay to speculate which acyl-HSL signals are
produced by this strain, but the exact signals were not identified (Poonguzhali et al., 2007). We identified
several features with the signature decrease of four m/z units when DSM2163 was grown on ^{13}C-methanol
+ ^{12}C-methionine compared to ^{13}C-methanol alone. Three of these features corresponded to the signals *N*-
octanoyl-*L*-homoserine lactone (C$_8$-HSL), *N*-(3-hydroxyoctanoyl)-*L*-homoserine lactone (3-OH-C$_8$-HSL),
and *N*-(3-oxooctanoyl)-*L*-homoserine lactone (3-oxo-C$_8$-HSL) (Figure 2 and Table 1), which we
confirmed by high-resolution mass spectrometry (Table S2) as well as by verifying these compounds have
identical LC-MS retention times to commercial standards (Figures S3A-C).
Our method also identified features in the DSM2163 culture corresponding to the signals 2*E*,7*Z*-
C$_{14:2}$-HSL and 7*Z*-C$_{14:1}$-HSL, and these signals had identical LC-MS retention times (Figures S3D and
S3E) and high-resolution MS/MS spectra (Table S3) compared to the signals produced by PA1. This is
consistent with the fact that one of the DSM2163 acyl-HSL synthases shares 91% amino acid identity
with MlaI from PA1. We were therefore able to use the inverse labeling method to simultaneously
identify a total of five acyl-HSL QS signals produced by the methylotroph DSM2163.
187

Figure 3. Linking *M. rhodinum* DSM2163 LuxI-family acyl-HSL synthases with their products. Extracted ion chromatograms of supernatant extracts for the listed strains for the *m/z* ranges (A) 228.0-228.5, corresponding to C_8-HSL, (B) 242.0-242.5, corresponding to 3-oxo-C_8-HSL, (C) 244.0-244.5, corresponding to 3-OH-C_8-HSL, (D) 308.0-308.5, corresponding to $2E,7Z$-$C_{14:2}$-HSL, and (E) 310.0-310.5, corresponding to $7Z$-$C_{14:1}$-HSL.

188
189 *Heterologous expression links acyl-HSL signals with their synthases*
190
191   Finally, we constructed a strain for heterologous expression of *luxI*-family acyl-HSL synthase
192 genes to link synthases with their signal products. We created an unmarked deletion strain of PA1 called
193 AWP227 that no longer produces acyl-HSL signals by knocking out the acyl-HSL receptor gene *mlaR*
194 and the synthase gene *mlaI*. We then used AWP227 to express each of the three DSM2136 synthase genes
195 separately on a plasmid to determine which signals are produced by each synthase.
196
197

When we heterologously expressed the DSM2163 synthase gene Ga0373200_3300, we observed
features corresponding to production of C_8-HSL (Figure 3A) and 3-oxo-C_8-HSL (Figure 3B) that were not
present in the no-plasmid AWP227 control. Heterologous expression of Ga0373200_1920 also resulted in
production of C_8-HSL (Figure 3A), as well as 3-OH-C_8-HSL (Figure 3C). We also confirmed that the
synthase with 91% amino acid identity to MlaI (Ga0373200_956) produces the signals $2E,7Z$-$C_{14:2}$-HSL
and $7Z$-$C_{14:1}$-HSL (Figures 3D and 3E). All gene locus tags refer to the Joint Genome Institute Integrated
Microbial Genomes & Microbiomes data management system (JGI IMG/M) (Chen et al., 2021). Vorholt
and coworkers previously named a LuxI-family synthase that produces N-hexanoyl-L-homoserine lactone
(C_6-HSL) and C_8-HSL MsaI for <u>M</u>ethylobacterium <u>s</u>hort-chain <u>a</u>cyl-HSLs (Nieto Penalver et al., 2006).
We therefore name Ga0373200_1920 MsaI2 and Ga0373200_3300 MsaI3 based on our results, and also
refer to Ga0373200_956 as MlaI in *M. rhodinum* DSM2163.
209
210   **Discussion**
211
We used an inverse stable isotopic labeling approach to rapidly identify acyl-HSL natural
products made by methylotrophic bacteria. This method was effective using three different ^{13}C-one-
carbon sources (methane, methanol, and methylamine) and for diverse methylotrophs including both a
methane-oxidizing gammaproteobacterium and two species of alphaproteobacteria. Together these results
show the broad applicability of this method for identifying natural products by growing different bacteria
on a ^{13}C-carbon source and feeding readily available ^{12}C-precursors.
The inverse stable isotopic labeling method can be applied in the future to identify other classes
of natural products based on a common precursor, including those which may not be readily available in
an isotopically labeled form. Feeding studies using isotopically enriched precursors are extremely
widespread, and the inverse stable isotopic labeling method is applicable in any situation where a
traditional method can be used. However, this method is especially well-suited for organisms that grow
on inexpensive ^{13}C-carbon sources, including methanotrophs and autotrophs such as cyanobacteria, which
are known to have significant biosynthetic potential (Dittmann et al., 2015).
A strength of stable isotopic labeling is that it can be performed using unit-resolution mass
spectrometers, which are widely available to researchers for routine analysis. This approach complements
existing LC-MS/MS based methods for acyl-HSL identification, as fragmentation patterns may change
based on signal structure. Notably, the characteristic *m/z* 102 HSL fragment was not one of the top twenty
signals detected in the tandem mass spectra of $2E,7Z$-$C_{14:2}$-HSL, likely because it contains an α, β
unsaturated acyl chain (Table S3). More broadly, the inverse stable isotopic labeling approach can also be
applied using LC-MS/MS and techniques such as feature-based molecular networking (Klitgaard et al.,
2015; Nothias et al., 2020) to identify precursor incorporation into specific molecular fragments. The
combination of these techniques in the future will enable researchers to gain richer information about
identified natural products early in the discovery process.
The increase in metagenomic sequencing of microbial communities has led to an increased
demand for predicting how these organisms interact in the environment. Methods for rapidly determining
the QS signal(s) produced by an organism enable researchers to link these compounds to specific gene
sequences, and in turn to hypothesize how these bacteria are interacting *in situ*. The acyl-HSL signals
produced by MsaI2 and MsaI3 could not be predicted on the basis of amino acid sequence alone. MsaI2
shares only 58% amino acid identity with RaiI from *Rhizobium etli* ISP42, which was reported to produce
C_8-HSL and 3-OH-C_8-HSL (Pérez-Montaño et al., 2011). MsaI3 does not share >50% amino acid identity
with any currently characterized LuxI-family synthases. During heterologous expression of MsaI2 and
MsaI3 we determined that both synthases produce significant amounts of C_8-HSL (Figures 3 and S3). It is
therefore unclear if one or both of these synthases are responsible for the C_8-HSL detected in the
DSM2163 culture. Together, these annotations will aid in the future prediction of QS signal production
from bacterial genome sequences. Furthermore, our heterologous expression strain can be used in the
future to identify QS signals produced by synthases from non-methylotrophic species using the inverse
labeling approach.

249
250 **Significance**
251
Natural products are a vetted source of bioactive compounds with medicinal and agricultural value.
Isotopic labeling is often used to identify and characterize natural products, however the labeled
precursors required for these studies can be difficult to obtain commercially or synthetically. Here we use
an inverse stable isotopic labeling approach to identify natural products via incorporation of unlabeled
precursors, which significantly expands precursor availability for natural product studies. This approach is
broadly applicable to bacteria that grow on inexpensive labeled carbon sources such as methylotrophs,
and can be used to rapidly identify natural products known to incorporate a particular precursor.
259

Methods

Key Reagents

^{13}C-labeled methanol, methane, and methylamine were purchased from Cambridge Isotope Laboratories. C_8-HSL was purchased from Millipore Sigma. All other acyl-HSLs were purchased from Cayman Chemical.

Routine bacterial culturing

Strains used in this study are listed in Table S4. *Escherichia coli* strains were grown in lysogeny broth (LB) at 37°C. *M. extorquens* PA1 and *M. rhodinum* DSM2163 were grown at 30 °C in modified ammonium mineral salts (AMS) medium (Whittenbury et al., 1970), with the addition of 0.1% (m/v) yeast extract for strain DSM2163. AMS contains 0.2 g/L $MgSO_4 \cdot 7H_2O$, 0.2 g/L $CaCl_2 \cdot 6H_2O$, 0.5 g/L NH_4Cl, 30 µM $LaCl_3$, and 1X trace elements. 500X trace elements contains 1.0 g/L Na_2-EDTA, 2.0 g/L $FeSO_4 \cdot 7H_2O$, 0.8 g/L $ZnSO_4 \cdot 7H_2O$, 0.03 g/L $MnCl_2 \cdot 4H_2O$, 0.03 g/L H_3BO_3, 0.2 g/L $CoCl_2 \cdot 6H_2O$, 0.6 g/L $CuCl_2 \cdot 2H_2O$, 0.02 g/L $NiCl_2 \cdot 6H_2O$, and 0.05 g/L $Na_2MoO \cdot 2H_2O$. Final concentrations of 4 mM phosphate buffer pH 6.8 and 50 mM ^{12}C- or ^{13}C-methanol were added prior to use and cultures were shaken at 200 rpm. *M. tundripaludum* 21/22 was grown in modified nitrate mineral salts (NMS) medium (Whittenbury et al., 1970), which is the same as AMS with 1 g/L KNO_3 substituted for the NH_4Cl. *M. tundripaludum* 21/22 was cultured at room temperature (22-24°C) in an atmosphere of 50% (v/v) methane in air. For routine culturing, plates were incubated in sealed jars while liquid cultures were grown in 18- by 150-mm tubes sealed with rubber stoppers and aluminum seals shaken at 200 rpm.

Plasmid construction

Plasmids used in this study are listed in Table S5. Primers used in this study are listed in Table S6. All plasmids were constructed using Gibson Assembly (Gibson et al., 2009) and selection was performed with kanamycin (50 µg/mL) both in *E. coli* and *M. extorquens* PA1 strains.

Genetic manipulation

Genetic manipulation of strain PA1Δ*cel* (CM2730) and the derivative strain AWP227 was performed at 30°C. Verified plasmids were conjugated into these strains using the *E. coli* donor S17-1 (Simon et al., 1983). 500 µL of exponentially growing cultures (OD 0.4-0.6) of the donor and recipient strains were pelleted at 16,100 rcf for one minute and resuspended in 500 µL sterile ultrapure H_2O. These strains were then pelleted again and the two pellets were combined in a total volume of 50 µL sterile ultrapure H_2O. Next, the entire mixture was spotted onto an AMS agar plate containing 50 mM methanol and 10% (v/v) nutrient broth and incubated for two days. Successful conjugants were selected on AMS plates containing kanamycin (50 µg/mL). To construct the unmarked deletion mutant AWP227, kanamycin-resistant integrants (single crossovers) were restreaked and then plated on an AMS plate containing 50 mM methanol and 1% (m/v) sucrose for counterselection. The resulting colonies were screened for double crossovers by kanamycin sensitivity and colony PCR before the final mutant was verified by Sanger sequencing.

Inverse labeling experiments

Exponentially growing bacterial cultures were pelleted at 16,100 rcf for one minute and resuspended in growth medium with no carbon source. Subsequently, three separate six milliliter cultures were inoculated with the resuspended strain at a starting OD of 0.02. The ^{12}C-carbon source was added to one culture, the ^{13}C-carbon source to the second, and the ^{13}C-carbon source plus 500 nM ^{12}C-methionine to the last culture. The carbon sources used were 50 mM methanol, 50 mM methylamine, or 50% (v/v) methane. Cultures were grown until reaching stationary phase (OD of approximately 0.8) and then were centrifuged at 4,800 rcf for ten minutes. The resulting supernatant was extracted twice with an equal volume of ethyl acetate containing 0.01% acetic acid, and the combined organic extract was evaporated to dryness using a nitrogen stream and stored at -20°C until analysis by LC-MS.

311
LC-MS for acyl-HSL signal detection
Dried culture supernatant extracts were resuspended in 200 microliters of 1:1 water:acetonitrile, and
subsequently 65 microliters were injected onto an Agilent 1260 Infinity liquid chromatography system
connected to an Agilent 6120 single quadrupole mass spectrometer operating with positive polarity and a
mass range of 150-1500 *m/z*. A Waters Xselect HSS T3 column (2.5 μm particle size, 2.1 mm x 50 mm)
held at 30 ºC was used for reverse phase separation with a flow rate of 0.4 mL/min. Solvent A: Water +
0.1 % formic acid, Solvent B: Acetonitrile + 0.1% formic acid. Gradient: 0-2 min, 0% B. 2-32 min, 0-
100% B. 32-35 min, 100% B. 35-36 min, 100-0% B. 36-38 min, 0% B. The limit of detection for known
standards on this setup was 125 nM (Figure S4), which corresponds to a limit of approximately 5 nM in
the original bacterial culture.
322
LC-MS analysis
Raw data files in netCDF format were exported using Agilent OpenLab CDS (rev C.01.07). Features were
detected using MZmine version 2.53 (Pluskal et al., 2010) using the following workflow: 1. Mass
detection (centroid, noise level 1.0E3). 2. ADAP chromatogram builder (minimum group size 5 scans,
group intensity threshold 1.0E3, min highest intensity 5.0E3), *m/z* tolerance 0.3) (Myers et al., 2017). 3.
Chromatogram deconvolution (local minimum search, chromatogram threshold 30%, search minimum
0.1 min, minimum relative height 10%, minimum absolute height 6.0E3, minimum ratio of peak top/edge
2, peak duration 0-2 min). 4. Adduct search (RT tolerance 0.1 min, adducts [M+Na-H] and [M+NH$_3$]
selected, *m/z* tolerance 0.2, max relative peak height 200%). 5. Feature list rows filter (remove identified
adducts). Subsequently, isotopes were removed from ^{12}C samples using the Isotopic peaks grouper (*m/z*
tolerance 0.2, retention time tolerance 0.1 min, monotonic shape required, maximum charge 3,
representative isotope most intense), and the three feature lists were aligned in the order ^{13}C-carbon
source + ^{12}C-methionine, ^{12}C-carbon source, ^{13}C-carbon source using the Join aligner (*m/z* tolerance 0.3,
weight of *m/z* 50, retention time tolerance 0.1 min, weight for retention time 50). The alignment was
exported in .csv format with the row retention time as a common element and peak *m/z* as the data file
element. Features containing the desired four *m/z* unit difference in the ^{13}C-carbon source and ^{13}C-carbon
source + ^{12}C-methionine samples were then detected using a custom Python script (available at
https://github.com/purilab/inverse).
341
HRMS
Mass spectrometry data were collected using a Waters Acquity I-class ultra-high pressure liquid
chromatograph coupled to a Waters Xevo G2-S quadrupole time-of-flight mass spectrometer. An Acquity
UPLC BEH C18 column (2.1 x 50 mm) was used for separation and resolving samples. Solvent A: Water
+ 0.1 % formic acid, Solvent B: Acetonitrile + 0.1% formic acid. The sample was eluted from the column
using a ten minute linear solvent gradient: 0-0.1 min, 1% B; 0.1 - 10 min, 100% B. The solvent flow rate
was 0.45 mL per minute. Mass spectra were collected in positive ion mode, with following parameters: 3
kV capillary voltage; 25 V sampling cone voltage; 150 °C source temperature; 500 °C desolvation
temperature; nitrogen desolvation at 800 L/hr. The fragmentation spectra were collected using the same
parameters with a 10-25 eV collision energy ramp. The lockspray solution was 200 pg/μL leucine
enkephalin. The lockspray flow rate was 6 μL/min. Sodium formate was used to calibrate the mass
spectrometer. The acquired mass spectra were processed using Masslynx 4.1 software.
354
355
356 **Acknowledgements**
357

358 This work is dedicated to Mary E. Lidstrom (University of Washington) for her 70[th] birthday. This work
359 was supported by National Institutes of Health grant R00 GM118762 (to AWP). We thank members of
360 the Puri Lab for reading the manuscript and for helpful discussions. We thank Yanfen Fu (Facebook, Inc.)
361 for useful discussions about methylotroph metabolism and isotopic labeling that led to this project. We

thank N. Cecilia Martinez-Gomez (UC Berkeley) for *M. extorquens* PA1 Δ*cel* strain CM2730, and Ming Hammond (University of Utah) for generously sharing her LC-MS system and other lab equipment. We thank Daniel Petras (University of Tübingen) for sharing Python code and for helpful discussions about LC-MS analysis.

Author Contributions

AWP and DAC designed the experiments. DAC, AIS, and AWP performed the experiments. AWP wrote the manuscript. AWP and DAC edited the manuscript. All authors read and approved of the final version of the manuscript.

Conflicts of Interest

The authors declare no conflicts of interest.

References

379 **References**
380
Ahlgren, N.A., Harwood, C.S., Schaefer, A.L., Giraud, E., and Greenberg, E.P. (2011). Aryl-homoserine
lactone quorum sensing in stem-nodulating photosynthetic bradyrhizobia. Proc. Natl. Acad. Sci. *108*,
7183–7188.

Batey, R.T., Cloutier, N., Mao, H., and Williamson, J.R. (1996). Improved large scale culture of
Methylophilus methylotrophus for ^{13}C/^{15}N labeling and random fractional deuteration of ribonucleotides.
Nucleic Acids Res. *24*, 4836–4837.

Brady, S.F., and Clardy, J. (2005). Systematic Investigation of the *Escherichia coli* Metabolome for the
Biosynthetic Origin of an Isocyanide Carbon Atom. Angew. Chem. Int. Ed. *44*, 7045–7048.

Cantrell, C.L., Dayan, F.E., and Duke, S.O. (2012). Natural products as sources for new pesticides. J. Nat.
Prod. *75*, 1231–1242.

Chen, I.-M.A., Chu, K., Palaniappan, K., Ratner, A., Huang, J., Huntemann, M., Hajek, P., Ritter, S.,
Varghese, N., Seshadri, R., Roux, S., Woyke, T., Eloe-Fadrosh, E.A., Ivanova, N.N., and Kyrpides, N.C.
(2021). The IMG/M data management and analysis system v.6.0: new tools and advanced capabilities.
Nucleic Acids Res. *49*, D751–D763.

Chistoserdova, L., Kalyuzhnaya, M.G., and Lidstrom, M.E. (2009). The Expanding World of
Methylotrophic Metabolism. Annu. Rev. Microbiol. *63*, 477–499.

Delaney, N.F., Kaczmarek, M.E., Ward, L.M., Swanson, P.K., Lee, M.-C., and Marx, C.J. (2013).
Development of an optimized medium, strain and high-throughput culturing methods for
Methylobacterium extorquens. PloS One *8*, e62957.

Dittmann, E., Gugger, M., Sivonen, K., and Fewer, D.P. (2015). Natural Product Biosynthetic Diversity
and Comparative Genomics of the Cyanobacteria. Trends Microbiol. *23*, 642–652.

Eberhard, A., Longin, T., Widrig, C.A., and Stranick, S.J. (1991). Synthesis of the *lux* gene autoinducer in
Vibrio fischeri is positively autoregulated. Arch. Microbiol. *155*, 294–297.

Fedorov, D.N., Doronina, N.V., and Trotsenko, Y.A. (2011). Phytosymbiosis of aerobic methylobacteria:
New facts and views. Microbiology *80*, 443–454.

Gibson, D.G., Young, L., Chuang, R.-Y., Venter, J.C., Hutchison, C.A., and Smith, H.O. (2009).
Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat. Methods *6*, 343–345.

Green, P.N., and Ardley, J.K. (2018). Review of the genus *Methylobacterium* and closely related
organisms: a proposal that some *Methylobacterium* species be reclassified into a new genus,
Methylorubrum gen. nov. Int. J. Syst. Evol. Microbiol. *68*, 2727–2748.

Gross, H., Stockwell, V.O., Henkels, M.D., Nowak-Thompson, B., Loper, J.E., and Gerwick, W.H.
(2007). The Genomisotopic Approach: A Systematic Method to Isolate Products of Orphan Biosynthetic
Gene Clusters. Chem. Biol. *14*, 53–63.

Hoang, T.T., Sullivan, S.A., Cusick, J.K., and Schweizer, H.P. (2002). β-Ketoacyl acyl carrier protein
reductase (FabG) activity of the fatty acid biosynthetic pathway is a determining factor of 3-oxo-
homoserine lactone acyl chain lengths. Microbiology *148*, 3849–3856.

Kalyuzhnaya, M.G., Lapidus, A., Ivanova, N., Copeland, A.C., McHardy, A.C., Szeto, E., Salamov, A.,
Grigoriev, I.V., Suciu, D., Levine, S.R., Markowitz, V.M., Rigoutsos, I., Tringe, S.G., Bruce, D.C.,
Richardson, P.M., Lidstrom, M.E., and Chistoserdova, L. (2008). High-resolution metagenomics targets
specific functional types in complex microbial communities. Nat. Biotechnol. *26*, 1029–1034.

Kinnel, R.B., Esquenazi, E., Leao, T., Moss, N., Mevers, E., Pereira, A.R., Monroe, E.A., Korobeynikov,
A., Murray, T.F., Sherman, D., Gerwick, L., Dorrestein, P.C., and Gerwick, W.H. (2017). A
Maldiisotopic Approach to Discover Natural Products: Cryptomaldamide, a Hybrid Tripeptide from the
Marine Cyanobacterium *Moorea producens*. J. Nat. Prod. *80*, 1514–1521.

Kleef, M.A.G. van, and Duine, J.A. (1988). L-tyrosine is the precursor of PQQ biosynthesis in
Hyphomicrobium X. FEBS Lett. *237*, 91–97.

Klitgaard, A., Nielsen, J.B., Frandsen, R.J.N., Andersen, M.R., and Nielsen, K.F. (2015). Combining
Stable Isotope Labeling and Molecular Networking for Biosynthetic Pathway Characterization. Anal.
Chem. *87*, 6520–6526.

Liao, L., Schaefer, A.L., Coutinho, B.G., Brown, P.J.B., and Greenberg, E.P. (2018). An aryl-homoserine
lactone quorum-sensing signal produced by a dimorphic prosthecate bacterium. Proc. Natl. Acad. Sci.
115, 7587–7592.

Lindemann, A., Pessi, G., Schaefer, A.L., Mattmann, M.E., Christensen, Q.H., Kessler, A., Hennecke, H.,
Blackwell, H.E., Greenberg, E.P., and Harwood, C.S. (2011). Isovaleryl-homoserine lactone, an unusual
branched-chain quorum-sensing signal from the soybean symbiont *Bradyrhizobium japonicum.* Proc.
Natl. Acad. Sci. *108*, 16765–16770.

Marx, C.J., Bringel, F., Chistoserdova, L., Moulin, L., Farhan Ul Haque, M., Fleischman, D.E., Gruffaz,
C., Jourand, P., Knief, C., Lee, M.-C., et al. (2012). Complete genome sequences of six strains of the
genus *Methylobacterium*. J. Bacteriol. *194*, 4746–4748.

May, D.S., Crnkovic, C.M., Krunic, A., Wilson, T.A., Fuchs, J.R., and Orjala, J.E. (2020). [15]N Stable
Isotope Labeling and Comparative Metabolomics Facilitates Genome Mining in Cultured Cyanobacteria.
ACS Chem. Biol. *15*, 758–765.

Moré, M.I., Finger, L.D., Stryker, J.L., Fuqua, C., Eberhard, A., and Winans, S.C. (1996). Enzymatic
Synthesis of a Quorum-Sensing Autoinducer Through Use of Defined Substrates. Science *272*, 1655–
1658.

Morin, D., Grasland, B., Vallée-Réhel, K., Dufau, C., and Haras, D. (2003). On-line high-performance
liquid chromatography–mass spectrometric detection and quantification of *N*-acylhomoserine lactones,
quorum sensing signal molecules, in the presence of biological matrices. J. Chromatogr. A *1002*, 79–92.

Myers, O.D., Sumner, S.J., Li, S., Barnes, S., and Du, X. (2017). One Step Forward for Reducing False
Positive and False Negative Compound Identifications from Mass Spectrometry Metabolomics Data:
New Algorithms for Constructing Extracted Ion Chromatograms and Detecting Chromatographic Peaks.
Anal. Chem. *89*, 8696–8703.

Newman, D.J., and Cragg, G.M. (2020). Natural Products as Sources of New Drugs over the Nearly Four
Decades from 01/1981 to 09/2019. J. Nat. Prod. *83*, 770–803.

Nieto Penalver, C.G., Morin, D., Cantet, F., Saurel, O., Milon, A., and Vorholt, J.A. (2006). *Methylobacterium extorquens* AM1 produces a novel type of acyl-homoserine lactone with a double unsaturated side chain under methylotrophic growth conditions. FEBS Lett. *580*, 561–567.

Nisbet, E.G., Dlugokencky, E.J., and Bousquet, P. (2014). Atmospheric science. Methane on the rise--again. Science *343*, 493–495.

Nothias, L.F., Petras, D., Schmid, R., Dührkop, K., Rainer, J., Sarvepalli, A., Protsyuk, I., Ernst, M., Tsugawa, H., Fleischauer, M., et al. (2020). Feature-based molecular networking in the GNPS analysis environment. Nat. Methods *17*, 905–908.

Papenfort, K., and Bassler, B.L. (2016). Quorum sensing signal–response systems in Gram-negative bacteria. Nat. Rev. Microbiol. *14*, 576–588.

Patel, N.M., Moore, J.D., Blackwell, H.E., and Amador-Noguez, D. (2016). Identification of Unanticipated and Novel *N*-Acyl *L*-Homoserine Lactones (AHLs) Using a Sensitive Non-Targeted LC-MS/MS Method. PLoS One. *11*, e0163469.

Pérez-Montaño, F., Guasch-Vidal, B., González-Barroso, S., López-Baena, F.J., Cubo, T., Ollero, F.J., Gil-Serrano, A.M., Rodríguez-Carvajal, M.Á., Bellogín, R.A., and Espuny, M.R. (2011). Nodulation-gene-inducing flavonoids increase overall production of autoinducers and expression of *N*-acyl homoserine lactone synthesis genes in rhizobia. Res. Microbiol. *162*, 715–723.

Pluskal, T., Castillo, S., Villar-Briones, A., and Orešič, M. (2010). MZmine 2: Modular framework for processing, visualizing, and analyzing mass spectrometry-based molecular profile data. BMC Bioinformatics *11*, 395.

Poonguzhali, S., Madhaiyan, M., and Sa, T. (2007). Production of acyl-homoserine lactone quorum-sensing signals is wide-spread in Gram-negative *Methylobacterium*. J. Microbiol. Biotechnol. *17*, 226-233.

Puri, A.W. (2019). Specialized Metabolites from Methylotrophic Proteobacteria. Curr. Issues Mol. Biol. 211–224.

Puri, A.W., Schaefer, A.L., Fu, Y., Beck, D.A.C., Greenberg, E.P., and Lidstrom, M.E. (2017). Quorum Sensing in a Methane-Oxidizing Bacterium. J. Bacteriol. *199*, e00773-16.

Radajewski, S., Ineson, P., Parekh, N.R., and Murrell, J.C. (2000). Stable-isotope probing as a tool in microbial ecology. Nature *403*, 646–649.

Schaefer, A.L., Val, D.L., Hanzelka, B.L., Cronan, J.E., and Greenberg, E.P. (1996). Generation of cell-to-cell signals in quorum sensing: acyl homoserine lactone synthase activity of a purified *Vibrio fischeri* LuxI protein. Proc. Natl. Acad. Sci. *93*, 9505–9509.

Schaefer, A.L., Greenberg, E.P., Oliver, C.M., Oda, Y., Huang, J.J., Bittan-Banin, G., Peres, C.M., Schmidt, S., Juhaszova, K., Sufrin, J.R., et al. (2008). A new class of homoserine lactone quorum-sensing signals. Nature *454*, 595–599.

Schaefer, A.L., Harwood, C.S., and Greenberg, E.P. (2018). "Hot Stuff": The Many Uses of a Radiolabel Assay in Detecting Acyl-Homoserine Lactone Quorum-Sensing Signals. In Quorum Sensing, L. Leoni, and G. Rampioni, eds. (New York, NY: Springer New York), pp. 35–47.

Simon, R., Priefer, U., and Pühler, A. (1983). A Broad Host Range Mobilization System for *in vivo*
Genetic Engineering: Transposon Mutagenesis in Gram Negative Bacteria. Nat Biotechnol *1*, 784–791.

Singh, B.K., Bardgett, R.D., Smith, P., and Reay, D.S. (2010). Microorganisms and climate change:
terrestrial feedbacks and mitigation options. Nat. Rev. Microbiol. *8*, 779–790.

Stein, L.Y., and Klotz, M.G. (2016). The nitrogen cycle. Curr. Biol. *26*, R94–R98.

Thornhill, S.G., and McLean, R.J.C. (2018). Use of Whole-Cell Bioassays for Screening Quorum
Signaling, Quorum Interference, and Biofilm Dispersion. In Quorum Sensing, L. Leoni, and G. Rampioni,
eds. (New York, NY: Springer New York), pp. 3–24.

Val, D.L., and Cronan, J.E. (1998). *in vivo* Evidence that *S*-Adenosylmethionine and Fatty Acid Synthesis
Intermediates Are the Substrates for the LuxI Family of Autoinducer Synthases. J. Bacteriol. *180*, 2644–
2651.

Volpe, J.J., and Vagelos, P.R. (1976). Mechanisms and regulation of biosynthesis of saturated fatty acids.
Physiol. Rev. *56*, 339–417.

Whiteley, M., Diggle, S.P., and Greenberg, E.P. (2017). Progress in and promise of bacterial quorum
sensing research. Nature *551*, 313–320.

Whittenbury, R., Phillips, K.C., and Wilkinson, J.F. (1970). Enrichment, isolation and some properties of
methane-utilizing bacteria. J. Gen. Microbiol. *61*, 205–218.