Title

Nuclear biocondensate-forming lncRNA and cellular stress surveillance shape host
cancer susceptibility

Anjali Bajpai[1,2*], Quazi Taushif Ahmad[1], Sushmita Kundu[1], Ravi Kant Pandey[1,3], Alan
Goodman[4], Apul Goel[5], Bushra Ateeq[1,2], Subhash C. Lakhotia[6], Pradip Sinha[1,2]

[1]Biological Sciences and Bioengineering, Indian Institute of Technology Kanpur, Kanpur
208016, India.
[2]The Mehta Family Centre for Engineering in Medicine, Indian Institute of Technology
Kanpur, Kanpur 208016, India.
[3]Current Address: PierianDx India Pvt Ltd, Chattushringi, Gokhalenagar, Pune 411016,
Maharashtra, India
[4]School of Molecular Biosciences, College of Veterinary Medicine, Washington State
University, Pullman, WA 99164, USA
[5]Department of Urology, King George's Medical University, Lucknow 226003, Uttar Pradesh,
India.
[6]Cytogenetics Laboratory, Department of Zoology, Banaras Hindu University,Varanasi,
221005, India.
* Corresponding author: Anjali Bajpai, anjalii@iitk.ac.in
22
23
ORCID:
A. Bajpai: 0000-0002-5645-2466
B. Ateeq: 0000-0003-4682-9773
A. G. Goodman: 0000-0001-6394-332X
S. C. Lakhotia: 0000-0003-1842-8411
P. Sinha: 0000-0002-9202-6050
30

32 *Running title:* Host genetic regulators of cancer susceptibility

34 *Key words*

35 lncRNA *hsrω*, omega speckles, nuclear condensate, haploinsufficiency, proteostasis, tumor
36 susceptibility, *dSTING*, *Keap1*

Abstract

Host genetics is known to influence cancer susceptibility. However, the specific candidate genes and molecular mechanisms that confer resistance remain poorly understood. Here, we demonstrate the power of haploinsufficiency screen to uncover host genetic regulators of cancer in *Drosophila* and identify the long noncoding RNA (lncRNA) *hsrω*, a structural component of nuclear biomolecular condensate known as omega speckles, as a key host cancer susceptibility locus. Loss of *hsrω* disrupts proteostasis and cell fitness, while its haploinsufficiency accelerates epithelial tumor progression driven by loss of the Lethal giant larvae (Lgl) tumor suppressor. Further validating the breadth of this screening strategy, we independently identified *Drosophila STING* (innate immunity) and *Keap1* (oxidative stress defense) as genetic modifiers of cancer. Moreover, in humans, copy number variations (CNV) in these genes and *Sat III* (a functional human homolog of *hsrω*) correlate with poor cancer prognosis, thereby revealing conserved stress pathways as potential host genetic susceptibility regulators.

Introduction

Genetic modifiers of cancer are genes that influence cancer risk by interacting with primary driver mutations or oncogenic processes without independently causing cancer. One of the central paradoxes of cancer biology is that, while a variety of oncogenes and tumor suppressors have been identified based on their causal underpinnings of cancer, modifier genes regulating host genetic susceptibility in humans remain underexplored (Castro-Giner et al., 2015; Turnbull et al., 2018). Among the known examples of host genetic modifiers identified by Genome-wide association studies (GWAS), notable are DNA repair genes for oxidative damage (*MUTYH* and *OGG1*, (Raetz and David, 2019), immune checkpoint genes (*CTLA4* and *PDCD1*, (Behrouzfar et al., 2021), or those affecting apoptosis pathways (*BIRC5*, (Siragusa et al., 2024). However, GWAS have limited capacity to identify functional cancer modifiers, which often require validation via haploinsufficiency or heterozygosity tests (Chenevix-Trench et al., 2007; Zeng et al., 2019). In *Drosophila*, genetic tractability enables a rapid assessment of such modifiers (Datta et al., 2023). A classic precedent is the *Minute* (Morata and Ripoll, 1975) class of ribosomal genes (Kongsuwan et al., 1985). Their haploinsufficiency compromises cellular proteostasis (Baumgartner et al., 2021) and disrupts tumor surveillance. Consequently, *Minute* +/- cells are outcompeted by fitter wild type cells or, conversely, fail to compete against tumors, thereby enhancing progression when tumor suppressors are lost (Agrawal et al., 1995; Froldi et al., 2010; Khan et al., 2013).

An emerging theme in cancer biology centres on the role of phase-separated biomolecular condensates, the dynamic cellular entities that regulate processes ranging from signalling and transcription to proteostasis and immune surveillance (Mehta and Zhang, 2022; Shin and Brangwynne, 2017). Disruptions of condensate integrity are increasingly linked to diverse pathologies including tumorigenesis (Banani et al., 2017; Boija et al., 2021). For instance, oncogene activation often utilizes super-enhancer condensates (Sabari et al., 2018) while the loss of PML condensates, for example, compromises p53 availability (Zhang et al., 2024).

In *Drosophila*, the long noncoding RNA-encoding *hsrω* gene codes for multiple transcripts ((Bendena et al., 1989; Lakhotia, 2011), also see **Figure 1A**), the largest nuclear-retained transcript, forms a scaffold for nucleoplasmic condensates known as omega speckles ((Prasanth et al., 2000), also see **Figure 1B**). These condensates dynamically sequester diverse heterogenous ribonuclear proteins (hnRNPs) and RNA-binding proteins (RBPs) to buffer cellular stress (Singh and Lakhotia, 2015). Crucially, loss of *hsrω* transcripts or the HRB87F hnRNP disrupts these condensates (Singh and Lakhotia, 2012). This biology is conserved since the functional human homolog, *Satellite III* (*Sat III*), similarly forms phase-separated nuclear stress bodies dependent on Heat shock factor 1 (HSF1) and sequester RNA-binding proteins (Goenka et al., 2016; Jolly et al., 2002; Jolly et al., 2004). Given the critical role of *hsrω* in stress buffering, we hypothesized that genetic perturbation of this lncRNA would profoundly impact cancer progression.

Here, we employed a hypothesis-driven candidate gene approach to uncover host genetic modifiers of cancer. We demonstrate that the omega speckle-associated lncRNA, *hsrω*, acts as a potent host genetic modifier of cancer susceptibility by regulating proteostasis. Furthermore, to validate the breadth of this haploinsufficiency screening approach, we expanded our search to other arms of the cellular stress response. We demonstrate that *Drosophila STING* (*dSTING*), a regulator of innate immune stress (Cai et al., 2020; Martin et al., 2018) and *Keap1*, a regulator of oxidative stress (Sykiotis and Bohmann, 2008) also serve as independent host genetic modifiers. Collectively, our findings highlight a conserved network of stress surveillance pathways that define tumor susceptibility.

Results

Study design

To systematically uncover how host genetics contributes to intrinsic epithelial tumor surveillance, we employed a candidate gene approach focusing on haploinsufficiency. We selected distinct but potentially overlapping stress regulators: the nuclear biomolecular condensate-associated lncRNA, *hsrω*, the innate immunity pathway component, *dSTING* and the redox regulator Keap1. Our strategy involved a stepwise characterization: first, we defined the consequences of *hsrω* loss on tissue growth, proteostasis, and cellular fitness in normal epithelium. Next, we tested whether *hsrω* haploinsufficiency facilitates the neoplastic transformation of somatic clones lacking the tumor suppressor *Lgl*. Finally, extending the logic of stress surveillance, we examined whether haploinsufficiency of *dSTING*—a key sentinel of the cGAS–STING pathway—mirrors the tumor-promoting effects of *hsrω* loss, thereby establishing a broader role for stress regulators in host cancer susceptibility.

hsrω mutants are growth-compromised and display disrupted proteostasis

Homozygosis for the *hsrω* near-null mutant, *hsrω^66* (Johnson et al., 2009), causes high embryonic lethality, with the escapers emerging as weak short-lived adults . Analysis of these larvae revealed undergrown wing imaginal discs with significantly fewer dividing cells per unit

area than controls (**Figure 1C–F**) and elevated cell death (**Figure 1G, H**). The $hsr\omega^{66}$ somatic clones were similarly smaller than their wild type twin spots, with cell death evident along clone borders (**Figure 1I, J**)—a phenotype characteristic of 'loser' cell behaviour in cell competition (de la Cova et al., 2004; Moreno et al., 2002). These growth defects prompted us to examine whether loss of $hsr\omega$ perturbs the protein homeostasis landscape in developing epithelial cells.

Proteomic profiling revealed a collapse in the translation machinery. Comparing the proteome of mutant and wild type (WT) larval wing epithelium, we noted $>\pm2$-fold ($p<0.05$) changes in 73 proteins in $hsr\omega^{66}$ mutants (**Figures 2A and S1A, B**). We observed a specific depletion of ribosomal structural proteins (RpS15Ab, mRpL50, and mRPL24; **Figure 2B**) alongside proteins involved in protein stability such as the heat shock protein Hsp23 and protein turnover, the SUMO-protease Ulp1 (**Figure S1C**). Rank-based enrichment analysis confirmed the overall downregulation of cytosolic and mitochondrial large ribosomal subunits (**Figure 2C**), as well as the cytoplasmic and mitochondrial translation machinery (**Figure 2D**). Furthermore, RNA Pol II transcription machinery components and the mediator complex were depleted (**Figures S1E**), indicating a systemic failure in gene expression and protein synthesis control. We noted that the $hsr\omega^{66}$ proteome showed an upregulation of stress response markers (**Figure S1D**), including the cytoplasmic Glutathione S-transferase E6 (GstE6) (Aloke et al., 2024) and mitochondria-associated Mgstl (Toba and Aigaki, 2000), indicative of redox imbalance. Such cellular changes likely prefigure DNA damage signatures, consistent with the observed elevation of Rpa1, a DNA double-strand break-associated protein (**Figures 2E**).

We further validated these proteomic signatures using cellular stress markers. The $hsr\omega^{66}$ mutant wing epithelium displayed elevated ROS levels (DHE staining; **Figure 2F**) and enhanced JNK signaling (**Figure 2G**). This oxidative stress was accompanied by augmented γH2AX staining, confirming the presence of DNA double-strand breaks (Mah et al., 2010 ; **Figures 2H**). We reasoned that a suboptimal proteome under such stress would lead to the accumulation of damaged proteins, necessitating increased lysosomal clearance (Settembre et al., 2013). In agreement, $hsr\omega^{66}$ wing discs showed increased acidic lysosomes detected by LysoTracker (**Figure 2I**) and an increase in Ubiquitin-positive foci (**Figure S2G**) indicating proteins marked for proteasomal degradation (Martínez-Férriz et al., 2022). Finally, we observed signatures of increased proteotoxicity in the wing epithelium owing to loss of proteostasis. For instance, the mutant proteome showed a marked elevation of ARC1 and ARC2 (**Figure 2A**), proteins associated with pathological aggregates (Schulz et al., 2023). Crucially, $hsr\omega^{66}$ wing epithelium exhibited marked increase in cellular uptake of ProteoStat (**Figure 2J**), a dye that fluoresces upon intercalation with protein aggregates (Baumgartner et al., 2021), and a pronounced increase in Thioflavin fluorescence in cell homogenates (**Figure 2K**), both telltale markers of elevated proteotoxicity (Kandel et al., 2024).

These findings indicate that $hsr\omega$ loss disrupts ribosome integrity and proteome surveillance, fostering oxidative stress and proteotoxicity. The resulting translational stress therefore compromises cellular fitness, predisposing $hsr\omega^{66}$ cells to elimination (**Figure 1I**) through competition with the neighboring wild type cells. Thus, the cell-competition phenotype of $hsr\omega$

mutants—and, by extension, the functional compromise of the omega speckle biomolecular
condensate—culminates in impaired cellular proteostasis and compromised stress adaptation.

hsrω displays haploinsufficiency for tumor suppression

We next tested the survival of _lgl_ somatic clones in _hsrω_^{66}/+ larvae, given that perturbed
proteostasis permits tumor progression in other contexts, such as the _Minute_ genetic
background (Baumgartner et al., 2021; Khan et al., 2013). In a wild type epithelium, somatic
clones lacking Lgl are recognized and eliminated by a tissue surveillance mechanism known
as cell competition (Khan et al., 2013; Menéndez et al., 2010; Suijkerbuijk et al., 2016). These
clones display increased cellular stress (**Figure S2**), exhibit elevated cell death, and are basally
extruded ((Khan et al., 2013), also see **Figure 3A**). By contrast, _lgl_ clones generated in
hsrω^{66}/+ wing imaginal discs evaded elimination and underwent neoplastic transformation,
marked by disrupted cortical F-Actin (**Figure 3B, C**). Consistent with the acquisition of a
'winner' cell state, we observed cell death in _lgl_$^+$/_lgl_ cells along the _lgl_ clone boundaries (**Figure**
3B). Furthermore, these transformed _lgl_ clones displayed elevated cell proliferation (pH3,
Figure 3D) and an increase in Matrix Metalloproteinase-1 (MMP1; **Figure 3E**), indicating
ECM degradation and the acquisition of invasive properties (Beaucher et al., 2007). We thus
hypothesized that the compromised proteostasis in the _hsrω_ mutant background (**Figure 2**)
would render the epithelium hypersensitive to proteosome inhibitor. Indeed, administration of
the classical proteasome inhibitor MG-132 (Chen et al., 2011) compromised the tumor
susceptibility of _hsrω_^{66}/+ larvae, reducing _lgl_ tumor burden in a dose-dependent manner
(**Figure 3F**). This suggests that the tumor-promoting environment of _hsrω_ haploinsufficiency
relies on a precarious proteostasis balance.

We next asked if this tumor susceptibility was driven by the loss of structural integrity of the
omega speckle. A hallmark of condensate-based regulatory systems is their sensitivity to
component dosage—both depletion and overexpression can perturb phase separation (Shin and
Brangwynne, 2017) also recapitulated by omega speckles that are sensitive to doses of _hsrω_
and omega speckles-associated hnRNPs such as Hrb87F (Lakhotia, 2011; Prasanth et al.,
2000). Consistent with this model, we found that _lgl_ somatic clones underwent neoplastic
transformation not only upon _hsrω_ knockdown (_UAS-hsrω-RNAi_; **Figure 3G-I**) but also upon
overexpression (_UAS-hsrω_^{EP3037}; **Figure 3J, K**). Under both conditions, _lgl_ clones gained a
growth advantage, induced cell death in the neighboring cells (**Figure 3G**), and upregulated
invasion markers (MMP1; **Figure 3I**).

To confirm that condensate integrity is the critical factor, we manipulated other structural
components. Heterozygosity of _Df(3R)Hrb87F_—which encodes an hnRNP localized to omega
speckles—similarly promoted _lgl_ transformation (**Figure S3A**). Moreover, downregulating the
chromatin remodeling ATPase _ISWI_—required for omega speckle structural integrity (Onorati
et al., 2011; Singh and Lakhotia, 2015)—also facilitated neoplastic transformation (**Figure**
S3C). Finally, _lgl_ somatic clones induced in the adult intestinal epithelium of _hsrω_^{66}/+ adult
flies also displayed neoplastic transformation (**Figure S4**), unlike those generated in a WT
background. This indicates that _hsrω_ haploinsufficiency creates a tumor-susceptible host

genetic background, likely through sensitivity to biocondensates disruption and ensuing
impaired protein homeostasis.

Tumor inhibitory role of redox regulators Nrf2-Keap1 and inflammatory protein
dSTING

To validate the versatility of haploinsufficiency screening in uncovering diverse host genetic
determinants, we extended our analysis to two other functional classes of stress regulators
identified in our screen: the redox sensor Nrf2-Keap1 and the viral immune sensor cGAS-
STING. While somatic variations in these pathways are suspected to influence cancer risk
(Michalak and Michalak, 2025) their direct impact on intrinsic tumor surveillance remains to
be functionally dissected in a host genetic model.

First, we examined the Nrf2-Keap1 oxidative stress pathway. This conserved system mediates
cellular responses to oxidative damage as Keap1 suppresses the transcription factor Nrf2
(*Drosophila* homolog, CncC) by targeting it for proteasomal degradation (Fuse and Kobayashi,
2017). In the absence of Keap1, CncC translocates to the nucleus to activate antioxidant and
proteostasis genes (Radhakrishnan et al., 2010). We reasoned that altering this redox balance
might shield emerging tumor cells from oxidative stress-induced elimination. Consistent with
this, *lgl* somatic clones generated in a *Keap1* haploinsufficient background (*Keap1^{EY5}/+*)
exhibited enhanced growth and neoplastic transformation (**Figure 4A**) when compared to *lgl*
clones generated in wild type genetic background. We observed a similar tumor-promoting
effect upon cell-autonomous downregulation of *Keap1* (*lgl; UAS-Keap1-RNAi*; **Figure 4B, C**)
or direct genetic gain of *CncC* (*lgl; UAS-CncC*; **Figure 4D**), with transformed *lgl* clones
displaying elevated cell death in abutting wild type neighbors (**Figure 4C**). This suggests that
constitutive activation of antioxidant defense (via *Keap1* loss) acts as a barrier to the
elimination of oncogenic clones.

Next, we investigated the cGAS–STING pathway. This pathway acts as a sentinel for aberrant
cytosolic DNA—a hallmark of genomic instability and cellular damage or of virus-induced
biotic stress (Hopfner and Hornung, 2020). In *Drosophila*, activation of dSTING triggers an
inflammatory response via the NF-κB/Relish pathway (Cai et al., 2020; Martin et al., 2018).
We hypothesized that haploinsufficiency of *dSTING* would impair the host's ability to sense
and eliminate early oncogenic clones. Indeed, *lgl* somatic clones generated in a *dSTING* mutant
background (*dSTING^{ΔRG5}/+*) showed enhanced survival and underwent neoplastic
transformation (**Figure 4E**), effectively phenocopying the *hsrω^{66}/+* tissue. Similarly, cell-
autonomous *dSTING* knockdown within *lgl*-deficient clones led to robust tumor formation
(**Figure 4F**), confirming that innate immune sensing is required for effective tumor
surveillance.

Finally, we explored the potential crosstalk between these mechanisms. While identified
independently, our data suggests a functional link between nuclear condensates and immune
surveillance. Notably, we observed that compared to wild type larval salivary glands, the
hsrω^{66} mutant cells displayed disrupted dSTING punctae with reduced diffuse distribution in

cytoplasm (**Figure 4G, H** and **Figure S5**). This suggests that the condensate defects in *hsrω*
mutants may compromise dSTING trafficking or assembly. This was confirmed by marked
suppression of *lgl* tumor growth following treatment with the dSTING agonist (**Figure 4I**) 2′3′-
cGAMP (Cai et al., 2020). This demonstrates that pharmacological reactivation of innate
immunity can compensate for the surveillance deficits caused by *hsrω* loss, highlighting a
functional intersection between nuclear condensates and immune signaling.

CNVs in human homologs of stress regulators and Sat III correlate with cancer severity

To evaluate the clinical relevance of our findings in *Drosophila*, we investigated whether
genetic alterations in the human homologs of these stress regulators correlate with patient
outcomes. First, we examined if variations in gene dosage affect the survival of cancer patients
using the Cancer Genome Atlas (TCGA). We analysed Copy Number Variations (CNVs) in
human *KEAP1* and *STING1* across diverse cancer types (**Figures S6A, B)**. Although TCGA
data primarily reflects somatic alterations of tumor origin, it provides a window into how gene
dosage impacts progression. Kaplan-Meier survival plots revealed a significant correlation
between CNVs in these genes (including deletions and amplifications) and poor patient
survival (**Figures S6A, B**). This confirms that, similar to our *Drosophila* models, the dosage of
these stress regulators is a critical determinant of human cancer prognosis.

Next, we focused on the human functional analog of *hsrω*, the *Satellite III* (*Sat III*) repeats.
While *hsrω* lacks a sequence homolog in mammals, the *Sat III* repeats at the pericentromeric
region of chromosome 9 (9q12) are its functional analog (Jolly and Lakhotia, 2006; Jolly et al.,
2004). Like *hsrω*, *Sat III* is actively transcribed during stress to form phase-separated nuclear
'stress bodies' (Jolly et al., 2004; Goenka et al., 2016). These bodies sequester Heat Shock
Factor 1 (HSF1) and specific RNA-binding proteins. Since repetitive *Sat III* sequences are not
resolved in standard TCGA exome data, we first looked at its obligate binding partner, HSF1.
Notably, TCGA analysis revealed that CNVs in HSF1 are also associated with poor patient
survival (**Figure 5A, B**), thereby supporting the importance of the stress body machinery.
Finally, we tested the hypothesis that germline variations in *Sat III* itself act as a host
susceptibility factor. Since *Sat III* resides in a large heterochromatic block of pentameric repeats
at 9q12 (**Figure 5C**), we utilized digital droplet PCR (ddPCR) to precisely quantify copy
number variations in *Sat III* using the 9q12-specific, pHuR98 probe (de Lima et al., 2021; Jolly
et al., 2002). We analysed genomic DNA from prostate cancer patients and compared them to
benign controls. Strikingly, we observed significant germline copy number variations in *Sat III*
in patients with high cancer severity (Gleason scores 6–10) compared to benign controls
(**Figure 5D**). Further, reminiscent of the tumor-promoting effects of *hsrω* haploinsufficiency
in flies, we also observed an overall decrease in Sat III transcript levels across different prostate
cancer cell lines compared to benign control (**Figure 5E**). This agrees with previous report that
Sat III is misregulated in cancer cell lines (Chatterjee and Sengupta, 2022).

Collectively, our findings in *Drosophila* and human cancers show that perturbations in the
dosage of stress surveillance components, whether the protein coding *KEAP1* and *STING1* or

the noncoding *hsrω*/*Sat III* architectural RNAs, are broadly associated with cancer
susceptibility, severity and poor prognosis.

Discussion

Our study establishes *Drosophila* haploinsufficiency screen as a powerful tool for identifying
host genetic regulators of cancer susceptibility **(Figure 6)**, a class of genes often overlooked
by traditional oncogene/tumor suppressor screens. While the roles of high-penetrance
mutations (e.g., TP53, PTEN) are well defined (Berger and Pandolfi, 2011), the genetic
architecture of host resistance remains underexplored. We demonstrate that subtle genetic
deficits (haploinsufficiency) in fundamental homeostatic regulators, specifically the nuclear
condensates, redox sensors, and immune sentinels, do not initiate tumors per se, but severely
erode the 'surveillance buffer' of cells. This renders the epithelium permissive to neoplastic
transformation upon a second hit, such as loss of the cell polarity protein Lgl.

A major finding of this study is the identification of the lncRNA-encoding *hsrω*, essential for
formation of omega speckle biomolecular condensates, as a key suppressor of susceptibility.
This expands the "Proteostasis Addiction" model of cancer (Dai and Sampson, 2016). We
propose that *hsrω*- dependent omega speckle condensates function as proteostasis regulator by,
sequestering various RNA-binding proteins, and buffering the translational stress. Loss of this
buffer, as in *hsrω* haploinsufficiency, creates a 'proteostasis-vulnerable' epithelium that cannot
cope with the physiological demands of emerging oncogenic clones, which effectively lowers
the threshold for tumorigenesis. This mirrors the classic Minute phenotype in *Drosophila*
(Baumgartner et al., 2021), but with a crucial distinction that in this case a non-coding RNA
rather than a ribosomal protein acts as the linchpin of surveillance.

Validating the breadth of our screening strategy, we independently identified *STING* (an innate
immunity factor) and *Keap1* (a redox regulatory factor) as host modifiers. While distinct in
function, these pathways likely converge. Our observations that *hsrω* loss disrupts the cellular
distribution of dSTING, and that STING agonists can rescue the *hsrω* tumor susceptibility
phenotype suggest a functional intersection. We postulate that the dynamics of omega speckle
nuclear condensates also regulate the nucleocytoplasmic distribution or availability of specific
immune or redox factors, such as STING. Thus, *hsrω*, *STING*, and *Keap1* may function as
nodes in an integrated 'Stress Surveillance Network.' Disabling any node, whether by
compromising activities of the hsrω or the sensors (STING/Keap1), collapses the network,
which in turn permits cells carrying tumorigenic hits overcome tumor suppressive cell
competition.

A significant implication of our study is the potential translational link to human *Sat III* repeats.
Although the *Sat III* repeats have long been considered "junk DNA" or mere markers of stress,
in context of the *Drosophila* model, we suggest a significant tumor-suppressive function to this
non-coding locus. Our finding that the germline Copy Number Variations (CNVs) in *Sat III*
correlate with prostate cancer severity suggests that the highly polymorphic repetitive regions
of the human genome, often ignored in standard exome sequencing, may harbor critical
determinants of cancer susceptibility. Just as *hsrω* dosage dictates fly cells' tumor resistance,

the intrinsic copy number of human *Sat III* may determine the robustness of a patient's "cellular stress response", thereby influencing clinical prognosis.

Finally, our findings resonate with the evolutionary concept that cancer resistance is a distinct trait from tumor suppression (Klein, 2009). Long-lived mammals, such as elephants or whales, have evolved redundant resistance mechanisms, including increased TP53 copy numbers (Sulak et al., 2016). We propose that *hsrω/Sat III*, STING, and Keap1 represent a similar layer of 'background' resistance. These genes do not drive the cancer, but they set the rules of engagement. A host with robust condensate dynamics and immune sensing (high resistance) forces oncogenes to overcome a steep barrier, whereas a host with haploinsufficient defects (low resistance) provides a more permissive environment. Combining *Drosophila* genetics with human CNV analysis opens a new avenue to systematically map these elusive 'host resistance' genes, moving the field beyond the tumor-centric driver model towards a more holistic understanding of cancer susceptibility.

Acknowledgement

We acknowledge Bloomington Drosophila Stock Centre (Indiana University) for fly stocks and Developmental Studies Hybridoma Bank (University of Iowa) for antibodies. Ethical approval for use of patient samples in this study was obtained from the Institutional Ethics Committee (IEC) at Indian Institute of Technology Kanpur (Approval Number: IITK/IEC/2020-21/II/05) issued on February 18, 2021 for five years. All patients participating in this study provided written informed consent. We thank S. Ganesh, Indian Institute of Technology Kanpur, for the gift of the FK2 anti-ubiquitin antibody. This work was supported by the Department of Biotechnology, Govt. of India and Wellcome Trust India Alliance (IA/E/13/1/501271) and Department of Science and Technology, DST/WOS-A/LS-75/2021 to AB. Support by the Science & Engineering Research Board (SERB), Department of Science and Technology, Govt. of India (CO/S5/AB-03/2017) to PS and DBT/ Wellcome Trust India Alliance (IA/S/19/2/504659) to BA is acknowledged. SCL is supported by the Department of Biotechnology, Ministry of Science and Technology (BT/PR32126/BRB/10/1775/2019) and the Science & Engineering Research Board, Govt. of India, as a Distinguished Fellow (SB/DF/009/2019).

The authors declare no competing interests.

Material and Methods

Fly Stocks

Fly stocks were maintained at 25º C on standard fly food containing corn powder, yeast, and sugar. The following mutant/transgenic lines were used in this study: *w; l(2)gl⁴ FRT40A/CyO* (#36289, BDSC); *w; w; hsrω⁶⁶/TM6B* (Johnson et al., 2011); *w;UAS-hsrω-RNAi(2x)*(Mallik and Lakhotia, 2009); *EP3037/ EP3037* (Mallik and Lakhotia, 2009); *Df(3R)Hrb87F/Tb* (#67701, BDSC); *y w hs-flp, tub-Gal4, UAS-GFP/; tub-Gal80 FRT40A/CyO; w; FRT82B* (#86313, BDSC); *y w hs-flp; FRT82B Ubi-GFP* (#5188, BDSC); *w; TRE-DsRed;+ (*#59011,

BDSC);*UAS-STING RNAi* (#31565, BDSC); *w;+;th-lacZ* (Bennett and Harvey, 2006); *STING^{ARG5}* (Martin et al., 2018); *Keap^{EY5}* (Gift from Dirk Bohmann (Sykiotis and Bohmann, 2008)). BDSC: Bloomington Drosophila Stock Center, Indiana University Bloomington.

Genotypes of clone and method clone induction:

Heat shock induced somatic clones were generated using the flp/FRT technique (Xu and Rubin, 1993) or MARCM technique (Lee and Luo, 1999). Heat shock was given at $37^{O}C$ for 30 minutes to early second instar progeny larvae (48 hours after egg laying, ARL) of the following parental genotypes. The larvae were then grown at $25^{O}C$ till 96 to 120 hours AEL.

- ***hsrω^{66}* mutant** clones **(**non-GFP) and its twin spot (two copies of GFP):
  FRT82B hsrω^{66}/ TM3b X *y w hs-flp; FRT82B Ubi-GFP*
- ***lgl⁻* clones in *hsrω^{66}/+* genetic background:**
  lgl^4 FRT40A/Cy-O;hsrω^{66}/ TM3b and *y w hs-flp tub Gal4 UAS-GFP; tub-Gal80 FRT40A/Cy-O*
- ***lgl⁻* clones in *Df(3R)Hrb87F)/+* genetic background:**
  lgl^4 FRT40A/Cy-O;Df(3R)Hrb87F/TM3b and *y w hs-flp tub Gal4 UAS-GFP; tub-Gal80 FRT40A/Cy-O*
- ***lgl⁻ UAS-hsrω-RNAi* and *lgl⁻ UAS-hsrω ^{EP3037}*:**
  lgl^4UAS-hsrω RNAi (2X) FRT40A; and *y w hs-flp tub Gal4 UAS-GFP; tub-Gal80 FRT40A/Cy-O.* or *lgl^4FRT40A; hsrω^{EP3037}* and *y w hs-flp tub Gal4 UAS-GFP; tub-Gal80 FRT40A/Cy-O* respectively.
- ***lgl⁻* in *STING^{Δ5}/+* genetic background:** *lgl^4 FRT40A, STING^{Δ5}/Cy-O* and *y w hs-flp tub Gal4 UAS-GFP; tub-Gal80 FRT40A/Cy-O*
- ***lgl⁻ UAS-STING-RNAi:*** *w;lgl^4 FRT40A UAS-STING RNAi/Cy-O;+* and *y w hs-flp tub Gal4 UAS-GFP; tub-Gal80 FRT40A/Cy-O*
- ***lgl⁻* in *Keap1^{EY5}/+* genetic background:** *w;lgl^4 FRT40A Keap1^{EY5}/Cy-O;+* and *y w hs-flp tub Gal4 UAS-GFP; tub-Gal80 FRT40A/Cy-O*
- ***lgl⁻ UAS-STING-RNAi:*** *w;lgl^4 FRT40A UAS-keap1RNAi/Cy-O;+* and *y w hs-flp tub Gal4 UAS-GFP; tub-Gal80 FRT40A/Cy-O*
- ***lgl⁻ UAS-CncC:*** *w;lgl^4 FRT40A UAS-CncC/Cy-O;+* and *y w hs-flp tub Gal4 UAS-GFP; tub-Gal80 FRT40A/Cy-O*

Immunofluorescence staining and microscopy

Larvae of desired genotypes were dissected in 1x Phosphate Buffer Saline (PBS), and imaginal discs were fixed in 4% paraformaldehyde in 1x PBS containing 0.02 % Triton-X-100, and incubated overnight in the desired primary antibody at $4^{0}C$, followed by washing and incubation with appropriate fluorescently tagged secondary antibodies for two hours at the room, and counterstained with SYTOX-Green (S33025, Invitrogen) or Alexa Fluor™ 633 Phalloidin (A22284, Invitrogen). The following primary antibodies were used: anti-Caspase-3, rabbit (1:500 dilution, MERK/Sigma Aldrich, C8487), Polyubiquitinylated conjugates monoclonal antibody FK1, mouse (1:500 dilution, Enzo Life Sciences, BML-PW8805-0500), Anti-phospho Histone-3 (pSer^{10}), rabbit (1:500 dilution, MERK /Sigma Aldrich H0412), anti- β Galactosidase, mouse (1:200 dilution, MERK/Sigma Aldrich, G6282). To detect MMP1, we used a cocktail of

monoclonal antibodies raised in mice (3A6B4, 3B8D12, 5H7B11, and 3B8D12; 1:1:1:1 ratio at a
final dilution of 1:300, Developmental Studies Hybridoma Bank (DSHB), University of Iowa),
anti-Fasciclin III, Fas III, mouse, (7G10, DSHB). Anti-dSTING, rabbit (1:100, (Martin et al.,
2018)). We used the following secondary antibodies from Invitrogen: Alexa Fluor™ 555 anti-
mouse (A32727) and Alexa Fluor 555 anti-rabbit (A32732).

Nuclear staining was carried out using either SYTOX-Green (Invitrogen #S33025) at 1:300 or To-
PRO (Invitrogen, #T3605) at 1:1000 dilution. Corticular F-Actin was labeled using Alexa Fluor™
633 Phalloidin (#A22284, Invitrogen). After washing, the discs were mounted using a Vectashield
anti-fade medium (Vector Laboratories, H-1000). Images were acquired with a Leica-SP5 confocal
microscope and processed using the Leica confocal software LAS AF. Images were acquired using
a 20x or a 40x (oil immersion) objective. Images of GFP-marked lgl; $hsr\omega^{66}$/+ tumor-bearing
larvae were acquired with a Leica M205FA stereomicroscope with epi-fluorescent illumination
(excitation filter 480 nm). All images were assembled using Adobe Photoshop CS5 software.

For detecting protein aggregates, the wing imaginal discs were dissected out from larvae of the
desired genotype and age in PBS and fixed in 4% formaldehyde in 1× PBS, followed by
permeabilization in 0.4% Triton X-100 and 3mM ethylenediaminetetraacetic acid (pH 8.0) in 1×
PBS. Fixed imaginal discs were subsequently incubated in ProteoStat detection reagent (1:10,000
dilution, Enzo Life Sciences, #ENZ-51023-KP050), washed in 1XPBS, and the wing imaginal
discs were mounted in glycerol-based anti-fade and imaged using Leica-SP5 confocal microscope.
We used LysoTracker® Red DND-99 dye (#L7528, Thermo-Fisher Scientific/Invitrogen) to
detect acidic lysosomes and MitoTracker™ Red CMXRos (#M22425, Thermo-Fisher
Scientific/Invitrogen). This cell-permeant, red-fluorescent dye stains mitochondria in live cells
and is a marker for altered membrane potential. Briefly, larvae of the desired genotype were
gently rinsed and dissected in 1x PBS to remove tissues like fat body, gut, etc. The flipped
larvae were incubated in freshly prepared 500nM of LysoTracker (in 1X PBS) or 500nM
MitoTracker (in 1X PBS) at room temperature in the dark for 15 minutes, followed by a
thorough rinse in 1x PBS. Unfixed wing discs were mounted in anti-fade-containing
Vectashield (Vector Laboratories, #H-1000), and images were acquired immediately using a
Leica-SP5 confocal microscope and processed using Leica confocal software-LAS AF and
assembled using Adobe Photoshop.

We used dihydroethidium (#D11347, Invitrogen), a free radical sensitive dye, to detect
ROS levels in unfixed larval tissues (Owusu-Ansah et al., 2008). Briefly, larvae of the desired
genotype were washed and dissected in 1xPBS. After removing the fat body, gut, and other
adherent tissues, the flipped larvae were incubated in freshly made 1:1000 DHE at room
temperature in the dark for 15 minutes. The wing imaginal discs were subsequently thoroughly
rinsed in 1XPBS, mounted in anti-fade containing Vectashield (Vector Laboratories, #H-1000),
and images were acquired with Leica-SP5 confocal microscope and processed using Leica
confocal software-LAS AF and AdobePhotoshop CS3.

Clone size and Fluorescence Quantification

ImageJ platform ((Schneider et al., 2012) , http://rsb.info.nih.gov/ij/) was used for image
analysis. For measuring size of wing discs, confocal projection images acquired with a

Leica-SP5 confocal microscope were projected as a single image and used as the input file. The Freehand selection tool was used to trace the outline of the imaginal discs marked by nuclear fluorescence, and the "Measure" option was used to obtain the area of the wing disc in arbitrary units. For clone size quantification freehand selection tool was used to trace the outline of the nuclear GFP-marked clones in individual discs. For quantification of $hsr\omega^{66}$ mutant clones (marked by complete loss of GFP) and their twin spots (hsr^+ GFP/ hsr^+GFP) characterized by their brighter GFP fluorescent was calculated using "Analyze Particle" option of Image J platform on confocal projection images. Images were first converted to an 8-bit format; signal thresholds were set to calculate the number of pixels with no signal (non-GFP, $hsr\omega^{66}$/$hsr\omega^{66}$ clones) and those with double GFP (twin spots, +/+). To quantify total fluorescence (for pH3/DHE/TRE-dsRED/γH2AX/Lysotracker/ProteoStat) confocal projection images of the red channel of individual wing imaginal discs was used as input file after converting to 8-bit formats. Image J was used to overlay a grid of uniform unit size, and the maximum fluorescence intensity per ROI was recorded. Graphs were plotted using MS Excel.

ThT Assay for estimating protein aggregates

Larval wing discs were homogenized in ice cold 1xPBS to release cellular contents, including any insoluble amyloid-like aggregates. Thioflavin T (ThT) at a final concentration of 10μM was added to the homogenate and incubate for 10 minutes at room temperature. ThT selectively binds to β-sheet-rich protein aggregates, such as amyloids, leading to a significant increase in its fluorescence. The homogenate was transferred to a black 96-well plate and fluorescence was measured at excitation of 440–450 nm and emission at 480–490 nm using a plate reader (Bio-Rad).

Proteomics of wing imaginal discs from *hsrω⁶⁶* mutant and *Canton^S* larvae

- **Protein extraction from wing discs of third instar larvae**

Wing imaginal discs from third instar larvae were dissected in cold 1× PBS put in 100 μl extraction buffer (6 M GnHCl in 50 mM Tris-HCl pH 7.4, 65 mM dithiothreitol) with 50 mM sodium acetate and protease inhibitors (1× protease inhibitor cocktail with 0.2 mM PMSF). Within 30 minutes of dissection the wing discs were sonicated with a Bioruptor (Diagenode) using 30 s ON and 30 s OFF for 5 cycles at 4°C. Cell debris was removed by centrifuging at 6000 g for 3 minutes and supernatant transferred to a new tube. The protein concentration was determined spectrophotometrically using Nanodrop. Five micrograms of the protein were used for LC-MS-MS analysis.

- **Sample preparation for LC-MS/MS**

Five micrograms of the protein samples were reduced with 5 mM tris(2-carboxyethyl) phosphine (TCEP), further alkylated with 50 mM iodoacetamide and digested with Trypsin (1:50, Trypsin/lysate ratio) for 16 h at 37°C. Digests were cleaned using a C18 silica cartridge to remove the Digests were cleaned using a C18 silica cartridge to remove the salt and dried using a speed vac. The dried pellet was resuspended in buffer A (2% acetonitrile, 0.1% formic acid).

- **Mass Spectrometric Analysis of Peptide Mixtures**

Experiments were performed on an Easy-nlc-1000 system coupled with an Orbitrap Exploris mass spectrometer. 1ug of peptide sample were loaded on C18 column 15 cm, 3.0μm Acclaim PepMap (Thermo Fisher Scientific) and separated with a 0–40% gradient of buffer B (80% acetonitrile, 0.1% formic acid) at a flow rate of 300 nl/min) and injected for MS analysis. LC gradients were run for 60 minutes. MS1 spectra were acquired in the Orbitrap (Max IT = 25ms, AGQ target = 300%; RF Lens = 70%; R=60K, mass range = 375−1500; Profile data). Dynamic exclusion was employed for 30s excluding all charge states for a given precursor. MS2 spectra were collected for top 12 peptides. MS2 (Max IT= 22ms, R= 15K, AGC target 200%).

- **Data Processing**

All samples were processed and RAW files generated were analyzed with Proteome Discoverer (v2.5) against the Uniprot Drosophila melanogaster database. For dual Sequest and Amanda search, the precursor and fragment mass tolerances were set at 10 ppm and 0.02 Da, respectively. The protease used to generate peptides, i.e., enzyme specificity was set for trypsin/P (cleavage at the C terminus of "K/R: unless followed by "P"). Carbamidomethyl on cysteine as fixed modification and oxidation of methionine and N-terminal acetylation were considered as variable modifications for database search. Both peptide spectrum match and protein false discovery rate were set to 0.01 FDR.

- **Proteome data analysis**

To identify biologically relevant protein signatures in $hsr\omega^{66}$ wing disc proteome versus wild type proteome we calculated the log2 abundance ratios, using mean abundance values for individual UniProt IDs. Only those with combined FDR confidence <0.05 (medium) or <0.01 (high) were taken into consideration; those with combined FDR >0.05 (low) were discarded. We further filtered out peptides that were not detected in either MS or MS/MS spectra depending on the peak calling. Overall 3125 Uniprot IDs were subjected to function enrichment analysis using STRING platform (https://string-db.org/, (Szklarczyk et al., 2021)) in conjunction with ssGSEA (single-sample Gene Set Enrichment Analysis).

Kaplan-Meier survival plots

Kaplan-Meier survival plots were generated using clinical data from TCGA (The cancer genome atlas), using cBioPortal platform (https://www.cbioportal.org/). Datasets were restricted to epithelial cancers only and queried for CNA (Copy Number Alterations) and mutations for individual genes: *KEAP1*, *HSF1* and *STING1*.

Study of *Sat III* germline copy number variations in PCa patients

- **Study population and patient recruitment**

The peripheral blood used in this study was collected from the biopsy-confirmed treatment-naïve prostate cancer (PCa) patients (n=18) and benign prostatic hyperplasia (BPH) patients (n=5), enrolled at the Department of Urology, King George's Medical University (KGMU,

Lucknow, India). Clinical specimens were collected after obtaining written informed consent

and institutional review board approvals from KGMU and Indian Institute of Technology

Kanpur (Kanpur, India). The study was initiated after obtaining approval from the

Institutional Ethics Committee.

Digital droplet PCR quantification of SatIII copy number in PCa patients

Blood was collected in K_2-EDTA 2ml tubes from BPH (benign prostatic hyperplasia) and

PCa (prostate cancer) patients. Genomic DNA was isolated using QIAamp® DNA Blood

mini-kit (#51104, Qiagen), which was further processed to identify *Sat III* copies. Droplet

digital PCR was performed with the QX200 Droplet Digital PCR System (#1864001,Bio-

Rad, USA). The ddPCR reaction mixture consisted of 10μL of 2XddPCR Supermix for

Probes (#1863026, Bio-Rad, USA), 10pg of genomic DNA, and 20Xprimers/probes mix in a

final volume of 20μL. The entire reaction mixture was loaded into a sample well of a DG8

Cartridge (Bio-Rad, USA) together with 70μL of droplet generation oil (Bio-Rad, USA) and

placed into the automated droplet generator (Bio-Rad, USA). After processing, the droplets

generated from each sample were transferred to a 96-well PCR plate (Bio-Rad,USA). PCR

amplification was carried out on a T100 Touch thermal cycler (Bio-Rad, USA) using a

thermal profile beginning at 95 °C for 10 min, followed by 40 cycles of 94 °C for 30 s, and

60 °C for 60 s, and ending of 98 °C for 10 min at a ramp rate of 2 °C/s. After PCR, the plate

was loaded on the droplet reader (Bio-Rad, USA); acquired data were analyzed with

QuantaSoft Analysis Pro software (Bio-Rad, USA). All samples had ≥10000 accepted

droplets to be considered for analysis. SatIII copies were determined by QuantaSoft, the

software coupled to the Droplet Reader QX200 that applies a Poisson algorithm to calculate

the initial concentration of DNA target molecules as units of copies/μL input based on

positive and negative droplets. Specifically, after setting an appropriate fluorescence

threshold to determine the number of positive droplets (containing at least one target

molecule), the software calculates the number of copies per μL of the reaction volume.

Student's *t*-test was applied to determine the statistical significance for independent samples

p-value. Violin plots show the median and quartiles.

Sat III expression analysis in cancer cell lines

• **Cancer cell lines and culture conditions**

22RV1, PC3, VCaP, LNCaP (Prostate cancer cell lines), were obtained from American Type

Culture Collection (ATCC) and were cultured as per the ATCC recommended guidelines.

Briefly, the cells were cultured in a growth medium supplemented with 10% fetal bovine serum

(FBS) and 0.5% Penicillin-Streptomycin (Gibco, Thermo-Fisher Scientific) in a cell culture

incubator (Thermo Fisher Scientific) supplied with 5% CO_2 at 37°C.

• **RNA isolation and Quantitative PCR (qPCR)**

The total RNA was extracted using RNAiso Plus (Takara), and 1μg of total RNA was reverse

transcribed into cDNA using First Strand cDNA synthesis kit (PureGene) according to the

manufacturer's protocol. Quantitative PCR (qPCR) was performed using cDNA template,

primer sets (sat III_FP: TATGAATTCAATCAACCCGAGTGCAATCGAA, sat III_RP: TATGGATCCTTCCATTCCATTCCTGTACTCG; GAPDH_FP: TGCACCACCAACTGCTTAGC, GAPDH_RP: GGCATGGACTGTGGTCATGAG) and SYBR Green PCR Master-Mix (PureGene) on the QuantStudio 5 Real-Time PCR System (Applied Biosystems). Relative sat III expression was calculated for each sample by the $\Delta\Delta C_T$ method. A two-tailed unpaired Student's t-test was applied to determine the statistical significance for independent samples *$P\leq 0.05$ and **$P\leq 0.001$. Error bars represent mean $\pm$ SEM.

References

Agrawal, N., Kango, M., Mishra, A. and Sinha, P. (1995). Neoplastic transformation and aberrant cell-cell interactions in genetic mosaics of lethal(2)giant larvae (lgl), a tumor suppressor gene of Drosophila. *Dev Biol* **172**, 218–229.

Aloke, C., Onisuru, O. O. and Achilonu, I. (2024). Glutathione S-transferase: A versatile and dynamic enzyme. *Biochem Biophys Res Commun* **734**, 150774.

Banani, S. F., Lee, H. O., Hyman, A. A. and Rosen, M. K. (2017). Biomolecular condensates: organizers of cellular biochemistry. *Nat Rev Mol Cell Biol* **18**, 285–298.

Baumgartner, M. E., Dinan, M. P., Langton, P. F., Kucinski, I. and Piddini, E. (2021). Proteotoxic stress is a driver of the loser status and cell competition. *Nat Cell Biol* **23**, 136–146.

Beaucher, M., Hersperger, E., Page-McCaw, A. and Shearn, A. (2007). Metastatic ability of *Drosophila* tumors depends on MMP activity. *Developmental Biology* **303**, 625–634.

Behrouzfar, K., Burton, K., Mutsaers, S. E., Morahan, G., Lake, R. A. and Fisher, S. A. (2021). How to Better Understand the Influence of Host Genetics on Developing an Effective Immune Response to Thoracic Cancers. *Front. Oncol.* **11**,.

Bendena, W. G., Garbe, J. C., Traverse, K. L., Lakhotia, S. C. and Pardue, M. L. (1989). Multiple inducers of the Drosophila heat shock locus 93D (hsr omega): inducer-specific patterns of the three transcripts. *J Cell Biol* **108**, 2017–2028.

Bennett, F. C. and Harvey, K. F. (2006). Fat Cadherin Modulates Organ Size in Drosophila via the Salvador/Warts/Hippo Signaling Pathway. *Current Biology* **16**, 2101–2110.

Berger, A. H. and Pandolfi, P. P. (2011). Haplo-insufficiency: a driving force in cancer. *J Pathol* **223**, 137–146.

Boija, A., Klein, I. A. and Young, R. A. (2021). Biomolecular Condensates and Cancer. *Cancer Cell* **39**, 174–192.

Cai, H., Holleufer, A., Simonsen, B., Schneider, J., Lemoine, A., Gad, H. H., Huang, J., Huang, J., Chen, D., Peng, T., et al. (2020). 2'3'-cGAMP triggers a STING- and NF-κB-dependent broad antiviral response in Drosophila. *Sci Signal* **13**, eabc4537.

Castro-Giner, F., Ratcliffe, P. and Tomlinson, I. (2015). The mini-driver model of polygenic cancer evolution. *Nat Rev Cancer* **15**, 680–685.

Chatterjee, M. and Sengupta, S. (2022). Human satellite III long noncoding RNA imparts survival benefits to cancer cells. *Cell Biol Int* **46**, 611–627.

Chen, S., Wei, H.-M., Lv, W.-W., Wang, D.-L. and Sun, F.-L. (2011). E2 Ligase dRad6 Regulates DMP53 Turnover in Drosophila. *J Biol Chem* **286**, 9020–9030.

Chenevix-Trench, G., Milne, R. L., Antoniou, A. C., Couch, F. J., Easton, D. F., Goldgar, D. E., and CIMBA (2007). An international initiative to identify genetic modifiers of cancer risk in BRCA1 and BRCA2 mutation carriers: the Consortium of Investigators of Modifiers of BRCA1 and BRCA2 (CIMBA). *Breast Cancer Res* **9**, 104.

Dai, C. and Sampson, S. B. (2016). HSF1: Guardian of Proteostasis in Cancer. *Trends Cell Biol* **26**, 17–28.

Datta, I., Vassel, T., Linkous, B., Odum, T., Drew, C., Taylor, A. and Bangi, E. (2023). A targeted genetic modifier screen in Drosophila uncovers vulnerabilities in a genetically complex model of colon cancer. *G3 (Bethesda)* **13**, jkad053.

de la Cova, C., Abril, M., Bellosta, P., Gallant, P. and Johnston, L. A. (2004). Drosophila myc regulates organ size by inducing cell competition. *Cell* **117**, 107–116.

Froldi, F., Ziosi, M., Garoia, F., Pession, A., Grzeschik, N. A., Bellosta, P., Strand, D., Richardson, H. E., Pession, A. and Grifoni, D. (2010). The lethal giant larvae tumour suppressor mutation requires dMyc oncoprotein to promote clonal malignancy. *BMC Biol* **8**, 33.

Fuse, Y. and Kobayashi, M. (2017). Conservation of the Keap1-Nrf2 System: An Evolutionary Journey through Stressful Space and Time. *Molecules* **22**, 436.

Goenka, A., Sengupta, S., Pandey, R., Parihar, R., Mohanta, G. C., Mukerji, M. and Ganesh, S. (2016). Human satellite-III non-coding RNAs modulate heat-shock-induced transcriptional repression. *J Cell Sci* **129**, 3541–3552.

Hopfner, K.-P. and Hornung, V. (2020). Molecular mechanisms and cellular functions of cGAS–STING signalling. *Nat Rev Mol Cell Biol* **21**, 501–521.

Johnson, T. K., Carrington, L. B., Hallas, R. J. and McKechnie, S. W. (2009). Protein synthesis rates in Drosophila associate with levels of the hsr-omega nuclear transcript. *Cell Stress Chaperones* **14**, 569–577.

Johnson, T. K., Cockerell, F. E. and McKechnie, S. W. (2011). Transcripts from the Drosophila heat-shock gene hsr-omega influence rates of protein synthesis but hardly affect resistance to heat knockdown. *Mol Genet Genomics* **285**, 313–323.

Jolly, C. and Lakhotia, S. C. (2006). Human sat III and Drosophila hsr omega transcripts: a common paradigm for regulation of nuclear RNA processing in stressed cells. *Nucleic Acids Res* **34**, 5508–5514.

Jolly, C., Konecny, L., Grady, D. L., Kutskova, Y. A., Cotto, J. J., Morimoto, R. I. and Vourc'h, C. (2002). In vivo binding of active heat shock transcription factor 1 to human chromosome 9 heterochromatin during stress. *J Cell Biol* **156**, 775–781.

Jolly, C., Metz, A., Govin, J., Vigneron, M., Turner, B. M., Khochbin, S. and Vourc'h, C. (2004). Stress-induced transcription of satellite III repeats. *J Cell Biol* **164**, 25–33.

Kandel, R., Jung, J. and Neal, S. (2024). Proteotoxic stress and the ubiquitin proteasome system. *Seminars in Cell & Developmental Biology* **156**, 107–120.

Khan, S. J., Bajpai, A., Alam, M. A., Gupta, R. P., Harsh, S., Pandey, R. K., Goel-Bhattacharya,
S., Nigam, A., Mishra, A. and Sinha, P. (2013). Epithelial neoplasia in Drosophila entails
switch to primitive cell states. *Proc Natl Acad Sci U S A* **110**, E2163-2172.

Klein, G. (2009). Toward a genetics of cancer resistance. *Proc Natl Acad Sci U S A* **106**, 859–863.

Kongsuwan, K., Yu, Q., Vincent, A., Frisardi, M. C., Rosbash, M., Lengyel, J. A. and Merriam,
J. (1985). A Drosophila Minute gene encodes a ribosomal protein. *Nature* **317**, 555–558.

Lakhotia, S. C. (2011). Forty years of the 93D puff of Drosophila melanogaster. *J Biosci* **36**, 399–
423.

Lee, T. and Luo, L. (1999). Mosaic analysis with a repressible cell marker for studies of gene
function in neuronal morphogenesis. *Neuron* **22**, 451–461.

Mah, L.-J., El-Osta, A. and Karagiannis, T. C. (2010). γH2AX: a sensitive molecular marker of
DNA damage and repair. *Leukemia* **24**, 679–686.

Mallik, M. and Lakhotia, S. C. (2009). The developmentally active and stress-inducible noncoding
hsromega gene is a novel regulator of apoptosis in Drosophila. *Genetics* **183**, 831–852.

Martin, M., Hiroyasu, A., Guzman, R. M., Roberts, S. A. and Goodman, A. G. (2018). Analysis of
Drosophila STING Reveals an Evolutionarily Conserved Antimicrobial Function. *Cell Rep*
23, 3537-3550.e6.

Martínez-Férriz, A., Ferrando, A., Fathinajafabadi, A. and Farràs, R. (2022). Ubiquitin-mediated
mechanisms of translational control. *Semin Cell Dev Biol* **132**, 146–154.

Mehta, S. and Zhang, J. (2022). Liquid–liquid phase separation drives cellular function and
dysfunction in cancer. *Nat Rev Cancer* **22**, 239–252.

Menéndez, J., Pérez-Garijo, A., Calleja, M. and Morata, G. (2010). A tumor-suppressing
mechanism in Drosophila involving cell competition and the Hippo pathway. *Proc Natl Acad*
Sci U S A **107**, 14651–14656.

Michalak, K. P. and Michalak, A. Z. (2025). Understanding chronic inflammation: couplings
between cytokines, ROS, NO, Cai 2+, HIF-1α, Nrf2 and autophagy. *Front Immunol* **16**,
1558263.

Morata, G. and Ripoll, P. (1975). Minutes: Mutants of Drosophila autonomously affecting cell
division rate. *Developmental Biology* **42**, 211–221.

Moreno, E., Basler, K. and Morata, G. (2002). Cells compete for decapentaplegic survival factor to
prevent apoptosis in Drosophila wing development. *Nature* **416**, 755–759.

Onorati, M. C., Lazzaro, S., Mallik, M., Ingrassia, A. M. R., Carreca, A. P., Singh, A. K.,
Chaturvedi, D. P., Lakhotia, S. C. and Corona, D. F. V. (2011). The ISWI chromatin
remodeler organizes the hsrω ncRNA-containing omega speckle nuclear compartments. *PLoS*
Genet **7**, e1002096.

Owusu-Ansah, E., Yavari, A., Mandal, S. and Banerjee, U. (2008). Distinct mitochondrial
retrograde signals control the G1-S cell cycle checkpoint. *Nat Genet* **40**, 356–361.

Prasanth, K. V., Rajendra, T. K., Lal, A. K. and Lakhotia, S. C. (2000). Omega speckles - a novel
class of nuclear speckles containing hnRNPs associated with noncoding hsr-omega RNA in
Drosophila. *J Cell Sci* **113 Pt 19**, 3485–3497.

Radhakrishnan, S. K., Lee, C. S., Young, P., Beskow, A., Chan, J. Y. and Deshaies, R. J. (2010).
Transcription factor Nrf1 mediates the proteasome recovery pathway after proteasome
inhibition in mammalian cells. *Mol Cell* **38**, 17–28.

Raetz, A. G. and David, S. S. (2019). When you're strange: Unusual features of the MUTYH
glycosylase and implications in cancer. *DNA Repair (Amst)* **80**, 16–25.

**Sabari, B. R., Dall'Agnese, A., Boija, A., Klein, I. A., Coffey, E. L., Shrinivas, K., Abraham, B.
J., Hannett, N. M., Zamudio, A. V., Manteiga, J. C., et al.** (2018). Coactivator condensation
at super-enhancers links phase separation and gene control. *Science* **361**, eaar3958.

Sahu, R. K., Mutt, E. and Lakhotia, S. C. (2020). Conservation of gene architecture and domains
amidst sequence divergence in the hsrω lncRNA gene across the Drosophila genus: an in
silico analysis. *J Genet* **99**, 64.

Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* **9**, 671–675.

Schulz, L., Ramirez, P., Lemieux, A., Gonzalez, E., Thomson, T. and Frost, B. (2023). Tau-
induced elevation of the activity-regulated cytoskeleton associated protein Arc1 causally
mediates neurodegeneration in the adult Drosophila brain. *Neuroscience* **518**, 101–111.

Settembre, C., Fraldi, A., Medina, D. L. and Ballabio, A. (2013). Signals from the lysosome: a
control centre for cellular clearance and energy metabolism. *Nat Rev Mol Cell Biol* **14**, 283–
296.

Shin, Y. and Brangwynne, C. P. (2017). Liquid phase condensation in cell physiology and disease.
Science **357**, eaaf4382.

Singh, A. K. and Lakhotia, S. C. (2015). Dynamics of hnRNPs and omega speckles in normal and
heat shocked live cell nuclei of Drosophila melanogaster. *Chromosoma* **124**, 367–383.

Siragusa, G., Tomasello, L., Giordano, C. and Pizzolanti, G. (2024). Survivin (BIRC5):
Implications in cancer therapy. *Life Sci* **350**, 122788.

Suijkerbuijk, S. J. E., Kolahgar, G., Kucinski, I. and Piddini, E. (2016). Cell Competition Drives
the Growth of Intestinal Adenomas in Drosophila. *Curr Biol* **26**, 428–438.

**Sulak, M., Fong, L., Mika, K., Chigurupati, S., Yon, L., Mongan, N. P., Emes, R. D. and Lynch,
V. J.** (2016). TP53 copy number expansion is associated with the evolution of increased body
size and an enhanced DNA damage response in elephants. *eLife* **5**, e11994.

Sykiotis, G. P. and Bohmann, D. (2008). Keap1/Nrf2 signaling regulates oxidative stress tolerance
and lifespan in Drosophila. *Dev Cell* **14**, 76–85.

**Szklarczyk, D., Gable, A. L., Nastou, K. C., Lyon, D., Kirsch, R., Pyysalo, S., Doncheva, N. T.,
Legeay, M., Fang, T., Bork, P., et al.** (2021). The STRING database in 2021: customizable
protein-protein networks, and functional characterization of user-uploaded gene/measurement
sets. *Nucleic Acids Res* **49**, D605–D612.

Toba, G. and Aigaki, T. (2000). Disruption of the *Microsomal glutathione S-transferase-like* gene
reduces life span of *Drosophila melanogaster*. *Gene* **253**, 179–187.

Turnbull, C., Sud, A. and Houlston, R. S. (2018). Cancer genetics, precision prevention and a call to
action. *Nat Genet* **50**, 1212–1218.

Xu, T. and Rubin, G. M. (1993). Analysis of genetic mosaics in developing and adult Drosophila
tissues. *Development* **117**, 1223–1237.

Zeng, H., Castillo-Cabrera, J., Manser, M., Lu, B., Yang, Z., Strande, V., Begue, D., Zamponi,
R., Qiu, S., Sigoillot, F., et al. (2019). Genome-wide CRISPR screening reveals genetic
modifiers of mutant EGFR dependence in human NSCLC. *Elife* **8**, e50223.

Zhang, Y., Lou, J., Liu, Y., Jin, P., Tan, Y., Song, H., Jin, W., Wang, D., Dong, F., Wu, S., et al.
(2024). Phase separation of PML/RARα and BRD4 coassembled microspeckles governs
transcriptional dysregulation in acute promyelocytic leukemia. *Proceedings of the National*
Academy of Sciences **121**, e2406519121.

Figure Legends

Figure 1. *hsrω* mutant imaginal epithelia display cell stress and undergrowth

(**A**) Diagrammatic depiction of *hsrω* gene and its multiple nuclear and cytoplasmic
transcripts (Sahu et al., 2020). The larger nuclear transcripts (RB, RG and RF) carry a long
stretch of tandem repeats, providing a physical scaffold for binding hnRNPs, such as Hrb87F
(Prasanth et al., 2000).

(**B**) Nuclear omega speckle biocondensates, marked by GFP-tagged Hrb87F (green) in wing
imaginal disc cells. Nucleus is marked by To-PRO (blue).

(**C–F**) Third instar wing imaginal discs of *hsrω*[66]/+ (C) and *hsrω*[66] (D) larvae immunostained
for phospho-Histone 3 (pH3, red) to mark dividing cells, and quantification for wing disc size
(E) and relative abundance of pH3-positive nuclei per arbitrary unit area (F).

(**G–I**) Expression of Caspase-3 (red) in *hsrω*[66]/+ (G) and *hsrω*[66] (H) wing discs and
quantification of Caspase-3 (fluorescence per ROI, I)

(**J-K**) Mosaic wing disc with *hsrω*[66] somatic clones (non-GFP) and their *hsrω*[66]/+ twins
(bright green, due to two copies of GFP). Boxed area in J is shown at a higher magnification
to reveal Caspase-3 expression (red) in *hsrω*[66] cells abutting the clone's border. Comparison
of *hsrω*[66] clone areas with the twin-spot regions (K).

Scale bars 100μm in all confocal images, except in B' which is 10 μm.

Figure 2. *hsrω*[66] epithelium reveals cellular and proteotoxic stress

(**A**) Clustered heatmap displaying status of proteins significantly altered ($\pm > \log_2$ 2-fold, $p < 0.05$, $q < 0.05$) in *hsrω*^{66} versus wild type wing imaginal disc proteome. On the right is the list of proteins upregulated (red) or downregulated (blue) in the *hsrω*^{66} proteome.

(**B**) Relative abundance ($\log_2$ fold change) of select ribosomal structural proteins in *hsrω*^{66} proteome compared to its wild type counterpart.

(**C-D**) Rank-based Gene Ontology (GO) enrichment analysis of cytosolic and mitochondrial ribosome proteins (C) and translation machinery (D). The x-axis represents the list of proteins in ascending order of fold change in *hsrω*^{66} versus wild type. Black bars along the x-axis mark the position of proteins being queried. The tables alongside give the protein levels of the top five mis-expressed proteins belonging to a given GO class.

(**E**) Relative abundance of key DNA damage-associated proteins in *hsrω*^{66} proteome.

(**F-H**) Representative examples of wing imaginal discs displaying signatures of cellular stress: ROS seen using superoxide-sensitive, Dihydroethidium (DHE, F); JNK signalling visualized using a bio-sensor, TRE-dsRED, for JNK-mediated AP-1 signalling (G); DNA double-strand breaks, marked by γH2AX (H), in *hsrω*^{66} and *hsrω*^{66}/+ counterpart and its quantification on the right. The Y-axis of the graphs represents the fluorescence per ROI.

(I-K) Signatures of increased proteotoxicity in *hsrω*^{66} versus *hsrω*^{66}/+ wing epithelium: Elevated LysoTracker fluorescence (I) and protein aggregates visualized using fluorescent dye ProteoStat (J) in *hsrω*^{66} and *hsrω*^{66} /+ wing imaginal discs. Quantification of Thioflavin fluorescence in cell homogenates of wing discs of *hsrω*^{66} and control larvae a measure of protein aggregates.

Two-tailed Student's t-test was applied to determine the statistical significance between test and controls in 2F-J; *$P \leq 0.05$ and **$P \leq 0.001$. Error bars represent mean ± SEM. ES: Enrichment Score. p-value is the Nominal p-value. A negative value of NES indicates poor enrichment of the queried geneset in *hsrω*^{66} mutant proteome versus control proteome. Scale bars: 100 μm.

Figure 3. Haploinsufficiency of *hsrω* is a tumor-permissive genetic background

(**A**) GFP-marked *lgl⁻* somatic clones (GFP, green) induced using heat shock flippase, generated in a wild type genetic background. *lgl* clones (*hsflp; lgl⁴FRT40*) display cell death (Caspase-3, red). Note enrichment of Caspase-positive *lgl* clones in basal compared to apical projections of select optical sections on the right panel.

(**B-C**) *lgl⁻* somatic clones (GFP, green) in a heterozygous *hsrω*^{66}/+ background (*hsflp; lgl⁴FRT40; hsrω*^{66}/+) undergo neoplastic transformation revealed by disrupted F-Actin. Note the characteristic large, dispersed nuclei (GFP) of transformed clones, and Caspase-3 expression along boundaries. Quantification of *lgl* clone area in wild type (WT) vs. *hsrω*^{66}/+ wing imaginal discs (C).

(D-E) *lgl⁻* clones in *hsrω⁶⁶/+* imaginal discs displaying excessive proliferation (pH3, D) and
elevated levels of Matrix Metalloproteinase (MMP1) expression (E).

(F) Graph displaying reduction in clone size of *lgl; hsrω⁶⁶/+* tumors, when larvae were fed
proteosome inhibitor MG-132 at two different concentrations.

(G-I) *lgl* somatic clones (marked in GFP) with co-down regulation *(hsflp; lgl⁴ UAS-hsrω-*
RNAi FRT40A (G-I) or up regulation (*lgl⁴ FRT40A UAS-hsrω^EP3039*; J) of in *WT* genetic
background. *lgl⁻* clones survive and undergo neoplastic transformation (stars), marked by
disrupted F-Actin (Phalloidin staining) and induce exaggerated Caspase-3 expression along
the clone boundary (G). Quantification of *lgl⁻* clone areas with loss (H) or gain (K) of *hsrω*.

Two-tailed Student's t-test was applied to determine the statistical significance between test
and controls in 3C, F. H and K; *$P \leq 0.05$ and **$P \leq 0.001$. Scale bars: 100 μm.

Figure 4. Redox regulator *Keap1/Nrf2* and inflammatory gene *dSTING* are
haploinsufficient for tumor suppression

(A-D) GFP-marked *lgl* somatic clones in a *Keap1* heterozygous genetic background (*lgl⁴;*
Keap^EY5/+, A), or upon cell autonomous downregulation of *Keap1* (*lgl⁴; UAS-Keap1 RNAi*,
B), or gain of downstream regulator CncC (*lgl⁴; UAS-CncC,* C), display neoplastic
transformation (disrupted F-Actin) and cell death along clone boundary (Caspase-3, C).

(E-F) GFP-marked *lgl* somatic clones in dSTING heterozygous (*lgl⁴; STING^ARG5/+*, E) or
STING down regulated (*lgl⁴; UAS-dSTING RNAi*, F) genetic background display neoplastic
transformation with disrupted F-Actin.
816

(G-I) Disrupted sub-cellular distribution of dSTING in cells of *hsrω⁶⁶* mutant salivary gland
(H) with enhanced punctae and reduced general cytoplasmic distribution compared with a
uniform distribution and fewer punctae in wild type gland (G). (I) Representative wing discs
from larvae bearing *lgl⁻* somatic tumors generated in *hsrω⁶⁶/+* genetic background and fed on
STING agonist, 2′3′-cGAMP, at two different concentrations; quantification of *lgl* clone size
from unfed and fed larvae (far right).

Two-tailed Student's t-test was applied to determine the statistical significance between test
and controls in 4I. NS: Not Significant.

Figure 5. Genetic variations in Sat III in human cancer patients

(A-B) Kaplan-Meier curves (generated on cBioPortal platform), comparing survival of
individuals with epithelial cancer carrying Copy Number Alterations (CNA) in *HSF1* with
individuals not carrying any alteration in *HSF1*. The y-axis represents the probability percent
survival, and the x-axis shows progress free survival in months. The plot was generated by
querying 91469 samples across 233 studies and included 16 cancer types, (A); Graph
displaying the distribution of various genetic alterations in *HSF1* across patients (B).

(C-E) Schema depicting *Sat III* locus on pericentromeric region of chromosome 9 (C),
expression of *Sat III* in benign and prostate cancer cell lines (D). Violin plot depicting
germline copy number variations of *Sat III* in PCa patients compared to benign control, BPH
(E).

Figure 6. Summary:

Schema displaying haploinsufficiency as a strategy to uncover host genetic modifiers of
cancer susceptibility. Moderate genetic variations (haploinsufficiency) transform a tumor-
resistant wild type epithelium to a genetically altered tumor-permissive epithelium (right
panel).

Supplementary data

Figure S1 Comparative Proteome of wing epithelium of *hsrω*[66] mutants versus wild type

(A-B) Box plot displaying abundance of proteins in *hsrω*[66] mutant and wild type control
proteomes (A); Volcano plot of 3198 variables depicting the comparative levels of mis-
regulated proteins in *hsrω*[66] proteome compared to wild type (B).

(C-D) Relative abundance ($\log_2$ fold change) of select proteins belonging to Protein
metabolism (C) and Stress response (D) functional classes in *hsrω*[66] proteome compared to
its wild type counterpart.

(E) Rank-based Gene Ontology (GO) enrichment analysis of RNA polII transcription
regulatory complex (GO: 00905757) and Mediator complex. The x-axis represents the list of
proteins in ascending order of fold change in *hsrω*[66] versus wild type wing disc proteome.
Black bars along the x-axis mark the position of proteins being queried. The tables alongside
give the protein levels of the top five mis-expressed proteins belonging to a given GO class.

(F) Representative examples of wing imaginal discs displaying increase in ubiquitinated
proteins using anti-Ubiquitin FK2 Antibody in *hsrω*[66] and *hsrω*[66]/+ wing imaginal discs.

Figure S2. *lgl* clones display cellular stress

(A-C) GFP-marked *lgl* mutant cells generated in a wild type epithelia exhibit signatures of
cellular stress. *lgl* clones (GFP, green) display elevated levels of ROS, (DHE, Red, A),
increase in MitoTracker staining, revealing perturbed mitochondrial activity (B), and elevated
number of acidic lysosomes as seen by LysoTracker staining (red, C). Area marked by yellow
box magnified on the right in each case. Scale bars: 100 μm.

Figure S3. Genetic disruption of omega biocondensates facilitates *lgl* tumor progression

(A-B) GFP-marked *lgl* clones generated in *Df(3R)Hrb87F/+* genetic background
(*lgl;Df(3R)Hrb87F/+,* A) or with co-downregulation of ISWI (*lgl;UAS-ISWI-RNAi,*) undergo

neoplastic transformation, as revealed by disrupted F-Actin. Transformed clones exhibit loss
of junction protein FasIII (**A**). Scale bars: 100 μm.

Figure S4 *hsrω*/+ host genetic background facilitates growth of *lgl* tumors in *Drosophila*
adult gut

(**A-C**) GFP-marked somatic control (wild type) (A) or with loss of *lgl* (B, C) clones generated
in wild type (A, B) or *hsrω^{66}*/+ (C) genetic background. Note significantly large *lgl* clones in
C compared to A or B, with characteristic small nuclei (red, stained with To-PRO) indicative
of intestinal stem cells.

Figure S5. Cellular Localization of STING

(**A**) Uniform punctae of dSTING in cells of larval salivary gland over expressing dSTING (*vg*
Gal4 >UAS-dSTING).

Figure S6. Genetic variations in *Sat III* in human cancer patients

(**A-B**) Kaplan-Meier survival curves generated on cBioPortal (cbioportal.org): comparing
survival of individuals with epithelial cancer carrying Copy Number Alterations (CNA) in
KEAP1 with individuals not carrying an alteration in *KEAP1*. The plot was generated by
querying 91469 samples across 233 studies, across 16 cancer types (A). The inset displays
distribution of various genetic alterations in *HSF1* across patients (B); comparing survival of
individuals with cancers of Lymphoid origin, carrying Copy Number Alterations (CNA) in
STING1. The plot was generated by querying 9672 samples across 17 studies.

Figure 1

A *hsrω* gene structure

Ex-1 Intron Ex-2 280bp Tandem Repeats
2382 bp 19328 bp

Transcripts
RD
RC
RA
RH
RB
RG
RF

B Hrb87F-GFP marked omega speckles

B' wing epithelium

B'' peripodium

10 uM

10 uM

Hrb87F-GFP To-PRO

Hrb87F-GFP

C *hsrω66/+*

D *hsrω66*

pH3

E Area (Arbitrary units)

**

$hsrω^{66}/+$ $hsrω^{66}$

F pH3 foci per ROI

**

$hsrω^{66}/+$ $hsrω^{66}$

G

H

Caspase-3

I fluorescence per ROI (arbitary units)

**

$hsrω^{66}/+$ $hsrω^{66}$

J *hsrω66* clones (non-GFP) and WT twin spot

Caspase-3 GFP

hsrω66 clones

wt twin clone (brighter GFP)

hsrω66/+ genetic background

K Ratio total area non-GFP/GFP (Arbitrary units)

**

+/+ $hsrω^{66}$

Figure 2

Clustered heatmap of differentially expressed proteins

A

wild type   hsrω⁶⁶

Row Z-Score

Column 1 (red):
Jon99Cii, CG5506, TotA, CG43085, Jon25Biii, CG2233, Acbp3, Arc1, Lcp65Ab2, Arc2, 18w, Lcp3, Pepck1, Jon65Aiv, Rpn7, Ldh, CG32176, CG18180, CG33306, yip7, Cyp6a23, S6k, Tsp42Ed, path, Tspo, CG10472

Column 2 (red):
vir-1, Hou, CG15784, mute, ck, CG14261, glob1, Eato, Hml, Tsf1, Idgf2, Gs1, Fbp2, Inos, Nt5b, Amy-d, CG4098, PI31, Adh, Psf2, uex, CG2076, regulcalcin

Column 3 (blue):
pst, jim, Kaz, PolA2, EG:8D8.4, CG7860, p8, Cpsf160, Ulp1, Hsp23, Esyt2, mRpL50, JIL-1, Lsp1gamma, mRpL24, gbb, GstD3, CG8034, AMPdeam, Sgs5, CG4382, RpS15Ab, ImpE2

Structural ribosomes

B — Ribosomal Proteins

Log fold (x-axis from -4 to 2):
RpS15Ab, mRpL30, mRpL54, mRpL17, mRpL3, mRpL22, mRpL39, mRpL20, mRpL24, mRpL51, mRpL13, mRpL1, mRpL45, mRpL4, mRpL50, mRpL47, mRpS30, mRpS14, mRpS4

C — GOCC:0022625 Cytosolic large ribosomal subunit

n=47
ES =0.868
FDR 1.73E-08

mRpL1	-1.697
RpL34b	-0.835
RpL36A	-0.825
RpL32	-0.767
RpL27A	-0.707

GO:0005762 Mitochondrial large ribosomal subunit

low — Ranked list — high

n=33
ES =1.869
FDR 1.26E-05

mRpL24	-2.685
mRpL50	-2.247
mRpL17	-1.795
mRpL1	-1.697
mRpL2	-1.577

Translation

D — GO:0002181: Cytoplasmic translation

low — Ranked list — high

n=99
ES =0.478
FDR 0.0001

RpS15Ab	-3.786
RpL34b	-0.835
RpL36A	-0.825
RpL32	-0.767
RpL27A	-0.707

DME-5389840: Mitochondrial translation elongation

low — Ranked list — high

n=99
ES =1.091
FDR 0.0043

mRpL50	-2.248
mRpL17	-1.795
mRpL1	-1.697
mRpL4	-1.577
mRpL51	-1.504

E — DNA Damage

Log fold (x-axis from -2 to 2):
Psf2, Irbp, RPA1 (positive, red)
JIL-1, PolA2 (negative, blue)

Cellular stress

hsrω⁶⁶   hsrω⁶⁶/+

F — Redox / DHE
**

G — JNK Signaling / TRE-DsRed
**
hsrω⁶⁶/+   hsrω⁶⁶

H — DNA Damage / γH2AX
**
hsrω⁶⁶/+   hsrω⁶⁶

Proteotoxicity

hsrω⁶⁶   hsrω⁶⁶/+

I — Lysotracker
**
hsrω⁶⁶/+   hsrω⁶⁶

J — ProteoStat
**
hsrω⁶⁶/+   hsrω⁶⁶

K — ThT Fluorescence (au) x100

Genotype:
1. Negative control
2. Positive control
3. hsrω⁶⁶/+ (10uL)
4. hsrω⁶⁶/+ (20uL)
5. hsrω⁶⁶/hsrω⁶⁶ (10uL)
6. hsrω⁶⁶/hsrω⁶⁶ (20uL)

Figure 3

lgl⁻ in *wt* genetic background

GFP Caspase-3 Actin

Apical   Basal

lgl⁻ in *hsrω⁶⁶/+* genetic background

GFP Caspase-3 Actin

GFP pH3 Actin

GFP MMP1 Actin

lgl⁻ UAS-hsrω RNAi

GFP Caspase-3 Actin

GFP Caspase-3

lgl⁻ UAS-hsrω RNAi

GFP MMP1 Actin

*lgl⁴ UAS-hsrω*EP3037

GFP Actin

X-Z

Figure 4

lgl clones (GFP) in *Keap1^{EY5}/+* epithelium

A

GFP Actin   F-Actin

lgl UAS-Keap1 RNAi (GFP)

B

GFP Actin   F-Actin

lgl UAS-Keap1 RNAi (GFP)

C

GFP Caspase-3 Actin

lgl UAS-CncC

D

GFP Actin   F-Actin

lgl clones (GFP) in *dSTING^{ΔRG5}/+* epithelium

E

GFP Actin   F-Actin

lgl UAS-dSTING RNAi (GFP)

F

GFP Actin   F-Actin

wild type   *hsrω^{66}*
Salivary Gland

G   H

To-PRO STING

I   Treatment with 2'3' cGAMP

lgl clones (GFP) in *hsrω^{66}/+* epithelium

untreated   2'3'cGAMP

100 uM   300 uM

GFP F-Actin

p=0.03
NS

Figure 5

A

HSF1

P=6.82e-10

■ Altered group
■ Unaltered group

Probability of overall survival

100%
90%
80%
70%
60%
50%
40%
30%
20%
10%
0%

Progress Free Survival (Months)

0 20 40 60 80 100 120 140 160 180 200 220 240

Altered group	639	279	133	64	34	15	9	6	2	1	1	1	0
Unaltered group	9046	4433	2225	1241	692	375	205	104	53	31	20	13	4

B

HSF1

Mutation count ((log2(value + 1))

1 2 3 4 5

1. Deep Deletion
2. Shallow Deletition
3. Diploid
4. Gain
5. Amplification

C

9q11-q12

D

Germline copy number variations in Sat III

$SatIII$ copies/µl

P=0.01

BPH(n=5) PCa(n=18)

E

Expression of Sat III in human cancer cell lines

Log_{10} $SatIII/GAPDH$ expression

10000
1000
100
10
1

PNT2 22RV1 PC3 VCaP LNCaP

Figure 7