1

A Scenic River in Rural Ohio Shows Elevated Antibiotic Resistance Genes, Including

***Acinetobacter* Tetracycline and Macrolide Resistance, Downstream of Wastewater**

Treatment Plant Effluent

By April Murphy, Daniel Barich, Siobhan Fennessy, and Joan L. Slonczewski

Address: Department of Biology, Kenyon College, Gambier, Ohio

Corresponding Author: Joan L. Slonczewski, slonczewski@kenyon.edu

ABSTRACT

Rivers in rural areas receive continual influx of wastewater carrying antibiotics originally

administered to humans and livestock. The entry of antibiotic resistance genes (ARGs) into

aquatic systems has been documented for large municipal wastewater treatment plants, but there

is less study of the impact of smaller plants that are situated on small rural rivers. We sampled

water metagenomes for ARG and taxa composition from the Kokosing State Scenic River, a

small rural river in Knox County, Ohio. Samples were obtained upstream, a few meters

downstream, and 6 km downstream from the effluent release of the Mount Vernon wastewater

treatment plant (WWTP). Through all seasons, the metagenome just downstream of the WWTP

effluent showed a substantial elevation of at least 15 different ARGs, including 6 ARGs from

Acinetobacter baumannii such as *msrE, mphE* (macrolide resistance) and *tet*(39) (tetracycline

resistance). The ARGs most prevalent near the effluent pipe persisted 6 km downstream. The

taxa distribution near the effluent showed elevation of *Acinetobacter* species as well as gut-

associated taxa, Bacteroides and Firmicutes. The ARG levels and taxa prevalence showed little

dependence on seasonal chlorination of effluent. Nitrogen and phosphorus were elevated near the

effluent pipe but had no consistent effect on ARG levels. We show that in a rural river

microbiome, year-round wastewater effluent substantially elevates ARGs including those of

multidrug-resistant *A. baumanii.*

IMPORTANCE

Antibiotic resistance is a growing problem worldwide, with frequent transmission between

pathogens and environmental organisms. Rural rivers support recreational use by people unaware

of inputs from treated wastewater. In our study, the river water proximal to wastewater effluent

shows increased prevalence of multidrug-resistant *Acinetobacter baumanii*, an opportunistic

pathogen of concern for hospitals but also widespread in natural environments. Our work

highlights the importance of wastewater effluent in management of environmental antibiotic

resistance.

INTRODUCTION

Environmental antibiotic resistance increasingly threatens global public health (1–3). Antibiotics from clinical use and livestock husbandry promote the development of resistant bacteria, and they readily pollute urban and rural waterways (4–6). Even sublethal concentrations of antimicrobial drugs select for resistance (7). Antibiotic resistance genes (ARGs) that enter environmental microbial communities have the potential to transfer to pathogenic bacteria (8). A major source of ARGs and antibiotics in aquatic systems is the effluent of wastewater treatment plants (WWTPs) (9–11). Wastewater treatment may actually select for increased antibiotic resistance of potential pathogens such as *Acinetobacter* species (12,13). It is important to understand the potential of WWTP to transfer ARGs as well as resistant microbes into rural streams, where they may disturb autochthonous microbial communities and spread drug resistance to human microbiomes. We investigated the impact of WWTP effluent on the taxa distribution and ARG counts in the Kokosing State Scenic River, a rural river designated "scenic" by the Ohio Department of Natural Resources.

The river microbiome may be affected by WWTP effluent in various ways: by elevation of phosphorus, nitrogen, and organic nutrients; by introduction of exogenous microbes; and by introduction of DNA including ARGs. The WWTP in our study chlorinates effluent only during the months of May through October, so we compared both conditions. While chlorination effectively decreases bacterial communities, it does not fully remove resistance genes from effluent. Some studies show partial decrease of ARGs by chlorine (14,15), whereas others show that chlorination may increase the effluent content of ARGs and promote their conjugative transfer (16,17). Various stress conditions in the WWTP can co-select antibiotic resistances and virulence properties (18). The release of heavy metals, antibiotics, and other compounds into

receiving rivers (such as nitrogen, phosphate, or ammonia) further propagates resistance by

selecting for multidrug efflux pumps (19–21).

The establishment of anthropogenic resistance in environmental microbial communities

can be controlled when municipalities reduce antibiotic usage (22). Therefore, understanding the

impact of ARG pollutants on rural river resistomes is important for understanding the lasting

potential of resistance in the environment. River resistomes offer the opportunity for surveillance

of opportunistic pathogens that move between environment and human host, such as the

ESKAPE pathogen *Acinetobacter baumanii* (23–25). The ESKAPE acronym comprises six

leading hospital-acquired pathogens with multidrug resistance (26). While *A. baumanii* is known

for hospital transmission, recent reports indicate community acquisition of strains that carry

ARGs on plasmids (27,28). In the Kokosing we examined evidence for *Acinetobacter* ARGs

such as *tet(39)* (29,30) and *msrE*, *mphE* (31).

To understand how WWTP effluent might alter rural river microbial communities, we

sampled sites upstream, midstream and downstream of the effluent release of the Mount Vernon

WWTP on the Kokosing River. The Kokosing river in east-central Ohio, USA, flows 92 km into

the Walhonding River, a part of the watershed of the Mississippi River (32). Our study focused

on a segment of the Kokosing in Knox County, proximal to the WWTP that serves the City of

Mount Vernon (pop. 17,000). The Mount Vernon catchment includes surrounding suburban and

rural homes as well as the Knox Community Hospital.

MATERIALS AND METHODS

Water sampling and metadata. Water samples were obtained at three sites on the Kokosing

river (**Fig. 1**). The Upstream site (coordinates 40.38368, -82.47042) lies approximately 1.5 km

upstream of the WWTP effluent discharge. The Midstream site, nearest the WWTP (40.378007,

-82.467822) is located 9 m downstream of the effluent release pipe. The downstream site

(40.376038, -82.40346) lies approximately 6 km downstream of the WWTP. The next nearest

site where wastewater enters the Kokosing is the Fredericktown WWTP, approximately 1.6 km

upstream of the Mount Vernon WWTP. The three river sites were sampled by identical

procedures at six dates of the year. Water samples were collected on October 27, November 3,

and December 3, 2019; and on April 13, May 28, and June 25, 2020. The WWTP effluent

undergoes chlorination before discharge only from May 1^{st} through October 31^{st}; thus, only the

October, May and June samples occurred during the time that effluent was chlorinated.

On each sampling date, the three sites were sampled within a 2-h period. At each site,

400 ml water was collected from the river using a dipper and sealed in a sterile bottle. The water

was transported to a lab for analysis. Within 24 h of each sample date, three 100-ml samples

were vacuum-filtered through a sterile 0.22-micron filter, 45 mm in diameter. Filter paper was

folded using sterile forceps and deposited in centrifuge tubes which were then frozen at -80°C to

preserve microbial DNA. Water was analyzed using a portable spectrophotometer, including pH,

temperature, DO, nitrogen concentration and phosphorus concentration (**Supplemental Table**

S1, Supporting Information). We chemically assayed nitrogen, phosphate and ammonia

concentrations using a Hach® DR900 Multiparameter Portable Colorimeter.

DNA isolation and sequencing. Metagenomic DNA was isolated using a ZymoBIOMICS DNA

Miniprep Kit. Each filter was cut into small pieces and transferred to a ZR BashingBead Lysis

Tube. 650 µl ZymoBIOMICS Lysis Solution was added, and all tubes were processed on a

Vortex Genie 2 for 40 min. The remainder of the preparation was performed according to the

manufacturer's protocol. Shotgun sequencing of DNA was performed by Admera Health

(www.admerahealth.com). Libraries for sequencing were prepared using Illumina's Nextera XT

DNA Library Preparation Kit, following manufacturer's instructions. Final libraries were then

pooled and sequenced on Illumina HiSeq X sequencer for 150-bp read length in paired-end

mode, with an output of 40 million reads per sample.

ARG marker analysis. Sequence reads were analyzed for ARG marker hits using ShortBRED,

a computational pipeline from Huttenhower Lab (33). ShortBRED-Identify was used to create a

database of short marker peptides specific to ARGs from the Comprehensive Antibiotic

Desistance database (CARD) (34). The CARD database of ARGs was grouped in families and

short consensus peptides were identified. The set of peptides was then filtered against the

Univeral Protein Database UNIREF90 (https://www.uniprot.org/uniref/) (data accessed October

23, 2019). This database was used to eliminate markers that match any sequences outside a

specific ARG. One additional marker, ARO_3002930 (vanRO, *Rhodococcus_hoagii*) was

removed from the marker set because inspection showed it lacked specificity. The final list of

markers (ShortBRED-2019) is presented in **Supplemental Table S2**.

The ShortBRED-2019 marker list was used to screen metagenomic reads from each of

the three river sites, from six sampling dates (**Supplemental Table S3**). Total read counts per

sample were determined using Trimmomatic (35) (**Supplemental Table S4**).

Taxa profiles. The microbial taxa were profiled using the Huttenhower lab pipeline MetaPhlAn2

(Metagenomic Phylogenetic Analysis) MetaPhlAn2 (36,37). MetaPhlAn2 assigns metagenomic

reads to taxa using a set of clade-specific marker genes. Taxa were grouped at the levels of

phylum, class, order, family, and genus (**Supplemental Table S5**).

Data Analysis. To generate an ARG heatmap from the ShortBRED data, we employed R Studio

® 1.3.073. For One-Way ANOVA analysis, we used JMP ® 14.2.0. One-Way ANOVA was

used to analyze the significance of ARG and taxa variances among the sites. ARG hits and

metadata were correlated by the Spearman rank correlation using R (**Supplemental Table S6**).

Data submission. For all DNA sequences, FASTQ files were submitted to NCBI, SRA

accession number PRJNA706754.

152
153

RESULTS

ARGs are elevated downstream of the WWTP effluent. We sought to determine how the

ARG distribution of the Kokosing River microbiomes was affected by the effluent from the

Mount Vernon WWTP. Microbial samples were obtained from three sites on the Kokosing

River, designated Upstream (1.5 km upstream of the WWTP), Midstream (just below the effluent

pipe) and Downstream (6 km downstream of the effluent pipe) (**Fig. 1**). From all sites the

metagenomic DNA sequences were analyzed for ARG prevalence using the ShortBRED pipeline

(33) applied to the CARD database (34). For each marker, the numbers of read hits were

summed across all samples and dates, and the markers were ranked according to total hits

(**Supplemental Table S3**). Results for the top 60 scoring markers are presented as a heat map

(**Fig. 2**).

Most of the top-scoring ARGs were elevated in the Midstream samples, compared to

samples from either Upstream or Downstream. The elevated ARGs include resistance

determinants from several organisms that are of clinical concern. Most striking, six of the

abundant ARGs come from the ESKAPE pathogen *Acinetobacter baumanii* and related strains:

msrE, *mphE*, *tet(39)*, CfxA6, *oxa280*, and *aadA4* (23–25). The three top-ranked ARGs (*msrE*,

mphE, *tet(39)*) are found together on *A. baumanii* plasmid pS30-1 (31). Overall, the four top *A.*

baumanii ARGs account for 37% of the total ARG hits found.

We tested whether the Downstream samples show evidence of carryover from the

Midstream site. First, the numbers of ARG hit reads were re-sorted by Midstream site ARG

totals (**Table 1**). For each of the top-ranked Midstream ARGs, we present the difference in ARG

hits between Upstream and Downstream. The top 27 most abundant Midstream ARGs, including

those from *Acinetobacter*, all show higher numbers at Downstream site than Upstream.

The overall percentage of reads that matched ARG markers ranged from 0.0015-0.0052%

for Midstream samples, and from 0.0002-0.0010% for Upstream and Downstream samples.

These numbers indicate roughly 5-fold elevation of ARG hits in the Midstream, compared to the

other two sites. We considered the possible effect of sample size, that is, whether the ARG hit

numbers reflect the number of reads in our samples (**Supplemental Table S4**). The read counts

from individual samples deviated less than 20% from the mean. There was no significant

difference in read numbers amongst the three collection sites Upstream, Midstream, and

Downstream. Thus, the elevated number of ARGs near the effluent pipe was independent of the

number of sequenced reads per sample.

Taxa profiles associated with WWTP effluent. We investigated whether the elevation of

ARGs by the WWTP was associated with specific microbial taxa. The taxa structure of our river

metagenomes was determined using the pipeline MetaPhlAn2 (36). The distribution of major

bacterial phyla and classes in our samples is shown in **Fig. 3A**, with p-values for Wilcoxon rank

sum test (**Fig. 4A**). The genus *Acinetobacter* showed a striking prevalence in the Midstream,

accounting for as high as 30% of predicted organisms (June sample, Midstream); and in some

months elevated levels persisted downstream (December and April). Given that *A. baumanii*

ARGs accounted for 37% of the total ARG hits, these results would be consistent with a high

prevalence of multidrug-resistant *A. baumanni* associated with the WWTP plant effluent.

The Midstream site showed proportions of several major taxa significantly higher than

those Upstream (**Fig. 3A**). The taxa with greater abundance include Bacteroidetes (p=0.002),

Epsilonproteobacteria (p=0.002), Gammaproteobacteria (p=0.002) and Firmicutes (p=0.005).

Downstream taxa appeared largely similar to those Upstream, with the exception of elevated

abundance of Bacteroidetes (p=0.015). These four taxa are consistent with a human fecal source,

although they were present during the period of effluent chlorination as well as during absence of

chlorination. The Upstream and Downstream sites showed higher proportions of Actinobacteria

relative to the Midstream. Alphaproteobacteria and Betaproteobacteria showed high prevalence

across all three sites. High prevalence of Actinobacteria and Betaproteobacteria are consistent

with metagenomic studies of freshwater oligotrophic lakes and rivers (38).

We considered whether the Midstream elevated ARGs might be associated with bacterial

clades that were enriched in Midstream samples. A Spearman rank correlation was performed

comparing ARG hits with the major taxa identified (**Fig. 4C**). ARGS were categorized as "Top

60" and "below 60" based on overall rank prevalence (**Fig. 2** and **Supplemental Table S3**). The

"Top 60" were those ARG classes showing relative elevation at the Midstream site near the

WWTP effluent, whereas ARGs "Below 60" (ranked below the top 60 ARGs) more likely

represent autochthonous genes commonly found in a relatively undisturbed river ecosystem. The

number of ARG hits at Midstream and Downstream showed a positive correlation with

Firmicutes and Epsilonproteobacteria, taxa that might be expected to arise from the WWTP

effluent. Negative correlations were seen between ARGs and Betaproteobacteria, which are most

likely native to the river.

If the source of "top 60" ARGs is the WWTP, are they carried by the genomes of effluent

bacteria, or do they enter the river in the form of environmental DNA? The answer is unclear

from our data. However, the occurrence of effluent chlorination (during the months of

November, December and April) shows no significant effect on the Midstream taxa profiles (**Fig**

3A). If live bacteria are responsible for ARGs elevation, significant numbers must be surviving

chlorination.

Nitrate, phosphate and ammonia levels show no correlation with elevated ARGs. The

Midstream water samples showed elevated levels of nitrate, phosphate and ammonia (**Fig. 5** and

Supplemental Table S1). We therefore looked for possible correlations between water

chemistry and ARG prevalence. Spearman rank correlations were performed for ARG levels and

various chemical and physical factors (**Supplemental Table S5**). Correlations were run

separately for the sums of Top 60 ARG hits and for the sums of Below 60 ARG hits. We

hypothesized that the top 60 ARGs are dominated by the WWTP effluent and would therefore

show stronger correlations with the Midstream chemistry.

In fact, the nitrate, phosphate and ammonia levels showed no consistent correlations with

ARGs, either Top 60 or Below 60. This finding suggests that, despite the elevation of these

minerals near the effluent pipe, nitrogen and phosphorus loading are not responsible for ARGs

elevation.

ARG numbers increased with pH and temperature. The strongest ARG correlations we did

find for water chemistry were for pH and temperature (**Supplemental Table S5**). At Midstream

and Downstream sites, pH showed a strong positive correlations with ARG hits, particularly the

Below 60 ARGs. These results suggest that low pH might select against ARGs that commonly

occur in metagenomes of the undisturbed river. In laboratory evolution experiments on

Escherichia coli, low pH and membrane-permeant aromatic acids select for loss of drug

resistance genes (39,40).

244   Temperature showed a strongly negative correlation with ARG levels, particularly those

245   Below 60. This finding would be consistent with high-temperature selection against ARGs

246   commonly found in the river community.

DISCUSSION

The footprint of the WWTP effluent release was evident across our data, including shifts in ARG

prevalence (**Fig. 2, Table 1**), microbial community taxa distribution (**Fig. 3**), and chemical

indicators (**Fig. 5**). The top three ARGs commonly occur together on *A. baumanii* plasmid pS30-

1 (31); and *A. baumanii* showed high prevalence near the WWTP effluent (**Fig. 3B**). It is

possible that the multidrug-resistant *A. baumanii* actually comes from the WWTP. Wastewater

treatment is known to increase the prevalence of multidrug resistance in *A. baumanii* from

influent to the final effluent (12). Thus, an effect of the wastewater effluent could be to increase

community exposure to drug-resistant strains of this ESKAPE pathogen.

Rural rivers have substantial economic and cultural significance for local communities, as

examplified by the state "scenic" designation and promotion of recreational activities. The public

is rarely aware of the potential impact of WWTP ARG exposure, despite the common presence

of WWTP plants along rural rivers. For example, a few km upstream of the Mount Vernon plant

is the Fredericktown WWTP; and just downstream of our sampled sites in Gambier another

small WWTP releases effluent to the Kokosing.

We found evidence that detectable levels of ARGs persist in the river microbial

community several kilometers past the effluent pipe. The Downstream site exhibited higher ARG

counts than Upstream for the top 27 ARGs elevated at Midstream (**Table 1**). Thus, WWTP-

associated ARGs persist in the environment at least 6 km downstream. These ARGs might

become established in the river microbial resistome and could propagate to pathogenic bacteria

in the future.

The WWTP-proximal site also showed substantial alteration of overall taxa distributions,

such as increased prevalence of Bacteroidetes and Firmicutes, findings consistent with previous

274 study (35). There is need for future study regarding efficient methods of ARG control from

275 WWTP, such as limiting the concentration of heavy metals in the plant to prevent co-selection of

276 multi-drug efflux pumps (19,42). Better awareness of the consequences of WWTP-effluent

277 release into rivers will improve our ability to sustain healthy microbial communities in our

278 freshwater systems.

280        **Acknowledgements**

281 This work was supported by the National Science Foundation award MCB-1923077. We

282 acknowledge contributions of students in the Kenyon College course BIOL 239 Experimental

283 Microbiology.

285
286
287

288   REFERENCES
289

1.   Bengtsson-Palme J, Larsson DGJ. Antibiotic resistance genes in the environment:

Prioritizing risks. Vol. 13, Nature Reviews Microbiology. 2015.

2.   Zhang T, Fukuda K, Topp E, Zhu YG, Smalla K, Tiedje JM, et al. Editorial: The

environmental dimension of antibiotic resistance. FEMS Microbiol Ecol. 2020;96(8).

3.   Yin X, Deng Y, Ma L, Wang Y, Chan LYL, Zhang T. Exploration of the antibiotic

resistome in a wastewater treatment plant by a nine-year longitudinal metagenomic study.

Environ Int [Internet]. 2019;133(October):105270. Available from:

https://doi.org/10.1016/j.envint.2019.105270

4.   Nesme J, Simonet P. The soil resistome: A critical review on antibiotic resistance origins,

ecology and dissemination potential in telluric bacteria. Vol. 17, Environmental

Microbiology. 2015.

5.   Pärnänen KMM, Narciso-Da-Rocha C, Kneis D, Berendonk TU, Cacace D, Do TT, et al.

Antibiotic resistance in European wastewater treatment plants mirrors the pattern of

clinical antibiotic resistance prevalence. Sci Adv. 2019;5(3).

6.   Sabri NA, Schmitt H, Van Der Zaan B, Gerritsen HW, Zuidema T, Rijnaarts HHM, et al.

Prevalence of antibiotics and antibiotic resistance genes in a wastewater effluent-receiving

river in the Netherlands. J Environ Chem Eng. 2020;8(1):102245.

7.   Gullberg E, Cao S, Berg OG, Ilbäck C, Sandegren L, Hughes D, et al. Selection of

resistant bacteria at very low antibiotic concentrations. PLoS Pathog. 2011;7(7):e1002158.

8.   Manaia CM. Assessing the Risk of Antibiotic Resistance Transmission from the

Environment to Humans: Non-Direct Proportionality between Abundance and Risk. Vol.

25, Trends in Microbiology. 2017.

9.   Rizzo L, Manaia C, Merlin C, Schwartz T, Dagot C, Ploy MC, et al. Urban wastewater

treatment plants as hotspots for antibiotic resistant bacteria and genes spread into the

environment: A review. Vol. 447, Science of the Total Environment. 2013.

10.   Karkman A, Do TT, Walsh F, Virta MPJ. Antibiotic-Resistance Genes in Waste Water.

Trends Microbiol. 2018;26(3):220–8.

11.   Hultman J, Tamminen M, Pärnänen K, Cairns J, Karkman A, Virta M. Host range of

antibiotic resistance genes in wastewater treatment plant influent and effluent. FEMS

Microbiol Ecol. 2018;94(4).

12.   Zhang Y, Marrs CF, Simon C, Xi C. Wastewater treatment contributes to selective

increase of antibiotic resistance among Acinetobacter spp. Sci Total Environ [Internet].

2009;407(12):3702–6. Available from: http://dx.doi.org/10.1016/j.scitotenv.2009.02.013

13.   Higgins PG, Hrenovic J, Seifert H, Dekic S. Characterization of Acinetobacter baumannii

from water and sludge line of secondary wastewater treatment plant. Water Res [Internet].

2018;140:261–7. Available from: https://doi.org/10.1016/j.watres.2018.04.057

14.   Quach-Cu J, Herrera-Lynch B, Marciniak C, Adams S, Simmerman A, Reinke RA. The

effect of primary, secondary, and tertiary wastewater treatment processes on antibiotic

resistance gene (ARG) concentrations in solid and dissolved wastewater fractions. Water

(Switzerland). 2018;10(1):13–8.

15.   Zheng J, Su C, Zhou J, Xu L, Qian Y, Chen H. Effects and mechanisms of ultraviolet,

chlorination, and ozone disinfection on antibiotic resistance genes in secondary effluents

of municipal wastewater treatment plants. Chem Eng J. 2017;317:309–16.

16.   Liu SS, Qu HM, Yang D, Hu H, Liu WL, Qiu ZG, et al. Chlorine disinfection increases

both intracellular and extracellular antibiotic resistance genes in a full-scale wastewater

treatment plant. Water Res. 2018;136:131–6.

17.   Guo MT, Yuan Q Bin, Yang J. Distinguishing effects of ultraviolet exposure and

chlorination on the horizontal transfer of antibiotic resistance genes in municipal

wastewater. Environ Sci Technol. 2015;49(9):5771–8.

18.   Di Cesare A, Eckert EM, Corno G. Co-selection of antibiotic and heavy metal resistance

in freshwater bacteria. J Limnol. 2016;75(2S).

19.   Seiler C, Berendonk TU. Heavy metal driven co-selection of antibiotic resistance in soil

and water bodies impacted by agriculture and aquaculture. Front Microbiol.

2012;3(DEC):1–10.

20.   Wang P, Wu D, You X, Su Y, Xie B. Antibiotic and metal resistance genes are closely

linked with nitrogen-processing functions in municipal solid waste landfills. J Hazard

Mater. 2021;403.

21.   Bernier SP, Létoffé S, Delepierre M, Ghigo JM. Biogenic ammonia modifies antibiotic

resistance at a distance in physically separated bacteria. Mol Microbiol. 2011;81(3).

22.   Berendonk TU, Manaia CM, Merlin C, Fatta-Kassinos D, Cytryn E, Walsh F, et al.

Tackling antibiotic resistance: The environmental framework. Vol. 13, Nature Reviews

Microbiology. 2015.

23.   Peleg AY, Seifert H, Paterson DL. Acinetobacter baumannii: Emergence of a successful

pathogen. Clin Microbiol Rev. 2008;21(3):538–82.

24.   Lee CR, Lee JH, Park M, Park KS, Bae IK, Kim YB, et al. Biology of Acinetobacter

baumannii: Pathogenesis, antibiotic resistance mechanisms, and prospective treatment

options. Front Cell Infect Microbiol. 2017;7(MAR).

25.   Harding CM, Hennon SW, Feldman MF. Uncovering the mechanisms of Acinetobacter

baumannii virulence. Nat Rev Microbiol. 2018;16(2):91–102.

26.   Santajit S, Indrawattana N. Mechanisms of Antimicrobial Resistance in ESKAPE

360   Pathogens. Biomed Res Int. 2016;2016.

361   27.   Chusri S, Chongsuvivatwong V, Silpapojakul K, Singkhamanan K, Hortiwakul T,

362   Charernmak B, et al. Clinical characteristics and outcomes of community and hospital-

363   acquired Acinetobacter baumannii bacteremia. J Microbiol Immunol Infect [Internet].

364   2019;52(5):796–806. Available from: https://doi.org/10.1016/j.jmii.2019.03.004

365   28.   Salgado-Camargo AD, Castro-Jaimes S, Gutierrez-Rios RM, Lozano LF, Altamirano-

366   Pacheco L, Silva-Sanchez J, et al. Structure and Evolution of Acinetobacter baumannii

367   Plasmids. Front Microbiol. 2020;11(June):1–21.

368   29.   Agersø Y, Guardabassi L. Identification of Tet 39, a novel class of tetracycline resistance

369   determinant in Acinetobacter spp. of environmental and clinical origin. J Antimicrob

370   Chemother. 2005;55(4):566–9.

371   30.   Rumbo C, Gato E, López M, Ruiz De Alegría C, Fernández-Cuenca F, Martínez-Martínez

372   L, et al. Contribution of efflux pumps, porins, and β-lactamases to multidrug resistance in

373   clinical isolates of Acinetobacter baumannii. Antimicrob Agents Chemother.

374   2013;57(11):5247–57.

375   31.   Blackwell GA, Hall RM. The Tet39 determinant and the msrE-mphE genes in

376   Acinetobacter plasmids are each part of discrete modules flanked by inversely oriented p

377   dif (XerC-XerD) sites. 2017;61(8):1–9.

378   32.   Payne JT, Millar JJ, Jackson CR, Ochs CA. Patterns of variation in diversity of the

379   Mississippi river microbiome over 1,300 kilometers. PLoS One. 2017;12(3):1–18.

33.   Kaminski J, Gibson MK, Franzosa EA, Segata N, Dantas G, Huttenhower C. High-

Specificity Targeted Functional Profiling in Microbial Communities with ShortBRED.

PLoS Comput Biol. 2015;11(12):e1004557.

34.   Alcock BP, Raphenya AR, Lau TTY, Tsang KK, Bouchard M, Edalatmand A, et al.

CARD 2020: Antibiotic resistome surveillance with the comprehensive antibiotic

resistance database. Nucleic Acids Res. 2020;48(D1):D517–25.

35.   Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible read trimming tool for Illumina

NGS data. Bioinformatics. 2014;btu170.

36.   Segata N, Waldron L, Ballarini A, Narasimhan V, Jousson O, Huttenhower C.

Metagenomic microbial community profiling using unique clade-specific marker genes.

Nat Methods. 2012;9:811–814.

37.   Truong DT, Franzosa EA, Tickle TL, Scholz M, Weingart G, Pasolli E, et al. MetaPhlAn2

for enhanced metagenomic taxonomic profiling. Nat Methods [Internet].

2015;12(10):902–3. Available from:

http://www.nature.com/doifinder/10.1038/nmeth.3589

38.   Newton RJ, Jones SE, Eiler A, McMahon KD, Bertilsson S. A Guide to the Natural

History of Freshwater Lake Bacteria. Vol. 75, Microbiology and Molecular Biology

Reviews. 2011. 14–49 p.

39.   Moore JP, Li H, Engmann ML, Bischof KM, Kunka KS, Harris ME, et al. Inverted

regulation of multidrug efflux pumps, acid resistance, and porins in benzoate-evolved

Escherichia coli K-12. Appl Environ Microbiol. 2019;85(16):1–21.

40.   He A, Penix SR, Basting PJ, Griffith JM, Creamer KE, Camperchioli D, et al. Acid

evolution of *Escherichia coli* K-12 eliminates amino acid decarboxylases and reregulates

catabolism. Appl Environ Microbiol. 2017;83:e00442-17.

41.   Price JR, Ledford SH, Ryan MO, Toran L, Sales CM. Wastewater treatment plant effluent

405   introduces recoverable shifts in microbial community composition in receiving streams.

406   Sci Total Environ. 2018;613–614.

407   42.   Barancheshme F, Munir M. Strategies to combat antibiotic resistance in the wastewater

408   treatment plants. Vol. 8, Frontiers in Microbiology. 2018.

411   TABLES

414   **Table 1. Top-ranked Midstream ARG hits: Difference in hit numbers between**

415   **Downstream and Upstream Sample Sites***

Mid Rank	ARG Family	Resistome where Found or Prevalent	Reads: Down - Up
1	*msrE*	*Acinetobacter baumannii*	60
2	*mphE*	*Acinetobacter baumannii*	64
3	*tet(39)*	*Acinetobacter baumannii*	51
4	CfxA6	*Acinetobacter baumannii*	70
5	OXA-256	*Enterobacter cloacae*	28
6	Mef(En2)	*Bacteroides fragilis*	36
7	AAC(3)-VIIa	*Streptomyces rimosus*	34
8	OXA-5	*Pseudomonas aeruginosa*	9
9	QnrS6	*Aeromonas hydrophila*	25
10	OXA-226	*Pseudomonas aeruginosa*	16
11	ANT(4_)-Ib	*Staphylococcus aureus*	38
12	*mtrA*	*Mycobacterium tuberculosis*	8
13	*sul1*	*Vibrio fluvialis*	12
14	OXA-280	*Acinetobacter johnsonii*	11
15	*tetE*	*Escherichia coli*	4
16	EreA2	*Providencia stuartii*	3
17	*mefC*	*Photobacterium damselae*	7
18	*tetQ*	*Bacteroides fragilis*	9
19	OXA-46	*Pseudomonas aeruginosa*	9
20	MOX-5	*Aeromonas caviae*	11
21	CblA-1	*Bacteroides uniformis*	2
22	ErmB	*Enterococcus faecium*	17
23	*(aadA4)*	*Acinetobacter baumannii*	1
24	*mphG*	*Photobacterium damselae*	6
25	ErmG	*Bacteroides thetaiotaomicron*	4
26	*dfrF*	*Enterococcus faecalis*	2
27	ErmF	*Bacteroides fragilis*	14

*ARGs shown represent the top 30 most abundant ARGs from Midstream sites. Values in right-

hand column indicate the difference between total Downstream reads and total Upstream reads

that match the marker shown.

FIGURES

Figure 1. Map of water sampling sites on the Kokosing River. The Upstream site is 1 km

upstream of the Mount Vernon City Wastewater Treatment Plant (WWTP). Midstream site is

located 9 m downstream of the WWTP. Downstream site is 6 km downstream of the WWTP.

Map was generated using the National Wild and Scenic Rivers System (2021).

#	ARG	Taxon of Origin or Prevalence	October			November			December			April			May			June			Total
		Family	Up	Mid	Down	Up	Mid	Down	Up	Mid	Down	Up	Mid	Down	Up	Mid	Down	Up	Mid	Down	
1	msrE	Acinetobacter baumannii	0	60	5	0	97	4	8	244	33	2	210	6	0	110	2	2	1011	22	1816
2	mphE	Acinetobacter baumannii	0	56	6	0	99	4	2	174	29	2	178	9	0	95	0	0	964	20	1638
3	tet(39)	Acinetobacter baumannii	0	26	6	0	100	2	3	101	30	0	83	11	0	124	2	0	154	3	645
4	CfxA6	Acinetobacter baumannii	0	21	10	0	36	0	0	207	50	0	63	6	2	48	2	0	85	4	534
5	AAC(3)-VIIa	Streptomyces rimosus	10	119	43	9	40	20	11	44	8	13	7	13	11	9	5	12	13	11	398
6	ANT(4)	Staphylococcus aureus	0	73	98	0	0	0	17	21	11	0	0	1	61	80	0	1	1	7	371
7	OXA-256	Enterobacter cloacae	3	65	16	0	48	5	2	97	11	0	29	2	3	32	2	0	54	0	369
8	Mef(En2)	Bacteroides fragilis	0	11	2	0	17	0	2	88	25	0	42	1	0	23	10	0	52	0	273
9	OXA-5	Pseudomonas aeruginosa	0	28	1	0	50	0	3	32	10	0	41	0	0	15	1	0	62	0	243
10	QnrS6	Aeromonas hydrophila	0	16	0	0	29	3	0	65	16	1	33	4	0	27	0	0	29	3	226
11	OXA-226	Pseudomonas aeruginosa	0	13	7	0	22	0	2	73	10	0	13	0	0	19	1	0	51	0	211
12	mtrA	Mycobacterium tuberculosis	0	14	2	2	12	3	3	20	1	0	24	0	0	9	2	3	95	8	198
13	sul1	Vibrio fluvialis	0	42	2	0	12	0	1	53	6	0	11	3	0	14	2	2	40	2	190
14	OXA-280	Acinetobacter johnsonii	0	7	0	0	23	1	0	34	10	0	8	0	1	21	0	0	67	1	173
15	tet(E)	Escherichia coli	0	2	0	0	32	2	1	40	5	2	25	0	0	1	0	0	43	0	153
16	mefC	Photobacterium damselae	0	14	2	0	32	0	0	22	4	0	8	0	0	20	1	0	31	0	134
17	EreA2	Providencia stuartii	0	58	0	0	32	2	0	13	1	0	0	0	0	5	0	0	19	0	130
18	ACT-35	Enterobacter cloacae	3	5	6	8	4	9	8	6	16	6	4	9	13	6	1	5	6	4	119
19	OXA-156	Pandoraea pulmonicola	3	10	4	2	0	7	3	14	7	12	9	10	3	13	5	1	3	7	113
20	tetQ	Bacteroides fragilis	0	0	0	0	6	0	0	59	7	0	11	2	0	3	0	0	14	0	102
21	OXA-46	Pseudomonas aeruginosa	0	12	0	0	16	0	0	29	4	0	12	4	1	4	0	0	17	2	101
22	ErmB	Enterococcus faecium	0	1	0	0	2	0	0	34	17	0	17	0	1	6	0	0	15	1	94
23	MOX-5	Aeromonas caviae	0	6	1	0	8	0	1	6	8	0	10	0	0	11	2	0	37	1	91
24	Erm(O)-lrm	Streptomyces lividans	0	2	0	3	3	3	2	2	3	11	9	3	2	18	1	9	9	3	83
25	CblA-1	Bacteroides uniformis	0	1	0	0	7	0	0	31	1	0	11	1	0	13	0	0	15	0	80
26	SHV-100	Klebsiella pneumoniae	2	13	7	2	2	3	2	4	1	4	5	4	5	7	2	3	10	2	78
27	mphG	Photobacterium damselae	0	6	2	0	10	0	0	19	3	0	7	0	0	6	0	0	21	1	75
28	OXA-443	Ralstonia mannitolilytica	5	3	4	5	7	5	2	15	0	0	5	1	3	4	4	2	7	3	75
29	aadA4	Acinetobacter baumannii	0	2	0	0	36	0	0	16	1	0	2	0	0	13	0	0	4	0	74
30	ErmF	Bacteroides fragilis	0	0	0	0	1	2	0	36	5	0	5	2	0	3	5	0	12	0	71
31	cmlA5	Gram-negative species	0	1	0	4	15	0	1	22	2	1	7	2	1	2	1	0	9	1	69
32	FOX-3	Klebsiella oxytoca	0	0	6	1	7	0	1	11	0	2	6	0	1	8	5	3	15	3	69
33	ErmG	Bacteroides thetaiotaomicron	0	1	0	0	5	1	0	38	3	0	14	0	0	3	0	0	2	0	67
34	bcrC	Bacillus licheniformis	0	3	0	6	3	0	2	29	2	1	5	0	0	2	0	2	8	1	64
35	OXA-137	Brachyspira pilosicoli	6	5	2	4	7	1	1	11	0	5	8	3	2	3	1	1	3	1	64
36	dfrF	Enterococcus faecalis	0	0	0	0	4	0	0	25	0	0	10	1	0	10	0	0	12	1	63
37	tetW	Butyrivibrio fibrisolvens	0	0	4	0	1	0	0	29	0	0	5	0	5	5	2	2	9	0	62
38	MOX-9	Citrobacter freundii	2	1	10	0	1	3	1	10	1	0	4	1	3	6	7	6	6	0	62
39	mefB	Escherichia coli	0	5	0	0	4	0	0	17	2	0	5	0	0	7	0	0	13	0	53
40	qacH	Vibrio cholerae	0	4	0	0	12	0	2	16	0	0	5	1	0	3	0	0	10	0	53
41	AER-1	Aeromonas hydrophila	0	8	2	0	5	0	0	13	1	0	4	0	0	0	0	0	17	0	50
42	ANT(3)	Serratia marcescens	0	2	1	0	7	0	0	26	0	0	1	0	0	7	0	0	4	1	49
43	EreD	Riemerella anatipestifer	0	2	0	0	14	0	0	7	0	0	0	0	2	8	0	0	14	2	49
44	aadA27	Acinetobacter lwoffii	0	2	0	0	7	2	4	8	1	3	3	5	0	5	0	3	4	2	49
45	GES-21	uncultured bacterium	2	1	0	0	1	0	0	19	0	0	9	0	0	1	0	0	15	0	48
46	lnuC	Streptococcus agalactiae	0	3	0	0	6	0	0	16	3	2	6	1	0	5	0	0	6	0	48
47	sul2	Vibrio cholerae	0	9	0	0	6	0	0	14	0	0	2	0	0	8	0	0	3	0	42
48	mphA	Escherichia coli	0	4	0	0	3	0	0	18	3	0	2	0	0	0	0	0	11	1	42
49	APH(6)-Id	Pseudomonas aeruginosa	0	6	0	0	4	0	0	16	2	0	5	0	0	0	0	0	8	0	41
50	OXA-31	Pseudomonas aeruginosa	0	7	0	0	7	0	2	13	1	0	1	0	0	5	0	0	5	0	41
51	tet(L)	Geobacillus stearothermophilus	0	12	7	0	0	0	0	2	2	0	0	0	9	9	0	0	0	0	41
52	lsaE	Enterococcus faecalis	0	0	1	0	3	0	0	14	2	0	7	0	0	1	0	0	12	0	40
53	imiS	Aeromonas veronii	0	0	0	0	5	0	2	14	4	0	3	0	0	5	0	0	4	0	37
54	tet(40)	Clostroides sp.	0	0	2	0	0	0	0	17	4	0	5	0	0	4	1	0	4	0	37
55	ANT(3)	Escherichia coli	0	5	1	0	0	2	0	9	2	0	4	1	0	5	0	0	4	1	34
56	OXA-45	Pseudomonas aeruginosa	0	2	0	0	1	0	0	16	0	0	1	0	0	5	0	0	7	0	32
57	mel	Streptococcus pyogenes	0	0	0	0	2	0	0	13	0	0	6	1	0	0	0	0	9	0	31
58	APH(3)-IIIa	Campylobacter coli CVM N29710	0	1	0	0	1	0	0	11	2	0	1	0	0	6	0	0	7	0	29
59	tetX	Bacteroides fragilis	0	1	0	0	1	0	0	6	1	0	5	0	0	4	1	0	8	0	27
60	aadS	Transposon Tn4551	0	2	0	0	1	0	0	10	2	0	0	1	0	9	0	0	2	0	27

434
435

436 **Figure 2**. **Heatmap of relative abundance of top 60 ARG marker hits.** Read hit numbers are

437 ranked in descending order by total hits across samples. Yellow represents highest abundance,

438 cyan represents lowest abundance.

A.

B.

440   **Figure 3. ARGs and bacterial taxa distribution across sampling sites and dates.**

441   **A.** Percentages of phyla and of proteobacterial classes predicted by MetaPhlAn2. Taxa with

442   prevalence too small to be quantified were grouped as "Other." Samples are sorted by site, then

443   by sampling date. Horizontal black bars indicate dates when effluent was unchlorinated.

444   **B.** Percentage of *Acinetobacter* species predicted by MetaPhlAn2.

A.

	Alphaproteobacteria	Betaproteobacteria	Epsilonproteobacteria	Gammaproteobacteria	Actinobacteria	Bacteroidetes	Cloacimonetes	Cyanobacteria	Firmicutes
Midstream -- Downstream	0.18	0.093	0.015	0.002	0.026	0.015	0.608	0.199	0.002
Upstream -- Downstream	0.699	0.818	0.093	0.093	0.589	0.015	0.074	0.937	0.375
Midstream -- Upstream	0.093	0.132	0.002	0.002	0.004	0.002	0.074	0.093	0.005

B.

Taxa and ARG prevalence: Top 60 and Below 60	Upstream Top 60	Upstream Below 60	Midstream Top 60	Midstream Below 60	Downstream Top 60	Downstream Below 60
Firmicutes	0.55	0.84	0.43	0.71	0.89	0.83
Bacteroidetes	-0.20	0.31	0.66	0.49	0.49	0.77
Cyanobacteria	0.54	0.26	0.03	-0.49	-0.03	0.20
Actinobacteria	-0.20	-0.14	-0.09	-0.26	0.03	-0.31
Alphaproteobacteria	-0.49	-0.20	-0.54	-0.54	0.20	0.60
Betaproteobacteria	0.14	0.31	-0.77	-0.94	-0.37	-0.20
Gammaproteobacteria	0.66	0.60	0.20	0.09	0.43	0.60
Epsilonproteobacteria	-0.09	0.54	0.60	0.31	0.60	0.83

Figure 4. Taxa correlations across river sites.

A. Wilcoxon rank sum test p-values for comparison of taxa percentages at different sites, grouped across all dates. p-values ≤ 0.02 indicate significant differences (highlighted).

B. Spearman rank correlations between bacterial taxa and ARG abundance.

Figure 5. Nitrate, phosphate and ammonia concentrations across sites and dates.

Concentrations of nitrate, phosphorus and ammonia were measured at each river site for each

month. Full metadata are presented in **Supplemental Table S1**; and metadata correlations with

ARG abundance in Supplemental **Table S6**.