Nasal BCG Exposure Accelerates Dural Lymphatics Development via Macrophages'

Role in Newborn Mice

Junhua Yang (ORCID: 0000-0001-8037-9644)[1#], Lifang Yuan(ORCID:

0000-0002-6315-2824)[2,#], Linyang Song(ORCID: 0000-0002-2770-9568)[1], Fangfang

Qi[3], Zejie Zuo[3], Jie Xu[3], Zhibin Yao(ORCID: 0000-0003-1613-9157)[3,*]

[1] Department of Anatomy, School of Biosciences & Biopharmaceutics, Guangdong

Pharmaceutical University, Guangzhou 510006, PR China.

[2] School of Public Health (Shenzhen), Sun Yat-sen University, Shenzhen 510275, PR

China.

[3] Department of Anatomy and Neurobiology, Zhongshan School of Medicine, Sun

Yat-Sen University, Guangzhou 510080, PR China.

[#]Junhua Yang and Lifang Yuan contributed equally to this study and share first

authorship.

[*]Corresponding author. Fax: +862087330709. E-mail address: yao.zb@163.com

(Zhibin Yao).

Abstract

The dural lymphatics develop mainly during the first postnatal month. Lymphatics may be shaped by immune activation when bacterial infection happens. BCG, a strong immune activator, is widely injected to newborns. Moreover, human beings are nasally exposed in daily life to bacterial stimuli. These background prompted us to investigate whether neonatal BCG injection combined with nasally exposure exerts an influence on dural lymphatics develop. Here, mice received a single dose intracutaneous (i.c.) BCG injection immediately after birth followed by repeated nasal BCG challenge once a day (i.c./nasal group). These mice had an accelerated dural lymphatics growth and increased levels of several cytokines and VEGFR-3. Furthermore, macrophages were identified as a key mediator of these alterations. Mice that received mere one dose i.c. BCG injection showed no significant alterations in these indexes. Additionally, mere repeated nasal BCG challenge induced similar effects to i.c./nasal challenge but with a slighter extent. Taken together, these findings show that repeated nasal BCG vaccination accelerates dural lymphatics development in neonatal mice, especially in the presence of neonatal i.c. BCG injection.

Keywords: Meningeal lymphatic vessels; Development; Vaccine; Central nervous system; Mucosal immunity

Introduction

Since two publications in 2015 describing the structure and function of the dural lymphatics (*1, 2*), these vessels have drawn increasing attention. A series of researches as well as review articles focused on them (*3, 4*). In 2017, Salli Antila et al. reported the development and plasticity of dural lymphatics in mice, revealing that the dural lymphatics network starts to develop after birth and they were already well developed at about four weeks old (*5*). As well known, lymphatics may be shaped by inflammatory factors produced in immune activation. It has been reported that immune activation induced by bacterial or virus infection led to lymphangiogenesis outside the central nervous system (*6-8*). BCG is a live attenuated vaccine containing live *Mycobacterium tuberculosis* and is widely used to newborns. Planned BCG vaccination is a strong and unnatural neonatal immune stimuli. About 90 percent of infants of the world were administrated neonatally with BCG (*9*).

We here imitated the immune challenge by BCG that happens to human infants, by means of a single dose i.c. BCG injection immediately after birth. Moreover, we also imitated the daily nasal exposure to microbiota that exist in the air by repeated nasal challenge with BCG performed once a day within the first 14, 19 or 21 postnatal days. For the nasal immune challenge with bacteria, BCG was used for three reasons: 1) this constitutes a boosting immunization when combined with neonatal i.c. BCG injection in spite of a different exposure pathway, 2) in the long period of human evolution and survival, *Mycobacterium tuberculosis*, as a kind of bacteria mainly transmitted through the air, has been exposed to almost all human individuals (*10*), 3)

despite advances in modern medicine, one third of the world's population is estimated

to have contracted the bacteria, with estimates of ten million new infections globally

each year (*11*). Meanwhile, mice receiving PBS, mere one dose i.c. neonatal BCG

injection or mere repeated nasal BCG challenge were also prepared and observed. We

found that neonatal BCG vaccination accelerates lymphatics development in mice,

especially in the presence of neonatal i.c. BCG exposure.

Materials and methods

Animals and study design

Newborn litters of C57Bl/6 mice were obtained from the SYSU Laboratory Animal

Center (Guangzhou, China) and were housed in a specific pathogen-free facility. This

study consisted of different tests, each of which used four groups of mice, namely

con-group, intracutaneous(i.c.)-group, nasal-group and i.c./nasal-group. All

experiments in the study were started on postnatal day 0 (P0).

For each test, six newborn litters of C57Bl/6 mice were used. Within each litter, four

male pups were used and randomly divided into four groups, receiving i.c. PBS

injection combined with nasal PBS administration (con-group), i.c. BCG injection

combined with nasal PBS administration (i.c.-group), i.c. PBS injection combined

with nasal BCG vaccination (nasal-group) or i.c. BCG injection combined with nasal

BCG vaccination (i.c./nasal-group). The pups that were sacrificed after P21 were

weaned on P21. The study was approved by the SYSU Institute Research Ethics

Committee and all animal experiments were conducted in compliance with the

relevant laws and institutional guidelines.

Immunization procedures

A single dose of BCG was administered (i.c.) to the mice in i.c.-group and

i.c./nasal-group in the back on P0. For nasal-group and i.c./nasal-group, each mouse

received nasal administration of BCG once a day from P0 to P14, P19 or P21.

Freeze-dried living BCG (D2-BP302 strain, Biological Institute of Shanghai, China)

was suspended in PBS. The dose both of i.c. injection and nasal administration is

about 10^5 CFU.

Immunofluorescence staining and quantification

To test the proliferating cells in dural lymphatics, the mice received five

intraperitoneal injections of BrdU (Sigma-Aldrich, 50 mg/kg) from P14 once a day to

label the dividing cells. At P19, the animals were deeply anesthetized with chloral

hydrate (10% solution) and then transcardially perfused with PBS. After removing the

mandibles and the skull rostral to the maxillae, the skull cap and the attached

dura/arachnoid containing superior sagittal sinus (SSS) and two transverse sinus (TS)

areas were removed with surgical scissors. The dura mater tissue used for all the other

tests were obtained from mice at P14 or P21 that were not treated with BrdU. Whole

mounted meninges that were still attached to the skull cap were placed in 4%

paraformaldehyde (PFA) in PBS and postfixed for 24 h at 4°C. The dura/arachnoid

were then carefully and gently separated from the skull cap using tweezers with a thin

head. Actually, the adhesion between dura mater and calvaria is relatively loose in

young mice, and therefore it is easy to keep them intact during peeling from calvaria.

The dura were then washed with PBS three times and processed for staining. The dura

were blocked in PBS containing 1% bovine serum albumin and 0.25% Triton X-100

(Sigma-Aldrich, St. Louis, MO, USA) for 1 h at 37°C before being incubated with

primary antibodies diluted in the same blocking solution overnight at 4°C. The

primary antibodies were as follows: goat anti-Lyve-1 (R&D Systems, Minneapolis,

MN, USA; 1:600), rabbit anti-Prox-1 (Abcam, Cambridge, MA, USA; 1:500), rat

anti-BrdU (Santa Cruz, CA, USA;1:500), goat anti-VEGFR-3 (R&D Systems,

Minneapolis, MN, USA; 1:50) and rat anti-F4/80 (Abcam, Cambridge, MA, USA;

1:200). Specimens were washed three times with PBS before being incubated with the

following secondary antibodies for 2 h at 37°C: Alexa Fluor 555-conjugated donkey

anti-goat, Alexa Fluor 647-conjugated donkey anti-rat and Alexa Fluor

488-conjugated donkey anti-rabbit antibodies. All secondary antibodies (Invitrogen

Molecular Probes®, Eugene, Oreg., USA) were diluted 1:400. Each of the dura

samples was then placed on a glass slide with the skull face attached on the slide. This

process was performed in PBS to ensure that the SSS area and two TS areas stretched

without folding.

Confocal micrographs of immunofluorescence staining were obtained using a LSM

780 confocal laser-scanning microscope (Zeiss, Heidelberg, Germany). The total

numbers of each kind of labeled cells showed in Results section within the selected

areas were quantified using a Stereo Investigator (MicroBrightField, Williston, USA)

stereology system by an investigator who was blinded to the groups.

Multiplex assay

Mice at P21 were anesthetized with chloral hydrate (10% solution). Then, 5-7 µl of cerebrospinal fluid (CSF) were obtained from the cisterna magna of each mouse, according to a previously described protocol (*12*). The mouse cytokine/chemokine panel (MCYTOMAG-70K-04; Millipore, Billerica, MA, USA) and bead-based Luminex system were used to detect the concentrations of IFN-γ, IL-4, TNF-α, IL-7, IL-17, IL-1β, IL-6, IL-10 and monocyte chemotactic protein-1 (MCP-1) in the CSF. We first added 25 µl of prepared CSF samples that had previously been diluted 1:5 in assay buffer to each well. Then, 25 µl of the magnetic beads coated with specific antibodies were added, and the mixture was incubated overnight at 4°C. Then, the beads were washed and incubated with 25 µl of the biotinylated detection antibody for 1 h at room temperature (RT). Afterwards, 25 µl of the streptavidin-phycoerythrin conjugate were added, and the reaction was incubated for 30 min at RT. The beads were then washed with 150 µl of sheath fluid for 5 min at RT. Data were collected on a Bio-Plex-200 system (Bio-Rad, Hercules, CA, USA) and analyzed using professional software (Bio-Plex Manager; Bio-Rad, CA, USA).

ELISA

Mice at P21 were anesthetized with chloral hydrate (10% solution). Then, 5-7 µl of CSF were obtained from the cisterna magna of each mouse, according to a previously described protocol (*12*). VEGF-C levels were assayed in the CSF of mice using an ELISA kit (VEGF-C, Cusabio, China), according to the manufacturer's instruction.

Western blot

At P21, levels of the VEGFR-3 proteins in the dura were examined by western blot

analyses. After the animals were anesthetized and perfused with PBS, the skull cap

and the attached dura/arachnoid were obtained. Before and after the dura/arachnoid

were separated from the skull cap, the tissues were washed three times with ice-cold

PBS to remove the CSF and intercellular fluid that may have adhered to the meninges.

Then, the dura/arachnoid were homogenized in ice-cold RIPA buffer. The homogenate

was centrifuged at 9,000 g for 15 minutes at 4°C, after which the supernatant was

analyzed by western blotting. Protein concentrations were quantified using the

bicinchoninic acid (BCA) protein assay kit (Biotime, China). Membrane homogenates

(50 μg) were separated on 6% SDS-PAGE gels and transferred onto presoaked PVDF

membranes (Bio-Rad, Pleasanton, CA, USA). Blots were then blocked with 10%

non-fat milk in PBST (100 nM phosphate buffer, pH 7.5, containing 150 nM NaCl

and 0.1% Tween-20) for 2 h at RT. The the membranes were incubated overnight at

4°C with rabbit anti-VEGFR-3 antibodies (Abcam, Cambridge, MA, USA; 1:50).

Horseradish peroxidase (HRP)-conjugated secondary goat anti-rabbit antibodies

(Sigma-Aldrich Co., USA) were used at a dilution of 1:4000. Protein bands were

visualized using a chemiluminescence detection system (Pierce, Milan, Italy). Optical

densities were measured by dividing the intensity of the bands by the intensity of the

housekeeping protein β-actin (Cell Signaling Technology, MA, USA, 1:1000). Data

from control samples were assigned a relative value of 100%.

Statistical analyses

All data were statistically analyzed using the Statistical Package for the Social

Sciences 25.0. The data were analyzed using randomly completely block design

ANOVA followed by Bonferroni's *post hoc* test or paired-samples *t* test. The

statistically significant level was set at $\alpha = 0.05$. The normality and homogeneity of

variance were conducted using the Shapiro-Wilk test and Levene's test, respectively.

Results

Accelerated dural lymphatics development in nasal-group and i.c./nasal-group in

TS zone at P14

At P14, lymphatics distribution around the TS was observed by

immunofluorescencely staining for Lyve-1. Lyve-1$^+$ vessels covered about 59%, 63%,

85% and 100% of the length of ipsilateral TS in con-group, i.c.-group, nasal-group

and i.c./nasal-group, respectively (Fig.1). Compared with con-group, both nasal-group

and i.c./nasal-group had a significant increase in the length of TS covered by Lyve-1$^+$

vessels, Lyve-1$^+$ vessels area, numbers of sprouting from Lyve-1$^+$ vessels and

numbers of cross points of Lyve-1$^+$ vessels (Fig.1). In all these four indexes, the

i.c./nasal-group possessed a significant larger extent of increase than the nasal-group

(Fig.1). Moreover, the i.c./nasal-group showed obvious sprouting and extending of

lymphatics from the TS zone toward both rostral (upward) and caudal side (downward)

in the dura mater (Fig.1D). All these findings indicated accelerated dural lymphatics

development by neonatal nasal exposure to BCG.

Accelerated dural lymphatics development with more macrophage-derived
193

lymphatic endothelial cell progenitors (M-LECPs) in SSS zone in nasal-group
194

and i.c./nasal-group at P21
195

Macrophages (F4/80$^+$) play a vital role in lymphangiogenesis induced by immune

stimuli because of their capacity to transform into lymphatic endothelial progenitors

(*13, 14*). F4/80$^+$/Lyve-1$^+$ cells, F4/80$^+$/Prox-1$^+$ cells and F4/80$^+$/Lyve-1$^+$/Prox-1$^+$ cells

have previously been reported as the macrophage-derived LEC progenitors (*13, 14*).

In the current study, lymphatics distribution and macrophages presence around the

SSS was observed after the dura mater was immunofluorescencely trible-stained for

Lyve-1/Prox-1/F4/80 at P21. Lyve-1$^+$ vessels covered about 30%, 50%, 70% and 100%

of the length of SSS in con-group, i.c.-group, nasal-group and i.c./nasal-group,

respectively (Fig.2). Compared with con-group, both nasal-group and i.c./nasal-group

had a significant increase in Lyve-1$^+$ vessels area%, numbers of sprouting from

Lyve-1$^+$ vessels and Prox-1$^+$ nucleus numbers (Fig.2). In all these three indexes, the

i.c./nasal-group possessed a significant larger extent of increase than the nasal-group

(Fig.2). These indexes also indicated accelerated dural lymphatics development by

BCG, especially considering the fact that lymphatics begin to cover 100% of the

length of SSS at P28 under normal condition (*5*).

Compared with con-group, both nasal-group and i.c./nasal-group had a significant

increase in the number of total F4/80$^+$ cells (Fig.2H), F4/80$^+$/Prox-1$^+$ cells and

F4/80$^+$/Prox-1$^+$/Lyve-1$^+$ cells (Fig.3M-O). In all these three indexes, the

i.c./nasal-group possessed a significant larger extent of increase than the nasal-group

(Fig.2H) (Fig.3M-O). In addition, M-LECPs in different stages were found, indicating

by different features of phenotype and location to lymphatics (See detailed description

in Fig.4 and the corresponding legend).

Elevated real time lymphangiogenesis level in SSS zone in nasal-group and

i.c./nasal-group at P19

The real time lymphangiogenesis level was also evaluated by

immunofluorescencely co-staining for Lyve-1/Prox-1/BrdU at P19 and the SSS zone

of dura mater was observed. Fig.S1A-D shows whole length graph of SSS area from

which representative higher scopes were presented in Fig.5. As expected, both

nasal-group and i.c./nasal-group had a significant increase in BrdU$^+$ cells (Fig. S1F),

BrdU$^+$/Prox-1$^+$ (Fig.5M) and BrdU$^+$/Prox-1$^+$/Lyve-1$^+$ (Fig.5N) cells within the entire

area of SSS compared with con-group besides the increased indexes that reflecting the

accelerated dural lymphatics development (Fig. S1E, G-H). In both indexes, the

i.c./nasal-group possessed a significant larger extent of increase than the nasal-group.

Pro-lymphangiogenesis and macrophage-recruting profile of cytokine expression

in the CSF from mice in nasal-group and i.c./nasal-group at P21

Lymphangiogenesis is a dynamic process that is regulated also by other cytokines,

such as vascular endothelial growth factor-C (VEGF-C), monocyte chemotactic

protein-1 (MCP-1), interleukin-1β (IL-1β), IL-4, IL-6, IL-7, IL-10, IL-17, interferon-γ

(IFN-γ), and transforming growth factor-β (TGF-β) (*6, 8, 15-20*). Therefore, we

determined their concentrations in the CSF and serum of each group of mice. In the

CSF, both nasal-group and i.c./nasal-group had a significant increase in the level of

VEGF-C, MCP-1, IFN-γ and TGF-β compared with con-group (Fig.6). The

i.c./nasal-group possessed a significant larger extent of increase than the nasal-group

(Fig.6). The other tested cytokines in the CSF were of no significant differences

among groups or undetectable in all groups. In serum, both nasal-group and

i.c./nasal-group had a significant increase in the level of MCP-1, IFN-γ, TGF-β, IL-4

and IL-10 compared with con-group (Fig.7). The i.c./nasal-group possessed a

significant larger extent of increase than the nasal-group (Fig.7). The other tested

cytokines in serum were of no significant differences among groups or undetectable in

all groups.

Besides, the level of VEGFR-3, the receptor for VEGF-C and mainly expressed by

lymphatics, was also detected in dura mater because of its vital role in

lymphangiogenesis (*21, 22*). WB analysis showed both nasal-group and

i.c./nasal-group had a significant increase in VEGFR-3 expression (Fig.8). In both

indexes, the i.c./nasal-group possessed a significant larger extent of increase than the

nasal-group (Fig.8).

Macrophage identified as a mediator of BCG-induced promotion in dural

lymphatics development.

VEGF-C, mainly secreted by macrophages, is the most important cytokine for

promoting lymphangiogenesis (*16, 17, 23*). MCP-1 is the most potential chemokine

for recruting macrophages (*24*). Therefore, the increased levels of them in the CSF in

nasal-group and i.c./nasal-group may explain the larger number of macrophages

counting in nasal-group and i.c./nasal-group as shown in Fig.2H, Fig.3 and Fig. S1.

To further confirm the mediating role of macrophages in BCG-induced promotion

in dural lymphatics development, each group of mice were intraperitoneally injected

with a macrophage-depleting agent clodronate liposomes (110 mg/kg, once every

three days). At P28, dura mater were stained for Lyve-1$^+$ vessels. The mean Lyve-1$^+$

vessels area in SSS zone in CL/con-group (Fig.S2D) is less than that of con-group at

P21 (Fig.2A, Fig. S1A). Moreover, there is no longer any significant difference

among four groups in either Lyve-1$^+$ vessels area in SSS zone or in the level of

VEGF-C in CSF after macrophage-depletion (Fig.S2). What's more, rare Lyve-1$^+$

cells out of Lyve-1$^+$ vessels could be observed in dura mater when macrophages were

depleted.

Discussion

This study for the first time revealed that repeated neonatal BCG vaccination, either

combined with or without i.c. injection of BCG, accelerated dural lymphatics

development in mice. Furthermore, macrophages were identified as the mediator of

BCG-induced dural lymphatics development acceleration.

A notable phenomenon found in this study is that a single dose of BCG i.c.

injection resulted in no significant influence in dural lymphatics development at P21.

This may be because of the lack of boosting immunization. The possible role of

boosting immunization was also supported by the fact that the i.c./nasal-group

consistently possessed a significant larger extent of alteration than the nasal-group in

nearly all indexes. The present study aimed mainly to explore the potential influence

of the immune activation pattern 'neonatal i.c. injection/nasal immunization of

mycobacterium tuberculosis' that happens to people worldwide. Therefore, boosting

immunization by repeating i.c. injection of BCG was not attempted in this study.

BCG immunization could not only induce Th1 cytokines production such as IFN-γ

(*25*) but also activate macrophages (*26*). This was confirmed by the results of

cytokines analysis in this study. Macrophages is the major source of VEGF-C protein

(*16*). In the nasal-group and i.c./nasal-group, the elevated VEGF-C levels in the CSF

(Fig.6A) may be explained by the increased numbers of macrophages in the dura

mater. In turn, the larger number of recruited macrophages in the dura mater may be

explained by the higher CSF level of MCP-1 that is a very potential chemokine for

monocyte/macrophages (*27*).

It is reported that BCG could induce systemic release of MCP-1 (*28*). However,

although the base level of MCP-1 both in serum and CSF indicated by the data from

con-group (Fig.6B and Fig.7A), the CSF MCP-1 level was about twice of its serum

level in nasal-group and i.c./nasal-group. This suggests that the MCP-1 in the CSF

may not derive from circulating blood, but rather produced by local cells. Activated

microglia was verified to serve as a possible producer of MCP-1 (*29*). Neonatal BCG

immunization has been shown to activate microglia (*25, 30*). This deduction provides

with an interest cue for future research to investigate the exactly mechanism under the

role of BCG in increasing the CSF level of MCP-1.

According to the previous report, the frontiers of the growth point of dural

lymphatics reach the scalp of skull such as the lateral parts of TS area at about two

weeks old (5). Given that the nasal BCG vaccination lasted for two weeks or more,

the dura mater of TS and SSS area, not skull base, was chosen for dural lymphatics

analysis. Moreover, the complex anatomical structure of the base of skull and the tight

attaching of dura mater there would make it difficult to analyze the lymphatics

precisely. It is similar with the part of dura mater overlapping attached to the frontal

bone overlying the olfactory bulb. In addition, the part of dura mater attached to the

scalp of skull is easy to be isolated from bone.

The dural lymphatics begin to cover the whole length area along the SSS but are

often discontinuous even in adult mice as reported previously (5). These vessels in

mice at P21 were seen in about 1/4 to 1/3 of the whole SSS length in con-group mice

(Fig. 2A). At P21, mere nasal BCG vaccination mildly increased the dural lymphatics

whereas mere neonatal i.c. BCG injection brought no significant effects. Both the

nasal-group and i.c./nasal-group showed an advanced development of the dural

lymphatics at P21, with their dural lymphatics distribution reaching the level

comparable with those in normal adult mice. This advanced development during the

critical period for CNS development is very likely to allow an advanced and more

effective drainage of CSF and thereby to produce an affect on brain development.

In natural condition, the microbiota to which human beings nasally expose are

certainly different from nasal BCG vaccination used in this study both in kind and in

dosage. Admittedly, it is not proper to estimate the status of dural lymphatics

development in human beings straightly according to the findings in this study

especially given that it is not clear whether the dural lymphatics development in

perinatal period differs in human beings from that in mice. What is vital and novel in

this study is that nasal BCG vaccination does promote dural lymphatics development.

Even if this promotion means only a matter of an advanced development or an earlier

coming of matured dural lymphatics, rather than a long-lasting elevation of

lymphatics structure and function, it is still important for brain development since it

occurs during the the early life period when is critical for CNS development

programming. In other words, even a transient promoted/advanced dural lymphatics

development is likely to exert a long-lasting effects on brain development and

function. In addition, the protocol used in this work to immunize mice may serve as

an animal model for experiments studying dural lymphatics-related issues.

Authors' contributions

Study design: Junhua Yang and Zhibin Yao. Performing the experiments: Junhua

Yang (primarily morphological observations and ELISA), Lifang Yuan (primarily

animal treatments and grouping, morphological observations and ELISA), Linyang

Song (primarily multiplex assays and western blotting). Data analyses (blinded to

groups): Junhua Yang. Writing the paper: Junhua Yang. Funding acquisition: Junhua

Yang, Zejie Zuo and Zhibin Yao. Fangfang Qi and Jie Xu participated in discussions,

assisted with data analysis, technical instructions and manuscript editing.

Conceptualization, Supervision All authors read, commented and approved the final

manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

This work was supported by grants from the starting fund for high-level talent

introduction into Guangdong Pharmaceutical University (No. 51355093) to Junhua

Yang, the Special Foundation of Education Department of Guangdong Province, the

Medical Scientific Research Foundation of Guangdong Province, China (2013-159) to

Zhibin Yao and the National Natural Science Foundation of China (Nos. 81801058) to

Zejie Zuo. The funding agencies had no role in study design and performance; in the

collection, analysis and interpretation of data; in the writing of the report; or in the

decision to submit the paper for publication. We thank Technician Yuanjun Guan

(Zhongshan School of Medicine, SYSU) for her kind assistance with experimental

techniques, particularly the confocal imaging and analyses.

References:

1.   A. Aspelund *et al.*, A dural lymphatic vascular system that drains brain interstitial fluid and
macromolecules. *J Exp Med* **212**, 991-999 (2015).

2.   A. Louveau *et al.*, Structural and functional features of central nervous system lymphatic
vessels. *Nature* **523**, 337-341 (2015).

3.   R. Tamura, K. Yoshida, M. Toda, Current understanding of lymphatic vessels in the central
nervous system. *Neurosurg Rev* **43**, 1055-1064 (2020).

4.   S. T. Proulx, Cerebrospinal fluid outflow: a review of the historical and contemporary
evidence for arachnoid villi, perineural routes, and dural lymphatics. *Cell Mol Life Sci* **78**,
2429-2457 (2021).

5.   S. Antila *et al.*, Development and plasticity of meningeal lymphatic vessels. *J Exp Med* **214**,
3645-3667 (2017).

6.   P. Baluk *et al.*, Transgenic overexpression of interleukin-1beta induces persistent
lymphangiogenesis but not angiogenesis in mouse airways. *Am J Pathol* **182**, 1434-1447
(2013).

7.   T. Okazaki *et al.*, alpha5beta1 Integrin blockade inhibits lymphangiogenesis in airway
inflammation. *Am J Pathol* **174**, 2378-2387 (2009).

8.   S. Fiorentini *et al.*, Human cytomegalovirus productively infects lymphatic endothelial cells
and induces a secretome that promotes angiogenesis and lymphangiogenesis through
interleukin-6 and granulocyte-macrophage colony-stimulating factor. *J Gen Virol* **92**, 650-660
(2011).

9.   A. N. Chard, M. Gacic-Dobo, M. S. Diallo, S. V. Sodha, A. S. Wallace, Routine Vaccination
Coverage - Worldwide, 2019. *MMWR Morb Mortal Wkly Rep* **69**, 1706-1710 (2000).

10. H. A. Fletcher, H. D. Donoghue, J. Holton, I. Pap, M. Spigelman, Widespread occurrence of
Mycobacterium tuberculosis DNA from 18th-19th century Hungarians. *Am J Phys Anthropol*
120, 144-152 (2003).

11. Z. Alzayer, Y. Al Nasser, in *StatPearls*. (Treasure Island (FL), 2021).

12. L. Liu, K. Duff, A technique for serial collection of cerebrospinal fluid from the cisterna magna
in mouse. *J Vis Exp*,   (2008).

13. P. Religa *et al.*, Presence of bone marrow-derived circulating progenitor endothelial cells in
the newly formed lymphatic vessels. *Blood* **106**, 4184-4190 (2005).

14. J. Y. Lee *et al.*, Podoplanin-expressing cells derived from bone marrow play a crucial role in
postnatal lymphatic neovascularization. *Circulation* **122**, 1413-1425 (2010).

15. M. Iolyeva *et al.*, Interleukin-7 is produced by afferent lymphatic vessels and supports
lymphatic drainage. *Blood* **122**, 2271-2281 (2013).

16. W. K. Wu, O. P. Llewellyn, D. O. Bates, L. B. Nicholson, A. D. Dick, IL-10 regulation of
macrophage VEGF production is dependent on macrophage polarisation and hypoxia.
Immunobiology **215**, 796-803 (2010).

17. H. Kinashi, Y. Ito, T. Sun, T. Katsuno, Y. Takei, Roles of the TGF-beta(-)VEGF-C Pathway in
Fibrosis-Related Lymphangiogenesis. *Int J Mol Sci* **19**,   (2018).

18. V. Gagliostro *et al.*, Pro-lymphangiogenic properties of IFN-gamma-activated human dendritic
cells. *Immunol Lett* **173**, 26-35 (2016).

19. X. Chen *et al.*, Increased IL-17-producing cells correlate with poor survival and

lymphangiogenesis in NSCLC patients. *Lung Cancer* **69**, 348-354 (2010).

20.   K. M. Bryant-Hudson, H. R. Gurung, M. Zheng, D. J. Carr, Tumor necrosis factor alpha and
interleukin-6 facilitate corneal lymphangiogenesis in response to herpes simplex virus 1
infection. *J Virol* **88**, 14451-14457 (2014).

21.   T. Sasahira *et al.*, Prox1 and FOXC2 act as regulators of lymphangiogenesis and angiogenesis
in oral squamous cell carcinoma. *PLoS One* **9**, e92534 (2014).

22.   M. J. Karkkainen, T. V. Petrova, Vascular endothelial growth factor receptors in the regulation
of angiogenesis and lymphangiogenesis. *Oncogene* **19**, 5598-5605 (2000).

23.   R. C. Ji, Macrophages are important mediators of either tumor- or inflammation-induced
lymphangiogenesis. *Cell Mol Life Sci* **69**, 897-914 (2012).

24.   K. R. Higgins, W. Kovacevic, L. Stokes, Nucleotides regulate secretion of the inflammatory
chemokine CCL2 from human macrophages and monocytes. *Mediators Inflamm* **2014**,
293925 (2014).

25.   J. Yang *et al.*, Neonatal BCG vaccination of mice improves neurogenesis and behavior in early
life. *Brain Res Bull* **120**, 25-33 (2016).

26.   A. N. Civello *et al.*, The effect of BCG vaccination on macrophage phenotype in a mouse
model of intranasal Mycobacterium bovis challenge. *Vaccine* **38**, 4755-4761 (2020).

27.   J. Chao, J. G. Wood, N. C. Gonzalez, Alveolar macrophages initiate the systemic microvascular
inflammatory response to alveolar hypoxia. *Respir Physiol Neurobiol* **178**, 439-448 (2011).

28.   M. Reale *et al.*, Production of MCP-1 and RANTES in bladder cancer patients after bacillus
Calmette-Guerin immunotherapy. *Cancer Immunol Immunother* **51**, 91-98 (2002).

29.   Y. Inose, Y. Kato, K. Kitagawa, S. Uchiyama, N. Shibata, Activated microglia in ischemic stroke
penumbra upregulate MCP-1 and CCR2 expression in response to lysophosphatidylcholine
derived from adjacent neurons and astrocytes. *Neuropathology* **35**, 209-223 (2015).

30.   M. J. Haley, D. Brough, J. Quintin, S. M. Allan, Microglial Priming as Trained Immunity in the
Brain. *Neuroscience* **405**, 47-54 (2019).

 Figure legends:

 Fig.1 Accelerated dural lymphatics development in nasal-group and

 i.c./nasal-group mice at P14. (A-D) Representative graphs show the differences in

 dLVs within the corresponding TS areas of the dura in each group of mice. Scale bar,

 200 μm. (E-H) Bars represent the length of TS covered by Lyve-1$^+$ vessels, Lyve-1$^+$

 vessels area%, numbers of sprouting from Lyve-1$^+$ vessels and numbers of cross

 points of Lyve-1$^+$ vessels dLVs within the unilateral TS area. $n = 6$ mice/group. *: $p <$

 0.05; **: $p < 0.01$; ***: $p < 0.001$. Randomly completely block design ANOVA

 followed by Bonferroni's *post hoc* test. Data are presented as the mean ± SEM.

 Fig.2 Accelerated dural lymphatics development with more M-LECPs in SSS

 zone in nasal-group and i.c./nasal-group at P21. (A-D) Representative graphs show

 the differences in dLVs distribution and macrophages number within the

 corresponding SSS areas of the dura in each group of mice. Scale bar, 400 μm. (E-H)

 Bars represent Lyve-1$^+$ vessels area, numbers of sprouting from Lyve-1$^+$ vessels,

 Prox-1$^+$ nucleus numbers and the number of total F4/80$^+$ cells. $n = 6$ mice/group. *: p

 < 0.05; **: $p < 0.01$; ***: $p < 0.001$. Randomly completely block design ANOVA

 followed by Bonferroni's *post hoc* test. Data are presented as the mean ± SEM.

 Fig.3 Selected representative scopes from graphs in Fig.2A-D, showing dural

 lymphatics and M-LECPs presence in SSS zone in nasal-group and

 i.c./nasal-group at P21. (A-D) Selected representative scopes that were next to the

 sinus confluence (the left end) of each SSS graph in Fig.2A-D. Scale bar, 100 μm.

(E-H) Selected representative scopes that were located at the middle of each SSS

graph in Fig.2A-D. Scale bar, 100 μm. (I-L) Representative high-magnified scopes

showing co-labelled Lyve-1$^+$(white), Prox-1$^+$(green) and F4/80$^+$ (red) signals. (M-N)

Bars represent the numbers of F4/80$^+$/Prox-1$^+$ cells and F4/80$^+$/Prox-1$^+$/Lyve-1$^+$ cells

within the whole SSS area of each group. (O) The ratio of F4/80$^+$/Prox-1$^+$/Lyve-1$^+$

cells numbers to F4/80$^+$/Prox-1$^+$ cells numbers within the whole SSS area of each

group. n = 6 mice/group. *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$. Randomly

completely block design ANOVA followed by Bonferroni's *post hoc* test. Data are

presented as the mean ± SEM.

Fig.4 Selected representative scopes showing the various stages of M-LECPs in

their transformation into dural lymphatics in SSS zone. (A-B) The same

representative scope showing co-labelled Lyve-1$^+$(white), Prox-1$^+$(green) and F4/80$^+$

(red) signals. (C) Another representative scope showing co-labelled Lyve-1$^+$(white),

Prox-1$^+$(green) and F4/80$^+$ (red) signals. (D) Another representative scope showing

co-labelled Lyve-1$^+$(purple), Prox-1$^+$(green) and F4/80$^+$ (red) signals.

Fig.5 Selected representative scopes from graphs in Fig. S1A-D, showing the

elevated real time lymphangiogenesis level in SSS zone in nasal-group and

i.c./nasal-group. (A-D) Selected representative scopes that were next to the sinus

confluence of each SSS graph in Fig. S1A-D. Scale bar, 50 μm. (E-H) Selected

representative scopes that were located at the middle of each SSS graph in Fig.

S1A-D. Scale bar, 50 μm. (I-L) Representative high-magnified scopes showing

co-labelled Lyve-1$^+$ (white), Prox-1$^+$(green) and BrdU$^+$ (red) signals. (M-N) Bars

represent the numbers of BrdU$^+$/Prox-1$^+$ cells and BrdU$^+$/Prox-1$^+$/Lyve-1$^+$ cells within

the whole SSS area of each group. $n = 6$ mice/group. (O) The ratio of

BrdU$^+$/Prox-1$^+$/Lyve-1$^+$ cells numbers to BrdU$^+$/Prox-1$^+$ cells numbers within the

whole SSS area of each group. *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$. Randomly

completely block design ANOVA followed by Bonferroni's *post hoc* test. Data are

presented as the mean $\pm$ SEM.

Fig.6 Elevated cytokine levels in CSF in nasal-group and i.c./nasal-group at P21

(A-B) Bars represent cytokines levels. $n = 6$ mice/group. **: $p < 0.01$; ***: $p < 0.001$.

Randomly completely block design ANOVA followed by Bonferroni's *post hoc* test.

(C-D) Representative bars represent cytokines levels. $n = 6$ mice/group. nd indicates

undetectable levels of one cytokine in certain group, such as con-group; ***: $p <$

0.001. Paired-samples t test between nasal-group and i.c./nasal-group. Data are

presented as the mean $\pm$ SEM.

Fig.7 Elevated cytokine levels in serum in nasal-group and i.c./nasal-group at

P21 (A-B) Bars represent cytokines levels. $n = 6$ mice/group. *: $p < 0.05$; **: $p < 0.01$;

***: $p < 0.001$. Randomly completely block design ANOVA followed by Bonferroni's

post hoc test. Data are presented as the mean $\pm$ SEM.

Fig.8 Increased VEGFR-3 protein level in dura mater of the mice in nasal-group

and i.c./nasal-group. (A) Representative western blots showing VEGFR-3 protein

level in dura mater of each group. (B) Quantitative data showing alteration in the

relative levels of the VEGFR-3 protein in the dura mater. $n = 6$ mice/group. *: $p <$

0.05; **: $p < 0.01$; ***: $p < 0.001$. Randomly completely block design ANOVA

followed by Bonferroni's *post hoc* test. Data are presented as the mean ± SEM. (C)

Representative graphs show VEGFR-3 expression in dLVs. Scale bar, 50 μm.

con   i.c. BCG   nasal BCG   i.c./nasal BCG   Prox-1/F4/80/Lyve-1

Prox-1/BrdU/Lyve-1   ● con ● i.c. BCG ● nasal BCG ● i.c./nasal BCG

A

con   i.c.BCG   nasal BCG   i.c./nasal BCG

VEGFR-3

β-actin

B

Relative VEGFR-3 protein level

***
***
*
*
*

C

VEGFR-3/Hoechst