# Sexual rejection modulates social interaction and reproductive physiology

Liora Omesi[1], Mali Levi[1], Elia Dayan[1], Yong-Kyu Kim[2a], Lital Barak-Buchris[1], Reza Azanchi[2b], Ulrike Heberlein[2] and Galit Shohat-Ophir[1,3]

[1]The Mina & Everard Goodman Faculty of Life Sciences, The Leslie and Susan Gonda Multidisciplinary Brain Research Center and The Nanotechnology Institute, Bar-Ilan University, Ramat Gan 5290002 Israel

[2]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA 20147, USA

[a]Current Address: Department of Neurobiology, Northwestern University, Evanston, IL 60208, USA

[b]Current Address: Department of Neuroscience, Brown University, Providence, USA

[3]Lead contact

Correspondence should be addressed to G.S.O. (galit.ophir@biu.ac.il)

**Abstract**

In highly polyandrous species, in which females mate with multiple males within a single fertility period, there is typically a high level of sperm competition. To cope with this challenge, males apply various behavioral and physiological strategies to maximize their reproductive success. Previous studies in *Drosophila melanogaster* established a link between the composition of the social environment and the reproductive success of individual male flies. While most studies until now focused on the adaptive responses of male flies to the presence of rival males, little is known about whether the outcomes of sexual interactions with female partners alter male-male social interactions. Here we show that repeated failures to mate promote coordinated physiological and behavioral responses that can serve to increase reproductive success over mating rivals in the future competition. We exposed male flies to sexual rejection, successful mating or no sexual experience, and analyzed the behavioral repertoires of individuals within groups and the structure of their emerging social networks. We discovered that failures to mate promote the formation of distinct emergent group interactions and structures, where rejected male flies form low density social networks and actively minimize their encounters with other group members, while increasing their aggressive behavior. In addition, sexually rejected male flies elevate the production of seminal fluid proteins and extend mating duration with receptive females, altogether leading to reduced re-mating rates. Our results demonstrate the existence of a flexible mating response as possible coping strategy for living in a highly dynamic and competitive environment as the social domain.

## Introduction

The ability to adapt to environmental changes is an essential feature of biological systems, achieved in multicellular organisms by a coordinated crosstalk between neuronal and hormonal programs that generate plastic physiological and behavioral responses to environmental challenges[1–3]. This is particularly important in a dynamic environment such as the social domain composed of many behaving animals, the interaction with ultimately determines the reproductive success of individuals[2,4–6]. The intricate nature of social interaction requires the ability to recognize other members of the group in the right context, season, sex, age and reproductive state, and integrate this information with prior experience to produce the appropriate and optimal behavioral response[5]. Plastic social responses are seen in diverse animals, and include modulation of competitive sexual behaviors such as mating preferences and aggressive displays, and also the regulation of social foraging and parental care[7–9]. A remarkable example of social plasticity is evident in the African cichlid fish *Astatotilapia burtoni*, which live in a highly complex social environment consisting of many rival males that compete over limited food, territorial resources and female partners. Such a complex biotic and social environment produces a small number of dominant male fish and a large number of submissive males that closely monitor the social landscape in a constant search for opportunities to improve their social status, taking over mating territories and females[9,10].

As a species with sociable lifestyle, *Drosophila melanogaster* exhibit communal living around freshly decaying fruits[11] and engage in diverse forms of social interactions[12]. This includes courtship and mating[13–16], fighting over resources[17], group interactions[18–20], coordinated responses to threats[18,21,22], cultural transmission of complex behaviors[23], learning from conspecifics[24,25], and synchronization of activity by social cues[26]. Although some of these behaviors are considered innate responses, there are striking examples of the ability of fruit flies to exhibit social plasticity as they modulate their behavior

and physiology in response to changes in their social environment. This includes the ability of male flies

to change their aggressive behavior in response to prior fighting experience[27–29], regulate sperm

composition and the duration of copulation events in response to perceived competition[30–32], and suppress

courtship efforts towards non-receptive female flies[33–35].

Recent studies in *Drosophila* demonstrate that fruit flies generate complex and rich group

structures that are sensitive to the density of the group, its composition, as well as to the prior experience

of its members[19,20,36,37]. We previously showed that sexual experience in male flies can modulate their

motivational state and, subsequently, their reward seeking behaviors[38,39]. However less is known about

the way by which prior sexual interactions that are experienced as success or failure to mate shape social

interaction of male flies in a group context. Furthermore, it is not clear whether sexually rejected male

flies exhibit loser-like responses, as in the case of social defeat[29], or rather actively increase their

competitive behavior to cope with mating rivals. Here we explored the effects of success or failure to

mate on the dynamics of social interaction in groups of male flies. We discovered that rejection promotes

the formation of discrete group interactions and structures, wherein rejected male flies actively minimize

their interactions with group members and at the same time enhance competitive behaviors and alter

reproductive physiology, presumably to increase their reproductive success.

## Results

**Failure to mate modifies action selection upon encounters with rival male flies**

We previously demonstrated that sexual experiences associated with different levels of mating success, such as repeated events of successful mating, or failures to mate in the form of repeated rejection events by non-receptive female flies, alter internal state and consequently motivational responses[38,39]. The negative valence of rejection, reflected by its capacity to induce courtship suppression and increase the consumption of ethanol, prompted us to explore the effects of sexual rejection on male-male social interaction and test whether failure to mate leads to loser-like state or rather facilitates competitive responses. To discriminate between these alternatives we analyzed the behavioral signature of male flies that experienced repeated failures to mate upon interaction with other male flies in a group context, compared to controls that experienced successful mating or lack of any sexual experience.

Wild type male flies (Canton S) were collected upon eclosion and kept in isolation for four days before being exposed to female flies. Rejected cohort (rejected-isolated) was generated by exposing 4 days old single male flies to 1h long interaction with a non-receptive female fly, 3 times a day for 4 days (Fig. 1A)[40]. As control, we used male flies that experienced successful mating, 1h long interactions with a receptive virgin female (1:1 ratio) , 3 times a day for 4 days (mated-isolated) and age matched male flies that lack any sexual experience (naïve isolated). At the end of this experience, their social group interactions was examined by introducing 10 flies from each cohort into the FlyBowl arenas, where they could move and interact in two dimensions. Their behavior was recorded for 30 min and analyzed using the FlyBowl suite of tracking and behavior analysis softwares[20,41,42] (Fig. 1A).

The tracking data obtained was used to calculate various kinetic features including velocities, distances and angles between flies and their relative differences across time (Fig. 1B). This set of per frame features served to train 8 types of behavior classifiers using JAABA (Fig. 1B)[20], and to analyze

their emergent social networks, by measuring the overall number or duration of interactions between pair

of flies. Interaction events were scored using the following parameters: >8mm between 2 flies and the

visual field of view of the focal fly is occupied by the other fly (angle subtended > 0), indicating that the

focal fly can see the other fly, and that these two requirements are maintained for at least 2 seconds. We

used these interaction criteria to calculate various properties of the emergent social networks of flies from

the rejected, mated, and naive cohorts.

Examining features that reflect activity levels such as time spent walking, turning, time spent

stopping and average speed, revealed similar activity levels in rejected and naive males, that are

significantly higher than those of mated males (Fig. 1 C-F). Analysis of specific social behaviors

indicated overall reduced degree of social interaction in rejected males than in naive males (Fig. 1 G-J).

This included low levels of approach towards other individuals (Fig. 1G), low levels of male-male chase

behavior (Fig. 1H), reduced formation of chains composed of multiple chasing males (Fig. 1I), and

reduced rates of close touch behavior (Fig. 1 G-J).

**Failure to mate promotes social avoidance**

Next, we investigated the network structures of groups composed of rejected, mated and naive

male flies using features that describe the properties of the network and the individuals within it (Fig.

2A)[20]. We calculated network weights according to the overall duration of interactions (emphasizing

long-lasting interactions) or the overall number of interactions (emphasizing short interactions) between

each pair of flies (Fig. 2B-I). Analysis by duration indicated that the social networks of rejected males

are characterized by lower density (Fig. 2B), and that the average strength of individual flies in these

networks is lower compared to mated and naive male flies (Fig. 2C). Likewise analysis of network

features by number of interactions identified reduced density and low strength of the individuals than

naive males (Fig. 2 F,G). These findings suggest that rejection promotes the formation of sparser groups

containing individuals with reduced social interaction.

The reduced network density of rejected male flies prompted us to compare the average distance

between individuals of each group as indicated by the average field of view occluded by another fly

(anglesub), a feature that increases when distance between individuals decreases (Fig. 2J). Analyzing the

average degree of anglesub along the experiment discovered that rejected male flies maintain

significantly low values of anglesub, suggestive of maintaining long distances between flies across the

experiment (Fig. 2J). This finding is also supported by reduced number of flies found near a focal fly

compared to mated and naive male flies (Fig. 2K). Given that rejected male flies exhibit similar activity

levels as naive male flies, and presumably higher probability to encounter other flies, their reduced

network density and high inter-individual distance suggest that rejected individuals actively avoid social

interactions with other flies, resulting in low density groups.

The behavioral signatures of the three cohorts across all 60 parameters are depicted in a scatter

plot of normalized differences and are divided into 4 main categories: activity, interaction, coordination

between individuals and social clustering where male aggregate to form clusters (Fig. 3). While the

cohort composed of mated male flies is characterized by overall reduced activity and the formation of

stable social clusters, the cohort composed of naïve male flies displayed increased activity, high levels

of social interaction and low levels of social clustering (Fig. 3). In contrast, the rejected cohort depicted

similar activity levels to that of the naïve cohort, reduced formation of social clusters and overall reduced

behaviors that are associated with social interaction (Fig. 3). The existing similarities and differences

between the three cohorts indicate that sexual rejection induce a discrete behavioral state that differs from

successful mating and the lack of mating and point to sexual rejection as the major contributor to reduced

social interaction.

**Sexual rejection modulates competitive behaviors**

The discrete group signature associated with sexual rejection led us to hypothesize that the behaviors exhibited by rejected males reflect behavioral response to coping with high sexual competition over mating partners, where repeated encounters with mated females are indicative of a high male to female sex ratio. To examine this hypothesis, we next tested whether rejection modulate competitive behaviors that can affect reproductive success such as aggressive responses toward other males and mating with virgin female flies. We compared rejected males to mated males, since both cohorts experienced interaction with females, unlike naïve flies that were socially isolated, a condition known by itself to promote aggression. We discovered that pairs of rejected male flies exhibited significantly higher aggressive displays in comparison to pairs of mated male flies (Fig. 4A), and that in mixed pairs, rejected males exhibited greater numbers of lunges compared to their mated opponents (Fig. 4B,C). To test whether rejection modulates investment in mating behavior, we compared the mating duration of rejected males to that of naïve male flies since mating investment of the mated cohort is shaped by their previous mating events. When allowed to mate with virgin female flies, rejected male flies extended the duration of copulation events by 25% (3.5 minutes longer) compared to naïve males (Fig. 4D). Thus, rejected male flies exhibited a significant increase in behaviors that may enhance their reproductive success in a competitive environment.

**Failure to mate induce changes in sperm and seminal fluid composition**

To determine whether prior rejection affects reproductive physiology such that it can promote reproductive successes, the expression levels of genes related to sperm production and reproduction were assessed. First, the expression of DON-JUAN (DJ), a protein that is specifically expressed in mature male sperm cells[32,33], was measured using a GFP-based reporter line in which a GFP sequence was inserted within the coding locus, so that the expression of GFP reflects the expression of the endogenous

DJ protein. The reliability of the DJ-GFP reporter as a sensitive measure for changes in sperm production

was first confirmed by comparing DJ-GFP levels in isolated and grouped housed male flies, since these

conditions are known to affect the amount of mature sperm[31,43](Supp Fig.1). The relative levels of GFP

were then measured in rejected and naïve male flies (Fig. 4E,F). We documented a two-fold decrease in

the levels of GFP in the rejected cohort compared to naïve males, suggesting that male flies decrease

their investment in sperm allocation in response to sexual rejection (Fig. 4E,F).

Next, the relative expression of the following reproductive related genes was directly assessed in

fly abdomens by qRT-PCR. We measured the expression of *Sex-Peptide (Acp70A), Acp63, Acp53,*

*Ovulin (Acp26Aa)*, which are responsible for the females' long-term post-mating responses and fertility[44].

We also measured the expression of genes encoding the *Ejaculatory bulb protein (Ebp),* which is

responsible for the posterior mating plug formation at the end of mating[45], *don-juan (dj)*[46], the *corazonin*

*(Crz)* neuropeptide, which promotes sperm and seminal fluid ejaculation in males and its receptor *Crz-*

*receptor*[39,47] (Fig. 4G). There was a two-fold increase in the levels of *Acp-70A (Sex-Peptide)* and *Acp-63*

in rejected male flies when compared to naïve males, suggesting that rejected male flies increase their

investment in the production of seminal fluid proteins that are transferred to females flies during

copulation (Fig. 4G). Nevertheless, in agreement with the observed reduction in DJ-GFP reporter levels,

there was a drastic decrease in the transcript levels of *don-Juan* in rejected males. The transcript levels

of *Ebp and Crz* and its receptor were similar in both cohorts (Fig. 4G). Overall, these results suggest that

rejected male flies respond to sexual rejection by elevating seminal fluid protein transcript levels,

presumably to maximize their fitness.

In addition to proteins associated with the male reproductive system, levels of several genes

expressed in the brain and antenna were also assessed. These included the neuropeptides *Crz*,

*Neuropeptide F* (*npf*) and its receptor (*npfr*), and two olfactory related genes associated with aggression

(the *Odorant binding protein 69a*[48], and *Cyp6a20*[49]. In agreement with previous studies, the levels of *npf*

were significantly lower in sexually rejected male flies[38]; we also observed a reduction in *npfr* (Fig. 4H).

Interestingly, sexually rejected male flies also exhibited reduced levels of *Cyp6a20* in comparison to

naïve male flies (Fig. 4H), consistent with their enhanced aggression (Fig. 3 A-C).

**Females that mate with rejected male flies exhibit reduced re-mating**

While the molecular changes associated with rejection support our initial hypothesis that rejected

male flies adjust their behavior and physiology to cope with high sexual competition, it is not clear

whether these and the extended copulation improve their reproductive success. To test this direction,

several aspects associated with female fecundity were measured. First, the fertility of female flies was

assessed after one mating event with either rejected or naïve male flies, by counting the number of eggs

laid (fecundity) and the corresponding number of adult progeny (productivity). There was no significant

difference in the number of eggs laid or emerging progeny between the two cohorts 2, 4 and 6 days after

initial mating (Fig. 5A). The lack of difference in the amount of progeny suggested that lower sperm

investment in rejected males (as reflected by reduced DJ levels) does not affect the total offspring

number, meaning that there is no link between sperm investment and the number of offspring.

Finally, we tested whether the increase in sex-peptide could facilitate enhanced post-mating

behavior (such as reduced receptivity) in females that mated with rejected male flies. Since the strongest

post-mating response is observed 24h after mating (data not shown), the proportion of female flies that

re-mated with fresh male flies 24h after they mated with either rejected or naïve male flies was measured.

A significant reduction was documented in the re-mating rates of females that mated initially with

rejected males (Fig. 5B), suggesting that extended copulation time and increase in seminal fluid proteins

can lead to a stronger reduction in female receptivity.

During copulation, male flies transfer to female flies seminal fluid proteins and also anti-

aphrodisiac pheromones such as cVA[34]. The extended copulation observed in rejected male flies may

facilitate the transfer of larger amounts of cVA as a means to delay further courtship and copulation with

subsequent male flies. As an indirect measure for possible changes in the amount of transferred cVA, we

analyzed the courtship behavior of naïve male flies towards females that previously mated with either

rejected or naive male flies 1h after the initial mating. No significant difference was observed in the

latency to court, i.e. the time it takes male flies to exhibit their first courtship action (wing vibration)

following introduction of the pair into the courtship arena (Fig. 5C). However, there was a significant

reduction in the number of naïve male flies that courted females previously mated with rejected males

(Fig. 5D), suggesting that mating experience with rejected male flies results in mated females that are

less attractive courtship targets than females previously mated with naïve males.

**Discussion**

In this study we used the FlyBowl as an agnostic tool to explore responses altered by sexual interaction, and discovered that rejected male flies cope with their failures to mate by changing their behavior and physiology, possibly to enhance their reproductive success. This is presumably achieved by avoiding interaction with potential rival male flies and competing over mating partners via increased aggression and prolonged copulation (known as mate guarding). The latter is strengthened by the increased production of certain seminal fluid proteins that facilitate stronger post-mating responses in female flies.

The behavior of sexually rejected male flies was examined in this study under behavioral contexts that illuminate different aspects of their action selection. Using the FlyBowl system, we analyzed their emergent group interactions and social networks, and discovered that although rejected males are highly active, they exhibit sparse networks and maintain large distance with other members, as if they were actively minimizing or avoiding interaction with rival male flies. When tested in a social context that promotes fighting over limited resources, rejected male flies exhibited enhanced aggression.

The increased aggression displayed by the rejected cohort is associated with a significant decrease in the levels of *Cyp6a20*. This is consistent with a previous study showing that *Cyp6a20* levels are reduced in social conditions that promote aggression and that this reduction is responsible for the observed increase in aggression[49]. Interestingly, exposure to female flies prior to male-male interactions was previously shown to suppress aggression[50]. However, our findings suggest that not all types of interactions with female flies are sufficient for suppressing aggression, but rather that the quality of the interactions (i.e., the male's mating success) determines aggression levels when encountering another male fly.

There are two possible explanations for the behavioral responses exhibited by rejected male flies.

First, failure to mate could enhance aggression to improve the chances of successful mating and, upon

eventual mating, the increased duration of copulation could increase the relative paternity share. Second,

repeated rejection experiences could be perceived by male flies as an indication for high density of sexual

competition over mating partners, where encountering mated females is suggestive of a high male to

female sex ratio. Consistent with the second hypothesis, several studies have described a link between

pre-exposure to rival male flies and an extension of copulation events[30,51]. One study also demonstrated

that male flies use multiple sensory cues such as auditory, olfactory and gustatory signals to estimate the

level of mating competition[51]. Although rejected males were not exposed directly to other male flies

during the training phase, the observed extension of their copulation events suggests that they can assess

the level of competition by evaluating the quality of their sexual interaction with female flies. Studies

performed in *Pieris rapae* butterflies, in which virgin males were shown to allocate their sperm

investment by assessing not only the mating status of the female but also her previous mating history[52],

are consistent with this hypothesis.

The behavioral responses to sexual rejection were accompanied by changes in the repertoire of

genes expressed in the brain and reproductive system in the form of increased expression of several

accessory gland protein genes (*Acps*). This, together with the increased copulation duration, supports the

idea that the observed extension in mating duration serves to transfer a higher amount of Acps to intensify

the females' post- mating responses[23,79]. Unlike previous studies that demonstrated a link between the

presence of rival male flies and an increase in both copulation duration and sperm allocation (measured

by increased number of sperm cells)[31], rejected male flies exhibited a significant reduction in the levels

of DJ expressed in mature sperm cells. Although this finding is limited to only one protein, this is

surprising in light of sperm competition theory, which predicts that males should strategically increase

their investment in sperm allocation when in competition[53]. This finding is also different from previous

studies that mating with non-preferred males produces smaller number of offspring and that non-

preferred males deliver larger amounts of sperm, presumably as a compensation mechanism[54–56].

Furthermore, our findings are different from studies in crickets, sunfish, birds rats and also flies, which

showed that the perceived risk of sperm competition, in the form of the presence of rival males or their

odors before and during mating, led to an increase in sperm investment[52,57,59,61,63]. The uncoupling

between the investment in sperm and non-sperm components and the regulation of investment in

copulation time demonstrates that sexually rejected male flies regulate each of these processes

independently.

Functionally, the observed decrease in sperm quantity with increasing seminal fluid protein *(Acp)*

expression in rejected males did not affect the amount of progeny produced in females. This observation

suggests that there is no link between the observed behavioral and physiological changes and the number

of progeny. Nevertheless, females that mated with rejected males were less attractive to naïve male flies,

as reflected by the reduced number of male flies that courted these females. The combination of reduced

female attractiveness in subsequent mating encounters, and reduced motivation of the female to re-mate,

may reduce the odds for a second mating and thus increase the rejected male's paternity rate despite the

lack of an effect on progeny number.

In summary, our results demonstrate a plastic mating strategy by males that experienced repeated

events of rejection which improves their reproductive success when competing with rival male flies. We

postulate that rejected males invest more energy in the production of seminal fluid proteins over sperm;

these Acps are known to have important roles in modulating different aspects of female mating

physiology and behavior. Furthermore, at low population density, the chances to meet a receptive female

are low, therefore an investment in sperm ejaculate may be more costly[58]. In this respect, Cameron et al.,

developed models that can explain the tradeoff between sperm and seminal fluid protein production and

their regulation under various conditions[60]. Their model predicts that competing males should evolve the

ability to adjust ejaculate composition based on competition, and whether they are the first or second

male to mate with a particular female[60]. Further research is needed to test ejaculate plasticity in the

context of sexual rejection, including whether mRNA levels of ACPs correlate with protein levels as

there are studies suggesting a weak correlation between the two[62], the proteome composition of ejaculate

that is transferred to females during copulation, and the ability of females to actively control this

process[64].

**Acknowledgments:** We thank all members of the Shohat-Ophir lab for fruitful discussions and technical

support. We specially thank Kristin Branson and Alice Robie (HHMI Janelia research campus) for their

valuable advice and guidance with the experimental and computational design. This work was supported

by the Israel Science Foundation grant 384/14 and Israel Science Foundation grant 174/19.

**Materials and methods:**

<u>Fly lines and culture</u>

Canton S flies were used as the wild-type strain. Flies were raised at 25°C in a 12-h light/12-h dark cycle in 60% relative humidity and maintained on cornmeal, yeast, molasses, and agar medium, and were tested as 3–4-day old adults, unless otherwise specified. The DJ-GFP and White Berlin (WB) lines were obtained from the HHMI Janelia Farm Research Campus.

<u>Sexual experience paradigm</u>

Male and female flies were anesthetized under $CO_2$ and isolated immediately after eclosion. Male flies were reared as single-housed in vials (23 mm by 94 mm) containing 7 ml of medium. Female flies were kept in groups of 10 flies/vial. Both males and females were aged for 3–4 day before the beginning of the experiment and kept in the incubator at 25°C, ~70% humidity, and light/dark of 12:12 hours. Mated females to be used as trainers, were generated by pair groups of 10 virgin female flies with 15 male flies 16 - 18 h before the training phase and were separated from male flies each morning, before the beginning of the training phase.

Experimental groups consisted of naive male, rejected and mated male flies: Rejected and mated cohorts were generated by pairing single male flies with either virgin female (mated cohort) or mated female (rejected cohort), for three one-hour sessions spaced by one hour rest, for consecutive days. During these one-hour sessions, pairs were monitored closely to make sure that the males interacting with virgin females mate successfully within the first hour, and that the males interacting with the previously-mated females do not mate. Males from the rejected cohort that managed to mate and males from the mated cohort that did not end up mating during all sessions were discarded. At the end of each session, the female fly was removed, and the males that experienced rejection were kept in the original vial for one hour rest. Naive male cohort was kept in the behavior chamber during the training phase, and the vials

containing single virgin males were handled as similar as possible to the rejected and mated cohorts,

without inserting female trainers. Detailed protocol is previously described[40].

<u>Social group interaction using the FlyBowl system</u>

At the end of the sexual experience phase, rejected and mated male flies were inserted in groups of 10

into Fly Bowl arenas[41], and their behavior was recorded for 30 minutes and analyzed using CTRAX,

FixTrax[20] and JAABA[41]. For kinetic features, scripts were written in MATLAB to use the JAABA code

to generate the statistical features as specified in Kabra et al. [41]. Time series graphs (per frame) were

created using JAABA Plot[41]. Quantification of complex behavios was done using JAABA Classifiers[41]

to identify specific behaviors: Walk, Stop, Turn, Approach, Touch, Chase, Chain, Song, Social

Clustering and Grooming. Bar graphs were created using JAABA Plot[41]. Network analysis was

performed using an interaction matrix according to the interaction parameters described previously[20].

Two interaction matrices were created for each movie, one with the total number of frames each pair of

flies were interacting divided by the number of frames in the movie and another with the number of

separate interactions between each pair of flies divided by the maximum number of possible interactions,

calculated as:

344
$$max \text{ \# of interaction possible } \frac{\text{\# of frames} - min\text{ \# of frames for interaction}}{min\text{ \# of frames for interaction} + min\text{ \# of gap frames}} + 1$$

The parameters to define an interaction are: angle subtended by the other fly > 0, distance between the

nose of current fly to any point on the other fly $\leq$ 8 mm, number of frames for interaction $\geq$ 60 and

number of gap frames $\geq$ 120. Interaction end is defined when distance or angle conditions are not

maintained for 4 seconds. Networks and their features were generated from the interaction matrix in R

using the igraph package. The function that was used to the generate networks is

"graph_from_adjacency_matrix" with parameters "mode = undirected" and "weighted = TRUE".

Density was calculated on all movies with the formula:

$$density = \frac{sum\ of\ weights}{[number\ of\ vertices * (number\ of\ vertices - 1)] * 0.5}$$

Modularity was calculated using the "modularity" function on output from the "cluster_walktrap"

function[65]. Strength was calculated using "strength" function and SD Strength was calculated on all

movies using "sd" function on the strength value. Betweenness Centrality was calculated on all flies

using the "betweenness" function and SD Betweenness Centrality was calculated on all movies using

"sd" function on the Betweenness Centrality value. Box plots were created using R.

Each feature of the FlyBwol experiment was standardized according to all values calculated in our

experiments for that feature to generate a z-score. Scatter plots were created using R.

<u>Aggression</u>

Pairs of rejected or mated male flies were introduced into aggression arenas (circular chambers, about

0.08 cm3 in volume), which contained a mixture of agarose and apple juice (1% agarose, 50% apple

juice) that was placed in arenas to enhance aggressive behavior. Flies were filmed for 30 min with Point-

Grey Flea3 (1080×720 pixels) at 60 fps. Aggressive behavior was later quantified by counting the number

of lunges for each pair using CADABRA software (ref). The $log_2$ ratio between the number of lunges in

rejected and mated flies was calculated for each pair, and then a one-sample t-test was performed to test

whether the mean ratio is significantly different from 0.

<u>Copulation duration</u>

Rejected and naïve male flies were put into courtship arenas (circular chambers, about 0.04 cm3 in

volume) with virgin females and were allowed to mate for 1 hour. They were recorded for the whole

experiment using a Point-Grey firefly camera. Courtship arenas consist of 25 flat arenas each arena

containing only one pair of male-female flies. The copulation duration was measured from the moment

the mating began until it ended. We calculated the time in seconds for each fly and the average for each

group.

Egg laying assay

Egg production was determined for females that had been allowed to copulate with rejected or naive

males for 1 hour at the end of the conditioning (as described above). Every female was put in a glass vial

containing fresh food every two days for 6 days in total and was kept in the incubator. The sum of the

number of eggs in the vials of each female was used for analysis.

Receptivity assay

3-4-day old White Berlin (WB) females were allowed to mate once with rejected or naïve males at the

end of the conditioning for 1 h. After mating, the males were disposed and the mated females were kept

in the incubator for 24h. Afterward, the mated females were exposed to 5-day-old WT naïve males for 1

h to measure their receptivity to mate. Approximately 40 pairs of each group (rejected or naive) were set

up in every biological repeat.

Latency to court assay

1 hour after allowing WB females to mate with rejected or naïve males, we transferred the females into

courtship arenas and paired them with new WT naïve males. The pairs were recorded for 15 min to

measure courtship latency. Latency was defined as the time elapsing between the introduction of the pair

into the chamber and the first appearance of wing vibration made by the courting male fly. We also

quantified the number of males who did and did not try to court in this assay.

Courtship Index

Courtship index for a given male is the fraction of time a male fly spent in courtship activity in the 10

min observation period (600 sec). It is calculated by dividing the number of seconds the male courted

over the total observation time and is been exhibit in percentage (CI = courtship behavior [sec] · 100 /

total observation [sec]).

<u>Molecular methods</u>

Western blot analysis: Sperm allocation in male flies carrying the DJ-GFP reporter was determined by

Western blotting. DJ protein size is ~29 kDa, and GFP size is ~25 kDa. We also determined the levels of

Sex-peptide (SP), a protein of size ~7 kDa, and the levels of Tubulin for normalization. The primary

antibodies used were mouse anti-GFP, rabbit anti-SP and rabbit anti-Tubulin, and the secondary

antibodies that were used are rabbit α-mouse HRP and mouse α-rabbit HRP, respectively. Virgin females

were used as negative controls.

<u>Quantitative Real-Time PCR analysis</u>

Frozen flies were placed on ice and decapitated using a scalpel. Total RNA was extracted from ~15

frozen heads and bodies (separately), using TRIZOL reagent according to the manufacturer's protocol.

mRNA was reverse transcribed using BIORAD cDNA synthesis kit. cDNA was analyzed by quantitative

real-time PCR (BIORAD CFX96) using specific primers for the head and for the body. Relative

expression was quantified by ΔΔCT method using RPL32[66] as a loading control. We run each sample in

triplicates. Each experiment was repeated four times using independent sets of experimental flies.

<u>Statistical analysis</u>

For each experiment, Shapiro–Wilk test was done on each experiment to test for normal distribution.

Statistical significance was determined by t-test for experiments that were distributed normally, and by

Wilcoxon test for experiments that were not distributed normally. For experiments with three conditions:

statistical significance determined by one-way ANOVA followed by Tukey's range test for experiments

that were distributed normally, and by Kruskal–Wallis test followed by Wilcoxon signed-rank test for

experiments that were not distributed normally. FDR correction for multiple testing was performed for

all analyses.

## References

1. Robinson, G.E., Fernald, R.D., and Clayton, D.F. (2008). Genes and Social Behavior. Science *322*, 896–900.

2. Oliveira, R.F. (2009). Social behavior in context: Hormonal modulation of behavioral plasticity and social
competence. Integr. Comp. Biol. *49*, 423–440.

3. Baker, M.R., Hofmann, H.A., and Wong, R.Y. (2017). Neurogenomics of Behavioural Plasticity in
Socioecological Contexts. In eLS, John Wiley & Sons, Ltd, ed. (Wiley), pp. 1–11.

4. Cardoso, S.D., Teles, M.C., and Oliveira, R.F. (2015). Neurogenomic mechanisms of social plasticity. J. Exp.
Biol. *218*, 140–149.

5. Taborsky, B., and Oliveira, R.F. (2012). Social competence: an evolutionary approach. Trends Ecol. Evol. *27*,
679–688.

6. Solomon-Lane, T.K., and Hofmann, H.A. (2019). Early-life social environment alters juvenile behavior and
neuroendocrine function in a highly social cichlid fish. Horm. Behav. *115*, 104552.

7. Jennions, M.D., and Petrie, M. (1997). Variation in Mate Choice and Mating Preferences: A Review of Causes
and Consequences. Biol. Rev. *72*, 283–327.

8. O'Connell, L.A., and Hofmann, H.A. (2012). Evolution of a Vertebrate Social Decision-Making Network.
Science *336*, 1154–1157.

9. Fernald, R.D. (2017). Cognitive skills and the evolution of social systems. J. Exp. Biol. *220*, 103–113.

10. Wallace, K.J., Choudhary, K.D., Kutty, L.A., Le, D.H., Lee, M.T., Wu, K., and Hofmann, H.A. (2022). Social
ascent changes cognition, behaviour and physiology in a highly social cichlid fish. Philos. Trans. R. Soc. B
Biol. Sci. *377*, 20200448.

11. Soto-Yéber, L., Soto-Ortiz, J., Godoy, P., and Godoy-Herrera, R. (2018). The behavior of adult Drosophila in
the wild. PloS One *13*, e0209917.

12. Sokolowski, M.B. (2010). Social interactions in "simple" model systems. Neuron *65*, 780–794.

13. Dickson, B.J. (2008). Wired for Sex: The Neurobiology of Drosophila Mating Decisions. Science *322*, 904–
909.

14. Anderson, D.J. (2016). Circuit modules linking internal states and social behaviour in flies and mice. Nat.
Rev. Neurosci. *17*, 692–704.

15. Cheriyamkunnel, S.J., Rose, S., Jacob, P.F., Blackburn, L.A., Glasgow, S., Moorse, J., Winstanley, M.,
Moynihan, P.J., Waddell, S., and Rezaval, C. (2021). A neuronal mechanism controlling the choice between
feeding and sexual behaviors in Drosophila. Curr. Biol. *31*, 4231-4245.e4.

16. von Philipsborn, A.C. (2020). Neuroscience: The Female Art of Saying No. Curr. Biol. *30*, R1080–R1083.

17. Wang, L., Han, X., Mehren, J., Hiroi, M., Billeter, J.-C., Miyamoto, T., Amrein, H., Levine, J.D., and
Anderson, D.J. (2011). Hierarchical chemosensory regulation of male-male social interactions in Drosophila.
Nat. Neurosci. *14*, 757–762.

18. Ramdya, P., Lichocki, P., Cruchet, S., Frisch, L., Tse, W., Floreano, D., and Benton, R. (2015).
Mechanosensory interactions drive collective behaviour in Drosophila. Nature *519*, 233–236.

19. Rooke, R., Rasool, A., Schneider, J., and Levine, J.D. (2020). Drosophila melanogaster behaviour changes in
different social environments based on group size and density. Commun. Biol. *3*, 304.

20. Bentzur, A., Ben-Shaanan, S., Benichou, J.I.C., Costi, E., Levi, M., Ilany, A., and Shohat-Ophir, G. (2020).
Early Life Experience Shapes Male Behavior and Social Networks in Drosophila. Curr. Biol. CB.

21. Elliott, K.H., Betini, G.S., and Norris, D.R. (2017). Fear creates an Allee effect: experimental evidence from
seasonal populations. Proc. Biol. Sci. *284*.

22. Ferreira, C.H., and Moita, M.A. (2020). Behavioral and neuronal underpinnings of safety in numbers in fruit
flies. Nat. Commun. *11*, 4182.

23. Danchin, E., Nöbel, S., Pocheville, A., Dagaeff, A.-C., Demay, L., Alphand, M., Ranty-Roby, S., van Renssen,
L., Monier, M., Gazagne, E., et al. (2018). Cultural flies: Conformist social learning in fruitflies predicts long-
lasting mate-choice traditions. Science *362*, 1025–1030.

24. Kacsoh, B.Z., Bozler, J., Ramaswami, M., and Bosco, G. (2015). Social communication of predator-induced
changes in Drosophila behavior and germ line physiology. eLife *4*.

25. Mery, F., Varela, S.A.M., Danchin, E., Blanchet, S., Parejo, D., Coolen, I., and Wagner, R.H. (2009). Public
versus personal information for mate copying in an invertebrate. Curr. Biol. CB *19*, 730–734.

26. Levine, J.D., Funes, P., Dowse, H.B., and Hall, J.C. (2002). Resetting the Circadian Clock by Social
Experience in Drosophila melanogaster. Science *298*, 2010–2012.

27. Yurkovic, A., Wang, O., Basu, A.C., and Kravitz, E.A. (2006). Learning and memory associated with
aggression in Drosophila melanogaster. Proc. Natl. Acad. Sci. U. S. A. *103*, 17519–17524.

28. Liu, W., Liang, X., Gong, J., Yang, Z., Zhang, Y.-H., Zhang, J.-X., and Rao, Y. (2011). Social regulation of
aggression by pheromonal activation of Or65a olfactory neurons in Drosophila. Nat. Neurosci. *14*, 896–902.

29. Kim, Y.-K., Saver, M., Simon, J., Kent, C.F., Shao, L., Eddison, M., Agrawal, P., Texada, M., Truman, J.W.,
and Heberlein, U. (2018). Repetitive aggressive encounters generate a long-lasting internal state in Drosophila
melanogaster males. Proc. Natl. Acad. Sci. U. S. A. *115*, 1099–1104.

30. Kim, W.J., Jan, L.Y., and Jan, Y.N. (2013). A PDF/NPF neuropeptide signaling circuitry of male Drosophila
melanogaster controls rival-induced prolonged mating. Neuron *80*, 1190–1205.

31. Garbaczewska, M., Billeter, J.-C., and Levine, J.D. (2013). Drosophila melanogaster males increase the
number of sperm in their ejaculate when perceiving rival males. J. Insect Physiol. *59*, 306–310.

32. Hopkins, B.R., Sepil, I., Thézénas, M.-L., Craig, J.F., Miller, T., Charles, P.D., Fischer, R., Kessler, B.M.,
Bretman, A., Pizzari, T., et al. (2019). Divergent allocation of sperm and the seminal proteome along a
competition gradient in Drosophila melanogaster. Proc. Natl. Acad. Sci. U. S. A. *116*, 17925–17933.

33. Siegel, R.W., and Hall, J.C. (1979). Conditioned responses in courtship behavior of normal and mutant
Drosophila. Proc. Natl. Acad. Sci. U. S. A. *76*, 3430–3434.

34. Keleman, K., Vrontou, E., Krüttner, S., Yu, J.Y., Kurtovic-Kozaric, A., and Dickson, B.J. (2012). Dopamine
neurons modulate pheromone responses in Drosophila courtship learning. Nature *489*, 145–149.

35. Ejima, A., Smith, B.P.C., Lucas, C., van der Goes van Naters, W., Miller, C.J., Carlson, J.R., Levine, J.D.,
and Griffith, L.C. (2007). Generalization of courtship learning in Drosophila is mediated by cis-vaccenyl
acetate. Curr. Biol. CB *17*, 599–605.

36. Schneider, J., Dickinson, M.H., and Levine, J.D. (2012). Social structures depend on innate determinants and
chemosensory processing in Drosophila. Proc. Natl. Acad. Sci. U. S. A. *109 Suppl 2*, 17174–17179.

37. Battesti, M., Pasquaretta, C., Moreno, C., Teseo, S., Joly, D., Klensch, E., Petit, O., Sueur, C., and Mery, F.
(2015). Ecology of information: social transmission dynamics within groups of non-social insects. Proc. Biol.
Sci. *282*, 20142480.

38. Shohat-Ophir, G., Kaun, K.R., Azanchi, R., Mohammed, H., and Heberlein, U. (2012). Sexual deprivation
increases ethanol intake in Drosophila. Science *335*, 1351–1355.

39. Zer-Krispil, S., Zak, H., Shao, L., Ben-Shaanan, S., Tordjman, L., Bentzur, A., Shmueli, A., and Shohat-
Ophir, G. (2018). Ejaculation Induced by the Activation of Crz Neurons Is Rewarding to Drosophila Males.
Curr. Biol. CB *28*, 1445-1452.e3.

40. Zer, S., Ryvkin, J., Wilner, H.J., Zak, H., Shmueli, A., and Shohat-Ophir, G. (2016). A Simple Way to Measure
Alterations in Reward-seeking Behavior Using Drosophila melanogaster. JoVE J. Vis. Exp., e54910.

41. Kabra, M., Robie, A.A., Rivera-Alba, M., Branson, S., and Branson, K. (2012). JAABA: interactive machine
learning for automatic annotation of animal behavior. Nat. Methods *10*, 64–67.

42. Branson, K., Robie, A.A., Bender, J., Perona, P., and Dickinson, M.H. (2009). High-throughput ethomics in
large groups of Drosophila. Nat. Methods *6*, 451–457.

43. Moatt, J.P., Dytham, C., and Thom, M.D.F. (2014). Sperm production responds to perceived sperm
competition risk in male Drosophila melanogaster. Physiol. Behav. *131*, 111–114.

44. Sirot, L.K., LaFlamme, B.A., Sitnik, J.L., Rubinstein, C.D., Avila, F.W., Chow, C.Y., and Wolfner, M.F.
(2009). Molecular social interactions: Drosophila melanogaster seminal fluid proteins as a case study. Adv.
Genet. *68*, 23–56.

45. Avila, F.W., Cohen, A.B., Ameerudeen, F.S., Duneau, D., Suresh, S., Mattei, A.L., and Wolfner, M.F. (2015).
Retention of Ejaculate by Drosophila melanogaster Females Requires the Male-Derived Mating Plug Protein
PEBme. Genetics *200*, 1171–1179.

46. Santel, A., Winhauer, T., Blümer, N., and Renkawitz-Pohl, R. (1997). The Drosophila don juan (dj) gene
encodes a novel sperm specific protein component characterized by an unusual domain of a repetitive amino
acid motif. Mech. Dev. *64*, 19–30.

47. Tayler, T.D., Pacheco, D.A., Hergarden, A.C., Murthy, M., and Anderson, D.J. (2012). A neuropeptide circuit
that coordinates sperm transfer and copulation duration in Drosophila. Proc. Natl. Acad. Sci. U. S. A. *109*,
20697–20702.

48. Bentzur, A., Shmueli, A., Omesi, L., Ryvkin, J., Knapp, J.-M., Parnas, M., Davis, F.P., and Shohat-Ophir, G.
(2018). Odorant binding protein 69a connects social interaction to modulation of social responsiveness in
Drosophila. PLoS Genet. *14*, e1007328.

49. Wang, L., Dankert, H., Perona, P., and Anderson, D.J. (2008). A common genetic target for environmental
and heritable influences on aggressiveness in Drosophila. Proc. Natl. Acad. Sci. U. S. A. *105*, 5657–5663.

50. Yuan, Q., Song, Y., Yang, C.-H., Jan, L.Y., and Jan, Y.N. (2014). Female contact modulates male aggression
via a sexually dimorphic GABAergic circuit in Drosophila. Nat. Neurosci. *17*, 81–88.

51. Bretman, A., Westmancoat, J.D., Gage, M.J.G., and Chapman, T. (2011). Males use multiple, redundant cues
to detect mating rivals. Curr. Biol. CB *21*, 617–622.

52. Wedell, N., and Cook, P.A. (1999). Butterflies tailor their ejaculate in response to sperm competition risk and
intensity. Proc. R. Soc. Lond. B Biol. Sci. *266*, 1033–1039.

53. Parker, G.A., Ball, M.A., Stockley, P., and Gage, M.J. (1997). Sperm competition games: a prospective
analysis of risk assessment. Proc. R. Soc. B Biol. Sci. *264*, 1793–1802.

54. Gowaty, P.A., Anderson, W.W., Bluhm, C.K., Drickamer, L.C., Kim, Y.-K., and Moore, A.J. (2007). The
hypothesis of reproductive compensation and its assumptions about mate preferences and offspring viability.
Proc. Natl. Acad. Sci. U. S. A. *104*, 15023–15027.

55. Gowaty, P.A., Kim, Y.-K., Rawlings, J., and Anderson, W.W. (2010). Polyandry increases offspring viability
and mother productivity but does not decrease mother survival in Drosophila pseudoobscura. Proc. Natl. Acad.
Sci. U. S. A. *107*, 13771–13776.

56. Anderson, W.W., Kim, Y.-K., and Gowaty, P.A. (2007). Experimental constraints on mate preferences in
Drosophila pseudoobscura decrease offspring viability and fitness of mated pairs. Proc. Natl. Acad. Sci. *104*,
4484–4488.

57. Gage, M.J.G. (1991). Risk of sperm competition directly affects ejaculate size in the Mediterranean fruit fly.
Anim. Behav. *42*, 1036–1037.

58. Parker, G.A., and Pizzari, T. (2010). Sperm competition and ejaculate economics. Biol. Rev. *85*, 897–934.

59. Nicholls, E.H., Burke, T., and Birkhead, T.R. (2001). Ejaculate Allocation by Male Sand Martins, Riparia
riparia. Proc. Biol. Sci. *268*, 1265–1270.

60. Cameron, E., Day, T., and Rowe, L. (2007). Sperm Competition and the Evolution of Ejaculate Composition.
Am. Nat. *169*, E158–E172.

61. Sirot, L.K. (2019). Modulation of seminal fluid molecules by males and females. Curr. Opin. Insect Sci. *35*,
109–116.

62. Smith, D.T., Sirot, L.K., Wolfner, M.F., Hosken, D.J., and Wedell, N. (2012). The consequences of genetic
variation in sex peptide expression levels for egg laying and retention in females. Heredity *109*, 222–225.

63. De Nardo, A.N., Roy, J., Sbilordo, S.H., and Lüpold, S. (2021). Condition-dependent interaction between
mating success and competitive fertilization success in Drosophila melanogaster. Evol. Int. J. Org. Evol. *75*,
2014–2026.

64. Kerwin, P., Yuan, J., and von Philipsborn, A.C. (2020). Female copulation song is modulated by seminal fluid.
Nat. Commun. *11*, 1430.

65. Pons, P., and Latapy, M. (2005). Computing communities in large networks using random walks. In Lecture
Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in
Bioinformatics) (Springer, Berlin, Heidelberg), pp. 284–293.

66. Zheng, Y.T., Li, H.B., Lu, M.X., and Du, Y.Z. (2014). Evaluation and validation of reference genes for qRT-
PCR normalization in *Frankliniella occidentalis* (Thysanoptera: Thripidae). PLoS ONE *9*.

Figure 1. Failure to mate modifies action selection upon encounters with rival male flies. A. Schematic representation of sexual experience and behavioral analysis. Rejected males experienced 1h long interaction with non-receptive female flies, 3 times a day for 4 days. Mated males experienced 1h long interactions with a receptive virgin female, 3 times a day for 4 days. Male flies with no sexual experience are kept in social isolation (naïve). The social group interaction of the 3 cohorts is recorded for 30 min using the FlyBowl system. B. List of kinetic features, 8 behaviors and network parameters produced using the FlyBowl system. C-J Average percentage of time mated (blue) naive (green) and rejected (red) took to perform walk (C), turn (D), stop (E), change orientation (F), approach (G), Chase (H), chain (I) and touch (J) behaviors. n = 15, 10, 15 for mated, naïve and rejected respectively. one-way ANOVA followed by Tukey's and FDR correction for multiple tests *p < 0.05, **p < 0.01, ***p < 0.001. Error bars signify SEM.

**Figure 2. Sexual rejection promotes social avoidance. A.** Illustration of network parameters. Density of networks represents how saturated they are compared to the maximum possible. Modularity is a measure of the division of a network into sub-networks. Strength is proportional to vertex size (high in red individual). Standard deviation (SD) strength is a measure of the heterogeneity of the connections between individuals. **B-I.** Social network analysis of groups composed of mated (blue) naive (green) and rejected (red). Network density, modularity, and SD strength calculated by network weights according to duration (B-E) or number of interactions (F-I). Kruskal–Wallis test followed by Wilcoxon signed-rank test and FDR correction for multiple tests *p < 0.05, **p < 0.01, ***p < 0.001, Box plot with median IQR. **J.** Rejected male flies maintain large distances between flies along (anglesub), statistical significance was tested on averages across time. one-way ANOVA followed by Tukey's and FDR correction for multiple tests *p < 0.05, **p < 0.01, ***p < 0.001. Error bars signify SEM. . **K.** Average number of flies close to a focal fly. n = 15, 10, 15 for mated, naïve and rejected respectively. one-way ANOVA followed by Tukey's and FDR correction for multiple tests *p < 0.05, **p < 0.01, ***p < 0.001. Error bars signify SEM.

**Figure 3.** Behavioral signatures of mated, rejected and naïve WT male flies. Data is represented as normalized Z scores of 60 behavioral parameters, n = 15, 10, 15 for mated, naïve and rejected respectively. statistical significance determined by one-way ANOVA followed by Tukey's range test for experiments that were distributed normally, and by Kruskal–Wallis test followed by Wilcoxon signed-rank test for experiments that were not distributed normally. FDR correction was used for multiple tests. LOI: calculated according to the length of interactions. NOI: calculated according to the number of interactions. Features marked in yellow exhibit statistically significant differences among 3 cohorts. Features colored in blue are different between mated and the two other conditions. Features colored in red are different between rejected and the two other conditions. Features colored in green are different between naïve and the two other conditions.

Figure 4. Sexual rejection modulates competitive behaviors and ejaculate composition. A.-C. Aggression display (number of lunges) was compared between pairs of rejected and mated male flies (n=16, statistical significance determined by T-test, $p< 0.005$ (A), and mixed pairs (n=12) (B-C). The $\log_2$ ratio between the number of lunges in rejected and mated flies was calculated for each pair, and then a one-sample T-test was performed to test whether the mean ratio was significantly different than 0, $p<0.005$. Data is presented as the mean ± SEM. D. Duration of copulation in rejected vs. naïve male flies. Statistical significance was determined by T-test, $p < 0.001$. Data is presented as mean ± SEM, n=25. E-F. Protein lysates prepared from abdomen of rejected and naïve male flies and were analyzed for the relative levels of Don-Juan-GFP using western blot, actin was used as a loading control. Expression levels of Don-Juan-GFP protein were quantified and normalized to actin levels (n=3), Statistical significance was determined by T-test, $p<0.05$ (F). G-H. Relative transcript levels of candidate genes expressed in abdomen (G) and heads (H) of rejected and naïve male flies were quantified by qRT-PCR, n = 6 independent experiments of 15–20 fly heads and abdomen. Statistical significance was determined by Student's T-test with Bonferroni correction for multiple comparisons. **, $p <0.01$ ***, $p < 0.005$.

**Figure 5. Effect of male rejection on female's fertility and remating tendencies**. **A.** Number of eggs laid by females that copulated with rejected or naïve male flies over the course of 6 days (left panel) and emerging progeny (right panel). Statistical significance was determined using two-way ANOVA repeated measure, n=30 naive, n=25 rejected $p>0.05$. **B.** Female receptivity to re-mate with male flies 24h after the first mating with rejected or naïve male flies was scored bycounting the precent of female flies that mated during 1 hour of test. Data is presented as the mean ± SEM, n=4 repeats. Statistical significance was determined by Cochran-Mantel-Haenszel Chi-square test, $p<0.005$. **C.** Mean courtship latencies of rejected or naïve male flies towards mated female flies (24 hours post first mating), n=25. Statistical significance was determined by Mann-Whitney U-test, N.S., $p>0.05$. **D.** Number of new males that courted females that were previously mated with rejected or naïve male flies, n=25. Statistical significance was determined by T-test, $p<0.05$.

629
630 **Supplementary Figure 1. The expression of Don-Juan protein in sensitive to the presence of rival male flies.**
631 Relative expression levels of Don-Juan-GFP in male flies in single or grouped housed male flies.