A re-analysis of the data in Sharkey et al.'s (2021) minimalist revision reveals that BINs do not deserve names, but BOLD Systems needs a stronger commitment to open science

Rudolf Meier[1,2], Bonnie B. Blaimer[2], Eliana Buenaventura[2], Emily Hartop[2], Thomas von Rintelen[2], Amrita Srivathsan[1], Darren Yeo[1]

[1] Department of Biological Sciences, National University of Singapore, Singapore

[2] Museum für Naturkunde, Leibniz Institute for Evolution and Biodiversity Science, Center for Integrative Biodiversity Discovery, Berlin, Germany

Abstract

Halting biodiversity decline is one of the most critical challenges for humanity, but

monitoring biodiversity is hampered by taxonomic impediments. One impediment is the

large number of undescribed species (here called "dark taxon impediment") while

another is caused by the large number of superficial species descriptions which can only

be resolved by consulting type specimens ("superficial description impediment").

Recently, Sharkey et al. (2021) proposed to address the dark taxon impediment for

Costa Rican braconid wasps by describing 403 species based on barcode clusters

("BINs") computed by BOLD Systems. More than 99% of the BINs (387 of 390) are

converted into species by assigning binominal names (e.g., BIN "BOLD:ACM9419"

becomes *Bracon federicomatarritai*) and adding a minimal diagnosis (usually consisting

only of a consensus barcode). We here show that many of Sharkey et al.'s species are

unstable when the underlying data are analyzed using different species delimitation

algorithms. Add the insufficiently informative diagnoses, and many of these species will

become the next "superficial description impediment" for braconid taxonomy because

they will have to be tested and redescribed after obtaining sufficient evidence for

confidently delimiting species. We furthermore show that Sharkey et al.'s approach of

using consensus barcodes as diagnoses is not functional because it cannot be

consistently applied. Lastly, we reiterate that COI alone is not suitable for delimiting and

describing species and voice concerns over Sharkey et al.'s uncritical use of BINs

because they are calculated by a proprietary algorithm (RESL) that uses a mixture of

public and private data. We urge authors, reviewers, and editors to maintain high

standards in taxonomy by only publishing new species that are rigorously delimited with

open-access tools and supported by publicly available evidence.

Running head: Reply to Sharkey et al.

Keywords: Integrative Taxonomy, DNA barcodes, BIN, BOLD Systems, Species

Delimitation, Taxonomic Impediment

After being principally an academic subject for many decades, biodiversity declines are

now also a major topic for decision makers (e.g., economists, CEOs, political leaders)

(World Economic Forum, 2020). This heightened interest has highlighted once more that

one of the most important tasks in science is unfinished: the discovery and description of

earth's biodiversity (May, 2011). Most of the species-level diversity and a significant

proportion of biomass are concentrated in taxonomically poorly known taxa that are

nowadays often referred to as "dark taxa" (e.g., bacteria, fungi, invertebrates)

(Hausmann et al., 2020). Such groups are usually avoided by taxonomists because they

combine high diversity with high abundance. Yet, dark taxa are a major cause for the

"taxonomic impediment" that is the result of a range of issues. One is the large number of

undescribed species (here called "dark taxon impediment") while another is species

descriptions that are too superficial by today's standards to allow for the identification of

the species without inspecting type specimens, collecting additional data, and

re-describing the species (here called "superficial description impediment"). Addressing

the superficial description impediment is particularly time-consuming because it requires

museum visits/loans, type digitization, and possibly lengthy searches for misplaced

specimens. Eventually, the species have to be re-described, making the total process of

getting to a useful species description take twice the amount of time needed if there had

not been a poor-quality legacy description. The reality is that new species descriptions

could be accelerated manifold if taxonomists did not have to spend so much time on

improving existing descriptions.

What the 21[st] century undoubtedly needs is more species descriptions, but what should

be avoided is creating new "superficial description impediments" in the form of large

numbers of poorly supported and described species. Yet, this is what Sharkey et al.

(2021) propose when they state: "… we view barcode-based descriptions as a first pass

in an iterative approach to solve the taxonomic impediment of megadiverse and

under-taxonomically resourced groups that standard technical and biopolitical

approaches have not been able to tackle." Sharkey et al. delegate the critical "iterative"

work to future generations of taxonomists. They will have to start their own revisions by

first revisiting the species descriptions, types, and specimens of Sharkey et al.'s (2021)

species to resolve the species boundaries based on data that should have been

collected and analyzed at the time of description.

Upon closer inspection, Sharkey et al.'s "minimalist revision" relies on three core

assumptions/techniques which are here shown to be problematic. The first is that COI

barcodes are sufficiently decisive and informative that they can be used as the only/main

data source for delimiting and describing species. The second is that species can be

diagnosed unambigously using only a consensus barcode. The third is that it is sufficient

to use a particular type of barcode cluster ("BIN") that is delimited by a proprietary

algorithm that is only installed on BOLD systems.

<u>(1) Are COI barcodes sufficiently informative for delimiting and describing species?</u>

Sharkey et al.'s revision covers 390 BINs obtained from BOLD. Of these, >99% (387) are

converted into species. This is surprising because virtually all studies that have looked

into the stability of barcode clusters have found that the cluster composition varies

depending on which barcode clustering algorithm is used (Virgilio et al., 2010; Kekkonen

and Hebert, 2014; Srivathsan et al., 2019; Yeo et al., 2020; Hartop et al., 2021); i.e.,

barcode data is not sufficiently decisive that they can be described as species – rather,

species delimitation must integrate multiple forms of data. We here first tested whether

the barcode data used in Sharkey et al. (2021) is unusually decisive and could be used

without validation from another source. Conventionally, such re-analyses of data from

published studies are straightforward because the data are made available by the

authors. However, the "molecular data" in the supplementary material of Sharkey et al.

(2021) consist only of neighbor-joining (NJ) trees. We thus proceeded to obtain the

sequences directly from BOLD by consulting the BIN numbers in the revision and NJ

trees in the supplement. Note, however, that BINs are calculated based on public and

private data with the ratio between the two apparently being approximately 3:1 (see

below). In addition, the NJ trees in the supplement contained non-Costa Rican BINs from

other Neotropical countries that were not covered by the revision. Consequently, we

ended up with three datasets of increasing size: (1) public barcodes for only those BINs

described by Sharkey et al. (2021), (2) public barcodes for all BINs found on the NJ trees

in the supplementary materials, and (3) all public barcodes in the BOLD database for the

11 braconid subfamilies covered by the revision of the Costa Rican fauna (see

supplementary data).

All three datasets were analyzed with several species delimitation methods as

recommended by Carstens et al. (2013). The overall analyses followed Yeo et al. (2020):

objective clustering with uncorrected p-distance thresholds at 1-5% using TaxonDNA

(Meier et al., 2006), ASAP – a new implementation of ABGD – where we only kept the

results obtained with the top five partitions (= lowest p-values) (Puillandre et al., 2012a;

Puillandre et al., 2021), and Single-rate PTP (Zhang et al., 2013) based on maximum

likelihood trees generated from the barcode data with RAxML v.8 (Stamatakis, 2014).

These re-analyses were used to test whether the different methods unambiguously

supported the BINs that were described as species by Sharkey et al. (2021). Arguably,

this is a minimum requirement for using BIN=Species as default because it is hard to

justify describing barcode clusters as species if the former are not consistent across

methods.

We find that in dataset 1, 131 of the 401 BINs (32.7%) conflict with at least one of the

species delimitation treatments. The instability of these BINs increases as more *COI*

evidence is included (Fig. 1), with 283 of the 615 BINs (46.0%) from dataset 2, and 2912

of the 3896 BINs (74.7%) from dataset 3 in conflict with the results of at least one of the

analyses. The corresponding numbers for only those 401 BINs described as species in

Sharkey et al. (2021) are 138 (34.4%: dataset with all Neotropical barcodes from BOLD)

and 276 (68.8% dataset with all braconid barcodes from BOLD). This means that as

braconid barcodes are sampled more densely across the Neotropics and the world, an

increasing number of the BINs described as species in Sharkey et al. (2021) contain

specimens that are assigned to barcode clusters that disagree with the BIN assignment

from BOLD. Overall, this reanalysis shows that COI barcodes contain ambiguous signal

when it comes to delimiting many species, and that this problem increases with

sampling.

Figure 1. Heatmaps showing incongruence between species delimitations for three

datasets of increasing size. In order to accommodate a large number of BINs, each

heatmap is split into 8 blocks that are separated by dark horizontal lines (number of bins

represented in each block is provided). Rows within each block represent species

delimitation algorithms (PTP, objective clustering: 1 – 5% p-distance, ASAP: top 5 priors)

and BINs in columns (blue = congruent BINs; pink = incongruent BIN).

To obtain lower-bound estimates for incongruence, we identified the parameters for each

species delimitation algorithm (objective clustering, ASAP) whose application yielded the

largest number of barcode clusters that are identical to BINs; i.e., the parameters from

each algorithm that maximized congruence with BINs. We find that 38 of the 401 BINs

(9.48%) described in the paper are incongruent even after the methods were optimized

to replicate BINs as well as possible (Fig. 2). Incongruence again increases as more of

the available barcodes and BINs are included [dataset 2: 82 of the 615 BINs incongruent

(13.33%), dataset 3: 1340 of the 3896 BINs (34.39%) incongruent]. These analyses

illustrate that the available *COI* data provide ambiguous support for 10-30% of the BINs

described as new species in Sharkey et al. (2021). This finding is congruent with virtually

all studies that have investigated the congruence between barcode clusters obtained

with different clustering methods (Virgilio et al., 2010; Kekkonen and Hebert, 2014;

Srivathsan et al., 2019; Yeo et al., 2020; Hartop et al., 2021). Furthermore, there are

good reasons to assume that 10-30% is an underestimate of the true number of clusters

that require other forms of data for resolving species boundaries because denser *COI*

sampling will close barcoding gaps between species, thus leading to further uncertainty

with regard to the boundaries of barcode clusters (Bergsten et al., 2012).

Sharkey et al. (2021) are aware that broader geographical sampling regularly leads to

BINs that contain multiple species: "Braconid specimens from the following New World

countries appear to be relatively well-sampled in BOLD: Canada, USA, Belize,

Argentina, French Guiana, and Mexico. There is a small number of cases where

specimens from these countries fall in the same BIN as one of our Costa Rican species,

but they were not studied. More sampling between these disparate localities, and more

genomic and/or morphological and behavioral data will help resolve these species-level

cases, which are beyond the scope of this paper." When they occasionally engage with

the readily available evidence, additional problems with BINs are immediately found. For

example, for BIN "BOLD:AAG8211" the authors state under "other material" that

"BIOUG07751-A11 (Malaise-trapped) from Honduras was examined and considered

conspecific, and it is included as a paratype. There are dozens of specimens in the same

BIN that are from widespread locations across Canada and the USA. Several of these

were examined and they differ significantly in color from the Mesoamerican specimens

and are not considered conspecific. See BOLD for details." In particular, the North

American specimens from Florida should have been consulted before delimiting the

Costa Rican species, but these specimens are nevertheless neglected.

194
Figure 2. Barcode cluster congruence across methods using optimal thresholds for BIN,

PTP, Objective Clustering (OC) (2%) and ASAP (d -3) for (a) dataset I, (b) dataset II, (c)

dataset III.

We suspect that Sharkey et al. (2021) overlooked the problems with delimiting species

with COI because they relied too much on BINs supplied by BOLD Systems. As in all

scientific studies, it is important that authors analyze the underlying data thoroughly in

order to present robust conclusions. For species descriptions involving molecular data,

the first step is to evaluate the decisiveness of the data by applying different species

delimitation algorithms (Virgilio et al., 2010; Carstens et al., 2013; Kekkonen and Hebert,

2014; Yeo et al., 2020). It also means avoiding publication of results obtained with

algorithms that are not publicly available (in this case: RESL) and operate on a mixture of

public and private data.

Evidence for the lack of decisiveness can also been seen when studying the composition

of BOLD BINs as more specimens are added to BOLD. This information is available

when one traces the "BIN history" of specimens. Here is the history for three BINs

described as species in Sharkey et al.'s (2021) minimalist revision (see suppl. materials

for details):

• BOLD:AAH8697 (*Heterogamus donstonei*) has 54 members of which 18 are public

and 14 are stable (have only been in one BIN). One unstable specimen

(BCLDQ0377) was originally placed in a fly BIN (BOLD:AAG1770: Diptera:

Muscidae: *Lispocephala varians*; 5 June 2010) before moving on to the current wasp

BIN (19 June 2010). The three remaining unstable barcodes were all originally placed

into the current BIN in September 2012, but subsequently shifted into two BINs

without public members (in May 2013). All three then shifted back into the current BIN

on 8 August 2015.

• BOLD:AAV3035 (*Pneumagathis erythrogastra*, new combination in Sharkey et al.

(2021)) has 14 members of which 12 are barcode compliant. Only two barcodes are

stable. The remaining 10 have similar histories. They were first placed in a stonefly

BIN (BOLD:AAC5216: Plecoptera: Chloroperlidae: *Alloperla severa*; March 2010)

before shifting to a BIN that is no longer available (January 2011). It is unclear when

these specimens shifted to the current BIN because this is not revealed by the delta

view tool in the BOLD Systems workbench. The two other barcodes were first placed

in BIN(s) that are no longer available. One (specimen H1170) then shifted to the

current BIN in September 2012 but later shifted to a BIN with a single private member

in May 2013. This specimen returned to the current BIN in June 2018. The other

specimen (specimen H7621) shifted from the original placement (BIN has ceased to

exist) to the current BIN in June 2018.

• BOLD:ABY5286 (*Chelonus scottmilleri*). The history for the 15 public barcodes is

detailed in Figure 3. It involves two BINs that are "no longer available"

(BOLD:AAA2380 and BOLD:AAK1017) and two others that could not be accessed as

they are "awaiting compliance with metadata requirements" (BOLD:AAD2009 and

BOLD:ABX7466). Note that the two currently valid BINs (BOLD:AAA7014 and

BOLD:ABX5499) contain species from different subfamilies of Braconidae

(Microgastrinae: *Apanteles anamartinezae* and Cheloninae: *Chelonus scottshawi*,

respectively).

All this highlights that relying heavily on BIN=Species for describing species is not

advisable. Indeed, depending on the time of BIN description, two of Sharkey et al.'s

(2021) wasp taxa would have been described as a fly or stonefly species, respectively.

Similarly, biologists using BOLD as an identification tool should be aware that using BINs

can lead to misleading conclusions. This makes it all the more important that BIN history

can be traced easily. However, this is time-consuming in BOLD Systems. One can only

reconstruct BIN evolution by tracing individual specimens – one at a time – and clicking

through pages of changes over time; yet, this will still only reveal information on publicly

available sequences. This is unfortunate because "taxon concepts" matter greatly in

taxonomy (Franz, 2005; Meier, 2016; Packer et al., 2018) and it would be particularly

straightforward to establish a "BIN-concept tracking" feature in BOLD Systems given that

it is a relatively new database and only deals with one type of data.

Figure 3. BIN membership through time for the 15 public barcodes in BOLD:ABY5286

(*Chelonus scottmilleri*). Two BINs are "no longer available"; two others cannot be

accessed as they are "awaiting compliance with metadata requirements"; the two

currently valid BINs contain species from different subfamilies (AAA7014 Microgastrinae:

Apanteles anamartinezae and ABX5499 Cheloninae: *Chelonus scottshawi*). They

contain 139 barcodes (all public) and 44 barcodes (although 45 listed as public),

respectively.

The instability of species estimates obtained with barcodes is also evident when

evaluating whether the original species diversity estimates in publications are consistent

with species richness estimates based on the number of BINs in BOLD. We initially

studied the data from publications that were highlighted in Sharkey et al. (2021). We then

added the data for additional publications as long as they allowed for a direct comparison

of species numbers (=authors provided a list of GenBank accession or BOLD sample

IDs). We obtained the sequences in February-April 2021 using these numbers by

querying GenBank, the Sequence-ID tool of GBIF, and BOLD Systems. Table 1

compares the number of species and BINs for those species that had molecular data.

We find that, for example, the barcodes for the 10 species in *Astraptes* belong to 5 BINs

(Hebert et al. 2004), the 32 species of *Belvosia* reported in Smith et al. (2006) belong to

20 BINs, and the barcodes in Smith et al. (2007) are in 61 BINs instead of the 73 species

reported in the paper. These data illustrate how ambiguous COI can be when it comes to

delimiting species. For the data from these studies, BINs are either not equal to species

or the species reported in the publications were not species.

Table 1: **Comparison of published species and BIN numbers (April 2021)**.

Study	Taxa	Journal	BINs in BOLD	Species or BINs published	Deviation
Hebert et al. 2004*	*Astraptes* (Lepidoptera: Hesperiidae)	PNAS	5	10 species	100%
Smith et al. 2006*	*Belvosia* (Diptera: Tachinidae)	PNAS	20	32 species	60.0%

Raupach et al. 2016	Carabidae (Coleoptera)	ZooKeys	51	69 BINs	35.3%
Chacón et al. 2013*	*Dunama* (Lepidoptera: Notodontidae, Nystaleinae)	ZooKeys	6	8 species	33.3%
Gibbs 2018	*Lasioglossum* (Hymenoptera: Halictidae)	Genome	85	110 species	29.4%
Fernandez-Triana et al. 2014*	*Apanteles* (Hymenoptera: Braconidae)	ZooKeys	240	180 species	25.0%
Smith et al. 2007*	various Tachinidae (Diptera)	PNAS	61	73 species	19.7%
Janzen et al. 2011*	Eudaminae and Pyrginae *(Hesperiidae) skippers*	PLoS One	294	347[§] species	18.0%
Hajibabaei et al. 2006*	Hesperiidae, Sphingidae, Saturniidae (Lepidoptera)	PNAS	600	521 species	13.2%
Fleming et al. 2019	*Hyphantrophaga* (Diptera, Tachinidae)	Biodiversity Data Journal	25	22 species	12.0%
Bartolini et al. 2020	*Anastrepha* (Diptera, Tephritidae)	PLoS One	17	15 species	11.8%
Pohjoismäki et al. 2016	Tachinidae (Diptera)	PLoS One	331	366 BINs	10.6%

Smith et al. 2008*	Braconidae (Hymenoptera)	PNAS	287	313 species	9.1%
Fleming et al. 2014*	*Houghia* (Diptera: Tachinidae)	Zootaxa	34	35 species	2.9%
Ortiz et al. 2017	Erebidae (Lepidoptera)	Biodiversity Data Journal	157	160 species	1.9%
Burns et al. 2008*	*Perichares* (Lepidoptera: Hesperiidae)	PNAS	5	5 species	0%

* Study cited in Sharkey *et al.*

§ number includes 44 putative barcode clusters.

In this context it is important to remember that there are good theoretical and empirical

reasons why one should not expect *COI* clusters to be equal to species (Meier et al.,

2006; Puillandre et al., 2012a; Puillandre et al., 2012b; Zhang et al., 2013). It has long

been known that introgression, lineage sorting, and maternal inheritance of mt DNA can

obfuscate the species-level signal in *COI* data (Will and Rubinoff, 2004; DeSalle et al.,

2005). In addition, we know that for closely related taxa the COI protein is under strong

stabilizing selection (Roe and Sperling, 2007; Kwong et al., 2012; Pentinsaari et al.,

2016), which explains why most of the evolutionary change at the DNA level is

synonymous and concentrated in 3[rd] positions. Nucleotide fixation in these positions is

likely caused by genetic drift; i.e., *COI* distances between BINs and closely related

species mostly measure time of divergence which is only correlated but not identical to

the probability of speciation.

This can be illustrated by the three datasets that we analyzed here. We find numerous

BINs whose barcodes translate into identical amino acid sequences (Table S1: dataset

1: 11 cases involving 26 BINs; dataset 2: 21 cases and 49 BINs; dataset 3: 28 cases and

57 BINs – see supplementary materials for details) and the proportion in other taxa can

be as high as 80-90% (Kwong et al., 2012). These BINs have no known, functionally

significant biological differences because the biology of these wasp "species" is not at all

affected by which triplet codons are used to code for identical proteins. Given that most

biologists associate speciation with the origin of biologically meaningful differences,

describing such BINs as species rests on the assumption that the correlation between

time of divergence and the origin of new species is sufficiently strong that biologically

meaningful differences will later be found. However, most biologists are rightfully

skeptical of results based on correlations. Correlations warrant more exploration; i.e.,

barcode clusters should only be treated as unnamed first-pass groupings until they are

validated with other data before being described as species.

An even more serious concern is the non-random nature of the error that is caused by

faulty species delimitation with COI barcodes. Species originating quickly via an adaptive

radiation will be systematically lumped while old species will be systematically split.

Evidence for these problems can be found throughout BOLD. Anyone who has worked

with the database will have found many cases where one BIN contains sequences that

were identified as coming from multiple species. Conversely, it is common to find species

that have barcodes in multiple BINs. This can be due to misidentification or genuine

lumping/splitting. Fortunately, there are taxa for which there is enough morphological,

genetic, and behavioral data that species boundaries are well understood. For these,

one can test whether the sequences of recently diverged species end up in one BIN and

the sequences for old species are found in several BINs. Such data are available, for

example, for many species of Sepsidae (Diptera) (Puniamoorthy et al., 2009;

Puniamoorthy et al., 2010; Tan et al., 2010; Ang et al., 2013a; Ang et al., 2013b; Araujo

et al., 2014; Rohner et al., 2014; Ang et al., 2017). Known pairs of closely related species

that are distinct with regard to morphology and behavior are routinely found lumped into

the same BIN (e.g., *Sepsis neocynipsea* and *S. cynipsea*: BOLD:AAC2855; *Sepsis*

orthocnemis and *S. fulgens*: BOLD:AAJ7599; *Themira lucida and T. flavicoxa*:

BOLD:AAD7140). *Sepsis punctum* is an example for the opposite. The populations in

North America and Europe can interbreed but are split into two BINs (BOLD:AAG5639;

BOLD:ACS2531). If taxonomists had applied BIN=Species as a default for delimiting

these species, several putatively young species would have been overlooked and some

old species would have been split. Some will argue that this degree of error is

acceptable, but we would counter that the non-random nature of the error is a major

concern because it systematically misinforms about speciation.

(2) Can species diagnoses consist of only a consensus barcode?

One of the key proposals of Sharkey et al. (2021) is that newly described species can be

quickly and efficiently diagnosed using only one *COI* consensus barcode. Indeed,

approximately 2/3rds of all newly described species in Sharkey et al. (2021) have such

diagnoses. This approach to diagnosis would work if BINs=Species and there were no

named species that were described based on morphological evidence. However, both

conditions are not met and the minimalist revision thus struggles with implementing the

"diagnosis=consensus barcode" concept. Indeed, one compromise follows another one

until it becomes clear that morphological data remain indispensable for many species

and highly desirable for others:

(1) Several BINs contain more than one species. The consensus barcodes for 11

species thus have to be amended by brief morphological diagnoses (e.g. *Bracon*

tihisiaboshartae can be differentiated from *B. alejandromasisi* by the color of the

metasomal terga: entirely yellow in *B. tihisiaboshartae* and partly black in *B.*

alejandromasisi"). The brevity of these amendments for all 11 species is reminiscent of

species descriptions from the 18th and 19th century, and it appears very unlikely that

these diagnoses will stand the test of time for species in a genus like *Bracon*, which "is

an enormous, polyphyletic, cosmopolitan genus with thousands of undescribed species

(Sharkey et al., 2021)." Such superficial diagnoses have the potential to become a major

"superficial description impediment" in future.

(2) Some species within the revision lack molecular data. Three new species are

described based on short morphological diagnoses: "Differing from the five other known

species (two undescribed and not included here) in many characters including a

hypopygium that is much more convex than the others. Body length 7.1 mm. Other

details can be gleaned from the images in Fig. 367". The description of these species

based on morphology means that all new, congeneric species delimited with DNA

barcodes also need morphological information in the diagnoses. Otherwise, it cannot be

shown that the new species are different. Sharkey et al. (2021) again amend the

consensus barcodes with very short morphological circumscriptions, which are unlikely

to be sufficiently informative once more Central American species are studied.

(3) Two new species are described based on *COI* sequences that are too short to be

barcodes according to the thresholds set by BOLD. Therefore neither has its own BIN

identifier. Sharkey et al. (2021) proceed to describe one species based on a consensus

mini-barcode which is again amended with morphological information ("Unique within

Rogadinae, this species possesses a strongly petiolate (clavate) hind femur. The

patchily patterned wings are also diagnostic within the *Colastomion-Cystomastax*

group."). For the other, the sequence information is ignored and only morphological

information is provided: "The COI barcode has only 293 bases and therefore was not

given a BIN. This species can be distinguished from all other species of *Zacremnops* by

red coloration restricted to the metapleuron (i.e., not on the propodeum) in combination

with the melanic hind tarsus."

(4) A special challenge to the "diagnosis=consensus barcode" concept are the 22 BINs

that contain non-Costa Rican specimens. These BINs and their consensus barcodes

were computed based on all data, but Sharkey et al.'s (2021) revision only covers the

Costa Rican species. This is problematic because the diagnoses based on consensus

barcodes are only correct when all specimens from outside of Costa Rica belong to the

Costa Rican species. This is unlikely for two reasons. Firstly, some of the non-Costa

Rican specimens are from geographically distant localities (e.g., Oklahoma, Kentucky,

Mexico, Columbia, Bolivia, Argentina). Secondly, Sharkey et al. (2021) occasionally

conclude based on images for non-Costa Rican specimens that some BINs contain >2

species. Sometimes, this leads to a diagnosis that consists of a consensus barcode and

a geographic amendment (e.g., *Phanerotoma angelsolisi* consists of those specimens in

BOLD:AAG8211 that are "Restricted to Central America"). In other cases, the

amendment is morphological (*Bracon mariamartachavarriae* consists of those

specimens of BOLD:AAV6295 that have a "Terminal flagellomeres black, concolorous

with basal flagellomeres"). However, the validity of the diagnoses is in question because

it is unclear whether critical specimens were studied (e.g., specimens from Florida for

BOLD:AAG8211; specimen from Argentina for BOLD:AAV6295). It is also unclear why

Sharkey et al. (2021) use consensus barcode for entire BINs when the authors only want

the species to consist of the Costa Rican specimens in a BIN. For BINs with wide

distributions, such diagnoses are very unlikely to pertain to the Costa Rican species.

Note that there are other cases where the diagnoses consisting only of a consensus

barcode are almost certainly incorrect because the authors indicate that the BINs appear

to contain more than one species (e.g., BOLD:AAH8705 and BOLD:ABZ7672).

There are additional oddities with regard to the use of consensus barcodes as

diagnoses, which highlight that relying only on one type of data is ill-advised.

BOLD:ABX6701 was described by Sharkey et al. (2021) as *Plesiocoelus vanachterbergi*.

The consensus barcode includes two single indels and was presumably obtained from

the seven "barcode-compliant" sequences in BOLD Systems (as of April 18, 2021). Only

one of the seven barcodes is translatable to amino acids with the remaining six having

deletions. Such sequences are typical for pseudogenes, and it is conceivable that *P.*

vanachterbergi was delimited and is now diagnosed based on paralogs. In addition, the

consensus barcode for *Pseudorhysipolis mailyngonzalezae* contains 104 indels, which

may be related to the fact that the BOLD fasta download for this BIN consists of data for

multiple genes (*COI, 28S, 16S, Ef1a*).

We here discuss these issues because diagnoses are critical elements of species

descriptions. They render species hypotheses testable (Ahrens et al., 2021). A diagnosis

consisting of only a consensus barcode cannot be tested with morphological data and

vice versa. This is one reason why integrative taxonomy is preferred over other

approaches. Sharkey et al.'s (2021) reliance on consensus barcodes is particularly

problematic because they are static summaries of dynamically evolving barcode

databases. More sequencing will almost certainly reveal synonymous nucleotide

changes in barcodes for the more than 200 species of braconids for which Sharkey et al.

(2021) only provide consensus barcodes as diagnoses; i.e., these new polymorphisms

will falsify the diagnoses. By omitting morphological information for these species,

Sharkey et al. also created a parallel taxonomy that does not allow for cross-referencing

the new species to the described fauna (Ahrens et al., 2021; Zamani et al., 2021). It

furthermore interferes with the work of those biologists, policymakers, and citizen

scientists in biodiverse countries who have no access to molecular data (Zamani et al.,

2021).

Our discussion of consensus barcodes as diagnoses also highlights Sharkey et al.'s

(2021) confusing relationship to morphological evidence. On the one hand, Sharkey et

al. (2021) argue that morphological evidence should not be used for braconid taxonomy.

On the other hand, a large number of diagnoses contain morphological information, and

in many cases the morphological characters are essential for the validity of the species.

Morphology was presumably also used to identify those BINs that consist of multiple

species. Despite this, no integrative process is described and the minimalist revision

failed to provide proper morphological diagnoses/description for many species.

(3) Should BINs obtained from BOLD Systems be used as species estimates?

There are three reasons why the BINs in the "minimalist revision" do not meet the

minimum standards for reproducible science. All three are related to open science issues

within BOLD Systems. Firstly, the algorithm for calculating BINs (RESL: Refined Single

Linkage) is not publicly available. Secondly, it is unclear how the underlying sequences

were aligned. Thirdly, BINs are calculated based on public and private barcodes:

RESL: Refined Single Linkage. The code for this algorithm that computes BINs is

proprietary. This makes it impossible to vary parameters such as the pairwise distances

used for the initial clustering, the thresholds used for merging neighbors, and the inflation

parameters for cluster refinement. This lack of user control can give the impression that

BIN boundaries are well defined, but this is achieved by not allowing users to vary

parameters; i.e., BINs are more akin to means that lack standard deviations, or a tree

with nodes that lack information on support. Barcode data should always be analyzed

using several species delimitation algorithms in order to investigate how robust different

barcode clusters are.

Overall, we know very little about RESL's performance because it was simultaneously

proposed and implemented based on only eight small datasets that comprised 18,843

barcodes (Ratnasingham and Hebert, 2013). The optimal clustering thresholds for these

datasets varied from 0.7% to 1.8% and eventually "2.2% was adopted as it represents

the upper 99% confidence limit for the optimal thresholds in the eight test datasets

(x☐=1.26), SD =0.40)" (Ratnasingham and Hebert, 2013). The same eight datasets were

also analyzed with four other species delimitation algorithms (ABGD, CROP, GMYC,

jMOTU). Overall, RESL and GMYC performed best with 89% congruence with

morphology. Note, however, that this level of congruence was only achieved after RESL

was applied to the same eight datasets for which it was optimized.

All remaining comparisons of RESL to other species delimitation algorithms are

compromised because RESL can only be used within BOLD Systems and the other

delimitation algorithms cannot be applied to BOLD System's private data. This means

that the comparative performance of RESL remains unclear. However, it is unlikely that it

is a superior method. As discussed by Ratnasingham and Hebert (2011), the algorithm

was developed with speed and scalability in mind and the authors anticipated that

"[f]uture research will undoubtedly reveal analytical approaches that are better at

recognizing species boundaries from sequence information than RESL." This means that

taxonomic revisions should not overly rely on BINs. Instead, the available data should be

analyzed using a range of species delimitation algorithms. As shown earlier, this would

have provided important insights into the robustness of BINs. Note that the use of

different analysis tools also allows users to gain control over the quality of the

alignments. This is important because this is another open science issue in Bold

Systems given that the "parameters for BOLD's alignment of sequences are not publicly

available" (Nugent et al., 2020) and cannot be changed.

Private data. The third reason why Sharkey et al. should not have relied on BINs is that

they are calculated based on public and private barcodes. The ratio of public and private

barcodes is difficult to determine because the relevant numbers vary between what is

listed on BOLD Systems under "Identification engine" and "Taxonomy", and what can be

downloaded from BOLD Systems via the "Public Data Portal". We here estimate that

approximately one quarter of all data are private. In order to obtain this estimate, we

downloaded the data for each taxonomic category listed under Animals in BOLD

Systems' "Taxonomy". Given the large size of the Arthropoda and Insecta datasets,

sub-taxa were downloaded separately. Downloading was done either directly from the

web browser or by using BOLD Systems' API (27 March-11 April 2021). Overall, we

obtained 6,165,928 COI barcodes >500 bp (6,508,300 overall) of the 8,480,276 (18 April

2021) barcodes that are listed as "All Barcode Records on BOLD". We estimate that

obtaining these data and setting up the facilities and bioinformatics pipelines cost >200

million C$, given that one project alone (Barcode500K: 2010-2015) had a budget of 125

million C$ (https://ibol.org/programs/barcode-500k/). It is very likely that some of these

funds were also used for generating private data. This includes private BINs that have

remained private for many years.

Use of potentially misleading tools. The value of Sharkey et al.'s (2021) minimalist

revision is not only diminished because it relies too heavily on BINs, but the authors were

also undiscerning in their choice of tools. Indeed, many are considered "deadly sins of

DNA barcoding" in Collins and Cruickshank (2013). This included the use of

Neighbor-Joining (NJ) trees which can yield biased results when the input order of taxa

is not randomized (Takezaki, 1998). BOLD Systems' K2P NJ trees feature prominently in

Sharkey et al. although they lack support values, information on how tie-trees are

broken, and use with K2P an inappropriate evolutionary model for *COI* evolution (Collins

et al., 2012; Srivathsan and Meier, 2012). Note that this model is not even used by RESL

when calculating BINs. Many authors – including Sharkey et al. (2021) – argue that the

use of K2P NJ trees is acceptable because they are "only" used for visualization.

However, poorly designed visualization tools can misinform about the validity and

support for groupings.

All of the issues herein should have been addressed during manuscript review. This is

why we urge reviewers, journals, and grantors to be more rigorous when reviewing

barcoding studies and projects. A lenient attitude toward BIN data and BOLD has

resulted in scientific publications that do not meet the minimum standards for

reproducible science. Some journals seem to neglect their own data and code

accessibility standards or else papers that are critically dependent on BINs would not be

published. Similarly, a major funder of DNA barcoding is Genome Canada, which has an

explicit data sharing policy that contradicts how RESL and alignment parameters are

handled (see https://www.genomecanada.ca/en/programs/guidelines-and-policies).

Final Remarks and Suggestions

Sharkey et al. (2021) are pessimistic about new 21st century solutions to the dark-taxon

impediment. We do not share this sentiment. Large throughput imaging and sequencing

are already available, these are not vague promises for the future (Hebert et al., 2018;

Ärje et al., 2020; Srivathsan et al., 2021; Wührl et al., 2021). Similarly, data can now be

analyzed with increasingly sophisticated algorithms that will provide taxonomists with a

solid foundation for species descriptions that can be based on multiple sources of data

(Puillandre et al., 2012b; Hartop et al., 2021). These data will be particularly suitable for

generating automatic species descriptions that are resilient and future ready. Therefore,

this is the worst time to propose "minimalist revisions" that trade scientific rigor for speed.

We would like to conclude our critique of Sharkey et al. (2021) with some suggestions for

BOLD:

•   The RESL algorithm must be published and access must be provided to the private

data, or these data must be excluded from BIN calculation.

•   BOLD Systems should implement additional clustering algorithms for barcode data.

This would provide users with much needed information on which barcode clusters

are congruent across methods.

•   BOLD Systems should include a BIN tracking tool for "BIN concepts." It is essential

for validating the results of studies that used BINs computed in the past.

•   The relationship between BINs and species in BOLD Systems should be revisited.

The original BOLD Systems publication highlighted the workbench character of the

database. BINs known to contain several verified species were supposed to be

labeled with decimal numbers ("BOLD:AAB2314.1"). However, this option has been

largely ignored. This means that the BIN matches of searches within BOLD Systems

or GBIF's SequenceID lead to BINs, even if they are known to contradict species

boundaries. BINs containing sequences for multiple species should be flagged.

Acknowledgements

We would like to thank Drs Roderic D. M. Page, Steve Marshall, and one additional

colleague for providing valuable comments on the manuscript. This work was supported

by a Ministry of Education grant on biodiversity discovery (R-154-000-A22-112).

Data accessibility statement

All alignments and scripts are available from

https://figshare.com/articles/dataset/BOLD_braconid_datasets/14452992.

References

Ahrens, D., Ahyong, S. T., Ballerio, A., Barclay, M. V. L., Eberle, J., Espelund, M., Huber,
B. A., Mengual, X., Pacheco, T. L., Peters, R. S., Rulik, B., Vaz-de-Mello, F.,
Wesener, T., K, rell, F. T. 2021. Is it time to describe new species without
diagnoses? - A comment on Sharkey et al. (2021). Zenodo,
10.5281/zenodo.4899151.
Ang, Y., Puniamoorthy, J., Pont, A. C., Bartak, M., Blanckenhorn, W. U., Eberhard, W.
G., Puniamoorthy, N., Silva, V. C., Munari, L., Meier, R. 2013a. A plea for digital
reference collections and other science-based digitization initiatives in taxonomy:
Sepsidnet as exemplar. Systematic Entomology 38, 637-644.
Ang, Y., Rajaratnam, G., Su, K. F., Meier, R. 2017. Hidden in the urban parks of New
York City: *Themira lohmanus*, a new species of Sepsidae described based on
morphology, DNA sequences, mating behavior, and reproductive isolation
(Sepsidae, Diptera). ZooKeys, 95-111.
Ang, Y. C., Wong, L. J., Meier, R. 2013b. Using seemingly unnecessary illustrations to
improve the diagnostic usefulness of descriptions in taxonomy-a case study on
Perochaeta orientalis (Diptera, Sepsidae). Zookeys, 9-27.
Araujo, D., Tuan, M., Yew, J., Meier, R. 2014. Analysing small insect glands with UV-LDI
MS: high-resolution spatial analysis reveals the chemical composition and use of
the osmeterium secretion in *Themira superba* (Sepsidae: Diptera). Journal of
Evolutionary Biology 27, 1744-1750.
Ärje, J., Melvad, C., Jeppesen, M. R., Madsen, S. A., Raitoharju, J., Rasmussen, M. S.,
Iosifidis, A., Tirronen, V., Gabbouj, M., Meissner, K., Høye, T. T. 2020. Automatic
image-based identification and biomass estimation of invertebrates. 11,
922-931.
Bartolini, I., Rivera, J., Nolazco, N., Olortegui, A. 2020. Towards the implementation of a
DNA barcode library for the identification of Peruvian species of *Anastrepha*
(Diptera: Tephritidae). Plos One 15.
Bergsten, J., Bilton, D. T., Fujisawa, T., Elliott, M., Monaghan, M. T., Balke, M.,
Hendrich, L., Geijer, J., Herrmann, J., Foster, G. N., Ribera, I., Nilsson, A. N.,

Barraclough, T. G., Vogler, A. P. 2012. The effect of geographical scale of
sampling on DNA barcoding. Systematic Biology 61, 851–869.
Burns, J. M., Janzen, D. H., Hajibabaei, M., Hallwachs, W., Hebert, P. D. N. 2008. DNA
barcodes and cryptic species of skipper butterflies in the genus *Perichares* in
Area de Conservacion Guanacaste, Costa Rica. Proceedings of the National
Academy of Sciences of the United States of America 105, 6350-6355.
Carstens, B. C., Pelletier, T. A., Reid, N. M., Satler, J. D. 2013. How to fail at species
delimitation. Molecular Ecology 22, 4369-4383.
Chacon, I. A., Janzen, D. H., Hallwachs, W., Sullivan, J. B., Hajibabaei, M. 2013. Cryptic
species within cryptic moths: new species of *Dunama* Schaus (Notodontidae,
Nystaleinae) in Costa Rica. Zookeys, 11-45.
Collins, R., Cruickshank, R. 2013. The seven deadly sins of DNA barcoding. Molecular
Ecology Resources 13, 969-975.
Collins, R. A., Boykin, L. M., Cruickshank, R. H., Armstrong, K. F. 2012. Barcoding's next
top model: an evaluation of nucleotide substitution models for specimen
identification. Methods in Ecology and Evolution 3, 457-465.
DeSalle, R., Egan, M. G., Siddall, M. 2005. The unholy trinity: taxonomy, species
delimitation and DNA barcoding. Philosophical Transactions of the Royal Society
B-Biological Sciences 360, 1905-1916.
Fernandez-Triana, J. L., Whitfield, J. B., Rodriguez, J. J., Smith, M. A., Janzen, D. H.,
Hallwachs, W. D., Hajibabaei, M., Burns, J. M., Solis, M. A., Brown, J., Cardinal,
S., Goulet, H., Hebert, P. D. N. 2014. Review of *Apanteles* sensu stricto
(Hymenoptera, Braconidae, Microgastrinae) from Area de Conservacion
Guanacaste, northwestern Costa Rica, with keys to all described species from
Mesoamerica. Zookeys, 1-565.
Fleming, A. J., Wood, D. M., Smith, M. A., Dapkeyl, T., Hallwachs, W., Janzen, D. 2019.
Twenty-two new species in the genus *Hyphantrophaga* Townsend (Diptera:
Tachinidae) from Area de Conservacion Guanacaste, with a key to the species of
Mesoamerica. Biodiversity Data Journal 7.
Fleming, A. J., Wood, D. M., Smith, M. A., Hallwachs, W., Janzen, D. H. 2014. Revision
of the New World species of *Houghia* Coquillett (Diptera, Tachinidae) reared from
caterpillars in Area de Conservacion Guanacaste, Costa Rica. Zootaxa 3858,
1-90.
Forum, W. E. 2020. World Economic Forum. The Global Risks Report 2020.
Franz, N. M. 2005. On the lack of good scientific reasons for the growing
phylogeny/classification gap. Cladistics 21, 495-500.
Gibbs, J. 2018. DNA barcoding a nightmare taxon: assessing barcode index numbers
and barcode gaps for sweat bees. Genome 61, 21-31.
Hajibabaei, M., Janzen, D. H., Burns, J. M., Hallwachs, W., Hebert, P. D. N. 2006. DNA
barcodes distinguish species of tropical Lepidoptera. Proceedings of the National
Academy of Sciences of the United States of America 103, 968-971.
Hartop, E., Srivathsan, A., Ronquist, F., Meier, R. 2021. Large-scale Integrative
Taxonomy (LIT): resolving the data conundrum for dark taxa. bioRxiv
10.1101/2021.1104.1113.439467.

Hausmann, A., Krogmann, L., Peters, R. S., Rduch, V., Schmidt, S. 2020. GBOL III: Dark
taxa. Barcode Bulletin 10.
Hebert, P. D., Braukmann, T. W., Prosser, S. W., Ratnasingham, S., DeWaard, J. R.,
Ivanova, N. V., Janzen, D. H., Hallwachs, W., Naik, S., Sones, J. E. 2018. A
Sequel to Sanger: amplicon sequencing that scales.   19, 1-14.
Hebert, P. D. N., Penton, E. H., Burns, J. M., Janzen, D. H., Hallwachs, W. 2004. Ten
species in one: DNA barcoding reveals cryptic species in the Neotropical skipper
butterfly *Astraptes fulgerator*. Proceedings of the National Academy of Sciences
of the United States of America 101, 14812-14817.
Janzen, D. H., Hallwachs, W., Burns, J. M., Hajibabaei, M., Bertrand, C., Hebert, P. D. N.
2011. Reading the complex skipper butterfly fauna of one tropical place. Plos One
6.
Kekkonen, M., Hebert, P. D. N. 2014. DNA barcode-based delineation of putative
species: efficient start for taxonomic workflows. Molecular Ecology Resources 14,
706-715.
Kwong, S., Srivathsan, A., Vaidya, G., Meier, R. 2012. Is the COI barcoding gene
involved in speciation through intergenomic conflict? Molecular Phylogenetics and
Evolution 62, 1009-1012.
May, R. M. 2011. Why worry about how many species and their loss? PLoS Biology 9.
Meier, R. 2016. Citation of taxonomic publications: the why, when, what and what not.
Systematic Entomology 42, 301-304.
Meier, R., Shiyang, K., Vaidya, G., Ng, P. K. L. 2006. DNA barcoding and taxonomy in
diptera: A tale of high intraspecific variability and low identification success.
Systematic Biology 55, 715-728.
Nugent, C. M., Elliott, T. A., Ratnasingham, S., Adamowicz, S. J. 2020. Coil: an R
package for cytochrome c oxidase I (COI) DNA barcode data cleaning,
translation, and error evaluation. Genome 63, 291-305.
Ortiz, A. S., Rubio, R. M., Guerrero, J. J., Garre, M. J., Serrano, J., Hebert, P. D. N.,
Hausmann, A. 2017. Close congruence between Barcode Index Numbers (bins)
and species boundaries in the Erebidae (Lepidoptera: Noctuoidea) of the Iberian
Peninsula. Biodiversity Data Journal 5.
Packer, L., Monckton, S. K., Onuferko, T. M., Ferrari, R. R. 2018. Validating taxonomic
identifications in entomological research. Insect Conservation and Diversity 11,
1-12.
Pentinsaari, M., Salmela, H., Mutanen, M., Roslin, T. 2016. Molecular evolution of a
widely-adopted taxonomic marker (COI) across the animal tree of life. Scientific
Reports 6, 35275.
Pohjoismaki, J. L. O., Kahanpaa, J., Mutanen, M. 2016. DNA barcodes for the Northern
European tachinid flies (Diptera: Tachinidae). Plos One 11.
Puillandre, N., Brouillet, S., Achaz, G. 2021. ASAP: assemble species by automatic
partitioning. Molecular Ecology Resources 21, 609-620.
Puillandre, N., Lambert, A., Brouillet, S., Achaz, G. 2012a. ABGD, Automatic Barcode
Gap Discovery for primary species delimitation. Molecular Ecology 21,
1864-1877.

Puillandre, N., Modica, M. V., Zhang, Y., Sirovich, L., Boisselier, M.-c., Cruaud, C., Holford, M., Samadi, S. 2012b. Large-scale species delimitation method for hyperdiverse groups. Molecular Ecology 21, 2671-2691.

Puniamoorthy, N., Ismail, M. R. B., Tan, D. S. H., Meier, R. 2009. From kissing to belly stridulation: comparative analysis reveals surprising diversity, rapid evolution, and much homoplasy in the mating behaviour of 27 species of sepsid flies (Diptera: Sepsidae). Journal of Evolutionary Biology 22, 2146-2156.

Puniamoorthy, N., Kotrba, M., Meier, R. 2010. Unlocking the "Black box": internal female genitalia in Sepsidae (Diptera) evolve fast and are species-specific. BMC Evolutionary Biology 10.

Ratnasingham, S., Hebert, P. D. N. 2011. BOLD's role in barcode data management and analysis: a response. Molecular Ecology Resources 11, 941-942.

Ratnasingham, S., Hebert, P. D. N. 2013. A DNA-based registry for all animal species: The Barcode Index Number (BIN) system. Plos One 8.

Raupach, M. J., Hannig, K., Moriniere, J., Hendrich, L. 2016. A DNA barcode library for ground beetles (Insecta, Coleoptera, Carabidae) of Germany: The genus *Bembidion* Latreille, 1802 and allied taxa. Zookeys, 121-141.

Roe, A. D., Sperling, F. A. H. 2007. Patterns of evolution of mitochondrial cytochrome c oxidase I and II DNA and implications for DNA barcoding. Molecular Phylogenetics and Evolution 44, 325-345.

Rohner, P. T., Ang, Y., Lei, Z., Puniamoorthy, N., Blanckenhorn, W. U., Meier, R. 2014. Genetic data confirm the species status of *Sepsis nigripes* Meigen (Diptera: Sepsidae) and adds one species to the Alpine fauna while questioning the synonymy of *Sepsis helvetica* Munari. Invertebrate Systematics 28, 555-563.

Smith, M. A., Rodriguez, J. J., Whitfield, J. B., Deans, A. R., Janzen, D. H., Hallwachs, W., Hebert, P. D. N. 2008. Extreme diversity of tropical parasitoid wasps exposed by iterative integration of natural history, DNA barcoding, morphology, and collections. Proceedings of the National Academy of Sciences of the United States of America 105, 12359-12364.

Smith, M. A., Wood, D. M., Janzen, D. H., Hallwachs, W., Hebert, P. D. N. 2007. DNA barcodes affirm that 16 species of apparently generalist tropical parasitoid flies (Diptera, Tachinidae) are not all generalists. Proceedings of the National Academy of Sciences of the United States of America 104, 4967-4972.

Smith, M. A., Woodley, N. E., Janzen, D. H., Hallwachs, W., Hebert, P. D. N. 2006. DNA barcodes reveal cryptic host-specificity within the presumed polyphagous members of a genus of parasitoid flies (Diptera: Tachinidae). Proceedings of the National Academy of Sciences of the United States of America 103, 3657-3662.

Srivathsan, A., Hartop, E., Puniamoorthy, J., Lee, W. T., Kutty, S. N., Kurina, O., Meier, R. 2019. Rapid, large-scale species discovery in hyperdiverse taxa using 1D MinION sequencing. BMC Biology 17, 96.

Srivathsan, A., Lee, L., Katoh, K., Hartop, E., Kutty, S. N., Wong, J., Yeo, D., Meier, R. J. b. 2021. MinION barcodes: biodiversity discovery and identification by everyone, for everyone. BioRxiv.

Srivathsan, A., Meier, R. 2012. On the inappropriate use of Kimura-2-parameter (K2P) divergences in the DNA-barcoding literature. Cladistics 28, 190-194.

Stamatakis, A. J. B. 2014. RAxML version 8: a tool for phylogenetic analysis and
post-analysis of large phylogenies.   30, 1312-1313.
Takezaki, N. 1998. Tie trees generated by distance methods of phylogenetic
reconstruction. Molecular Biology and Evolution 15, 727-737.
Tan, D. S., Ang, Y., Lim, G. S., Ismail, M. R. B., Meier, R. 2010. From 'cryptic species' to
integrative taxonomy: an iterative process involving DNA sequences, morphology,
and behaviour leads to the resurrection of Sepsis pyrrhosoma (Sepsidae:
Diptera). Zoologica Scripta 39, 51-61.
Virgilio, M., Backeljau, T., Nevado, B., De Meyer, M. J. B. b. 2010. Comparative
performances of DNA barcoding across insect orders. BMC Bioinformatics 11,
1-10.
Will, K. W., Rubinoff, D. 2004. Myth of the molecule: DNA barcodes for species cannot
replace morphology for identification and classification. Cladistics 20, 47–55.
Wührl, L., Pylatiuk, C., Giersch, M., Lapp, F., von Rintelen, T., Balke, M., Schmidt, S.,
Cerretti, P., Meier, R. 2021. DiversityScanner: Robotic discovery of small
invertebrates with machine learning methods. 2021.2005.2017.444523.
Yeo, D., Srivathsan, A., Meier, R. 2020. Longer is not always better: Optimizing barcode
length for large-scale species discovery and Identification. Systematic Biology 69,
999-1015.
Zamani, A., Vahtera, V., Sääksjärvi, I. E., Scherz, M. D. 2021. The omission of critical
data in the pursuit of 'revolutionary' methods to accelerate the description of
species. Systematic Entomology 46, 1-4.
Zhang, J., Kapli, P., Pavlidis, P., Stamatakis, A. 2013. A general species delimitation
method with applications to phylogenetic placements. Bioinformatics 29,
2869-2876.

729