Differential tracking of linguistic vs. mental state content in naturalistic stimuli by language and Theory of Mind (ToM) brain networks

Alexander M. Paunov[1,2], Idan A. Blank[1,2,3], Olessia Jouravlev[1,2,4], Zachary Mineroff[1,2], Jeanne Gallée[5] & Evelina Fedorenko[1,2,5]

[1] Department of Brain and Cognitive Sciences, MIT, Cambridge MA 02139
[2] McGovern Institute for Brain Research, MIT, Cambridge MA 02139
[3] Department of Psychology, UCLA, Los Angeles CA 90095
[4] Institute for Cognitive Science, Carleton University, Ottawa, Canada K1S 5B6
[5] Program in Speech and Hearing Biosciences and Technology, Harvard University, Boston, MA, 02115

Corresponding authors: Alex Paunov (apaunov@mit.edu; 43 Vassar Street, Room 3037, Cambridge, MA 02139) and Ev Fedorenko (evelina9@mit.edu; 43 Vassar Street, Room 3037, Cambridge, MA 02139).

Acknowledgments
We would like to acknowledge the Athinoula A. Martinos Imaging Center at the McGovern Institute for Brain Research at MIT, and its support team (Steve Shannon and Atsushi Takahashi). We thank former and current EvLab members (especially Brianna Pritchett and Caitlyn Hoeflin) for their help with data collection, Amaya Arcelus for help with creating the Heider & Simmel style animation, Caroline Arnold for help with recording the dialog condition, and Jessica Chen and Yotaro Sueoka for help with manuscript preparation. This research and EF were supported by R01 awards DC016607 and DC016950 from NIDCD, a grant from the Simons Foundation to the Simons Center for the Social Brain at MIT, and the Newton Brain Science Award from the Brain and Cognitive Sciences Department at MIT.

Contributions
AP, IB and EF developed the concept for the study. AP, IB and JG selected / created experimental materials. AP, IB, OJ and ZM collected and analyzed the data. AP performed the final version of the statistical analyses for the current study that are reported here. AP, IB, and EF wrote the manuscript with input from other co-authors.

Conflict of interest
Authors report no conflict of interest.

38 **Differential tracking of linguistic vs. mental state content in naturalistic stimuli by**
39 **language and Theory of Mind (ToM) brain networks**
40
41 **Short title:** Language vs. ToM Networks in Naturalistic Stimulus Tracking
42
43 **Keywords:** language, theory of mind, social cognition, intersubject correlations, naturalistic
44 fMRI

45
Abstract

Language and social cognition, especially the ability to reason about mental states, known as

Theory of Mind (ToM), are deeply related in development and everyday use. However, whether

these cognitive faculties rely on distinct, overlapping, or the same mechanisms remains debated.

Some evidence suggests that, by adulthood, language and ToM draw on largely distinct—though

plausibly interacting—cortical networks. However, the broad topography of these networks is

similar, and some have emphasized the importance of social content / communicative intent in

the linguistic signal for eliciting responses in the language areas. Here, we combine the power of

individual-subjects functional localization with the naturalistic-cognition inter-subject correlation

approach to illuminate the language-ToM relationship. Using fMRI, we recorded neural activity

as participants (n=43) listened to stories and dialogs with mental state content (+linguistic,

+ToM), viewed silent animations and live action films with mental state content but no language

(-linguistic, +ToM), or listened to an expository text (+linguistic, -ToM). The ToM network

robustly tracked stimuli rich in mental state information regardless of whether mental states were

conveyed linguistically or non-linguistically, while tracking a +linguistic/-ToM stimulus only

weakly. In contrast, the language network tracked linguistic stimuli more strongly than a) non-

linguistic stimuli, and than b) the ToM network, and showed reliable tracking even for the

linguistic condition devoid of mental state content. These findings suggest that in spite of their

indisputably close links, language and ToM dissociate robustly in their neural substrates—and

thus plausibly cognitive mechanisms—including during the processing of rich naturalistic

materials.

69
1. Introduction

Language and social cognition, especially the ability to reason about mental states, known as

Theory of Mind (ToM), are deeply related in human development, everyday use, and possibly

evolution. After all, language use is a communicative behavior, which is a kind of cooperative

behavior, and cooperative behaviors are, in turn, a kind of social behavior (e.g., Grice, 1968,

1975, Sperber & Wilson, 1986). Construed this way, language can hardly be encapsulated from

social cognition (cf. Fodor, 1983); it is subsumed *within* social cognition. Interpreting linguistic

signals bears key parallels to the interpretation of other intentional behaviors (e.g., Grice, 1975;

Sperber & Wilson, 1986). Communicative utterances, like other behaviors, are assumed to have

goals, and conversation partners are assumed to pursue these goals rationally. Furthermore,

everyday discourse appears to be dominated by information about other people (e.g., Dunbar,

Marriott & Duncan, 1997), and the need to keep track of others' social record has been proposed

as a key driver of language evolution (e.g., Dunbar, 2004; Nowak & Sigmund, 2005;

Sommerfield, Krambeck & Milinski, 2008; Nowak & Highfield, 2011). Lastly, evidence of

others' mental states conveyed through language is arguably richer and certainly more direct /

less ambiguous than what can be inferred from non-linguistic intentional behavior alone: trying

to infer the beliefs guiding someone's actions can be obviated by their telling you what those

beliefs are.

Yet in the relatively brief history of cognitive neuroscience, the study of language and of social

cognition have, for the most part, proceeded independently. The two cognitive domains have

been treated, often implicitly, as components of a "nearly decomposable system" (Simon,

1962)—a system in which interactions between the component subsystems (language and social

cognition here) are sufficiently less common than interactions among each subsystem's parts to

warrant a "divide and conquer" strategy (Saxe, Brett, & Kanwisher, 2006). On such a strategy,

each subsystem can first be investigated on its own, leaving questions of potential dependencies

between subsystems for a later stage. The application of this strategy to language and social

cognition has been remarkably successful, and even this later stage is now coming within reach.

But is near-decomposability a true property of the relationship between language and social

cognition, or merely a simplifying assumption? It is difficult to say because most studies to date

have deliberately aimed to isolate one or the other component subsystem rather than probe their

relationship (cf. Deen et al., 2015; Paunov et al., 2019; Braga et al., 2020). So we know that

language and social cognition *can* dissociate, under appropriate experimental conditions. But do

they, in fact, dissociate in everyday, naturalistic cognition? This is the question we tackle in the

current study.

Before we proceed to describe the current study, we first describe two ways in which language

and ToM are related in adult cognition and clarify our research aims with respect to these. Then

we review what is currently known about the neural bases of linguistic processing and of Theory

of Mind.

1.1. The role of ToM in inferring communicative intent vs. processing linguistic content.

Two aspects of the relationship between language and social cognition can be identified in the

context of communication: i) the role of mental state inference in language comprehension

generally, whether or not the content concerns mental states; and ii) the use of language to

express information about the mental states of agents, either directly or through descriptions of

physical events, which prompt mental state attribution (e.g., Fletcher et al., 1995; Gallagher et

al., 2000; Saxe & Kanwisher, 2003; Saxe, Schulz, & Jiang, 2006; Jacoby et al., 2016). In the

present study, we focus primarily on the second of these aspects.

The first aspect of the language-ToM relationship is generally subsumed in the broad category of

pragmatics, which is traditionally concerned with how communicative intent—a form of ToM

inference—guides linguistic interpretation (e.g., Grice, 1957, 1968, 1975). A key starting point

of pragmatics is that the intended meaning of utterances is vastly underdetermined by linguistic

inputs (Wittgenstein, 1953; Sperber & Wilson, 1986), and ToM must be recruited to draw

inferences about what a speaker intends to convey, based on evidence contained in the speaker's

utterances and the extended context (linguistic, visual, social, etc.). However, delimiting the

scope of pragmatic inference is a long-standing challenge, raising the question of whether it is

possible to draw a boundary between decoded (literal) and inferred meaning, that is, between

semantics and pragmatics (e.g., Jackendoff, 2009).

The lack of clearly defined boundaries for the construct of pragmatics poses empirical

challenges. For one, pragmatic inference need not be limited to linguistic communication: it can

equally be present in other forms of cooperative information transfer between individuals, such

as the interpretation of communicative gestures or, more relevant in the present context,

understanding non-linguistic 'stories' such as non-verbal animated or live-action films. At the

same time, within the narrower context of linguistic communication, it is implausible that all

forms of context-based inference of meaning recruit mental state reasoning. Phenomena that

require context-based inferences include not just the paradigmatic instances of non-literal

meaning, such as irony, indirect requests, hyperbole, and other 'conversational implicatures'

(Grice, 1975), but also relatively 'low-level' and ubiquitous phenomena such as pronoun

resolution, or lexical and syntactic disambiguation.

Further, different situational contexts plausibly differ in the nature and extent of their social-

cognitive demands. In **Figure 1**, we schematically illustrate three contexts for language

processing: reading or listening to narratives or expository texts, reading or listening to a

conversation, and directly participating in face-to-face conversation. For each, we outline the

ToM cognitive demands associated with, on the one hand, a) understanding the *communicative*

intent of the person generating the linguistic output, and, on the other, b) the processing of the

content of the linguistic materials. Past work has investigated both kinds of ToM demands. For

example, a number of studies have manipulated the difficulty of inferring the communicative

intent of a speaker by examining the kinds of paradigmatic cases of non-literal language

mentioned above, from irony (e.g., Spotorno et al., 2012), to indirect speech (e.g., Feng et al,

2017; van Ackeren et al., 2012) and other forms of conversational implicature (e.g., Feng, Yu, &

Zhou, 2021; Jang et al., 2013; see Hagoort & Levinson, 2014 for review). These studies have

reported stronger responses in the ToM network for the critical, non-literal stimuli compared to

literal controls. Other studies have varied the amount of mental state content in verbal vignettes

or stories (e.g., Ferstl & von Cramon, 2002; Saxe & Powell, 2006) and reported stronger activity

in the ToM network for materials rich in mental state content (although even expository

connected texts appear to engender responses in the ToM network relative to sets of unconnected

sentences; Jacoby et al., 2020). So, both kinds of demands appear capable of recruiting mental

state reasoning.

In the current study, we focus on the presence or absence of ToM content in rich naturalistic

materials—in line with current emphasis in the field on the importance of going beyond carefully

controlled experimental materials in the testing of hypotheses about human cognitive

architecture (e.g., Blank, Kanwisher & Fedorenko, 2014; Huth et al., 2016; Hasson et al, 2018)—

and evaluate the evidence for *differential recruitment* of language and ToM cortical regions

across these materials. Establishing robustly dissociable neural substrates for language

understanding and mental state inference is an important step toward bringing neural evidence to

bear on the differentiation of pragmatics into distinct sub-domains of context-based inference,

which may differentially entail mental state attribution (e.g., Sperber & Wilson, 2002; Hagoort &

Levinson, 2014; Bosco, Tirassa, & Gabbatore, 2018).

Figure 1. *Cognitive demands on ToM processing across different contexts of discourse-level*

linguistic processing with respect to understanding the communicative intent of the producer,

and the linguistic content of the materials. The comprehender is circled in red. Whereas specific

demands differ across contexts, any form of language use arguably involves attribution of

communicative intent (pragmatics) in the service of comprehension and / or production.

1.2. Neural bases of language and Theory of Mind.

Language network. Language processing engages a network of left-lateralized brain regions in

lateral frontal and temporal cortex. These regions support lexico-semantic processing (word

meanings) and combinatorial morpho-syntactic and semantic processing (e.g., Fedorenko et al.,

2010, 2012, 2016, 2020; Bautista & Wilson, 2016; Blank et al., 2016; Anderson et al., 2018;

Reddy & Wehbe, 2020). Responses to linguistic stimuli generalize across different materials,

tasks, modalities of presentation (reading vs. listening), languages, and developmental

trajectories (e.g., Fedorenko et al., 2010; Bedny et al., 2011; Fedorenko, 2014; Scott et al., 2017;

Diachek, Blank, Siegelman et al., 2020; Ayyash, Malik-Moraleda et al., in prep.; Ivanova et al.,

in prep.). In contrast to these robust and consistent responses to linguistic stimuli, these regions

do not respond to a wide range of non-linguistic cognitive processes, including arithmetic

processing, music perception, executive function tasks, action or gesture observation, non-verbal

social-cognitive tasks, and the processing of computer code (e.g., Fedorenko et al., 2011; Monti

et al., 2012; Pritchett et al., 2018; Almaric & Dehaene, 2018; Paunov, 2018; Jouravlev et al.,

2019; Ivanova et al., 2020; Shain, Paunov et al., in prep.; see Fedorenko & Varley, 2016 for a

review).

ToM network. Attribution of mental states, such as beliefs, desires, and intentions, engages a

network of brain regions in bilateral temporo-parietal cortex and anterior and posterior regions

along the cortical midline. These responses generalize across the type of mental state, its specific

content or format (linguistic vs. pictorial), and the source of evidence for it (e.g., Fletcher et al.,

1995; Castelli et al., 2000; Gallagher et al., 2000; Vogeley et al., 2001; Ruby & Decety, 2003;

Saxe & Kanwisher, 2003; Saxe, Schulz, & Jiang, 2006; Kandylaki et al., 2015; Jacoby et al.,

2016; see Koster-Hale & Saxe, 2013 for review) as well as across populations with diverse

cultural backgrounds or developmental experiences (e.g., Callaghan et al., 2005; Sabbagh et al.,

2006; Shahaeian et al., 2011; Koster-Hale, Bedny, & Saxe, 2014; Richardson et al., 2018, 2020).

However, by adulthood, these regions, and especially the most selective component of the ToM

network, the right temporo-parietal junction (RTPJ), do not respond to social stimuli, like faces,

voices, or biological motion (e.g., Deen et al., 2015), to general executive demands (e.g., Saxe,

Schulz, & Jiang, 2006), to physical or broadly social attributes of agents, or to attribution of

bodily sensations of pain and hunger (e.g., Saxe & Powell, 2006; Bruneau, Pluta & Saxe, 2012;

Jacoby et al., 2016).

Together, these two lines of work support the notion that language and ToM are distinct

cognitive faculties, each mapping onto functionally specialized neural structures. Investigations

of developmental and acquired disorders have provided convergent support for the dissociability

of language and ToM. Individuals with even severe aphasia appear to retain the capacity for

mental state reasoning as long as non-verbal materials are used (e.g., Varley & Siegal, 2000;

Varley, Siegal & Want, 2001; Apperly et al., 2006; Willems et al., 2011; see Fedorenko &

Varley, 2016 for review). And at least some individuals with social, ToM-related impairments

(e.g., some individuals with autism spectrum disorders) show preservation of lexical and

syntactic linguistic abilities (e.g., Frith & Happe, 1994; Wilkinson, 1998; Tager-Flusberg, Paul &

Lord, 2005; Tager-Flusberg, 2006). But as alluded to above, most of this evidence has come

from traditional, task-based experimental paradigms, and such paradigms are well-suited to the

divide-and-conquer enterprise, but may overestimate the degree of separation. Indeed, Paunov et

al. (2019) recently examined inter-regional functional correlations within and between the

language and ToM networks during naturalistic cognition paradigms, like story comprehension,

and found that in spite of their dissociability (stronger within- than between-network

correlations), the language and ToM networks also showed a significant amount of

synchronization in their neural activity. These results point to some degree of functional

integration between the networks. Furthermore, several prior studies have argued for the

importance of social content / communicative intent in the linguistic signal for eliciting

responses in the language areas (e.g., Mellem et al., 2016; Redcay et al., 2016). Here, we adopt

an approach that relies on the use of rich naturalistic stimuli but allows us to not only examine

the structure of and the relationship between the networks, but to also ask what stimulus features

each network "tracks" through the inter-subject correlation approach (e.g., Hasson et al., 2004,

2008).

2. Materials and Methods

2.1. General approach.

Following Blank & Fedorenko (2017, 2020), we combine two powerful methodologies, which

have previously been productively applied separately in the domains of language and social

cognition. In particular, we use *functional localization* (e.g., Brett et al., 2002; Saxe et al., 2006;

Fedorenko et al., 2010) to identify the two networks of interest in individual subjects, and *inter-*

subject correlations (ISCs; e.g., Hasson et al., 2004, 2008) to examine the degree to which these

networks 'track' different stimulus features during the processing of rich naturalistic materials.

Here, we highlight key strengths of each approach and consider their synergistic advantages in

the context of our research goals.

Naturalistic paradigms have become a crucial complement to traditional, task-based studies in

cognitive neuroscience. The obvious advantage of naturalistic paradigms is their high ecological

validity: by giving up a measure of experimental control, it becomes possible to study cognition

"in the wild" (Blank & Fedorenko, 2017; see e.g., Sonsukare et al., 2019 and Nastase et al., 2020

for general discussions). In particular, one can examine how coherent and structured mental

representations are extracted from rich and noisy perceptual inputs, which is what happens in

everyday cognition. This is in contrast to artificially isolating various features of these perceptual

inputs, as is typically done in constrained experimental tasks. Naturalistic paradigms have been

argued to elicit more reliable responses compared to traditional, task-based paradigms (Hasson et

al., 2010), perhaps because they are generally more engaging, and can enable discoveries of

functional relationships among brain regions and networks that are altogether missed in more

constrained settings (e.g., Gallivan, Cavina-Pratesi & Culham, 2009). Another advantage of

naturalistic paradigms is their hypothesis-free nature. Through the use of naturalistic materials,

researchers impose minimal design constraints to investigate the domain of interest in a manner

that is maximally unbiased by prior theoretical assumptions. In effect, they are letting the data

speak for itself.

However, naturalistic paradigms also come with an inherent analytic challenge: how do we make

sense of data acquired without the typical constraints of standard hypothesis-driven modeling

approaches? Hasson et al. (2004) pioneered an approach to tackle this challenge, known as the

inter-subject correlation (ISC) approach (see Hasson et al., 2008, for an overview), which we

adopt in the current study. The key insight behind the ISC approach is that we can model any

given participant's fMRI signal time series using another participant's or other participants' time

series: if a voxel, brain region, or brain network 'tracks' features of the stimulus during which

the time series were obtained, then fMRI signal fluctuations will be stimulus-locked, resulting in

similar time courses across participants (i.e., high inter-subject correlations).

ISCs have been used in several studies of narrative comprehension (e.g., Wilson et al., 2007;

Lerner et al., 2011; Honey et al., 2012; Regev et al, 2013; Silber et al., 2014; Schmälzle et al.,

2015; Blank & Fedorenko, 2017, 2020), and whole-brain voxel-wise analyses have revealed high

inter-subject correlations across large swaths of cortex that resemble the union of the language

and ToM networks. On their own, these results might be taken as *prima facie* evidence for non-

dissociability of language and ToM, given that the two networks appear to be jointly recruited.

And insofar as the mental processes recruited in narrative comprehension recapitulate those used

in everyday abstract cognition—an assumption that, we take it, partially justifies the interest in

narratives in cognitive science and neuroscience (e.g., Finlayson & Winston, 2011; Willems et

al., 2020)—the results may be taken to suggest the non-dissociability of language and ToM more

generally.

However, it is difficult to draw inferences from these studies about the *relative* contributions of

the language and ToM networks to narrative comprehension for two reasons. First, in whole-

brain analyses, ISCs are computed on a voxel-wise basis: individual brains are normalized to a

stereotaxic template, and one-to-one voxel correspondence across individuals is then assumed in

computing the ISCs. This approach is problematic because i) inter-individual variability is well

established in the high-level association cortex (e.g., Fischl et al., 2008; Frost & Goebel, 2012;

Tahmasebi et al., 2012), so any given voxel may belong to functionally distinct regions across

participants; and ii) there is no independent criterion based on which an anatomical location can

be interpreted as belonging to the language vs. the ToM network, thus necessitating reliance on

the fallacious "reverse inference" (Poldrack, 2006) to interpret the resulting topography (see

Fedorenko, 2021, for discussion). And second, traditional whole-brain analyses typically include

all voxels that showed significant (above baseline) ISCs, thus potentially obscuring large

differences in effect sizes (cf. Blank & Fedorenko, 2017; see Chen, Taylor & Cox, 2017, for a

general discussion of the importance of considering effect sizes in interpreting fMRI findings).

Combining the ISC approach with individual-participant functional localization enables us to

identify and directly compare the networks of interest (including with respect to effect sizes), as

well as to relate the findings straightforwardly to the prior literature on the language and ToM

networks. In the current study, we therefore identified the language and ToM networks using

well-established functional localizers (Saxe & Kanwisher, 2003; Fedorenko et al., 2010), and

then examined the degree of inter-subject synchronization in those regions during the processing

of diverse naturalistic linguistic and non-linguistic conditions varying in the presence of mental

state content. If the language and ToM networks are dissociable during naturalistic cognition, we

would expect the language regions to track linguistic stimuli, including those that lack mental

state content, and the ToM regions to track stimuli that have mental state content, including both

linguistic and non-linguistic ones.

2.2. Overall experimental design and statistical analyses.

Our overall design and analytic strategy were as follows: participant-specific regions that

responded more strongly during the reading of sentences compared with lists of nonwords were

defined as regions of interest comprising the language network. Similarly, regions that responded

more strongly to stories about others' beliefs vs. stories about physical reality were defined as

regions of interest comprising the ToM network (see ***Section 2.4*** for details). Whereas the

precise anatomical locations of these regions were allowed to vary across participants, their

overall topography was constrained by independently derived criteria to establish functional

correspondence across brain regions of different participants (e.g., Fedorenko et al., 2010; Julian

et al., 2012).

Activity in these two sets of brain regions was recorded with fMRI while participants listened to

or watched a series of naturalistic stimuli, as detailed below. For each region in each network,

our critical dependent variable was the strength of the correlation between each participant's

time series and the average time series from the rest of the participants. The group-averaged ISC

in each region was tested for significance via a permutation test of the time series data. For our

critical analysis, all individual ISC values were modeled using a linear mixed-effects regression

with participant, brain region, and stimulus (what we call 'condition' below) as random effects.

2.3. Participants.

Forty-seven native English speakers (age 19-48, M = 24.5, SD = 5.08; 30 female) from MIT and

the surrounding Boston community participated for payment. Forty participants were right-

handed, as determined by the Edinburgh handedness inventory (Oldfield 1971) or by self-report

(n=1). All seven left-handed participants showed typical left lateralization in the language

localizer task described below (see Willems et al., 2014, for arguments to include left handers in

cognitive neuroscience research). Four participants were excluded from the analyses due to poor

quality of the localizer data (2 for ToM localizer, 1 for language localizer, and 1 for both), with

the exclusion criterion defined as fewer than 100 suprathreshold voxels (at the $p < 0.001$

uncorrected whole-brain threshold) across the respective network's masks (see below), bringing

the number of participants included in the critical analyses to 43. All participants gave informed

written consent in accordance with the requirements of MIT's Committee on the Use of Humans

as Experimental Subjects (COUHES).

2.4. Stimuli and procedure.

Each participant completed a language localizer, a ToM localizer, a localizer for the domain-

general Multiple Demand (MD) system (used in a replication analysis, as described below), and

a subset, or all, of the critical naturalistic stimuli ('conditions') (between 1 and 7) due to scan

duration constraints (18 participants completed all 7 conditions of interest, 1 participant

completed 6 conditions, 10 participants completed 5 conditions, 1 participant completed 4

conditions, 1 participant completed 3 conditions, 7 participants completed 2 conditions, and 5

participants completed 1 condition). Each condition was presented to between 28 and 32

participants (see **Table 1**). Each stimulus, lasting ~5-7 minutes (see **Table 1** for precise

durations), was preceded and followed by 16 s of fixation. Finally, ten participants performed a

resting state scan, used in one of the reality-check analyses, as described below. For the language

localizer, 36/43 participants completed it in the same session as the critical conditions, the

remaining 7 participants completed it in an earlier session. Similarly, for the ToM and MD

localizers, 37/43 participants completed them in the same session as the critical conditions, the

remaining 6 participants completed them in an earlier session. We will now describe the

localizers and the critical experiment in more detail.

Language localizer task. The task used to localize the language network is described in detail in

Fedorenko et al. (2010) and targets brain regions that support high-level language processing.

Briefly, we used a reading task contrasting sentences and lists of unconnected, pronounceable

nonwords (**Figure 2**) in a blocked design with a counterbalanced condition order across runs.

Stimuli were presented one word / nonword at a time. Participants read the materials passively

(we included a button-pressing task at the end of each trial, to help participants remain alert). As

discussed in the introduction, this localizer is robust to task manipulations (e.g., Fedorenko et al.,

2010; Scott et al., 2017; Ivanova et al., in prep.). Moreover, this localizer identifies the same

regions that are localized with a broader contrast, between listening to natural speech and its

acoustically-degraded version (Scott et al., 2017; Ayyash, Malik-Moraleda et al., in prep.). All

participants completed two runs, each lasting 358 s and consisting of 8 blocks per condition and

5 fixation blocks. (A version of this localizer is available from http://evlab.mit.edu/funcloc/.)

ToM localizer task. The task used to localize the ToM network is described in detail in Saxe &

Kanwisher (2003) and targets brain regions that support reasoning about others' mental states.

Briefly, the task was based on the classic false belief paradigm (Wimmer and Perner, 1983), and

contrasted verbal vignettes about false beliefs (e.g., a protagonist has a false belief about an

object's location; the critical condition) and linguistically matched vignettes about false physical

states (physical representations depicting outdated scenes, e.g., a photograph showing an object

that has since been removed; the control condition) (**Figure 2**) in a long-event-related design

with a counterbalanced order across runs (when multiple runs were administered). Stimuli were

presented one at a time. Participants read these vignettes and answered a true / false

comprehension question after each one. Forty-one participants completed two runs and two

completed one run due to time limitations, each lasting 262 s and consisting of 5 vignettes per

condition. (A version of this localizer is available from http://saxelab.mit.edu/use-our-efficient-

false-belief-localizer.)

An alternative, non-verbal ToM localizer task. One of the naturalistic conditions in this study

("Partly Cloudy", an animated short film; see next section) has been previously used as a non-

verbal ToM localizer (Jacoby et al., 2016; see also Richardson et al., 2018). To that end, it has

been coded into 'mental', 'physical', 'social', and 'pain' segments, and the regions defined by

the mental > pain contrast have been validated against the traditional ToM localizer described

above (see Jacoby et al., 2016 for details). Examples of mental content include a character falsely

believing they have been abandoned by a companion and a character observing others interacting

happily after experiencing pain (4 events, 44 sec total). Following a reviewer's suggestion, we

used this localizer as an alternative ToM localizer in some of the analyses. We used the mental >

physical contrast rather than mental > pain to maintain conceptual similarity with the verbal

localizer's false belief > false physical contrast. The activations obtained with the two different

control conditions were qualitatively similar. Examples of physical content include a wide shot

of clouds and birds flying (3 events, 22 sec total). The main goal was to ensure that the language-

ToM dissociation is not due to an overly narrow definition of Theory of Mind in terms of false

beliefs, implicit in the use of this particular type of mental state attribution in the standard, verbal

localizer. Notably, many previous studies have shown that the ToM network defined with the

false belief localizer responds to a wide range of mental state content besides (false) beliefs,

including intentions, sources of evidence about others' minds, emotional pain, and the 'minds' of

group agents (e.g., Young & Saxe, 2008; Bruneau, Pluta & Saxe, 2012; Koster-Hale, Bedny &

Saxe, 2014; Jenkins et al., 2014). Nevertheless, analyses that use the non-verbal ToM localizer

should confirm that the results of the present study generalize beyond a particular way of

localizing the ToM network.

Figure 2. *Sample trials from the functional localizer paradigms.* ***Language****: reading of*

sentences was contrasted with reading of sequences of pronounceable non-words (Fedorenko et

al., 2010). ***ToM****: reading of vignettes about false mental states was contrasted with reading of*

vignettes about false physical states, each followed by a true/false statement (Saxe & Kanwisher,

2003).

Multiple Demand (MD) localizer task (used in a replication analysis, as described below). The

task used to localize the MD network is described in detail in Fedorenko et al. (2011) and targets

brain regions that support goal-directed effortful behaviors (e.g., Duncan, 2010, 2013). Briefly,

we used a spatial working-memory task contrasting a harder version with an easier version

(**Supplemental Figure A1**) in a blocked design with a counterbalanced condition order across

runs (when multiple runs were administered). On each trial, participants saw a 3x4 grid and kept

track of eight (hard version) or four (easy version) randomly generated locations that were

sequentially flashed two at a time or one at a time, respectively. Then, participants indicated their

memory for these locations in a two-alternative, forced-choice paradigm via a button press, and

received feedback. Of the thirty-two participants included in the replication analysis (i.e., non-

overlapping with those used in the original study in Blank & Fedorenko, 2017), twenty-three

participants completed two runs of the localizer, and nine completed one run, each lasting 448 s

and consisting of 6 blocks per condition and 4 fixation blocks.

The critical naturalistic task. In the main experiment, each participant listened to (over scanner-

safe Sensimetrics headphones) and/or watched a set of naturalistic stimuli (varying between 4

min 50 s and 7 min 6 s in duration). Four of the conditions used linguistic materials: i) a story

("Elvis" from the Natural Stories corpus; Futrell et al., 2020), ii) an audio play (a segment from

an HBO miniseries, "Angels in America", from Chapter 1: "Bad News",

https://www.imdb.com/title/tt0318997/, audio only), iii) a naturalistic dialog—a casual

unscripted conversation between two female friends (recorded by JG), and iv) a non-narrative

expository text (a text about trees adapted from Wikipedia;

https://en.wikipedia.org/wiki/Tree) (recorded by JG). The first three of the linguistic conditions

were rich in mental state content; the fourth was meaningful naturalistic discourse with little/no

mental state content (see below for additional discussion). The three remaining conditions were

videos with no linguistic content: i) an animated short film ("Partly Cloudy" by Pixar), ii) a clip

from a live action film ("Mr. Bean", https://www.youtube.com/watch?v=bh__g-ZZ6WA), and

iii) a custom-created Heider and Simmel style animation (Heider & Simmel, 1944) consisting of

simple geometric shapes moving in ways as to suggest intentional interactions designed to tell a

story (e.g., a shape gets locked up inside a space, another shape goes on a quest to get help to

release it, etc.). All three non-linguistic conditions were rich in mental state content. (Five

additional conditions—included in some participants for another study—are of no interest to the

current study.) All the materials are available on OSF (except in cases where copyright issues

will prevent us from doing so): https://osf.io/prghx/. In the resting state scan, used for one of our

reality-check analyses, as described below, participants were instructed to close their eyes and let

their mind wander but to remain awake while resting in the scanner for 5 min (the scanner lights

were dimmed and the projector was turned off).

It is important to note that although we classify these naturalistic conditions into 'types' in a

binary way (i.e., either involving ToM or not, and either involving language or not), this should

not be taken to suggest that there cannot be gradation within each category. Indeed, given the

richness of the stimuli, there almost certainly is, at least for the ToM dimension. However, we do

not pursue this question further given the small number of stimuli within each category, and their

complexity. Another challenge is coding the materials, especially for mental state content. In

particular, linguistically mediated mental state attribution often proceeds not from explicit

mentions of mental states but from action descriptions, and in non-verbal settings, ToM

attribution proceeds exclusively from action observation. As a result, it is difficult to specify,

especially in complex naturalistic materials, when mental state attribution is prompted.

Conversely, ToM vocabulary need not lead to stronger mental state attribution (e.g., "Alan thinks

that this is a nice house" *vs.* "Alan says that this is a nice house"; presumably, a mental state is

ascribed to Alan in both cases). Thus, we do not attempt to quantify degrees of mental state

attribution beyond the overall presence or (near-) absence of mental state content.

Another important point to acknowledge is that there may be a certain degree of "contamination"

across categories. Specifically, as discussed in the Introduction, the '-ToM' condition (expository

text) plausibly involves some degree of pragmatic inference, and such texts have previously been

shown to elicit responses in parts of the ToM network (e.g., Jacoby & Fedorenko, 2020; Ferstl &

von Cramon, 2002). In the case of language, even though binary classification is relatively

straightforward (linguistic input is either present or not), one might argue that the language

network may nevertheless be recruited to some extent owing to the communicative nature or rich

semantics of the non-linguistic stimuli. Although prior work suggests that non-verbal

communication does not recruit the language network (e.g., Deen et al., 2015; Pritchett et al.,

2018; Jouravlev et al., 2019), some studies have found the language network is activated during

the processing of visual event semantics (e.g., Ivanova et al., 2020). Critically for present

purposes, however, to the extent that there is 'contamination' in either direction, it should work

against finding a language-ToM dissociation; i.e., our results might underestimate the true

degree of dissociation. The rich and graded nature of the stimuli might help account for some

results inconsistent with a *complete* dissociation, and we return to this question in the Discussion.

Condition	Duration	N	
+Lang +ToM			500
Story	5m 16s	31	501
Audio play	6m 14s	29	502
Dialog	5m 35s	28	
-Lang +ToM			503
Animated short film	5m 48s	32	504
Live action movie clip	6m 10s	30	
Intentional shapes animation	4m 50s	30	505
+Lang -ToM			506
Expository text	7m 6s	28	
-Lang -ToM (control)			507
Resting state	5m 0s	10	508

Table 1. *Naturalistic conditions in each of the three condition types of interest and a reality-check condition (resting state), including durations, and number of participants per condition. Durations include 16 s fixations at the beginning and end of the scan (32 s total).*

2.5. Data acquisition and preprocessing.

Data acquisition. Whole-brain structural and functional data were collected on a whole-body 3

Tesla Siemens Trio scanner with a 32-channel head coil at the Athinoula A. Martinos Imaging

Center at the McGovern Institute for Brain Research at MIT. T1-weighted structural images were

collected in 176 axial slices with 1 mm isotropic voxels [repetition time (TR) = 2530 ms; echo

time (TE) = 3.48 ms]. Functional, blood oxygenation level-dependent (BOLD) data were

acquired using an EPI sequence with a 90° flip angle and using GRAPPA with an acceleration

factor of 2; the following parameters were used: thirty-one 4-mm-thick near-axial slices acquired

in an interleaved order (with 10% distance factor), with an in-plane resolution of 2.1 x 2.1 mm,

FoV in the phase encoding (A >> P) direction 200 mm and matrix size 96 x 96 mm, TR = 2000

ms and TE = 30 ms. The first 10 s of each run were excluded to allow for steady-state

magnetization.

Spatial preprocessing. Data preprocessing was performed with SPM12 (using default

parameters, unless specified otherwise) and supporting, custom scripts in MATLAB.

Preprocessing of anatomical data included normalization into a common space [Montreal

Neurological Institute (MNI) template], resampling into 2 mm isotropic voxels, and

segmentation into probabilistic maps of the gray matter, white matter (WM), and cerebro-spinal

fluid (CSF). Preprocessing of functional data included motion correction (realignment to the

mean image using second-degree b-spline interpolation), normalization (estimated for the mean

image using trilinear interpolation), resampling into 2 mm isotropic voxels, smoothing with a 4

mm FWHM Gaussian filter and high-pass filtering at 200 s.

Temporal preprocessing. Additional preprocessing of data from the story comprehension runs

was performed using the CONN toolbox (Whitfield-Gabrieli and Nieto-Castañón, 2012;

http://www.nitrc.org/projects/conn) with default parameters, unless specified otherwise. Five

temporal principal components of the BOLD signal time courses extracted from the WM were

regressed out of each voxel's time course; signal originating in the CSF was similarly regressed

out. Six principal components of the six motion parameters estimated during offline motion

correction were also regressed out, as well as their first time derivative. Next, the residual signal

was bandpass filtered ($0.01 – 0.1$ Hz) to preserve only low-frequency signal fluctuations (Cordes

et al., 2001). This filtering did not influence the results reported below.

2.6. Participant-specific functional localization of the language and ToM (and MD, for a

replication analysis) networks.

Modeling localizer data. For each localizer task, a standard mass univariate analysis was

performed in SPM12 whereby a general linear model estimated the effect size of each condition

in each experimental run. These effects were each modeled with a boxcar function (representing

entire blocks) convolved with the canonical hemodynamic response function. The model also

included first-order temporal derivatives of these effects, as well as nuisance regressors

representing entire experimental runs, offline-estimated motion parameters, and timepoints

classified as outliers (i.e. where the scan-to-scan differences in global BOLD signal are above 5

standard deviations, or where the scan-to-scan motion is above 0.9mm). The obtained weights

were then used to compute the functional contrast of interest: for the language localizer,

sentences > nonwords, for the ToM localizer false belief > false photo, and for the MD localizer

(replication analysis; see **Section 2.4**), hard > easy spatial working memory.

Defining fROIs. Language and ToM (and MD, in the replication analysis) functional regions-of-

interest (fROIs) were defined individually for each participant based on functional contrast maps

from the localizer experiments (a toolbox for this procedure is available online;

https://evlab.mit.edu/funcloc/). These maps were first restricted to include only gray matter

voxels by excluding voxels that were more likely to belong to either the WM or the CSF based

on SPM's probabilistic segmentation of the participant's structural data.

Then, fROIs in the language network were defined using group-constrained, participant-specific

localization (Fedorenko et al., 2010). For each participant, the map of the sentences > nonwords

contrast was intersected with binary masks that constrained the participant-specific language

network to fall within areas where activations for this contrast are relatively likely across the

population. These masks are based on a group-level representation of the contrast obtained from

a previous sample of 220 participants. We used five such masks in the left-hemisphere, including

regions in the mid-to-posterior and anterior temporal lobe, as well as in the middle frontal gyrus,

the inferior frontal gyrus, and its orbital part (**Figure 4**). A version of these masks is available

online (https://evlab.mit.edu/funcloc/). In each of the resulting 5 masks, a participant-specific

language fROI was defined as the top 10% of voxels with the highest contrast values. This top

n% approach ensures that fROIs can be defined in every participant and that their sizes are the

same across participants, allowing for generalizable results (Nieto-Castañon and Fedorenko,

2012).

For the ToM fROIs, we used masks derived from a group-level representation for the false belief

> false physical contrast in an independent group of 462 participants (Dufour et al., 2013). These

masks included regions in the left and right temporo-parietal junction (L/RTPJ), precuneus /

posterior cingulate cortex (L/RPC), and dorsal medial prefrontal cortex (**Figure 4**). A version of

these masks is available online (http://saxelab.mit.edu/use-our-theory-mind-group-maps), but the

masks were edited as follows: the RSTS (right superior temporal sulcus) mask was excluded, as

it covers the entire STS, which is known to show complex functional organization, with reduced

ToM selectivity (Deen et al., 2015). The middle- and ventral-medial prefrontal cortex (MMPFC

and VMPFC) masks were also excluded to reduce the number of statistical comparisons in per-

fROI analyses, but the dorsal MPFC and PC masks were split into left- and right-hemispheres,

for a total of 6 masks.

Additionally, for the replication analysis, fROIs in the MD network were defined based on the

hard > easy contrast in the spatial working memory task. Here, instead of using binary masks

based on group-level functional data, we used anatomical masks (Tzourio-Mazoyer et al., 2002;

Fedorenko et al., 2013; Blank et al., 2014; Blank & Fedorenko, 2017). Nine masks were used in

each hemisphere, including regions in the middle frontal gyrus and its orbital part, the opercular

part of the inferior frontal gyrus, the precentral gyrus, the superior and inferior parts of the

parietal lobe, the insula, the supplementary motor area, and the cingulate cortex (**Supplemental**

Figure A2). (We note that functional masks derived for the MD network based on 197

participants were largely overlapping with the anatomical masks; we chose to use the anatomical

masks to maintain comparability between our functional data and data from previous studies that

have used these masks.)

In line with prior studies (e.g., Blank, Fedorenko & Kanwisher, 2014; Blank & Fedorenko, 2017;

Paunov, Blank, & Fedorenko, 2019), the resulting fROIs showed small pairwise overlaps within

individuals across networks, and overlapping voxels were excluded in fROI definition. In the

current sample, the language-MD and ToM-MD overlaps were negligible, with a median overlap

of 0 and average percentage overlap of fewer than 3% of voxels, on average across participants,

relative to the total size of all fROIs in either network. Similarly, the language-ToM overlaps

were small relative to all fROIs in either network (6.3% of voxels, on average across

participants, relative to the total number of voxels in language fROIs and 3.2% relative to all

ToM fROIs). This overlap was localized entirely to one pair of fROIs: the LPostTemp language

fROI and the LTPJ ToM fROI, and was more substantial relative to the total sizes of just these

two fROIs: 38.6 voxels, on average across participants, i.e., 13.1% out of 295 total LPostTemp

voxels, and 11.6% out of 332 total LTPJ voxels. We therefore repeated all key analyses without

excluding these voxels in defining the fROIs. The results of these alternative analyses were

qualitatively and statistically similar.

2.7. Reality check and replication analyses.

Prior to performing our critical analyses, we conducted two reality-check analyses and—in line

with increasing emphasis in the field on robust and replicable science (e.g., Poldrack et al.,

2017)—an analysis aimed at replicating and extending a previous ISC-based finding from our

lab (Blank & Fedorenko, 2017).

ISCs in perceptual cortices. Anatomical ROIs were additionally defined in early visual and

auditory cortex in all participants. For visual cortex, regions included inferior, middle, and

superior occipital cortex bilaterally (six ROIs in total; masks available from

http://fmri.wfubmc.edu/software/PickAtlas). For auditory cortex, regions included posteromedial

and anterolateral sections of Heschl's gyrus bilaterally (4 ROIs in total; Morosan et al., 2001;

these regions are based on post-mortem histology, and have been used in a number of previous

fMRI papers). Signal extraction, ISC estimation, and inferential statistics were performed

identically to the critical analyses (see *Section 2.8*). ISCs from these regions were used in a

reality check (see *Section 3.1*), to ensure a double-dissociation obtains between visually and

auditorily presented conditions in these perceptual regions.

Resting state ISCs. A subset of ten participants (who completed 1-7 of the critical conditions)

completed a resting state scan, which was included to ensure that data acquisition, preprocessing,

and modelling procedures do not induce spurious ISCs. To this end, signal extraction, ISC

estimation, and inferential statistics were performed identically to the critical analyses (see

Section 2.8).

Replication analysis: Closer tracking of linguistic input by language regions than by domain-

general Multiple Demand (MD) regions. Blank & Fedorenko (2017) reported stronger ISCs

during the processing of naturalistic linguistic materials in the language regions, compared to

domain-general Multiple Demand (MD) regions. The MD network has been implicated in

executive processes and goal-directed behavior (e.g., Duncan, 2010, 2013), including in the

domain of language (e.g., see Fedorenko, 2014, for a review; cf. Diachek, Blank, Siegelman et

al., 2020; Fedorenko & Shain, under review). We sought to replicate Blank & Fedorenko's key

result in a new set of participants and to extend it to different types of linguistic materials.

Specifically, the original study used narratives, including the narrative used in the present study

along with three others. We expected the results to generalize to non-narrative linguistic

conditions. Ten participants in our dataset (n=6 in the narrative condition) who also participated

in the original study were excluded from this analysis. Again, signal extraction, ISC estimation,

and inferential statistics were identical to the critical analyses (see *Section 2.8*).

2.8. Critical analyses.

Computing ISCs. For each participant and fROI, BOLD signal time courses recorded during each

naturalistic condition were extracted from each voxel beginning 6 s following the onset of the

stimulus (to exclude an initial rise in the hemodynamic response relative to fixation, which could

increase ISCs) and averaged across voxels. For each fROI, participant, and condition we

computed an ISC value, namely, Pearson's moment correlation coefficient between the z-scored

time course and the corresponding z-scored and averaged time course across the remaining

participants (Lerner et al., 2011). ISCs were Fisher-transformed before statistical testing to

improve normality (Silver and Dunlap, 1987).

Statistical testing. In each fROI, ISCs were then tested for significance against an empirical null

distribution based on 1,000 simulated signal time courses that were generated by phase-

randomization of the original data (Theiler et al., 1992). Namely, we generated null distributions

for individual participants, fit each distribution with a Gaussian, and analytically combined the

resulting parameters across participants. The true ISCs, also averaged across participants, were

then z-scored relative to these empirical parameters and converted to one-tailed *p*-values.

ISCs were compared across networks and condition types using linear, mixed-effects

regressions, implemented in Matlab 2020a. ISCs were modeled with maximal random effects

structure appropriate for each analysis (Barr et al., 2013; Baayen et al., 2008), including random

intercepts for participants, with random slopes for the effects of interest, and crossed random

intercepts for fROI and condition. Hypothesis testing was performed with two-tailed tests over

the respective model coefficients, with Satterthwaite approximation for the degrees of freedom.

Further analyses performed within networks across condition types or against the theoretical null

distribution (i.e., testing the intercept term), as well as those per fROI within a network or per

condition across networks, also always included maximal random effects on the remaining

grouping variables. P-values in these analyses are reported following false discovery rate (FDR)

correction for multiple comparisons (Benjamini and Yekutieli, 2001).

Lastly, in comparisons against baseline, per-fROI analyses against empirical null distributions

are also reported, which aim to ensure that, at the finest grain (each individual fROI and

condition, across participants), differences from baseline are independent of assumptions

regarding data normality. These tests were also FDR corrected for multiple comparisons for all

fROIs within a network, and across all seven conditions of interest.

3. Results

3.1. Results of reality check and replication analyses.

Reality check #1: ISCs in perceptual cortices. We examined ISCs for the conditions of interest in

early auditory and visual cortex, grouping the

conditions by presentation modality. As

expected, we observed stronger ISCs for the

auditory conditions in the auditory cortex, and

stronger ISCs for the visual conditions in the

visual cortex (**Figure 3**). The linear mixed-

effects (LME) regression (see *Section 2.2*)

revealed a strong crossover interaction (beta =

0.648, SE = 0.046, $t(89.79) = 13.970$, $p = 10^{-25}$).

Figure 3. *A reality-check analysis showing the expected double dissociation in ISCs in*

perceptual (visual and auditory) cortices. Bars correspond to Fisher-transformed inter-subject

correlation (ISC) coefficients (Pearson's r) in early visual and auditory cortex to the conditions

of interest, grouped by modality of presentation (all +Language conditions: story, audio play,

dialog, expository text were auditorily presented; the remaining conditions – animated short

film, live action movie clip, and intentional shapes animation – were visually presented). Error

bars are standard errors of the mean by participants. Black dots correspond to the individual

participants' values. Large unfilled circles correspond to individual condition averages. Vertical

curves are Gaussian fits to empirical null distributions.

Notably, we also observed that the visual areas weakly but reliably tracked the auditory

conditions (beta = 0.066, SE = 0.012, $t(38.10) = 5.398$, $p = 10^{-7}$), and the early auditory areas

reliably tracked the visual conditions (beta = 0.175, SE = 0.020, $t(24.15) = 8.573$, $p = 10^{-10}$). We

return to the interpretation of these effects in the Discussion. For the time being, we note that

care must be taken in interpreting deviations of the ISCs during 'active' (cf. resting state)

conditions from baseline.

Reality check #2: Resting state ISCs. To exclude the possibility that the ISCs in the critical

analyses are driven by scanner noise or preprocessing/analysis procedures, we measured ISCs

across a subset of ten participants who were scanned in a 5-minute resting state condition

(Hasson et al., 2004). The ISCs during rest did not significantly differ from baseline in either the

language or ToM networks, as assessed with an LME regression, or against the empirical null

distribution. This analysis suggests that any above-baseline ISCs for our critical conditions are

not an artifact of data acquisition, preprocessing, or analysis procedures.

Replication analysis: Closer tracking of linguistic input by language regions than by domain-

general Multiple Demand (MD) regions. We successfully replicated the key finding for the

narrative condition (beta = 0.181, SE = 0.039, $t(34.89) = 4.664$, $p = 10^{-5}$; p values are FDR-

corrected for the four linguistic conditions), and extended it to the audio play (beta = 0.307, SE =

0.052, $t(34.23) = 5.944$, $p = 10^{-7}$), the dialog (beta = 0.157, SE = 0.039, $t(33.05) = 4.014$, $p =$

10^{-5}), and the expository text (beta = 0.129, SE = 0.036, $t(36.75) = 3.543$, $p = 10^{-4}$) conditions

(**Supplemental Figure A3**). These results suggest that across diverse kinds of linguistic stimuli,

the language network's activity is more tightly coupled with the inputs, compared to the domain-

general MD regions' activity.

3.2. Results of critical analyses.

Evidence for dissociation between the language and ToM networks in direct network

comparisons.

We tested three key predictions, which, if supported, provide evidence in favor of language-ToM

dissociability in naturalistic settings. First, the ToM network should track conditions rich in ToM

content irrespective of whether these conditions are linguistic or non-linguistic, whereas the

language network should track linguistic conditions more strongly than non-linguistic ones.

Indeed, we found a network (language, ToM) x condition type (*linguistic*: narrative, audio play,

dialog, expository text; *non-linguistic*: animated film, live action film clip, Heider & Simmel-

style animation) interaction (beta = 0.191, SE = 0.059, $t(84.06) = 3.270$, $p = 0.002$). We also

found main effects of condition type and network: the linguistic conditions were—on average

across networks—tracked more strongly than the non-linguistic conditions (beta = 0.124, SE =

0.0459, $t(96.47) = 2.700$, $p = 0.008$), and the language network tracked the conditions more

strongly, on average, than the ToM network (beta = 0.096, SE = 0.039, $t(86.87) = 2.487$, $p =$

0.015).

Second, for non-linguistic conditions, the ToM network should exhibit stronger tracking of

conditions with mental state content than the language network. Indeed, the ToM network

showed higher ISCs than the language network (beta = 0.095, SE = 0.043, $t(35.59) = 2.197$, $p =$

0.035).

And third, for the linguistic condition without mental state content (the expository text), the

language network should exhibit stronger tracking than the ToM network. Indeed, the language

network showed higher ISCs than the ToM network (beta = 0.111, SE = 0.044, $t(19.10) = 2.507$,

$p = 0.021$).

The same qualitative pattern obtains when the non-verbal ToM localizer is used to define the

ToM fROIs (**Supplemental Figure C1**).

Figure 4. Left. *Masks within which individual functional regions of interest (fROIs) were defined*

for each network: Language (Top, red): IFGorb, inferior frontal gyrus, orbital portion; IFG,

inferior frontal gyrus; MFG, middle frontal gyrus; AntTemp, anterior temporal cortex;

PostTemp, posterior temporal cortex (only the classic, left-hemisphere language regions were

included in all analyses). ToM (Bottom, green): TPJ, temporoparietal junction; DMPFC,

dorsomedial prefrontal cortex; PC, posterior cingulate cortex and precuneus. Both the right-

hemisphere (shown) and left-hemisphere ToM regions were included (six regions total). ***Middle.***

Average inter-subject correlations (ISCs) per condition type in the language (top) and ToM

(bottom) networks. Bars correspond to Fisher-transformed ISC coefficients (Pearson's r),

averaged across regions of interest within each network, separately per condition. Colors

represent condition types: +Lang+ToM, orange, -Lang+ToM, green, +Lang-ToM, red. Error

bars are standard errors of the mean by participants. Black dots correspond to the individual

participants' values. Large unfilled circles correspond to individual condition averages, shown

individually in the right-most panels. Vertical curves are Gaussian fits to empirical null

distributions. The key pattern is as follows. The ToM network tracks +ToM materials in both

+Linguistic and -Linguistic conditions, but shows weak tracking of the -ToM stimulus. The

language network preferentially tracks +Linguistic materials over -Linguistic ones, and it tracks

+Linguistic materials in both +ToM and -ToM conditions. **Right.** *ISCs per individual*

naturalistic stimuli ('conditions'); conventions are as in middle panels.

A more detailed characterization of the two networks' ISC profiles.

In this section, we examine more closely the detailed pattern of ISCs in the two networks of

interest. The first aim of these analyses is to establish that the observed dissociation is not driven

by particular conditions or regions within the networks, but rather, that different aspects of the

data provide convergent support for the dissociation. The second aim is to highlight aspects of

the ISC pattern that are not consistent with a complete language-ToM dissociation, and thus to

evaluate the strength of counter-evidence in favor of the null hypothesis, that language and ToM

are not dissociable in naturalistic cognition (see **Supplemental Figure B1** for ISCs per fROI for

each network and condition).

ToM network. First, the ToM network reliably tracked each of the six conditions with mental

state content: story (beta = 0.363, SE = 0.037, $t(26.55) = 9.693$, $p = 10^{-10}$); p values are FDR-

corrected for seven conditions—we are including all conditions in the correction, not only the

+ToM conditions), audio play (beta = 0.295, SE = 0.028, $t(10.54) = 10.716$, $p = 10^{-7}$), dialog

(beta = 0.203, SE = 0.026, $t(16.40) = 7.797$, $p = 10^{-7}$), animated short (beta = 0.472, SE = 0.053,

$t(9.35) = 8.830$, $p = 10^{-6}$), live action film (beta = 0.241, SE = 0.035, $t(11.18) = 6.869$, $p = 10^{-6}$),

and Heider & Simmel (H&S) style animation (beta = 0.205, SE = 0.026, $t(15.72) = 7.838$, $p =$

10^{-7}). Moreover, in tests against the empirical null distributions these effects were significant in

every ToM fROI with the exception of the Dialog condition in the RH Posterior

Cingulate/Precuneus, (all other $ps < 0.04$, FDR-corrected for the six fROIs and seven

conditions).

Second, the ToM network showed no preference for linguistic *vs.* non-linguistic conditions with

mental state content (beta = 0.018, SE = 0.043, $t(49.74) = 0.675$, *n.s.*), consistent with these

regions' role in representing mental states irrespective of how this information is conveyed (e.g.,

Jacoby et al., 2016).

And third, the ToM network tracked the linguistic conditions with mental state content (story,

audio play, dialog) more strongly than the one without mental state content (expository text)

(beta = 0.199, SE = 0.037, $t(30.69) = 5.344$, $p = 10^{-7}$), suggesting that the network represents

mental state information in linguistic signals, rather than the linguistic signal itself. However, the

ToM network did exhibit weaker but significantly above-baseline tracking of the expository text

(beta = 0.093, $SE = 0.022$, $t(22.22) = 4.165$, $p = 0.001$). In per-fROI tests against the empirical

null distributions, this effect was only reliable in the LTPJ ($p = 0.015$; all other $ps > 0.05$; FDR-

corrected for the six fROIs). We consider possible explanations in the Discussion.

Language network. First, the language network reliably tracked each of the three linguistic

conditions with mental state content: story (beta = 0.374, SE = 0.041, $t(13.27) = 9.171$, $p = 10^{-7}$;

p values are FDR-corrected for seven conditions), audio play (beta = 0.479, *SE* = 0.046, $t(8.75) =$

10.370, $p = 10^{-6}$), and dialog ($r = 0.294$, *SE* = 0.045, $t(9.72) = 6.491$, $p = 10^{-4}$). Moreover, these

effects were significant in every language fROI, in tests against the empirical null distributions

(ps < 0.01, FDR-corrected for the five fROIs and seven conditions).

Second, importantly, the language network also reliably tracked the linguistic condition with no

mental state content (beta = 0.204, *SE* = 0.044, $t(8.72) = 4.668$, $p = 10^{-4}$), and this effect, too, was

significant in every language fROI ($ps < 0.03$, FDR-corrected for the five fROIs). This result

suggests that mental state content is not necessary to elicit reliable ISCs in the language network.

And third, the language network showed stronger tracking of linguistic relative to non-linguistic

conditions with mental state content, (beta = 0.177, SE = 0.043, $t(43.54) = 4.155$, $p = 10^{-5}$). This

result suggests a special role for linguistic input in driving the network's responses.

However, the language network exhibited some patterns that might be taken to suggest that

mental state content—or social information more generally—is, to some extent, important for

linguistic processing. First, the language network tracked the linguistic conditions with mental

state content more strongly than the linguistic condition with no mental state content, i.e., the

expository text (beta = 0.188, SE = 0.061, $t(25.67) = 3.050$, $p = 0.005$). This result may be taken

to suggest that mental state content contributes to the language network's input tracking over and

above the linguistic content alone. This interpretation warrants caution, however. In particular,

reflecting the general challenges of naturalistic stimuli (see Discussion), the linguistic condition

with no mental state content is not matched to the linguistic conditions with mental state content

on various potentially relevant features, from how engaging they are, which could influence the

depth of linguistic encoding, to specifically linguistic properties (e.g., lexical and syntactic

complexity), which could also affect the strength of ISCs (e.g., Shain, Blank et al., 2020; Wehbe

et al., 2021). Furthermore, only a single linguistic condition with no mental state content was

included in the current study, making it difficult to rule out idiosyncratic features driving the

difference.

And second, the above-baseline ISCs in the language network for the non-linguistic conditions—

although weaker than those for the linguistic conditions—are also notable, suggesting some

degree of reliable tracking in the language network for non-linguistic meaningful information

(see also Ivanova et al., 2020 for evidence of reliable responses in the language network to visual

events).

4. Discussion

Much prior work in cognitive neuroscience has suggested—based on traditional controlled

experimental paradigms—that the network of brain regions that support linguistic interpretation

and the brain regions that support mental state reasoning are distinct (e.g., Saxe & Kanwisher,

2003; Mason & Just, 2009; Fedorenko et al., 2011; Mar, 2011; Deen et al., 2015; Paunov, 2019).

However, such paradigms differ drastically from real-world cognition, where we process rich

and complex information. And linguistic and social cognition seem to be strongly intertwined in

everyday life. Here, we tested whether the language and Theory of Mind networks are

dissociated in their functional profiles as assessed using the inter-subject correlation (ISC)

approach, where neural activity patterns are correlated across individuals during the processing

of naturalistic materials (e.g., Hasson et al., 2004, 2008). Following Blank & Fedorenko (2017),

we combined the ISC approach with the power of individual-participant functional localization

(e.g., Brett et al., 2002; Saxe et al., 2006; Fedorenko et al., 2010; Nieto-Castañón & Fedorenko,

2012). This synergistic combination has two key advantages over the whole-brain voxel-wise

ISC approach, where individual brains are first anatomically aligned and, then, each stereotaxic

location serves as a basis for comparing signal time courses across participants. First, relating the

resulting cortical topography of ISCs to the topography of known functional brain networks can

only proceed only through "reverse inference" based on anatomy (Poldrack, 2006; Fedorenko,

2021). Instead, evaluating signal time courses from functionally defined regions ensures

interpretability, and allows us to straightforwardly link our findings to the wealth of prior studies

characterizing the response profiles of our two networks of interest. And second, this approach

allows us to directly test the correlations in the language network against those in the ToM

network. Such an explicit comparison between networks allows for stronger inferences compared

with those licensed when each network is separately tested against a null baseline and differences

across networks are indirectly inferred (e.g., see Nieuwenhuis et al., 2011, for discussion).

We examined the ISCs in the language and ToM networks during the processing of seven

naturalistic conditions: three linguistic conditions with mental state content (+linguistic, +ToM),

three non-linguistic conditions (silent animations and live action films) with social content but no

language (-linguistic, +ToM), and a linguistic condition with no social content (+linguistic, -

ToM). We found reliable differences in the ISC patterns between the language and ToM

networks, in support of the hypothesis that language and Theory of Mind are dissociable even

during the processing of rich and complex naturalistic materials. In particular, the ToM network

tracked materials rich in mental state content irrespective of whether this content was presented

linguistically or non-linguistically (see also Jacoby et al., 2016), but it showed only weak

tracking of the stimulus with no mental state content. In contrast, the language network

preferentially tracked linguistic materials over non-linguistic ones, and it did so regardless of

whether these materials contained information about mental states.

These results expand on the existing body of knowledge about language and social cognition,

with both theoretical and methodological implications. Critically, the observed dissociation

extends prior findings of dissociable functional profiles between the language and the ToM

networks during task-based paradigms to rich naturalistic conditions. This result suggests that the

two networks represent *different kinds of information*. (They may also perform *distinct*

computations on the perceptual inputs, though the idea of a canonical computation carried out

across the cortex is gaining ground (e.g., Keller & Mrsic-Flogel, 2018; Fedorenko & Shain, in

press), and predictive processing seems like one likely candidate (e.g., Koster-Hale & Saxe,

2013; Shain, Blank et al., 2020).) In particular, the language regions appear to track linguistic

features of the input (see also Shain, Blank et al., 2020; Shain et al., 2021; Wehbe et al., 2021).

Our results extend prior findings from ISC paradigms (Wilson et al., 2008; Lerner et al., 2011;

Honey et al., 2012; Regev et al., 2013; Silbert et al., 2014; Schma □lzle et al., 2015; Blank &

Fedorenko, 2017), which all used materials rich in mental state content, as is typical of linguistic

information, to a stimulus that is largely devoid of information about mental states—an

expository text. Strong tracking of the latter stimulus aligns with prior findings from task-based

paradigms of robust responses to linguistic materials with little or no mental state content (e.g.,

Deen et al., 2015; Jacoby & Fedorenko, 2020).

The ToM regions, in contrast, appear to track some features related to representing mental states

across diverse kinds of representations (linguistic materials, animations, including highly abstract

and minimalistic ones, and live action movies), again aligning with prior findings from task-

based paradigms (e.g., Fletcher et al., 1995; Gallagher et al., 2000; Castelli et al., 2000; Vogeley

et al., 2001; Ruby & Decety, 2003; Saxe & Kanwisher, 2003; Saxe, Schulz, & Jiang, 2006;

Jacoby et al., 2016). It is important to keep in mind that the fact that the two networks are

dissociable does not imply that they do not interact. Indeed, Paunov et al. (2019) reported

reliably above chance correlations in the patterns of inter-regional synchronization between the

language and ToM networks, suggesting some degree of functional integration.

On the methodological level, these results vindicate the divide-and-conquer strategy in general,

and the functional localization approach (e.g., Brett et al., 2002; Saxe et al., 2006; Fedorenko et

al., 2010) in particular. Language and Theory of Mind appear to be distinct, supported by

dissociable cortical networks for the processing of linguistic vs. mental state information, at least

in adulthood (see also Braga et al., 2020). It is therefore justifiable to study each cognitive

faculty and each network separately, although further probing the mechanisms of their potential

interactions is equally important.

Although the overall pattern clearly supports a language-ToM dissociation between the two

networks, some aspects of the results are not in line with a complete dissociation. In particular, i)

the language regions show reliable tracking of non-linguistic conditions with mental state

content; ii) the language regions show stronger tracking of linguistic conditions with than

without mental state content; and iii) the ToM regions show weak but reliable tracking of the

linguistic condition with no mental state content. These findings may be due to methodological

limitations: the above-baseline ISCs (especially the relatively weak ones) may not reflect

stimulus tracking. Although we have ruled out the possibility that the above-baseline ISCs are

driven by acquisition, preprocessing, or analysis artifacts in our reality-check analysis of resting

state data, they could be driven by other, non-mutually-exclusive, factors. One possibility is that

inter-network interactions could induce ISCs. In particular, given that the language and the ToM

network show some degree of synchronization in activity during naturalistic cognition (Paunov

et al., 2019), the ToM network's tracking of a linguistic stimulus with no mental state content,

for example, may be due do the fact that the language system is tracking this stimulus, and there

is some 'leakage' of this tracking to the ToM network through inter-network synchronization.

Similarly, the language network's preference of linguistic stimuli with mental state content over

those without mental state content may be due to the leakage from the ToM network.

Another possibility is that the incomplete dissociation between language and Theory of Mind (at

least with respect to (i) and (iii) above) may be at least partly attributable to pragmatic

processing, arguably present across all 'conditions': the non-linguistic ToM conditions are still

'story-like' and hence communicative and the linguistic non-ToM stimulus arguably still

requires attribution of communicative intentions, as discussed in the introduction. The division of

labor between the language and ToM networks in pragmatic inference is an exciting future

direction. Demonstrating that these two networks are, in the first place, dissociable, even during

rich naturalistic cognition—the goal of the present study—is an important step to pursuing this

line of research. With this groundwork in place, neuroimaging evidence can be increasingly

brought to bear on the question of which aspects of pragmatics require mental state reasoning, as

evidenced by the engagement of the ToM network. Given the broad scope of pragmatics,

encompassing diverse heterogeneous phenomena, an empirically motivated pragmatic taxonomy

may be developed by investigating whether some classes of pragmatic inference are resolved

within the language network proper (e.g., 'lower-level' inferences about lexical or syntactic

ambiguity) whereas others (e.g., establishing discourse coherence, or understanding irony)

require the ToM network.

Finally, the stimuli themselves may be too confounded to fully dissociate the respective

contributions of language- and ToM-related components. For example, in linguistic stimuli,

mental state attribution often requires particular syntactic structures—sentential complements.

More generally, the use of naturalistic materials, despite its advantages (e.g., Hasson et al., 2018;

Sonsukare et al., 2019; Nastase et al., 2020), is associated with a host of challenges. The key one,

mentioned above, is that certain features are necessarily confounded in naturalistic settings, and

can only be dissociated through careful experimentation and altering the natural statistics of the

input. Relying on naturalistic materials alone can lead to wrong conclusions about the cognitive

and neural architecture. This problem is especially pronounced in studying the relationship

between language and social cognition given that language is primarily used in social settings

and to share socially relevant information. The use of linguistic materials with no social

information, and of non-linguistic mental-state-rich materials has been critical, here and in earlier

studies, to uncover the dissociation that holds between the language and ToM systems. Another

challenge associated with the use of naturalistic materials is that they are difficult, or altogether

impossible, to match for diverse properties bound to affect neural responses. Again, this problem

presents a particular challenge in comparing responses to materials rich in social information vs.

devoid of such information given that the former are, almost by definition, going to be more

engaging and exciting given the social nature of primates, including humans (e.g., Aronson,

1980; Cheney & Seyfarth, 1990; Tomasello, 2014). Possible ways to address these concerns

could involve a) characterizing the natural statistics of co-occurrences between linguistic and

social processing in order to better understand how well the naturalistic stimuli reflect those

statistics and perhaps altering naturalistic conditions to allow dissociating features that

commonly co-occur in life; b) developing novel neural analysis methods to isolate the

components of neural signals attributable to a particular cognitive process / brain network (e.g.,

using analytic methods well-suited to high-dimensional data such as independent components

analysis across both cortical space and large feature spaces representative of naturalistic

environments (e.g., Norman-Haignere et al., 2015), or across both cortical space and time (e.g.,

probabilistic ICA, PICA; see Beckmann et al., 2005 for overview); and c) carefully annotating

naturalistic materials and performing reverse correlation analyses (e.g., Hasson et al., 2004;

Richardson et al., 2018) in an effort to understand the precise features that elicit increases in

neural responses in different brain regions. The latter approach may be particularly informative

with respect to the question of possible *gradation* of demands on language and ToM processing

both within naturalistic stimuli and across 'conditions' that we here grouped in the same 'type'

or general categories of +/-ToM and +/-Language. Our results seem to suggest that considerable

within-category heterogeneity exists in the degree of stimulus tracking (e.g., in the language

network, the dialog is tracked less strongly than the narrative, and in the ToM network, the

Heider & Simmel-style animation is tracked less strongly than the animated film. Our dataset is

not ideally suited for such investigation because it includes only a single instance of each

'condition', and no attempt was made to match the conditions on at least *some* dimensions that

may improve homogeneity, but this is a promising direction for future work.

1017

A few other exciting future directions are worth highlighting, some of which may build directly

on the approaches introduced and the findings reported in the current study. First, the language

and ToM networks appear to be dissociable in the adult mind and brain. However, it is

possible—perhaps even plausible—that this dissociation emerges over the course of

development. Prior neuroimaging work has shown that the ToM network becomes gradually

more specialized for mental state attribution (e.g., Saxe, 2009; Gweon et al., 2012; Richardson et

al., 2018), and this specialization appears to be protracted with delayed language acquisition

(Richardson et al., 2020). Very little is known about how specialization for linguistic processing

(e.g., Fedorenko et al., 2011; Monti et al., 2012) emerges. Perhaps early on in development, a set

of lateral frontal, lateral temporal, and midline cortical areas are tuned to *any* socially relevant

information, and these areas later fractionate into those specialized for processing linguistic

signals vs. those for mental state attribution vs. those that support many other kinds of social

signals, both visual (e.g., eye gaze, facial expressions, gestures) and auditory (e.g., non-verbal

vocalizations, prosodic information, speech acoustics). This fractionation is likely driven by

computational and metabolic advantages of localized processing (e.g., Barlow, 1995; Foldiak &

Young, 1995; Chklovskii & Koulakov, 2004; Olshausen & Field, 2004; see Kanwisher, 2010, for

discussion). Probing linguistic and social cognition across the lifespan will be critical to

understand how the two networks form and develop, leading to the segregation we observe in the

adult brain.

1037

Second, as noted above, given the likely frequent interactions between the language and ToM

networks (including their most strongly dissociated components), searching for possible

mechanisms of those interactions (e.g., Paunov et al., 2019) seems critical. This would require a

combination of studies characterizing the patterns of anatomical connections for the language

and ToM regions (e.g., Saur et al., 2010; Wiesmann et al., 2017) and studies probing online

interactions using methods with high temporal resolution, like MEG or intracranial recordings.

1044

Third, we are still a long way away from a mechanistic-level understanding of what the language

or the ToM regions do. The use of naturalistic stimuli, including in the context of the ISC

approach, is promising. In particular, by examining the points in the stimulus where most

participants show increases in neural activity can help generate (and subsequently test) specific

hypotheses about the necessary and sufficient features of the input that are required to elicit

neural responses in the relevant brain regions.

1051

1052   To conclude, we have demonstrated that the dissociation between the language and ToM

1053   networks that has been previously reported based on traditional task paradigms, robustly

1054   generalizes to rich naturalistic conditions. However, the precise nature of each network's

1055   representations and computations, the emergence of these networks in development, and the

1056   mechanisms for information sharing between them remain to be discovered.

1057

1058

1059

1060

1061 **References**

1062

Amalric, M., & Dehaene, S. (2018). Cortical circuits for mathematical knowledge: evidence for a

major subdivision within the brain's semantic networks. *Philosophical Transactions of the*

Royal Society B: Biological Sciences, *373*(1740), 20160515.

Anderson, A. J., Lalor, E. C., Lin, F., Binder, J. R., Fernandino, L., Humphries, C. J., ... & Wang,

X. (2019). Multiple regions of a cortical network commonly encode the meaning of words in

multiple grammatical positions of read sentences. *Cerebral cortex, 29*(6), 2396-2411.

Apperly, I. A., Samson, D., Carroll, N., Hussain, S., & Humphreys, G. (2006). Intact first-and

second-order false belief reasoning in a patient with severely impaired grammar. *Social*

neuroscience, 1(3-4), 334-348.

Aronson E. The social animal. New York: Palgrave Macmillan; 1980.

Ayyash, D., Malik-Moraleda, S., Gallée, J., Affourtit, J., Mineroff, Z., Jouravlev, O., &

Fedorenko, E. (in prep). The universal language network: A cross-linguistic investigation

spanning 45 languages and 11 language families.

Baayen, R. H., Davidson, D. J., & Bates, D. M. (2008). Mixed-effects modeling with crossed

random effects for subjects and items. *Journal of memory and language, 59*(4), 390-412.

Barlow, H. (1995). The neuron doctrine in perception.

Barr, D. J., Levy, R., Scheepers, C., & Tily, H. J. (2013). Random effects structure for

confirmatory hypothesis testing: Keep it maximal. Journal of memory and language, 68(3),

255-278.

Bautista, A., & Wilson, S. M. (2016). Neural responses to grammatically and lexically degraded

speech. *Language, cognition and neuroscience*, *31*(4), 567-574.

Beckmann, C. F., DeLuca, M., Devlin, J. T., & Smith, S. M. (2005). Investigations into resting-

state connectivity using independent component analysis. *Philosophical Transactions of the*

Royal Society B: Biological Sciences, *360*(1457), 1001-1013.

Bedny, M., Pascual-Leone, A., Dodell-Feder, D., Fedorenko, E., & Saxe, R. (2011). Language

processing in the occipital cortex of congenitally blind adults. *Proceedings of the National*

Academy of Sciences, *108*(11), 4429-4434.

Benjamini, Y., & Yekutieli, D. (2001). The control of the false discovery rate in multiple testing

under dependency. *The annals of statistics*, *29*(4), 1165-1188.

Blank, I. A., & Fedorenko, E. (2017). Domain-general brain regions do not track linguistic input

as closely as language-selective regions. *Journal of Neuroscience*, *37*(41), 9999-10011.

Blank, I., Balewski, Z., Mahowald, K., & Fedorenko, E. (2016). Syntactic processing is

distributed across the language system. *Neuroimage*, *127*, 307-323.

Blank, I., Kanwisher, N., & Fedorenko, E. (2014). A functional dissociation between language

and multiple-demand systems revealed in patterns of BOLD signal fluctuations. *Journal of*

neurophysiology, *112*(5), 1105-1118.

Bosco, F. M., Tirassa, M., & Gabbatore, I. (2018). Why pragmatics and Theory of Mind do not

(completely) overlap. *Frontiers in Psychology*, *9*, 1453.

Bower, G. H., & Morrow, D. G. (1990). Mental models in narrative comprehension. *Science*,

247(4938), 44-48.

Braga, R. M., DiNicola, L. M., Becker, H. C., & Buckner, R. L. (2020). Situating the left-

lateralized language network in the broader organization of multiple specialized large-scale

distributed networks. *Journal of neurophysiology, 124*(5), 1415-1448.

Brett, M., Johnsrude, I. S., & Owen, A. M. (2002). The problem of functional localization in the

human brain. *Nature reviews neuroscience*, *3*(3), 243.

Bruneau, E. G., Pluta, A., & Saxe, R. (2012). Distinct roles of the 'shared pain' and 'theory of

mind'networks in processing others' emotional suffering. *Neuropsychologia*, *50*(2), 219-231.

Callaghan, T., Rochat, P., Lillard, A., Claux, M. L., Odden, H., Itakura, S., ... & Singh, S. (2005).

Synchrony in the onset of mental-state reasoning: Evidence from five cultures. *Psychological*

Science, *16*(5), 378-384.

Castelli, F., Happé, F., Frith, U., & Frith, C. (2000). Movement and mind: a functional imaging

study of perception and interpretation of complex intentional movement

patterns. *Neuroimage*, *12*(3), 314-325.

Chen, G., Taylor, P. A., & Cox, R. W. (2017). Is the statistic value all we should care about in

neuroimaging?. *Neuroimage*, *147*, 952-959.

Cheney, D., & Seyfarth, R. (1990). Attending to behaviour versus attending to knowledge:

Examining monkeys' attribution of mental states. *Animal Behaviour*, *40*(4), 742-753.

Chklovskii, D. B., & Koulakov, A. A. (2004). Maps in the brain: what can we learn from

them?. *Annu. Rev. Neurosci.*, *27*, 369-392.

Cordes, D., Haughton, V. M., Arfanakis, K., Carew, J. D., Turski, P. A., Moritz, C. H., ... &

Meyerand, M. E. (2001). Frequencies contributing to functional connectivity in the cerebral

cortex in "resting-state" data. *American Journal of Neuroradiology*, *22*(7), 1326-1333.

Cuevas, P., Steines, M., He, Y., Nagels, A., Culham, J., & Straube, B. (2019). The facilitative

effect of gestures on the neural processing of semantic complexity in a continuous

narrative. *NeuroImage*, *195*, 38-47.

Deen, B., Koldewyn, K., Kanwisher, N., & Saxe, R. (2015). Functional organization of social

perception and cognition in the superior temporal sulcus. *Cerebral Cortex*, bhv111.

Diachek, E., Blank, I., Siegelman, M., Affourtit, J., & Fedorenko, E. (2020). The domain-general

multiple demand (MD) network does not support core aspects of language comprehension: a

large-scale fMRI investigation. *Journal of Neuroscience, 40*(23), 4536-4550.

Downing, P. E., Peelen, M. V., Wiggett, A. J., & Tew, B. D. (2006). The role of the extrastriate

body area in action perception. Social Neuroscience, 1(1), 52-62.

Dufour, N., Redcay, E., Young, L., Mavros, P. L., Moran, J. M., Triantafyllou, C., ... & Saxe, R.

(2013). Similar brain activation during false belief tasks in a large sample of adults with and

without autism. *PloS one*, *8*(9), e75468.

Dunbar, R. I. (2004). Gossip in evolutionary perspective. *Review of general psychology*, *8*(2),

100-110.

Dunbar, R. I., Marriott, A., & Duncan, N. D. (1997). Human conversational behavior. *Human*

nature, 8(3), 231-246.

Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: mental programs for

intelligent behaviour. *Trends in cognitive sciences*, *14*(4), 172-179.

Duncan, J. (2013). The structure of cognition: attentional episodes in mind and brain. *Neuron*,

80(1), 35-50.

Fedorenko, E. (2021). The early origins and the growing popularity of the individual-subject

analytic approach in human neuroscience. *Current Opinion in Behavioral Sciences, 40*, 105-

112.

Fedorenko, E. (2014). The role of domain-general cognitive control in language comprehension.

Frontiers in psychology, *5*, 335.

Fedorenko, E., Behr, M. K., & Kanwisher, N. (2011). Functional specificity for high-level

linguistic processing in the human brain. *Proceedings of the National Academy of Sciences*,

108(39), 16428-16433.

Fedorenko, E., Blank, I. A., Siegelman, M., & Mineroff, Z. (2020). Lack of selectivity for syntax

relative to word meanings throughout the language network. *Cognition, 203*, 104348.

Fedorenko, E., Duncan, J., & Kanwisher, N. (2013). Broad domain generality in focal regions of

frontal and parietal cortex. *Proceedings of the National Academy of Sciences*, *110*(41), 16616-

16621.

Fedorenko, E., Hsieh, P. J., Nieto-Castañón, A., Whitfield-Gabrieli, S., & Kanwisher, N. (2010).

New method for fMRI investigations of language: defining ROIs functionally in individual

subjects. *Journal of neurophysiology*, *104*(2), 1177-1194.

Fedorenko, E., & Kanwisher, N. (2009). Neuroimaging of language: why hasn't a clearer picture

emerged?. *Language and Linguistics Compass*, *3*(4), 839-865.

Fedorenko, E., Mineroff, Z., Siegelman, M., & Blank, I. (2017). The distinction between lexico-

semantic and syntactic processing is not an organizing dimension of the human language

system. In *30th CUNY Conference on Human Sentence Processing, Cambridge, MA*.

Fedorenko, E., Nieto-Castanon, A., & Kanwisher, N. (2012b). Lexical and syntactic

representations in the brain: an fMRI investigation with multi-voxel pattern analyses.

Neuropsychologia, *50*(4), 499-513.

Fedorenko, E., Scott, T. L., Brunner, P., Coon, W. G., Pritchett, B., Schalk, G., & Kanwisher, N.

(2016). Neural correlate of the construction of sentence meaning. *Proceedings of the National*

Academy of Sciences, *113*(41), E6256-E6262.

Fedorenko & Shain (under review). Local implementation of general computations: The case of

human language comprehension.

Fedorenko, E., & Thompson-Schill, S. L. (2014). Reworking the language network. *Trends in*

cognitive sciences, *18*(3), 120-126.

Fedorenko, E., & Varley, R. (2016). Language and thought are not the same thing: evidence

from neuroimaging and neurological patients. *Annals of the New York Academy of*

Sciences, 1369(1), 132.

Feng, W., Wu, Y., Jan, C., Yu, H., Jiang, X., & Zhou, X. (2017). Effects of contextual relevance

on pragmatic inference during conversation: an fMRI study. *Brain and language*, *171*, 52-61.

Feng, W., Yu, H., & Zhou, X. (2021). Understanding particularized and generalized

conversational implicatures: Is theory-of-mind necessary?. *Brain and Language*, *212*, 104878.

Ferstl, E. C., & von Cramon, D. Y. (2002). What does the frontomedian cortex contribute to

language processing: coherence or theory of mind?. *NeuroImage*, *17*(3), 1599-1612.

Finlayson, M. A., & Winston, P. H. (2011). Narrative is a Key Cognitive Competency. In *BICA*

(p. 110).

Fischl, B., Rajendran, N., Busa, E., Augustinack, J., Hinds, O., Yeo, B. T., ... & Zilles, K. (2007).

Cortical folding patterns and predicting cytoarchitecture. *Cerebral cortex*, *18*(8), 1973-1980.

Fletcher, P. C., Happe, F., Frith, U., Baker, S. C., Dolan, R. J., Frackowiak, R. S., & Frith, C. D.

(1995). Other minds in the brain: a functional imaging study of "theory of mind" in story

comprehension. *Cognition*, *57*(2), 109-128.

Fodor, J. A. (1983). *The modularity of mind*. MIT press.

Foldiak, P., & Young, M. (1995). Sparse coding in the primate cortex. The Handbook of Brain

Theory and Neural Networks, 895-898.

Frith, U., & Happé, F. (1994). Autism: beyond "theory of mind". *Cognition*, *50*(1), 115-132.

from neuroimaging and neurological patients. *Annals of the New York Academy of*

Sciences, 1369(1), 132-153.

Frost, M. A., & Goebel, R. (2012). Measuring structural–functional correspondence: spatial

variability of specialised brain regions after macro-anatomical alignment. *Neuroimage*, *59*(2),

1369-1381.

Frost, M. A., & Goebel, R. (2012). Measuring structural–functional correspondence: spatial

variability of specialised brain regions after macro-anatomical alignment. *Neuroimage*, *59*(2),

1369-1381.

Futrell, R., Gibson, E., Tily, H., Blank, I., Vishnevetsky, A., Piantadosi, S. T., & Fedorenko, E.

(2017). The natural stories corpus. *arXiv preprint arXiv:1708.05763*.

Gallagher, H. L., Happé, F., Brunswick, N., Fletcher, P. C., Frith, U., & Frith, C. D. (2000).

Reading the mind in cartoons and stories: an fMRI study of 'theory of mind' in verbal and

nonverbal tasks. *Neuropsychologia*, *38*(1), 11-21.

Gallivan, J. P., Cavina-Pratesi, C., & Culham, J. C. (2009). Is that within reach? fMRI reveals

that the human superior parieto-occipital cortex encodes objects reachable by the

hand. *Journal of Neuroscience*, *29*(14), 4381-4391.

Grice, H. P. (1957). Meaning. Reprinted in. *Studies in the Way of Words*, 213-223.

Grice, H. P. (1968). Utterer's meaning, sentence-meaning, and word-meaning. In *Philosophy,*

Language, and Artificial Intelligence (pp. 49-66). Springer, Dordrecht.

Grice, H. P. (1975). Logic and conversation. In *Speech acts* (pp. 41-58). Brill.

Gweon, H., Dodell□Feder, D., Bedny, M., & Saxe, R. (2012). Theory of mind performance in

children correlates with functional specialization of a brain region for thinking about thoughts.

Child development, *83*(6), 1853-1868.

Hagoort, P., & Levinson, S. C. (2014). Neuropragmatics. In *The cognitive neurosciences* (pp.

667-674). MIT Press.

Hasson, U., Egidi, G., Marelli, M., & Willems, R. M. (2018). Grounding the neurobiology of

language in first principles: The necessity of non-language-centric explanations for language

comprehension. *Cognition*, *180*, 135-157.

Hasson, U., Furman, O., Clark, D., Dudai, Y., & Davachi, L. (2008). Enhanced intersubject

correlations during movie viewing correlate with successful episodic encoding. *Neuron*, *57*(3),

452-462.

Hasson, U., Malach, R., & Heeger, D. J. (2010). Reliability of cortical activity during natural

stimulation. *Trends in cognitive sciences*, *14*(1), 40-48.

Hasson, U., Nir, Y., Levy, I., Fuhrmann, G., & Malach, R. (2004). Intersubject synchronization

of cortical activity during natural vision. *Science*, *303*(5664), 1634-1640.

HBO (Producer). (2003, December 7). Chapter 1: Bad News [Audio only]. *Angels in America*.

Retrieved from https://www.imdb.com/title/tt0318997/

Heider, F., & Simmel, M. (1944). An experimental study of apparent behavior. *The American

journal of psychology*, *57*(2), 243-259.

Honey, C. J., Thompson, C. R., Lerner, Y., & Hasson, U. (2012). Not lost in translation: neural

responses shared across languages. *Journal of Neuroscience*, *32*(44), 15277-15283.

Huth, A. G., De Heer, W. A., Griffiths, T. L., Theunissen, F. E., & Gallant, J. L. (2016). Natural

speech reveals the semantic maps that tile human cerebral cortex. *Nature*, *532*(7600), 453-

458.

Isik, L., Koldewyn, K., Beeler, D., & Kanwisher, N. (2017). Perceiving social interactions in the

posterior superior temporal sulcus. *Proceedings of the National Academy of*

Sciences, *114*(43), E9145-E9152.

Ivanova, A. A., Mineroff, Z., Zimmerer, V., Kanwisher, N., Varley, R., & Fedorenko, E. (2021).

The language network is recruited but not required for nonverbal event

semantics. *Neurobiology of Language, 2*(2), 176-201.

Ivanova, A. A., Srikant, S., Sueoka, Y., Kean, H. H., Dhamala, R., O'reilly, U. M., ... &

Fedorenko, E. (2020). Comprehension of computer code relies primarily on domain-general

executive brain regions. *Elife, 9*, e58906.

Jackendoff, R. S. (2009). *Language, consciousness, culture: Essays on mental structure*. MIT

Press.

Jacoby, N., & Fedorenko, E. (2020). Discourse-level comprehension engages medial frontal

Theory of Mind brain regions even for expository texts. *Language, Cognition and*

Neuroscience, *35*(6), 780-796.

Jacoby, N., Bruneau, E., Koster-Hale, J., & Saxe, R. (2016). Localizing Pain Matrix and Theory

of Mind networks with both verbal and non-verbal stimuli. *Neuroimage*, *126*, 39-48.

Jang, G., Yoon, S. A., Lee, S. E., Park, H., Kim, J., Ko, J. H., & Park, H. J. (2013). Everyday

conversation requires cognitive inference: neural bases of comprehending implicated

meanings in conversations. *NeuroImage*, *81*, 61-72.

Jenkins, A. C., Dodell-Feder, D., Saxe, R., & Knobe, J. (2014). The neural bases of directed and

spontaneous mental state attributions to group agents. *PLoS One*, *9*(8), e105341.

Jouravlev, O., Zheng, D., Balewski, Z., Pongos, A. L. A., Levan, Z., Goldin-Meadow, S., &

Fedorenko, E. (2019). Speech-accompanying gestures are not processed by the language-

processing mechanisms. *Neuropsychologia*, 107132.

Julian, J. B., Fedorenko, E., Webster, J., & Kanwisher, N. (2012). An algorithmic method for

functionally defining regions of interest in the ventral visual pathway. *Neuroimage, 60*(4),

2357-2364.

Kanwisher, N. (2010). Functional specificity in the human brain: a window into the functional

architecture of the mind. *Proceedings of the National Academy of Sciences*, *107*(25), 11163-

11170.

Keller, G. B., & Mrsic-Flogel, T. D. (2018). Predictive processing: a canonical cortical

computation. *Neuron*, *100*(2), 424-435.

Kandylaki, K. D., Nagels, A., Tune, S., Wiese, R., Bornkessel□Schlesewsky, I., & Kircher, T.

(2015). Processing of false belief passages during natural story comprehension: an fMRI

study. *Human brain mapping*, *36*(11), 4231-4246.

Koster-Hale, J., & Saxe, R. (2013). Functional neuroimaging of theory of mind. *Understanding

other minds: Perspectives from developmental social neuroscience*, 132-163.

Koster-Hale, J., Bedny, M., & Saxe, R. (2014). Thinking about seeing: Perceptual sources of

knowledge are encoded in the theory of mind brain regions of sighted and blind

adults. *Cognition*, *133*(1), 65-78.

Lerner, Y., Epshtein, B., Ullman, S., & Malach, R. (2008). Class information predicts activation by object fragments in human object areas. *Journal of Cognitive Neuroscience*, *20*(7), 1189-1206.

Lerner, Y., Honey, C. J., Silbert, L. J., & Hasson, U. (2011). Topographic mapping of a hierarchy of temporal receptive windows using a narrated story. *Journal of Neuroscience*, *31*(8), 2906-2915.

Mar, R. A. (2011). The neural bases of social cognition and story comprehension. *Annual review of psychology*, *62*, 103-134.

Mason, R. A., & Just, M. A. (2009). The role of the theory‐of‐mind cortical network in the comprehension of narratives. *Language and Linguistics Compass*, *3*(1), 157-174.

Mellem, M. S., Jasmin, K. M., Peng, C., & Martin, A. (2016). Sentence processing in anterior superior temporal cortex shows a social-emotional bias. *Neuropsychologia*, *89*, 217-224.

Monti, M. M., Parsons, L. M., & Osherson, D. N. (2012). Thought beyond language neural dissociation of algebra and natural language. *Psychological Science*, 0956797612437427.

Morosan, P., Rademacher, J., Schleicher, A., Amunts, K., Schormann, T., & Zilles, K. (2001). Human primary auditory cortex: cytoarchitectonic subdivisions and mapping into a spatial reference system. *Neuroimage*, *13*(4), 684-701.

Mr Bean. (2009, Aug 25). *Falling Asleep in Church | Funny Clip | Mr Bean Official* [Video file]. Retrieved from https://www.youtube.com/watch?v=bh__g-ZZ6WA

Nastase, S. A., Goldstein, A., & Hasson, U. (2020). Keep it real: rethinking the primacy of experimental control in cognitive neuroscience. *NeuroImage, 222,* 117254.

Nieto-Castañón, A., & Fedorenko, E. (2012). Subject-specific functional localizers increase sensitivity and functional resolution of multi-subject analyses. *Neuroimage*, *63*(3), 1646-1669.

Nieuwenhuis, S., Forstmann, B. U., & Wagenmakers, E. J. (2011). Erroneous analyses of

interactions in neuroscience: a problem of significance. *Nature neuroscience*, *14*(9), 1105.

Norman-Haignere, S., Kanwisher, N. G., & McDermott, J. H. (2015). Distinct cortical pathways

for music and speech revealed by hypothesis-free voxel decomposition. *Neuron*, *88*(6), 1281-

1296.

Nowak, M. A., & Sigmund, K. (2005). Evolution of indirect reciprocity. *Nature*, *437*(7063),

1291.

Nowak, M., & Highfield, R. (2011). *Supercooperators: Altruism, evolution, and why we need*

each other to succeed. Simon and Schuster.

Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory.

Neuropsychologia, *9*(1), 97-113.

Olshausen, B. A., & Field, D. J. (2004). Sparse coding of sensory inputs. *Current opinion in*

neurobiology, *14*(4), 481-487.

Paunov A.M., Shain, C., Chen, J., Affourtit, J. & Fedorenko, E. (in prep). Spatial overlap

between language and Theory of Mind networks with verbal but not with non-verbal tasks:

fMRI evidence for dissociability.

Paunov, A. M., Blank, I. A., & Fedorenko, E. (2019). Functionally distinct language and Theory

of Mind networks are synchronized at rest and during language comprehension. *Journal of*

neurophysiology, *121*(4), 1244-1265.

Paunov, A.M. (2019). fMRI studies of the relationship between language and theory of mind in

adult cognition. PhD Dissertation. Massachusetts Institute of Technology. Cambridge, MA,

USA.

Piantadosi, S. T., Tily, H., & Gibson, E. (2011). Word lengths are optimized for efficient

communication. *Proceedings of the National Academy of Sciences*, *108*(9), 3526-3529.

Poldrack, R. A. (2006). Can cognitive processes be inferred from neuroimaging data?. *Trends in*

cognitive sciences, *10*(2), 59-63.

Poldrack, R. A., Baker, C. I., Durnez, J., Gorgolewski, K. J., Matthews, P. M., Munafò, M. R., ...

& Yarkoni, T. (2017). Scanning the horizon: towards transparent and reproducible

neuroimaging research. *Nature Reviews Neuroscience*, *18*(2), 115.

Pritchett, B. L., Hoeflin, C., Koldewyn, K., Dechter, E., & Fedorenko, E. (2018). High-level

language processing regions are not engaged in action observation or imitation. *Journal of*

neurophysiology, *120*(5), 2555-2570.

Redcay, E., Velnoskey, K. R., & Rowe, M. L. (2016). Perceived communicative intent in gesture

and language modulates the superior temporal sulcus. *Human brain mapping*, *37*(10), 3444-

3461.

Reddy, A. J., & Wehbe, L. (2020). Syntactic representations in the human brain: beyond effort-

based metrics. *bioRxiv*.

Regev, M., Honey, C. J., Simony, E., & Hasson, U. (2013). Selective and invariant neural

responses to spoken and written narratives. *Journal of Neuroscience*, *33*(40), 15978-15988.

Richardson, H., Koster-Hale, J., Caselli, N., Magid, R., Benedict, R., Olson, H., ... & Saxe, R.

(2020). Reduced neural selectivity for mental states in deaf children with delayed exposure to

sign language. *Nature communications, 11*(1), 1-13.

Richardson, H., Lisandrelli, G., Riobueno-Naylor, A., & Saxe, R. (2018). Development of the

social brain from age three to twelve years. *Nature communications, 9*(1), 1-12.

Sabbagh, M. A., Xu, F., Carlson, S. M., Moses, L. J., & Lee, K. (2006). The development of

executive functioning and theory of mind: A comparison of Chinese and US

preschoolers. *Psychological science*, *17*(1), 74-81.

Saur, D., Schelter, B., Schnell, S., Kratochvil, D., Küpper, H., Kellmeyer, P., ... & Mader, W.

(2010). Combining functional and anatomical connectivity reveals brain networks for auditory

language comprehension. *Neuroimage*, *49*(4), 3187-3197.??

Saxe, R. R., Whitfield☐Gabrieli, S., Scholz, J., & Pelphrey, K. A. (2009). Brain regions for

perceiving and reasoning about other people in school☐aged children. *Child*

development, *80*(4), 1197-1209.

Saxe, R., & Kanwisher, N. (2003). People thinking about thinking people: the role of the

temporo-parietal junction in "theory of mind". Neuroimage, 19(4), 1835-1842.

Saxe, R., & Powell, L. J. (2006). It's the thought that counts: specific brain regions for one

component of theory of mind. Psychological science, 17(8), 692-699.

Saxe, R., Brett, M., & Kanwisher, N. (2006). Divide and conquer: a defense of functional

localizers. Neuroimage, 30(4), 1088-1096.

Saxe, R., Schulz, L. E., & Jiang, Y. V. (2006). Reading minds versus following rules:

Dissociating theory of mind and executive control in the brain. *Social neuroscience*, *1*(3-4),

284-298.

Schmälzle, R., Häcker, F. E., Honey, C. J., & Hasson, U. (2015). Engaged listeners: shared

neural processing of powerful political speeches. *Social cognitive and affective*

neuroscience, *10*(8), 1137-1143.

Scott, T. L., Gallée, J., & Fedorenko, E. (2017). A new fun and robust version of an fMRI

localizer for the frontotemporal language system. *Cognitive neuroscience*, *8*(3), 167-176.

Shahaeian, A., Peterson, C. C., Slaughter, V., & Wellman, H. M. (2011). Culture and the

sequence of steps in theory of mind development. *Developmental psychology*, *47*(5), 1239.

Shain, C., Blank, I. A., Fedorenko, E., Gibson, E., & Schuler, W. (2021). Robust effects of

working memory demand during naturalistic language comprehension in language-selective

cortex. *bioRxiv*.

Shain, C., Blank, I., Van Shijndel, M., Schuler, W. & Fedorenko, E. (2020). fMRI reveals

language-specific predictive coding during naturalistic sentence comprehension.

Neuropsychologia, 138, 107307.

Silbert, L. J., Honey, C. J., Simony, E., Poeppel, D., & Hasson, U. (2014). Coupled neural

systems underlie the production and comprehension of naturalistic narrative

speech. *Proceedings of the National Academy of Sciences*, *111*(43), E4687-E4696.

Silver, N. C., & Dunlap, W. P. (1987). Averaging correlation coefficients: should Fisher's z

transformation be used?. *Journal of Applied Psychology*, *72*(1), 146.

Simon, H. A. (1991). The architecture of complexity. In Facets of systems science (pp. 457-476).

Springer, Boston, MA.

Smith, E. A. (2010). Communication and collective action: language and the evolution of human

cooperation. *Evolution and human behavior*, *31*(4), 231-245.

Sommerfeld, R. D., Krambeck, H. J., & Milinski, M. (2008). Multiple gossip statements and

their effect on reputation and trustworthiness. *Proceedings of the Royal Society B: Biological

Sciences*, *275*(1650), 2529-2536.

Sonkusare, S., Breakspear, M., & Guo, C. (2019). Naturalistic stimuli in neuroscience: critically

acclaimed. *Trends in cognitive sciences, 23*(8), 699-714.

Sperber, D., & Wilson, D. (1986). *Relevance: Communication and cognition* (Vol. 142).

Cambridge, MA: Harvard University Press.

Sperber, D., & Wilson, D. (2002). Pragmatics, modularity and mind□reading. *Mind &*

Language, *17*(1□2), 3-23.

Spotorno, N., Koun, E., Prado, J., Van Der Henst, J. B., & Noveck, I. A. (2012). Neural evidence

that utterance-processing entails mentalizing: The case of irony. *NeuroImage*, *63*(1), 25-39.

Tager-Flusberg, H., Paul, R., & Lord, C. (2005). Handbook of autism and pervasive

developmental disorders, Vol. 1: Diagnosis, development, neurobiology, and behavior.

Tager-Flusberg, H., Paul, R., Lord, C., Volkmar, F., Paul, R., & Klin, A. (2005). Language and

communication in autism. *Handbook of autism and pervasive developmental disorders*, *1*,

335-364. –

Tahmasebi, A. M., Davis, M. H., Wild, C. J., Rodd, J. M., Hakyemez, H., Abolmaesumi, P., &

Johnsrude, I. S. (2011). Is the link between anatomical structure and function equally strong at

all cognitive levels of processing?. *Cerebral Cortex*, *22*(7), 1593-1603.

Theiler, J., Eubank, S., Longtin, A., Galdrikian, B., & Farmer, J. D. (1992). Testing for

nonlinearity in time series: the method of surrogate data. *Physica D: Nonlinear Phenomena*,

58(1-4), 77-94.

Tomasello, M. (2014). The ultra□social animal. *European journal of social psychology*, *44*(3),

187-194.

Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, Mazoyer B,

Joliot M (2002) Automated anatomical labeling of activations in SPM using a macroscopic

anatomical parcellation of the MNI MRI single-subject brain. Neuroimage 15:273–289.

Van Ackeren, M. J., Casasanto, D., Bekkering, H., Hagoort, P., & Rueschemeyer, S. A. (2012).

Pragmatics in action: indirect requests engage theory of mind areas and the cortical motor

network. *Journal of Cognitive Neuroscience*, *24*(11), 2237-2247.

Varley, R., & Siegal, M. (2000). Evidence for cognition without grammar from causal reasoning

and 'theory of mind' in an agrammatic aphasic patient. *Current Biology*, *10*(12), 723-726.

Varley, R., Siegal, M., & Want, S. C. (2001). Severe impairment in grammar does not preclude

theory of mind. *Neurocase*, *7*(6), 489-493.

Wehbe, L., Blank, I., Shain, C., Futrell, R., Levy, R., Malsburg, T. Smith, N., Gibson, E.,

Fedorenko, E. (in press). Incremental language comprehension difficulty predicts activity in

the language network but not the multiple demand network. *Cerebral Cortex.*

Willems, R. M., Nastase, S. A., & Milivojevic, B. (2020). Narratives for neuroscience. *Trends in*

neurosciences, 43(5), 271-273.

Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: a functional connectivity toolbox for

correlated and anticorrelated brain networks. *Brain connectivity*, *2*(3), 125-141.

Wiesmann, C. G., Schreiber, J., Singer, T., Steinbeis, N., & Friederici, A. D. (2017). White

matter maturation is associated with the emergence of Theory of Mind in early

childhood. *Nature communications*, *8*, 14692.

Wilkinson, K. M. (1998). Profiles of language and communication skills in autism. *Mental*

Retardation and Developmental Disabilities Research Reviews, *4*(2), 73-79.

Willems, R. M., Benn, Y., Hagoort, P., Toni, I., & Varley, R. (2011). Communicating without a

functioning   language   system: implications   for   the   role   of   language   in

mentalizing. *Neuropsychologia*, *49*(11), 3130-3135.

Willems, R. M., Van der Haegen, L., Fisher, S. E., & Francks, C. (2014). On the other hand:

including left-handers in cognitive neuroscience and neurogenetics. *Nature Reviews*

Neuroscience, *15*(3), 193.

Wilson, S. M., Molnar-Szakacs, I., & Iacoboni, M. (2007). Beyond superior temporal cortex:

intersubject correlations in narrative speech comprehension. *Cerebral cortex*, *18*(1), 230-242.

Wimmer, H., & Perner, J. (1983). Beliefs about beliefs: Representation and constraining function

of wrong beliefs in young children's understanding of deception. *Cognition*, *13*(1), 103-128.

Wittgenstein, L. (2001). Philosophical Investigations: The German Text, With a Revised English

Translation. 1953. *Trans. GEM Anscombe. 3rd ed. Blackwell.*

Young, L., & Saxe, R. (2008). The neural basis of belief encoding and integration in moral

judgment. *NeuroImage*, *40*(4), 1912-1920.

ToM cognitive demands

	Non-conversation linguistic inputs	Conversation (observer)	Conversation (participant)

Non-conversation linguistic inputs

COMPREHENSION

Conversation (observer)

COMPREHENSION

Conversation (participant)

COMPREHENSION + PRODUCTION

Linguistic content

- Representing the mental states (stated or inferred) of ***characters*** to interpret their actions
- Representing the mental state of the ***narrator*** to interpret attitudes towards events and characters

- Representing the mental states of ***persons under discussion***
- Representing the mental states of potentially ***affected parties***: conversation-external, conversants, self

- Representing the mental states of ***persons under discussion***
- Representing the mental states of potentially ***affected parties***: conversation-external, conversants (including self)

Communicative intent

- Representing the mental state of the ***author*** to interpret structural decisions (why is X said / not-said at time Y?)

- Representing the mental states of the ***conversants*** to interpret their linguistic utterances (and other actions)

- Representing the mental states of the ***conversant(s)*** to
  - interpret their utterances (and other actions)
  - generate appropriate responses

Language

Sentences

| + | THE | SPEECH | ... | WAS | TOO | LONG | FOR | THE | MEETING | + |

Non-words

| + | LAS | TUPING | ... | CRE | POME | VILLPA | OLP | PRONT | CHO | + |

| 100ms | 450ms | 450ms | | 450ms | 450ms | 450ms | 450ms | 450ms | 450ms | 500ms |

Theory of Mind

False Belief

| + | The morning of the high school dance Sarah placed her high heel shoes under her dress and then went shopping. That afternoon, her sister borrowed the shoes and later put them under Sarah's bed. | Sarah gets ready assuming her shoes are under the dress.
True   False | + |

False Physical

| + | A large oak tree stood in front of City Hall from the time the building was built. Last year the tree fell down and was replaced by a stone fountain. | An antique drawing of City Hall shows a fountain in front.
True   False | + |

| 12s | 12s | 4s Response | 12s |