Bioremediation of industrial pollutants by insects expressing a fungal laccase

Michael Clark[1ŧ], Kate Tepper[1ŧ], Kerstin Petroll[2], Sheemal Kumar[1], Anwar Sunna[2,3], Maciej

Maselko[1,3,4,5]*

Affiliations

[1] Applied BioSciences, Macquarie University, Sydney, NSW, Australia
[2] Department of Molecular Sciences, Macquarie University, Sydney, NSW, Australia
[3] Biomolecular Discovery Research Centre, Macquarie University, Sydney, NSW, Australia
[4] CSIRO Synthetic Biology Future Science Platform, Brisbane, QLD, Australia
[5] CSIRO Health and Biosecurity, Canberra, ACT, Australia
[ŧ] Authors contributed equally to this research.
*Correspondence and requests for materials should be sent to maciej.maselko@mq.edu.au
12

Abstract

Inadequate management of household and industrial wastes pose major challenges to human

and environmental health. Advances in synthetic biology may help address these challenges

by engineering biological systems to perform new functions such as biomanufacturing of high-

value compounds from low-value waste streams and bioremediation of industrial pollutants.

The current emphasis on microbial systems for biomanufacturing, which often require highly

pre-processed inputs and sophisticated infrastructure, is not feasible for many waste streams.

Concerns about transgene biocontainment have limited the release of engineered microbes

or plants for bioremediation. Engineering animals may provide opportunities for utilizing

various waste streams that are not suitable for microbial biomanufacturing while effective

transgene biocontainment options should enable *in situ* bioremediation. Here, we engineer

the model insect *Drosophila melanogaster* to express a functional laccase from the fungus

Trametes trogii. Laccase expressing flies reduced concentrations of the endocrine disruptor

bisphenol A by more than 50% when present in their growth media. A lyophilized powder

made from engineered adult flies retained substantial enzymatic activity, degrading more

than 90% of bisphenol A and the textile dye indigo carmine in aqueous solutions. Our results

demonstrate that transgenic animals may be used to bioremediate environmental

contaminants *in vivo* and serve as novel production platforms for industrial enzymes. These

results support further development of insects, and possibly other animals, as bioproduction

platforms and their potential use in bioremediation.

Main Text

Developing economically viable approaches for sustainable waste management and

mitigation of environmental pollution are important for ensuring a high quality of life[1]. At

present, over half of all municipal solid waste is either landfilled[2], where it is estimated to

contribute to 11% of global methane emissions[3,4], or openly dumped, where it contaminates

drinking water and disrupts natural ecosystems[5]. Pollution from industrial wastes includes a

variety of persistent compounds which have been linked to neurological defects, cancer, and

species extinction[6–10].

Regulatory changes including prohibiting discharge of industrial pollutants and reducing the

landfilling of food waste are helpful, however, the cost of compliance can be substantial[11].

Generating value from a variety of waste-streams and developing inexpensive methods to

treat pollutants *in situ* will reduce the costs of achieving sustainability targets and help

support a circular economy[12].

We propose developing solutions by expanding the enzymatic repertoire of select animals to

consume waste, degrade xenobiotic chemicals, and produce high-value compounds (Fig. 1).

Microbial and fungal enzymes can enable animals to breakdown pollutants *in situ*. These may

include herbivores that degrade pollutants accumulating in plants, fish that secrete enzymes

to degrade aquatic pollutants, or invertebrates engineered to interrupt the biomagnification

of xenobiotics in food webs. Effective transgene biocontainment safeguards are currently a

challenge for plants/microbes[13,14], but are possible in animals via surgical sterilization or

engineered genetic incompatibilities[15]. Developing insects as a biomanufacturing platforms

for industrial enzymes, lipids, or other molecules is also an unexplored avenue for waste

valorisation.

Insects have advantages over bacteria and fungi for processing many waste-streams. Their

complex digestive systems, including mastication, allow them to consume a variety of low-

value wastes, including municipal organic waste[16] and livestock manure[17]. Crude waste can

be fed directly to insects without pre-processing, refinement, and sterilization necessary for

microbial fermentation[18,19]. Some insect species, such as Black soldier flies (*Hermetia

illucens*), which are already used to manage waste have excellent tolerance to bacteria and

fungi[20] and are easily separated from the organic waste for downstream processing[21]. Insect

biotechnology is readily scalable for mass production, as the infrastructure can be relatively

unsophisticated, and the installation costs and land use requirements are low[22].

Furthermore, utilising organic waste means crops that could otherwise be used for food do

not need to be diverted as feedstocks for fermentation, and will not threaten food security.

There is a long history of using lepidopteran insect larvae to make recombinant proteins

starting with *Bombyx mori* expressing human α-interferon[23]. However, these utilize transient

baculoviral systems to infect the larvae. This approach is not practical for large-scale or long-

term, multi-generational use. Here we explore the potential of stably engineered insects as

platforms for the production of an industrially relevant fungal laccase and bioremediation of

polluted sites. Laccases are multicopper oxidases that are present in many taxa. Fungal

laccases were selected due to their demonstrated utility for bioremediation of a wide variety

of industrial pollutants such as bisphenols[24], industrial dyes[25], pharmaceuticals[26], phenolics[27],

polycyclic aromatic hydrocarbons[28], perfluoroalkyl and polyfluoroalkyl substances (PFAS)[29,30],

plastics[31], pesticides[32], and mycotoxin contaminants such as aflatoxins[33]. Fungal laccases also

have diverse industrial applications in the textile, paper and pulp, food, pharmaceutical,

chemical synthesis, and forestry industries[34–36]. They have been applied to treat distillery[37],

paper and pulp[38], and olive mill[39] industry wastewater. Ligninolytic enzymes, such as fungal

laccases, also show potential application in the delignification of feedstocks for biofuel

production[40]. However, fungal laccases have not yet reached appreciable commercial scale

production. This is because the native hosts for laccases often produce low yields and

although this has been improved using heterologous hosts, the production levels are still too

low for many commercial applications[41].

We engineered the common fruit fly (*Drosophila melanogaster*) to express a laccase from the

fungus *Trametes trogii*. Transgenic flies degraded bisphenol A (BPA), a widespread endocrine

disruptor[42], when present in their diet. We also demonstrate that aqueous solutions of BPA

and indigo carmine (IC), a textile dye and aquatic pollutant[43], can be treated with a lyophilized

powder from adult transgenic flies to degrade both compounds. Our results describe the first

instance of engineering an animal to perform an ecosystem service by heterologous

expression of a fungal enzyme and demonstrate the potential for using transgenic strains of

insects for enzyme bioproduction.

Fig. 1: Applications of transgenic animals for bioremediation and bioproduction.

Results

Screening for fungal laccase activity in transgenic *D. melanogaster*

D. melanogaster and other insects express endogenous laccases involved in immune defense[44]. However, it was unknown if an insect could heterologously express a functional fungal laccase or if its expression would be toxic. We were also uncertain which tissues would provide the optimal expression of a functional laccase for either *in vitro* or *in vivo* activity. We selected a short tubulin promoter, previously used to express a toxic gene[45], with the goal of achieving moderate expression across many tissues. The native fungal signal peptides were replaced with the *D. melanogaster* larval cuticle protein 9[46] signal peptide to facilitate extracellular secretion.

We generated homozygous transgenic strains expressing laccases from two fungal species, *T. trogii* and *Polyporus brumalis*, known to have a broad substrate range and high redox potential (Table 1). Fly lysates from both were assayed for oxidation of the chromogenic laccase substrate 2,2'-Azino-bis(3-ethylbenzthiazoline-6-sulfonic acid) (ABTS). The

engineered *D. melanogaster* strain Dm/Tt.Lcc1, which expresses a laccase from *T. trogii,*

oxidized ABTS (Fig. 2a). There was no activity observed in the wild-type or from the

engineered Dm/Pb.Lac1 strain expressing a laccase from *P. brumalis* (Fig. 2a). Moving

forward, we focused on characterizing Dm/Tt.Lcc1.

We next determined if Dm/Tt.Lcc1 flies are capable of degrading bisphenol A (BPA), an

endocrine disruptor and emerging environmental contaminant found in many consumer

products[47]. BPA has been shown to be a substrate of *T. trogii* laccase when expressed from

its native host[48]. We incubated lysates from flies with 25 µg/mL BPA for 21 hours and

measured BPA concentrations by liquid-chromatography mass spectrometry (LC-MS). Lysates

from Dm/Tt.Lcc1 reduced BPA concentrations by 95% compared to the no-fly controls and

the wild-type fly lysate (Fig. 2b-c).

Table 1. The fungal laccases engineered into *D. melanogaster*

Gene name	Native host	Genbank accession	Redox potential (V vs. NHE, pH 5)	Laccase substrates
Lcc1	*Trametes trogii*	AJ294820.1	$E\circ$ 0.79 V [49]	Aminophenols, polyphenols, industrial dyes[50–52].
Lac1	*Polyporus brumalis*	ABN13591.1	$E\circ$ 0.80 V [53]	Industrial dyes, aromatic hydrocarbons [53].

Fig. 2: *In vitro* laccase substrate oxidation in fly lysates. (A) Fly lysate oxidation of 1 mg/mL

ABTS for 2 hours. *n* = 2 biologically independent replicates for WT, *n* = 6 for all others. **(B)** Fly

lysate degradation of 25 µg/mL BPA for 21 h. *n* = 3 biologically independent replicates. **(C)**

Overlaid representative extracted ion chromatograms (EIC) (m/z 227.15) displaying the BPA

peak for samples shown in **B**. Dm/Tt.Lcc1 = *D. melanogaster* engineered with *T. trogii* laccase,

Dm/Pb.Lac1 = *D. melanogaster* engineered with *P. brumalis* laccase, WT = wild-type, NFC =

No-Fly control. The statistical analysis was conducted using one-way ANOVA using Dunnett's

method compared to WT, where ns = not significant.

***In vivo* laccase activity**

We evaluated the potential of laccase expressing insects to oxidize substrates present in their

environment by secreted enzymes. Flies were reared on a minimal media to avoid binding of

substrates to insoluble components present in standard media. The minimal media included

1 mM $CuSO_4$ to supply the laccase co-factor. First, we reared 3 adult males and 3 adult females

on minimal diet including 1 mg/mL ABTS for 2 days, after which they were removed. Larvae

of the mated flies were grown for an additional 5 days. Dm/Tt.Lcc1 flies oxidized ABTS as

indicated by blue colorization of the fly media, however, no oxidation was observed with wild-

type flies (Fig. 3a). We controlled for the possibility that the ABTS oxidation was due to laccase

that may have been leaking from dead larvae by incubating 6 male flies in ABTS media. After

11 days, only the Dm/Tt.Lcc1 males had oxidized ABTS (Fig. 3b). This result indicates that the

oxidation was due to enzyme secreted into the media.

Next, we reared 5 male and 5 female flies in a minimal diet containing 25 µg/mL BPA. Adults

were removed after three days. On day 10, BPA was extracted from their diet and quantified

by Ultra-High-Performance Liquid Chromatography coupled with fluorescence detection

(UHPLC-FLD). Both Dm/Tt.Lcc1 and wild-type flies had reduced BPA concentrations by 36%

and 45%, respectively, compared to no-fly controls (Fig. 3c). There was no significant

difference between Dm/Tt.Lcc1 and wild-type indicating that the reduction in BPA

concentration was not due to the activity of the laccase transgene. The reduction in BPA

concentration for wild-type flies relative to the no-fly control may be in part due to BPA

absorbed by the adult flies which were removed after three days. Another possibility is that

BPA detoxification pathways have been characterized in mammals[54] and it is possible *D.*

melanogaster also contains endogenous pathways capable of detoxifying BPA.

Some laccase substrates, such as ABTS, can function as redox mediators to facilitate the

oxidation of additional compounds[55]. When *in vivo* BPA degradation was tested in the

presence of ABTS, Dm/Tt.Lcc1 flies effectively reduced BPA concentration by 68% and 56%

compared to the no-fly control and wild-type, respectively (Fig. 3d). It is not clear why ABTS

was necessary for the *in vivo* assay and not the *in vitro* lysate assays. Adding lysate directly to

the media also required ABTS for efficient degredation, suggesting that there is some

component of the diet which may be interfering (Fig. S1).

Fig. 3: *In vivo* laccase enzyme activity. (A) *In vivo* oxidation of 1 mg/mL ABTS after 7 days. (**B**) *In vivo* oxidation of 1 mg/mL ABTS after 11 days using only male flies. **(C, D)** *In vivo* degradation of 25 µg/mL BPA without ABTS **(C)** and with 1 mg/mL ABTS **(D).** *n* = 3 biologically

172   independent replicates. Dm/Tt.Lcc1 = *D. melanogaster* expressing *T. trogii* laccase, WT = wild-

173   type, NFC = No-Fly control. The statistical analysis was conducted using one-way ANOVA using

174   Dunnett's method compared to WT, where ns = not significant.

Lyophilized insect powder laccase activity

We next examined the potential of insects as platforms to produce a lyophilized laccase which

can be easily stored and transported before use. A crude lyophilized powder from Dm/Tt.Lcc1

was compared to a commercially available purified laccase from *Trametes versicolor* using

ABTS as a substrate. The lyophilized Dm/Tt.Lcc1 displayed a total laccase activity of 1.4 U/g ±

0.1 SEM dry weight, while the commercial purified laccase displayed a specific activity of 75.3

U/g ± 6 SEM dry weight.

We next tested if Dm/Tt.Lcc1 lyophilized whole-fly powder had activity against indigo

carmine, a textile industry dye and pollutant[43]. Incubating 5 mg/mL of Dm/Tt.Lcc1 lyophilized

powder in water containing 100 mg/L indigo carmine resulted in 90% decolorization after 48

h; however, WT fly lyophilized powder also decolorized 55% of the dye (Fig. 4a). We observed

that insoluble material in the powder appeared to adsorb the dye. However, incubating dye

using supernatant from centrifuged lyophilized Dm/Tt.Lcc1 in water did not appear to

decolorize indigo carmine compared to controls (Fig. 4b). When the Cu^{2+} co-factor was

included, supernatant from centrifuged lyophilized powder in water decolorized nearly 100%

of indigo carmine in Dm/Tt.Lcc1 and 25% from WT flies after 90 hours (Fig. 4c-d).

We also evaluated Dm/Tt.Lcc1 lyophilized powder activity against BPA. Incubating 12.5 µg/mL

BPA with soluble enzyme from the lyophilized powder in the presence of Cu^{2+} co-factor

resulted in a 96% reduction of BPA from Dm/Tt.Lcc1 fly powder after 90 hours. Incubation

with WT fly powder resulted in a 3% reduction of BPA (Fig. 4e-f).

A. IC decolorization (%) in whole lyophilized powder

B. Time course IC decolorization (%) in the soluble fraction of lyophilized powder

C. Time course IC decolorization (%) in the soluble fraction of lyophilized powder with copper

D. IC decolorization (%) in the soluble fraction of lyophilized powder with copper

E. Time course BPA oxidation in the soluble fraction of lyophilized powder with copper

F. BPA oxidation in the soluble fraction of lyophilized powder with copper

Fig. 4: *In vitro* activity of lyophilized Dm/Tt.Lcc1 whole-fly powder on laccase substrates. (A)

Decolorization (%) of 100mg/L indigo carmine at 610 nm in whole lyophilized powder

resuspended in water after 48 hours. *n* = 3 biologically independent replicates. **(B-D)** Time

course decolorization (%) of 100 mg/mL indigo carmine in the soluble fraction of lyophilized

powder in water **(B)** or supplemented with 10 mM $CuSO_4$ **(C)**. Statistical analysis of **C** was

performed at end point **(D)**. *n* = 3 biologically independent replicates. **(E,F)** Time course

degradation of 12.5 μg/mL BPA in the soluble fraction of lyophilized samples in water

supplemented with 10 mM $CuSO_4$ **(E)**. Statistical analysis of **E** was performed at the end point

(F). *n* = 3 biologically independent replicates. Dm/Tt.Lcc1 = *D. melanogaster* engineered with

T. trogii laccase, WT = wild-type, NFC = No-Fly control. The statistical analysis was conducted

using one-way ANOVA using Dunnett's method compared to WT, where ns = not significant.

Discussion

Engineering animals to heterologously express fungal/microbial enzymes may improve

sustainable waste management, bioremediation, and facilitate the use of low-value waste

streams as inputs for the production of high-value products such as industrial enzymes. Here,

we engineer *D. melanogaster* to express a functional laccase from the fungus *T. trogii*. Lysates

of this engineered fly (Dm/Tt.Lcc1) oxidized the endocrine disruptor BPA by 95% *in vitro*

compared to the WT flies. Engineered flies also degraded BPA by 56% *in vivo* when added to

their diet. Further, a powder prepared from lyophilized engineered flies also degraded BPA

by 96% and the textile dye indigo carmine by almost 100% *in vitro*.

Mediator compounds are often used to improve the substrate range and activity of laccases[56].

Although Dm/Tt.Lcc1 lyophilized powder did not require the addition of ABTS as a mediator

for degrading BPA, *in vivo* BPA degradation did. White rot fungi such as *Phanerochaete*

chrysosporium and *Pycnoporus cinnabarinus* synthesize and secrete endogenous redox

mediators[57,58] and it may be possible to engineer animals to secrete redox mediators to also

enhance their bioremediation activities. Alternatively, colloidal lignin or derivatives of lignin

also show potential to act as redox mediators[59,60], and including them may be feasible to assist

with *in vivo* activity.

Using transgenic insects as platforms to convert waste to high-value products, such as

industrial enzymes, will require high production titres. Dm/Tt.Lcc1 lyophilized powder made

from whole flies had 1.4 U/g of laccase activity without any attempt at optimization compared

to 75.3 U/g of activity from a commercially available purified *T. versicolor* laccase. This is a

very promising start and developing tools to genetically manipulate more relevant hosts and

strain engineering is needed for insects to reach their potential as bioproduction platforms.

In summary, our findings demonstrate for the first time that insects can be stably engineered

to express a functional fungal laccase, which demonstrated *in vivo and in vitro* laccase activity.

Transgenic flies were capable of oxidizing laccase substrates in their environment and they

can be used as a lyophilized powder to oxidize substrates in aqueous solutions. These results

provide strong support for further research in using transgenic animals as platforms for

bioremediation, waste stream management, and recombinant protein production.

Methods

Construct design and assembly

NEBuilder® HiFi DNA Assembly Master Mix (New England Biolabs) was used to assemble all

constructs. pMBO2744[15] was used as a backbone to generate the enzyme expression

plasmids. A gBLOCK (IDT®) was synthesized that encodes a short alpha-tubulin promoter, a

downstream SV40 polyadenylation sequence, and a NotI restriction site in between. This

gBLOCK was installed into NotI linearized pMB02744 by HiFi assembly to produce pMC-1-1-1

(Supplementary Table 4). This plasmid was then linearized with NotI, and gBLOCKs that

encode for *T. trogii* laccase or *P. brumalis* laccase enzyme were installed with HiFi assembly

to generate pMC-1-2-3 and pMC-1-2-6, respectively (Supplementary Table 4). The DNA

sequences were codon optimized for expression in *D. melanogaster* using the IDT® codon

optimization tool. The native fungal signal peptides were replaced with the larval cuticle

protein 9 signal peptide. The Kozak sequence, AATCTTACAAA, was upstream of the start

codon. See Supplementary Table 2 for plasmid descriptions and Supplementary Table 3 for

primers used in this study.

Insect rearing and generation of transgenic fly strains

Canton S wild-type flies were from the Bloomington Stock Centre (64349, RRID: BDSC 64349).

Enzyme expression plasmids were purified using ZymoPURE™ II Plasmid Midiprep Kit (Zymo

Research #D4200) and sent to BestGene Inc (Chino Hills, Ca) for ϕC31 mediated integration.

All strains were maintained on a cornmeal diet based on the Bloomington standard Nutri-Fly

formulation (catalog number 66-113; Genesee Scientific). In preparation for the *in vitro* or *in*

vivo laccase enzyme activity assay, the diet for both the transgenic and wild-type strains was

supplemented with $CuSO_4 \cdot 5H_2O$ (1 mM, Merck, CAS-no: 7758-99-9). Flies were reared in a

controlled environment room at 25.0˚C, 75% humidity, and a 12 hour light/dark cycle with a

30 minute transition period. See Supplementary Table 1 for all of the fly strains used in this

study. Biosafety approval was granted by the Institutional Biosafety Committee of Macquarie

University (#5049).

***In vitro* enzyme activity assay**

To obtain a fly lysate, ten frozen adult flies were homogenized with a motorized pestle in the

stated buffer supplemented with 10 mM $CuSO4 \cdot 5H2O$ and acid-washed glass beads. The

samples were then agitated along a tube rack 10 times. Each tube was incubated for 15

minutes at 25 °C and centrifuged twice to remove insoluble debris.

For the ABTS oxidation assay, fly lysates were prepared in citrate-phosphate buffer (pH 5) and

added to 1 mM ABTS. The tubes were incubated at 25 ˚C for 2.5 hours. The reaction was

quenched by adding 5% trichloroacetic acid (TCA, Sigma Aldrich, catalogue #T0699, CAS-

no:76-03-9). A UV-Vis spectrophotometer (Jasco V-760 UV-Vis spectrophotometer) was used

to measure the absorbance at 420 nm. The measurements were blanked to an equivalent

sample that did not contain ABTS.

For the BPA degradation assay, fly lysates or controls were prepared in sodium acetate buffer

(pH 5) and added to 25 µg/mL BPA. The assay mixtures were incubated at room temperature

for 21 hours. The reaction was quenched by adding 5% (v/v) trichloroacetic acid (Sigma

Aldrich, catalog #T0699, CAS-no:76-03-9). BPA concentration was analyzed by LC-MS as

described below.

***In vivo* laccase activity assays**

Minimal media:

The minimal media components were for 100 mL: 1.58 g yeast, 0.66 g agar, 5 g sugar, 4 mL 1

M propionic acid (pH 4.4), 12.4 mg $CuSO_4 \cdot 5H_2O$ (1 mM final concentration). A buffer of 20

mM citrate-phosphate (pH 5) was added for the initial ABTS assays, 25 mM sodium acetate

buffer (pH 5) was used for all BPA experiments.

ABTS only *in vivo* experiments:

To minimal media 100 mg ABTS (1 mg/mL final concentration) was added. Three female and

three male adult flies were incubated on the media for 48 hours, after which they were

removed. For the all-male ABTS assay, the adult flies remained in the vial for 11 days. The

appearance of the dark green radical cation was monitored and recorded with photographs.

For the ABTS + BPA experiments:

To minimal media, BPA was added (25 µg/mL final concentration). ABTS (1 mg/mL final

concentration) was added for the +BPA, + ABTS condition.

For each condition, the liquid media was pipetted into a 14 mL polypropylene round-bottom

culture tube. 5 male and 5 female adult flies were incubated on the media for 3 days and

then removed. For the lysate condition, 50 µL of lysate from ~100 flies was added to the

tube. After a total of 10 days the enzymatic reaction was stopped with TCA (5% v/v final

concentration). The media was melted in a water bath, after which methanol (making ~1:1

methanol: aqueous TCA for a two-fold dilution of BPA) was added, followed by vortexing.

The tubes were then incubated at 300 rpm and 37 ˚C for 30 minutes. After incubation, the

tubes were centrifuged twice with the final supernatant reserved for BPA quantification by

UHPLC-FLD as described below.

Lyophilized Dm/Tt.Lcc1 enzyme activity assays

Frozen Dm/Tt.Lcc1 or wild-type flies were lyophilized (Alpha 1-4 LDplus, Christ) at -45 °C at

0.1 mBar. The freeze-dried flies were homogenized in a porcelain mortar and pestle and

stored at 4 °C.

For the activity assays, lyophilized Dm/Tt.Lcc1 and commercial *T. versicolor* laccase (Sigma-

Aldrich, Catalogue 38429) were resuspended in 25 mM sodium acetate buffer (pH 5)

supplemented with 10 mM $CuSO_4 \bullet 5H_2O$. Lyophilized Dm/Tt.Lcc1 did not dissolve, and the

soluble fraction was extracted at 25 °C for 15 minutes at 800 rpm and centrifuged twice to

remove the insoluble debris. Enzyme serial dilutions were added in duplicate to 1 mM ABTS

and ABTS oxidation was monitored on a PHERAStar FS plate reader (BMG Labtech) at 420 nm

at 25 °C. One unit of enzyme activity was defined as the amount of enzyme that catalyzes the

oxidation of 1 μmol ABTS min^{-1} using the molar extinction coefficient 36 mM^{-1} cm^{-1} [61].

For the indigo carmine decolorization assay, 5 mg/mL of lyophilized powder or controls in

Millipore water were added to 100 mg/L indigo carmine (Sigma-Aldrich catalog 131164) and

incubated at RT. At 0 and 48 hours, indigo carmine decolorization was monitored on a

PHERAStar FS plate reader (BMG Labtech) at 610 nm.

As indigo carmine appeared to adsorb to the lyophilized powder, the soluble fraction was

tested for enzymatic activity. 5 mg/mL lyophilized powder or controls in Millipore water

supplemented with and without 10 mM $CuSO_4 \bullet 5H_2O$ were extracted at 25 °C for 15 minutes

at 800 rpm. The samples were centrifuged twice to remove the insoluble debris. The

supernatant of the lyophilized powder and controls were added to 100 mg/L indigo carmine

and incubated at room temperature. Indigo carmine decolorization was monitored on a

PHERAStar FS plate reader (BMG Labtech) at 610 nm at 0, 21, 48, 72, and 90 hours.

For the BPA degradation assay, the soluble fraction of the lyophilized powder was extracted

as described in the indigo carmine assay. The supernatant of the lyophilized powder and

controls were added to 12.5 μg/mL BPA and incubated at room temperature. At 0, 48, and 96

hours, 100 μL aliquots were removed and the reaction was stopped using a 3 KDa nominal

molecular weight limit (NMWL) centrifuge filter (Merck Amicon Ultra 0.5mL Centrifugal

Filters, UFC500396). BPA concentration was analyzed by UHPLC-FLD as described below.

BPA quantification

BPA in samples was either quantified by UHPLC-FLD analysis or by LC-MS. UHPLC-FLD analysis

was performed using an Agilent RRHD ZORBAX Eclipse Plus C18 column (3.0 x 150 mm 1.8 μm,

p/n 959759-302) equipped with a UHPLC Eclipse Plus C18 Guard Column, (2.1 mm 1.8 μm,

p/n 821725-901). The isocratic mobile phase consisted of acetonitrile (ACN):10 mM KH_2PO_4

(50:50 v/v) at a flow rate of 0.5 mL/min. A gradient wash step was introduced after each run

with eluent B 100% ACN and eluent A 10mM KH_2PO_4, starting with 45% B1 from 0 - 1 minute,

70% B from 1 – 6 minutes, 5% B from 6 – 11 minutes, and finally 50% B from 11 - 16 minutes

at 0.3 mL/min. BPA was detected by setting FLD excitation wavelength at 229 nm, and

emission wavelength at 316 nm and a gain setting of 12. Linear regression of 7-point

standards from 12.5 ng/mL to 12,500 ng/mL BPA was performed to calibrate BPA

concentration. All samples and standards were either filtered using syringe filters (Filtropur S

0.2 μm, Sarstedt, 83.1826.001) or centrifuge filters (Merck Amicon Ultra 0.5mL Centrifugal

Filters, UFC500396) prior to injection into the UHPLC.

LC-MS validation and quantification of BPA was performed externally at the Mass

Spectrometry Facility at the University of Sydney. LC separation was performed using a

Waters Sunfire C18 column (5 μm, 2.1 mm ID x 15 mm) coupled to a Bruker amaZon SL mass

spectrometer. Gradient elution was performed with a mobile phase consisting of (A)

methanol and (B) water with 0.1% acetic acid at a flow rate of 0.3 mL/min. The gradient

consisted of 20% A from 0 - 9.3 minutes, 76% A from 9.3 - 17.3 minutes, 92% A from 17.3 –

26 minutes, and finally 20% A from 26 - 35 minutes. MS analysis was performed using

Atmospheric Pressure Chemical Ionisation (ACPI) under positive and negative ionization mode

over a range of 200-1400 m/z. The operation parameters were: Nebulizer 27.3 psi, Dry Gas

4.0 L/min, Dry temperature 180 °C, APCI Vaporisation temperature: 400 °C, APCI Corona

needle current: 6000 nA, Mass mode: enhanced resolution, SPS (Smart parameter setting)

Target mass 150 m/z, compound stability 100%. Linear regression of 6-point BPA standards

from 20 ng/mL to 10 µg/mL BPA was performed to calibrate BPA concentration. An internal

deuterated BPA-D$_{16}$ standard was spiked into the samples prior to injection to calibrate

ionization in the sample matrix compared to the standard curve matrix prepared in methanol.

All samples were centrifuge filtered (Merck Amicon Ultra 0.5mL Centrifugal Filters,

UFC500396) prior to injection into the LC-MS.

Statistical analysis

Figures were generated in Prism 9 by GraphPad. For statistical analysis, a one-way ANOVA

using Dunnet's method was conducted. The significance level for the analysis was 0.05, and

all error bars represent the standard error of the mean. The exact *P* values, statistical tests

used, and sample numbers (*n*) can be found in the figure legend.

Acknowledgements:

We are grateful Dr. Nick Proschogo from the Mass Spectrometry Facility at the University of

Sydney. **Funding:** This material is based upon work supported by the International

Technology Center Pacific (ITC-PAC) under Contract No. FA520920P0100, a Macquarie

University seeding grant, and the CSIRO Synthetic Biology Future Science Platform. **Author**

Contributions: MC, KT, and MM conceived of this study. MC, KT, KP, AS, and MM designed

experiments and analyzed data. MC, KT, KP, and SK performed the experiments. All authors

contributed to writing the manuscript. **Competing Interests:** The authors have filed a patent

application for aspects of this work. **Data and materials availability:** All data is available

upon request.

Supplementary Materials:

Supplementary Figure 1
Supplementary Tables 1-4
Supplementary Data File
386

<u>**Supplementary information**</u>

Supplementary figure 1: *In vivo* BPA degradation conditions comparing Dm/Tt.Lcc1 lysate to
NFC. Fly lysate was added to media that contained 25 µg/mL BPA with and without 1 mg/mL
ABTS. Dm/Tt.Lcc1 lysate was unable to degrade BPA without ABTS. *n* = 3 biologically
independent replicates. The statistical analysis was a one-way ANOVA with Dunnett's
method. ns= not significant, Dm/Tt.Lcc1 = *D. melanogaster* engineered with *T. trogii* laccase,
NFC = no fly control.

Supplementary Table 1: *D. melanogaster* strains used in this study

Strain	Description	Reference	BDSC #
Wild Type	Canton S	[62]	64349
2nd Chr. attP3	y^1 $w^{1118;}$ PBac{y[+]-attP-3B}VK00037	[63]	9752
2nd Chr. Bal, Tm2/Tm6b	w^{1118}; Gla/CyO; Tm2/Tm6b	This study	
2nd Chr. Bal, Tm2/Tm6b	w^{1118}; Bl/CyO; Tm2/Tm6b	This study	
Dm/Tt.Lcc1	y^1 $w^{1118;}$ PBac{Tt.Lcc1}attP-3B	This study	
Dm/Pb.Lac1	y^1 $w^{1118;}$ PBac{Pb.Lac1}attP-3B	This study	

Supplementary Table 2: Plasmids used in this study

Plasmid	Integrated in Fly Strain	Description	Reference/ GenBank Accession #
pMB02744	NA	Contains a mini-white gene for *D. melanogaster* selection, an *AttB* site for genomic integration, and an ampicillin resistance cassette for cloning in *E. coli*.	[15]
pMC-1-1-1	NA	Empty vector, with short alpha-tubulin promoter and SV40 regulatory sequences	This study
pMC-1-2-3	Dm/Tt.Lcc1	Short alpha-tubulin promoter driving expression of *Lcc1* from *Trametes trogii*	This study
pMC-1-2-6	Dm/Pb.Lac1	Short alpha-tubulin promoter driving expression of *Lac1* from *Polyporus brumalis*	This study

Supplementary table 3: Primers used in this study		
Name	Sequence (5' → 3')	Notes
1stconstruct_F_2.0	GCTTGGATTTCACTGGAACTAG	pMC-1-1-1 sequencing
1stconstruct_R	TCGAGACCGTGACCTACATC	pMC-1-1-1 sequencing
2ndconstruct_F_2.0	cgtttgtcaagcctcatagc	pMC-1-2-3/pMC-1-2-6 sequencing
2ndconstruct_R_2.0	ttcattttatgtttcaggttcaggg	pMC-1-2-3/pMC-1-2-6 sequencing
P.bru_F	TGACTTTCAGGTGCCCGATC	pMC-1-2-6 sequencing
P.bru_R	GCCGGGAATGTGAGTTCGATG	pMC-1-2-6 sequencing

| T.tro_F | ATGGCCCTGCATTCGTCAAC | pMC-1-2-3 sequencing |
| T.tro_R | CGTTGCTGGCAAGCTGATTTC | pMC-1-2-3 sequencing |

Supplementary table 4: gblocks used to construct laccase expression plasmids:	
Name and description	**Sequence (5' → 3')**
<u>1st construct</u> Gblock that includes short alpha-tubulin and SV40 regulatory sequences for expression of the enzymes in *Drosophila melanogaster*. Contains a NotI site between the promoter and terminator for assembly of enzyme gene gblocks. Generates pMC-1-1-1 from pMBO2744	AGCGTAAGCTTCGTACGTAGCGACCGTCTCAAAGTACTGCCTTTCTGCGTTGGAAAAC ATCGCCTTTTTCGTCCAAAAGGAGTCCCCAGGTTCGATCCGCATGGCGTTGTGCGTGC GTGCCTTTCTTTTCAAATGATTACGGCTATTAACTTGGGGGCGTTAAGTTGGAAACAC GTAAATTGCAGACTGCGATTAGAGTGACCATGAGTAGGAGTTCAAAATCTCCTGACA TCATTTTCTTAAAACCTGCTTTGTTTTTTACATTTCTATTTAATATAACTCCTATTTGAAT AAAAAAACAAAACAAGTTTAGATGTTAAGATATTAACTACATCCTTTGCTCCAAAGGG AGAGGGGAAGTTATGGAGTTAATTAATTTGCTGTTGGAAATCAATATGGAGTCAGAA ATATAATGATTTACTAAACCTTATTGAATCGGTAACGATGCGAATTTATATTAAAATA GCTTTTATGAAACATTCAACAAAAATATTATTAATGTTGGCCCACTTTAGCAACCGGTT AGGTCTACCGGTTGGGCAAGCAAAGATTCACGCCCTGGTTCGAGTCCCAACTAGTCC TGCAAAATACCGCAGCAAGTTTTAGAGAGACCAAGTGCCATTACCTCTCCCACTTCAG TTATCGGTTATGCGGCGTTTAAGTCGACAGCTTGCCGTCTCTAGCTCCGGTGCCTATA TAAAGCAGCCCGCTTTCCACATTTCATATTCGTTTTACGTTTGTCAAGCCTCATAGCCG GCAGTTCGAACGTATACGCTCTCTGAGTCAGACCTCGAAATCGTAGCTCTACACAATT CTGTGAATTTTCCTTGTCGCGTGTGAAACACTTCCAAGGCGGCCGCCCGACTCTAGAT CATAATCAGCCATACCACATTTGTAGAGGTTTTACTTGCTTTAAAAAACCTCCCACACC TCCCCCTGAACCTGAAACATAAAATGAATGCAATTGTTGTTGTTAACTTGTTTATTGCA GCTTATAATGGTTACAAATAAAGCAATAGCATCACAAATTTCACAAATAAAGCATTTT TTTCACTGCATTCTAGTTGTGGTTTGCCCAAACTCATCAATGTATCTTAGCGGCTCGAG GGTACCTCTAG
<u>P. Bru Lac</u> *Polyporus brumalis* laccase gene, codon optimised for *D. melanogaster* with overlaps to pMC-1-1-1 and larval cuticle 9 signal peptide sequence. Generates pMC-1-2-6	gtgtgaaacacttccaagGCaatcttacaaaATGAAGTTTGTTATTGTTCTGGCTTGCCTGTTGG CAGTTGTCTTTGCCGCTATTGGCCCTGTTACCGACTTGACGGTTACGGATGCCAACAT CAGTCCCGACGGATTTACGCGCGCCGGTATAGTGGTCAACAACGTGTTCCCCGCTCC CCTGATTACGGGACAAAAAGGCGACAGGTTTCAGCTGAATCTGGTCAACCAAATGAG TAATCATACGATGTTGAAGACAACTTCGATACACTGGCACGGTTTTTTTCAAAAGGGA ACTAATTGGGCGGATGGACCCGCATTCGTGAATCAGTGTCCTATCGCTAGCGGTAAT TCGTTTCTGTATGACTTTCAGGTGCCCGATCAAGCTGGAACTTTCTGGTACCACTCCCA CCTGTCCACGCAGTATTGTGACGGTCTCCGCGGACCTTTCGTAGTTTATGACCCAACT GACCCTCACCTGTCGCTGTATGACGTGGATGACGATAGCACAGTTATAACGCTGGCG GATTGGTACCATGTGGCCGCACGCCTGGGACCGCGCTTCCCGTTGGGTGCTGACTCC ACTCTGATCAACGGTTTGGGCAGGTCCACCGCGACTCCGACTGCTGATCTGGCTGTG ATCAGTGTAACCAAAGGTAAACGTTACCGGTTTAGGCTGGTGAGTATTTCCTGCGAT CCGAACCATACCTTCAGCATTGACGGTCACAAGCTCACCGTCATCGAAGCTGATGGC ATTTCCACTCAACCCGTCACTGGCATTGACTCCATTCAAATTTTTGCCGCACAGAGGTA CAGCTTCGTCCTCACCGCCGACCAGGATGTAGACAATTACTGGGTTAGGGCTAACCCT AATTTTGGTACTACTGGATTTGCCGGAGGCATTAACAGCGCGATTCTCCGTTATGATG GTGCACCTGCGGTTGAACCACTACCTCCCAAACCGGTACGAATCTGCTCGTGGAGA CAGATCTGCATCCTCTGAGCACAATGCCCGTTCCGGGACTGCCGACACAAGGTGGTG CCGATTTTAATCTCAATCTCGCCTTTAACTTTAACGGTTCGGACTTCTTTATTAACGGT GCTTCGTTCGTTCCACCGACCGTCCCTGTCCTCCTCCAGATAATCAGCGGTGCGAACA GCGCCCAAGATTTGCTCCCAAGTGGTTCGGTATATGCACTCCCCTCGAATAGCAGCAT CGAACTCACATTCCCGGCCACTGCTGCAGCGCCAGGCGCCCCCCATCCATTCCACTTG CATGGCCACGCTTTTGCCGTAGTACGCAGTGCTGGAAGTACTGTTTATAACTACGACA

		ACCCGGTATTCAGGGATGTAGTTTCCACTGGTACGCCCGCTGCGGGCGACAACGTCACCATCCGATTCCAAACGGACAATCCTGGACCATGGTTTTTGCATTGTCACATCGACTTCCACCTCGATGCTGGCTTCGCTGTCGTGTTCGCGGAAGACCTCCCGGACGTTGTAAGTGCAAACCCCGTGCCCCAAGCGTGGAGCGATTTGTGCCCCATCTATAACGCTCTCGATCCGTCGGATCAGTAAGCccgactctagatcataatc
T. tro. Lac *Trametes trogii* laccase gene, codon optimised for *D. melanogaster* with overlaps to pMC-1-1-1 and larval cuticle 9 signal peptide sequence. Generates pMC-1-2-3	gtgtgaaacacttccaagGCaatcttacaaaATGAAGTTTGTTATTGTTCTGGCTTGCCTGTTGGCAGTTGTCTTTGCCGCTATTGGTCCAGTTGCAGATCTCACTATCAGCAACGGAGCGGTTAGTCCTGATGGATTCTCCCGCCAAGCCATATTGGTGAACGATGTGTTCCCGTCGCCGTTGATAACGGGAAATAAAGGTGACAGGTTTCAGCTCAACGTAATAGACAACATGACAAATCACACGATGCTCAAATCCACAAGTATCCATTGGCATGGTTTTTTCCAACATGGCACAAATTGGGCTGATGGCCCTGCATTCGTCAACCAATGTCCCATTAGTACGGGCCACGCTTTTCTGTATGATTTTCAGGTTCCGGATCAGGCGGGTACCTTCTGGTATCACTCGCATTTGAGCACTCAGTACTGCGACGGCCTGCGTGGTCCAATCGTAGTCTATGATCCCCAGGACCCGCATAAAAGTCTGTATGATGTTGATGATGACAGTACAGTGATAACCCTCGCCGACTGGTATCATTTGGCAGCTAAGGTCGGCTCGCCGGTACCAACAGCCGACGCAACACTCATAAATGGCCTGGGTAGGTCCATAGACACCCTGAACGCAGATTTGGCGGTTATTACCGTTACAAAAGGCAAGAGGTATCGCTTCCGTTTGGTCAGTTTGAGTTGCGACCCAAATCATGTTTTCTCCATTGATGGTCATTCGTTGACCGTAATTGAGGCGGATTCCGTAAACTTGAAGCCTCAAACGGTAGATAGTATCCAAATCTTTGCGGCCCAACGGTATAGCTTCGTCCTGAATGCTGATCAAGATGTGGGCAACTACTGGATTCGAGCTCTGCCGAATAGTGGCACACGCAACTTTGACGGAGGTGTTAATTCGGCGATCCTCCGCTATGATGGCGCCGCGCCCGTCGAACCGACAACCAGTCAAACGCCCAGCACGAATCCACTCGTAGAGTCGGCCCTGACAACATTGGAGGGAACAGCAGCGCCGGGTAGCCCTGCTCCGGGAGGTGTAGACCTCGCTCTGAACATGGCGTTTGGCTTTGCTGGCGGCAAGTTTACCATAAATGGAGCAAGCTTTACTCCCCCGACCGTCCCTGTGCTGTTGCAGATACTCTCCGGTGCGCAGTCGGCCCAAGATCTGCTCCCGTCGGGCTCCGTTTATTCCCTGCCCGCCAATGCAGACATCGAAATCAGCTTGCCAGCAACGGCTGCCGCTCCCGGTTTCCCACACCCGTTTCATCTCCACGGTCATACTTTTGCTGTGGTTCGCAGCGCAGGCTCCAGTACATACAATTATGAGAATCCCGTCTACCGTGACGTCGTTTCGACTGGCTCGCCCGGCGATAATGTCACCATCCGTTTCCGAACGGATAACCCGGGCCCATGGTTTCTGCATTGTCATATAGACTTTCATTTGGAGGCGGGTTTTGCAGTGGTTATGGCCGAGGATATCCCGGAGGTTGCCGCCACTAACCCCGTACCACAAGCATGGTCGGATCTGTGCCCCACATACGACGCGCTGTCGCCGGACGATCAGTAAGCccgactctagatcataatc

References:

1. UN General Assembly. *Transforming our world : the 2030 Agenda for Sustainable Development*. (2015).

2. Silpa Kaza, Yao, L., Bhada-Tata, P. & Woerden, F. Van. *What a Waste 2.0: A Global Snapshot of Solid Waste Management to 2050*. (Urban Dev. World Bank, Washington, DC, 295, 2018).

3. Weitz, M., Coburn, J. B. & Salinas, E. Estimating National Landfill Methane Emissions: An Application of the 2006 Intergovernmental Panel on Climate Change Waste Model in Panama. *J. Air Waste Manage. Assoc.* **58**, 636–640 (2008).

4. Du, M. *et al.* Quantification of methane emissions from municipal solid waste landfills in China during the past decade. *Renewable and Sustainable Energy Reviews* **78**, 272–279 (2017).

5. Przydatek, G. & Kanownik, W. Impact of small municipal solid waste landfill on groundwater quality. *Environ. Monit. Assess.* **191**, 1–14 (2019).

6. Landrigan, P. J. *et al.* The Lancet Commission on pollution and health. *Lancet* **391**, 462–512 (2018).

7. Fuller, R. Hazardous Waste and Toxic Hotspots. in *Textbook of Children's Environmental Health* 254–261 (Oxford University Press, 2014). doi:10.1093/med/9780199929573.003.0027

8. Lanphear, B. P. *et al.* Low-level environmental lead exposure and children's intellectual function: An international pooled analysis. *Environ. Health Perspect.* **113**, 894–899 (2005).

9. Kingsford, R. T. *et al.* Major Conservation Policy Issues for Biodiversity in Oceania. *Conserv. Biol.* **23**, 834–840 (2009).

10. Malaj, E. *et al.* Organic chemicals jeopardize the health of freshwater ecosystems on the continental scale. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 9549–54 (2014).

11. (CIE), C. of I. E. NSW waste regulation. (2014).

12. Stahel, W. R. The circular economy. *Nature* **531**, 435–438 (2016).

13. Clark, M. & Maselko, M. Transgene Biocontainment Strategies for Molecular Farming. *Frontiers in Plant Science* **11**, 210 (2020).

14. Lee, J. W., Chan, C. T. Y., Slomovic, S. & Collins, J. J. Next-generation biocontainment systems for engineered organisms. *Nat. Chem. Biol.* **14**, 530–537 (2018).

15. Maselko, M. *et al.* Engineering multiple species-like genetic incompatibilities in insects. *Nat. Commun.* **11**, 1–7 (2020).

16. Diener, S., Studt Solano, N. M., Roa Gutiérrez, F., Zurbrügg, C. & Tockner, K. Biological treatment of municipal organic waste using black soldier fly larvae. *Waste and Biomass Valorization* **2**, 357–363 (2011).

17. Parodi, A. *et al.* Black soldier fly larvae show a stronger preference for manure than

for a mass-rearing diet. *J. Appl. Entomol.* **144**, 560–565 (2020).

18.   Meyer, H.-P., Minas, W. & Schmidhalter, D. Industrial-Scale Fermentation. in
Industrial Biotechnology 1–53 (Wiley-VCH Verlag GmbH & Co. KGaA, 2016).
doi:10.1002/9783527807833.ch1

19.   Dahman, Y., Syed, K., Begum, S., Roy, P. & Mohtasebi, B. 14 - Biofuels: Their
characteristics and analysis. in *Woodhead Publishing Series in Composites Science and*
Engineering (eds. Verma, D., Fortunati, E., Jain, S. & Zhang Biopolymer-Based
Materials, and Bioenergy, X. B. T.-B.) 277–325 (Woodhead Publishing, 2019).
doi:https://doi.org/10.1016/B978-0-08-102426-3.00014-X

20.   Joosten, L. *et al.* Review of insect pathogen risks for the black soldier fly (Hermetia
illucens) and guidelines for reliable production. *Entomol. Exp. Appl.* **168**, 432–447
(2020).

21.   Dortmans, B., Diener, S., Verstappen, B. & Zurbrugg, C. *Black Soldier Fly Biowaste*
Processing. (2017).

22.   Kim, C.-H. *et al.* Use of Black Soldier Fly Larvae for Food Waste Treatment and Energy
Production in Asian Countries: A Review. *Processes* **9**, (2021).

23.   Maeda, S. *et al.* Production of human α-interferon in silkworm using a baculovirus
vecto. *Nature* **315**, 592–594 (1985).

24.   Zeng, S., Zhao, J. & Xia, L. Simultaneous production of laccase and degradation of
bisphenol A with Trametes versicolor cultivated on agricultural wastes. *Bioprocess*
Biosyst. Eng. **40**, 1237–1245 (2017).

25.   Chivukula, M. & Renganathan, V. Phenolic Azo Dye Oxidation by Laccase from
Pyricularia oryzae. *Appl. Environ. Microbiol.* **61**, 4374 LP – 4377 (1995).

26.   Alharbi, S. K. *et al.* Degradation of diclofenac, trimethoprim, carbamazepine, and
sulfamethoxazole by laccase from Trametes versicolor: Transformation products and
toxicity of treated effluent. *Biocatal. Biotransformation* **37**, 399–408 (2019).

27.   Collins, P. J., Kotterman, M., Field, J. A. & Dobson, A. Oxidation of Anthracene and
Benzo[a]pyrene by Laccases from Trametes versicolor. *Appl. Environ. Microbiol.* **62**,
4563 LP – 4567 (1996).

28.   Cañas, A. I. *et al.* Transformation of Polycyclic Aromatic Hydrocarbons by Laccase Is
Strongly Enhanced by Phenolic Compounds Present in Soil. *Environ. Sci. Technol.* **41**,
2964–2971 (2007).

29.   Luo, Q. *et al.* Laccase-Catalyzed Degradation of Perfluorooctanoic Acid. *Environ. Sci.*
Technol. Lett. **2**, 198–203 (2015).

30.   Luo, Q., Yan, X., Lu, J. & Huang, Q. Perfluorooctanesulfonate Degrades in a Laccase-
Mediator System. *Environ. Sci. Technol.* **52**, 10617–10626 (2018).

31.   Sumathi, T., Viswanath, B., Sri Lakshmi, A. & SaiGopal, D. V. R. Production of Laccase
by *Cochliobolus* sp. Isolated from Plastic Dumped Soils and Their Ability to Degrade
Low Molecular Weight PVC. *Biochem. Res. Int.* **2016**, 9519527 (2016).

32. Jin, X. *et al.* Conditions Optimizing and Application of Laccase-mediator System (LMS)
for the Laccase-catalyzed Pesticide Degradation. *Sci. Rep.* **6**, 35787 (2016).

33. Alberts, J. F., Gelderblom, W. C. A., Botha, A. & van Zyl, W. H. Degradation of aflatoxin
B1 by fungal laccase enzymes. *Int. J. Food Microbiol.* **135**, 47–52 (2009).

34. Rodríguez Couto, S. & Toca Herrera, J. L. Industrial and biotechnological applications
of laccases: A review. *Biotechnol. Adv.* **24**, 500–513 (2006).

35. Widsten, P. & Kandelbauer, A. Laccase applications in the forest products industry: A
review. *Enzyme Microb. Technol.* **42**, 293–307 (2008).

36. Witayakran, S. & Ragauskas, A. J. Synthetic Applications of Laccase in Green
Chemistry. *Adv. Synth. Catal.* **351**, 1187–1209 (2009).

37. Kharayat, Y. Distillery wastewater: bioremediation approaches. *J. Integr. Environ. Sci.*
9, 69–91 (2012).

38. Baghel, S., Sahariah, B. P. & Anandkumar, J. Bioremediation of Lignin-Rich Pulp and
Paper Industry Effluent BT - Combined Application of Physico-Chemical &
Microbiological Processes for Industrial Effluent Treatment Plant. in (eds. Shah, M. &
Banerjee, A.) 261–278 (Springer Singapore, 2020). doi:10.1007/978-981-15-0497-
6_12

39. Ahmed, P. M., Fernández, P. M., de Figueroa, L. I. C. & Pajot, H. F. Exploitation
alternatives of olive mill wastewater: production of value-added compounds useful
for industry and agriculture. *Biofuel Res. J.* **6**, 980–994 (2019).

40. Kudanga, T. & Le Roes-Hill, M. Laccase applications in biofuels production: current
status and future prospects. *Appl. Microbiol. Biotechnol.* **98**, 6525–6542 (2014).

41. Viswanath, B., Rajesh, B., Janardhan, A., Kumar, A. P. & Narasimha, G. Fungal laccases
and their applications in bioremediation. *Enzyme Res.* **2014**, (2014).

42. Rochester, J. R. Bisphenol A and human health: A review of the literature.
Reproductive Toxicology **42**, 132–155 (2013).

43. Chowdhury, M. F. *et al.* Current treatment technologies and mechanisms for removal
of indigo carmine dyes from wastewater: A review. *Journal of Molecular Liquids* **318**,
114061 (2020).

44. Gorman, M. J., Dittmer, N. T., Marshall, J. L. & Kanost, M. R. Characterization of the
multicopper oxidase gene family in Anopheles gambiae. *Insect Biochem. Mol. Biol.* **38**,
817–24 (2008).

45. Waters, A. J., Capriotti, P., Gaboriau, D. C. A., Papathanos, P. A. & Windbichler, N.
Rationally-engineered reproductive barriers using CRISPR & CRISPRa: an
evaluation of the synthetic species concept in Drosophila melanogaster. *Sci. Rep.* **8**,
13125 (2018).

46. Charles, J. P., Chihara, C., Nejad, S. & Riddiford, L. M. Identification of proteins and
developmental expression of RNAs encoded by the 65A cuticle protein gene cluster in
Drosophila melanogaster. *Insect Biochem. Mol. Biol.* **28**, 131–8 (1998).

47. Almeida, S., Raposo, A., Almeida-González, M. & Carrascosa, C. Bisphenol A: Food
Exposure and Impact on Human Health. *Compr. Rev. Food Sci. Food Saf.* **17**, 1503–
1517 (2018).

48. Atacag Erkurt, H. Biodegradation and Detoxification of BPA: Involving Laccase and a
Mediator. *CLEAN - Soil, Air, Water* **43**, 932–939 (2015).

49. Garzillo, A. M. *et al.* Structural and Kinetic Characterization of Native Laccases from
Pleurotus ostreatus, Rigidoporus lignosus, and Trametes trogii. *J. Protein Chem.* **20**,
191–201 (2001).

50. Garzillo, A. M. V *et al.* Laccase from the white-rot fungus Trametes trogii. *Appl.*
Microbiol. Biotechnol. **49**, 545–551 (1998).

51. Atacag Erkurt, H. Biodegradation and Detoxification of BPA: Involving Laccase and a
Mediator. *CLEAN – Soil, Air, Water* **43**, 932–939 (2015).

52. Colao, M. C., Lupino, S., Garzillo, A. M., Buonocore, V. & Ruzzi, M. Heterologous
expression of lcc1 gene from Trametes trogii in Pichia pastoris and characterization of
the recombinant enzyme. *Microb. Cell Fact.* **5**, 31 (2006).

53. Nakade, K., Nakagawa, Y., Yano, A., Sato, T. & Sakamoto, Y. Characterization of an
extracellular laccase, PbLac1, purified from Polyporus brumalis. *Fungal Biol.* **114**, 609–
618 (2010).

54. Shmarakov, I. O., Borschovetska, V. L. & Blaner, W. S. Hepatic Detoxification of
Bisphenol A is Retinoid-Dependent. *Toxicol. Sci.* **157**, 141–155 (2017).

55. Bourbonnais, R. & Paice, M. G. Oxidation of non-phenolic substrates: An expanded
role for laccase in lignin biodegradation. *FEBS Lett.* **267**, 99–102 (1990).

56. Morozova, O. V., Shumakovich, G. P., Shleev, S. V. & Yaropolov, Y. I. Laccase-mediator
systems and their applications: A review. *Appl. Biochem. Microbiol.* **43**, 523–535
(2007).

57. Jensen, K. A., Evans, K. M., Kirk, T. K. & Hammel, K. E. Biosynthetic Pathway for
Veratryl Alcohol in the Ligninolytic Fungus Phanerochaete chrysosporium. *Appl.*
Environ. Microbiol. **60**, 709–714 (1994).

58. Eggert, C., Temp, U., Dean, J. F. D. & Eriksson, K.-E. L. A fungal metabolite mediates
degradation of non-phenolic lignin structures and synthetic lignin by laccase. *FEBS*
Lett. **391**, 144–148 (1996).

59. Felby, C., Nielsen, B. R., Olesen, P. O. & Skibsted, L. H. Identification and
quantification of radical reaction intermediates by electron spin resonance
spectrometry of laccase-catalyzed oxidation of wood fibers from beech (Fagus
sylvatica). *Appl. Microbiol. Biotechnol.* **48**, 459–464 (1997).

60. Camarero, S., Ibarra, D., Martínez, M. J. & Martínez, A. T. Lignin-derived compounds
as efficient laccase mediators for decolorization of different types of recalcitrant
dyes. *Appl. Environ. Microbiol.* **71**, 1775–1784 (2005).

61. Aslam, M. S., Aishy, A., Samra, Z. Q., Gull, I. & Athar, M. A. Identification, Purification

and Characterization of a Novel Extracellular Laccase from Cladosporium
Cladosporioides. *Biotechnol. Biotechnol. Equip.* **26**, 3345–3350 (2012).

62. Lindsley, D. L., Grell, E. H. & Bridges, C. B. Genetic Variations of Drosophila
melanogaster. *Science (80-.).* **162**, 993–993 (1968).

63. Venken, K. J. T., He, Y., Hoskins, R. A. & Bellen, H. J. P[acman]: A BAC transgenic
platform for targeted insertion of large DNA fragments in D. melanogaster. *Science
(80-.).* **314**, 1747–1751 (2006).

586